

Biases in the Experimental Annotations of Protein Function and their Effect on Our Understanding of Protein Function Space

1 Alexandra M. Schnoes Department of Bioengineering and Therapeutic Sciences, University of California, San Francisco, San Francisco, CA, USA

2 David C. Ream Department of Microbiology, Miami University, Oxford, OH, USA

3 Alexander W. Thorman, Department of Microbiology, Miami University, Oxford, OH, USA

4 Patricia C. Babbitt Department of Bioengineering and Therapeutic Sciences, University of California, San Francisco, San Francisco, CA, USA

5 Iddo Friedberg, Departments of Microbiology and Computer Science & Software engineering, Miami University, Oxford, OH, USA

* E-mail: corresponding i.friedberg@miamioh.edu

1 Abstract

2 The ongoing functional annotation of proteins relies upon the work of curators to capture
3 experimental findings from scientific literature and apply them to protein sequence and
4 structure data. However, with the increasing use of high-throughput experimental assays,
5 a small number of experimental studies dominate the functional protein annotations col-
6 lected in databases. Here we investigate just how prevalent is the “few articles – many
7 proteins” phenomenon. We examine the experimentally validated annotation of proteins
8 provided by several groups in the GO Consortium, and show that the distribution of pro-
9 teins per published study is exponential, with 0.14% of articles providing the source of

10 annotations for 25% of the proteins in the UniProt-GOA compilation. Since each of the
11 dominant articles describes the use of an assay that can find only one function or a small
12 group of functions, this leads to substantial biases in what we know about the function
13 of many proteins. Mass-spectrometry, microscopy and RNAi experiments dominate high
14 throughput experiments. Consequently, the functional information derived from these
15 experiments is mostly of the subcellular location of proteins, and of the participation
16 of proteins in embryonic developmental pathways. For some organisms, the information
17 provided by different studies overlap by a large amount. We also show that the informa-
18 tion provided by high throughput experiments is less specific than those provided by low
19 throughput experiments. Given the experimental techniques available, certain biases in
20 protein function annotation due to high-throughput experiments are unavoidable. Know-
21 ing that these biases exist and understanding their characteristics and extent is important
22 for database curators, developers of function annotation programs, and anyone who uses
23 protein function annotation data to plan experiments.

24 **Author Summary**

25 Experiments and observations are the vehicles used by science to understand the world
26 around us. In the field of molecular biology, we are increasingly relying on high-throughput,
27 genome-wide experiments to provide answers about the function of biological macro-
28 molecules. However, any experimental assay is essentially limited in the type of infor-
29 mation it can discover. Here we show that our increasing reliance on high-throughput
30 experiments biases our understanding of protein function. While the primary source of
31 information is experiments, the functions of many proteins are computationally annotated
32 by sequence-based similarity, either directly or indirectly, to proteins whose function is

33 experimentally determined. Therefore, any biases in experimental annotations can get
34 amplified and entrenched in the majority of protein databases. We show here that high-
35 throughput studies are biased towards certain aspects of protein function, and that they
36 provide less information than low-throughput studies. While there is no clear solution to
37 the phenomenon of bias from high-throughput experiments, recognizing its existence and
38 its impact can help take steps to mitigate its effect.

39 **Introduction**

40 Functional annotation of proteins is an open problem and a primary challenge in molecular
41 biology today [1–4]. The ongoing improvements in sequencing technology have shifted
42 the emphasis shifting from realizing the \$1,000 genome to realizing the 1-hour genome [5].
43 The ability to rapidly and cheaply sequence genomes is creating a flood of sequence
44 data, but to make these data useful, extensive analysis is needed. A large portion of
45 this analysis involves assigning biological function to newly determined gene sequences,
46 a process that is both complex and costly [6]. To aid current annotation procedures and
47 improve computational function prediction algorithms, high-quality and experimentally
48 derived data are necessary. Currently, one of the few repositories of such data is the
49 UniProt-GOA database [7], which is a compilation of data contributed by several member
50 groups of the GO consortium. UniProt-GOA contains functional information derived
51 from literature, and by computational means. The information derived from literature is
52 extracted by human curators who capture functional data from publications, assign the
53 data to their appropriate place in the Gene Ontology hierarchy [8] and label them with
54 appropriate functional evidence codes. UniProt-GOA is compiled from annotations made
55 by several member groups of the GO consortium, and as such presents the current state of

56 our view of protein function space. It is therefore important to understand any trends and
57 biases that are encapsulated in UniProt-GOA, as those impact well-used sister databases
58 and consequently a large number of users worldwide.

59 One concern surrounding the capture of functional data from articles is the propensity
60 for high-throughput experimental work to become a large fraction of the data in the GO
61 Consortium database, thus having a small number of experiments dominate the protein
62 function landscape. In this work we analyzed the relative contribution of peer-reviewed
63 articles describing all the experimentally derived annotations in UniProt-GOA. We found
64 some striking trends, stemming from the fact that a small fraction of articles describing
65 high-throughput experiments disproportionately contribute to the pool of experimental
66 annotations of model organisms. Consequently we show that: 1) annotations coming
67 from high-throughput experiments are overall less informative than those provided by
68 low-throughput experiments; 2) annotations from high-throughput experiments are biased
69 towards a limited number of functions, and, 3) many high-throughput experiments overlap
70 in the proteins they annotate, and in the annotations assigned. Taken together, our
71 findings offer a picture of how the protein function annotation landscape is generated
72 from scientific literature. Furthermore, due to the biases inherent in the current system
73 of sequence annotations, this study serves as a caution to the producers and consumers
74 of biological data from high-throughput experiments.

75 **Results**

76 **Articles and Proteins**

77 The increase in the number of high-throughput experiments used to determine protein
78 functions may introduce biases into experimental protein annotations, due to the inher-

ent capabilities and limitations of high-throughput assays. To test the hypothesis that such biases exist, and to study their extent if they do, we compiled the details of all experimentally annotated proteins in UniProt-GOA. This included all proteins whose GO annotations have the GO experimental evidence codes EXP, IDA, IPI, IMP, IGI, IEP (See Methods for an explanation of GO evidence codes). We first examined the distribution of articles that are the source of experimentally validated annotations by the number of proteins they annotate. As can be seen in Figure 1, the distribution of the number of proteins annotated per article follows a power-law distribution. $f(x) = ax^k$. Using linear regression over the log values of the axes we obtained a fit with $p < 1.18 \times 10^{-8}$ and $R^2 = -0.72$. We therefore conclude that there is indeed a substantial bias in experimental annotations, in which there are few articles that annotate a large number of proteins.

To better understand the consequences of such a distribution, we divided the annotating articles into four cohorts, based on the number of proteins each article annotates. *Single-throughput* articles are those articles that annotate only one protein; *low throughput* articles annotate 2-9 proteins; *moderate throughput* articles annotate 10-99 proteins and *high throughput* articles annotate over 99 proteins. The results are shown in Table 1. The most striking finding is that high throughput articles are responsible for 25% of the annotations that the GO Consortium creates, even though they are found only in 0.14% of the articles. 96% of the articles are single-throughput and low-throughput, however those annotate only 53% of the proteins. So while moderate-throughput and high-throughput studies account for almost 47% of the annotations in Uniprot-GOA, they constitute only 3.66% of the studies published.

To understand how the log-odds distribution affects our understanding of protein function, we examined different aspects of the annotations in the four article cohorts. Also, we examined in greater detail the top-50 high-throughput annotating articles. “Top-50

104 high throughput annotating articles” are the articles describing experimental annotations
105 that are top ranked by the number of proteins annotated per article. An initial charac-
106 terization of these articles is shown in Table S1. As can be seen, most of the articles are
107 specific to a single species (typically a model organism) and to a single assaying pipeline
108 that is used to assign function to the proteins in that organism. With one exception, only
109 one ontology of the three GO ontologies was used for annotation in any single experiment.
110 The three ontologies are Molecular Function (MF), Biological Process (BP) and Cellular
111 Component (CC). These are separate ontologies within GO, describing different aspects
112 of function as detailed in [8]. As we show later, for some species this means that a single
113 functional aspect (MF, BP or CC) of a species can be dominated by a single study.

114 **The Impact of High Throughput Studies on the Annotation of** 115 **Model Organisms**

116 We examined the relative contribution of the top-50 articles to the entire corpus of ex-
117 perimentally annotated proteins in each species. Unsurprisingly, all the species found in
118 the top-50 articles were either common model organisms or human. For each species,
119 we examined the five most frequent terms in the top-50 articles. We then examined
120 the contribution of this term by the top-50 articles to the general annotations of that
121 species. The *contribution* is the number of annotations by any given GO term in the
122 top 50 articles divided by the number of annotations by that GO term in all of UniProt-
123 GOA. For example, as seen in Figure 2 in *D. melanogaster* 88% of the annotations using
124 the term “precatalytic spliceosome” in articles experimentally annotating this species are
125 contributed by the top-50 articles.

126 For most organisms annotated by the top-50 articles, the annotations were within the

127 cellular component or biological process ontologies. Notable exceptions are *D. melanogaster*
128 and *C. elegans* where the dominant terms were from the Biological Process ontology, and
129 in mouse, where “protein binding” and “identical protein binding” are from the Molecular
130 Function Ontology. *D. melanogaster*’s annotation for the top terms is dominated (over
131 50% contribution) by the top-50 articles.

132 The term frequency bias described here can be viewed more broadly within the ontol-
133 ogy bias. The proteins annotated by the cohorts of single-protein articles, low-throughput
134 articles, and moderate-throughput articles have similar ratios of the fraction of proteins
135 annotated. Twenty-two to twenty-six percent of assigned terms are in the Molecular
136 Function Ontology, and 51-57% are in the Biological Process Ontology and the remaining
137 17-25% are in the Cellular Component ontology. These ratios change dramatically with
138 high-throughput articles (over 99 terms per article). In the high-throughput articles, only
139 5% of assigned terms are in the Molecular Function Ontology, 38% in the Biological Pro-
140 cess Ontology and 57% in the Cellular Compartment Ontology, ostensibly due to a lack of
141 high-throughput assays that can be used for generating annotations using the Molecular
142 Function Ontology.

143 **Repetition and Consistency in Top-50 Annotations**

144 How many of the top-50 articles actually annotate the same set of proteins? Answering
145 this question will tell us how repetitive experiments are in identifying the same set of
146 proteins to annotate. However, even when annotating the same set of proteins and within
147 the same ontology, different experiments may provide different results, lacking consistency.
148 Therefore, the annotation consistency was also checked. Repetition is given as $\frac{n}{N}$ with n
149 being the number of proteins annotated by two or more articles, and N being the total
150 number of proteins. The results of the repetition analysis are shown in Figure 3 and in

151 Table 2. As can be seen, the highest repetition (65%) is in the 12 articles annotating
152 *C. elegans*. Of course, a higher number of articles is expected to increase repetitive
153 annotations simply due to increased sampling of the genome. However, the goal of this
154 analysis is to present the degree of repetition, rather than to try to rank and normalize
155 it. As an additional repetition metric, Table 2 also lists the mean number of sequences
156 per cluster. When normalized by number of annotating articles, the highest repetition
157 is found in Mouse (15.33% in three articles) closely followed by *M. tuberculosis* (14% in
158 two articles). Taken together, these results show that there is repetition in choosing the
159 proteins that are to be annotated in most model organisms using high-throughput assays,
160 although the rate of this repetition varies widely.

161 Consistency analysis took place as described in Methods. The consistency measure
162 is normalized on a 0-1 scale, with 1 being most consistent, meaning that all annotations
163 from all sources are identical. Table 3 shows the results of this analysis. In *A. thaliana*,
164 1941 proteins are annotated by 15 articles and 18 terms in the Cellular Component on-
165 tology. The mean maximum-consistency is 0.251. The highest mean consistency is for
166 the annotation of 807 mouse proteins annotated in Cellular Component ontology with
167 an annotation consistency 0.832. However, that is not surprising given that there are
168 only three annotating articles, and two annotating terms. We omitted the ontology and
169 organism combinations that were annotated by less than three articles or two GO terms,
170 or both.

171 **Quantifying Annotation Information**

172 A common assumption holds that while high-throughput experiments do annotate more
173 protein functions than low-throughput experiments, the former also tend to be more
174 shallow in the predictions they provide. The information provided, for example, by a

175 large-scale protein binding assay will only tell us if two proteins are binding, but will
176 not reveal whether that binding is specific, will not provide an exact K_{bind} , will not say
177 under what conditions binding takes place, or whether there is any enzymatic reaction
178 or signal-transduction involved. Having on hand data from experiments with different
179 “throughputness” levels, we set out to investigate whether there is indeed a difference in
180 the information provided by high-throughput experiments vs. low-throughput ones. We
181 examined the information provided by GO terms in each paper cohort using two methods:
182 edge-count, and information-content. See Methods for details.

183 The results of both analyses are shown in Figure 4. In general, the results from the
184 edge count analysis and the information-content based analysis are in agreement when
185 compared across annotation cohorts. For the Molecular Function ontology, the distribu-
186 tion of edge counts and log-frequency scores decreases as the number of annotated proteins
187 per-article increases. For the Biological Process ontology, the decrease is significant. How-
188 ever the contributors to the decrease are the high-throughput articles while there is little
189 change in the first three article cohorts. Finally, there is no significant trend of GO-depth
190 decrease in the Cellular Component Ontology. However, using the information-content
191 metric, there is also a significant decrease in information-content in the high-throughput
192 article cohort.

193 **Exclusive High Throughput Annotations**

194 Of interest is the fraction of proteins that are exclusively annotated by high-throughput
195 experiments. The question here is: from the experimentally annotated proteins in an or-
196 ganism, how much do we know of their function *only* using high-throughput experiments?
197 We have seen that high-throughput experiments annotate a large number of proteins, but
198 still some 80% of experimentally determined proteins are annotated via medium-, low- and

199 single-throughput experiments. Given the lower information-content of high-throughput
200 experiments, it is important to know which organisms have a substantial fraction of the
201 proteins experimentally annotated by high throughput studies only. To do so, we ana-
202 lyzed all species with more than 200 genes in the NCBI taxa database for the fraction
203 of the genes that are exclusively annotated by high throughput studies. The results are
204 shown in Table 4.

205 As can be seen, although the fraction of high-throughput annotated proteins is large,
206 not many species are affected with a large fraction of proteins that are exclusively anno-
207 tated by high-throughput studies. However, the few species that are affected are important
208 study and model species. It is important to note that some redundancy due to isoforms,
209 mutants and duplications may exist.

210 **Frequently Used High-Throughput Experiments**

211 The twenty GO evidence codes, discussed above, encapsulate the means by which the
212 function was inferred, but they do not capture all the necessary information. For example,
213 “Inferred by Direct Assay” (IDA) informs that some experimental assay was used, but
214 does not say which type of assay. This information is often needed, since knowing which
215 experiments were performed can help the researcher establish the reliability and scope
216 of the produced data. RNA, used in an RNAi experiment does not traverse the blood-
217 brain-barrier, meaning that no data from the central nervous system can be drawn from an
218 RNAi experiment. The Evidence Code Ontology, or ECO, seeks to improve upon the GO-
219 attached evidence codes. ECO provides more elaborate terms than “Inferred by Direct
220 Assay”: ECO also conveys which assay was used, for example “microscopy” or “RNA
221 interference”. In addition to evidence terms, the ECO ontology provides *assertion terms*
222 in which the nature of the assay is given. For example, an enzyme-linked immunosorbent

223 assay (ELISA) provides quantitative protein data *in vitro* while an immunogold assay may
224 provide the same information, and cellular localization information *in situ*. We manually
225 assigned Evidence Codes Ontology (ECO) assertion and evidence terms to the top-50
226 articles. The assignment is shown in detail in Table S2. Table S3 shows the sorted count
227 of ECO terms in the top-50 papers.

228 The most frequent ECO term used is ECO:0000160 “protein separation followed by
229 fragment identification evidence”: this fits the 27 papers that essentially describe mass-
230 spectrometry studies. Consequently this means that the assignment procedure is limited
231 to the cellular compartments that can be identified with the fractionation methods used.
232 So while Cellular Component is the most frequent annotation used, fractionation and
233 mass-spectrometry is the most common method used to localize proteins in subcellu-
234 lar compartments. A notable exception to the use of fractionation and MS for protein
235 localization is in the top annotating article [9], which uses microscopy for subcellular
236 localization.

237 The second most frequent experimental ECO term is “Imaging assay evidence” (ECO:000044).
238 Several types of studies fall under this ECO. Those include microscopy, RNAi, some of the
239 mass-spectrometry studies that used microscopy, and a yeast-2-hybrid study. As imaging
240 information is used in a variety of studies, this ECO term is not informative of the chief
241 method used in any study, but rather the importance of imaging assays in a variety of
242 methods. The third most frequent experimental ECO term used was “Cell fractionation
243 evidence” which is closely associated with the top term, “Imaging assay evidence”. The
244 fourth and fifth most frequent ECO term used were “loss-of-function mutant phenotype
245 evidence” (ECO:0000016) and “RNAi evidence” (ECO:000019). These two terms are also
246 closely associated, in RNAi whole-genome gene knockdowns in *C. elegans*, *D. melanogaster*
247 and one in *C. albicans*. RNAi experiments use targeted dsRNA which is delivered to the

248 organism and silences specific genes. Typically the experiments here used libraries of
249 RNAi targeted to the whole exome (for example [10–13]). The phenotypes searched for
250 were mostly associated with embryonic and post-embryonic development. Some studies
251 focused on mitotic spindle assembly [14], lipid storage [15] and endocytic traffic [16]. One
252 study used RNAi to identify mitochondrial protein localization [17]. These studies mostly
253 use the same RNAi libraries, and target the whole *C. elegans* genome using common data
254 resources. Hence the large redundancy observed for *C. elegans* in Table 2. It should be
255 noted that all experiments are associated with computational ECO terms, which describe
256 sequence similarity and motif recognition techniques used to identify the sequences found:
257 “sequence similarity evidence”, “transmembrane domain prediction evidence”, “protein
258 BLAST evidence” etc. These terms are all bolded in Table S3. A strong reliance on
259 computational annotation is therefore an integral part of high throughput experiments.
260 It should be noted that computational annotation here is not used directly for functional
261 annotation, but rather for identifying the protein by a sequence or motif similarity search.
262 The third most frequently used assertion in the top experimental articles was not an exper-
263 imental assertion, but rather a computational one: the term ECO:00053 “computational
264 combinatorial evidence” is defined as “A type of combinatorial analysis where data are
265 combined and evaluated by an algorithm.” This is not a computational prediction *per se*,
266 but rather a combination of several experimental lines of evidence used in a article.

267 Discussion

268 We have identified several annotation biases in GO annotations provided by the GO
269 consortium. These biases stem from the uneven number of annotations produced by dif-
270 ferent types of experiments. It is clear that results from high-throughput experiments

271 contribute substantially to the function annotation landscape, as up to 20% of experi-
272 mentally annotated proteins are annotated by high-throughput assays. At the same time,
273 high throughput experiments produce less information per protein than moderate-, low-
274 and single- throughput experiments as evidenced by the type of GO terms produced in
275 the Molecular Function and Biological Process ontologies. Furthermore, the number of
276 total GO terms used in the high-throughput experiments is much lower than that used in
277 low and medium throughput experiments. Therefore, while high throughput experiments
278 provide a high coverage of protein function space, it is the low throughput experiments
279 that provide more specific information, as well as a larger diversity of terms.

280 We have also identified several types of biases that are contributed by high throughput
281 experiments. First, there is the enrichment of low-information-content GO terms, which
282 means that our understanding of the protein function as provided by high-throughput
283 experiments is more limited than that provided by low-throughput experiments. Second,
284 there is the small number of terms used, when considering the large number of proteins
285 that are being annotated. Third is the general ontology bias towards the cellular com-
286 ponent ontology and, to a lesser extent, the Biological Process ontology: there are very
287 few articles that deal with the Molecular Function ontology. These biases all stem from
288 the inherent capabilities and limitations of the high-throughput experiments. A fourth,
289 related bias is the organism studied: taken together, studies of *C. elegans* and *A. thaliana*
290 studies comprise 36 of the top-50 annotating articles, or 72%.

291 **Information Capture and Scope of GO**

292 We have discussed the information loss that is characteristic of high-throughput experi-
293 ments, as shown in Figure 4. However, another reason for information loss is the inability
294 to capture certain types of information using the Gene Ontology. GO is purposefully

295 limited to three aspects (MF, BP and CC) of biological function, which are assigned per
296 protein. However, other aspects of function may emerge from experiments. Of note is
297 the study, “Proteome survey reveals modularity of the yeast cell machinery” [9]. In this
298 study, the information produced was primarily of protein complexes, and the relationship
299 to cellular compartmentalization and biological networks. At the same time, the only GO
300 term captured in the curation of proteins from this study was “protein binding”. Some,
301 but not all of this information can be captured more specifically using the children of
302 the term “protein binding”, but such a process is arguably laborious by manual curation
303 of the information from a high throughput article. Furthermore, the main information
304 conveyed by this article, namely the types of protein complexes discovered and how they
305 relate to cellular networks, is outside the scope of GO. It is important to realize that while
306 high-throughput experiments do convey less information per protein within the functional
307 scope as defined by GO, they still convey composite information such as possible pathway
308 mappings – information which needs to be captured into annotation databases by means
309 other than GO. In the example above, the information can be captured by a protein in-
310 teraction database, but not by GO terms. Methods such as the Statistical Tracking of
311 Ontological Phrases [18] can help in selecting the appropriate ontology for better infor-
312 mation capture.

313 **Conclusions**

314 Taken together, the annotation trends in high-throughput studies affect our understand-
315 ing of protein function space. This, in turn, affects our ability to properly understand the
316 connection between predictors of protein function and the actual function – the hallmark
317 of computational function annotation. As a dramatic example, during the 2011 Critical
318 Assessment of Function Annotation experiment [19] it was noticed that roughly 20% of

319 the proteins participating in the challenge and annotated with the Molecular Function
320 Ontology were annotated as “protein binding”, a GO term that conveys little informa-
321 tion. Furthermore, it was shown that the major contribution of “protein binding” term
322 to the CAFA challenge data set was due to high-throughput assays. This illustrates how
323 the concentration of a large number of annotations in a small number of studies provides
324 only a partial picture of the function of these proteins. As we have seen, the picture
325 provided from high throughput experiments is mainly of: 1) subcellular localization cell
326 fractionation and MS based localization and 2) developmental phenotypes. While these
327 data are important, we should be mindful of this bias when examining protein function in
328 the database, even those annotations deemed to be of high quality, those with experimen-
329 tal verification. Furthermore, such a large bias in prior probabilities can adversely affect
330 programs employing prior probabilities, as most machine-learning programs do. If the
331 training set for these programs has included a disproportional number of annotations by
332 high-throughput experiments, the results these programs provide will be strongly biased
333 towards a few frequent and shallow GO terms.

334 To remedy the bias created by high throughput annotations, the provenance of an-
335 notations should be described in more detail by curators and curation software. Many
336 function annotation algorithms rely on homology transfer as part of their pipeline to an-
337 notate query sequences [1,19]. Knowing the annotation provenance, including the number
338 of proteins annotated by the original paper can create less biased benchmarks or otherwise
339 incorporate that information into the annotation procedure. The ECO ontology can be
340 used to determine the source of the annotation, and the user or the algorithm can decide
341 whether to rely upon any combinations of “throughputness” and experimental type. Of
342 course, such approaches should be taken cautiously, as sweeping measures can cause the
343 unintended loss of information. We hereby call upon the communities of annotators, com-

344 putational biologists and experimental biologists to be mindful of the phenomenon of the
345 experimental biases described in this study, and to work to understand its implications
346 and impact.

347 **Methods**

348 We used the UniProt-GOA database from December 2011. Data analyses were performed
349 using Python scripts. The following tools were used in the analyses: Biopython [20], mat-
350 plotlib [21]. ECO terms classifying the proteins in the top 50 experiments were assigned
351 to the proteins manually after reading the articles. All data and scripts are available on:
352 <http://github.com/idoerg/Uniprot-Bias/> and on <http://datadryad.org> (the latter upon
353 acceptance).

354 **Use of GO evidence codes**

355 Proteins in UniProt-GOA are annotated with one or more GO terms using a procedure
356 described in Dimmer *et al.* (2012). Briefly, this procedure consists of six steps which
357 include sequence curation, sequence motif analyses, literature-based curation, reciprocal
358 BLAST [22] searches, attribution of all resources leading to the included findings, and
359 quality assurance. If the annotation source is a research article, the attribution includes
360 its PubMed ID. For each GO term associated with a protein, there is also an *evidence code*
361 which the curator assigns to explain how the association between the protein and the GO
362 term was made. Experimental evidence codes include such terms as: *Inferred by Direct As-*
363 *say* (IDA) which indicates that “a direct assay was carried out to determine the function,
364 process, or component indicated by the GO term” or *Inferred from Physical Interaction*
365 (IPI) which “Covers physical interactions between the gene product of interest and another

366 molecule.” (All GO evidence code definitions were taken from the GO site, geneontol-
367 ogy.org.) Computational evidence codes include terms such as *Inferred from Sequence or*
368 *Structural Similarity* (ISS) and *Inferred from Sequence Orthology* (ISO). Although the ev-
369 idence in computational evidence codes is non-experimental, the proteins annotated with
370 these evidence codes are still assigned by a curator, rendering a degree of human oversight.
371 Finally, there are also computational, non-experimental evidence codes, the most preva-
372 lent being *Inferred from Electronic Annotation* (IEA) which is “used for annotations that
373 depend directly on computation or automated transfer of annotations from a database”.
374 IEA evidence means that the annotation is electronic, and was not made or checked by a
375 person. Different degrees of reliability are associated with different evidence codes, with
376 experimental codes generally considered to be of higher reliability than non-experimental
377 codes. (For details see: <http://www.ebi.ac.uk/GOA/ElectronicAnnotationMethods>)

378 **Quantifying GO-term Information**

379 We used two methods to quantify the information given by GO terms. First we used
380 edge counting , where the information contained in a term is dependent on the edge
381 distance of that term from the root. The term “catalytic activity”(one edge distance
382 from the ontology root node) would be less informative than “hydrolase activity” (two
383 edges) and the latter will be less informative than “haloalkane dehalogenase activity”
384 (five edges). We therefore counted edges from the ontology root term to the GO term
385 to determine term information. The larger the number of edges, the more specific –and
386 therefore informative– is the annotation. In cases where several paths lead from the root
387 to the examined GO term, we used the minimal path. We did so for all the annotating
388 articles split into groups by the number of proteins each article annotates.

389 While edge counting provides a measure of term-specificity, this measure is imperfect.

390 The reason is that each of the three GO ontologies is constructed as a directed acyclic
 391 graph (DAG) where different areas of the GO DAG have different connectivities, and terms
 392 may have different depths unrelated to the intuitive specificity of a term. For example “D-
 393 glucose transmembrane transporter activity”, (GO:0055056) is 10 terms deep, while “L-
 394 tryptophan transmembrane transporter activity”, (GO:0015196) is fourteen terms deep.
 395 It is hard to discern whether these differences are meaningful. For this reason, information
 396 content, the logarithm of the inverse of the GO term frequency in the corpus, is generally
 397 accepted as a measure of GO term information content [23,24]. To account for the possible
 398 bias created by the GO-DAG structure, we also used the log-frequency of the terms in
 399 the experimentally annotated proteins in Uniprot-GOA. However, it should be noted that
 400 the log-frequency measure is also imperfect because, as we see throughout this study, a
 401 GO term’s frequency may be heavily influenced by the top annotating articles, injecting
 402 a circularity problem into the use of this metric. Since no single metric for measuring the
 403 information conveyed by a GO term is wholly satisfactory, we used both edge-counting
 404 and information-content in this study.

405 **Annotation Consistency**

406 To examine annotation consistency, we employed the following method: given a protein
 407 P , let G be the terminal (leaf) GO terms g_1, g_2, \dots, g_m that annotate that protein in all
 408 top-50 articles for a single ontology $O \in \{BPO, MFO, CCO\}$. The count of each of these
 409 GO terms per protein per ontology is n_1, n_2, \dots, n_m with n_i being the number of times
 410 GO term g_i annotates protein P .

411 The number of total annotations for a protein in an ontology is $\sum_1^m n_i$. The *maximum*
 412 *annotation consistency* for protein P in ontology O $0 \leq k_{P,O} \leq 1$ is calculated as:

$$k_{P,O} = \frac{\max(n_1, n_2, \dots, n_m)}{\sum^m n_i} \text{ for } \max(n_1, n_2, \dots, n_m) \geq 2$$

413 For example, the protein “Oleate activated transcription factor 3” (UniProtID: P36023)
 414 in *S. cerevisiae* is annotated four times by three articles using the Cellular Component
 415 ontology:

PubMedID	UniProt ID	Ontology	GO term	description
14562095	P36023	CCO	GO:0005634	nucleus
14562095	P36023	CCO	GO:0005737	cytoplasm
16823961	P36023	CCO	GO:0005739	mitochondrion
14576278	P36023	CCO	GO:0005739	mitochondrion

416 The annotation consistency for P36023 is therefore the maximum count of identical
 417 GO terms (*mitochondrion*, 2), divided by the total number of annotations, 4: 0.5.

418 When choosing a measure for annotation consistency, we favored a simple and inter-
 419 pretable measure. We therefore examined identity among leaf terms only, rather than
 420 use a more complex comparison of multiple subgraphs in the GO ontology DAG (Di-
 421 rected Acyclic Graph). Doing so without manual curation is unreliable, and may skew
 422 the perception of similarity [25].

423 Acknowledgments

424 We thank Predrag Radivojac, Nives Škunca, Cristophe Dessimoz, Yanay Ofran, Marc
 425 Robinson-Rechavi, Wyatt Clark, Tony Sawford, Chris Mungall, Rama Balakrishnan, and
 426 members of the Friedberg and Babbitt labs for insightful discussions. We are especially
 427 grateful to Rachael Huntley for careful reading of the manuscript and for providing de-
 428 tailed explanations of annotation provenance and methodology in UniProt.

429 Funding

430 This research was funded, in part, by NSF DBI 1146960 award to IF and NIH R01
431 GM60595 and NSF DBI 0640476 to PCB.

432 References

- 433 1. Friedberg I (2006) Automated protein function prediction—the genomic challenge.
434 Brief Bioinform 7: 225–242.
- 435 2. Schnoes AM, Brown SD, Dodevski I, Babbitt PC (2009) Annotation error in public
436 databases: Misannotation of molecular function in enzyme superfamilies. PLoS
437 Comput Biol 5: e1000605+.
- 438 3. Erdin S, Lisewski AM, Lichtarge O (2011) Protein function prediction: towards
439 integration of similarity metrics. Current Opinion in Structural Biology 21: 180 -
440 188.
- 441 4. Rentzsch R, Orengo CA (2009) Protein function prediction the power of multiplic-
442 ity. Trends in Biotechnology 27: 210 - 219.
- 443 5. Sthl PL, Lundeberg J (2012) Toward the single-hour high-quality genome. Annual
444 Review of Biochemistry 81: 359-378.
- 445 6. Sboner A, Mu X, Greenbaum D, Auerbach R, Gerstein M (2011) The real cost of
446 sequencing: higher than you think! Genome Biology 12: 125+.
- 447 7. Dimmer EC, Huntley RP, Alam-Faruque Y, Sawford T, O'Donovan C, et al. (2012)
448 The uniprot-go annotation database in 2011. Nucleic Acids Research 40: D565–
449 D570.

- 450 8. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, et al. (2000) Gene ontology:
451 tool for the unification of biology. *Nature Genetics* 25: 25–29.
- 452 9. Barbe L, Lundberg E, Oksvold P, Stenius A, Lewin E, et al. (2008) Toward a
453 confocal subcellular atlas of the human proteome. *Mol Cell Proteomics* 7: 499–
454 508.
- 455 10. Fraser AG, Kamath RS, Zipperlen P, Martinez-Campos M, Sohrmann M, et al.
456 (2000) Functional genomic analysis of *C. elegans* chromosome I by systematic RNA
457 interference. *Nature* 408: 325–330.
- 458 11. Maeda I, Kohara Y, Yamamoto M, Sugimoto A (2001) Large-scale analysis of gene
459 function in *Caenorhabditis elegans* by high-throughput RNAi. *Curr Biol* 11: 171–
460 176.
- 461 12. Piano F, Schetter AJ, Morton DG, Gunsalus KC, Reinke V, et al. (2002) Gene
462 clustering based on RNAi phenotypes of ovary-enriched genes in *C. elegans*. *Curr*
463 *Biol* 12: 1959–1964.
- 464 13. Kamath RS, Fraser AG, Dong Y, Poulin G, Durbin R, et al. (2003) Systematic
465 functional analysis of the *Caenorhabditis elegans* genome using RNAi. *Nature* 421:
466 231–237.
- 467 14. Goshima G, Wollman R, Goodwin SS, Zhang N, Scholey JM, et al. (2007) Genes
468 required for mitotic spindle assembly in *Drosophila* S2 cells. *Science* 316: 417–421.
- 469 15. Ashrafi K, Chang FY, Watts JL, Fraser AG, Kamath RS, et al. (2003) Genome-
470 wide RNAi analysis of *Caenorhabditis elegans* fat regulatory genes. *Nature* 421:
471 268–272.

- 472 16. Balklava Z, Pant S, Fares H, Grant BD (2007) Genome-wide analysis identifies
473 a general requirement for polarity proteins in endocytic traffic. *Nat Cell Biol* 9:
474 1066–1073.
- 475 17. Hughes JR, Meireles AM, Fisher KH, Garcia A, Antrobus PR, et al. (2008) A
476 microtubule interactome: complexes with roles in cell cycle and mitosis. *PLoS Biol*
477 6: e98.
- 478 18. Wittkop T, TerAvest E, Evani U, Fleisch K, Berman A, et al. (2013) STOP using
479 just GO: a multi-ontology hypothesis generation tool for high throughput experi-
480 mentation. *BMC Bioinformatics* 14: 53+.
- 481 19. Radivojac P, Clark W, Ronnen-Oron T, Schnoes A, Wittkop T, et al. (2013) A large-
482 scale evaluation of computational protein function prediction. *Nature Methods in*
483 *press*.
- 484 20. Cock P, Antao T, Chang J, Chapman B, Cox C, et al. (2009) Biopython: freely
485 available python tools for computational molecular biology and bioinformatics.
486 *Bioinformatics* 25: 1422.
- 487 21. Hunter JD (2007) Matplotlib: A 2d graphics environment. *Computing In Science*
488 *& Engineering* 9: 90–95.
- 489 22. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, et al. (1997) Gapped
490 blast and psi-blast: a new generation of protein database search programs. *Nucleic*
491 *acids research* 25: 3389–3402.
- 492 23. Lord PW, Stevens RD, Brass A, Goble CA (2003) Investigating semantic simi-
493 larity measures across the gene ontology: the relationship between sequence and
494 annotation. *Bioinformatics* 19: 1275–1283.

- 495 24. Pesquita C, Faria D, Falcão AO, Lord P, Couto FM (2009) Semantic similarity in
496 biomedical ontologies. *PLoS Comput Biol* 5: e1000443+.
- 497 25. Faria D, Schlicker A, Pesquita C, Bastos H, Ferreira AEN, et al. (2012) Mining GO
498 annotations for improving annotation consistency. *PLoS ONE* 7: e40519+.
- 499 26. Matsuyama A, Arai R, Yashiroda Y, Shirai A, Kamata A, et al. (2006) ORFeome
500 cloning and global analysis of protein localization in the fission yeast *Schizosaccha-*
501 *romyces pombe*. *Nat Biotechnol* 24: 841–847.
- 502 27. Pagliarini DJ, Calvo SE, Chang B, Sheth SA, Vafai SB, et al. (2008) A mitochon-
503 drial protein compendium elucidates complex I disease biology. *Cell* 134: 112–123.
- 504 28. Simmer F, Moorman C, van der Linden AM, Kuijk E, van den Berghe PV, et al.
505 (2003) Genome-wide RNAi of *C. elegans* using the hypersensitive *rrf-3* strain reveals
506 novel gene functions. *PLoS Biol* 1: E12.
- 507 29. Huh WK, Falvo JV, Gerke LC, Carroll AS, Howson RW, et al. (2003) Global
508 analysis of protein localization in budding yeast. *Nature* 425: 686–691.
- 509 30. Zybaylov B, Rutschow H, Friso G, Rudella A, Emanuelsson O, et al. (2008) Sorting
510 signals, N-terminal modifications and abundance of the chloroplast proteome. *PLoS*
511 *ONE* 3: e1994.
- 512 31. Sonnichsen B, Koski LB, Walsh A, Marschall P, Neumann B, et al. (2005) Full-
513 genome RNAi profiling of early embryogenesis in *Caenorhabditis elegans*. *Nature*
514 434: 462–469.

- 515 32. Mootha VK, Bunkenborg J, Olsen JV, Hjerrild M, Wisniewski JR, et al. (2003)
516 Integrated analysis of protein composition, tissue diversity, and gene regulation in
517 mouse mitochondria. *Cell* 115: 629–640.
- 518 33. Benschop JJ, Mohammed S, O’Flaherty M, Heck AJ, Slijper M, et al. (2007) Quan-
519 titative phosphoproteomics of early elicitor signaling in Arabidopsis. *Mol Cell Pro-*
520 *teomics* 6: 1198–1214.
- 521 34. Mawuenyega KG, Forst CV, Dobos KM, Belisle JT, Chen J, et al. (2005) My-
522 *cobacterium tuberculosis* functional network analysis by global subcellular protein
523 profiling. *Mol Biol Cell* 16: 396–404.
- 524 35. Ito J, Batth TS, Petzold CJ, Redding-Johanson AM, Mukhopadhyay A, et al. (2011)
525 Analysis of the Arabidopsis cytosolic proteome highlights subcellular partitioning
526 of central plant metabolism. *J Proteome Res* 10: 1571–1582.
- 527 36. Rual JF, Ceron J, Koreth J, Hao T, Nicot AS, et al. (2004) Toward improving
528 *Caenorhabditis elegans* phenome mapping with an ORFeome-based RNAi library.
529 *Genome Res* 14: 2162–2168.
- 530 37. Reinders J, Zahedi RP, Pfanner N, Meisinger C, Sickmann A (2006) Toward the
531 complete yeast mitochondrial proteome: multidimensional separation techniques
532 for mitochondrial proteomics. *J Proteome Res* 5: 1543–1554.
- 533 38. Fernandez-Calvino L, Faulkner C, Walshaw J, Saalbach G, Bayer E, et al. (2011)
534 Arabidopsis plasmodesmal proteome. *PLoS ONE* 6: e18880.
- 535 39. Gu S, Chen J, Dobos KM, Bradbury EM, Belisle JT, et al. (2003) Comprehensive
536 proteomic profiling of the membrane constituents of a *Mycobacterium tuberculosis*
537 strain. *Mol Cell Proteomics* 2: 1284–1296.

- 538 40. Ferro M, Brugiere S, Salvi D, Seigneurin-Berny D, Court M, et al. (2010)
539 ATCHLORO, a comprehensive chloroplast proteome database with subplastidial
540 localization and curated information on envelope proteins. *Mol Cell Proteomics* 9:
541 1063–1084.
- 542 41. Kleffmann T, Russenberger D, von Zychlinski A, Christopher W, Sjolander K, et al.
543 (2004) The *Arabidopsis thaliana* chloroplast proteome reveals pathway abundance
544 and novel protein functions. *Curr Biol* 14: 354–362.
- 545 42. Sasseti CM, Boyd DH, Rubin EJ (2003) Genes required for mycobacterial growth
546 defined by high density mutagenesis. *Mol Microbiol* 48: 77–84.
- 547 43. Mitra SK, Gantt JA, Ruby JF, Clouse SD, Goshe MB (2007) Membrane proteomic
548 analysis of *Arabidopsis thaliana* using alternative solubilization techniques. *J Pro-*
549 *teome Res* 6: 1933–1950.
- 550 44. Ceron J, Rual JF, Chandra A, Dupuy D, Vidal M, et al. (2007) Large-scale RNAi
551 screens identify novel genes that interact with the *C. elegans* retinoblastoma path-
552 way as well as splicing-related components with synMuv B activity. *BMC Dev Biol*
553 7: 30.
- 554 45. Sickmann A, Reinders J, Wagner Y, Joppich C, Zahedi R, et al. (2003) The pro-
555 teome of *Saccharomyces cerevisiae* mitochondria. *Proc Natl Acad Sci USA* 100:
556 13207–13212.
- 557 46. Gavin AC, Aloy P, Grandi P, Krause R, Boesche M, et al. (2006) Proteome survey
558 reveals modularity of the yeast cell machinery. *Nature* 440: 631–636.

- 559 47. Green RA, Kao HL, Audhya A, Arur S, Mayers JR, et al. (2011) A high-resolution
560 *C. elegans* essential gene network based on phenotypic profiling of a complex tissue.
561 *Cell* 145: 470–482.
- 562 48. Simpson JC, Wellenreuther R, Poustka A, Pepperkok R, Wiemann S (2000) Sys-
563 tematic subcellular localization of novel proteins identified by large-scale cDNA
564 sequencing. *EMBO Rep* 1: 287–292.
- 565 49. Marmagne A, Ferro M, Meinnel T, Bruley C, Kuhn L, et al. (2007) A high content
566 in lipid-modified peripheral proteins and integral receptor kinases features in the
567 arabidopsis plasma membrane proteome. *Mol Cell Proteomics* 6: 1980–1996.
- 568 50. Dunkley TP, Hester S, Shadforth IP, Runions J, Weimar T, et al. (2006) Mapping
569 the Arabidopsis organelle proteome. *Proc Natl Acad Sci USA* 103: 6518–6523.
- 570 51. Jaquinod M, Villiers F, Kieffer-Jaquinod S, Hugouvieux V, Bruley C, et al. (2007)
571 A proteomics dissection of Arabidopsis thaliana vacuoles isolated from cell culture.
572 *Mol Cell Proteomics* 6: 394–412.
- 573 52. Heazlewood JL, Tonti-Filippini JS, Gout AM, Day DA, Whelan J, et al. (2004) Ex-
574 perimental analysis of the Arabidopsis mitochondrial proteome highlights signaling
575 and regulatory components, provides assessment of targeting prediction programs,
576 and indicates plant-specific mitochondrial proteins. *Plant Cell* 16: 241–256.
- 577 53. Carter C, Pan S, Zouhar J, Avila EL, Girke T, et al. (2004) The vegetative vacuole
578 proteome of Arabidopsis thaliana reveals predicted and unexpected proteins. *Plant*
579 *Cell* 16: 3285–3303.

- 580 54. Da Cruz S, Xenarios I, Langridge J, Vilbois F, Parone PA, et al. (2003) Proteomic
581 analysis of the mouse liver mitochondrial inner membrane. *J Biol Chem* 278: 41566–
582 41571.
- 583 55. Rual JF, Venkatesan K, Hao T, Hirozane-Kishikawa T, Dricot A, et al. (2005)
584 Towards a proteome-scale map of the human protein-protein interaction network.
585 *Nature* 437: 1173–1178.
- 586 56. Bakthavatsalam D, Gomer RH (2010) The secreted proteome profile of developing
587 *Dictyostelium discoideum* cells. *Proteomics* 10: 2556–2559.
- 588 57. Froehlich JE, Wilkerson CG, Ray WK, McAndrew RS, Osteryoung KW, et al.
589 (2003) Proteomic study of the *Arabidopsis thaliana* chloroplastic envelope mem-
590 brane utilizing alternatives to traditional two-dimensional electrophoresis. *J Pro-*
591 *teome Res* 2: 413–425.
- 592 58. Stroschein-Stevenson SL, Foley E, O’Farrell PH, Johnson AD (2006) Identification
593 of *Drosophila* gene products required for phagocytosis of *Candida albicans*. *PLoS*
594 *Biol* 4: e4.
- 595 59. Rutschow H, Ytterberg AJ, Friso G, Nilsson R, van Wijk KJ (2008) Quantitative
596 proteomics of a chloroplast SRP54 sorting mutant and its genetic interactions with
597 CLPC1 in *Arabidopsis*. *Plant Physiol* 148: 156–175.
- 598 60. Kumar A, Agarwal S, Heyman JA, Matson S, Heidtman M, et al. (2002) Subcellular
599 localization of the yeast proteome. *Genes Dev* 16: 707–719.
- 600 61. Gonczy P, Echeverri C, Oegema K, Coulson A, Jones SJ, et al. (2000) Functional
601 genomic analysis of cell division in *C. elegans* using RNAi of genes on chromosome
602 III. *Nature* 408: 331–336.

- 603 62. Suzuki H, Fukunishi Y, Kagawa I, Saito R, Oda H, et al. (2001) Protein-protein
604 interaction panel using mouse full-length cDNAs. *Genome Res* 11: 1758–1765.
- 605 63. Sarry JE, Kuhn L, Ducruix C, Lafaye A, Junot C, et al. (2006) The early re-
606 sponses of *Arabidopsis thaliana* cells to cadmium exposure explored by protein and
607 metabolite profiling analyses. *Proteomics* 6: 2180–2198.
- 608 64. Chen D, Toone WM, Mata J, Lyne R, Burns G, et al. (2003) Global transcriptional
609 responses of fission yeast to environmental stress. *Mol Biol Cell* 14: 214–229.
- 610 65. Herold N, Will CL, Wolf E, Kastner B, Urlaub H, et al. (2009) Conservation of the
611 protein composition and electron microscopy structure of *Drosophila melanogaster*
612 and human spliceosomal complexes. *Mol Cell Biol* 29: 281–301.
- 613 66. Bayer EM, Bottrill AR, Walshaw J, Vigouroux M, Naldrett MJ, et al. (2006) Ara-
614 bidopsis cell wall proteome defined using multidimensional protein identification
615 technology. *Proteomics* 6: 301–311.


616 **Figures**

Figure 1. Distribution of the number of proteins annotated per article. X-axis: number of annotating articles. Y-axis: number of annotated proteins. The distribution was found to be logarithmic with a significant ($R^2 = 0.72$; $p < 1.10 \times 10^{-18}$) linear fit to the log-log plot. The data came from 76137 articles annotating 256033 proteins with GO experimental evidence codes, in Uniprot-GOA 12/2011.




Figure 2. Relative contribution of top-50 articles to the annotation of major model organisms. The length of each bar represents the percentage of proteins annotated by the top-50 articles in a given organism by a given GO term. GO terms that are present in more than one species are highlighted.





Figure 3. Redundancy in proteins described by the top-50 articles. A circle represents the sum total of articles annotating each organism. Each colored arch is composed of all the proteins in a single article. A line is drawn between any two points on the circle if the proteins they represent have 100% sequence identity. A black line is drawn if they are annotated with a different ontology (for example, in one article the protein is annotated with the MFO, and in another article with BPO); a red line if they are annotated in the same ontology. Example: *S. pombe* is described by two articles, one with few protein (light arch on bottom) and one with many (dark arch encompassing most of circle). Many of the same proteins are annotated by both articles. See Table 2 for numbers.



617

4B




Figure 4. Information provided by articles depending on the number of proteins the articles annotate. Articles are grouped into cohorts: 1: one protein annotated by article; < 10: more than 1, up to 10 annotated; < 100: more than 10, less than 100 annotated; ≥ 100 : 100 or more proteins annotated per article. Blue bars: Molecular Function ontology; Green bars: Biological Process ontology; Red bars: Cellular Component ontology. Information is gauged by **A**: Information Content and **B**: GO depth. See text for details.

618

Table 1. Annotation Cohorts

Articles annotating the following number of proteins	1	$1 < n < 10$	$10 \leq n < 100$	$n \geq 100$	SUM
Number of proteins annotated	20699	46383	26485	31411	124978
Number of annotating articles	41156	32201	2672	108	76137
Percent of proteins annotated	16.56	37.11	21.19	25.13	100
Percent of annotating articles	54.09	42.32	3.51	0.14	100

Number of proteins and annotating articles assigned to each article annotation cohort. Columns: 1: articles annotating a single protein (singletons); $1 < n < 10$ articles annotating more than 1 and less than 10 proteins (low throughput); $10 \leq n < 100$: medium throughput; $n \geq 100$: articles annotating 100 proteins and more (high throughput). As can be seen, high-throughput articles comprise 0.14% of the total articles used for experimental annotations, but annotate 25.13% of the proteins in UniProt-GOA.

Table 2. Sequence Redundancy in Top-50 Annotating Articles

Species	num. articles	num. prot	Clusters at 100%	% redundancy	Mean genes/cluster
<i>C. elegans</i>	12	8416	3338	60	3.74
<i>A. thaliana</i>	16	8879	4694	47	3.92
<i>M. musculus</i>	3	4220	2273	46	2.75
<i>M. tuberculosis</i>	2	2351	1702	28	2.22
<i>S. cerevisiae</i>	5	3542	2550	28	2.33
<i>H. sapiens</i>	4	5593	4509	19	2.36
<i>D. melanogaster</i>	3	1217	1003	18	2.17
<i>S. pombe</i>	2	4502	4281	5	2.00

Species: annotated species; **num. articles** number of annotating articles; **num. prot:** number of proteins annotated by top-50 articles for that species; **Clusters at 100%:** number of clusters of 100% identical proteins; **% redundancy:** the product of column 4 by column 3: this is the percentage of proteins annotated more than once for a given species in the top 50 articles; **Mean genes/cluster:** the mean number of genes per cluster, for clusters having more than a single gene.

Table 3. Annotation Consistency in Top 50 articles

Species	Ont.	num prot	mean $k_{P,O}$	stdv	stderr	num articles	num terms
A. thaliana	CCO	1941	0.251	0.328	0.007	15	18
C. elegans	BPO	1847	0.388	0.239	0.006	12	41
D. melanogaster	BPO	76	0.086	0.22	0.025	3	8
D. melanogaster	CCO	81	0.068	0.234	0.026	3	5
H. sapiens	CCO	167	0.285	0.365	0.028	2	20
M. musculus	CCO	807	0.832	0.291	0.01	3	2
S. cerevisiae	CCO	744	0.759	0.379	0.014	4	15
B. tuberculosis	CCO	532	0.309	0.41	0.018	2	3

Species: annotated species; **Ontology:** annotating GO ontology; **num prot:** number of annotated proteins in that species & ontology that are annotated by more than one paper. **mean, stdv, stderr:** mean number of consistent annotations for a protein in that species and ontology, standard deviation from the mean and standard error. **num articles:** number of annotating articles **num terms** number of annotating terms. Annotations by less than two articles or two terms (or both) for the same protein/ontology combination have been omitted.

Table 4. Fraction of Proteins Exclusively Annotated by High Throughput Studies

Taxon ID	Taxon	XHT	Total Proteins	%XHT
284812	<i>Schizosaccharomyces pombe</i>	2781	4507	61.704
1773	<i>Bacillus tuberculosis</i>	1224	2317	52.8269
6239	<i>Caenorhabditis elegans</i>	2493	5302	47.02
9606	<i>Homo sapiens</i>	4016	11521	34.8581
44689	<i>Dictyostelium discoideum</i>	425	1256	33.8376
3702	<i>Arabidopsis thaliana</i>	3199	10153	31.5079
237561	<i>Candida albicans</i> SC5314	327	1243	26.3073
10090	LK3 transgenic mice	2567	22068	11.6322
7227	<i>Drosophila melanogaster</i>	735	7501	9.7987
559292	<i>Saccharomyces cerevisiae</i>	439	5086	8.6315
83333	<i>Escherichia coli</i> K-12	83	1606	5.1681
7955	<i>Brachidanio rerio</i>	117	4633	2.5254
10116	Buffalo rat	11	4634	0.2374

Taxon ID: NCBI Taxon ID number; **Species:** annotated species; **XHT:** number of proteins exclusively annotated by high-throughput experimental studies (100 or more proteins annotated per study); **Total proteins:** Total number of experimentally annotated proteins in that species. **%XHT:** percentage of proteins in that species that are annotated exclusively by HT studies.

620 **Supplementary Material Legends**

621 **Table S1: The top 50 annotating articles.**

622 **N:** article rank; **Proteins:** number of proteins annotated in this article; **Annotations:**
623 number of annotating GO terms; **Species:** annotated species; **ref.** annotating article;
624 **MFO/BPO/CCO:** number of proteins annotated in the Molecular Function, Biological
625 Process and Cellular Component ontologies, respectively.

626

627 **Table S2: The Top-50 studies and the ECO terms we have assigned to them.**

628 **PMID:** Articles' PubMed ID; **ECO terms/ECO ID's:** terms and ID's we assigned to
629 the articles.

630

631 **Table S3: ECO terms were assigned by us to the top-50 annotating papers.**

632 The table entries are ranked by the frequency of the assignments, i.e. 27 papers are as-
633 signed with term ECO:0000160, 21 were assigned ECO:0000004, etc. Entries in **boldface**
634 are for computational methods, which were used in many papers in combination with
635 experimental methods to assign function. Table S2 lists the ECO terms.

Table S1. Top 50 Annotating Articles

N	Proteins	Annotations	Species	ref.	MFO	BPO	CCO
1	4937	11050	<i>H. sapiens</i>	[9]	0	0	11050
2	4247	7046	<i>S. pombe</i>	[26]	0	0	7046
3	2412	2412	<i>H. sapiens</i>	[27]	0	0	2412
4	1791	5918	<i>C. elegans</i>	[28]	0	5918	0
5	1406	1863	<i>S. cerevisiae</i>	[29]	0	0	1863
6	1251	1251	<i>A. thaliana</i>	[30]	0	0	1251
7	1205	1476	<i>C. elegans</i>	[31]	0	1476	0
8	1186	1213	<i>M. musculus</i>	[32]	0	0	1213
9	1136	1136	<i>A. thaliana</i>	[33]	0	0	1136
10	1101	2269	<i>C. elegans</i>	[13]	0	2269	0
11	1043	1365	<i>M. tuberculosis</i>	[34]	0	0	1365
12	1041	1041	<i>A. thaliana</i>	[35]	0	0	1041
13	865	1533	<i>C. elegans</i>	[36]	0	1533	0
14	845	845	<i>S. cerevisiae</i>	[37]	0	0	845
15	784	784	<i>A. thaliana</i>	[38]	0	0	784
16	735	735	<i>M. tuberculosis</i>	[39]	0	0	735
17	724	882	<i>A. thaliana</i>	[40]	0	0	882
18	634	634	<i>A. thaliana</i>	[41]	0	0	634
19	613	613	Mycobacter sp.	[42]	0	613	0
20	607	661	<i>C. elegans</i>	[16]	0	659	2
21	577	577	<i>A. thaliana</i>	[43]	0	0	577

Continued on next page

N	Proteins	Annotations	Species	ref.	MFO	BPO	CCO
22	553	884	<i>C. elegans</i>	[11]	0	884	0
23	516	5972	<i>C. elegans</i>	[44]	0	5972	0
24	503	503	<i>S. cerevisiae</i>	[45]	0	0	503
25	498	638	<i>S. cerevisiae</i>	[46]	638	0	0
26	479	848	<i>C. elegans</i>	[47]	0	848	0
27	465	468	<i>H. sapiens</i>	[48]	0	0	468
28	436	436	<i>A. thaliana</i>	[49]	0	0	436
29	430	513	<i>A. thaliana</i>	[50]	0	0	513
30	413	456	<i>D. melanogaster</i>	[17]	0	39	417
31	401	401	<i>A. thaliana</i>	[51]	0	0	401
32	392	392	<i>A. thaliana</i>	[52]	0	0	392
33	392	639	<i>C. elegans</i>	[15]	0	639	0
34	383	917	<i>C. elegans</i>	[12]	0	917	0
35	380	380	<i>A. thaliana</i>	[53]	0	0	380
36	375	375	<i>M. musculus</i>	[54]	0	0	375
37	343	509	<i>H. sapiens</i>	[55]	509	0	0
38	338	338	Ddiscoideum	[56]	0	0	338
39	328	328	<i>A. thaliana</i>	[57]	0	0	328
40	319	329	<i>C. albicans</i>	[58]	1	328	0
41	305	312	<i>A. thaliana</i>	[59]	0	0	312
42	290	331	<i>S. cerevisiae</i>	[60]	0	0	331
43	285	761	<i>C. elegans</i>	[10]	0	761	0

Continued on next page

N	Proteins	Annotations	Species	ref.	MFO	BPO	CCO
44	283	499	<i>C. elegans</i>	[61]	0	499	0
45	266	433	<i>M. musculus</i>	[62]	433	0	0
46	260	260	<i>A. thaliana</i>	[63]	0	260	0
47	258	259	<i>S. pombe</i>	[64]	0	259	0
48	244	397	<i>D. melanogaster</i>	[14]	0	367	30
49	242	397	<i>D. melanogaster</i>	[65]	0	0	397
50	241	263	<i>A. thaliana</i>	[66]	0	0	263

636 **The top 50 annotating articles.** **N:** article rank; **Proteins:** number of proteins
637 annotated in this article; **Annotations:** number of annotating GO terms; **Species:**
638 annotated species; **ref.** annotating article; **MFO/BPO/CCO:** number of proteins
639 annotated in the Molecular Function, Biological Process and Cellular Component
640 ontologies, respectively.

Table S2. ECO Terms Assigned to Top-50 Papers

PMID	Ref	ECO terms/ECO ID's
18029348	[9]	imaging assay evidence/ECO:0000324 immunofluorescence evidence/ECO:0000007 immunolocalization evidence/ECO:0000087
16823372	[26]	imaging assay evidence/ECO:0000324 yellow fluorescent protein fusion protein localization evidence/ECO:0000128 enzyme inhibition experiment evidence/ECO:0000184
18614015	[27]	imaging assay evidence/ECO:0000324 protein separation followed by fragment identification evidence/ECO:0000160 sequence similarity evidence/ECO:0000044 cell fractionation evidence/ECO:0000004 GFP fusion protein localization evidence/ECO:0000126 computational combinatorial evidence/ECO:0000053 motif similarity evidence/ECO:0000028 targeting sequence prediction evidence/ECO:0000081 protein BLAST evidence/ECO:0000208
14551910	[28]	imaging assay evidence/ECO:0000324 RNAi evidence/ECO:0000019 loss-of-function mutant phenotype evidence/ECO:0000016 nucleotide BLAST evidence/ECO:0000207 sequence alignment evidence/ECO:0000200
14562095	[29]	imaging assay evidence/ECO:0000324 GFP fusion protein localization evidence/ECO:0000126 fusion protein localization evidence/ECO:0000124 affinity chromatography evidence/ECO:0000079

Continued on next page

PMID	Ref	ECO terms/ECO ID's
18431481	[30]	protein separation followed by fragment identification evidence/ECO:0000160 targeting sequence prediction evidence/ECO:0000081 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 imported information/ECO:0000311
15791247	[31]	imaging assay evidence/ECO:0000324 RNAi evidence/ECO:0000019 loss-of-function mutant phenotype evidence/ECO:0000016 protein BLAST evidence/ECO:0000208
14651853	[32]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 targeting sequence prediction evidence/ECO:0000081 sequence similarity evidence/ECO:0000044 protein BLAST evidence/ECO:0000208 nucleotide BLAST evidence/ECO:0000207 Affymetrix array experiment evidence/ECO:0000101 imported information/ECO:0000311
17317660	[33]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 transmembrane domain prediction evidence/ECO:0000083 sequence similarity evidence/ECO:0000044

Continued on next page

PMID	Ref	ECO terms/ECO ID's
12529635	[13]	imaging assay evidence/ECO:0000324 RNAi evidence/ECO:0000019 loss-of-function mutant phenotype evidence/ECO:0000016 motif similarity evidence/ECO:0000028 protein BLAST evidence/ECO:0000208 nucleotide BLAST evidence/ECO:0000207 computational combinatorial evidence/ECO:0000053
15525680	[34]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 transmembrane domain prediction evidence/ECO:0000083 sequence similarity evidence/ECO:0000044 computational combinatorial evidence/ECO:0000053 biological system reconstruction/ECO:0000088 imported information/ECO:0000311 protein BLAST evidence/ECO:0000208
21166475	[35]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 computational combinatorial evidence/ECO:0000053 imported information/ECO:0000311 transmembrane domain prediction evidence/ECO:0000083 sequence alignment evidence/ECO:0000200 motif similarity evidence/ECO:0000028
15489339	[36]	imaging assay evidence/ECO:0000324 RNAi evidence/ECO:0000019 loss-of-function mutant phenotype evidence/ECO:0000016 nucleotide BLAST evidence/ECO:0000207

Continued on next page

PMID	Ref	ECO terms/ECO ID's
16823961	[37]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 imported information/ECO:0000311
21533090	[38]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 imported information/ECO:0000311 computational combinatorial evidence/ECO:0000053 transmembrane domain prediction evidence/ECO:0000083 sequence alignment evidence/ECO:0000200 motif similarity evidence/ECO:0000028 targeting sequence prediction evidence/ECO:0000081
14532352	[39]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 transmembrane domain prediction evidence/ECO:0000083
20061580	[40]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 transmembrane domain prediction evidence/ECO:0000083 imported information/ECO:0000311 targeting sequence prediction evidence/ECO:0000081 protein expression level evidence/ECO:0000046

Continued on next page

PMID	Ref	ECO terms/ECO ID's
15028209	[41]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 targeting sequence prediction evidence/ECO:0000081 Affymetrix array experiment evidence/ECO:0000101 protein expression level evidence/ECO:0000046 protein BLAST evidence/ECO:0000208 computational combinatorial evidence/ECO:0000053 motif similarity evidence/ECO:0000028 transmembrane domain prediction evidence/ECO:0000083
12657046	[42]	mutant phenotype evidence/ECO:0000015 nucleic acid hybridization evidence/ECO:0000026 imported information/ECO:0000311 sequence similarity evidence/ECO:0000044 combinatorial evidence/ECO:0000212
17704769	[16]	imaging assay evidence/ECO:0000324 RNAi evidence/ECO:0000019 loss-of-function mutant phenotype evidence/ECO:0000016
17432890	[43]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 transmembrane domain prediction evidence/ECO:0000083 imported information/ECO:0000311 targeting sequence prediction evidence/ECO:0000081 protein BLAST evidence/ECO:0000208 computational combinatorial evidence/ECO:0000053

Continued on next page

PMID	Ref	ECO terms/ECO ID's
11231151	[11]	imaging assay evidence/ECO:0000324 RNAi evidence/ECO:0000019 loss-of-function mutant phenotype evidence/ECO:0000016
17417969	[44]	imaging assay evidence/ECO:0000324 RNAi evidence/ECO:0000019 loss-of-function mutant phenotype evidence/ECO:0000016
14576278	[45]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 transmembrane domain prediction evidence/ECO:0000083
16429126	[46]	protein separation followed by fragment identification evidence/ECO:0000160 sequence similarity evidence/ECO:0000044 affinity chromatography evidence/ECO:0000079 protein BLAST evidence/ECO:0000208 imported information/ECO:0000311
21529718	[47]	imaging assay evidence/ECO:0000324 RNAi evidence/ECO:0000019 loss-of-function mutant phenotype evidence/ECO:0000016 computational combinatorial evidence/ECO:0000053
11256614	[48]	GFP fusion protein localization evidence/ECO:0000126 yellow fluorescent protein fusion protein localization evidence/ECO:0000128 imaging assay evidence/ECO:0000324 motif similarity evidence/ECO:0000028 protein BLAST evidence/ECO:0000208 nucleotide BLAST evidence/ECO:0000207

Continued on next page

PMID	Ref	ECO terms/ECO ID's
17644812	[49]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 transmembrane domain prediction evidence/ECO:0000083 targeting sequence prediction evidence/ECO:0000081 computational combinatorial evidence/ECO:0000053
16618929	[50]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 transmembrane domain prediction evidence/ECO:0000083
18433294	[17]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 imaging assay evidence/ECO:0000324 RNAi evidence/ECO:0000019 loss-of-function mutant phenotype evidence/ECO:0000016 immunofluorescence evidence/ECO:0000007
17151019	[51]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 imported information/ECO:0000311 transmembrane domain prediction evidence/ECO:0000083

Continued on next page

PMID	Ref	ECO terms/ECO ID's
14671022	[52]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 protein BLAST evidence/ECO:0000208 targeting sequence prediction evidence/ECO:0000081
12529643	[15]	imaging assay evidence/ECO:0000324 RNAi evidence/ECO:0000019 loss-of-function mutant phenotype evidence/ECO:0000016
12445391	[12]	imaging assay evidence/ECO:0000324 RNAi evidence/ECO:0000019 loss-of-function mutant phenotype evidence/ECO:0000016 BLAST evidence/ECO:0000206
15539469	[53]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 targeting sequence prediction evidence/ECO:0000081 transmembrane domain prediction evidence/ECO:0000083 motif similarity evidence/ECO:0000028 protein BLAST evidence/ECO:0000208 computational combinatorial evidence/ECO:0000053
12865426	[54]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 transmembrane domain prediction evidence/ECO:0000083

Continued on next page

PMID	Ref	ECO terms/ECO ID's
16189514	[55]	yeast 2-hybrid evidence/ECO:0000068 imaging assay evidence/ECO:0000324 motif similarity evidence/ECO:0000028 co-purification evidence/ECO:0000022 combinatorial evidence/ECO:0000212
20422638	[56]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 combinatorial evidence/ECO:0000212
12938931	[57]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 sequence similarity evidence/ECO:0000044 nucleotide BLAST evidence/ECO:0000207 imported information/ECO:0000311 transmembrane domain prediction evidence/ECO:0000083
16336044	[58]	imaging assay evidence/ECO:0000324 RNAi evidence/ECO:0000019 loss-of-function mutant phenotype evidence/ECO:0000016
18633119	[59]	protein separation followed by fragment identification evidence/ECO:0000160 cell fractionation evidence/ECO:0000004 Western blot evidence/ECO:0000112
11914276	[60]	imaging assay evidence/ECO:0000324 immunofluorescence evidence/ECO:0000007 epitope-tagged protein immunolocalization evidence/ECO:0000092 transmembrane domain prediction evidence/ECO:0000083 imported information/ECO:0000311

Continued on next page

PMID	Ref	ECO terms/ECO ID's
11099033	[10]	imaging assay evidence/ECO:0000324 RNAi evidence/ECO:0000019 loss-of-function mutant phenotype evidence/ECO:0000016 protein BLAST evidence/ECO:0000208 computational combinatorial evidence/ECO:0000053
11099034	[61]	imaging assay evidence/ECO:0000324 RNAi evidence/ECO:0000019 loss-of-function mutant phenotype evidence/ECO:0000016 nucleotide BLAST evidence/ECO:0000207 protein BLAST evidence/ECO:0000208
11591653	[62]	hybrid interaction evidence/ECO:0000025 imaging assay evidence/ECO:0000324
16502469	[63]	protein separation followed by fragment identification evidence/ECO:0000160 sequence similarity evidence/ECO:0000044 protein BLAST evidence/ECO:0000208 Northern assay evidence/ECO:0000106 reverse transcription polymerase chain reaction transcription evidence/ECO:0000108
12529438	[64]	microarray RNA expression level evidence/ECO:0000104 sequence orthology evidence used in manual assertion/ECO:0000266 motif similarity evidence/ECO:0000028
17412918	[14]	RNAi evidence/ECO:0000019 loss-of-function mutant phenotype evidence/ECO:0000016 imaging assay evidence/ECO:0000324

Continued on next page

PMID	Ref	ECO terms/ECO ID's
18981222	[65]	protein separation followed by fragment identification evidence/ECO:0000160 sequence similarity evidence/ECO:0000044 protein BLAST evidence/ECO:0000208 in vitro assay evidence/ECO:0000181 affinity chromatography evidence/ECO:0000079 imaging assay evidence/ECO:0000324 mutant phenotype evidence/ECO:0000015
16287169	[66]	protein separation followed by fragment identification evidence/ECO:0000160 sequence similarity evidence/ECO:0000044 transmembrane domain prediction evidence/ECO:0000083 sequence alignment evidence/ECO:0000200 computational combinatorial evidence/ECO:0000053 motif similarity evidence/ECO:0000028 targeting sequence prediction evidence/ECO:0000081

641 The Top-50 studies and the ECO terms we have assigned to them. **PMID**: Articles'

642 PubMed ID; **ECO terms/ECO ID's**: terms and ID's we assigned to the articles.

Table S3. Count of ECO terms in top-50 papers

N	ECO term	ECO ID	Articles
1	protein separation followed by fragment identification evidence	ECO:0000160	27
2	sequence similarity evidence	ECO:0000044	27
3	imaging assay evidence	ECO:0000324	24
4	cell fractionation evidence	ECO:0000004	23
5	transmembrane domain prediction evidence	ECO:0000083	17
6	loss-of-function mutant phenotype evidence	ECO:0000016	15
7	protein BLAST evidence	ECO:0000208	15
8	RNAi evidence	ECO:0000019	15
9	imported information	ECO:0000311	13
10	computational combinatorial evidence	ECO:0000053	11
11	targeting sequence prediction evidence	ECO:0000081	11
12	motif similarity evidence	ECO:0000028	10
13	nucleotide BLAST evidence	ECO:0000207	7
14	sequence alignment evidence	ECO:0000200	4
15	GFP fusion protein localization evidence	ECO:0000126	3
16	immunofluorescence evidence	ECO:0000007	3
17	affinity chromatography evidence	ECO:0000079	3
18	computational combinatorial evidence	ECO:0000053	2
19	Affymetrix array experiment evidence	ECO:0000101	2

Continued on next page

N	ECO term	ECO ID	Articles
20	protein expression level evidence	ECO:0000046	2
21	mutant phenotype evidence	ECO:0000015	2
22	combinatorial evidence	ECO:0000212	2
23	co-purification evidence	ECO:0000022	1
24	epitope-tagged protein immunolocalization evidence	ECO:0000092	1
25	sequence orthology evidence used in manual assertion	ECO:0000266	1
26	YFP fusion protein localization evidence	ECO:0000128	2
27	in vitro assay evidence	ECO:0000181	1
28	biological system reconstruction	ECO:0000088	1
29	reverse transcription polymerase chain reaction transcription evidence	ECO:0000108	1
30	Northern assay evidence	ECO:0000106	1
31	Western blot evidence	ECO:0000112	1
32	microarray RNA expression level evidence	ECO:0000104	1
33	fusion protein localization evidence	ECO:0000124	1
34	BLAST evidence	ECO:0000206	1
35	nucleic acid hybridization evidence	ECO:0000026	1
36	enzyme inhibition experiment evidence	ECO:0000184	1
37	immunolocalization evidence	ECO:0000087	1
38	hybrid interaction evidence	ECO:0000025	1
39	yeast 2-hybrid evidence	ECO:0000068	1

643 ECO terms were assigned by us to the top-50 annotating papers. The table entries are
644 ranked by the frequency of the assignments, i.e. 27 papers are assigned with term
645 ECO:0000160, 21 were assigned ECO:0000004, etc. Entries in **boldface** are for
646 computational methods, which were used in many papers in combination with
647 experimental methods to assign function. TableS2 lists the ECO terms.