Title: Mandated data archiving greatly improves access to research data Authors: Timothy H. Vines^{1,2,*}, Rose L. Andrew¹, Dan G. Bock¹, Michelle T. Franklin^{1,3}, Kimberly J. Gilbert¹, Nolan C. Kane^{1,4}, Jean-Sébastien Moore¹, Brook T. Moyers¹, Sébastien Renaut¹, Diana J. Rennison¹, Thor Veen¹, Sam Yeaman¹ Affiliations: ¹Biodiversity Department, University of British Columbia, 6270 University Blvd Vancouver BC, Canada, V6T 1Z4. ²Molecular Ecology Editorial Office, 6270 University Blvd Vancouver BC, Canada, V6T 1Z4. ³Department of Biological Sciences, Simon Fraser University, 8888 University Drive, Burnaby, British Columbia, Canada, V5A 1S6. 4Ecology and Evolutionary Biology Department, University of Colorado at Boulder, Boulder, CO, USA, 80309 *Author for correspondence: vines@zoology.ubc.ca. Tel: 1 778 989 8755 Fax: 1 604 822 8982 Short title: Archiving policies and access to research data

- 18 Non standard abbreviations:
- 19 PLoS Public Library of Science
- 20 BMC BioMed Central
- 21 BJLS Biological Journal of the Linnean Society
- 22 TAG Theoretical and Applied Genetics
- 23 IF Impact Factor

Abstract: The data underlying scientific papers should be accessible to researchers both now and in the future, but how best can we ensure that these data are available? Here we examine the effectiveness of four approaches to data archiving: no stated archiving policy, recommending (but not requiring) archiving, and two versions of mandating data deposition at acceptance. We control for differences between data types by trying to obtain data from papers that use a single, widespread population genetic analysis, STRUCTURE. At one extreme, we found that mandated data archiving policies that require the inclusion of a data availability statement in the manuscript improve the odds of finding the data online almost a thousand-fold compared to having no policy. However, archiving rates at journals with less stringent policies were only very slightly higher than those with no policy at all. We also assessed the effectiveness of asking for data directly from authors and obtained over half of the requested datasets, albeit with about 8 days' delay and some disagreement with authors. Given the long-term benefits of data accessibility to the academic community, we believe that journal-based mandatory data archiving policies and mandatory data availability statements should be more widely adopted. Keywords: Science policy, reproducibility, Joint Data Archiving Policy, population genetics, STRUCTURE

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

45 Archiving the data underlying scientific papers is an essential component of scientific 46 publication and its subsequent reproducibility [1-3], but very few papers actually make the 47 underlying data available [4]. In response to this gap between the needs of science and author 48 behavior, a number of journals have introduced data archiving policies. Here, we evaluate the 49 effectiveness of these policies by comparing journals that have no stated data archiving policy, 50 journals that recommend data archiving, and journals that mandate archiving prior to publication. 51 Journals that mandate data archiving fall into two further subgroups: those that require an 52 explicit data availability statement and those that do not. We ask two questions: (1) does having 53 any kind of data archiving policy improve the likelihood of the data being available online, and 54 (2) does the type of data archiving policy have any effect the likelihood of obtaining the data? 55 56 We recently assembled datasets from a range of journals for a study of the reproducibility of 57 commonly used population genetic analyses [5]. Here, we use this opportunity to examine 58 whether data archiving policy (or lack thereof) was associated with the proportion of datasets we 59 were able to obtain from a journal. As papers within even a single journal contain many different 60 types of data, we restricted both this and our reproducibility study to articles using the population 61 genetics program STRUCTURE [6]. We chose STRUCTURE because it is widely used in ecology and 62 evolution, and because the underlying data is a table of microsatellite, Amplified Fragment 63 Length Polymorphism or Single Nucleotide Polymorphism genotypes, and for the ease of 64 archiving this type of dataset online. For example, the data could be uploaded as supplemental 65 material, or archived on the Dryad repository [7]. Dryad was established in 2010 for the 66 preservation of a wide range of data types associated with ecology or evolution articles, and is 67 often used to archive STRUCTURE datasets. 68 69 Data collection 70 71 We used Web of Science to identify papers published in 2011 or 2012 that cited the original 72 description of STRUCTURE [6]. We selected journals for each of the four journal categories 73 described above, and excluded those that had less than five eligible papers. We complemented 74 our list of papers by searching for additional articles that used STRUCTURE on the journal website.

75 Papers that used DNA sequence data were excluded, as preparing raw sequence data from e.g. 76 GenBank for re-analysis with STRUCTURE was found to be very time consuming. 77 78 We found four eligible journals with no stated data archiving policy: Conservation Genetics, 79 *Crop Science, Genetica, and Theoretical and Applied Genetics (TAG).* 80 81 There were four eligible journals that had some sort of data archiving policy but stopped short of 82 mandating archiving for all data (BMC Evolutionary Biology, Biological Journal of the Linnean 83 Society (BJLS), Journal of Heredity and PLoS One). These policies were retrieved from the 84 author guidelines in mid 2011 and are available on Dryad (doi:10.5061/dryad.6bs31). The latter 85 three journals ask that the data be placed onto an online archive whenever one exists. For 86 STRUCTURE data, Dryad is the most commonly used repository, and indeed the policies of the last 87 two journals (J. Heredity and PLoS One) explicitly mention Dryad. There is thus an expectation 88 for three of these four journals the data should be available somewhere online, most likely on 89 Dryad. For BMC Evolutionary Biology the data will only be online if the authors have decided to 90 share it at publication. The individual policies are as follows: 91 92 First, BMC Evolutionary Biology states that "submission ... implies that readily reproducible 93 materials described in the manuscript, including all relevant raw data, will be freely available to 94 any scientist wishing to use them for non-commercial purposes", and at that time did not require 95 that data appear in an online archive. This policy has been in place since 2009. 96 97 Second, the Biological Journal of the Linnean Society has the policy "we recommend that data 98 for which public repositories are widely used, and are accessible to all, should be deposited in 99 such a repository prior to publication." This policy was introduced in January 2011 and we hence 100 only considered papers submitted after this date. 101 102 Third, J. Heredity "endorses the principles of the Joint Data Archiving Policy [see below] in 103 encouraging all authors to archive primary datasets in an appropriate public archive, such as 104 Dryad, TreeBASE, or the Knowledge Network for Biocomplexity". As with *BJLS*, this policy 105 was introduced in January 2011 and we hence only considered papers submitted after this date.

106 107 Fourth, PLoS One has had a policy on data sharing in place since 2008, and one statement is as 108 follows: "If an appropriate repository does not exist, data should be provided in an open access 109 institutional repository, a general data repository such as Dryad, or as Supporting Information 110 files with the published paper." 111 112 Finally, there were four journals that adopted a mandatory data archiving policy (known as the 113 Joint Data Archiving Policy or JDAP [1]), which states "[Journal X] requires, as a condition for 114 publication, that data supporting the results in the paper should be archived in an appropriate 115 public archive". For these journals we excluded papers submitted before the policy came into 116 force: January 2011 for Molecular Ecology, Journal of Evolutionary Biology, and Evolution, and 117 March 2011 for *Heredity*. Of these four, two (*Molecular Ecology* and *Heredity*) additionally 118 require that authors include a data availability statement within each accepted manuscript; these 119 sections describe the location (typically the database and accession numbers) of all publicly 120 available data. 121 122 For all 229 eligible papers we then checked whether the STRUCTURE genotype data was available 123 either as supplemental material or elsewhere online, such as on the Dryad archive [7]. Our results 124 are shown in Table 1 and Figure 1, and the data and R code used in the analysis are archived at 125 doi:10.5061/dryad.6bs31. 126 127 Statistical analysis 128 129 To evaluate the statistical support for an association between the presence/absence of an 130 archiving policy and whether or not the STRUCTURE data could be found online, we fitted a 131 mixed effects logistic regression. The response variable was whether or not the data from a paper 132 was available online, coded as '0' for not available and '1' for available. The predictor variable 133 was either 'no policy' or 'archiving policy', and journals were included as a random effect within 134 each category. 135

136 Having any sort of archiving policy did lead to a significant improvement in the probability of the data being online (likelihood ratio test statistic = 4.27, p= 0.038), such that the odds of 137 138 getting the data were about 25 times higher (95% confidence interval: 1.5 to 416.7). 139 140 We then tested how well each type of archiving policy compared to having no policy at all. As 141 above, we used a mixed effects logistic regression. Again, the response variable was whether or 142 not the data from a paper was available online, coded as '0' for not available and '1' for 143 available. The predictor variable was policy type, and the categories were 'no policy', 144 'recommend archiving', 'mandate archiving, no data statement' and 'mandate archiving, with 145 data statement'. Journals were a random effect within each policy type. The overall model found 146 that policy type did have a very significant effect on data availability (likelihood ratio test 147 statistic = 28.06, p<0.001). 148 149 Since this is a logistic model, we can readily calculate the effect that the different policy types 150 have on the likelihood that the data will be available. We explore these odds for each type of 151 policy below, using 'no policy' as the baseline. 152 153 Having a 'recommend archiving' policy made it 3.6 times more likely that the data were online 154 compared to having no policy. However, the 95% confidence interval overlapped with 1 (0.96 to 155 13.6), and hence this increase in the odds is not significant. Overall, recommending data 156 archiving is only marginally more effective than having no policy at all. 157 158 The data was 17 times more likely to be available online for journals that had adopted a 159 mandatory data archiving policy but did not require a data accessibility statement in the 160 manuscript. This odds ratio was significantly greater than 1 (95% confidence interval: 3.7 to 161 79.6). 162 163 For 'mandate archiving' journals where a data accessibility statement is required in the 164 manuscript, the odds of finding the data online were 974 times higher compared to having no policy. The 95% confidence interval on these odds is very wide (97.9 to 9698.8), but nonetheless 165

166 shows that the combination of a mandatory policy and an accessibility statement is much more 167 effective than any other policy type. 168 169 170 Requesting data directly from authors 171 172 A number of the 'recommend archiving' policies state that the data should also be freely 173 available from the authors by request (see the 'Journal Policies' file at doi:10.5061/dryad.6bs31), 174 and hence we wanted to evaluate whether obtaining data directly from authors is an effective 175 approach. Part of the dataset collection for our reproducibility study [5] involved emailing 176 authors of papers from two of the 'recommend archiving' journals (BMC Evolutionary Biology 177 and *PLoS One*) and requesting their STRUCTURE input files. Here, we examine how often these 178 requests led to us obtaining the data. We did not email the authors of articles where the data were 179 already available online. A detailed description of our data request process appears on Dryad 180 (doi:10.5061/dryad.6bs31), but we essentially contacted corresponding and senior authors of 181 each paper up to three times over a three week period, and recorded if and when the data were 182 received. 183 184 We obtained data directly from the authors for seven of the 12 eligible papers in BMC 185 Evolutionary Biology, and 27 datasets from 45 papers from PLoS One (Table 1). All seven of the 186 BMC Evolutionary Biology datasets arrived between eight and 14 days after our initial request. 187 Ten of the *PLoS One* datasets came within a week, 13 came between eight and 14 days, and four 188 arrived between 15 and 21 days. Unlike the online data, which could generally be obtained 189 within a few minutes, the requested datasets took a mean of 7.7 days to arrive, with one author 190 responding that the dataset had been lost in the year since publication. More than one email had 191 to be sent to the corresponding and/or senior author for 53% of papers, and the authors of 29% of 192 the papers did not respond to any of our requests. No data were received more than 21 days after 193 our initial request. We also note that requesting data via email did upset some authors, 194 particularly when they were reminded of the journal's data archiving policy or when multiple 195 emails were sent.

197 Our average return of 59% in an average of 7.7 days is markedly better than has been reported in 198 similar studies: Wicherts et al. [8] received only 26% of requested datasets after six months of 199 effort with authors of 141 psychology articles, and Savage and Vickers [9] received only one of 200 ten datasets requested from papers in *PLoS Medicine* and *PLoS Clinical Trials*. In a 1999 study, 201 Leberg and Neigel [10] emailed the authors of 30 papers that contained an incomplete 202 description of their sequence dataset, but received the requested data from just one of them. 203 Since the latter study and ours both involved the evolutionary biology community, it appears that 204 attitudes to data sharing have improved dramatically over the last decade. However, the two 205 more recent studies that used human data still had low success rates, perhaps because privacy 206 and consent issues are a significant impediment to data sharing in these fields. 207 208 209 Conclusions 210 211 Our results demonstrate that journal-based data archiving policies can be very effective in 212 ensuring that research data are available to the scientific community, especially when journals 213 require that a data accessibility statement appear in the manuscript. The 'recommend archiving' 214 group of journals encompassed the broadest spread of policy types, yet as a whole only had 10 of 89 datasets available. The policies range from a simple "Submission ... implies that ... all 215 216 relevant raw data, will be freely available to any scientist wishing to use them for non-217 commercial purposes" at BMC Evolutionary Biology to an endorsement of the full Joint Data 218 Archiving Policy at J. Heredity. However, none of these policies led to more than 23% of the 219 data being available online (at BJLS), and there was no significant difference between the 220 success of this policy type and having no policy at all. 221 222 Interestingly, *PLoS One*'s very comprehensive policy, which is over 1000 words long and 223 contains statements like "data should be provided in an open access institutional repository, a 224 general data repository such as Dryad, or as Supporting Information files with the published 225 paper" was only marginally more effective than BMC Evolutionary Biology's simple request that 226 the data be freely available, with 11% and 7% of the data online, respectively. 227

228 The difference between *PLoS One* and the 'mandate archiving' journals may arise because the 229 wide breadth of subject areas in *PLoS One* precludes having a policy with the bald simplicity of 230 the Joint Data Archiving Policy: "[Journal X] requires, as a condition for publication, that data 231 supporting the results in the paper should be archived in an appropriate public archive". Even though the portion of *PLoS One*'s author community that uses STRUCTURE broadly overlaps with 232 233 the authors of the papers in the JDAP journals, it may be that the lack of a single strong 234 statement leads to much lower compliance. One simple remedy for this situation might be the 235 introduction of a mandatory data accessibility statement in all manuscripts. For fields where 236 archiving is not (yet) standard practice, this could state that the data were available from the 237 authors, but in fields where archiving is expected the authors would indicate where their data 238 were available online. 239 More broadly, a study by Piwowar and Chapman [11] on 397 microarray datasets from 20 240 241 journals also found that having a 'strong' (i.e. close to mandatory) data archiving policy led to a 242 high proportion (>50%) of the datasets being available online. Journals that had a 'weak' policy 243 (i.e. recommended archiving) had just over 30% of microarray datasets available, and journals 244 with no policy had only about 20% availability. Furthermore, they also found that a journal with 245 an Impact Factor (IF) of 15 was 4.5 times more likely to have the microarray data online than a 246 journal with an IF of 5. We find a similar effect in our data: using the 2010 Impact Factors, we 247 were 3.2 times more likely to find the data online for a journal with an IF of 5.0 (the average IF 248 of the JDAP journals) compared to those with an IF of 2.2 (the average IF of the 'no policy' 249 journals); details of this analysis are available at doi:10.5061/dryad.6bs31. We are able to 250 exclude higher Impact Factor as the primary cause of the high rate of data archiving in the JDAP 251 journals: in 2010 (before the mandatory archiving policy was introduced), none of the 27 eligible 252 papers in the Journal of Evolutionary Biology, Evolution or Heredity had archived their data, 253 even though their Impact Factors were essentially the same in 2010 and 2011 (Molecular 254 Ecology recommended archiving in 2010 and was excluded from this comparison). This result 255 suggests that the introduction of the JDAP policy in 2011 was primarily responsible for the 256 abrupt rise in the proportion of articles in these three journals that archived their data. However,

it is possible that Impact Factor still plays a role, as only journals with a high IF may feel able to

258 introduce stringent archiving policies. The positive effects of a strongly worded data archiving 259 statement were also confirmed by a much larger study involving 11603 microarray datasets [12]. 260 261 Requesting data directly from authors can also provide access to research data, but this approach 262 can be hampered by delays and the potential for disagreement between requester and the authors. 263 Furthermore, the availability of datasets directly from authors will only decrease as time since 264 publication increases. This is particularly true when researchers leave science or when data that 265 are stored on lab computers or websites get misplaced [13, 14]. 266 267 Even though our results strongly emphasize the value of public databases for archiving scientific 268 data, these databases do require ongoing financial support; this money may come from funding 269 agencies, journal publishers, libraries or even individual researchers. A recent study put the cost 270 of running the Dryad database at around \$400,000 per annum; these costs include the 271 maintenance of their archive and the addition and curation of an extra 10,000 datasets per year. 272 For comparison, the same amount spent by a funding agency on basic research would generate 273 about 16 new publications [15]. Given that the long-term availability of these data allows for 274 meta-analyses, the checking of previous results, and not collecting the same data again, money 275 spent on data archiving is extremely cost effective. In light of all these advantages, we believe 276 that journal-based mandatory data archiving policies and data accessibility statements should be 277 more widely adopted. 278


- 279 References:
- 280 1. Whitlock, M.C., McPeek, M.A., Rausher, M.D., Rieseberg, L., and Moore, A.J. (2010) Data
- 281 archiving. Am. Nat. 175, 145-146
- 282 2. Wolkovich, E.M., Regetz, J., and O'Connor, M.I. (2012) Advances in global change
- research require open science by individual researchers. Global Change Biology 18, 2102-
- 284 2110
- 285 3. Peng, R.D. (2011) Reproducible Research in computational science. *Science* **334**, 1226
- 286 4. Alsheikh-Ali, A.A., Qureshi, W., Al-Mallah, M.A., and Ioannidis, J.P.A. (2011) Public
- availability of published research data in high-impact journals. *PLoS One* 6(9), e24357,
- 288 doi:10.1371/journal.pone.0024357
- 5. Gilbert, K.J., Andrew, R.L., Bock, D.G., Franklin, M.T., Kane, N.C., Moore, J.-S., Moyers,
- B.T., Renaut, S., Rennison, D.J., Veen, T. and Vines, T.H. (2012) Recommendations for
- 291 utilizing and reporting population genetic analyses: the reproducibility of genetic clustering
- using the program STRUCTURE. *Molecular Ecology* **21**, 4925-4930
- 293 6. Pritchard, J.K., Stephens, M., and Donnelly, P. (2000) Inference of population structure
- using multilocus genotype data. *Genetics* **155**, 945-959
- 295 7. Vision, T. (2010) Open data and the social contract of scientific publishing. *Bioscience* **60**,
- 296 330-331
- 8. Wicherts, J.M., Borsboom, D., Kats, J., and Molenaar, D. (2006) The poor availability of
- psychological research data for reanalysis. *Am. Psychol.* **61**, 726–728
- 299 9. Savage, C.J. and Vickers A.J. (2009) Empirical study of data sharing by authors publishing
- in PLoS journals. *PLoS One* **4**, e7078
- 301 10. Leberg, P.L. and Neigel, J.E. (1999) Enhancing the retrievability of population genetic
- 302 survey data? An assessment of animal mitochondrial DNA studies. *Evolution* **53**,1961-1965
- 303 11. Piwowar, H.A. and Chapman W.W. (2010) Public sharing of research datasets: A pilot
- 304 study of associations. J. Informetrics 4, 148-156
- 305 12. Piwowar, H.A. (2011) Who shares? Who doesn't? Factors associated with openly archiving
- raw research. *PLoS One* **6**, e18657
- 307 13. Wren, J.D., Grissom, J.E., and Conway, T. (2006) E-mail decay rates among corresponding
- authors in MEDLINE. The ability to communicate with and request materials from authors
- is being eroded by the expiration of e-mail addresses. *EMBO Rep.* 7, 122-127

- 310 14. Michener, W.K., Brunt J.W., Helly J.J., Kirchner, T.B., and Stafford, S.G. (1997)
- Nongeospatial metadata for the ecological sciences. *Ecol. Appl.* 7, 330–342
- 312 15. Piwowar, H.A., Vision, T.J., and Whitlock, M.C. (2011) Data archiving is a good
- 313 investment. *Nature* **473**, 285-285

- 315 Acknowledgments: We thank Heather Piwowar, Loren Rieseberg, Phil Davis and Mike Whitlock
- 316 for comments on an earlier version of the manuscript, and Arianne Albert for help with the
- 317 statistics. We would also like to express our gratitude to the many authors who shared their data
- 318 with us.

Table 1. The number of eligible articles per journal, and the number for which data were obtained from online databases.

Policy	Journal	No. eligible papers	No. data online
No Policy	Conservation Genetics	47	1
	Crop Science	12	1
	Genetica	8	1
	T.A.G.	21	0
Recommend data archiving	BMC Evolutionary Biol.	13	1
	B.J.L.S.	13	3
	J. Heredity	12	0
	PLoS One	51	6
Mandatory data archiving	J. Evolutionary Biology	10	3
	Evolution	6	3
	Heredity	7	7
	Molecular Ecology	28	27



325