

Spectral instability of some non-selfadjoint  
anharmonic oscillators  
Instabilité spectrale de certains oscillateurs  
anharmoniques non-autoadjoints

Raphaël HENRY

1<sup>er</sup> octobre 2018

**Résumé**

Notre objectif est de mettre en évidence l'instabilité spectrale de certains opérateurs différentiels non-autoadjoints, via l'étude de la croissance des normes des projecteurs spectraux  $\Pi_n$  associés à leurs valeurs propres. Nous nous intéressons à certains *oscillateurs anharmoniques*

$$\mathcal{A}(m, \theta) = -\frac{d^2}{dx^2} + e^{i\theta}|x|^m$$

avec

$$|\theta| < \min \left\{ \frac{(m+2)\pi}{4}, \frac{(m+2)\pi}{2m} \right\},$$

définis sur  $L^2(\mathbb{R})$ . Nous étendons les résultats de [4] et [5] en donnant un développement asymptotique de la norme des projecteurs spectraux associés aux grandes valeurs propres pour les opérateurs  $\mathcal{A}(1, \theta)$  et  $\mathcal{A}(2k, \theta)$ ,  $k \geq 1$ .

The purpose of this Note is to highlight the spectral instability of some non-selfadjoint differential operators, by studying the growth rate of the norms of the spectral projections  $\Pi_n$  associated with their eigenvalues. More precisely, we are concerned with some *anharmonic oscillators*

$$\mathcal{A}(m, \theta) = -\frac{d^2}{dx^2} + e^{i\theta}|x|^m$$

with

$$|\theta| < \min \left\{ \frac{(m+2)\pi}{4}, \frac{(m+2)\pi}{2m} \right\},$$

defined on  $L^2(\mathbb{R})$ . We get asymptotic expansions for the norm of the spectral projections associated with the large eigenvalues of  $\mathcal{A}(1, \theta)$  and  $\mathcal{A}(2k, \theta)$ ,  $k \geq 1$ , extending the results of [4] and [5].

## 1 Spectral instability and pseudospectra

It is well-known that the spectral theorem implies some control of stability for the spectrum of selfadjoint operators : if  $\mathcal{A}$  is a selfadjoint operator acting on

the Hilbert space  $\mathcal{H}$ , the spectrum of its perturbations  $\mathcal{A} + \varepsilon\mathcal{B}$ , with  $\varepsilon > 0$  and any  $\mathcal{B} \in \mathcal{L}(\mathcal{H})$ ,  $\|\mathcal{B}\| \leq 1$ , lies entirely inside an  $\varepsilon$ -neighborhood of the spectrum  $\sigma(\mathcal{A})$ . In other words, the norm of the resolvent of  $\mathcal{A}$  near the spectrum blows up like the inverse distance to the spectrum. It has also been known for several years (see for instance [13]) that such a behavior could not be expected in general in the case of non-selfadjoint operators. One can understand it thanks to the notion of  $\varepsilon$ -pseudospectra of an operator  $\mathcal{A}$ , defined as the family of sets  $\sigma_\varepsilon(\mathcal{A})$ , indexed by  $\varepsilon > 0$ ,

$$\sigma_\varepsilon(\mathcal{A}) = \left\{ \xi \in \rho(\mathcal{A}) : \|(\mathcal{A} - \xi)^{-1}\| > \frac{1}{\varepsilon} \right\} \cup \sigma(\mathcal{A}).$$

The link between spectral instability and pseudospectra appears more clearly in the following equivalent formulation, which is a weak version of the Roch and Silbermann theorem [11] :

$$\sigma_\varepsilon(\mathcal{A}) = \bigcup_{\substack{\omega \in \mathcal{L}(\mathcal{H}), \\ \|\mathcal{B}\| \leq 1}} \sigma(\mathcal{A} + \varepsilon\mathcal{B})$$

(See also [12] and references therein).

In the following, we deal with the instability indices associated with an isolated eigenvalue  $\lambda \in \sigma(\mathcal{A})$ . The instability index associated with  $\lambda$  is defined as

$$\kappa(\lambda) = \|\Pi(\lambda)\|,$$

where  $\Pi(\lambda)$  denotes the spectral projection associated with  $\lambda$ . Of course  $\kappa(\lambda) \geq 1$  in any case, and  $\kappa(\lambda) = 1$  when  $\mathcal{A}$  is selfadjoint. These numbers  $\kappa(\lambda)$  are closely related to the size of  $\varepsilon$ -pseudospectra around  $\lambda$ . Indeed, if  $\sigma_\varepsilon^\lambda$  denotes the connected component of  $\sigma_\varepsilon(\mathcal{A})$  containing  $\lambda$ , and if we assume for simplicity that  $\sigma_\varepsilon^\lambda \cap \sigma(\mathcal{A}) = \{\lambda\}$  and  $\sigma_\varepsilon^\lambda$  is bounded, then the perimeter  $|\partial\sigma_\varepsilon^\lambda|$  of  $\sigma_\varepsilon^\lambda$  satisfies (see [3])

$$|\partial\sigma_\varepsilon^\lambda| \geq 2\pi\varepsilon\kappa(\lambda). \quad (1.1)$$

In the finite dimensional setting at least, instability indices give a better description of pseudospectra : if  $\mathcal{A} \in \mathcal{M}_n(\mathbb{C})$  is a diagonalisable matrix with distinct eigenvalues  $\lambda_1, \dots, \lambda_n$ , Embree and Trefethen show [13] that there exists  $\varepsilon_0 > 0$  such that, for all  $\varepsilon < \varepsilon_0$ ,

$$\bigcup_{\lambda_k \in \sigma(\mathcal{A})} D(\lambda_k, \varepsilon\kappa(\lambda_k) + \mathcal{O}(\varepsilon^2)) \subset \sigma_\varepsilon(\mathcal{A}) \subset \bigcup_{\lambda_k \in \sigma(\mathcal{A})} D(\lambda_k, \varepsilon\kappa(\lambda_k) + \mathcal{O}(\varepsilon^2)). \quad (1.2)$$

In the case of an infinite dimensional space, the validity of this statement should be investigated, as well as the dependance on  $\lambda_k$  of the  $\mathcal{O}(\varepsilon^2)$  terms.

In the following, we study the instability indices of simple non-selfadjoint differential operators introduced by Davies in [4], for which the instability phenomenon described above will appear clearly. Let us define the *anharmonic oscillators*

$$\mathcal{A}(m, \theta) = -\frac{d^2}{dx^2} + e^{i\theta}|x|^m \quad (1.3)$$

with

$$|\theta| < \min \left\{ \frac{(m+2)\pi}{4}, \frac{(m+2)\pi}{2m} \right\}, \quad (1.4)$$

defined on  $L^2(\mathbb{R})$  in [4] by taking the closure of the associated quadratic form defined on  $C_0^\infty(\mathbb{R})$ , which is sectorial if  $\theta$  satisfies (1.4). According to [4], its spectrum consists of a sequence of discrete simple eigenvalues, denoted in nondecreasing modulus order by  $\lambda_n = \lambda_n(m, \theta)$ ,  $|\lambda_n| \rightarrow +\infty$ . The associated spectral projections are of rank 1, and E.-B. Davies showed in [4] that for all  $m \in ]0, +\infty[$  and  $\theta \neq 0$  satisfying (1.4), for all  $\alpha > 0$ , there exists  $N = N(m, \theta, \alpha) \geq 0$  such that the instability indices  $\kappa_n(m, \theta)$  of  $\mathcal{A}(m, \theta)$  satisfy  $\kappa_n(m, \theta) \geq n^\alpha$  for  $n \geq N$ . This statement has been improved in the case  $m = 2$  of the harmonic oscillator (sometimes referred as the Davies operator), since E.-B. Davies and A. Kuijlaars showed [5] that  $\kappa_n(2, \theta)$  grows exponentially fast as  $n \rightarrow +\infty$ , with an explicit rate  $c(\theta)$  : there exists an explicit  $c(\theta)$  such that

$$\lim_{n \rightarrow +\infty} \frac{1}{n} \log \kappa_n(2, \theta) = c(\theta). \quad (1.5)$$

The purpose of this Note is to prove that this statement actually holds for the so-called *complex Airy operator*  $\mathcal{A}(1, \theta)$  and for the even anharmonic oscillators  $\mathcal{A}(2k, \theta)$ ,  $k \geq 1$ .

### Acknowledgments

I am greatly indebted to Professor Bernard Helffer for his help, advice and comments ; I am also grateful to Thierry Ramond, Christian Gérard and André Martinez for their valuable discussions. I acknowledge the support of the ANR NOSEVOL.

## 2 Non-selfadjoint anharmonic oscillators

We first deal with the complex Airy operator  $\mathcal{A}(1, \theta)$  defined in (1.3). We show that the corresponding instability indices  $\kappa_n(1, \theta)$  grow like in (1.5) as  $n \rightarrow +\infty$ . More precisely, we get asymptotic expansions in powers of  $n^{-1}$  as  $n \rightarrow +\infty$ .

**Theorem 2.1.** *Let  $0 < |\theta| < 3\pi/4$ . There exists a real sequence  $(\alpha_j(\theta))_{j \geq 1}$  such that the instability indices  $\kappa_n(1, \theta)$  of  $\mathcal{A}(1, \theta)$  satisfy, as  $n \rightarrow +\infty$ ,*

$$\exp(-C(\theta)(n - 1/2)) \kappa_n(1, \theta) = \frac{K(\theta)}{\sqrt{n}} \left( 1 + \sum_{j=1}^{+\infty} \alpha_j(\theta) n^{-j} \right) + \mathcal{O}(n^{-\infty}), \quad (2.1)$$

where

$$C(\theta) = \pi m_\theta^{3/2} |\sin \theta| \quad \text{and} \quad K(\theta) = \frac{1}{2\sqrt{3} |\sin \theta| m_\theta^{1/4}},$$

with

$$m_\theta = \sqrt{1 + \frac{\sin^2(2\theta/3)}{\sin^2 \theta} - 2 \frac{\cos(\theta/3) \sin(2\theta/3)}{\sin \theta}} > 0.$$

**Sketch of the proof :** Let us first recall that all the eigenvalues of  $\mathcal{A}(m, \theta)$ ,  $m \in \mathbb{N}$ , have associated spectral projections of rank 1, see Lemma 5 in [4]. Hence, one can easily check that [3]

$$\kappa_n(m, \theta) = \frac{\|u_n\|^2}{\langle u_n, \bar{u}_n \rangle}, \quad (2.2)$$

where  $u_n$  denotes an eigenfunction associated with the  $n$ -th eigenvalue of  $\mathcal{A}(m, \theta)$ . We get rid of the singularity of the potential at  $x = 0$  by decomposing  $\mathcal{A}(1, \theta)$  into its Dirichlet and Neumann realizations  $\mathcal{A}^D(1, \theta)$  and  $\mathcal{A}^N(1, \theta)$  on  $\mathbb{R}^+$ . We then compute their instability indices

$$\kappa_n^{D/N}(1, \theta) = \frac{\int_{\mathbb{R}^+} |Ai(\mu_n^{D/N} + e^{i\theta/3}x)|^2 dx}{\left| \int_{\mathbb{R}^+} Ai(\mu_n^{D/N} + e^{i\theta/3}x)^2 dx \right|}, \quad (2.3)$$

given by formula (2.2), where  $x \mapsto Ai(\mu_n^{D/N} + e^{i\theta/3}x)$  is the  $n$ -th eigenfunction of  $\mathcal{A}^{D/N}(1, \theta)$ ,  $\mu_n^D$  (resp.  $\mu_n^N$ ) being the  $n$ -th (negative) zero (resp. critical point) of the Airy function  $Ai$  (see [2], [8]). We estimate the numerator in (2.3) by using the well-known asymptotic expansion of  $Ai$  at infinity in the complex plane (see [1]), and the Laplace method brings an  $\exp(c_\theta |\mu_n^{D/N}|^{3/2})$  term in  $\kappa_n^{D/N}(1, \theta)$ ,  $c_\theta > 0$ . The integral in the denominator of (2.3), after deformation of the path of integration by homotopy, is equal to

$$\int_{\mu_n^{D/N}}^{+\infty} Ai^2(x) dx = Ai'^2(\mu_n^{D/N}) \quad (2.4)$$

(it is indeed straightforward, using Airy equation, to check that  $x \mapsto xAi^2(x) - Ai'^2(x)$  is a primitive for  $Ai^2$ ). Hence the expansion of  $Ai'(-z)$  as  $z \rightarrow +\infty$ , given in [1], provides an asymptotic expansion for (2.4) in powers of  $|\mu_n^{D/N}|^{-3/2}$ . The statement follows from the behavior of  $\mu_n^{D/N}$  as  $n \rightarrow +\infty$ , since we have asymptotic expansions for  $(n-1/4)^{-2/3}|\mu_n^D|$  (resp.  $(n-3/4)^{-2/3}|\mu_n^N|$ ) in powers of  $(n-1/4)^{-2}$  (resp.  $(n-3/4)^{-2}$ ).  $\square$

Notice that the exponential instability appears as soon as  $\theta \neq 0$ . We have a similar statement for even anharmonic oscillators :

**Theorem 2.2.** *Let  $k \in \mathbb{N}^*$  and  $\theta$  be such that  $0 < |\theta| < \frac{(k+1)\pi}{2k}$ . If  $\kappa_n(2k, \theta)$  denotes the  $n$ -th instability index of  $\mathcal{A}(2k, \theta) = -\frac{d^2}{dx^2} + e^{i\theta}x^{2k}$ , then there exist  $K(2k, \theta) > 0$  and a real sequence  $(C^j(2k, \theta))_{j \geq 1}$  such that*

$$e^{-c_k(\theta)n} \kappa_n(2k, \theta) = \frac{K(2k, \theta)}{\sqrt{n}} \left( 1 + \sum_{j=1}^{+\infty} C^j(2k, \theta) n^{-j} \right) + \mathcal{O}(n^{-\infty}) \quad (2.5)$$

as  $n \rightarrow +\infty$ , with

$$c_k(\theta) = \frac{2(k+1)\sqrt{\pi}\Gamma\left(\frac{k+1}{2k}\right)\varphi_k(\xi_k)}{\Gamma\left(\frac{1}{2k}\right)} > 0, \quad (2.6)$$

where

$$\xi_k = \left( \frac{\tan(|\theta|/(k+1))}{\sin(k|\theta|/(k+1)) + \cos(k|\theta|/(k+1))\tan(|\theta|/(k+1))} \right)^{\frac{1}{2k}} \quad (2.7)$$

$$\varphi_k(\xi) = \operatorname{Im} \int_0^{\xi e^{i\frac{\theta}{2(k+1)}}} (1-t^{2k})^{1/2} dt. \quad (2.8)$$

**Sketch of the proof :** We first perform an analytic dilation and a scale change to recover the semiclassical selfadjoint anharmonic oscillator  $\mathcal{P}_h(2k) = -\hbar^2 \frac{d^2}{dx^2} + x^{2k} - 1$ , with

$$h = h_n = |\lambda_n(2k, \theta)|^{-\frac{k+1}{2k}}.$$

The  $n$ -th instability index of  $\mathcal{A}(2k, \theta)$  then writes

$$\kappa_n(2k, \theta) = \frac{\int_{\mathbb{R}} |\psi_h(e^{i\frac{\theta}{2(k+1)}x})|^2 dx}{\int_{\mathbb{R}} \psi_h^2(x) dx} \quad (2.9)$$

where  $\psi_h$  solves  $\mathcal{P}_h(2k)\psi_h = 0$ ,  $\psi_h \in L^2(\mathbb{R})$  (see (2.2), after deformation of the integration path in the denominator). The complex WKB method (see [10], [14], [6]) and the analysis of the Stokes lines of  $\mathcal{P}_h(2k)$  provide an asymptotic expansion of  $\psi_h(e^{i\frac{\theta}{2(k+1)}x})$  as  $h \rightarrow 0$ , which enables us to determine the asymptotic behaviour of the numerator in (2.9), using again the Laplace method.

On the real axis,  $\psi_h$  is treated separately in its oscillatory region  $[-1 + \delta, 1 - \delta]$ ,  $\delta > 0$ , and in the neighbourhood of the turning points  $\pm 1$ . Hence, the stationary phase method leads to an asymptotic expansion in powers of  $h$  of the denominator in (2.9). Finally, the statement follows from the Bohr-Sommerfeld quantization rule for  $h_n$  (see [7], Exercise 12.3) or Weyl formula [9].  $\square$

In the harmonic case  $k = 1$  (Davies operator), the first term in (2.5) yields the Davies-Kuijlaars theorem [5] :

$$\lim_{n \rightarrow +\infty} \frac{1}{n} \log \|\Pi_n\| = c_1(\theta) = 4\varphi_1 \left( \frac{1}{\sqrt{2 \cos(\theta/2)}} \right) = 2\operatorname{Re} f \left( \frac{e^{i\theta/4}}{\sqrt{2 \cos(\theta/2)}} \right)$$

where  $f(z) = \log(z + \sqrt{z^2 - 1}) - z\sqrt{z^2 - 1}$ .

### 3 Eigenfunctions and semigroups

The following theorem has been proved in [2] in the case of complex Airy operator  $\mathcal{A}(1, \theta)$ , and in [4] in the harmonic case ( $k = 1$ ), as well as for  $\mathcal{A}(2k, \theta)$ ,  $k \geq 2$ ,  $|\theta| < \frac{\pi}{2}$ . The proof actually extends to any operator  $\mathcal{A}(2k, \theta)$  with  $|\theta| < \frac{(k+1)\pi}{2k}$  :

**Theorem 3.1.** *For any  $m = 1, 2k$ ,  $k \geq 1$ , and any  $\theta$  satisfying (1.4), the eigenfunctions of  $\mathcal{A}(m, \theta)$  form a complete set of the space  $L^2(\mathbb{R})$ .*

Notice however that the eigenfunctions of  $\mathcal{A}(1, \theta)$  and  $\mathcal{A}(2k, \theta)$ ,  $k \geq 1$ , can not form a Riesz basis because of the growth of the instability indices as  $n \rightarrow +\infty$ .

Theorem 3.1 and the previous estimates enable us to study the convergence of the operator series defining the semigroup  $e^{-t\mathcal{A}(m, \theta)}$  associated with  $\mathcal{A}(m, \theta)$  when decomposed along the projections  $\Pi_n$ .

The following statement extends the result of [5] in the harmonic case.

**Corollary 3.2.** *Let  $|\theta| \leq \pi/2$  and  $e^{-t\mathcal{A}(m, \theta)}$  be the semigroup generated by  $\mathcal{A}(m, \theta)$ ,  $\lambda_n = \lambda_n(m, \theta)$  the eigenvalues of  $\mathcal{A}(m, \theta)$ , and  $\Pi_n = \Pi_n(m, \theta)$  the*

associated spectral projections.

Let  $T(\theta) = c_1(\theta)/\cos(\theta/2)$ , where  $c_1(\theta)$  is the constant in (2.6). The series

$$\Sigma_{m,\theta}(t) = \sum_{n=1}^{+\infty} e^{-t\lambda_n(m,\theta)} \Pi_n(m,\theta)$$

is not normally convergent in cases  $m = 1$  for any  $t > 0$ , and  $m = 2$  for  $t < T(\theta)$ ; in cases  $m = 2$  for  $t > T(\theta)$ , and  $m = 2k$  for any  $t > 0$ ,  $k \geq 2$ , the series converges normally towards  $e^{-t\mathcal{A}(m,\theta)}$  and, for  $N$  sufficiently large and for some constants  $C_1 = C_1(k,\theta)$  and  $C_2 = C_2(\theta)$ , the following estimate holds :

$$\|e^{-t\mathcal{A}(m,\theta)}(I - \Pi_{<N})\| \leq \begin{cases} \frac{C_1}{\sqrt{N}} e^{c_k(\theta)n} \exp(-t\operatorname{Re} \lambda_N), & k \geq 2 \\ \frac{C_2}{\sqrt{N}} \exp(-2\cos(\theta/2)(t - T(\theta))N), & k = 1, t > T \end{cases} \quad (3.1)$$

where  $\Pi_{<N} = \Pi_1 + \dots + \Pi_{N-1}$  denote the projection on the first  $N - 1$  eigenspaces.

## Références

- [1] M. Abramowitz and I. Stegun. *Handbook of mathematical functions*, National bureau of standards, 1964.
- [2] Y. Almog. *The stability of the normal state of superconductors in the presence of electric currents*, Siam J. Math. Anal. 40 (2) (2008) 824-850.
- [3] A. Aslanyan and E.-B. Davies. *Spectral instability for some Schrödinger operators*, Numer. Math. 85 (2000) 525-552.
- [4] B. Davies. *Wild spectral behaviour of anharmonic oscillators*, Bull. London. Math. Soc. 32 (2000) 432-438.
- [5] B. Davies and A. Kuijlaars. *Spectral asymptotics of the non-self-adjoint harmonic oscillator*, J. London Math. Soc. (2) 70 (2004) 420-426.
- [6] C. Gérard and A. Grigis. *Precise estimates of tunneling and eigenvalues near a potential barrier*, Journal of differential equations 72 (1988) 149-177.
- [7] A. Grigis and J. Sjöstrand. *Microlocal analysis for differential operators, an introduction*, London Math. Soc. Lecture Note Series 196, 1994.
- [8] B. Helffer. *On pseudo-spectral problems related to a time dependent model in superconductivity with electric current*, Confluentes Math. 3, n. 2 (2011) 237-251.
- [9] B. Helffer and D. Robert. *Asymptotique des niveaux d'énergie pour des hamiltoniens à un degré de liberté*, Duke Math. J. Volume 49, n. 4 (1982) 853-868.
- [10] F. W. J. Olver. *Asymptotics and special functions*, Academic Press, 1974.
- [11] S. Roch and B. Silbermann. *C\*-algebras techniques in numerical analysis*, J. Oper. Theory 35 (1996) 241-280.

- [12] J. Sjöstrand. *Lecture notes : Spectral properties of non-self-adjoint operators*, J. E. D. P., Exposé n. I (2009) 111p.
- [13] L. N. Trefethen and M. Embree. *Spectra and pseudospectra. The behavior of nonnormal matrices and operators*, Princeton University Press. Princeton, N. J. 2005.
- [14] A. Voros. *Spectre de l'équation de Schrödinger et méthode BKW*, Publications mathématiques d'Orsay, 81.09, 1981.