

# MULTI-PARAMETER RESOLVENT TRACE EXPANSION I. FOR ELLIPTIC BOUNDARY PROBLEMS

BORIS VERTMAN

ABSTRACT. Various aspects of semi-classical analysis require discussion of asymptotic expansions in two parameters weighted with smooth functions, which in the setting of manifolds with boundary does not follow in an obvious way from the classical parametric elliptic theory. In the present paper we establish multi-parameter resolvent trace expansions for elliptic boundary value problems, polyhomogeneous both in the resolvent and the auxiliary parameters. Such multi-parameter resolvent trace expansions have been used in the setting of revolution surfaces with unexpected applications to regularized sums of zeta-determinants. Another example where these ideas play a prominent role is the asymptotics of determinants for discrete Laplacians on tori under growing discretization parameter.

## CONTENTS

1. Introduction and formulation of the main result	1
1.1. Introduction and motivation	1
1.2. Formulation of the main result	2
2. Resolvent kernel as a polyhomogeneous conormal distribution	3
2.1. Resolvent of an elliptic boundary value problem	3
2.2. Construction of the resolvent blowup space	4
3. Composition of polyhomogeneous Schwartz kernels	7
4. Multi-parameter resolvent trace expansion	11
4.1. Initial parametrix and Neumann series	11
4.2. Trace norm estimates and expansions	12
Acknowledgements	16
References	16

## 1. INTRODUCTION AND FORMULATION OF THE MAIN RESULT

**1.1. Introduction and motivation.** Interest in the multi-parameter resolvent trace expansions arises in view of their various applications. A particular example of such an application is a joint project with Lesch [LEVE13] where we compare the zeta-determinant of the Laplace Beltrami operator on a surface of revolution with a regularized sum of zeta-determinants for scalar operators arising from a spectral decomposition on the cross section of the revolution surface.

More generally, consider a manifold with fibered boundary  $\mathbb{S}^d \times M$ , with an elliptic boundary problem  $(\mathcal{A}, \mathcal{B})$  represented after the eigenspace decomposition

---

*Date:* This document was compiled on: November 13, 2021.

The author was supported by the Bonn Hausdorff Center and University Münster.

on  $\mathbb{S}^d$  by an infinite sum of elliptic boundary problems  $(A_n, B_n), n \in \mathbb{N}_0$  on  $M$ . Assume the boundary problems admit well defined zeta-determinants. Then the methods elaborated in [LEVE13] together with our main theorem here equate the zeta-determinant of  $\mathcal{A}_{\mathcal{B}}$  to the regularized sum of zeta determinants for  $A_n, B_n$ , up to a locally computable error term.

Another application of multi-parameter resolvent trace expansions appears in the study of asymptotics for determinants of graph Laplacians on discretized tori. In [VE15] we show that the constant term in the asymptotics of the combinatorial determinant for the graph Laplacian on the discretized tori is given in terms of the zeta-determinant of the Laplace Beltrami operator on the smooth torus manifold.

Finally, we expect a multi-parameter resolvent trace expansion to play a role in establishing an asymptotic expansion of the Bergmann kernel in the setting of complex manifolds with singular metric structure along divisors.

**1.2. Formulation of the main result.** Consider a compact manifold  $M$  of dimension  $m$  with boundary  $\partial M$ , equipped with a Hermitian vector bundle  $(E, h^E)$  of rank  $p$ . We recall the notion of an elliptic boundary problem from Seeley [SEE69]. Let  $A \in \text{Diff}^q(M, E)$  denote a differential operator on  $M$  with values in  $E$  of  $q$ -th order with  $pq \in 2\mathbb{N}_0$ . The operator  $A$  is elliptic if its principal symbol  $\sigma(A)(p, \zeta)$  is invertible for  $(p, \zeta) \in T^*M \setminus \{0\}$ . Assume,  $A$  satisfies the Agmon condition in a fixed cone  $(-\Gamma')$  with  $\Gamma' = \{z \in \mathbb{C} \mid \arg(z) \in (\theta_1, \theta_2)\}$  of the complex plane, i.e.  $(\sigma(A) + z^q)$  is invertible for  $z^q \in \Gamma'$ . Consider a system of differential operators  $B = (B_1, \dots, B_{pq/2})$  on  $\partial M$ , such that  $(A, B)$  defines an elliptic boundary problem satisfying the Agmon condition on  $\Gamma'$  in the sense of [SEE69, Def. 1,2].

Under this setup,  $(A, B)$  defines a closed unbounded operator  $A_B$  on  $L^2(M, E)$ , obtained as the graph closure of  $A$  acting on  $u \in C^\infty(M, E)$  satisfying the boundary conditions  $Bu = 0$ . Moreover,  $(A_B + z^q)$  is invertible for  $z^q \in \Gamma'$  sufficiently large, cf. [SEE69, p. 911], and the seminal work of Seeley [SEE69] establishes an expansion of the resolvent  $(A_B + z^q)^{-1}$  as  $|z| \rightarrow \infty$ . In the present paper we fix a finite collection of scalar smooth potentials  $V_1, \dots, V_n \in C^\infty(M)$  which are assumed to be nowhere vanishing along the boundary  $\partial M$ . Set  $\Gamma := \{z \in \mathbb{C} \mid z^q \in \Gamma'\}$  and write  $\Gamma^n$  for its  $n$ -th Cartesian product. Consider  $\lambda = (\lambda_1, \dots, \lambda_n) \in \Gamma^n$  and the corresponding multi-parameter family

$$A_B(\lambda) := A_B + \sum_{k=1}^n \lambda_k^q V_k.$$

For  $z \in \Gamma$  sufficiently large,  $(A_B(\lambda) + z^q)$  is invertible. For  $qN > m$  the  $N$ -th power of the resolvent  $(A_B(\lambda) + z^q)^{-N}$  is trace class and our main result establishes a multi-parameter expansion of the resolvent trace  $\text{Tr}(A_B(\lambda) + z^q)^{-N}$ , polyhomogeneous in  $(z, \lambda) \in \Gamma^{n+1}$ .

**Theorem 1.1.** *Consider any multiindex  $\alpha \in \mathbb{N}_0^n$  and  $\beta \in \mathbb{N}_0$ . Fix  $N \in \mathbb{N}$  such that  $qN > m$ . Then there exist  $e_i \in C^\infty(M \times (\Gamma^{n+1} \cap \mathbb{S}^n))$ , such that*

$$\partial_\lambda^\alpha \partial_z^\beta \text{Tr}(A_B(\lambda) + z^q)^{-N} \sim \sum_{j=0}^{\infty} e_j \left( \frac{(\lambda, z)}{|\lambda, z|} \right) |(\lambda, z)|^{-1-qN-j-|\alpha|-\beta+m}.$$

Let us mention that the statement has a straightforward extension to the case of additional summands in the expression for  $A_B(\lambda)$  of the form  $\lambda_k^\alpha D_k$ , where  $D_k \in \text{Diff}^p(M, E)$  and  $\alpha + p < q$ . Moreover the arguments extend to the case of matrix-valued potentials  $V_1, \dots, V_n \in C^\infty(M, \text{End}(E))$ , assuming that the action of their restrictions to the boundary is scalar.

This paper is organized as follows. In §2 we recast the symbolic expansion of the resolvent  $(A_B + z^q)^{-1}$  in terms of polyhomogeneity properties of its Schwartz kernel lifted to an appropriate blowup of  $\mathbb{R}^+ \times M^2$ . In §3 we establish a composition result for the polyhomogeneous conormal distributions on the blowup of  $\mathbb{R}^+ \times M^2$ . In §4 we employ this microlocal characterization of the resolvent kernel to establish the multi-parameter resolvent trace expansion. While the resolvent exists by standard arguments, its microlocal description is obtained by constructing the corresponding Schwartz kernel as a polyhomogeneous conormal distribution on the blowup space using an iterative Neumann series argument

## 2. RESOLVENT KERNEL AS A POLYHOMOGENEOUS CONORMAL DISTRIBUTION

**2.1. Resolvent of an elliptic boundary value problem.** Seeley [SEE69] provides a careful construction of the resolvent for the elliptic boundary problem  $A_B$ . More precisely, in view of [SEE69, Theorem 1, Lemma 2, (25), (32), (49)] we may state the following

**Theorem 2.1.** *Let  $\Gamma_R := \{\mu \in \Gamma \mid |\mu| \geq R\}$ . Consider a local coordinate neighborhood  $\mathcal{U} \subset [0, 1)_x \times \partial M$  in the collar of the boundary, with local coordinates  $\{x, y = (y_1, \dots, y_{m-1})\}$ . Then for any  $j \in \mathbb{N}_0$  there exist  $d_{-q-j} \in C^\infty(\mathcal{U} \times \mathbb{R}^m \times \Gamma, \text{Hom}(E \upharpoonright \mathcal{U}))$ , homogeneous of order  $(-q - j)$  in  $(x^{-1}, \zeta, \gamma, \mu)$ , where  $(x, y, \zeta, \gamma, \mu) \in \mathcal{U} \times \mathbb{R}_\zeta^{m-1} \times \mathbb{R}_\gamma \times \Gamma_\mu$ , such that for  $R > 0$  sufficiently large and  $\mu \in \Gamma_R$  the Schwartz kernel<sup>1</sup>*

$$\begin{aligned} & (A_B + \mu^q)^{-1}(x, y, \tilde{x}, \tilde{y}) - (A + \mu^q)^{-1}(x, y, \tilde{x}, \tilde{y}) \\ & - \sum_{j=0}^{N-1} (2\pi)^{-m} \int_{\mathbb{R}^{m-1}} \int_{\mathbb{R}} e^{i\langle y - \tilde{y}, \zeta \rangle} e^{-i\tilde{x}\gamma} d_{-q-j}(x, y, \zeta, \gamma, \mu) d\gamma d\zeta \\ & =: K_{A_B}(x, y, \tilde{x}, \tilde{y}; \mu) - K_A(x, y, \tilde{x}, \tilde{y}; \mu) - \sum_{j=0}^{N-1} \text{Op}(d_{-q-j}), \end{aligned}$$

is uniformly  $O(|\mu|^{m-q-N})$  as  $|\mu| \rightarrow \infty$ . Here, the first term  $K_{A_B}$  denotes the resolvent kernel of  $A_B$  near the boundary, whereas the second term  $K_A$  refers to an interior symbolic parametrix<sup>2</sup> defined without taking into account the boundary conditions. Note however that the symbol corresponding to  $K_A$  determines the choice of  $d_{-q-j}$ ,  $j \in \mathbb{N}_0$  by a system of differential equations [SEE69, (9), (10), (11)].

<sup>1</sup>The notation  $\text{Op}$  denotes the integral kernel of the operator rather than the operator itself.

<sup>2</sup>The interior symbolic parametrix  $K_A$  is obtained by inverting the symbol of  $(A + \mu^q)$  algebraically by an iteration as in [SEE69, (5), (6)], and studying the corresponding operator. Such a parametrix is defined independently of the boundary conditions and provides a good parametrix for  $(A + \mu^q)^{-1}$  away from boundary.

**2.2. Construction of the resolvent blowup space.** Assume for simplicity that  $\mu$  varies along a ray within  $\Gamma$ , which we identify with  $\mathbb{R}^+$ . Then the Schwartz kernel  $K_{A_B}$  of the resolvent  $(A_B + \mu^q)^{-1}$  is a distribution on  $\mathbb{R}_{1/\mu}^+ \times M^2$ . Choose local coordinates  $(x, y)$  and  $(\tilde{x}, \tilde{y})$  on the two copies of  $M$  in a collar neighborhood of the boundary, where  $x$  and  $\tilde{x}$  are the boundary defining functions. The Schwartz kernel  $K_{A_B}$  admits a non-uniform behaviour at

$$\begin{aligned}\mathcal{C} &:= \{\mu = \infty, x = \tilde{x} = 0, y = \tilde{y}\}, \\ \mathcal{D} &:= \{\mu = \infty, (x, y) = (\tilde{x}, \tilde{y})\}.\end{aligned}\tag{2.1}$$

This non-uniform behaviour is resolved by considering an appropriate blowup  $\mathcal{M}_b^2$  of  $\mathbb{R}^+ \times M^2$  at  $\mathcal{C}$  and  $\mathcal{D}$ , a procedure introduced by Melrose, see [MEL93], such that the kernels  $K_{A_B}, K_A$  lift to polyhomogeneous distributions on the manifold with corners  $\mathcal{M}_b^2$  in the sense of the following definition.

**Definition 2.2.** *Let  $X$  be a manifold with corners, with all boundary faces embedded, and  $\{(H_i, \rho_i)\}_{i=1}^N$  an enumeration of its boundaries and corresponding defining functions. For any multi-index  $b = (b_1, \dots, b_N) \in \mathbb{C}^N$  we write  $\rho^b = \rho_1^{b_1} \dots \rho_N^{b_N}$ . Denote by  $\mathcal{V}_b(X)$  the space of smooth vector fields on  $X$  which are tangent to all boundary faces. All distributions on  $X$  will be assumed to be restrictions of distributions defined on a closed manifold extending  $X$ . A distribution  $\omega$  on  $X$  is said to be conormal if  $\omega \in \rho^b L^\infty(X)$  for some  $b \in \mathbb{C}^N$ , and  $V_1 \dots V_\ell \omega \in \rho^b L^\infty(X)$ , for all  $V_j \in \mathcal{V}_b(X)$  and for every  $\ell \geq 0$ . An index set  $E_i = \{(\gamma, p)\} \subset \mathbb{C} \times \mathbb{N}$  satisfies the following hypotheses:*

- (1) if  $(\gamma_j, p_j) \in E_i$  and  $|(\gamma_j, p_j)| \rightarrow \infty$  then  $\operatorname{Re}(\gamma_j) \rightarrow \infty$ ,
- (2) if  $(\gamma, p) \in E_i$ , then  $(\gamma + j, p') \in E_i$  for all  $j \in \mathbb{N}_0$  and  $0 \leq p' \leq p$ ,

An index family  $E = (E_1, \dots, E_N)$  is an  $N$ -tuple of index sets. Finally, we say that a conormal distribution  $w$  is polyhomogeneous on  $X$  with index family  $E$ , we write  $\omega \in \mathcal{A}_{\text{phg}}^E(X)$ , if  $\omega$  is conormal and if in addition, near each  $H_i$ ,

$$\omega \sim \sum_{(\gamma, p) \in E_i} a_{\gamma, p} \rho_i^\gamma (\log \rho_i)^p, \text{ as } \rho_i \rightarrow 0,\tag{2.2}$$

with coefficients  $a_{\gamma, p}$  conormal on  $H_i$ , polyhomogeneous with index  $E_j$  at any  $H_i \cap H_j$ <sup>3</sup>.

We also need to consider polyhomogeneous distributions on a manifold with corners  $X$ , conormal to an embedded submanifold  $Y \subset X$ . The basic space  $I^m(\mathbb{R}^n, \{0\})$  consists of compactly supported distributions whose Fourier transform is given by a symbol of order  $(m - n/4)$ .  $I^m(\mathbb{R}^n, \{0\})$  is invariant under local diffeomorphisms and thus makes sense on any manifold around an isolated point.

For an embedded  $k$ -submanifold  $S \subset X$ , any point in  $S$  admits an open neighborhood  $\mathcal{V}$  in  $X$  which can be locally decomposed as a product  $\mathcal{V} = X' \times X''$  so that  $\mathcal{V} \cap S = X' \times \{p\}, p \in X''$ . The space  $I^m(X, S)$  is defined (locally) as the space of smooth functions on  $X'$  with values  $I^{m+\dim X'/4}(X'', \{p\})$ . The normalization is chosen to give pseudo-differential operators their expected orders. All distributions

<sup>3</sup>The asymptotic expansion for  $\omega$  means in precise terms that the difference between  $\omega$  and any finite portion of the expansion vanishes at the rate of the next term in the expansion, with a corresponding property for all higher derivatives, and with decay order at other boundary faces determined by the index family  $E$ .

in  $I^m(X, S)$  are locally restrictions of distributions on an ambient space, which are conormal to any smooth extension of  $S$  across  $\partial X$ .

Choosing now index sets  $E$  for each boundary face of  $X$  as in Definition 2.2, we define a space  $\mathcal{A}_{\text{phg}}^E(X, Y)$  as the space of distributions conormal to  $Y$ , with polyhomogeneous expansions as in Eq. (2.2) at all boundary faces and with coefficients conormal to the intersection of  $Y$  with each boundary face.

We now continue with the definition of a blown-up space<sup>4</sup>  $\mathcal{M}_b^2$ , to which the Schwartz kernels  $K_A$  and  $\text{Op}(d_{-q-j})$  lift to polyhomogeneous distributions, possibly conormal to an embedded submanifold. Blowing up  $\mathbb{R}^+ \times M^2$  at  $\mathcal{C}$  and  $\mathcal{D}$  amounts in principle to introducing polar coordinates in  $\mathbb{R}^+ \times (\mathbb{R}^+)^2$  at

$$\begin{aligned} \mathcal{C}' &:= \{(1/\mu, x, y) \in \mathbb{R}^+ \times (\mathbb{R}^+)^2 \mid \mu = \infty, x = \tilde{x} = 0, y = \tilde{y}\}, \\ \mathcal{D}' &:= \{(1/\mu, x, y) \in \mathbb{R}^+ \times (\mathbb{R}^+)^2 \mid \mu = \infty, (x, y) = (\tilde{x}, \tilde{y})\}, \end{aligned} \quad (2.3)$$

together with a unique minimal differential structure with respect to which these coordinates are smooth.

We first perform a blowup of  $\mathcal{C}$ . The resulting space  $[\mathbb{R}^+ \times M^2; \mathcal{C}]$  is defined as the union of  $\mathbb{R}^+ \times M^2 \setminus \mathcal{C}$  with the inward-pointing spherical normal bundle of  $\mathcal{C}$  in  $\mathbb{R}^+ \times M^2$ . The blowup  $[\mathbb{R}^+ \times M^2; \mathcal{C}]$  is endowed with the unique minimal differential structure with respect to which smooth functions in the interior of  $\mathbb{R}^+ \times M^2$  and polar coordinates on  $\mathbb{R}^+ \times M^2$  around  $\mathcal{C}$  are smooth. This blowup introduces a new boundary hypersurface, which we refer to as the front face  $\text{ff}$ . The other boundary faces are as follows. The union of  $\text{ff}$  and the right face  $\text{rf}$  is the lift of  $\{x = 0\}$ . The union of  $\text{ff}$  and the left face  $\text{lf}$  is the lift of  $\{\tilde{x} = 0\}$ . The union of  $\text{ff}$  and the temporal face  $\text{tf}$  is the lift of  $\{\mu = \infty\}$ .

The actual blowup space  $\mathcal{M}_b^2$  is obtained by a blowup of  $[\mathbb{R}^+ \times M^2; \mathcal{C}]$  along the lift of the diagonal  $\mathcal{D}$ . The resulting blowup space  $\mathcal{M}_b^2$  is defined as before by cutting out the submanifold and replacing it with its inward-pointing spherical normal bundle. This second blowup introduces an additional boundary hypersurface  $\text{td}$ , the temporal diagonal.  $\mathcal{M}_b^2$  is a manifold with boundaries and corners, illustrated below.




FIGURE 1. The blowup  $\mathcal{M}_b^2 = [[\mathbb{R}^+ \times M^2, \mathcal{C}], \mathcal{D}]$ .

The blowup  $\mathcal{M}_b^2$  is equipped with the canonical ‘blow-down map’  $\beta : \mathcal{M}_b^2 \rightarrow \mathbb{R}^+ \times M^2$ , which we discuss below.

<sup>4</sup>Do not confuse  $\mathcal{M}_b^2$  with the b-stretched double space in [MEL93]

Denote by  $Y := \{(\mu, p, \tilde{p}) \in \mathbb{R}^+ \times M^2 \mid p = \tilde{p}\}$  the diagonal submanifold and denote by  $\beta^*Y$  its lift to  $\mathcal{M}_b^2$ . We denote by  $\mathcal{A}_{\text{phg}}^{l,p}(\mathcal{M}_b^2, \beta^*Y)$  the space of distributions that lift<sup>5</sup> to polyhomogeneous conormal distributions on the blowup space  $\mathcal{M}_b^2$ , with leading order  $(-m + l)$  at the front face  $\text{ff}$ , leading order  $(-m + p)$  at the temporal diagonal  $\text{td}$ , index sets  $(\mathbb{N}_0, \mathbb{N}_0)$  at the left and right boundary faces, vanishing to infinite order at  $\text{tf}$ , conormal at the interior submanifold  $\beta^*Y$ . The space of such distributions without a conormal singularity is denoted by  $\mathcal{A}_{\text{phg}}^{l,p}(\mathcal{M}_b^2)$ .

A choice of projective coordinates on  $\mathcal{M}_b^2$  is given as follows. Near the top corner of  $\text{ff}$  away from  $\text{tf}$  projective coordinates are given by

$$\rho = \frac{1}{\mu}, \quad \xi = \frac{x}{\rho}, \quad \tilde{\xi} = \frac{\tilde{x}}{\rho}, \quad w = \frac{y - \tilde{y}}{\rho}, \quad \tilde{y}, \quad (2.4)$$

where in these coordinates  $\rho, \xi, \tilde{\xi}$  are the defining functions of the faces  $\text{ff}$ ,  $\text{rf}$  and  $\text{lf}$  respectively. For the bottom corner of  $\text{ff}$  near  $\text{rf}$  projective coordinates are given by

$$\tau = (\mu\tilde{x})^{-1}, \quad s = \frac{x}{\tilde{x}}, \quad u = \frac{y - \tilde{y}}{\tilde{x}}, \quad \tilde{x}, \quad \tilde{y}, \quad (2.5)$$

where in these coordinates  $\tau, s, \tilde{x}$  are defining functions of  $\text{tf}$ ,  $\text{rf}$  and  $\text{ff}$  respectively. For the bottom corner of  $\text{ff}$  near  $\text{lf}$  projective coordinates are obtained by interchanging the roles of  $x$  and  $\tilde{x}$ . Projective coordinates on  $\mathcal{M}_b^2$  near the top of  $\text{td}$  away from  $\text{tf}$  are given by

$$\eta = \tau, \quad S = \frac{s - 1}{\eta}, \quad U = \frac{u}{\eta}, \quad \tilde{x}, \quad \tilde{y}. \quad (2.6)$$

The temporal face  $\text{tf}$  is not in this coordinate chart but corresponds to the limit  $|(S, U)| \rightarrow \infty$ . The boundary faces  $\text{ff}$  and  $\text{td}$  are defined by  $\tilde{x}, \eta$ , respectively. The blowup  $\mathcal{M}_b^2$  is related to the original space  $\mathbb{R}^+ \times M^2$  via the obvious ‘blow-down map’

$$\beta : \mathcal{M}_b^2 \rightarrow \mathbb{R}^+ \times M^2,$$

which is in local coordinates simply the coordinate change back to  $(1/\mu, (x, y), (\tilde{x}, \tilde{y}))$ . The blowup  $\mathcal{M}_b^2$  is similar to the blowup space construction for incomplete conical singularities by Mooers [Moo99] with the difference that here the blowup is not parabolic in  $\mu^{-1}$ -direction. It will be crucial in the argument below to note that the restriction of  $\beta$  to  $\text{ff}$  is a fibration over the diagonal of  $\partial M \times \partial M$  with the fibre equal to the quarter sphere  $\mathbb{S}_{++}^m$ . Similarly, the restriction of  $\beta$  to  $\text{td}$  is a fibration over the diagonal of  $M \times M$  with the fibre equal to the hemisphere  $\mathbb{S}_+^m$ .

**Theorem 2.3.** *Consider the elliptic boundary value problem  $(A, B)$  with a differential operator  $A$  of order  $q$  on a compact manifold  $M$  of dimension  $m$  with boundary  $\partial M$ . Then the resolvent kernel  $K_{AB}$  and the Schwartz kernel  $K_A$  corresponding to an interior symbolic parametrix kernel are both elements of  $\mathcal{A}_{\text{phg}}^{q,q}(\mathcal{M}_b^2, \beta^*Y)$ , where  $Y := \{(\mu, p, \tilde{p}) \in \mathbb{R}^+ \times M^2 \mid p = \tilde{p}\}$  denotes the diagonal submanifold. Moreover, their difference  $K_{AB} - K_A =: K \in \mathcal{A}_{\text{phg}}^{q,\infty}(\mathcal{M}_b^2)$ .*

<sup>5</sup>A distribution  $\omega$  on  $\mathbb{R}^+ \times M^2$  lifts to a distribution  $\beta^*\omega$  on  $\mathcal{M}_b^2$  by requiring  $(\beta^*\omega)(f \circ \beta) := \omega(f)$  on any test function  $f \in C_0^\infty(\mathbb{R}^+ \times M^2)$ . Note that a (compactly supported) test function on  $\mathcal{M}_b^2$  can always be written as  $f \circ \beta$  for some test function  $f$  on  $\mathbb{R}^+ \times M^2$ .

*Proof.* For  $R > 0$  sufficiently large and  $\mu \in \Gamma_R$ , we may write according to [SEE69, (28)]

$$\text{Op}(d_{-q-j}) = (2\pi)^{-m} \int_{\mathbb{R}^{m-1}} e^{i\langle y-\tilde{y}, \zeta \rangle} \tilde{d}_{-q-j}(x, y, \zeta, \tilde{x}, \mu) d\zeta, \quad (2.7)$$

where  $\tilde{d}_{-q-j}$  is homogeneous of degree  $(-q-1)$  in  $(x^{-1}, \tilde{x}^{-1}, \zeta, \mu)$ . Moreover, [SEE69, (29)] asserts the following estimate

$$\begin{aligned} \left| x^i \tilde{x}^k \partial_x^\alpha \partial_{\tilde{x}}^\beta \partial_\zeta^\gamma \partial_\mu^\delta \tilde{d}_{-q-j}(x, y, \zeta, \tilde{x}, \mu) \right| &\leq C \exp(-c(x + \tilde{x})(|\zeta| + \mu)) \\ &\times (|\zeta| + \mu)^{1-q-j-k-i+\alpha+\beta-|\gamma|-\delta}, \end{aligned} \quad (2.8)$$

with constants  $c, C > 0$ . This estimate is stable under differentiation in  $y \in \mathbb{R}^{m-1}$ . We may now study the asymptotics of the lift  $\beta^* \text{Op}(d_{-q-j})$  in the various projective coordinates near the front face of  $\mathcal{M}_b^2$ . For instance, in coordinates (2.5) we find

$$\beta^* \text{Op}(d_{-q-j}) = (2\pi)^{-m} \tilde{x}^{-m+q+j} \int_{\mathbb{R}^{m-1}} e^{i\langle u, \nu \rangle} \tilde{d}_{-q-j}(s, y, \nu, 1, \tau^{-1}) d\nu. \quad (2.9)$$

Hence the lift  $\beta^* \text{Op}(d_{-q-j})$  is of order  $(-m+q+j)$  at the front face  $\text{ff}(\tilde{x} \rightarrow 0)$ , and smooth at  $\text{rf}(s \rightarrow 0)$ . In view of the estimate (2.8) the expression is also vanishing to infinite order at the temporal face  $\text{tf}(\tau \rightarrow 0)$ . Similarly, in coordinates (2.6)

$$\beta^* \text{Op}(d_{-q-j}) = (2\pi)^{-m} (\eta \tilde{x})^{-m+q+j} \int_{\mathbb{R}^{m-1}} e^{i\langle U, \nu \rangle} \tilde{d}_{-q-j}(S + \eta^{-1}, y, \nu, \eta^{-1}, 1) d\nu. \quad (2.10)$$

Hence the lift  $\beta^* \text{Op}(d_{-q-j})$  is of order  $(-m+q+j)$  at the front face  $\text{ff}(\tilde{x} \rightarrow 0)$ . In view of the estimate (2.8) the expression is also vanishing to infinite order at the temporal face  $\text{tf}(|(S, U)| \rightarrow \infty)$  as well as temporal diagonal  $\text{td}(\eta \rightarrow 0)$ .

Summarizing we have shown  $\beta^* \text{Op}(d_{-q-j}) \in \mathcal{A}_{\text{phg}}^{q+j, \infty}(\mathcal{M}_b^2)$ . Similar arguments applied to the classical symbol expansion of the interior parametrix  $K_A = (A + \mu^q)^{-1}$  justify  $K_A \in \mathcal{A}_{\text{phg}}^{q,q}(\mathcal{M}_b^2, \beta^* Y)$ . Since  $\beta^* \mu = \rho_{\text{ff}} \rho_{\text{td}} \rho_{\text{tf}}$ , up to multiplication with a bounded function, we infer from Theorem 2.1

$$\beta^* K_{AB} = \beta^* K_A + \sum_{j=0}^{N-1} \beta^* \text{Op}(d_{-q-j}) + (\rho_{\text{ff}} \rho_{\text{td}} \rho_{\text{tf}})^{-m+q+N} Q_N, \quad (2.11)$$

as  $(\rho_{\text{ff}}, \rho_{\text{td}}, \rho_{\text{tf}}) \rightarrow 0$ , where  $Q_N$  denotes some conormal distribution on  $\mathcal{M}_b^2$  which is bounded uniformly in  $N$  at all the boundary faces. Taking the limit  $N \rightarrow \infty$  proves the statement.  $\square$

### 3. COMPOSITION OF POLYHOMOGENEOUS SCHWARTZ KERNELS

Let  $X$  and  $X'$  be two compact manifolds with corners, and let  $f : X \rightarrow X'$  be a smooth map. Let  $\{H_i\}_{i \in I}$  and  $\{H'_j\}_{j \in J}$  be enumerations of the codimension one boundary faces of  $X$  and  $X'$ , respectively, and let  $\rho_i, \rho'_j$  be global defining functions for  $H_i$ , resp.  $H'_j$ . We say that the map  $f$  is a  $b$ -map if for all  $j \in J$  there exists a smooth positive function  $a_j$  such that

$$f^* \rho'_j = a_j \prod_{i \in I} \rho_i^{e(i,j)}, \quad e(i,j) \in \mathbb{N} \cup \{0\}.$$

The map  $f$  is called a  $b$ -submersion if  $f_*$  induces a surjective map between the  $b$ -tangent bundles of  $X$  and  $X'$ . The notion of  $b$ -tangent bundles has been introduced

in [MEL93, Lemma 2.5]. Assume moreover that for each  $j$  there is at most one  $i$  such that  $e(i, j) \neq 0$ . In other words no submanifold in  $X$  gets mapped to a corner in  $X'$ . Under this condition the  $b$ -submersion  $f$  is called a  $b$ -fibration.

Suppose that  $\nu_0$  is a density on  $X$  which is smooth up to all boundary faces and everywhere nonvanishing. A smooth  $b$ -density  $\nu_b$  is, by definition, any density of the form  $\nu_b = \nu_0(\Pi\rho_i)^{-1}$ . Let us fix smooth  $b$ -densities  $\nu_b$  on  $X$  and  $\nu'_b$  on  $X'$ .

**Proposition 3.1.** [MEL92, The Pushforward Theorem] *Let  $f_b : X \rightarrow X'$  be a  $b$ -fibration. Let  $u$  be a polyhomogeneous function on  $X$  with index sets  $E_i$  the faces  $H_i$  of  $X$ . Suppose that each  $(z, p) \in E_i$  has  $\operatorname{Re} z > 0$  if  $e(i, j) = 0$  for all  $j \in J$ . Then the pushforward  $f_*(u\nu_b)$  is well-defined and equals  $h\nu'_b$  where  $h$  is polyhomogeneous on  $X'$  and has an index family  $f_b(\mathcal{E})$  given by an explicit formula in terms of the index family  $\mathcal{E}$  for  $X$ .*

Rather than giving the formula for the image index set in general, we provide the index image set in a specific setup, enough for the present situation. If  $H_{i_1}$  and  $H_{i_2}$  are both mapped to a face  $H'_j$ , and if  $H_{i_1} \cap H_{i_2} = \emptyset$ , then they contribute the index set  $E_{i_1} + E_{i_2}$  to  $H'_j$ . If they do intersect, however, then the contribution is the extended union  $E_{i_1} \bar{\cup} E_{i_2}$

$$E_{i_1} \bar{\cup} E_{i_2} := E_{i_1} \cup E_{i_2} \cup \{(z, p + q + 1) : \exists (z, p) \in E_{i_1}, \text{ and } (z, q) \in E_{i_2}\}.$$

We now employ the Pushforward theorem to establish the following fundamental composition result. Let  $Y := \{(\mu, p, \tilde{p}) \in \mathbb{R}^+ \times M^2 \mid p = \tilde{p}\}$  denote the diagonal submanifold. Consider any  $K_a \in \mathcal{A}_{\text{phg}}^{\ell, k}(\mathcal{M}_b^2, \beta^*Y)$  and  $K_b \in \mathcal{A}_{\text{phg}}^{\ell', \infty}(\mathcal{M}_b^2)$ . Their composition is defined by

$$K_c(p, \tilde{p}; \mu) = \int_M K_a(p, p'; \mu) K_b(p', \tilde{p}; \mu) \operatorname{dvol}_M(p'). \quad (3.1)$$

**Proposition 3.2.**

$$\mathcal{A}_{\text{phg}}^{\ell, k}(\mathcal{M}_b^2, \beta^*Y) \circ \mathcal{A}_{\text{phg}}^{\ell', \infty}(\mathcal{M}_b^2) \subset \mathcal{A}_{\text{phg}}^{\ell+\ell', \infty}(\mathcal{M}_b^2). \quad (3.2)$$

*Proof.* Consider  $K_a \in \mathcal{A}_{\text{phg}}^{\ell, k}(\mathcal{M}_b^2, \beta^*Y)$  and  $K_b \in \mathcal{A}_{\text{phg}}^{\ell', \infty}(\mathcal{M}_b^2)$ . Their composition  $K_c = K_a \circ K_b$  is defined in (3.1). This expression can be rephrased in geometric terms. Consider the space  $\mathbb{R}_{1/\mu}^+ \times M_{(p, p', \tilde{p})}^3$ , and the three projections

$$\begin{aligned} \pi_c &: \mathbb{R}_{1/\mu}^+ \times M_{(p, p', \tilde{p})}^3 \rightarrow \mathbb{R}_{1/\mu}^+ \times M_{(p, \tilde{p})}^2, \\ \pi_a &: \mathbb{R}_{1/\mu}^+ \times M_{(p, p', \tilde{p})}^3 \rightarrow \mathbb{R}_{1/\mu}^+ \times M_{(p, p')}^2, \\ \pi_b &: \mathbb{R}_{1/\mu}^+ \times M_{(p, p', \tilde{p})}^3 \rightarrow \mathbb{R}_{1/\mu}^+ \times M_{(p', \tilde{p})}^2. \end{aligned} \quad (3.3)$$

We reinterpret  $K_a, K_b$  and  $K_c$  as ‘right densities’

$$\begin{aligned} K_a &\equiv K_a(p, p'; \mu) \operatorname{dvol}_M(p'), \\ K_b &\equiv K_b(p', \tilde{p}; \mu) \operatorname{dvol}_M(\tilde{p}), \\ K_c &\equiv K_c(p, \tilde{p}; \mu) \operatorname{dvol}_M(\tilde{p}). \end{aligned}$$

Then we can rewrite (3.1) as

$$K_c = (\pi_c)_* (\pi_a^* K_a \cdot \pi_b^* K_b).$$

The basic idea in the proof of polyhomogeneity of  $K_c$  is a construction of a triple-space  $\mathcal{M}_b^3$  which is a blowup of  $\mathbb{R}_{1/\mu}^+ \times M^3$  obtained by a sequence of blowups, designed such that there are maps

$$\Pi_a, \Pi_c, \Pi_b : \mathcal{M}_b^3 \longrightarrow [\mathbb{R}_{1/\mu}^+ \times M^2; \mathcal{C}] =: \mathcal{M}_{rb}^2$$

which ‘cover’ the three projections defined above. The construction is reminiscent of the triple space construction for the heat space calculus for conical singularities, see [Moo99], but differs from the latter since there is no convolution in the parameter  $\mu^{-1}$  variable and the blowups are not parabolic in the  $\mu^{-1}$  direction. On each copy of  $M$  we use the local coordinates  $p = (x, y), p' = (x', y'), \tilde{p} = (\tilde{x}, \tilde{y}) \in \mathbb{R}^+ \times \mathbb{R}^{m-1}$  near the boundary  $\partial M$  with  $(x, x', \tilde{x})$  being the three copies of the boundary defining function. First we blow up the submanifold

$$\begin{aligned} F &= \{(1/\mu, x, y, x', y', \tilde{x}, \tilde{y}) \mid \mu = \infty, x = x' = \tilde{x} = 0, y = y' = \tilde{y}\} \\ &= \pi_a^{-1}\mathcal{C} \cap \pi_b^{-1}\mathcal{C} \cap \pi_c^{-1}\mathcal{C}, \end{aligned}$$

which is the intersection of all highest codimension corners  $\mathcal{C}$  introduced in (2.1), pulled back under the three projections to  $\mathbb{R}^+ \times M^3$ . Then we blow up the resulting space  $[\mathbb{R}^+ \times M^3; F]$  at the interior lifts of each of the three submanifolds

$$\begin{aligned} F_c &= \pi_c^{-1}\mathcal{C} = \{(1/\mu, x, y, x', y', \tilde{x}, \tilde{y}) \mid \mu = \infty, x = \tilde{x} = 0, y = \tilde{y}\}, \\ F_a &= \pi_a^{-1}\mathcal{C} = \{(1/\mu, x, y, x', y', \tilde{x}, \tilde{y}) \mid \mu = \infty, x = x' = 0, y = y'\}, \\ F_b &= \pi_b^{-1}\mathcal{C} = \{(1/\mu, x, y, x', y', \tilde{x}, \tilde{y}) \mid \mu = \infty, x' = \tilde{x} = 0, y' = \tilde{y}\}. \end{aligned} \quad (3.4)$$

Identifying notationally each  $F_{a,b,c}$  with their lifts to  $[\mathbb{R}^+ \times M^3, F]$ , we may altogether define the triple space

$$\mathcal{M}_b^3 := [[\mathbb{R}^+ \times M^3; F]; F_a; F_b; F_c].$$

If we ignore the  $\mu^{-1}$ -direction, the spacial part of  $\mathcal{M}_b^3$  can be visualized as below.




FIGURE 2. The spacial component of the triple space  $\mathcal{M}_b^3$ .

Here, (101), (011) and (110) label the boundary faces created by blowing up  $F_c$ ,  $F_b$  and  $F_a$ , respectively. The face (111) is the front face introduced by blowing up  $F$ . We denote the defining function for the face  $(ijk)$  by  $\rho_{ijk}$ . The triple space comes with a natural blowdown map  $\beta^{(3)} : \mathcal{M}_b^3 \rightarrow \mathbb{R}^+ \times M^3$ , which as in the discussion of  $\mathcal{M}_b^2$  amounts in local coordinates to a coordinate change back to  $(x, y, x', y', \tilde{x}, \tilde{y}, \mu)$ .

Now consider the projections  $\pi_c$ ,  $\pi_a$  and  $\pi_b$  introduced in (3.3). These induce projections  $\Pi_c$ ,  $\Pi_a$  and  $\Pi_b$  from  $\mathcal{M}_b^3$  to the reduced blowup space  $\mathcal{M}_{rb}^2$ . It is not hard to check that the choice of submanifolds  $F, F_{a,b,c}$  that have been blown up ensures that these projections are in fact  $b$ -fibrations.

Denote defining functions for the right, front and left faces of each copy of  $\mathcal{M}_{rb}^2$  by  $\{\rho_{10}, \rho_{11}, \rho_{01}\}$ , respectively. These lift via the projections according to the following rules (modulo multiplication by a non-vanishing function)

$$\begin{aligned}\Pi_c^*(\rho_{ij}) &= \rho_{i0j}\rho_{i1j}, \\ \Pi_a^*(\rho_{ij}) &= \rho_{ij0}\rho_{ij1}, \\ \Pi_b^*(\rho_{ij}) &= \rho_{0ij}\rho_{1ij}.\end{aligned}\tag{3.5}$$

Now consider the behaviour in the parameter  $\mu^{-1}$ -direction. Let  $\tau$  be a defining function for the boundary face in  $\mathcal{M}_b^3$  which is mapped onto  $\{\mu = \infty\}$  by the blowdown map. Since each  $F, F_{a,b,c}$  is a submanifold of  $\{\mu = \infty\}$ , we find

$$\beta_{(3)}^*\mu^{-1} = \tau\rho_{111}\rho_{110}\rho_{101}\rho_{011}.\tag{3.6}$$

Let  $\beta_{(2)} : \mathcal{M}_{rb}^2 \rightarrow \mathbb{R}^+ \times M^2$  be the blowdown map for the reduced blowup space. Then  $\beta_{(2)}^*\mu^{-1} = T\rho_{11}$ , where  $T$  is a defining function for the temporal face  $tf$  in  $\mathcal{M}_{rb}^2$ . Note that  $\beta_{(2)} \circ \Pi_{a,b,c} = \pi_{a,b,c} \circ \beta_{(3)}$  and hence acting on functions on  $\mathbb{R}^+ \times M^2$  we have

$$\Pi_{a,b,c}^* \circ \beta_{(2)}^* = \beta_{(3)}^* \circ \pi_{a,b,c}^*.$$

Consequently, in view of Eq. (3.5) and Eq. (3.6), we conclude (modulo multiplication by a non-vanishing function)

$$\begin{aligned}\Pi_c^*(T) &= \tau\rho_{110}\rho_{011}, \\ \Pi_a^*(T) &= \tau\rho_{101}\rho_{011}, \\ \Pi_b^*(T) &= \tau\rho_{101}\rho_{110}.\end{aligned}\tag{3.7}$$

Using these data, we may now prove the anticipated composition formula. Consider the ‘right densities’,  $K_a(x, y, x', y'; \mu)dx'dy'$  and  $K_b(x', y', \tilde{x}, \tilde{y}; \mu)d\tilde{x}d\tilde{y}$ . Their product is given by

$$K_a(x, y, x', y'; \mu) \cdot K_b(x', y', \tilde{x}, \tilde{y}; \mu) dx' dy' d\tilde{x} d\tilde{y}$$

Its integral over  $dx'dy'$  gives  $K_c(x, y, \tilde{x}, \tilde{y}; \mu)d\tilde{x}d\tilde{y}$ . To put this into the same form required in the pushforward theorem, write  $t = \mu^{-1}$  and multiply this expression by  $dt dx dy$ .

Blowing up a submanifold of codimension  $n$  amounts in local coordinates to introducing polar coordinates, so that the coordinate transformation of a density leads to  $(n-1)^{\text{st}}$  power of the radial function, which is a defining function of the corresponding front face. Hence we compute the lift

$$\begin{aligned}\beta_{(3)}^*(dt dx dy dx' dy' d\tilde{x} d\tilde{y}) \\ = \rho_{111}^{3+2(m-1)} \rho_{101}^{2+(m-1)} \rho_{110}^{2+(m-1)} \rho_{011}^{2+(m-1)} \nu^{(3)} \\ = \rho_{111}^{3+2(m-1)} \rho_{101}^{2+(m-1)} \rho_{110}^{2+(m-1)} \rho_{011}^{2+(m-1)} \tau (\Pi\rho_{ijk}) \nu_b^{(3)},\end{aligned}\tag{3.8}$$

where  $\nu^{(3)}$  is a density on  $\mathcal{M}_b^3$ , smooth up to all boundary faces and everywhere nonvanishing;  $\nu_b^{(3)}$  is a  $b$ -density, obtained from  $\nu^{(3)}$  by dividing by a product of all

defining functions on  $\mathcal{M}_b^3$ ; and  $(\Pi\rho_{ijk})$  is a product over all  $(ijk) \in \{0,1\}^3$ . Set  $\kappa_a = \beta_{(2)}^* K_a$  and  $\kappa_b = \beta_{(2)}^* K_b$ . Since  $\kappa_b$  is vanishing to infinite order as  $T \rightarrow 0$ , its lift  $\Pi_b^* \kappa_b$  vanishes to infinite order in  $\tau\rho_{110}\rho_{101}$  by Eq. (3.7).

Since  $\kappa_a$  is not polyhomogeneous on the reduced blowup space  $\mathcal{M}_{rb}^2$ , the lift  $\Pi_a^* \kappa_a$  is not polyhomogeneous in  $\tau$ . However, due to infinite order vanishing of  $\Pi_b^* \kappa_b$  in  $\tau$ , their product  $\Pi_a^* \kappa_a \cdot \Pi_b^* \kappa_b$  is polyhomogeneous and vanishing to infinite order in  $\tau\rho_{110}\rho_{101}\rho_{011}$ . We obtain

$$\Pi_a^* \kappa_a \cdot \Pi_b^* \kappa_b \beta_{(3)}^* (dt dx dy dx' dy' d\tilde{x} d\tilde{y}) = \rho_{111}^{\ell+\ell'+1} (\Pi\rho_{ijk}) G \nu_b^{(3)},$$

where  $G$  is a bounded polyhomogeneous function on  $\mathcal{M}_b^3$ , vanishing to infinite order in  $(\tau\rho_{110}\rho_{101}\rho_{011})$ , with index sets  $\mathbb{N}_0$  at the faces (001), (100) and (010). Applying the Pushforward Theorem now gives

$$\begin{aligned} & (\Pi_c)_* (\Pi_a^* \kappa_a \cdot \Pi_b^* \kappa_b \beta_{(3)}^* (dt dx dy dx' dy' d\tilde{x} d\tilde{y})) \\ &= \beta_{(2)}^* (K_c dt dx dy d\tilde{x} d\tilde{y}) = \rho_{11}^{2+\ell+\ell'} G' \nu_b^{(2)}, \end{aligned} \quad (3.9)$$

where  $\nu_b^{(2)}$  is a  $b$ -density on  $\mathcal{M}_{rb}^2$  and  $G'$  is a bounded polyhomogeneous function on  $\mathcal{M}_{rb}^2$ , which vanishes to infinite order in  $T$ , and has the index set  $\mathbb{N}_0$  at the left and right boundary faces. By [EMM91, Proposition B7.20] the pushforward is smooth across  $\beta^* Y$ .

Note also that the pushforward by  $\Pi_c$  does not introduce logarithmic terms in the front face expansion of  $\kappa_c$ , since the kernel on  $\mathcal{M}_{rb}^2$  is vanishing to infinite order at (101). Hence, for  $\kappa_A$  and  $\kappa_B$  with integer exponents in their front face expansions, the same holds for their composition.

By an argument similar to Eq. (3.8), we compute

$$\beta_{(2)}^* (dt dx dy d\tilde{x} d\tilde{y}) = \rho_{11}^{m+1} (\rho_{10}\rho_{11}\rho_{01}T) \nu_b^{(2)}. \quad (3.10)$$

Consequently, combining Eq. (3.9) and Eq. (3.10), we deduce that  $\beta_{(2)}^* K_c$  vanishes to infinite order in  $T$ , is of leading order  $(-m + \ell + \ell')$  at the front face and has the index sets  $\mathbb{N}_0$  at the left and right boundary faces. This proves the statement.  $\square$

#### 4. MULTI-PARAMETER RESOLVENT TRACE EXPANSION

**4.1. Initial parametrix and Neumann series.** In this section we construct the Schwartz kernel of  $(A_B(\lambda) + z^q)^{-1}$  as a polyhomogeneous distribution in an open neighborhood of ff in  $\mathcal{M}_b^2$ . The microlocal description of the resolvent kernels in Theorem 2.3 does not apply directly to  $A_B(\lambda)$ , since the smooth potentials  $V_1, \dots, V_n$  need not be constant along  $\partial M$ . Hence we consider for each  $y_0 \in \partial M$  the operator  $A_B(\lambda)_{y_0} := A_B + \sum_{k=1}^n \lambda_k^q V_k(y_0)$ , with the corresponding resolvent for  $z \in \Gamma$  sufficiently large

$$K_{y_0} := (A_B(\lambda)_{y_0} + z^q)^{-1}(x, \tilde{x}, y, \tilde{y}). \quad (4.1)$$

The front face ff and its open  $\varepsilon$ -neighborhood in  $\mathcal{M}_b^2$  are fibrations over  $\partial M$  with fibres  $\mathbb{S}_{++}^m$  and  $[0, \varepsilon) \times \mathbb{S}_{++}^m$ , respectively. Theorem 2.3 asserts that  $K_{y_0}(\cdot, y_0)$  lifts to a polyhomogeneous conormal distribution on the  $y_0$ -fibre with  $\mu = |(\lambda, z)|^{-1}$ . Consequently we may define a distribution  $K_0 \in \mathcal{A}_{\text{phg}}^{q,q}(\mathcal{M}_b^2, \beta^* Y)$  by (e.g. in projective

coordinates (2.5))

$$K_0(\tau, s, \tilde{x}, u, y_0) := K_{y_0}(s\tilde{x}, \tilde{x}, y_0 + \tilde{x}u, y_0). \quad (4.2)$$

By construction,  $BK_0 = 0$  and  $K_0$  solves the resolvent equation<sup>6</sup>  $(A_B(\lambda) + z^q)K_0 = I + P$  up to an error term  $P$  that vanishes at ff to one order higher, more precisely is of order  $(q+1)$  at ff. As in Theorem 2.3, we may separate  $K_0$  into an interior and boundary contribution

$$K_0 = K_i + K_b, \quad K_i \in \mathcal{A}_{\text{phg}}^{q,q}(\mathcal{M}_b^2, \beta^*Y), K_b \in \mathcal{A}_{\text{phg}}^{q,\infty}(\mathcal{M}_b^2). \quad (4.3)$$

Formally,  $K_0$  is corrected to an exact solution of  $(A_B(\lambda) + z^q)K = I$  by convolution with a Neumann series of the remainder  $R$

$$\begin{aligned} K &= K_0(I + P)^{-1} = K_0 \sum_{j=0}^{\infty} (-1)^j P^j \\ &= \sum_{j=0}^{\infty} (-1)^j (K_0 P^j - K_i P^j) + \sum_{j=0}^{\infty} (-1)^j K_i P^j \\ &=: \sum_{j=0}^{\infty} (-1)^j R_0^j + \sum_{j=0}^{\infty} (-1)^j R_1^j =: R_0 + R_1. \end{aligned} \quad (4.4)$$

The Neumann series converge in the operator norm for  $z \in \Gamma$  sufficiently large. Note that the second term  $R_1$  is an interior parametrix  $(A(\lambda) + z^q)^{-1}$ . Its multi-parameter expansion follows from the classical calculus of pseudo-differential operators with parameter, for a survey type exposition see for example [LES10, Sec. 4 and 5].

More precisely, the differential expression  $(A(\lambda) + z^q)$  is of order  $q$ , elliptic in the parametric sense with parameter  $(\lambda, z) \in \Gamma^{n+1}$ . We write  $(A(\lambda) + z^q) \in \text{CL}^q(M; \Gamma^{n+1})$ . By [SHU01, Sec. II.9], its parametrix  $R_1 \in \text{CL}^{-q}(M; \Gamma^{n+1})$ , and the  $N$ -th power  $R_1^N \in \text{CL}^{-qN}(M; \Gamma^{n+1})$ . Consider  $N \in \mathbb{N}$ , such that  $qN > m$ . Fix any multiindex  $\alpha \in \mathbb{N}_0^n$  and  $\beta \in \mathbb{N}_0$ . Then the Schwartz kernel of  $R_1^N$ , which we do not distinguish notationally from the operator, is continuous with an asymptotic expansion on the diagonal as  $|(\lambda, z)| \rightarrow \infty$ ,  $(\lambda, z) \in \Gamma^{n+1}$

$$\partial_\lambda^\alpha \partial_z^\beta R_1^N(p, p; \lambda, z) \sim \sum_{j=0}^{\infty} f_j \left( p, \frac{(\lambda, z)}{|(\lambda, z)|} \right) |(\lambda, z)|^{-qN-j-|\alpha|-\beta+m}, \quad (4.5)$$

see [LES10, Theorem 5.1]. The functions  $f_j$  are smooth on  $M \times (\Gamma^{n+1} \cap \mathbb{S}^n)$  and the expansion Eq. (4.5) is uniform over the compact manifold  $M$ .

**4.2. Trace norm estimates and expansions.** It remains to study  $R_0$ . Trace norm estimates of  $R_0^j$  depend on the following basic result.

**Proposition 4.1.** *Consider  $R \in \mathcal{A}_{\text{phg}}^{\ell,\infty}(\mathcal{M}_b^2)$ . Then  $R$  defines a Hilbert Schmidt operator with the Hilbert Schmidt norm*

$$\|R(\cdot, \mu)\|_{HS} = O(\mu^{-\ell+m/2}), \quad \mu \rightarrow \infty.$$

<sup>6</sup> $I$  denotes the identity operator.

*Proof.* We make the argument explicit in projective coordinates (2.5).  $R$  lifts to a polyhomogeneous function on  $\mathcal{M}_{rb}^2$  and

$$\begin{aligned}\beta^*(R^2 dx dy d\tilde{x} d\tilde{y}) &= \tilde{x}^{-m+2\ell} G ds d\tilde{x} du dy \\ &= \mu^{-2\ell+m} \tau^{m-2\ell} G ds d\tilde{x} du dy,\end{aligned}$$

where  $G$  is bounded in  $(\tilde{x}, s, u, y)$  and vanishing to infinite order as  $\tau \rightarrow \infty$ . Similar estimates hold at the other two corners of the front face in the reduced blowup space  $\mathcal{M}_{rb}^2$ . Consequently we obtain as  $\mu \rightarrow \infty$

$$\|R(\cdot, \mu)\|_{HS}^2 = \int_M \int_M R(p, \tilde{p}; \mu)^2 d\text{vol}_M(p) d\text{vol}_M(\tilde{p}) = O(\mu^{-2\ell+m}).$$

□

**Corollary 4.2.** *Let  $M \in \mathbb{N}$  and  $\beta \in \mathbb{N}_0$  be fixed. Write for any multiindex  $\alpha \in \mathbb{N}_0^n$   $\partial_\lambda^\alpha = \partial_{\lambda_1}^{\alpha_1} \cdots \partial_{\lambda_n}^{\alpha_n}$ . Then for  $\mu_0$  sufficiently large there exist constants  $C > 0$  and  $0 < q < 1$  such that for  $N \geq M$  and  $\mu \geq \mu_0$  the trace norms satisfy the following estimates*

$$\begin{aligned}\|\partial_\lambda^\alpha \partial_z^\beta R_0^N\|_{\text{tr}} &\leq C \cdot N \cdot q^{N-M} \cdot \mu^{-M-|\alpha|-\beta-q+\frac{m}{2}}, \\ \|\partial_\lambda^\alpha \partial_z^\beta \sum_{j=M}^{\infty} R_0^j\|_{\text{tr}} &= O(\mu^{-M-|\alpha|-\beta-q+\frac{m}{2}}), \text{ as } \mu \rightarrow \infty.\end{aligned}\tag{4.6}$$

*Proof.* It suffices to consider the case  $\alpha = 0^n$  and  $\beta = 0$ . For elements  $a$  and  $b$  in a not necessarily commutative ring we have by induction the following identity

$$(a+b)^N - b^N = \sum_{k=0}^{N-1} b^k a (a+b)^{N-k-1}.\tag{4.7}$$

By construction,  $P = \lambda(V)K_0$ , where in local coordinates  $(x, y) \in \mathcal{U}$  in an open collar neighborhood of the boundary  $\lambda(V) = \sum_{k=1}^n \lambda_k^q (V_k(x, y) - V_k(0, y))$ . Consequently, we may write

$$\begin{aligned}R_0^N &= K_0 \circ (\lambda(V)K_0)^N - K_i \circ (\lambda(V)K_i)^N \\ &= \sum_{k=0}^{N-1} K_i \circ (\lambda(V)K_i)^k \circ K_b \lambda(V) \circ (\lambda(V)K_0)^{N-k-1} \\ &\quad + K_b \circ (\lambda(V)K_0)^N.\end{aligned}\tag{4.8}$$

Each of the  $N$  summands of  $R_0^N$  is of the form

$$P_N = Q_0 \circ \left( \prod_{j=1}^N \lambda(V)Q_j \right),\tag{4.9}$$

where each  $Q_j, j = 0, \dots, N$ , is either  $K_0, K_i$  or  $K_b$ . By construction, at least one of the kernels equals  $K_b$ .

By the classical resolvent decay, we have for  $z \in \Gamma$  sufficiently large and for each  $j$  an estimate of the operator norm

$$\|\lambda(V)Q_j\| \leq c < 1.\tag{4.10}$$

Thus for any fixed  $M \leq N$  we may estimate the trace norm of each summand  $P_N$  by

$$\|P_N\|_{\text{tr}} \leq c^{N-M} \cdot \|Q_0 \left( \prod_{j=1}^M \lambda(V) Q_j \right)\|_{\text{HS}}, \quad (4.11)$$

where  $\|\cdot\|_{\text{tr}}, \|\cdot\|_{\text{HS}}$  denote the trace norm resp. the Hilbert-Schmidt norm. Note that for  $\delta V_k(x, y) := V_k(x, y) - V_k(0, y)$  the lift  $\beta^* \delta V_k$  vanishes to first order at ff and hence writing for any  $\alpha \in \mathbb{N}_0^n$   $(\lambda^q)^\alpha := \lambda_1^{q\alpha_1} \cdots \lambda_n^{q\alpha_n}$  and  $|\alpha| := \alpha_1 + \cdots + \alpha_n$  we find

$$Q_0 \left( \prod_{j=1}^M \lambda(V) Q_j \right) \in \bigoplus_{|\alpha|=M} (\lambda^q)^\alpha \mathcal{A}_{\text{phg}}^{M(q+1)+q, \infty}(\mathcal{M}_b^2). \quad (4.12)$$

By Proposition 4.1 we find

$$\|Q_0 \left( \prod_{j=1}^M \lambda(V) Q_j \right)\|_{\text{HS}} \leq C_M \mu^{-M-q+m/2}. \quad (4.13)$$

This proves the first and subsequently the second statements.  $\square$

We may now separate

$$R_0 = \sum_{j=0}^{\infty} (-1)^j R_0^j = \sum_{j=0}^{M-1} (-1)^j R_0^j + \sum_{j=M}^{\infty} (-1)^j R_0^j,$$

and in view of Corollary 4.2, an asymptotic expansion of  $\text{Tr} R_0$  follows by studying the first sum and then taking  $M \rightarrow \infty$  to improve the remainder estimate. Asymptotics of the first sum follows from the next proposition.

**Proposition 4.3.** *For any  $R \in \mathcal{A}_{\text{phg}}^{\ell, \infty}(\mathcal{M}_b^2)$  we find*

$$\text{Tr} R(\cdot, \mu) \sim \sum_{j=0}^{\infty} a_j \mu^{-1-j-\ell+m}, \quad \mu \rightarrow \infty. \quad (4.14)$$

*Proof.* Consider  $Y := \{(\mu, p, \tilde{p}) \in \mathbb{R}^+ \times M^2 \mid p = \tilde{p}\}$ . The lift  $\beta^* R$  does not have a conormal singularity and restricts to a polyhomogeneous distribution on  $\beta^* Y \subset \mathcal{M}_b^2$ , which itself is a blowup of  $\mathbb{R}_{1/\mu}^+ \times M$  at the highest codimension corner, with the blowdown map denoted by  $\beta_Y$ . We refer to the restrictions of ff, lf and td in  $\mathcal{M}_b^2$  to  $\beta^* Y$  as the front face, left face and temporal diagonal again.

The restriction of  $\beta^* R$  to  $\beta^* Y$  is polyhomogeneous of leading order  $(-m+\ell)$  at the front face, index set  $\mathbb{N}_0$  at the left face and vanishes to infinite order at the temporal diagonal. Consider the obvious projection  $\pi : \mathbb{R}^+ \times M \rightarrow \mathbb{R}^+$ . Then  $(t = 1/\mu)$

$$\text{Tr} R(\cdot, \mu) dt = (\pi \circ \beta_Y)_* (\beta_Y^* R|_Y dt d\text{vol}_M).$$

Note that

$$\beta_Y^* (R|_Y dt d\text{vol}_M) = \rho_{\text{ff}}^{-m+\ell+2} G \nu_b.$$

where  $\nu_b$  is a  $b$ -density on  $\beta^* Y$ ,  $\rho_{\text{ff}}$  is the defining function of the front face,  $G$  is a bounded polyhomogeneous distribution with index set  $\mathbb{N}_0$  at the left face and

vanishes to infinite order at the temporal diagonal in  $\beta^*Y$ . By the Theorem 3.1 we find as  $t \rightarrow 0$

$$(\pi \circ \beta_Y)_*(\beta_Y^* R|_Y dt \, \text{dvol}_M) \sim \sum_{j=0}^{\infty} t^{-m+\ell+2+j} (t^{-1} dt).$$

This proves the statement.  $\square$

**Corollary 4.4.** *Consider any multiindex  $\alpha \in \mathbb{N}_0^n$  and  $\beta \in \mathbb{N}_0$ . For each  $i \in \mathbb{N}_0$  there exist  $h_i \in C^\infty(\Gamma^{n+1} \setminus \{0\})$ , homogeneous in  $(\lambda, z)$  of order  $(-1 - q + m - i)$ , such that in the notation of Corollary 4.2*

$$\partial_\lambda^\alpha \partial_z^\beta \text{Tr} R_0 \sim \sum_{i=0}^{\infty} h_{i+|\alpha|+\beta}(\lambda, z), \text{ as } \mu \rightarrow \infty.$$

*Proof.* As before in Corollary 4.2 it suffices to consider  $\alpha = 0^n, \beta = 0$ . As observed in (4.12), we find by construction

$$R_0^j = \sum_{|\alpha|=j}^n Q_{j,\alpha} \in \bigoplus_{|\alpha|=M} (\lambda^q)^\alpha \mathcal{A}_{\text{phg}}^{j(q+1)+q,\infty}(\mathcal{M}_b^2).$$

For each  $Q_{j,\alpha} \in (\lambda^q)^\alpha \mathcal{A}_{\text{phg}}^{j(q+1)+q,\infty}(\mathcal{M}_b^2)$  we find by Proposition 4.3 as  $\mu \rightarrow \infty$

$$\text{Tr} Q_{j,\alpha}(\mu) \sim \sum_{i=0}^{\infty} b_i \left( \frac{(\lambda^q)^\alpha}{\mu^{q|\alpha|}} \right) \mu^{-1-i-j-q+m}.$$

Altogether we obtain the following polyhomogeneous multi-parameter trace expansion for any  $M \in \mathbb{N}$

$$\text{Tr} \sum_{j=0}^{M-1} (-1)^j R_0^j \sim \sum_{i=0}^{\infty} h'_i(\lambda, z), \text{ as } \mu \rightarrow \infty,$$

where each  $h'_i(\lambda, z)$  is homogeneous in  $(\lambda, z)$  of order  $(-1 - q + m - i)$ . Taking  $M \rightarrow \infty$  we derive by Corollary 4.2 the stated full polyhomogeneous expansion.  $\square$

Together with the asymptotics of the interior contribution (4.5) this leads to our final main result.

**Theorem 4.5.** *Consider any multiindex  $\alpha \in \mathbb{N}_0^n$  and  $\beta \in \mathbb{N}_0$ . Fix  $N \in \mathbb{N}$  such that  $qN > m$ . Then there exist  $e_i \in C^\infty(M \times (\Gamma^{n+1} \cap \mathbb{S}^n))$ , such that*

$$\partial_\lambda^\alpha \partial_z^\beta \text{Tr}(A_B(\lambda) + z^q)^{-N} \sim \sum_{j=0}^{\infty} e_j \left( \frac{(\lambda, z)}{|(\lambda, z)|} \right) |(\lambda, z)|^{-1-qN-j-|\alpha|-\beta+m},$$

where  $e_0$  is purely a boundary contribution.

*Proof.* The statement follows from Corollary 4.4, Eq. (4.5) and

$$\begin{aligned} \partial_\lambda^\alpha \partial_z^\beta (A_B(\lambda) + z^q)^{-N} &= \partial_\lambda^\alpha \partial_z^\beta \left( -\frac{1}{q} z^{1-q} \partial_z \right)^{N-1} (A_B(\lambda) + z^q)^{-1} \\ &= \partial_\lambda^\alpha \partial_z^\beta \left( -\frac{1}{q} z^{1-q} \partial_z \right)^{N-1} (R_0 + R_1). \end{aligned} \tag{4.15}$$

$\square$

## ACKNOWLEDGEMENTS

The interest in multi-parameter resolvent trace expansions arose in a joint project with Matthias Lesch, whom the author thanks for various central discussions. The author also gratefully acknowledges helpful discussions with Benedikt Sauer, Leonid Friedlander and Rafe Mazzeo. The author is also indebted to the anonymous referee for the very detailed reading of the manuscript. The author was supported by the Hausdorff Center for Mathematics.

## REFERENCES

- [EMM91] C. L. EPSTEIN, R. B. MELROSE, and G. A. MENDOZA, *Resolvent of the Laplacian on strictly pseudoconvex domains*, Acta Math. **167** (1991), no. 1-2, 1–106. MR 1111745 (92i:32016)
- [LEVE13] M. LESCH AND B. VERTMAN *Regularized sum of zeta-determinants*, arXiv:1306.0780 [math.SP] (2013), accepted, to appear in Math. Annalen (2014)
- [LES10] M. LESCH, *Pseudodifferential operators and regularized traces*, Motives, quantum field theory, and pseudodifferential operators, Clay Math. Proc., vol. 12, Amer. Math. Soc., Providence, RI, 2010, pp. 37–72. arXiv:0901.1689 [math.OA]
- [MEL92] R. MELROSE *Calculus of conormal distributions on manifolds with corners* Intl. Math. Research Notices, No. 3 (1992), 51-61.
- [MEL93] R. MELROSE *The Atiyah-Patodi-Singer index theorem* Research Notes in Math., Vol. 4, A K Peters, Massachusetts (1993)
- [MOO99] E. A. MOOERS, *Heat kernel asymptotics on manifolds with conic singularities*, J. Anal. Math. **78** (1999), 1–36. MR 1714065 (2000g:58039)
- [SEE69] R. SEELEY *The resolvent of an elliptic boundary problem*, Amer. J. Math., No. 4 (1969), 889-920.
- [SHU01] M. A. SHUBIN, *Pseudodifferential operators and spectral theory*, second ed., Springer-Verlag, Berlin, 2001, Translated from the 1978 Russian original by Stig I. Andersson. MR 1852334 (2002d:47073)
- [VE15] B. VERTMAN *Regularized limit of determinants on discrete tori*, preprint arXiv:1502.04541 [math.SP], (2015)

UNIVERSITY MÜNSTER, EINSTEINSTRASSE 62, 48149 MÜNSTER, GERMANY  
*E-mail address:* vertman@uni-muenster.de