

A Bayesian Nonparametric Meta-Analysis Model¹

George Karabatsos², Elizabeth Talbott³, & Stephen G. Walker⁴

In a meta-analysis, it is important to specify a model that adequately describes the effect-size distribution of the underlying population of studies. The conventional normal fixed-effect and normal random-effects models assume a normal effect-size population distribution, conditionally on parameters and covariates. For estimating the mean overall effect size, such models may be adequate, but for prediction they surely are not if the effect size distribution exhibits non-normal behavior. To address this issue, we propose a Bayesian nonparametric meta-analysis model, which can describe a wider range of effect-size distributions, including unimodal symmetric distributions, as well as skewed and more multimodal distributions. We demonstrate our model through the analysis of real meta-analytic data arising from behavioral-genetic research. We compare the predictive performance of the Bayesian nonparametric model against various conventional and more modern normal fixed-effects and random-effects models.

Keywords: Meta-analysis; Bayesian nonparametric regression; meta-regression; effect-sizes; publication bias.

Short title: Bayesian Nonparametric Meta-Analysis.

¹This research is supported by grant SES-1156372 from the National Science Foundation, program in Methodology, Measurement, and Statistics.

²Corresponding author, Professor George Karabatsos, Department of Educational Psychology, Program in Measurement, Evaluation Statistics, and Assessments, College of Education, University of Illinois-Chicago, 1040 W. Harrison St. (MC 147), Chicago, IL 60607. E-Mail: georgek@uic.edu, gkarabatsos1@gmail.com.

³Associate Professor and Chair, Department of Special Education, College of Education, University of Illinois-Chicago. E-mail: etalbott@uic.edu.

⁴Professor of Statistics, The University of Texas at Austin, Division of Statistics and Scientific Computation. E-mail: s.g.walker@math.utexas.edu

1 Introduction

A research synthesis aims to integrate results from empirical research so as to produce generalizations (Cooper & Hedges, 2009). Meta-analysis, also referred to as the analysis of analyses (Glass, 1976), provides a quantitative synthesis of statistics that are reported by multiple research studies. Specifically, each study reports an effect-size statistic, and provides information about its sampling variance, while the features of the study may be described by one or more covariates. Typical examples of effect-size statistics include the unbiased standardized mean difference between two independent groups (Hedges, 1981), among others (Konstantopoulos, 2007). Given a sample of effect-size data, the primary aim of a meta-analysis is to infer the overall effect-size distribution from the given study population, as well as to infer the heterogeneity of the effect sizes. Conventional summaries of the overall effect-size distribution include the mean, which is often referred to as the "overall effect-size," and the variance which describes the heterogeneity of reported effect sizes in the overall effect-size distribution. Heterogeneity can be further investigated in a meta-regression analysis, in order to investigate how the mean effect-size relates to key study-level covariates (e.g., Berkey et al., 1995; Thompson & Sharp, 1999; Thompson & Higgins, 2002; Higgins & Thompson, 2004). Also, meta-regression analysis can be used to investigate and test for publication bias in the data, by relating effect-sizes with their standard errors (precisions; the square-root of the effect-size sampling variances) (Thompson & Sharp, 1999). This actually provides a regression analysis for the funnel plot (Egger et al., 1997).

The normal fixed-effects model and the normal random-effects model provide two traditional and alternative approaches to meta-analysis (Hedges & Vevea, 1998; Konstantopoulos, 2007; Borenstein et al. 2010). Each model is a weighted linear regression model, which treats study-reported effect-size as the dependent variable, weighs each reported effect-size by the inverse of its sampling variance, assumes normally-distributed regression errors, and represents the overall (mean) effect size by the intercept parameter. In other words, each of these models assume that the effect-size distribution is a unimodal and symmetric, normal distribution, conditionally on all model parameters, and conditionally on any set of chosen values of covariates in the model. The fixed-effects model is an ordinary (weighted) linear regression model. The normal random-effects model is a two-level model that extends the fixed-effects model, by allowing for between-study variance in the effect-sizes, through the addition of random intercept parameters that are assumed to have a normal distribution over the given study population. Specifically, effect-sizes at the first level are nested within studies at the second level. A three-level meta-analysis model can accommodate data structures where, for example, studies (level 2) are themselves nested within units, such as school districts (level 3) (for more details, see Konstantopoulos, 2011). Given a sample set of meta-analytic data, the parameters of a normal fixed-effects or normal random-effects model can be estimated via maximum-likelihood. Alternatively, a Bayesian inference approach can be taken, which involves the specification of a prior distribution on all parameters of the given meta-analytic model (e.g., Higgins et al., 2009). Then, Bayesian inference of the data is based on the posterior distribution of the parameters, formed by combining the data (likelihood) information of the model, with the information from the prior distribution.

Our motivation for the current paper involves the meta-analysis of 71 effect sizes, which are heritability estimates that were reported by a collection of behavioral-genetic studies of

the heritability of antisocial behavior (Talbot et al., 2012). Figure 1 presents a simple kernel probability density estimate of the effect sizes. This estimate, which is skewed and has at least two modes, clearly exhibits non-normality in the effect size distribution. According to the standard Anderson-Darling test of normality (Anderson & Darling, 1952), the data rejects the null hypothesis that the effect sizes are non-normal, at a .05 level of significance.

— Figure 1 —

Arguably, a normal model (fixed-effects or random-effects) can provide an adequate analysis of these data, specifically for the purposes of estimating the overall mean of the effect size, and possibly also the overall variance. However, such a model may not be adequate for predictive purposes, given the lack of normality in the data. The importance of predicting from meta-analytic data has been detailed in Higgins et al. (2009). As they state: "Predictions are one of the most important outcomes of a meta-analysis, since the purpose of reviewing research is generally to put knowledge gained into future application. Predictions also offer a convenient format for expressing the full uncertainty around inferences, since both magnitude and consistency of effects may be considered." Accurate prediction with non-normal data requires flexible models that support a wider range of distributions, beyond the normal distribution. This has already been recognized by Burr and Doss (2005) and Branscum and Hanson (2008), who proposed Bayesian nonparametric models for meta-analysis. More generally speaking, Bayesian nonparametric models are "nonparametric" in the sense that they avoid the more restrictive assumptions of "parametric" models, namely, that the data distribution can be fully-described by a few finite number of parameters. For example, a normal model assumes that the data distribution is unimodal and symmetric, and therefore can be entirely described by a mean parameter and a variance parameter. A Bayesian nonparametric model assigns a prior distribution to an infinite number (or a very large number) of model parameters. This is done for the purposes of defining a very flexible model that describes a wide range of data distributions, including unimodal distributions, and more multimodal distributions (Müller & Quintana, 2004).

Consequently, we propose a Bayesian nonparametric model for meta-analysis, which is more flexible than the normal, fixed-effects and random-effects models. Our model is a special case of the general model introduced by Karabatsos and Walker (2012), which was studied in more general regression settings. The new model specifies the effect-size distribution by an infinite random-intercept mixture of normal distributions, conditional on any covariate(s) of interest, with covariate-dependent mixture weights. Therefore, the model is flexible enough to describe a very wide range of effect-size distributions, including all normal distributions, as well as all (smooth) distributions that are more skewed and/or multimodal. Also, the model avoids the empirically-falsifiable assumption that effect-sizes arise strictly from symmetric unimodal distributions, such as normal distributions. Furthermore, the model's high flexibility encourages a rich and graphical inference of the whole effect-size distribution, as previously recommended for meta-analytic practice (Higgins et al., 2009).

Also, in the spirit of meta-regression analysis, our Bayesian nonparametric meta-analysis model allows the whole effect-size distribution to change flexibly and non-linearly as a function of key study-level covariates. This feature permits a rich and flexible meta-regression analysis, whereas the previous Bayesian nonparametric regression models do not account for

covariate information (see Burr & Doss, 2005; Branscum & Hanson, 2008). Moreover, for a given meta-analysis, our Bayesian model can automatically identify the subset of covariates that significantly predict changes in the mean effect-size, in a model-based and non-ad-hoc fashion. Specifically, the model makes use of spike-and-slab priors for the regression coefficients that allow for automatic covariate (predictor) selection in the posterior distribution. Such priors were developed for Bayesian normal linear regression models (George & McCulloch, 1997). Moreover, under either a normal fixed-effects or normal random-effects model inferred under a non-Bayesian (Frequentist) framework of maximum-likelihood estimation, the identification or selection of significant study-level covariates (predictors) is challenging because it deals with the standard issues of multiple hypothesis testing over predictors (e.g., Thompson & Higgins, 2002). The often-used stepwise procedures of covariate selection are known to be ad-hoc and sub-optimal.

We now describe the layout of the rest of the paper. Since the paper covers various key statistical concepts, it is necessary to first give them a brief review in Section 2. In Section 2.1, we review the basic data framework of meta-analysis, including effect sizes. In Section 2.2 we review of the conventional normal fixed-effects and normal random-effects models, and in Section 2.3 we briefly review the Bayesian statistical inference framework. In Section 2.4 we review the traditional Bayesian meta-analytic models, including conventional and more modern versions of normal fixed-effects and normal random-effects models. In Section 3, we describe our new Bayesian meta-analysis model. In Section 4, we review a standard criterion for comparing the predictive performance between different Bayesian models that are fit to a common data set, for the purposes of identifying the single model that has best predictive-fit, i.e., of identifying the single model that best describes the underlying population distribution of the sample data. In Section 5, illustrates our Bayesian meta-analytic model through the analysis of the large meta-analytic data set of behavioral genetic studies, which was briefly described above, and which involves 24 covariates. In that section, we also compare the predictive accuracy of our Bayesian nonparametric model, against the predictive accuracy of conventional and more modern normal fixed-effects and normal random-effects models. Section 6 ends with conclusions.

2 Review of Meta-Analytic Modeling Concepts

Before we review the key concepts underlying the various approaches to meta-analysis, we describe some notation that we use in the remainder of this paper. Following the standard notation of statistics, \sim will mean "distributed as"; $n(\cdot|\mu, v)$ denotes the ("bell-shaped") probability density function (p.d.f.) of the normal distribution having mean and variance (μ, v) ; the p.d.f. of the n -variate normal distribution with mean vector $\boldsymbol{\mu}$ and (symmetric and positive-definite) variance-covariance matrix $\boldsymbol{\Sigma}$ is denoted by $n_n(\cdot|\boldsymbol{\mu}, \boldsymbol{\Sigma})$; the p.d.f. of a gamma distribution with shape and rate parameters (a, b) is denoted by $ga(\cdot|a, b)$; and the the p.d.f. of a uniform distribution with minimum and maximum parameters (a, b) is denoted by $un(\cdot|a, b)$. Also, we denote a cumulative distribution function (c.d.f.) by a capital letter, such as G . Finally, $\delta_\theta(\cdot)$ denotes the degenerate distribution that assigns probability 1 (full support) to the number θ .

2.1 Data Framework of Meta-Analysis

In a typical meta-analysis context, data are available on n study reports, indexed by $i = 1, \dots, n$. Each study reports an effect-size y_i of interest, based on n_i observations, and provides information on the sampling variance of the effect size. Also, each study provides information about p study characteristics, which are described by p covariates $\mathbf{x}_i = (1, x_{1i}, \dots, x_{pi})^\top$, in addition to a constant (1) term for future notational convenience. A full meta-analytic data set is denoted by $\mathcal{D}_n = \{(y_i, \mathbf{x}_i, \hat{\sigma}_i^2)\}_{i=1}^n$.

Effect-size Description	Effect-size (y_i)	Variance ($\hat{\sigma}_i^2$)
Unbiased standardized mean difference, two independent groups (Hedges, 1981).	$\frac{\hat{\mu}_{1i} - \hat{\mu}_{2i}}{\sqrt{\frac{(n_{1i}-1)\hat{\sigma}_{1i} + (n_{2i}-1)\hat{\sigma}_{2i}}{n_{1i} + n_{2i} - 2}}} c^*$	$\left(\frac{n_{1i} + n_{2i}}{n_{1i}n_{2i}} + \frac{y_i^2}{2(n_{1i} + n_{2i})}\right) c^*$; $c^* = 1 - \frac{3}{4(n_{1i} + n_{2i} - 2) - 1}$
Fisher z transformation of the correlation $\hat{\rho}_i$.	$\frac{1}{2} \log \frac{1 + \hat{\rho}_i}{1 - \hat{\rho}_i}$	$\frac{1}{n_i + 3}$
Log odds ratio for two binary (0-1) variables.	$\log \left(\frac{n_{11i}/n_{10i}}{n_{01i}/n_{00i}} \right)$	$\frac{1}{n_{11i}} + \frac{1}{n_{10i}} + \frac{1}{n_{01i}} + \frac{1}{n_{00i}}$

Table 1: Examples of effect size statistics, along with corresponding variances.

Table 1 presents some typical examples of effect-size statistics that are often used in meta-analysis, along with their sampling variances ($\hat{\sigma}_i^2$) (Konstantopoulos, 2007; Borenstein, 2009, Fleiss & Berlin, 2009). More generally, we may consider a sampling covariance matrix for the n study reports, $\hat{\Sigma}_n = (\hat{\sigma}_{il})_{n \times n}$, having diagonal elements $\hat{\sigma}_{ii} = \hat{\sigma}_i^2$, where each off diagonal element is the sampling covariance for a given effect-size pair (y_k, y_l) , with $k \neq l$ (Gleser & Olkin, 2009). To maintain notational simplicity throughout the paper, we will present the meta-analytic models under the common assumption that $\hat{\Sigma}_n$ is a diagonal matrix (i.e., $\hat{\Sigma}_n = \text{diag}(\sigma_1^2, \dots, \sigma_n^2)$), implying the assumption of zero sampling covariances. Though, as we discuss in Section 3, this diagonal matrix assumption can be made for the Bayesian nonparametric meta-analytic model, without loss of generality in terms of being able to model covariances between distinct pairs of study effect size reports.

2.2 Traditional Meta-Analytic Models

A traditional meta-analysis model assumes that, for a given set of data $\mathcal{D}_n = \{(y_i, \mathbf{x}_i, \hat{\sigma}_i^2)\}_{i=1}^n$, the effect-size distribution follows the general form:

$$f(y_i | \mathbf{x}_i, \hat{\sigma}_i^2; \zeta) = n(y_i | \mathbf{x}_i^\top \boldsymbol{\beta} + \mu_{0i} + \mu_{00t(i)}, \hat{\sigma}_i^2), \quad i = 1, \dots, n; \quad (1a)$$

$$\mathbf{x}_i^\top \boldsymbol{\beta} = \beta_0 + \beta_1 x_{i1} + \dots + \beta_p x_{ip}; \quad (1b)$$

$$(\mu_{01}, \dots, \mu_{0n}) | \Sigma_0 \sim n_n(\mathbf{0}, \sigma_0^2 \mathbf{I}_n + \psi \mathbf{M}_n); \quad (1c)$$

$$\mu_{00t} | \sigma_{00}^2 \sim n(0, \sigma_{00}^2), \quad t = 1, \dots, T. \quad (1d)$$

Typical meta-analytic models are special cases of the general normal model shown in equation (1). In all, the general traditional model (1) has likelihood density $f(y | \mathbf{x}, \hat{\sigma}^2; \zeta)$ with

parameters $\boldsymbol{\zeta} = (\boldsymbol{\beta}, \boldsymbol{\mu}_0, \boldsymbol{\mu}_{00}, \sigma_0^2, \sigma_{00}^2, \psi)^\top$. These parameters are explained as follows.

The intercept parameter β_0 is interpreted as the *mean effect-size* over the given population of studies (Louis & Zelterman, 1994). This interpretation holds true, provided that each of the p covariates has data observations $(x_{k1}, \dots, x_{kn})^\top$ that have already been centered to have mean zero, as we assume throughout. Also, the p covariates are respectively parameterized by linear slope coefficients $(\beta_1, \dots, \beta_p)$.

Also, $\boldsymbol{\mu}_0 = (\mu_{01}, \dots, \mu_{0n})^\top$ are the level-2 random intercept parameters. Similarly, $\boldsymbol{\mu}_{00} = (\mu_{001}, \dots, \mu_{00T})^\top$ are the *level-3 random intercept parameters*, with each of the n study reports being nested within exactly one of $T \leq n$ study reports ($t = 1, \dots, T$), and with $\mu_{00t(i)}$ meaning that the level-3 intercept μ_{00t} is assigned to the i th study report. As shown in the model equations (1c) and (1d), the random intercepts $\boldsymbol{\mu}_0$ and $\boldsymbol{\mu}_{00}$ are each assumed to have a multivariate normal distribution (probability density).

A normal fixed-effects model assumes that all the random intercept parameters $(\boldsymbol{\mu}_0, \boldsymbol{\mu}_{00})$ are zero. In terms of the general normal meta-analytic model (1), this corresponds to the assumption of zero variances, i.e., $\sigma_0^2 = \sigma_{00}^2 = \psi = 0$ (Hedges & Vevea, 1998). A 2-level normal random-effects model allows for non-zero random intercepts $\boldsymbol{\mu}_0$, by allowing for nonzero variances (σ_0^2, ψ) , as shown in equation (1c). Typical normal random-effects models assume that the random intercepts $\boldsymbol{\mu}_0$ are uncorrelated, with n -variate normal density $n_n(\boldsymbol{\mu}_0 | \mathbf{0}, \sigma_0^2 \mathbf{I}_n)$ (Hedges & Vevea, 1998). However, it is possible to model correlations between the level-2 random intercepts $\boldsymbol{\mu}_0$. For example, Stevens and Taylor (2009) consider a 2-level normal random-effects model, which assumes that the level-2 random intercepts have a n -variate normal density $n_n(\boldsymbol{\mu}_0 | \mathbf{0}, \boldsymbol{\Sigma}_0)$, with a more general covariance structure $\boldsymbol{\Sigma}_0 = \sigma_0^2 \mathbf{I}_n + \psi \mathbf{M}_n$. Here, the parameter ψ represents the covariance between pairs of study reports. Also, \mathbf{M}_n is a fixed $n \times n$ indicator (0-1) matrix, with 1s specified in the off-diagonal to reflect a-priori beliefs as to which pairs of the n study reports have correlated level-2 random intercepts, and with zeros specified for all the other entries of \mathbf{M}_n . A 3-level normal random-effects model allows for non-zero random intercepts $\boldsymbol{\mu}_{00}$, by allowing for a positive variance σ_{00}^2 , as shown in equation (1d) of the general normal model (Konstantopoulos, 2011).

The general normal model described in equation (1) can be written explicitly as a normal mixture of multivariate normal model:

$$f(\mathbf{y} | \mathbf{X}, \widehat{\boldsymbol{\Sigma}}_n; \boldsymbol{\beta}, \sigma_0^2, \psi, \sigma_{00}^2) = \int n_n(\mathbf{y} | \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\mu}_0 + \boldsymbol{\mu}_{00}^*, \widehat{\boldsymbol{\Sigma}}_n) dG_2(\boldsymbol{\mu}_0) dG_3(\boldsymbol{\mu}_{00}),$$

for effect-size data $\mathbf{y} = (y_1, \dots, y_n)^\top$, given the n -by- $(p+1)$ matrix \mathbf{X} of row vectors \mathbf{x}_i^\top ($i = 1, \dots, n$), $\boldsymbol{\mu}_{00}^* = (\mu_{00t(1)}, \dots, \mu_{00t(n)})^\top$, and $\widehat{\boldsymbol{\Sigma}}_n = \text{diag}(\sigma_1^2, \dots, \sigma_n^2)$. Specifically, the mixture model presented above assumes that the mixture distribution $G_2(\boldsymbol{\mu}_0)$ is a multivariate distribution with probability density $n_n(\boldsymbol{\mu}_0 | \mathbf{0}, \sigma_0^2 \mathbf{I}_n + \psi \mathbf{M}_n)$, and that $G_3(\boldsymbol{\mu}_{00})$ is a multivariate normal distribution with probability density $n_T(\boldsymbol{\mu}_{00} | \mathbf{0}, \sigma_{00}^2 \mathbf{I}_n)$.

For any of the models described in this section, full maximum likelihood methods can be used estimate the parameters $\boldsymbol{\zeta} = (\boldsymbol{\beta}, \boldsymbol{\mu}_0, \boldsymbol{\mu}_{00}, \sigma_0^2, \sigma_{00}^2, \psi)^\top$, from a given data set \mathcal{D}_n . Alternatively, the parameters of a random-effects model can also be estimated by the restricted maximum-likelihood method, which focuses estimation on the variance parameters (Harville, 1977; Raudenbush & Bryk, 2002, Ch. 3, 13-14; Stevens & Taylor, 2009). For any one of the individual models that is described in this section, the estimate of the effect-size distribution

of the underlying population is given by the density $f_n(y|\mathbf{x}_0, \hat{\sigma}^2; \hat{\zeta}) = \mathfrak{n}(y|\hat{\beta}_0, \hat{\sigma}^2 + \hat{\sigma}_0^2 + \hat{\sigma}_{00}^2)$ of the normal distribution, given a maximum-likelihood estimate $\hat{\zeta}$ obtained from a sample data set \mathcal{D}_n , and after controlling for all p covariates via the covariate specification $\mathbf{x} = \mathbf{x}_0 = (1, 0, \dots, 0)^\top$.

As an alternative to the maximum-likelihood approach, parameter estimation can be performed in a fully-Bayesian framework (e.g., Higgins et al., 2009), which is described next.

2.3 Review of Bayesian Inference

For a general meta-analytic model, let $f(y|\mathbf{x}, \hat{\sigma}_i^2; \zeta)$ denote the likelihood density of the effect-size data point y , conditionally on covariates \mathbf{x} and model parameter ζ , which has space Ω_ζ . A given meta-analytic data set $\mathcal{D}_n = \{(y_i, \mathbf{x}_i, \hat{\sigma}_i^2)\}_{i=1}^n$ has likelihood $L(\mathcal{D}_n; \zeta) = \prod_{i=1}^n f(y_i|\mathbf{x}_i, \hat{\sigma}_i^2; \zeta)$ under the model. In the Bayesian approach to statistical inference, the model parameter ζ is assigned a prior probability density $\pi(\zeta)$ on the parameter space Ω_ζ , and this density reflects pre-experimental beliefs about the plausible values of the parameter, for the meta-analytic data set \mathcal{D}_n at hand. Then according to Bayes' theorem, the data \mathcal{D}_n , via the model's likelihood $L(\mathcal{D}_n; \zeta)$, combines with the prior density $\pi(\zeta)$, to yield a posterior density for ζ , defined by:

$$\pi(\zeta|\mathcal{D}_n) = \frac{L(\mathcal{D}_n; \zeta)\pi(\zeta)}{\int_{\Omega_\zeta} L(\mathcal{D}_n; \zeta)d\Pi(\zeta)} = \frac{\prod_{i=1}^n f(y_i|\mathbf{x}_i, \hat{\sigma}_i^2; \zeta)\pi(\zeta)}{\int_{\Omega_\zeta} \prod_{i=1}^n f(y_i|\mathbf{x}_i, \hat{\sigma}_i^2; \zeta)d\Pi(\zeta)}, \quad (2)$$

where $\Pi(\zeta)$ denotes the c.d.f. of the prior density $\pi(\zeta)$. The posterior density describes the plausible values of the model parameters ζ , given prior beliefs about ζ and data \mathcal{D}_n .

The specification of the prior density $\pi(\zeta)$ is an important step in a Bayesian analysis, and the posterior $\pi(\zeta|\mathcal{D}_n)$ can be quite sensitive to the choice of prior, especially when the sample size (n) is not large. Also, often in practice, there is a lack of prior information about the parameters ζ of the given model. This lack of prior information is reflected by a "diffuse" prior probability density $\pi(\zeta)$ that has high variance, and assigns rather-equal but broad support over the parameter space Ω_ζ . Such priors are often referred to as "non-informative", even though technically speaking, a prior cannot be fully non-informative. As a consequence of specifying a diffuse prior $\pi(\zeta)$, the posterior density $\pi(\zeta|\mathcal{D}_n)$ becomes mostly determined by the data \mathcal{D}_n likelihood $L(\mathcal{D}_n; \zeta)$, relative to the prior.

Prediction is a basic function of statistical modeling. A Bayesian meta-analytic model makes predictions of Y , given a chosen \mathbf{x} , on the basis of the posterior predictive density:

$$f_n(y|\mathbf{x}, \hat{\sigma}^2) = \int f(y|\mathbf{x}, \hat{\sigma}^2; \zeta)\pi(\zeta|\mathcal{D}_n)d\zeta, \quad (3)$$

and this density has posterior predictive mean (expectation) $E_n(Y|\mathbf{x}, \hat{\sigma}^2) = \int y f_n(y|\mathbf{x}, \hat{\sigma}^2)dy$, and posterior predictive variance $\text{Var}_n(Y|\mathbf{x}, \hat{\sigma}^2) = \int \{y - E_n(Y|\mathbf{x}, \hat{\sigma}^2)\}^2 f_n(y|\mathbf{x}, \hat{\sigma}^2)dy$. The posterior predictive density $f_n(y|\mathbf{x}, \hat{\sigma}^2)$ provides an estimate of the true effect-size density (distribution) for the underlying study population, given sample data \mathcal{D}_n and covariates \mathbf{x} of interest, under squared-error loss (Aitchison, 1975).

In most Bayesian meta-analytic models, ζ is a high-dimensional parameter vector, and

then the direct evaluation of the posterior equations (2) and (3) require prohibitive high-dimensional integrations. In such situations, one may use Markov chain Monte Carlo (MCMC) sampling methods to estimate such posterior densities of the given model. Such methods include the Gibbs sampler (Gelfand & Smith, 1990), Metropolis-Hastings algorithm (Chib & Greenberg, 1995), and other sampling algorithms (see for e.g., Brooks, Gelman, Jones, & Meng, 2011).

2.4 Review of Bayesian Normal Meta-Analytic Models

In relation to the Bayesian inference framework described in Section 2.3, consider the general normal meta-analytic model (1), which has likelihood density given by (1a), and which has parameters $\boldsymbol{\zeta} = (\boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\mu}_0, \boldsymbol{\mu}_{00}, \sigma_0^2, \sigma_{00}^2, \psi)^\top$. Here, $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_p)^\top$ are included as parameters which respectively indicate (0-1) whether or not the p covariates are included ($\gamma = 1$) or excluded ($\gamma = 0$) from the model. In typical practice involving such a model (including special cases), the prior has the general form:

$$\pi(\boldsymbol{\zeta}) = n(\boldsymbol{\beta}|\mathbf{0}, \mathbf{V}_\gamma)\pi(\boldsymbol{\gamma})n_n(\boldsymbol{\mu}_0|\mathbf{0}, \sigma_0^2\mathbf{I}_n + \psi\mathbf{M}_n)n_T(\boldsymbol{\mu}_{00}|0, \sigma_{00}^2\mathbf{I}_T)\pi(\sigma_0^2, \psi)\pi(\sigma_{00}^2). \quad (4)$$

In Bayesian meta-analytic modeling, it is common practice to specify a diffuse normal prior density for the coefficients $\boldsymbol{\beta}$ by taking $\mathbf{V}_\gamma = v\mathbf{I}_{p+1}$, with $v \rightarrow \infty$ (e.g., DuMouchel & Normand, 2000), with the implicit assumption that $\pi(\boldsymbol{\gamma} = \mathbf{1}) = 1$.

Also, it is common in the practice of Bayesian random-effects modeling to attempt to assign a non-informative prior for $(\sigma_0^2, \sigma_{00}^2)$ via the specification of inverse-gamma priors $\pi(\sigma_0^2) = \text{ga}(\sigma_0^{-2}|\epsilon, \epsilon)$ and $\pi(\sigma_{00}^2) = \text{ga}(\sigma_{00}^{-2}|\epsilon, \epsilon)$, for small choice of constant $\epsilon > 0$ (Gelman, 2006), implying a prior density of the form $\pi(\sigma_0^2, \psi) = \pi(\sigma_0^2)\delta_0(\psi)$. Though, recall from Section 2.2 that a more general multivariate normal $n_n(\boldsymbol{\mu}_0|\mathbf{0}, \sigma_0^2\mathbf{I}_n + \psi\mathbf{M}_n)$ mixture distribution can be specified (Stevens & Taylor, 2009). This mixture distribution prior allows for correlated level-2 random intercepts $\boldsymbol{\mu}_0$ via a parameter ψ that measures the covariance between specific pairs of the total n study reports, and where $\mathbf{M}_n = (m_{il})_{n \times n}$ is a fixed matrix which indicates (0-1) which pairs of the study reports are expected to yield correlated level-2 random intercepts $\boldsymbol{\mu}_0$, with zeros in the diagonal. For the parameters (σ_0^2, ψ) , Stevens and Taylor (2009) propose the rather non-informative prior density

$$\pi(\sigma_0^2, \psi) = (c_0/(c_0 + \sigma_0^2)^2)\text{un}(\psi | -\sigma_0^2/(K-1), \sigma_0^2), \quad (5)$$

where the first term in the product gives a log-logistic prior density for σ_0^2 , where $c_0 = \{n/\text{tr}([\text{diag}(\widehat{\boldsymbol{\Sigma}}_n)]^{-1})\}^{1/2}$ is the harmonic mean of the sampling variances $\widehat{\sigma}_i^2$ ($i = 1, \dots, n$), and $K = \max_i\{\sum_{l=1}^n m_{il}\}$ is the largest group of related study reports. This log-logistic prior is right-skewed, highly-dispersed, with quartiles $(c_0/3, c_0, 3c_0)$.

Simpler versions of the general normal random-effects model (1) can be specified via appropriate straightforward modifications of the prior density (4). A Bayesian 2-level normal random-effects model which assumes $\sigma_{00}^2 = 0$ (i.e., $\boldsymbol{\mu}_{00} = \mathbf{0}$), but allows for correlated level-2 random intercepts $\boldsymbol{\mu}_0$ assigns the prior density $\pi(\sigma_{00}^2) = \delta_0(\sigma_{00}^2)$; a 2-level normal random-effects model which assumes $\sigma_{00}^2 = \psi = 0$ (i.e., $\boldsymbol{\mu}_{00} = \mathbf{0}$) and assume independent level-2 random intercepts $\boldsymbol{\mu}_0$ assigns the prior density $\pi(\sigma_{00}^2, \psi) = \delta_0(\sigma_{00}^2)\delta_0(\psi)$; and a fixed-effects

model which assumes $\sigma_0^2 = \sigma_{00}^2 = \psi = 0$ (i.e., $\boldsymbol{\mu}_0 = \mathbf{0}$ and $\boldsymbol{\mu}_{00} = \mathbf{0}$) assigns the prior density $\pi(\sigma_0^2, \sigma_{00}^2, \psi) = \delta_0(\sigma_0^2)\delta_0(\sigma_{00}^2)\delta_0(\psi)$.

Other simple modifications of the prior density (4) can be used to specify other important versions of the general normal random-effects model (1), which consider different priors for the random intercepts $(\boldsymbol{\mu}_0, \boldsymbol{\mu}_{00})$. For example, Gelman (2006) notes that posterior inference of the parameter σ_0^{-2} , under the often-used "non-informative" gamma $\text{ga}(\sigma_0^{-2}|\epsilon, \epsilon)$ prior, is very sensitive to the choice of small ϵ , especially when the data support small values of σ_0^2 . Similarly for the gamma $\text{ga}(\sigma_{00}^{-2}|\epsilon, \epsilon)$ prior for the parameter σ_{00}^2 . Therefore, in a situation where there is little prior information available about the parameters $(\sigma_0^2, \sigma_{00}^2)$, he alternatively recommends the specification of uniform prior densities $\pi(\sigma_0) = \text{un}(\sigma_0|0, b_0)$ and $\pi(\sigma_{00}) = \text{un}(\sigma_{00}|0, b_{00})$ for reasonably-large values (b_0, b_{00}) , whenever n and T are both at least 5. When more prior information is desired, say when n is less than 5, he recommends the half- t prior density of the general form $\pi(\sigma_0^2) \propto (1 + a_0^{-1}(\sigma_0/b_0)^2)^{-(a+1)/2}$, and similarly for the level-3 variance parameter σ_{00} .

Finally, while it is common practice to assume a diffuse prior for the regression coefficients $\boldsymbol{\beta} = (\beta_0, \beta_1, \dots, \beta_p)^\top$, along with $\pi(\boldsymbol{\gamma} = \mathbf{1}) = 1$, in principle one may specify spike-and-slab priors for the slope parameters $(\beta_1, \dots, \beta_p)$ in order to enable automatic variable (covariate) selection via posterior inference (George & McCulloch, 1997), along with a diffuse prior $\beta_0 \sim n(0, v \rightarrow \infty)$. These spike-and-slab priors are defined by independent normal and Bernoulli prior densities, so that the $n(\boldsymbol{\beta}|\mathbf{0}, \mathbf{V}(\boldsymbol{\gamma}))\pi(\boldsymbol{\gamma})$ prior in (4) is based on:

$$\mathbf{V}_\gamma = \text{diag}(v \rightarrow \infty, v_0(1 - \gamma_1) + v_1\gamma_1, \dots, v_0(1 - \gamma_p) + v_1\gamma_p) \quad (6a)$$

$$\pi(\boldsymbol{\gamma}) = \prod_{k=1}^p \text{Pr}(\gamma_k = 1)^{\gamma_k} [1 - \text{Pr}(\gamma_k = 1)]^{1-\gamma_k}, \quad (6b)$$

where v_0 is a small prior variance (e.g., $v_0 = .001$), v_1 is a large prior variance (e.g., $v_1 = 10$), and $\text{Pr}(\gamma_k = 1)$ is the Bernoulli probability parameter that is often set to .5 in practice (George & McCulloch, 1997). So on the one hand, with prior probability $\text{Pr}[\gamma_k = 1] = .5$, the k th covariate is included in the model as a "significant" predictor of the effect-size, by assigning its regression coefficient β_k a normal $n(\beta_k|0, v_1)$ prior density that supports a large range of β_k values. On the other hand, with prior probability $\text{Pr}(\gamma_k = 0) = .5$, that covariate is excluded from the model, by assigning its regression coefficient β_k a normal $n(\beta_k|0, v_0)$ prior that places all its support on values $\beta_k \approx 0$. Given MCMC samples from the posterior, $\pi(\boldsymbol{\zeta}|\mathcal{D}_n)$, the k th covariate can be viewed as a "significant predictor," when the posterior inclusion probability of the covariate, $\text{Pr}[\gamma_k = 1|\mathcal{D}_n]$, is at least .5 (Barbieri & Berger, 2004). We assume throughout that the spike-and-slab prior specifications are given by $v_0 = .001$ and $v_1 = 10$. These specifications are consistent with the previous recommendation that the ratio v_1/v_0 be no greater than 10,000, for the purposes of reliably estimating the parameters $(\boldsymbol{\beta}, \boldsymbol{\gamma})$ via MCMC methods (George & McCulloch, 1997, p. 368).

For any one of the models described in this section, the posterior predictive density $f_n(y|\mathbf{x}_0, \hat{\sigma}^2)$ provides an estimate of the "overall" effect-size distribution of the underlying population, and is a symmetric and unimodal distribution, conditionally on covariates $\mathbf{x} = \mathbf{x}_0 = (1, 0, \dots, 0)^\top$. For example, under a Bayesian normal fixed-effects model, the posterior predictive density $f_n(y|\mathbf{x}_0, \hat{\sigma}^2)$ estimate of the overall population effect-size distribution is a normal density (distribution). Under a Bayesian normal 2-level random-effects model with

$\text{ga}(\sigma_0^{-2}|a, b)$ prior for independent level-2 random intercepts $\boldsymbol{\mu}_0$, the posterior predictive estimate $f_n(y|\mathbf{x}_0, \widehat{\sigma}^2)$ of the population effect-size distribution is given by a student density (distribution) (e.g., Denison et al. 2002, Appendix). The student distribution is very similar to a normal distribution, except that the student distribution has thicker tails.

For a given Bayesian normal meta-analytic model with prior density having the general form (4), the posterior density $\pi(\boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\mu}_0, \boldsymbol{\mu}_{00}, \sigma_0^2, \sigma_{00}^2, \psi|\mathcal{D}_n)$, the posterior predictive density $f_n(y|\mathbf{x}, \widehat{\sigma}^2)$, and any functional of these densities, can be estimated through the use of standard MCMC Gibbs and Metropolis sampling algorithms for normal linear and random-effects models (e.g., Gilks et al. 1993; Denison et al. 2002).

3 The Bayesian Nonparametric Meta-Analysis Model

For effect-size data, our Bayesian nonparametric meta-analysis model is defined by an infinite random-intercepts mixture of regressions. The model assumes the data likelihood:

$$f(y_i|\mathbf{x}_i, \widehat{\sigma}_i^2; \boldsymbol{\zeta}) = \int n(y_i|\mu_0 + \mathbf{x}_i^\top \boldsymbol{\beta}, \phi \widehat{\sigma}_i^2) dG_{\mathbf{x}}(\mu_0) \quad (7)$$

$$= \sum_{j=-\infty}^{\infty} n(y_i|\mu_{0j} + \mathbf{x}_i^\top \boldsymbol{\beta}, \phi \widehat{\sigma}_i^2) \omega_j(\mathbf{x}_i^\top \boldsymbol{\beta}_\omega, \sigma_\omega), \quad i = 1, \dots, n. \quad (8)$$

Using standard terminology for discrete mixture models (e.g., McLachlan & Peel, 2000), $G_{\mathbf{x}}$ is the (discrete) mixing distribution which depends on covariates \mathbf{x} ; the component indices are given by $j = 0, \pm 1, \pm 2, \dots$, the component (kernel) probability densities are given by the normal densities, $(n(y_i|\mu_{0j} + \mathbf{x}_i^\top \boldsymbol{\beta}, \phi \widehat{\sigma}_i^2))_{j=-\infty}^{\infty}$; the component parameters are given by $(\mu_{0j})_{j=-\infty}^{\infty}$; and the mixing weights are given by $(\omega_j(\mathbf{x}_i^\top \boldsymbol{\beta}_\omega, \sigma_\omega))_{j=-\infty}^{\infty}$ which sum to 1 at every $\mathbf{x} \in \mathcal{X}$. Specifically, in the model, the mixture weights $\omega_j(\mathbf{x}_i^\top \boldsymbol{\beta}_\omega, \sigma_\omega)$ are each defined by the difference between two standard Normal(0,1) cumulative distribution functions (c.d.f.s):

$$\omega_j(\mathbf{x}_i^\top \boldsymbol{\beta}_\omega, \sigma_\omega) = \Phi(\{j - \mathbf{x}_i^\top \boldsymbol{\beta}_\omega\}/\sigma_\omega) - \Phi(\{j - 1 - \mathbf{x}_i^\top \boldsymbol{\beta}_\omega\}/\sigma_\omega).$$

Thus, the mixture weights can be viewed as the categorical probabilities of a cumulative-probits regression model (e.g., McCullagh, 1980), for infinitely-many ordered categories $j = 0, \pm 1, \pm 2, \dots$. Thus, it is easy to see that the mixture weights $\omega_j(\mathbf{x}_i^\top \boldsymbol{\beta}_\omega, \sigma_\omega)$ sum to 1 at every $\mathbf{x} \in \mathcal{X}$. Finally, our mixture model (7) provides a flexible (infinite) mixture of normal densities, conditional on any covariates \mathbf{x} of interest. The development of our model is, in part motivated, by the well known fact that any smooth probability density can be approximated arbitrarily-well by a suitable discrete mixture of normal densities (e.g., Lo, 1984).

— Figure 2 —

As shown in equation (7), for a set of data $\mathcal{D}_n = \{(y_i, \mathbf{x}_i, \widehat{\sigma}_i^2)\}_{i=1}^n$, the model assumes that each effect-size y_i is distributed by according to a probability density $f(y_i|\mathbf{x}_i, \widehat{\sigma}_i^2; \boldsymbol{\zeta})$ that is constructed by a mixture of an infinite number of normal densities $n(y_i|\mu_{0j} + \mathbf{x}_i^\top \boldsymbol{\beta}, \phi \widehat{\sigma}_i^2)$, having corresponding means $\mu_{0j} + \mathbf{x}_i^\top \boldsymbol{\beta}$ and mixture weights $\omega_j(\mathbf{x}_i^\top \boldsymbol{\beta}_\omega, \sigma_\omega)$ (for $j = 0, \pm 1, \pm 2, \dots$).

Therefore, given covariates \mathbf{x} and model parameters $\boldsymbol{\zeta}$, the model is flexible enough to allow the shape of the effect-size distribution (density) $f(y|\mathbf{x}, \hat{\sigma}^2; \boldsymbol{\zeta})$ to take on virtually any form; this density can be unimodal symmetric, or skewed, or more multimodal. Moreover, the model allows the entire shape and location of the effect-size distribution (density) $f(y|\mathbf{x}, \hat{\sigma}^2; \boldsymbol{\zeta})$ to change flexibly with the covariates \mathbf{x} . The model has these flexibilities, because it models the effect-size density $f(y|\mathbf{x}, \hat{\sigma}^2; \boldsymbol{\zeta})$ by infinitely-many random intercept parameters $\boldsymbol{\mu}_0 = (\mu_{0j})_{j=-\infty}^{\infty}$, corresponding to infinitely-many covariate-dependent mixture weights $\{\omega_j(\mathbf{x}_i^\top \boldsymbol{\beta}_\omega, \sigma_\omega) : j = 0, \pm 1, \pm 2, \dots\}$.

In the Bayesian nonparametric model, the parameter σ_ω controls the level of multimodality of $f(y|\mathbf{x}, \hat{\sigma}^2; \boldsymbol{\zeta})$. To explain, assume for the moment that $\mathbf{x}_i^\top \boldsymbol{\beta} = 0$ and $f(y|\mathbf{x}) = f(y|\mathbf{x}, \hat{\sigma}^2 = 1; \boldsymbol{\zeta})$, for simplicity, and with no loss of generality. On the one hand, a small value of σ_ω indicates that $f(y|\mathbf{x})$ is unimodal, i.e., modeled as a unimodal normal density $n(y_i|\mu_j, \hat{\sigma}_i^2)$ for a j satisfying $j - 1 < \mathbf{x}^\top \boldsymbol{\beta}_\omega < j$, with mixture weight $\omega_j(\mathbf{x}^\top \boldsymbol{\beta}_\omega, \sigma_\omega)$ near 1. This is because the function $\Phi(\mathbf{x}^\top \boldsymbol{\beta}_\omega / \sigma_\omega)$ is approximately 0 for $\mathbf{x}^\top \boldsymbol{\beta}_\omega < 0$, while it is approximately 1 for $\mathbf{x}^\top \boldsymbol{\beta}_\omega > 0$. As σ_ω approaches infinity, the mixture weights become more spread out, and then $f(y|\mathbf{x})$ becomes multimodal, with each mixture weight $\omega_j(\mathbf{x}^\top \boldsymbol{\beta}_\omega, \sigma_\omega)$ above zero and much less than 1. These ideas are illustrated in Figure 2, which plots the mixture weights and the corresponding density of our model, $f(y|\mathbf{x})$, for a range of σ_ω , given $\mathbf{x}^\top \boldsymbol{\beta}_\omega = .7$, given samples of (μ_j, σ_j^2) from a normal-gamma distribution. As shown, the conditional density $f(y|\mathbf{x})$ is unimodal when σ_ω is small, and $f(y|\mathbf{x})$ becomes more multimodal as σ_ω increases. The level of multimodality in the data is indicated by the posterior distribution of σ_ω , under the model.

The Bayesian nonparametric meta-analytic model (7) is completed by the specification of a joint proper prior density $\pi(\boldsymbol{\zeta})$ for the infinitely-many model parameters $\boldsymbol{\zeta} = (\boldsymbol{\beta}, \gamma, (\mu_{0j})_{j=-\infty}^{\infty}, \phi, \boldsymbol{\beta}_\omega, \sigma_\omega)$, according to the joint prior distributions:

$$\mu_{0j} | \sigma_0^2 \sim n(\mu_{0j} | 0, \sigma_0^2), \quad j = 0, \pm 1, \pm 2, \dots; \quad (9a)$$

$$\beta_0 \sim n(\beta_0 | 0, v \rightarrow \infty); \quad (9b)$$

$$(\beta_k, \gamma_k) | \phi \sim n(\beta | 0, \phi 10^{\gamma_k} .001^{1-\gamma_k}) .5^{\gamma_k} (1 - .5)^{1-\gamma_k}, \quad k = 1, \dots, p; \quad (9c)$$

$$\phi^{-1} \sim \text{ga}(\sigma_\omega^{-2} | a_\phi/2, a_\phi/2) \quad (9d)$$

$$\sigma_0^2 \sim \text{un}(\sigma_0 | 0, b_0) \quad (9e)$$

$$(\boldsymbol{\beta}_\omega, \sigma_\omega) \sim n_{p+1}(\boldsymbol{\beta}_\omega | \mathbf{0}, \sigma_\omega^2 10^5 \mathbf{I}_{p+1}) \text{ga}(\sigma_\omega^{-2} | 1, 1). \quad (9f)$$

As shown, a diffuse prior is assigned to the overall mean effect-size parameter β_0 . Also, as shown in (9c), we adapt the default spike-and-slab priors, to enable automatic covariate (predictor) selection in the posterior distribution of our model (George & McCulloch, 1997). Also, the gamma prior for the inverse dispersion parameter ϕ^{-1} has mean $E(\phi^{-1}) = 1$ and variance $\text{Var}(\phi^{-1}) = \frac{2}{a_\phi}$, with a_ϕ indicating the degree of ‘belief’ in this prior (Nam, et al., 2003). Furthermore, we specify uniform prior density $\text{un}(\sigma_0 | 0, b_0)$ for the variance σ_0^2 of the random intercepts $(\mu_{0j})_{j=-\infty}^{\infty}$, for a reasonably-large value b_0 , as consistent with previous recommendations (Gelman, 2006, Section 7.1). Alternatively, one may specify a half- t prior density for σ_0 . Most of the prior distributions in (9) represent default and rather diffuse choices of prior, which can be used in general meta-analytic applications where prior information is typically limited. Of course, if for a given meta-analytic data set, there is more

prior (e.g., scientific) information available about one or more of the model parameters, then the prior distributions can be modified accordingly.

It is instructive to relate the parameters of the general normal meta-analytic model of equation (1) that are assigned a general prior density of equation (4) (Sections 2.2 and 2.4), with the parameters of the Bayesian nonparametric meta-analytic model. Across both models, the linear regression coefficients $\boldsymbol{\beta}$, including the overall effect-size mean β_0 , have the same interpretation of how the mean effect-size depends on covariates; the parameters $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_p)^\top$ have the same interpretation as (random) indicators of which covariates are included as significant predictors of the models; the infinitely-many random intercepts $(\mu_{0j})_{j=-\infty}^{\infty}$ of the Bayesian nonparametric model have the same interpretation as the level-2 random intercepts $\boldsymbol{\mu}_0$ of a normal random-effects meta-analytic model; and the parameter σ_0^2 has the same interpretation as the variance of the level-2 random intercepts. A key difference is that the Bayesian nonparametric model is a discrete mixture model which specifies a covariate-dependent infinite-mixture distribution $G_{\mathbf{x}}$ for the random intercept parameter μ_0 , as opposed to a hierarchical model or a random effects model. In contrast, a normal random-effects model is a hierarchical model which specifies a normal mixture distribution G for the random intercept parameter, such that the mixture distribution is not covariate dependent. Moreover, for the Bayesian nonparametric model, the discrete mixture distribution $G_{\mathbf{x}}$ induces (random) clusterings among the n study reports y_i (via the posterior distribution of the model), in terms of the random intercept parameter μ_0 . This clustering feature of the Bayesian nonparametric model enables the model to account for correlations among the study reports y_i ($i = 1, \dots, n$), lessening the need to specify a non-diagonal sampling covariance matrix $\widehat{\boldsymbol{\Sigma}}_n$ to account for correlated effect-sizes.

Following Bayes' theorem, the data likelihood $L(\mathcal{D}_n) = \prod_{i=1}^n f(y_i | \mathbf{x}_i, \widehat{\sigma}_i^2; \boldsymbol{\zeta})$ updates the prior density $\pi(\boldsymbol{\zeta})$, to a posterior density $\pi(\boldsymbol{\zeta} | \mathcal{D}_n)$, given by equation (2). Then the posterior predictive density is given by equation (3), which gives an estimator of the true effect-size density in the study population, given data \mathcal{D}_n and covariates \mathbf{x} of interest. Also, recall that for the task of covariate selection, the k th covariate can be viewed as a "significant predictor," when the posterior inclusion probability of the covariate, $\Pr[\gamma_k = 1 | \mathcal{D}_n]$, is at least .5 (Barbieri & Berger, 2004). Finally, the level of multimodality in the density $f(y | \mathbf{x}, \widehat{\sigma}^2, \boldsymbol{\zeta})$ is indicated by the posterior distribution of σ_ω .

Karabatsos and Walker (2012) describe the Markov Chain Monte Carlo (MCMC) methods can be used to perform inference of the posterior density $\pi(\boldsymbol{\zeta} | \mathcal{D}_n)$, of the posterior predictive density $f_n(y | \mathbf{x}, \widehat{\sigma}^2)$ of the model, and to perform inference of any functional of these densities.

4 Bayesian Predictive Model Assessment Methods

Model selection is the practice of comparing different models that are fitted to a common sample data set, and then identifying the single model that best describes or predicts the underlying population distribution of the sample data. In meta-analytic practice, it is often of interest to perform model selection (e.g., Sutton, 2000, Section 11.7.3). For example, model selection is used in meta-analysis to choose between the fixed-effects and random-effects model (Borenstein et al. 2010), or to select important predictors of the effect-size in

a regression setting (Higgins & Thompson, 2004).

After M meta-analytic models are fit to a data set, \mathcal{D}_n , the predictive performance of each Bayesian model $m \in \{1, \dots, M\}$ can be assessed by the mean-square posterior predictive-error criterion

$$D(m) = \sum_{i=1}^n \{y_i - E_n(Y_i|\mathbf{x}_i, \hat{\sigma}_i^2, m)\}^2 + \sum_{i=1}^n \text{Var}_n(Y_i|\mathbf{x}_i, \hat{\sigma}_i^2, m) \quad (10)$$

$$= \sum_{i=1}^n \int (y - y_i)^2 f_n(y|\mathbf{x}_i, \hat{\sigma}_i^2, m) dy = \sum_{i=1}^n D_i(m) \quad (11)$$

(Laud & Ibrahim, 1995; Gelfand & Ghosh, 1998). The criterion (10) is a standard criterion that is often used for the assessment and comparison of Bayesian models (e.g., Gelfand & Banerjee, 2010). Among the M Bayesian meta-analytic models that are compared, the model with the smallest value of $D(m)$ is identified as the one that best describes the underlying population distribution of the given sample data set \mathcal{D}_n . The first term of (10) measures data goodness-of-fit, and the second term is a penalty that is large for models which either over-fit or under-fit the data, as in other classical model selection criteria. Taking the square root, $\sqrt{D(m)}$, makes the criterion interpretable on the original scale of the effect size (y). Similarly, the individual square-root quantities $\sqrt{D_i(m)}$ (for $i = 1, \dots, n$) can provide a detailed assessment about a model's predictive performance. A large value of $\sqrt{D_i(m)}$ would indicate that the observed effect-size y_i is an outlier under the model.

5 Illustration

In this section, we illustrate all the methods that we presented in Sections 2-4, through the meta-analysis of a large real data set involving 24 covariates. In this data analysis, we use the $D(m)$ predictive mean-square error criterion to compare the predictive accuracy of the Bayesian nonparametric meta-analysis model, and the various Bayesian normal fixed-effects and normal random-effects models. We also compare some of the parameter estimates between these Bayesian models, as well as the parameter estimates of normal fixed-effects and random-effects models estimated either under full maximum likelihood (MLE) or restricted maximum likelihood (REML). For each data analysis, each covariate was previously z-standardized to have mean 0 and variance 1, by taking $x_{ki} = (x'_{ki} - \hat{\mu}'_k) / \hat{\sigma}'_{X(k)}$ for $i = 1, \dots, n$, given the mean and standard deviation ($\hat{\mu}'_k, \hat{\sigma}'_{X(k)}$) of the original covariate data (x'_{k1}, \dots, x'_{kn}). Then the the estimate of the intercept parameter (β_0) is interpretable as the mean study effect-size, and the β coefficients are all interpretable on a common scale.

In total, we will consider a total of 16 Bayesian meta-analytic models for the data set, including the Bayesian nonparametric model, along with various fixed effects models and various 2-level or 3-level normal-effects models, which among other things, differ as to whether or not they have covariates, whether or not they have spike-and-slab priors for covariate selection. For all of these models, we specified the same prior densities for parameters that the models shared in common, in order to place the Bayesian model comparisons on a rather equal-footing. These priors were generally consistent with the recommendations of the pre-

vious literature (see Sections 2.4, 3). Specifically, for all Bayesian models, as follows: we assigned the normal prior density $\pi(\beta_0) = \text{n}(\beta_0|0, v \rightarrow \infty)$ (with $v = 10^5$); we assigned diffuse normal priors $\pi(\beta_k) = \text{n}(\beta_k|0, v \rightarrow \infty)$ (with $v = 10^5$), $k = 1, \dots, p = 24$, to the slope coefficients of all models with covariates and without spike-and-slab priors; we assigned hyper-prior variances $v_0 = .001$ and $v_1 = 10$, and Bernoulli prior parameters $\text{Pr}(\gamma_k = 1) = \frac{1}{2}$ ($k = 1, \dots, p = 24$), for all models with covariates and spike-and-slab priors (in terms of equation 6); we assigned the uniform prior density $\pi(\sigma_0) = \text{un}(\sigma_0|0, 100)$ with large scale (100) for the variance parameter of the Bayesian nonparametric model and all normal 2-level random effects models; we specified the rather non-informative prior for (σ_0^2, ψ) (equation (5), Section 2.4), for all Bayesian 2-level normal random-effects models that allow for correlated random intercepts (Stevens & Taylor, 2009); and we specified the uniform prior density $\pi(\sigma_{00}) = \text{un}(\sigma_{00}|0, 100)$ with large scale (100) for the level-3 variance parameter σ_{00}^2 , for all 3-level normal random-effects models. For the Bayesian nonparametric model in particular, we also specified rather diffuse (high-variance) priors $\phi^{-1} \sim \text{ga}(\sigma_\omega^{-2} | .5/2, .5/2)$, $\beta_\omega | \sigma_\omega^2 \sim \text{n}_{p+1}(\beta_\omega | \mathbf{0}, \sigma_\omega^2 10^5 \mathbf{I}_{p+1})$, and $\sigma_\omega^{-2} \sim \text{ga}(\sigma_\omega^{-2} | 1, 1)$ (see equations (9d) and (9f), Section 3).

For the application of each Bayesian model, the posterior distribution of the parameters was estimated on the basis of 200,000 MCMC samples, after trace plots indicated that MCMC samples of key model parameters and of the $D(m)$ criterion stabilized and mixed well, and after 95% Monte Carlo confidence intervals (MCCIs) of these quantities attained half-widths that were small for practical purposes, i.e., that typically ranged between .01 and .05, and not exceeding .1. Again, these procedures accord with previous recommendations for checking the quality of MCMC-based posterior estimates, based on a single MCMC run (Geyer, 1992; 2011, Chapter 1). Our Bayesian nonparametric meta-analytic model was estimated using a program code we wrote in the MATLAB (Natick, VA) software language, for an earlier paper (Karabatsos & Walker, 2012). The code for the meta-analytic model is provided along with this paper. The Bayesian normal fixed-effects and random-effects models were each estimated using code we wrote in MATLAB. Finally, each of the twelve 2-level and 3-level random-effects models, assuming uncorrelated random intercepts, were also estimated by full maximum likelihood and restricted maximum likelihood, using the nmle package (Pinheiro, et al. 2010) of the R statistical software (R Development Core Team, 2012).

5.1 Behavioral Genetics Data

Antisocial behavior, which includes aggression, willingness to violate rules and laws, defiance of adult authority, and violation of social norms (Walker et al. 2003), is the most frequent reason why children are referred for mental health services in schools (Adelman & Taylor, 2010). Yet, it is the most intractable of all behavior and mental health problems, is challenging to treat, and must be addressed across the lifespan (Moffitt, 1993; 2005).

— Figure 3 —

To advance understanding and treatment, many behavioral genetic studies have investigated the heritability of antisocial behavior, by correlating ratings of antisocial behavior among monozygotic (MZ) identical twin pairs, and among dizygotic (DZ) fraternal twin

pairs. In each study, the ratings were done either by the mother, father, teacher, self, or an observer. Then the heritability, defined as the proportion of phenotypic variance explained by genetic factors, is estimated by twice the difference between the MZ correlation and DZ correlation, for twins of the same sex. Specifically, for a given gender, suppose that n_{MZ} monozygotic (MZ) identical twin pairs yield a correlation $\hat{\rho}_{MZ}$ on an antisocial behavior trait, such as conduct disorder, aggression, delinquency, and externalizing behavior. Also, suppose that n_{DZ} dizygotic (DZ) fraternal twin pairs yield a correlation $\hat{\rho}_{DZ}$ on the same trait. Then the heritability of the antisocial behavior trait is estimated by:

$$\hat{h}^2 = 2(\hat{\rho}_{MZ} - \hat{\rho}_{DZ}) \quad (12)$$

(Falconer & Mackay, 1996). This effect-size statistic (12) has sampling variance:

$$\hat{\sigma}^2 = 4\{[(1 - \hat{\rho}_{MZ}^2)^2/n_{MZ}] + [(1 - \hat{\rho}_{DZ}^2)^2/n_{DZ}]\}.$$

We identified 29 independent studies that provided the information necessary to estimate antisocial behavior heritability, for the $n = 71$ independent samples (study reports) of MZ-DZ twin comparisons (Talbot, et al. 2012). These studies were published during years 1966 through 2009, and their full references are listed in the Appendix. There were 2-3 heritability estimates per study on average, and each study provided between 1 to 10 estimates. The left panel of Figure 3 presents the heritability estimates of antisocial behavior (i.e., the effect-size observations), stratified by gender, by rater type, and by the studies which were numerically identified by publication year order (see Appendix). The one slightly-negative estimate ($-.06$) may have resulted from sampling error (Gill & Jensen, 1968), as suggested by its relatively-large variance.

Here, it is of interest to perform a meta-analysis of the studies, to learn about the overall heritability (effect-size) distribution for the underlying study population, as well as to learn how heritability changes with key study-level covariates. Again, analyses will be performed using the various fixed-effects and random-effects models under maximum likelihood estimation, and using our Bayesian nonparametric model. A total of 24 covariates were identified. They include publication year, the square root of the heritability variance (denoted SE(ES)) to provide an investigation of publication bias; indicators (0-1) of female status (49%) versus male; ten indicators of antisocial behavior ratings done by mother (mean=.53), father (.06), teacher (.24), self (.15), independent observer (.04), and ratings done on conduct disorder (.03), aggression (.40), delinquency (.10), and externalizing (.48) antisocial behavior; an indicator of whether a weighted average of heritability measures was taken within study over different groups of raters who rated the same twins (28% of cases, for which the ten indicator covariates are scored as group proportions), mean age of the study subjects in months (overall mean=119.8, s.d.=49.5); indicators of hi-majority ($\geq 60\%$) white twins in study (94%), zygosity obtained by questionnaire (80%), or through DNA samples (68%), study inclusion of low socioeconomic (SES) status level subjects (20%) and mid-to-high SES subjects (90%), missing SES information (10%), representative sample (85%), longitudinal sample (85%), and location of the study in terms of latitude and longitude.

Model	$D(m)$	Model	$D(m)$
BNP-ss	0.6	D2L-x	5.5
D2L-0	4.8	3L-x	5.5
2L-0, by MZ-DZ	4.8	2L-0, by Study	5.8
3L-0	4.8	FE-0	5.9
D2L-ss	5.4	2L-ss, by Study	6.0
2L-ss, by MZ-DZ	5.4	FE-ss	6.0
3L-ss	5.4	2L-x, by Study	6.0
2L-x, by MZ-DZ	5.5	FE-x	6.0

Table 2: For the behavioral genetics data, a comparison of the predictive mean square error criterion, between the Bayesian nonparametric model, and various normal fixed-effects (FE) models, and various normal 2-level (2L), dependent 2-level (D2L), and 3-level random effects (RE) models, each of which either has no covariates (0), or 24 covariates (x), or 24 covariates with stochastic search variable selection (ss). Also, each 2L or D2L model groups the random intercept parameters by the 29 studies (by Study), or groups them by the 71 independent samples of twins (by MZ-DZ).

Given the structure of the data, with the $n = 71$ heritability (effect-size) estimate reports nested within the 29 studies, any one of at least 15 normal fixed-effects models or normal random-effects models can be considered for the purposes of meta-analysis. Specifically, they include: fixed-effects models; 2-level random-effects models, with level-2 random intercepts $\boldsymbol{\mu}_0$ assumed to be independent via the specification of a multivariate normal $n_n(\boldsymbol{\mu}_0|\mathbf{0}, \sigma_0^2\mathbf{I}_n)$ distribution, with either a structure that has each of the 71 independent samples of MZ-DZ twin comparisons defining its own group, or a grouping structure has each of the 29 studies defining its own group; 2-level random-effects models that allow for dependent level-2 random intercepts via the specification of a multivariate normal $n_n(\boldsymbol{\mu}_0|\mathbf{0}, \sigma_0^2\mathbf{I}_n + \psi\mathbf{M}_n)$ distribution (Stevens & Taylor, 2009) with the binary (0-1) matrix \mathbf{M}_n indicating which pairs of the 71 study reports belong in the same study; 3-level random-effects models, with the 71 heritability (effect-size) estimate reports (level 2) nested within the 29 studies (level 2), and with random intercepts modeled respectively by the multivariate normal densities $n_n(\boldsymbol{\mu}_0|\mathbf{0}, \sigma_0^2\mathbf{I}_n)$ and $n_n(\boldsymbol{\mu}_{00}|\mathbf{0}, \sigma_{00}^2\mathbf{I}_n)$; with each model having either no covariates, or having all 24 covariates with no spike-and-slab priors for automatic covariate selection, or having all 24 covariates along with spike-and-slab priors for automatic covariate selection via the posterior distribution. Finally, we also analyze the data set using the Bayesian meta-analytic model, which includes all 24 covariates, and which assigns spike-and-slab priors for automatic covariate selection via the posterior distribution.

For the behavioral genetics data, Table 2 compares the estimate of the mean-squared predictive error criterion, $D(m)$, between the 15 Bayesian normal fixed-effects models and normal random-effects models, and the Bayesian nonparametric meta-analytic model. Among all the models compared, the Bayesian nonparametric model attained the smallest value of the predictive mean-square error criterion $D(m)$, by a relatively-large margin. Hence, among all the models compared, the Bayesian nonparametric model provides the best description of the study-population effect-size distribution that underlies the (sample) behavioral genetics data set. The meta-analytic models that attained the second best value of the criterion had

about 8 times the mean squared predictive error, compared to our Bayesian nonparametric model. Also, for the Bayesian nonparametric model, the 5-number summary of the estimates of the predictive residuals $\sqrt{D_i(m)}$ was (.0, .0, .1, .1, .3), over the 71 heritability effect-size observations y_i . So none of the observations appeared to be outliers under the model.

The mean heritability (effect-size) estimate ($\hat{\beta}_0$) was quite similar among all the 16 Bayesian models, ranging from .49 to .51; while the posterior mean estimates ($\hat{\sigma}_0^2, \hat{\sigma}_{00}^2, \hat{\psi}$) of the random intercept variances ranged between .00 to .02. Similar estimates were obtained from normal fixed-effects and random-effects models that were estimated either under maximum likelihood, or by restricted maximum likelihood. For all the Bayesian models assigned spike-and-slab priors, the posterior inclusion probabilities $\Pr[\gamma = 1 | \mathcal{D}_{71}]$ did not exceed .05 for all 24 covariates, well below the significance threshold of .5. For the Bayesian nonparametric model, the marginal posterior mean (standard deviation) estimate of the dispersion parameter ϕ was .09 (.03).

As mentioned, the Bayesian nonparametric model is an infinite mixture model that is able to account for all possible shapes and locations of effect-size distributions, including all normal distributions. Meanwhile, in terms of the predictive mean-square error criterion $D(m)$, the Bayesian nonparametric model far-outperformed all other normal fixed-effects and normal random-effects models, which assume more strictly assume normal effect-size densities. These facts together suggest that the data set violates the assumptions of effect-size normality. For the Bayesian nonparametric model, the right panel of Figure 4 presents the posterior predictive estimate of the overall heritability (effect-size) distribution, for the underlying population of studies. This estimate is conditional on the covariates $\mathbf{x} = \mathbf{x}_0 = (1, 0, \dots, 0)^\top$, and also, it is conditioned on the minimum effect-size variance σ_i^2 of .0001 over all 71 heritability reports, so that this distribution reflects information from a large-sample study. According to this estimate of the overall heritability (effect-size) distribution, there is some evidence of skewness (-.1), with the overall mean (.50) and median (.51) heritability (effect-size) being slightly different. Moreover, there seems to be two modes in this distribution, one at about the mean of .50, and the other at about .35, suggesting there are about two "significant" heritabilities (effect sizes) in the population, not only one. Upon closer inspection, the modes appear to be at heritability values of .38 and .51. So both modes can provide information that contribute to the accumulation of evidence about the overall heritability (effect-size) for the substantive researchers of behavioral genetics.

— Figure 4 —

The first panel of Figure 4 shows the median (50%ile) and interquartile range (i.e., 25%ile and 75%ile) of the posterior predictive estimate of the heritability (effect-size) distribution, by SE(ES) (and by corresponding effect-size variance $\{\text{SE(ES)}\}^2 = \hat{\sigma}^2$). As shown, the median effect-size has a rather nonlinear relationship with SE(ES), but the lack of strong relationship of the median effect-size with SE(ES) further confirms a lack of evidence of publication bias in the data.

Finally, another important issue in this area of behavioral genetics deals with the issue of informant discrepancy; that is, the issue of whether the heritability (effect-size) estimates are the same across raters, or whether they depend on rater type, and further the heritability estimates may also depend on ratee age. Recall that each of the 71 heritability (effect-size)

estimates were obtained from ratings of antisocial behavior that were made by one of 5 rater types, namely, the mother, the father, the teacher, the self, or an independent observer. To address this research issue in detail, the remaining five panels of Figure 4 present the posterior predictive estimates of the effect-size distributions, conditional on covariates \mathbf{x} that indicate rater type and ratee age, while controlling for all other non-constant covariates by setting them to zero, and while conditioning on the effect-size variance estimate $\hat{\sigma}_i^2 = .0001$ that reflects a large-sample study. As shown in these panels, the heritability-age correlation seems to be slightly negative for all rater types. Moreover, the heritability distributions seemed to be similar among mother, father, self, and independent observer raters, while the teacher raters have noticeably different heritability distributions.

6 Discussion

In this paper, we have proposed a Bayesian nonparametric model for meta-analysis, and demonstrated its suitability for meta-analytic data sets that give rise to asymmetric and more multimodal effect-size population distributions. As mentioned, the traditional normal fixed- and random-effects models, while frequently used for meta-analysis, are not fully satisfactory because they make empirically-falsifiable assumptions about the data. They include the assumption that the effect-sizes are normally-distributed (conditionally on model parameters). As we have shown, empirical violations of such an assumption can negatively affect the accuracy of prediction of meta-analytic data.

In contrast to the traditional models, our proposed Bayesian nonparametric model flexibly accounts for all distributions of the effect-sizes, including all normal distributions. For the real data set that was analyzed in this paper, this flexibility enabled the Bayesian model to provide a better description of the underlying effect-size distribution of the underlying study population. At the same time, the model provides a richer description of meta-analytic data, by allowing the data analyst to infer the whole distribution of effect-sizes, over studies, and to infer how the whole distribution changes as a function of key study-level covariates. Thus, the model goes beyond the mean as the measure of an overall effect-size. Furthermore, for the given meta-analytic data set at hand, the Bayesian nonparametric model automatically identifies important study-level predictors of the mean effect-size.

Acknowledgements

This research is supported by grant SES-1156372, from the NSF program in Methodology, Measurement, and Statistics. The author gives thanks to the Editor, and to anonymous Associate Editor and two reviewers, for detailed suggestions that have helped improve the presentation of this manuscript.

References

- Adelman, H. & Taylor, L. 2010. *Mental health in schools: Engaging learners, preventing problems, improving schools*, Corwin Press, Thousand Oaks, CA.
- Aitchison, J. 1975. Goodness of prediction fit, *Biometrika* **62**: 547-554.

- Anderson, T. & Darling, D. 1952. Asymptotic theory of certain "goodness-of-fit" criteria based on stochastic processes, *Annals of Mathematical Statistics* **23**: 193-212.
- Barbieri, M. & Berger, J. 2004. Optimal predictive model selection, *Annals of Statistics* **32**: 870-897.
- Berkey, C., Hoaglin, D., Mosteller, F. & Colditz, G. 1995. A random-effects regression model for meta-analysis, *Statistics in Medicine* **14**: 395-411.
- Borenstein, M. 2009. Effect sizes for continuous data, in H. Cooper, L. Hedges & J. Valentine (eds), *The Handbook of Research Synthesis and Meta-Analysis (Second Edition)*, Russell-Sage Foundation, New York, pp. 221-235.
- Borenstein, M., Hedges, L., Higgins, J. & Rothstein, H. 2010. A basic introduction to fixed-effect and random-effects models for meta-analysis, *Research Synthesis Methods* **1**: 97-111.
- Branscum, A. J. & Hanson, T. E. 2008. Bayesian nonparametric meta-analysis using Pólya tree mixture models, *Biometrics* **64**: 825-833.
- Brooks, S., Gelman, A., Jones, G. & Meng, X. 2011. *Handbook of Markov Chain Monte Carlo*, Chapman and Hall/CRC, Boca Raton, FL.
- Burr, D. & Doss, H. 2005. A Bayesian semiparametric model for random-effects meta-analysis, *Journal of the American Statistical Association* **100**: 242-251.
- Chib, S. & Greenberg, E. 1995. Understanding the Metropolis-Hastings algorithm, *American Statistician* **49**: 327-335.
- Cooper, H. & Hedges, L. 2009. Research synthesis as a scientific process, in H. Cooper, L. Hedges & J. Valentine (eds), *The Handbook of Research Synthesis and Meta-Analysis (Second Edition)*, Russell Sage Foundation, New York, pp. 3-18.
- Denison, D., Holmes, C., Mallick, B. & Smith, A. 2002. *Bayesian Methods for Nonlinear Classification and Regression*, John Wiley and Sons, New York.
- DuMouchel, W. & Normand, S.-L. 2000. Computer-modeling and graphical strategies for meta-analysis, in D. Stangl & D. Berry (eds), *Meta-analysis in medicine and health policy*, Marcel Dekker, New York, pp. 127-178.
- Egger, M., Smith, G. D., Schneider, M. & Minder, C. 1997. Bias in meta-analysis detected by a simple, graphical test, *British Medical Journal* **315**: 629-634.
- Falconer, D. & Mackay, T. 1996. *Introduction to quantitative genetics*, Longman, London.
- Fleiss, J. & Berlin, J. 2009. Effect size for dichotomous data, in H. Cooper, L. Hedges & J. Valentine (eds), *The Handbook of Research Synthesis and Meta-Analysis (Second Edition)*, Russell Sage Foundation, New York, pp. 237-253.

- Gelfand, A. & Banerjee, S. 2010. Multivariate spatial process models, in A. Gelfand, P. Diggle, P. Guttorp & M. Fuentes (eds), *Handbook of Spatial Statistics*, Chapman and Hall/CRC, Boca Raton, pp. 495-515.
- Gelfand, A. & Ghosh, J. 1998. Model choice: A minimum posterior predictive loss approach, *Biometrika* **85**: 1-11.
- Gelfand, A. & Smith, A. 1990. Sampling-based approaches to calculating marginal densities, *Journal of the American Statistical Association* **85**: 398-409.
- Gelman, A. 2006. Prior distributions for variance parameters in hierarchical models, *Bayesian Analysis* **3**: 515-533.
- George, E. & McCulloch, R. 1997. Approaches for Bayesian variable selection, *Statistica Sinica* **7**: 339-373.
- Geyer, C. 1992. Practical Markov chain Monte Carlo (with discussion), *Statistical Science* **7**: 473-511.
- Geyer, C. 2011. Introduction to MCMC, in S. Brooks, A. Gelman, G. Jones & X. Meng (eds), *Handbook of Markov Chain Monte Carlo*, CRC, Boca Raton, FL, pp. 3-48.
- Gilks, W., Wang, C., Yvonnet, B. & Coursaget, P. 1993. Random-effects models for longitudinal data using Gibbs sampling, *Biometrics* **49**: 441-453.
- Gill, J. & Jensen, E. 1968. Probability of obtaining negative estimates of heritability, *Biometrics* **24**: 517-526.
- Glass, G. 1976. Primary, secondary, and meta-analysis of research, *Educational Researcher* **5**: 3-8.
- Gleser, L. & Olkin, I. 2009. Stochastically dependent effect sizes, in H. Cooper, L. Hedges & J. Valentine (eds), *The Handbook of Research Synthesis and Meta-Analysis* (Second Edition), Russell Sage Foundation, New York, pp. 357-376.
- Harville, D. 1977. Maximum likelihood approaches to variance component estimation and to related problems, *Journal of the American Statistical Association* **72**: 320-338.
- Hedges, L. 1981. Distribution theory for Glass's estimator of effect size and related estimators, *Journal of Educational and Behavioral Statistics* **6**: 107-128.
- Hedges, L. & Vevea, J. 1998. Fixed and random effects models in meta analysis, *Psychological Methods* **3**: 486-504.
- Higgins, J. & Thompson, S. 2004. Controlling the risk of spurious findings from meta-regression, *Statistics in Medicine* **23**: 1663-1682.
- Higgins, J., Thompson, S. & Spiegelhalter, D. 2009. A re-evaluation of random-effects meta-analysis, *Journal of the Royal Statistical Society: Series A* **172**: 137-159.

- Karabatsos, G. & Walker, S. 2012. Adaptive-modal Bayesian nonparametric regression, *Electronic Journal of Statistics* **6**: 2038-2068.
- Konstantopoulos, S. 2007. Introduction to meta-analysis, in J. Osbourne (ed.), *Best Practices in Quantitative Methods*, Sage, Thousand Oaks, CA, pp. 177-194.
- Konstantopoulos, S. 2011. Fixed effects and variance components estimation in three-level meta-analysis, *Research Synthesis Methods* **2**: 61-76.
- Laud, P. & Ibrahim, J. 1995. Predictive model selection, *Journal of the Royal Statistical Society, Series B* **57**: 247-262.
- Lo, A. 1984. On a class of Bayesian nonparametric estimates, *Annals of Statistics* **12**: 351-357.
- Louis, T. & Zelterman, D. 1994. Bayesian approaches to research synthesis, in H. Cooper & L. Hedges (eds), *The Handbook of Research Synthesis*, Russell Sage Foundation, New York, pp. 411-422.
- McCullagh, P. 1980. Regression models for ordinal data (with discussion), *Journal of the Royal Statistical Society, Series B* **42**: 109-142.
- McLachlan, G. & Peel, D. 2000. *Finite Mixture Models*, John Wiley and Sons, New York.
- Moffitt, T. 1993. Adolescence-limited and life-course persistent antisocial behavior: A developmental taxonomy, *Psychological Review* **100**: 674-701.
- Moffitt, T. 2005. The new look of behavioral genetics in developmental psychopathology: Gene-environment interplay in antisocial behaviors, *Psychological Bulletin* **131**: 533-554.
- Müller, P. & Quintana, F. 2004. Nonparametric Bayesian data analysis, *Statistical Science* **19**: 95-110.
- Nam, I.-S., Mengersen, K. & Garthwaite, P. 2003. Multivariate meta-analysis, *Statistics in Medicine* **22**: 2309-2333.
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. & R Development Core Team 2010. *Nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1-97*.
- R Development Core Team 2012. *R: A Language and Environment for Statistical Computing*, R Foundation for Statistical Computing, Vienna, Austria.
- Raudenbush, S. & Bryk, A. 2002. *Hierarchical Linear Models: Applications and Data Analysis Methods (Second Edition)*, Sage, Thousand Oaks, CA.
- Stevens, J. & Taylor, A. 2009. Hierarchical dependence in meta-analysis, *Journal of Educational and Behavioral Statistics* **34**: 46-73.
- Sutton, A. 2000. *Methods for meta-analysis in medical research*, John Wiley, Chichester.

- Talbott, B., Karabatsos, G. & Zurheide, J. 2012. *Sensitivity of raters and the assessment of heritability of antisocial behavior: A meta-analysis of twin, adoption, and sibling studies*, Technical report, University of Illinois-Chicago.
- Thompson, S. & Higgins, J. 2002. How should meta-regression analyses be undertaken and interpreted?, *Statistics in Medicine* **21**: 1559-1573.
- Thompson, S. & Sharp, S. 1999. Explaining heterogeneity in meta-analysis: A comparison of methods, *Statistics in Medicine* **18**: 2693-2708.
- Walker, H., Ramsey, E. & Gresham, F. 2004. *Antisocial behavior in schools: Evidence-based practices*, Wadsworth Publishing, Belmont, CA.

Appendix. Behavioral-Genetic Studies

The following numerical list provides citations for all 29 behavioral-genetic studies, which were subject to a meta-analysis. The list is given in the order of publication year. Each item of the list presents the identification number assigned to the given study. The identification numbers of the studies are also shown in Figure 3.

1. Scarr, S. 1966. Genetic factors in activity motivation. *Child Development* **37**: 663-673.
2. Owen, D.R., & Sines, J.O. 1970. Heritability of personality in children. *Behavior Genetics* **1**: 235-248.
3. Loehlin, J.C., & Nichols, R.C. 1976. *Heredity, environment, and personality*. Austin: University of Texas Press.
4. Rowe, D.C. 1983. Biometrical genetic models of self-reported delinquent behavior: A twin study. *Behavior Genetics* **13**: 473-489.
5. Graham, P., & Stevenson, J. 1985. A twin study of genetic influences on behavioral deviance. *Journal of the American Academy of Child Psychiatry* **24**: 33-41.
6. Lytton, H., Watts, D., & Dunn, B.E. 1988. Stability of genetic determination from age 2 to age 9: A longitudinal twin study. *Social Biology* **35**: 62-73.
7. Eaves, L.J., Silberg, J.L., Meyer, J M., Maes, H.H., Simonoff, E., Pickles, A., Rutter, M., Neale, M.C., Reynolds, C.A., Erikson, M.T., Heath, A.C., Loeber, R., Truett, K.R., & Hewitt, J.K. 1997. Genetics and developmental psychopathology: 2. The main effects of genes and environment on behavioral problems in the Virginia twin study of adolescent behavioral development. *Journal of Child Psychology and Psychiatry* **38**: 965-980.
8. Gjone, H., & Stevenson, J. 1997. The association between internalizing and externalizing behavior in childhood and early adolescence: Genetic or environmental common influences? *Journal of Abnormal Child Psychology* **25**: 277-286.

9. Eley, T.C., Lichtenstein, P., & Stevenson, J. 1999. Sex differences in the etiology of aggressive and nonaggressive antisocial behavior: Results from two twin studies. *Child Development* **70**: 155-168.
10. Hudziak, J.J., Rudiger, L.P., Neale, M.C., Heath, A.C., & Todd, R.D. 2000. A twin study of inattentive, aggressive, and anxious/depressed behaviors. *Journal of the American Academy of Child and Adolescent Psychiatry* **39**: 469-476.
11. Arseneault, L., Moffitt, T.E., Caspi, A., Taylor, A., Rijdsdijk, F.V., & Jaffee, S.R. et al. 2003. Strong genetic effects on cross-situational antisocial behavior among 5-year-old children according to mothers, teachers, examiner-observers, and twins' self reports. *Journal of Child Psychology and Psychiatry*, **44**: 832-848.
12. Eley, T.C., Lichtenstein, P., & Moffitt, T.E. 2003. A longitudinal behavioral genetic analysis of the etiology of aggressive and nonaggressive antisocial behavior. *Development and Psychopathology*, **15**: 383-402.
13. Van Beijsterveldt, C.E.M., Bartels, M., Hudziak, J.J., & Boomsma, D.I. 2003. Causes of stability of aggression from early childhood to adolescence: A longitudinal genetic analysis in Dutch twins. *Behavior Genetics* **33**: 591-605.
14. Vierikko, E., Pulkkinen, L., Kaprio, J., Viken, R., & Rose, R.J. 2003. Sex differences in genetic and environmental effects on aggression. *Aggressive Behavior* **29**, 55-68.
15. Derks, E.M., Hudziak, J.J., Van Beijsterveldt, C.E.M., Dolan, C.V., & Boomsma, D.I. 2004. A study of genetic and environmental influences on maternal and paternal CBCL syndrome scores in a large sample of 3-year-old Dutch Twins. *Behavior Genetics* **34**: 571-583.
16. Gregory, A.M., Eley, T.C., & Plomin, R. 2004. Exploring the association between anxiety and conduct problems in a large sample of twins aged 2-4. *Journal of Abnormal Child Psychology* **32**: 111-122.
17. Van Beijsterveldt, C.E.M., Verhulst, F.C., Molenaar, P.C.M., & Boomsma, D.I. 2004. The genetic basis of problem behavior in 5-year-old Dutch twin pairs. *Behavior Genetics* **34**: 229-242.
18. Blonigen, D.M., Hicks, B.M., Krueger, R.F., Patrick, C.J., & Iacono, W.G. 2005. Psychopathic personality traits: Heritability and genetic overlap with internalizing and externalizing psychopathology. *Psychological Medicine* **35**, 637-648.
19. Burt, S.A., McGue, M., Krueger, R.F., & Iacono, W.G. 2005. Sources of covariation among the child-externalizing disorders: Informant effects and the shared environment. *Psychological Medicine* **35**, 1133-1144.
20. Haberstick, B.C., Schmitz, S., Young, S.E., & Hewitt, J.K. 2005. Contributions of genes and environments to stability and change in externalizing and internalizing problems during elementary and middle school. *Behavior Genetics* **35**: 381-396.

21. Ligthart, L., Bartels, M., Hoekstra, R.A., Hudziak, J.J., & Boomsma, D.I. 2005. Genetic contributions to subtypes of aggression. *Twin Research and Human Genetics* **8**: 483-491.
22. Saudino, K.J., Ronald, A., & Plomin, R. 2005. The etiology of behavior problems in 7-year-old twins: Substantial genetic influence and negligible shared environmental influence for parent ratings and ratings by same and different teachers. *Journal of Abnormal Child Psychology*, **33**: 113-130.
23. Tuvblad, C., Grann, M., & Lichtenstein, P. 2006. Heritability for adolescent antisocial behavior differs with socioeconomic status: gene-environment interaction. *Journal of Child Psychology and Psychiatry* **47**: 734-743.
24. Vierikko, E., Pulkkinen, L., Kaprio, J., & Rose, R.J. 2006. Genetic and environmental sources of continuity and change in teacher-rated aggression during early adolescence. *Aggressive Behavior* **32**: 308-320.
25. Van Hulle, C.A., Lemery-Chalfant, K., & Goldsmith, H.H. 2007. Genetic and environmental influences on socio-emotional behavior in toddlers: an initial twin study of the infant-toddler social and emotional assessment. *Journal of Child Psychology and Psychiatry* **48**: 1014-1024.
26. Ball, H.A., Arseneault, L., Taylor, A., Maughan, B., Caspi, A., & Moffitt, T.E. 2008. Genetic and environmental influences on victims, bullies, and bully-victims in childhood. *Journal of Child Psychology and Psychiatry*, **49**: 104-112.
27. Button, T.M. M., Lau, J.Y.F., Maughan, B., & Eley, T.C. 2008. Parental punitive discipline, negative life events and gene-environment interplay in the development of externalizing behavior. *Psychological Medicine* **38**: 29-39.
28. Saudino, K.J., Carter, A.S., Purper-Ouakil, D., & Gorwood, P. 2008. The etiology of behavioral problems and competencies in very young twins. *Journal of Abnormal Psychology* **117**: 48-62.
29. Tuvblad, C., Raine, A., Zheng, M., & Baker, L.A. 2009. Genetic and environmental stability differs in reactive and proactive aggression. *Aggressive Behavior* **35**: 437-452.

Figure Captions

Figure 1. Kernel density estimate of the effect size (heritability).

Figure 2. Effect size density $f(y|\mathbf{x}) = f(y|\mathbf{x}; \hat{\sigma} = 1; \zeta)$ for $\sigma_\omega = 1/20, 1/2, 1, 2$, given $\mathbf{x}^\top \boldsymbol{\beta} = .7$ and given sampled values of (μ_j, σ_j^2) .

Figure 3. Left panel: Heritability estimate, and its variance (+), for each of the 71 independent samples of MZ and DZ twin comparisons, provided by 29 studies. A circle refers to females, square refers to males. Also, red refers to mother rater, black to teacher rater, blue to self-rater, magenta to observer rater, and green to mixed raters. The study identification number is located in the given square or circle, and this numbering is according to the order of publication date. Right panel: Bayesian estimate of the heritability distribution, over the universe of studies. Med: median; Var: variance; Skew: skewness; Kurt: kurtosis.

Figure 4. For each of the 6 panels, the estimated posterior predictive median (solid line) and interquartile range (dashed lines) of antisocial behavior heritability (effect size), given values of a covariate, while controlling for all other covariates by fixing their values to zero. Mom: mother rater; Dad: father rater; Teacher: teacher rater; Self: self rater; Observer: observer rater.







