arXiv:1302.0130v2 [astro-ph.CO] 31 Mar 2014 [arXiv:1302.0130v2 \[astro-ph.CO\] 31 Mar 2014](http://arxiv.org/abs/1302.0130v2)

Symmetries and Consistency Relations in the Large Scale Structure of the Universe

A. Kehagias^a and A. Riotto^b

^a Physics Division, National Technical University of Athens, 15780 Zografou Campus, Athens, Greece

 b Department of Theoretical Physics and Center for Astroparticle Physics (CAP) 24 quai E. Ansermet, CH-1211 Geneva 4, Switzerland

Abstract

We study the symmetries enjoyed by the Newtonian equations of motion of the non-relativistic dark matter fluid coupled to gravity which give rise to the phenomenon of gravitational instability. We also discuss some consistency relations involving the soft limit of the $(n + 1)$ -correlator functions of matter and galaxy overdensities.

1 Introduction

There is no doubt that symmetries play a crucial role in high energy physics [1]. They allow, for instance, to derive Ward identities among correlation functions which remain valid even after renormalization [2]. Symmetries are also relevant in the cosmological setting and have been the subject of a recent and intense activity. They are particularly useful in characterizing the properties of the cosmological perturbations generated by an inflationary stage [3]. During inflation the de Sitter isometry group acts as conformal group on \mathbb{R}^3 when the fluctuations are on super-Hubble scales. In such a regime, the SO(1,4) isometry of the de Sitter background is realized as conformal symmetry of the flat \mathbb{R}^3 sections and correlators are constrained by conformal invariance [4–7]. This applies in the case in which the cosmological perturbations are generated by light scalar fields other than the inflaton (the field that drives inflation). In the opposite case in which the inflationary perturbations originate from only one degree of freedom, conformal consistency relations among the inflationary correlators have also been recently investigated [8–11]. The fluctuations in single-field inflation are Goldstone bosons of a spontaneously broken dilation symmetry. Being non-linearly realized, the broken symmetry is still respected in Ward identities and leads to a relation between the variation of the n-point function of the comoving curvature perturbation ζ under dilation and the squeezed limit of the $(n + 1)$ -point function [12, 13]. These identities will be extremely useful in discriminating among the various mechanisms for the generation of the cosmological perturbations. For instance, the detection of a sizable primordial three-point correlator in the squeezed limit would rule out all single-field models where inflation is driven by a single scalar field with canonical kinetic energy and an initial Bunch-Davies vacuum.

When perturbations re-enter the horizon, they provide the seeds for the the large scale structure of the universe which grows via the gravitational instability [14]. At early epochs, the growth of the density perturbations can be described by linear perturbation theory and the perturbation Fourier modes evolve independently from one another, thus conserving the statistical properties of the primordial perturbations. When the perturbations become nonlinear, the coupling between the different Fourier modes become relevant, inducing nontrivial correlations that modify the statistical properties of the cosmological fields. At intermediate quasi-linear scales the evolution of matter may be described analytically by extending the standard perturbation theory [14], where one defines a series solution to the fluid equations in powers of the initial density field. The *n*-th order term of the series for the density contrast grows as the *n*-th power of the scale factor a (for a pressureless fluid), thus affecting its convergence properties.

The need for improving theoretical predictions for the next generation of very large galaxy surveys has spurred many efforts to go beyond the standard perturbation theory. For instance, the renormalized perturbation theory [15] reorganizes the perturbation expansion in terms of different fundamental objects, the so-called non-linear propagator and non-linear vertices, to improve the convergence. The renormalization group method [16] represents an alternative possibility where truncating the renormalization equation at the level of some n-point correlator leads to a solution that corresponds to the summation of an infinite class of perturbative corrections. Other methods have been proposed in Refs. [17–20].

The goal of this paper is to investigate the underlying symmetries of the Newtonian equations of motion which describe the gravitational instability and their consequences. Consider the non-relativistic fluid equations in the presence of gravity

$$
\frac{\partial \delta(\vec{x}, \tau)}{\partial \tau} + \nabla (1 + \delta(\vec{x}, \tau)) \vec{v}(\vec{x}, \tau) = 0, \tag{1.1}
$$

$$
\frac{\partial \vec{v}(\vec{x},\tau)}{\partial \tau} + \mathcal{H}(\tau)\vec{v}(\vec{x},\tau) + [\vec{v}(\vec{x},\tau)\cdot\nabla]\vec{v}(\vec{x},\tau) = \nabla\Phi(\vec{x},\tau),\tag{1.2}
$$

$$
\nabla^2 \Phi(\vec{x}, \tau) = \frac{3}{2} \Omega \mathcal{H}^2(\tau) \delta(\vec{x}, \tau), \qquad (1.3)
$$

where we have denoted by \vec{x} the comoving spatial coordinates, $\tau = \int dt/a$ the conformal time, a the scale factor in the FRW metric, $\mathcal{H} = d \ln(a)/d\tau$ the conformal expansion rate and, in addition, $\delta(\vec{x}, \tau) =$ $(\rho(\vec{x}, \tau)/\bar{\rho}-1)$ is the overdensity over the mean matter density density $\bar{\rho}$, $\vec{v}(\vec{x}, \tau)$ is the velocity of the fluid away from the Hubble flow and $\Phi(\vec{x}, \tau)$ is the gravitational potential due to density fluctuations. Finally $\Omega = 8\pi G \bar{\rho} a^2 / 3\mathcal{H}^2$ is the density parameter.

We are interested in exploring possible symmetries of these equations (which are expected to correctly describe the gravitational instability when vorticity and multi streaming are not present). In other words, we are looking for which (possibly) non-linearly realized transformations of the coordinates and fields

$$
\tau \rightarrow \tau',
$$

\n
$$
\vec{x} \rightarrow \vec{x}',
$$

\n
$$
\delta(\vec{x}, \tau) \rightarrow Z_{\delta}(\vec{x}', \tau')\delta(\vec{x}', \tau') + \xi_{\delta}(\vec{x}', \tau'),
$$

\n
$$
\vec{v}(\vec{x}, \tau) \rightarrow Z_{\vec{v}}(\vec{x}', \tau')\vec{v}(\vec{x}', \tau') + \xi_{\vec{v}}(\vec{x}', \tau'),
$$

\n
$$
\Phi(\vec{x}, \tau) \rightarrow Z_{\Phi}(\vec{x}', \tau')\Phi(\vec{x}', \tau') + \xi_{\Phi}(\vec{x}', \tau'),
$$

the Newtonian equations (1.1-1.3) describing the gravitational instability are invariant. The symmetries of the equations depend on the explicit form of $\mathcal{H}(\tau)$. We will explore two particular cases, a namely matter-dominated era and ΛCDM.

Very much similar to what happens in quantum field theory, these symmetries lead to relations among the correlation functions which are valid at any order in perturbation theory. As such, they might be represent useful consistency checks of the various analytical approaches and possibly testable by observations.

One example of such a symmetry and its utility is represented by the Galilean symmetry: being the Newtonian equations those of a classical non-relativistic field theory, they are invariant under Galilean transformations. This invariance is indeed the underlying reason for the cancellation of the leading infrared divergences in the computation of the matter power spectrum observed in various perturbative schemes to arbitrary number of loops [21–23].

In fact, we will show that the Galilean transformations are only a special case of a more general set of transformations enjoyed by the non-relativistic fluid equations coupled to gravity. This general set contains also the acceleration transformations where in the new system of coordinates the observer is uniformly accelerated. Furthermore, if the universe is matter-dominated, the gravitational instability equations are invariant also under a Lifshitz scaling symmetry with a generic exponent z [24]. The discussion of these symmetries will be performed in section 2 under the assumption that the universe is matter-dominated, while section 3 is devoted to the case in which the universe contains a non vanishing cosmological constant. A more elegant discussion of the symmetries will be given in section 4 based on the derivation of the four-dimensional non-relativistic fluid equations from the dimensional reduction of a five-dimensional scalar field theory. Finally, section 5 will contain a discussion of the consistency relations which may obtained on the correlators from the underlying symmetries.

2 Symmetries during matter-domination

Let us start from the simplest possibility, a matter-dominated universe. In such a case the scale factor a scales like τ^2 and we have $\mathcal{H} = 2/\tau$. We will also assume $\Omega = 1$. Under these assumptions, the fluid equations Eqs. $(1.1-1.3)$ are obviously invariant under $SO(3)$ space rotations as well as under the Galilean boosts

$$
\tau' = \tau, \quad \vec{x}' = \vec{x} + \vec{u}\,\tau,\tag{2.1}
$$

$$
\delta'(\vec{x},\tau) = \delta(\vec{x}',\tau'),\tag{2.2}
$$

$$
\vec{v}'(\vec{x},\tau) = \vec{v}(\vec{x}',\tau') - \vec{u},\tag{2.3}
$$

$$
\Phi'(\vec{x},\tau) = \Phi(\vec{x}',\tau') - \frac{2}{\tau}\vec{u}\cdot\vec{x},\tag{2.4}
$$

where \vec{u} is a constant three-dimensional vector. Indeed, since

$$
\frac{\partial}{\partial \tau} = \frac{\partial}{\partial \tau'} + \vec{u} \cdot \nabla', \quad \nabla = \nabla', \tag{2.5}
$$

we have, for example for Eq. (1.2),

$$
0 = \frac{\partial \vec{v}'(\vec{x}, \tau)}{\partial \tau} + \mathcal{H}(\tau) \vec{v}'(\vec{x}, \tau) + [\vec{v}'(\vec{x}, \tau) \cdot \nabla] \vec{v}'(\vec{x}, \tau) - \nabla \Phi'(\vec{x}, \tau)
$$

\n
$$
= \frac{\partial \vec{v}(\vec{x}', \tau')}{\partial \tau'} + \mathcal{H}(\tau') \vec{v}(\vec{x}', \tau') - \mathcal{H}(\tau) \vec{u} + [\vec{v}(\vec{x}', \tau') \cdot \nabla'] \vec{v}(\vec{x}', \tau') - \nabla \Phi'(\vec{x}, \tau)
$$

\n
$$
= -\mathcal{H}(\tau) \vec{u} + \nabla' \Phi(\vec{x}', \tau') - \nabla \Phi'(\vec{x}, \tau). \tag{2.6}
$$

Therefore, Eq. (1.2) is invariant if

$$
\nabla \Phi'(\vec{x}, \tau) = \nabla' \Phi(\vec{x}', \tau') - \mathcal{H}(\tau)\vec{u}
$$
\n(2.7)

or

$$
\Phi'(\vec{x},\tau) = \Phi(\vec{x}',\tau') - \mathcal{H}(\tau)\,\vec{u}\cdot\vec{x}.\tag{2.8}
$$

This is just Eq. (2.4) for $\mathcal{H} = 2/\tau$. Using the transformation (2.8) into (1.3), we get

$$
0 = \nabla^2 \Phi'(\vec{x}, \tau) - \frac{3}{2} \Omega \mathcal{H}^2(\tau) \delta'(\vec{x}, \tau) = \nabla'^2 \Phi(\vec{x}', \tau') - \frac{3}{2} \Omega \mathcal{H}^2(\tau) \delta'(\vec{x}, \tau)
$$

=
$$
\frac{3}{2} \Omega \mathcal{H}(\tau')^2 \delta(\vec{x}', \tau') - \frac{3}{2} \Omega \mathcal{H}^2(\tau) \delta'(\vec{x}, \tau),
$$
 (2.9)

from which the transformation (2.2) follows immediately. Finally, a simple inspection of Eq. (1.1) shows that it is also invariant under (2.2-2.4), thus recovering the well-known invariance of the fluid equations under Galilean transformations. In the infinitesimal form, the transformations (2.1-2.4) take the form (with $\delta \vec{u} = \vec{b}$)

$$
\delta_{\rm g}\tau = 0, \quad \delta_{\rm g}\vec{x} = \vec{b}\,\tau,\tag{2.10}
$$

$$
\delta_{g}\delta(\vec{x},\tau) = \tau \vec{b} \cdot \nabla \delta(\vec{x},\tau), \qquad (2.11)
$$

$$
\delta_{g}\vec{v}(\vec{x},\tau) = \tau \vec{b} \cdot \nabla \vec{v}(\vec{x},\tau) - \vec{b},\tag{2.12}
$$

$$
\delta_{g}\Phi(\vec{x},\tau) = \tau \vec{b} \cdot \nabla \Phi(\vec{x},\tau) - \frac{2}{\tau} \vec{b} \cdot \vec{x}.\tag{2.13}
$$

The Newtonian equations are also invariant under the acceleration transformations

$$
\tau' = \tau, \quad \vec{x}' = \vec{x} + \frac{1}{2}\vec{a}\tau^2,\tag{2.14}
$$

$$
\delta'(\vec{x},\tau) = \delta(\vec{x}',\tau'),\tag{2.15}
$$

$$
\vec{v}'(\vec{x},\tau) = \vec{v}(\vec{x}',\tau') - \vec{a}\tau,\tag{2.16}
$$

$$
\Phi'(\vec{x},\tau) = \Phi(\vec{x}',\tau') - 3\vec{a}\cdot\vec{x},\tag{2.17}
$$

where \vec{a} is a constant three-dimensional vector. Indeed, using

$$
\frac{\partial}{\partial \tau} = \frac{\partial}{\partial \tau'} + \tau \, \vec{a} \cdot \nabla', \quad \nabla = \nabla',\tag{2.18}
$$

we get

$$
0 = \frac{\partial \vec{v}'(\vec{x}, \tau)}{\partial \tau} + \mathcal{H}(\tau) \vec{v}'(\vec{x}, \tau) + [\vec{v}'(\vec{x}, \tau) \cdot \nabla] \vec{v}'(\vec{x}, \tau) - \nabla \Phi'(\vec{x}, \tau) = \frac{\partial \vec{v}(\vec{x}', \tau')}{\partial \tau'} - \vec{a} + \mathcal{H}(\tau') \vec{v}(\vec{x}', \tau') - \mathcal{H}(\tau') \tau \vec{a} + [\vec{v}(\vec{x}', \tau') \cdot \nabla'] \vec{v}(\vec{x}', \tau') - \nabla \Phi'(\vec{x}, \tau) = -\vec{a} - \mathcal{H}(\tau') \vec{a} \tau + \nabla' \Phi(\vec{x}', \tau') - \nabla \Phi'(\vec{x}, \tau)
$$
(2.19)

Therefore Eq. (1.2) is invariant if

$$
\nabla \Phi'(\vec{x}, \tau) = \nabla' \Phi(\vec{x}', \tau') - \vec{a} - \mathcal{H}\vec{a}\tau
$$
\n(2.20)

or

$$
\Phi'(\vec{x}, \tau) = \Phi(\vec{x}', \tau') - \vec{a} \cdot \vec{x} - \tau \mathcal{H} \vec{a} \cdot \vec{x}.\tag{2.21}
$$

Again, for $\mathcal{H} = 2/\tau$, Eq. (2.16) follows. From the Poisson equation we get

$$
0 = \nabla^2 \Phi'(\vec{x}, \tau) - \frac{3}{2} \Omega \mathcal{H}^2(\tau) \delta'(\vec{x}, \tau) = \nabla'^2 \Phi(\vec{x}', \tau') - \frac{3}{2} \Omega \mathcal{H}^2(\tau) \delta'(\vec{x}, \tau)
$$

\n
$$
= \frac{3}{2} \Omega \mathcal{H}^2(\tau') \delta(\vec{x}', \tau') - \frac{3}{2} \Omega \mathcal{H}^2(\tau) \delta'(\vec{x}, \tau), \tag{2.22}
$$

so that $\delta(\vec{x}, \tau)$ is an invariant scalar as Eq. (2.15) shows. Finally, using Eq. (2.15-2.17), we find that (1.1) is also invariant. The corresponding infinitesimal form of the acceleration transformation is (with $\delta \vec{a} = \vec{b}$)

$$
\delta_{\mathbf{a}}\tau = 0 \quad \delta_{\mathbf{a}}\vec{x} = \frac{1}{2}\vec{b}\,\tau^2,\tag{2.23}
$$

$$
\delta_a \delta(\vec{x}, \tau) = \frac{1}{2} \tau^2 \vec{b} \cdot \nabla \delta(\vec{x}, \tau), \qquad (2.24)
$$

$$
\delta_{a}\vec{v}(\vec{x},\tau) = \tau\vec{b}\cdot\nabla\vec{v}(\vec{x},\tau) - \vec{b}\tau,
$$
\n(2.25)

$$
\delta_{\mathbf{a}}\Phi(\vec{x},\tau) = \frac{1}{2}\tau^2 \vec{b} \cdot \nabla \Phi(\vec{x},\tau) - 3\vec{b} \cdot \vec{x}.\tag{2.26}
$$

In fact, Galilean and acceleration transformations are special cases of a most general transformation of the non-relativistic fluid equations

$$
\tau' = \tau, \quad \vec{x}' = \vec{x} + \vec{n}(\tau), \tag{2.27}
$$

$$
\delta'(\vec{x},\tau) = \delta(\vec{x}',\tau'),\tag{2.28}
$$

$$
\vec{v}'(\vec{x},\tau) = \vec{v}(\vec{x}',\tau') - \dot{\vec{n}}(\tau),\tag{2.29}
$$

$$
\Phi'(\vec{x},\tau) = \Phi(\vec{x}',\tau') - \left(\ddot{\vec{n}}(\tau) + \frac{2}{\tau}\dot{\vec{n}}(\tau)\right) \cdot \vec{x}.\tag{2.30}
$$

Note that we are using the dot to denote differentiation with respect to conformal time τ . In order to prove the invariance of the fluid equations under the transformations (2.27-2.30), we note that

$$
\frac{\partial}{\partial \tau} = \frac{\partial}{\partial \tau'} + \dot{\vec{n}} \cdot \nabla', \quad \nabla = \nabla',\tag{2.31}
$$

which means that the operator

$$
D_{\tau} = \frac{\partial}{\partial \tau} + \vec{v}(\vec{x}, \tau) \cdot \nabla \tag{2.32}
$$

is invariant under the transformation (2.27) . From Eq. (1.2) we therefore get

$$
0 = \frac{\partial \vec{v}'(\vec{x}, \tau)}{\partial \tau} + \mathcal{H}(\tau) \vec{v}'(\vec{x}, \tau) + [\vec{v}'(\vec{x}, \tau) \cdot \nabla] \vec{v}'(\vec{x}, \tau) - \nabla \Phi'(\vec{x}, \tau)
$$

\n
$$
= \frac{\partial \vec{v}(\vec{x}', \tau')}{\partial \tau'} - \ddot{\vec{n}} + \mathcal{H}(\tau') \vec{v}(\vec{x}', \tau') - \mathcal{H}(\tau') \dot{\vec{n}} + [\vec{v}(\vec{x}', \tau') \cdot \nabla'] \vec{v}(\vec{x}', \tau') - \nabla \Phi'(\vec{x}, \tau)
$$

\n
$$
= -\ddot{\vec{n}} - \mathcal{H}(\tau') \dot{\vec{n}} + \nabla' \Phi(\vec{x}', \tau') - \nabla \Phi'(\vec{x}, \tau),
$$
\n(2.33)

from which we deduce the transformation

$$
\nabla \Phi'(\vec{x}, \tau) = \nabla' \Phi(\vec{x}', \tau') + \ddot{\vec{n}} + \mathcal{H}(\tau')\dot{\vec{n}} \tag{2.34}
$$

or

$$
\Phi'(\vec{x},\tau) = \Phi(\vec{x}',\tau') - (\dot{\vec{n}} + \mathcal{H}(\tau')\dot{\vec{n}})\cdot\vec{x}.\tag{2.35}
$$

It is a straightforward exercise to check the invariance of Eqs. (1.1) and (1.3) under the general transformations (2.28),(2.29), and (2.30). Their infinitesimal form are given by (with $\delta \vec{n} = \vec{b}$)

$$
\delta_{\text{gen}} \tau = 0, \quad \delta_{\text{gen}} \vec{x} = \vec{b}(\tau), \tag{2.36}
$$

$$
\delta_{\text{gen}}\delta(\vec{x},\tau) = \vec{b}(\tau) \cdot \nabla \delta(\vec{x},\tau),\tag{2.37}
$$

$$
\delta_{\text{gen}}\vec{v}(\vec{x},\tau) = \vec{b}(\tau) \cdot \nabla \vec{v}(\vec{x},\tau) - \dot{\vec{b}}(\tau),\tag{2.38}
$$

$$
\delta_{\text{gen}}\Phi(\vec{x},\tau) = \vec{b}(\tau) \cdot \nabla \Phi(\vec{x},\tau) - \left(\ddot{\vec{b}}(\tau) + \frac{2}{\tau}\dot{\vec{b}}(\tau)\right) \cdot \vec{x}.\tag{2.39}
$$

Fluid equations during the matter-dominated period are also invariant under Lifshitz scalings of the form

$$
\tau' = \lambda^z \tau, \quad \vec{x}' = \lambda \vec{x}, \tag{2.40}
$$

$$
\delta'(\vec{x},\tau) = \delta(\vec{x}',\tau'),\tag{2.41}
$$

$$
\vec{v}'(\vec{x},\tau) = \lambda^{z-1}\vec{v}(\vec{x}',\tau'),\tag{2.42}
$$

$$
\Phi'(\vec{x},\tau) = \lambda^{2(z-1)} \Phi(\vec{x}',\tau') \tag{2.43}
$$

for a generic Lifshitz weight z. Indeed, using

$$
\frac{\partial}{\partial \tau} = \lambda^z \frac{\partial}{\partial \tau'}, \quad \nabla = \lambda \nabla', \tag{2.44}
$$

we get

$$
0 = \frac{\partial \vec{v}'(\vec{x}, \tau)}{\partial \tau} - \vec{a} + \mathcal{H}(\tau) \vec{v}'(\vec{x}, \tau) + [\vec{v}'(\vec{x}, \tau) \cdot \nabla] \vec{v}'(\vec{x}, \tau) - \nabla \Phi'(\vec{x}, \tau)
$$

= $\lambda^{2z-1} \frac{\partial \vec{v}(\vec{x}', \tau')}{\partial \tau'} + \lambda^{z-1} \mathcal{H}(\tau' \lambda^{-z}) \vec{v}(\vec{x}', \tau') + \lambda^{2z-1} [\vec{v}(\vec{x}', \tau') \cdot \nabla'] \vec{v}(\vec{x}', \tau') - \nabla \Phi'(\vec{x}, \tau). \tag{2.45}$

We see immediately that only when H scales like $1/\tau$, the above equation is covariant. In this case we get

$$
0 = \lambda^{2z-1} \left(\frac{\partial \vec{v}(\vec{x}', \tau')}{\partial \tau'} - \frac{2}{\tau'} \vec{v}(\vec{x}', \tau') + [\vec{v}(\vec{x}', \tau') \cdot \nabla'] \vec{v}(\vec{x}', \tau') \right) - \nabla \Phi'(\vec{x}, \tau) = \lambda^{2z-1} \nabla' \Phi(\vec{x}', \tau') - \nabla \Phi'(\vec{x}, \tau),
$$
\n(2.46)

that is

$$
\nabla \Phi'(\vec{x}, \tau) = \lambda^{2z - 1} \nabla' \Phi(\vec{x}', \tau')
$$
\n(2.47)

or

$$
\Phi'(\vec{x}, \tau) = \lambda^{2z - 2} \Phi(\vec{x}', \tau'). \tag{2.48}
$$

Using this transformation in Poisson equation we find

$$
0 = \nabla^2 \Phi'(\vec{x}, \tau) + \frac{3}{\tau^2} \Omega \delta'(\vec{x}, \tau) = \lambda^{2z} \nabla'^2 \Phi(\vec{x}', \tau') - \lambda^z \frac{3}{\tau'^2} \delta'(\vec{x}, \tau)
$$

=
$$
-\lambda^{2z} \frac{3}{\tau'^2} \Omega \delta(\vec{x}', \tau') - \lambda^{2z} \frac{3}{\tau'^2} \delta'(\vec{x}, \tau),
$$
 (2.49)

from which the transformation (2.41) immediately follows. Then, it is straightforward to check the invariance of Eq. (1.1) under the transformations $(2.41),(2.42)$ and (2.43) . The infinitesimal forms of the Lifshitz scalings read (with λ to be intended as an infinitesimal parameter)

$$
\delta_{\rm l}\tau = z\lambda\tau\,,\quad \delta_{\rm l}\vec{x} = \lambda\vec{x},\tag{2.50}
$$

$$
\delta_1 \delta(\vec{x}, \tau) = \lambda \, \vec{x} \cdot \nabla \delta(\vec{x}, \tau) + z \, \lambda \, \tau \frac{\partial}{\partial \tau} \delta(\vec{x}, t), \tag{2.51}
$$

$$
\delta_{\rm I}\vec{v}(\vec{x},\tau) = \lambda \,\vec{x} \cdot \nabla \vec{v}(\vec{x},\tau) + z\lambda\tau \frac{\partial}{\partial \tau}\vec{v}(\vec{x},t) + (z-1)\lambda \vec{v}(\vec{x},\tau),\tag{2.52}
$$

$$
\delta_1 \Phi(\vec{x}, \tau) = \lambda \, \vec{x} \cdot \nabla \Phi(\vec{x}, \tau) + z \lambda \tau \frac{\partial}{\partial \tau} \Phi(\vec{x}, \tau) + 2(z - 1) \lambda \Phi(\vec{x}, \tau). \tag{2.53}
$$

Notice that in the linear regime, when the gravitational potential $\Phi(\vec{x}, \tau)$ does not evolve as function of time, $\Phi(\vec{x}, \tau)$ has a Lifshitz weight equal to $2(z - 1)$. Going beyond the linear order and writing $\Phi(\vec{x}, \tau) = \sum_{n\geq 0} \Phi^{(n)}(\vec{x}) \tau^{2n}$, one finds that $\Phi^{(n)}(\vec{x})$ has Lifshitz weight $(2n+1)z - 2$.

3 Symmetries in ΛCDM

In this section we investigate the symmetries if the universe, besides a matter component, is characterized by the presence of a non vanishing cosmological constant. Having learnt that in the matter-dominated case there is a set of transformations which contain Galilean and acceleration transformations as particular case, let us consider the more general set of transformations

$$
\tau' = \tau, \quad \vec{x}' = \vec{x} + \vec{n}(T), \tag{3.1}
$$

where

$$
T(\tau) = \frac{1}{a(\tau)} \int^{\tau} d\eta a(\eta).
$$
 (3.2)

Following the same steps of the previous section, it can be checked that the non-relativistic fluid equations are invariant under

$$
\delta'(\vec{x},\tau) = \delta(\vec{x}',\tau'),\tag{3.3}
$$

$$
\vec{v}'(\vec{x},\tau) = \vec{v}(\vec{x}',\tau') - \dot{\vec{n}}(T),\tag{3.4}
$$

$$
\Phi'(\vec{x},\tau) = \Phi(\vec{x}',\tau') + \left(\mathcal{H}\dot{\vec{n}}(T) + \ddot{\vec{n}}(T)\right) \cdot \vec{x},\tag{3.5}
$$

where again we remind the reader that the dot denotes differentiation with respect to conformal time τ . The proof proceeds as in Eq. (2.33) after using (2.31). These transformation include the Galilean boosts for which

$$
\vec{n}(T) = T\vec{u}, \ \frac{\partial T}{\partial \tau} = 1 - \mathcal{H}T, \qquad \frac{\partial^2 T}{\partial \tau^2} = -\dot{\mathcal{H}}T - \mathcal{H}(1 - \mathcal{H}T). \tag{3.6}
$$

Eqs. (3.3-3.5) become

$$
\delta'(\vec{x},\tau) = \delta(\vec{x}',\tau'),\tag{3.7}
$$

$$
\vec{v}'(\vec{x},\tau) = \vec{v}(\vec{x}',\tau') - \vec{u}(1-\mathcal{H}T),\tag{3.8}
$$

$$
\Phi'(\vec{x}, \tau) = \Phi(\vec{x}', \tau') - T \frac{\partial \mathcal{H}}{\partial \tau} \vec{u} \cdot \vec{x}.\tag{3.9}
$$

Acceleration transformations are also included as a special case. It is enough to take

$$
\vec{n}(T) = \frac{1}{2}\vec{u}\,T^2,\tag{3.10}
$$

and Eqs. (3.3-3.5) become

$$
\delta'(\vec{x},\tau) = \delta(\vec{x}',\tau'),\tag{3.11}
$$

$$
\vec{v}'(\vec{x},\tau) = \vec{v}(\vec{x}',\tau') - \frac{1}{2}\vec{u}\,T(1-\mathcal{H}T),\tag{3.12}
$$

$$
\Phi'(\vec{x},\tau) = \Phi(\vec{x}',\tau') + \left((1 - \mathcal{H}T)^2 - T^2 \frac{\partial \mathcal{H}}{\partial \tau} \right) \vec{u} \cdot \vec{x}.\tag{3.13}
$$

One can also easily verify that the Lifshitz scaling is no longer a symmetry of the non-relativistic equations when the universe is of the ΛCDM type.

4 Fluid dynamics by dimensional reduction and symmetries

The purpose of this section is to show how the symmetries of the non-relativistic equations describing the gravitational instability may be though of as symmetries of higher dimensional theory on which dimensional reduction is applied [25–27]. We are going to do it step by step, starting from a flat geometery and adding gravity and the expansion of the universe later on. Consider first the five-dimensional Bargmann spacetime with metric

$$
ds^2 = 2dt d\xi + d\vec{x}^2,\tag{4.1}
$$

where ξ is a null direction. The dynamics of a free complex scalar field ϕ in such a space-time is described by the Lagrangian

$$
\mathcal{L}_5 = -\partial_\mu \phi^\dagger \partial^\mu \phi,\tag{4.2}
$$

which can explicitly be written as

$$
\mathcal{L}_5 = -\partial_t \phi^\dagger \partial_\xi \phi - \partial_\xi \phi^\dagger \partial_t \phi - |\nabla \phi|^2. \tag{4.3}
$$

For $\phi(t, \xi, \vec{x})$ of the form

$$
\phi(t,\xi,\vec{x}) = e^{im\xi}\psi(t,\vec{x})\tag{4.4}
$$

the Lagrangian becomes

$$
\mathcal{L}_5 = im\left(\psi^{\dagger}\partial_t\psi - \psi\partial_t\psi^{\dagger}\right) - |\nabla\psi|^2. \tag{4.5}
$$

The corresponding equation of motion for $\psi(t, \vec{x})$ is

$$
-\nabla^2 \psi(t, \vec{x}) = 2im\partial_t \psi(t, \vec{x}),\tag{4.6}
$$

which is the well-known Schrodinger equation. The theory (4.5) is invariant under the full Schrödinger group Sch(3), that is the symmetry group of a free non-relativistic theory. In general, Sch(d) is defined as the subgroup of $SO(2,d+2)$ that leaves invariant the momentum along a null direction. Note that $Sch(3)$ contains Lifshitz scaling symmetry with $z = 2$, as one may directly verify.

We may also express $\psi(t, \vec{x})$ as

$$
\psi(t,\vec{x}) = \sqrt{\frac{\rho(t,\vec{x})}{m}} e^{im\omega(t,\vec{x})},\tag{4.7}
$$

where $\rho(t, \vec{x})$ and $\omega(t, \vec{x})$ are real functions and they will play the role of the matter density and the velocity potential, respectively. The Lagrangian (4.3) written in terms of $\rho(t, \vec{x})$ and $\omega(t, \vec{x})$ becomes

$$
\mathcal{L}_5 = -2m\rho \,\partial_t \omega - \frac{1}{4m\rho} (\nabla \rho)^2 - m\rho (\nabla \omega)^2.
$$
\n(4.8)

If we now take the limit of the momentum m along the ξ -direction to infinity, $m \to \infty$, we get

$$
\mathcal{L}_5 = -2m\left(\rho \,\partial_t \omega + \frac{1}{2}\rho \, (\nabla \omega)^2\right) + \mathcal{O}\left(\frac{1}{m}\right). \tag{4.9}
$$

The theory described by the Lagrangian (4.9) is invariant under the Galilei group

$$
t \to t' = t, \quad \vec{x} \to \vec{x}' = \vec{x} + \vec{v}t,
$$
\n
$$
(4.10)
$$

if ρ and ω transform as

$$
\rho \to \rho' = \rho, \quad \omega \to \omega' = \omega - \vec{v} \cdot \vec{x} - \frac{1}{2}v^2t. \tag{4.11}
$$

Furthermore, the theory is invariant under the arbitrary Lifshitz scaling $t' = \lambda^z t$ and $\vec{x}' = \lambda \vec{x}$ if we assign the following weights

$$
\omega \to \lambda^{2-z}\omega \,, \quad \rho \to \lambda^{-5+z}\rho \tag{4.12}
$$

to $\rho(t, \vec{x})$ and $\omega(t, \vec{x})$. Note that for finite m, the theory (4.8) is invariant under the full Sch(3) group which, however, is broken down to the Galilei group plus arbitrary Lifshitz scalings in the $m \to \infty$ limit.

Let us now include gravity in the theory (4.2) . For this, we perturb the flat metric (4.1) as

$$
ds^{2} = -2\Phi(t, \vec{x})dt^{2} + 2(1 + \Phi(t, \vec{x})) dt d\xi + (1 + \Phi(t, \vec{x})) d\vec{x}^{2},
$$
\n(4.13)

where $\Phi(t, \vec{x})$ is the Newtonian potential. Then the dynamics is described by the Lagrangian

$$
\mathcal{L}_5 = \frac{M_*^3}{2} R - \partial_\mu \phi^\dagger \partial^\mu \phi,\tag{4.14}
$$

where M_* plays the role of the five-dimensional reduced Planck mass. Replacing the ansatz (4.4,4.7) for the scalar $\phi(t, \vec{x})$ we have

$$
\mathcal{L}_5 = -\left(-im\left(\psi^{\dagger}\partial_t\psi - \psi\partial_t\psi^{\dagger}\right) + |\nabla\psi|^2\Phi + m^2\psi^{\dagger}\psi\Phi + \frac{1}{2}M_*^3(\nabla\Phi)^2\right) \tag{4.15}
$$

and, after using the ansatz (4.7), the Lagrangian becomes in terms of the fields $\rho(t, \vec{x})$ and $\omega(t, \vec{x})$

$$
\mathcal{L}_g = -\left(2m\rho \partial_t \omega + \frac{1}{4m\rho} (\nabla \rho)^2 + m\rho (\nabla \omega)^2 + 2m\rho \Phi + \frac{1}{2} M_*^3 (\nabla \Phi)^2\right). \tag{4.16}
$$

Upon taking the $m \to \infty$ limit, we finally get

$$
\mathcal{L}_5 = -2m \left\{ \rho \left(\partial_t \omega + \frac{1}{2} \left(\nabla \omega \right)^2 + \Phi \right) + M_p^2 (\nabla \Phi)^2 \right\} + \mathcal{O}\left(\frac{1}{m}\right),\tag{4.17}
$$

which describes a fluid in gravitational field after the following identification

$$
M_*^3 = 4mM_{\rm p}^2,\tag{4.18}
$$

where M_p is the four-dimensional reduced Planck mass. The term of order $\mathcal{O}(1/m)$ we are neglecting is $(\nabla \rho)^2/(m\rho)$ and therefore the Lagrangian we have derived is valid as long as $|\nabla \rho/m\rho \nabla \omega| \ll 1$. Note that the theory (4.17) is invariant only under the Galilei group as the presence of the gravitational potential breaks the the arbitrary Lifshitz scaling invariance.

Finally, we introduce the cosmological expansion. As we shall see, this is essential in restoring the arbitrary Lifshitz scaling invariance. Consider the metric

$$
ds^{2} = -2\Phi(t, \vec{x})dt^{2} + 2(1 + \Phi(t, \vec{x})) dt d\xi + a^{2}(t) (1 + \Phi(t, \vec{x})) d\vec{x}^{2}.
$$
 (4.19)

Upon dimensional reduction with $\phi(t, \xi, \vec{x})$ of the form given in Eqs. (4.4,4.7) we obtain in the $m \to \infty$ limit

$$
S = \int dt d^3 \vec{x} \left\{ -2m \left\{ a^3 \rho \left(\partial_t \omega + \frac{1}{2a^2} |\nabla \omega|^2 + \Phi \right) + aM_\text{p}^2 (\nabla \Phi)^2 \right\} + \mathcal{O}\left(\frac{1}{m}\right) \right\}.
$$
 (4.20)

The equations of motions for $\rho(t, \vec{x})$, $\omega(t, \vec{x})$ and $\Phi(t, \vec{x})$ are then precisely the Newtonian equations describing the fluid gravitationally instability of a non-relativistic fluid with density ρ and velocity $\vec{v}(t, \vec{x}) = \nabla \omega(t, \vec{x})/a$, using cosmic time.

Let us try now to identify symmetries of the action (4.20) . It can easily be checked that the following transformations

$$
\delta_{\text{gen}}t = 0 \quad \delta_{\text{gen}}\vec{x} = \vec{n}(t),\tag{4.21}
$$

$$
\delta_{\text{gen}}\omega(t,\vec{x}) = -a^2 \frac{\partial \vec{n}(t)}{\partial t} \cdot \vec{x} + \vec{n} \cdot \nabla \omega,\tag{4.22}
$$

$$
\delta_{\text{gen}}\Phi(t,\vec{x}) = \frac{\partial}{\partial t}\left(a^2 \frac{\partial \vec{n}(t)}{\partial t}\right) \cdot \vec{x} + \vec{n} \cdot \nabla \Phi \tag{4.23}
$$

$$
\delta_{\text{gen}}\rho(t,\vec{x}) = \vec{n} \cdot \nabla \rho,\tag{4.24}
$$

leave (4.20) invariant. Of course, the above transformation includes Galilean boosts ($\vec{n} = \vec{n}_0 \tau$, \vec{n}_0 constant vector) as well as acceleration $(\vec{n} = \vec{n}_0 \tau^2)$. In addition, there is also a Lifshitz symmetry for the action (4.20). Assuming the rescalings

$$
t \to t' = \lambda_t t,\tag{4.25}
$$

$$
a \to a' = \lambda_a a,\tag{4.26}
$$

$$
\vec{x} \to \vec{x}' = \lambda \vec{x},\tag{4.27}
$$

$$
\omega \to \omega' = \lambda_{\omega}\omega,\tag{4.28}
$$

$$
\rho \to \rho' = \lambda_{\rho} \rho,\tag{4.29}
$$

$$
\Phi \to \Phi' = \lambda_{\Phi} \Phi, \tag{4.30}
$$

and that the scale factor a is a homogeneous function of time (so that $a \to a' = \lambda_a a$), it is easy to verify that (4.20) is invariant for

$$
\lambda_{\omega} = \lambda^{(z-1)/2}, \quad \lambda_{\Phi} = \lambda^{2-2z}, \quad \lambda_{\rho} = \lambda^{5-5z}, \quad \lambda_a = \lambda^{(3z-5)/2}, \quad \lambda_t = \lambda^{(5z-5)/2}
$$
\n(4.31)

We have taken conformal time to change as $\tau \to \tau' = \lambda^z \tau$, so that $\lambda_a = \lambda_t \lambda^{-z}$. Note that when a is not a homogeneous function of time, as in the ΛCDM case form example, the Lifshitz scaling is not a symmetry.

5 Consistency relations and conclusions

As the gravitational instability equations posses a set of symmetries, it is natural to ask what are their consequences. The popular choice for the initial conditions in cosmology is that of a (nearly) Gaussian random field which is statistically homogeneous and isotropic in space. The distribution in any one realization is therefore not expected to be satisfying, e.g., the Lifshitz scaling we have discussed for the matter-dominated period. The symmetries should be intended to play a role only at the statistical level in the sense that the correlators of a given observable, for example the matter density contrast, should have the same statistical properties of its transform under a given symmetry. This means that the correlators should satisfy appropriate Ward identities, reflecting the invariance under the given symmetries. For example, for the n-point connected correlators of the gravitational potential at equal time

$$
G_{\Phi}^{(n)}(\vec{x}_1, \vec{x}_2, \dots, \vec{x}_n, \tau) = \left\langle \Phi(\vec{x}_1, \tau), \Phi(\vec{x}_2, \tau) \cdots \Phi(\vec{x}_n, \tau) \right\rangle_c,
$$
\n(5.1)

the Ward identity follows from

$$
0 = \delta G_{\Phi}^{(n)}(\vec{x}_1, \vec{x}_2, \dots, \vec{x}_n, \tau) = \left\langle \delta \Phi(\vec{x}_1, \tau), \Phi(\vec{x}_2, \tau) \cdots \Phi(\vec{x}_n, \tau) \right\rangle_c + \text{permutations.}
$$
 (5.2)

For the Galilean and acceleration symmetry it is easy to see that Eq. (5.2) is satisfied for $G_c^{(n)}$ = $G_c^{(n)}(\vec{x}_{ij},\tau)$ where $\vec{x}_{ij} = (\vec{x}_i - \vec{x}_j)$. In addition, rotational invariance imposes that $G_c^{(n)} = G_c^{(n)}(x_{ij},\tau)$, where $x_{ij} = |\vec{x}_i - \vec{x}_j|$. The Lifshitz scaling symmetry on the other hand gives

$$
\left(2n(z-1) + z\tau \frac{\partial}{\partial \tau} + \sum_{i < j} \vec{x}_{ij} \cdot \nabla_{ij}\right) G_{\Phi}^{(n)}(x_{ij}, \tau) = 0. \tag{5.3}
$$

For the two-point correlator one finds therefore that $G_c^{(2)}$ should satisfy

$$
\left(4(z-1) + z\,\tau\frac{\partial}{\partial\tau} + \vec{x}_{12} \cdot \nabla_{12}\right) G_{\Phi}^{(2)}(x_{12}, \tau) = 0. \tag{5.4}
$$

Solving the above equation, we get that

$$
G_{\Phi}^{(2)}(\vec{x}_1, \vec{x}_2, \tau) = \frac{1}{x_{12}^{4(z-1)}} \mathcal{F}_{\Phi}\left(\frac{\tau}{x_{12}^z}\right). \tag{5.5}
$$

where \mathcal{F}_{Φ} is a function of only τ/x_{12}^z . Similarly, for the three-point correlator we get the equation

$$
\left(6(z-1) + z\,\tau\frac{\partial}{\partial\tau} + \vec{x}_{12}\cdot\nabla_{12} + \vec{x}_{13}\cdot\nabla_{13} + \vec{x}_{23}\cdot\nabla_{23}\right)G_{\Phi}^{(3)}(\vec{x}_1, \vec{x}_2, \vec{x}_3, \tau) = 0.\tag{5.6}
$$

The solution of the above equation which is also invariant under the permutation $\vec{x}_i \rightarrow \vec{x}_j$, $(i, j = 1, 2, 3)$ is

$$
G_{\Phi}^{(3)}(\vec{x}_1, \vec{x}_2, \vec{x}_3, \tau) = \frac{1}{x_{12}^{2(z-1)} x_{13}^{2(z-1)} x_{23}^{2(z-1)}} \mathcal{G}_{\Phi}\left(\frac{\tau}{x_{12}^z}, \frac{\tau}{x_{13}^z}, \frac{\tau}{x_{23}^z}\right). \tag{5.7}
$$

Similarly, one finds that the two-point connected correlator of the matter over density at equal time must satisfy the following relation because of the Lifshitz scaling symmetry [14, 24]

$$
G_{\delta}^{(2)}(\vec{x}_1, \vec{x}_2, \tau) = \left\langle \delta(\vec{x}_1, \tau), \delta(\vec{x}_2, \tau) \right\rangle_c = \mathcal{F}_{\delta} \left(\frac{\tau}{x_{12}^z} \right). \tag{5.8}
$$

If one matches this behavior with the linear one at early times $G_{\delta r}^{(2)}$ $\delta_L^{(2)}(\vec{x}_1, \vec{x}_2, \tau) \sim \tau^4/x_{12}^{3+n}$, one finds $z = 4/(n+3)$. This means that the power spectrum in momentum space $\Delta_{\delta}(k, \tau) = (k^3/2\pi^2)P_{\delta}(k, \tau)$ at any order in perturbation theory must be a function of $\Delta_{\delta}(k,\tau) = \Delta_{\delta}(k/k_*(\tau))$, where $k_*(\tau) \sim \tau^{-4/(n+3)}$ and it can be interpreted as the momentum scale at which perturbations become non-linear at any given time.

Another, and maybe more interesting, consequence of the symmetries we have discussed in the ΛCDM model is to produce consistency relations which correlators should satisfy in the squeezed limit, that is in the case in which one of the modes is a long wavelength mode and therefore it may be assumed to evolve in the linear regime. This allows to relate $(n + 1)$ -correlation functions containing a soft mode to the to n-point correlation functions of the short modes.

Consider, for instance, the *n*-point correlation function of short modes density contrasts $\langle \delta_{\vec{k}_1} \delta_{\vec{k}_2} \cdots \delta_{\vec{k}_n} \rangle$. The points are supposed to be contained in a sphere of radius R much smaller than the long wavelength mode of size $\sim 1/q$ and centered at the origin of the coordinates. According to what we have discussed in section 3, the non-relativistic equation of motions are invariant under the generic transformation $\tau' \to t$ and $\vec{x} \to \vec{x} + \vec{n}(T(\tau))$. This means that we can generate a long wavelength mode for the velocity perturbation $\vec{v}_L(\tau, \vec{0})$ just by choosing properly the vector $\vec{n}(\tau)$

$$
\vec{n}(\tau) = -\int^{\tau} d\eta \,\vec{v}_L(\eta, \vec{0}) + \mathcal{O}(qRv_L^2). \tag{5.9}
$$

In other words, the correlator of the short wavelength modes in the background of the long wavelength mode perturbation should satisfy the relation

$$
\left\langle \delta(\tau_1, \vec{x}_1) \delta(\tau_2, \vec{x}_2) \cdots \delta(\tau_n, \vec{x}_n) \right\rangle_{v_L} = \left\langle \delta(\tau'_1, \vec{x}'_1) \delta(\tau'_2, \vec{x}'_2) \cdots \delta(\tau'_n, \vec{x}'_n) \right\rangle. \tag{5.10}
$$

This is nothing else that the statement that the effect of a physical long wavelength velocity perturbation onto the short modes should be indistinguishable from the long wavelength mode velocity generated by the transformation $\delta x^i = n^i(\tau)$. In momentum space one therefore obtains

$$
\left\langle \delta_{\vec{q}}(\tau)\delta_{\vec{k}_1}(\tau_1)\cdots\delta_{\vec{k}_n}(\tau_n) \right\rangle_{q\to 0} = \left\langle \delta_{\vec{q}}(\tau)\left\langle \delta_{\vec{k}_1}(\tau_1)\cdots\delta_{\vec{k}_n}(\tau_n) \right\rangle_{v_L} \right\rangle. \tag{5.11}
$$

The variation of the *n*-point correlator under such a transformation is given by

$$
\delta_n \left\langle \delta(\tau_1, \vec{x}_1) \cdots \delta(\tau_n, \vec{x}_n) \right\rangle = \int \frac{d^3 \vec{k}_1}{(2\pi)^3} \cdots \frac{d^3 \vec{k}_n}{(2\pi)^3} \left\langle \delta_{\vec{k}_1}(\tau_1) \cdots \delta_{\vec{k}_n}(\tau_n) \right\rangle
$$

$$
\times \sum_{a=1}^n \delta x_a^i (ik_a^i) e^{i(\vec{k}_1 \cdot \vec{x}_1 + \cdots \vec{k}_n \cdot \vec{x}_n)}
$$

$$
= \int \frac{d^3 \vec{k}_1}{(2\pi)^3} \cdots \frac{d^3 \vec{k}_n}{(2\pi)^3} \left\langle \delta_{\vec{k}_1}(\tau_1) \cdots \delta_{\vec{k}_n}(\tau_n) \right\rangle
$$

$$
\times \sum_{a=1}^n n^i(\tau_a) (ik_a^i) e^{i(\vec{k}_1 \cdot \vec{x}_1 + \cdots \vec{k}_n \cdot \vec{x}_n)}.
$$
(5.12)

Then we find that

$$
\left\langle \delta_{\vec{q}}(\tau) \delta_{\vec{k}_{1}}(\tau_{1}) \cdots \delta_{\vec{k}_{n}}(\tau_{n}) \right\rangle_{q \to 0} = \left\langle \delta_{\vec{q}}(\tau) \left\langle \delta_{\vec{k}_{1}}(\tau_{1}) \cdots \delta_{\vec{k}_{n}}(\tau_{n}) \right\rangle_{v_{L}} \right\rangle
$$

$$
= i \sum_{a=1}^{n} \left\langle \delta_{\vec{q}}(\tau) n^{i}(\tau_{a}) \right\rangle k_{a}^{i} \left\langle \delta_{\vec{k}_{1}}(\tau_{1}) \cdots \delta_{\vec{k}_{n}}(\tau_{n}) \right\rangle.
$$
 (5.13)

As for the ΛCDM model we have

$$
\int^{\tau} d\eta \, \vec{v}_{\vec{q}}(\eta) = i \frac{q^i}{q^2} \int^{\tau} d\eta \, \mathcal{H} f(\eta) \, \delta_{\vec{q}}(\eta) = i \frac{q^i}{q^2} \int^{\tau} d\eta \, \mathcal{H} \, \frac{1}{\mathcal{H}} \frac{d \ln D_{+}(\eta)}{d\eta} \frac{D_{+}(\eta)}{D_{+}(\eta_{\text{in}})} \delta_{\vec{q}}(\eta_{\text{in}}) = i \frac{q^i}{q^2} \delta_{\vec{q}}(\tau), \tag{5.14}
$$

where D_+ is the linear growth factor, we finally get

$$
\left\langle \delta_{\vec{q}}(\tau)\delta_{\vec{k}_{1}}(\tau_{1})\cdots\delta_{\vec{k}_{n}}(\tau_{n}) \right\rangle_{q\to 0}^{\prime} = -P_{\delta_{L}}(q,\tau) \sum_{a=1}^{n} \frac{D_{+}(\tau_{a})}{D_{+}(\tau)} \frac{\vec{q} \cdot \vec{k}_{a}}{q^{2}} \left\langle \delta_{\vec{k}_{1}}(\tau_{1})\cdots\delta_{\vec{k}_{n}}(\tau_{n}) \right\rangle^{\prime},
$$
\n(5.15)

where the primes indicate that one should remove the Dirac delta's coming from the momentum conservation and $P_{\delta_L}(q,\tau) = (D_+(\tau)/D_+(\tau_{\rm min}))^2 P_{\delta}(q,\tau_{\rm in})$ is the linear matter power spectrum. Of course, similar consistency relations may be found involving the other quantities, the velocity perturbation and the gravitational potentials, in various combinations. Notice that, if the correlators are computed all at equal times, the right-hand side of eq. (5.15) vanishes by momentum conservation and the $1/q^2$ infrared divergence will not appear when calculating invariant quantities. For the three-point correlator, we obtain

$$
\left\langle \delta_{\vec{q}}(\tau)\delta_{\vec{k}_{1}}(\tau_{1})\delta_{\vec{k}_{2}}(\tau_{2}) \right\rangle_{q\to 0}^{\prime} = -P_{\delta}(q,\tau)\left(\frac{D_{+}(\tau_{1})}{D_{+}(\tau)} - \frac{D_{+}(\tau_{2})}{D_{+}(\tau)} \right) \frac{\vec{q} \cdot \vec{k}_{1}}{q^{2}} \left\langle \delta_{\vec{k}_{1}}(\tau_{1})\delta_{\vec{k}_{2}}(\tau_{2}) \right\rangle^{\prime}.
$$
 (5.16)

One can easily check this result holds at second-order in perturbation theory in the matter-dominated era when [14]

$$
\delta_{\vec{k}}^{(2)}(\tau) = \int \frac{\mathrm{d}^3 k_1}{(2\pi)^3} \frac{\mathrm{d}^3 k_2}{(2\pi)^3} \left[\frac{5}{7} + \frac{1}{2} (\vec{k}_1 \cdot \vec{k}_2) \frac{k_1^2 + k_2^2}{k_1^2 k_2^2} + \frac{2}{7} \frac{(\vec{k}_1 \cdot \vec{k}_2)^2}{k_1^2 k_2^2} \right] \delta^{(3)}(\vec{k} - \vec{k}_1 - \vec{k}_2) \, \delta_{\vec{k}_1}^{(1)}(\tau) \delta_{\vec{k}_2}^{(1)}(\tau). \tag{5.17}
$$

Indeed, in the squeezed limit

$$
\left\langle \delta_{\vec{q}}(\tau)\delta_{\vec{k}_{1}}(\tau_{1})\delta_{\vec{k}_{2}}(\tau_{2}) \right\rangle_{q \to 0} \simeq \left\langle \delta_{\vec{q}}^{(1)}(\tau)\delta_{\vec{k}_{1}}^{(2)}(\tau_{1})\delta_{\vec{k}_{2}}^{(1)}(\tau_{2}) \right\rangle_{q \to 0} + \left\langle \delta_{\vec{q}}^{(1)}(\tau)\delta_{\vec{k}_{1}}^{(1)}(\tau_{1})\delta_{\vec{k}_{2}}^{(2)}(\tau_{2}) \right\rangle_{q \to 0},
$$
(5.18)

and recalling that $\delta_{\vec{k}}(\tau) = (D_+(\tau)/D_+(\tau_{\rm min}))\delta_{\vec{q}}(\tau_{\rm in})$ with $D_+(\tau) = a(\tau)$, one recovers (5.16) by accounting for the appropriate permutations and taking the leading middle term in the squared parenthesis of Eq. $(5.18).$

Let us close with some comments. The consistency relation (5.15) is true at any order in perturbation theory. As such, it might represent a useful too to check the findings of the various schemes dealing analytically with the problem of structure formation beyond the standard perturbation theory. Nevertheless, the consistency relation might be of more practical use and tested in future galaxy surveys which are divided into multiple redshift bins. Indeed, cosmic tomography makes it possible to map out the three-dimensional distribution of mass and thus to observe correlators at different epochs. Of course, one needs the galaxy correlators and not the underlying dark matter ones. However, galaxies, once formed, obey the following equations on sub-Hubble scales

$$
\frac{\partial \delta_{\mathbf{g}}(\vec{x},\tau)}{\partial \tau} + \nabla (1 + \delta_{\mathbf{g}}(\vec{x},\tau)) \vec{v}_{\mathbf{g}}(\vec{x},\tau) = 0, \tag{5.19}
$$

$$
\frac{\partial \vec{v}_{g}(\vec{x},\tau)}{\partial \tau} + \mathcal{H}(\tau)\vec{v}_{g}(\vec{x},\tau) + [\vec{v}_{g}(\vec{x},\tau)\cdot\nabla]\vec{v}_{g}(\vec{x},\tau) = \nabla\Phi(\vec{x},\tau),\tag{5.20}
$$

$$
\nabla^2 \Phi(\vec{x}, \tau) = \frac{3}{2} \Omega \mathcal{H}^2(\tau) \delta(\vec{x}, \tau), \tag{5.21}
$$

where $\delta_{g}(\vec{x}, \tau)$ and $\vec{v}_{g}(\vec{x}, \tau)$ are the galaxy overdensity and peculiar velocity, respectively, while $\delta(\vec{x}, \tau)$ is the underlying dark matter overdensity. Following the same steps in sections 2 and 3, one can show that the set of equations (5.19-5.21) are invariant under the transformations

$$
\delta'_{g}(\vec{x},\tau) = \delta_{g}(\vec{x}',\tau'),\tag{5.22}
$$

$$
\vec{v}_{\mathrm{g}}'(\vec{x},\tau) = \vec{v}_{\mathrm{g}}(\vec{x}',\tau') - \dot{\vec{n}}(T),\tag{5.23}
$$

$$
\Phi'(\vec{x},\tau) = \Phi(\vec{x}',\tau') + \left(\mathcal{H}\dot{\vec{n}}(T) + \ddot{\vec{n}}(T)\right) \cdot \vec{x}.\tag{5.24}
$$

This is true even if the we do not assume $\vec{v}_{g}(\vec{x}, \tau) = \vec{v}(\vec{x}, \tau)$, that is the galaxy peculiar velocity is unbiased, as is often done. Therefore the consistency relation (5.15) should be true also for the galaxy overdensities, independently of the bias between $\delta_{\rm g}(\vec{x}, \tau)$ and $\delta(\vec{x}, \tau)$.

Finally, it is possible that gravity at large distances is modified by, for instance, a Yukawa-like modification of the Poisson equation

$$
(\nabla^2 - a^2 m^2) \Phi(\vec{x}, \tau) = \frac{3}{2} \Omega \mathcal{H}^2(\tau) \delta(\vec{x}, \tau), \qquad (5.25)
$$

where $1/m$ defines some new infrared scale where gravity gets modified (the factor a^2 in front of m^2 is such that the Yukawa correction becomes small in the early universe, but some other time dependence is possible). In such a case, the symmetries (3.3-3.5) fail in general due to the mass term. The only case (5.25) could be invariant is to choose a specific $\vec{n}(T)$ which will leave Φ itself invariant. This possibility is provided by

$$
\vec{n}(\tau) = \vec{n}_0 + \vec{n}_1 \int^{\tau} \frac{d\eta}{a(\eta)},\tag{5.26}
$$

where \vec{n}_0, \vec{n}_1 are constant vectors. In this case, the induced transformation for the velocity $\vec{v}(\tau, \vec{x})$ field will be

$$
\vec{v}'(\vec{x},\tau) = \vec{v}(\vec{x}',\tau') + \frac{1}{a(\tau)}\vec{n}_1,\tag{5.27}
$$

corresponding to a decaying mode. As the latter is of limited importance, we can safely say that the symmetries we have discussed in this paper do not hold any longer and a violation of the consistency relations might be a signal of modification of gravity.

Acknowledgments

A.R. is supported by the Swiss National Science Foundation (SNSF), project 'The non-Gaussian Universe" (project number: 200021140236).

Note added

When completing this work, we became aware of a similar work by M. Peloso and M. Pietroni. Our results, when overlap is possible, agree with theirs. We thank them for useful discussions.

References

- [1] See, for instance, 'spects of Symmetry: Selected Erice Lectures" by S. Coleman, Cambridge University press (1988).
- [2] Y. Takahashi, Nuovo Cimento, Ser. 10, 6 (1957) 370; J.C. Ward, Phys. Rev. 78, (1950) 182.
- [3] D. H. Lyth and A. Riotto, Phys. Rept. 314, 1 (1999) [hep-ph/9807278].
- [4] I. Antoniadis, P. O. Mazur and E. Mottola, JCAP 1209, 024 (2012) [arXiv:1103.4164 [gr-qc]].
- [5] P. Creminelli, Phys. Rev. D 85, 041302 (2012) [arXiv:1108.0874 [hep-th]].
- [6] A. Kehagias and A. Riotto, Nucl. Phys. B 864, 492 (2012) [arXiv:1205.1523 [hep-th]].
- [7] A. Kehagias and A. Riotto, Nucl. Phys. B 868, 577 (2013) [arXiv:1210.1918 [hep-th]].
- [8] P. Creminelli, J. Norena and M. Simonovic, JCAP 1207, 052 (2012) [arXiv:1203.4595 [hep-th]].
- [9] K. Hinterbichler, L. Hui and J. Khoury, JCAP 1208, 017 (2012) [arXiv:1203.6351 [hep-th]].
- [10] V. Assassi, D. Baumann and D. Green, JCAP 1211, 047 (2012) [arXiv:1204.4207 [hep-th]].
- [11] V. Assassi, D. Baumann and D. Green, arXiv:1210.7792 [hep-th].
- [12] J. M. Maldacena, JHEP 0305, 013 (2003) [astro-ph/0210603].
- [13] P. Creminelli and M. Zaldarriaga, JCAP 0410, 006 (2004) [astro-ph/0407059].
- [14] For a review, see F. Bernardeau, S. Colombi, E. Gaztanaga and R. Scoccimarro, Phys. Rept. 367, 1 (2002) [astro-ph/0112551].
- [15] M. Crocce and R. Scoccimarro, Phys. Rev. D 73, 063519 (2006) [astro-ph/0509418].
- [16] S. Matarrese and M. Pietroni, JCAP 0706, 026 (2007) [astro-ph/0703563].
- [17] P. Valageas, Astron. Astrophys. 421, 23 (2004) [astro-ph/0307008].
- [18] T. Matsubara, Phys. Rev. D 77, 063530 (2008) [arXiv:0711.2521 [astro-ph]].
- [19] R. Brustein and A. Riotto, JCAP 1111, 006 (2011) [arXiv:1105.4411 [astro-ph.CO]].
- [20] J. J. M. Carrasco, M. P. Hertzberg and L. Senatore, JHEP 1209, 082 (2012) [arXiv:1206.2926 [astro-ph.CO]].
- [21] B. Jain and E. Bertschinger, Astrophys. J. 456, 43 (1996) [astro-ph/9503025].
- [22] R. Scoccimarro and J. Frieman, Astrophys. J. Suppl. 105, 37 (1996) [astro-ph/9509047].
- [23] R. Scoccimarro and J. Frieman, Astrophys. J. 473, 620 (1996) [astro-ph/9602070].
- [24] M. Davis and P.J.E. Peebles, Astrpphys. Journal 34 (1977), 425.
- [25] C. Duval, G. Burdet, H. P. Kunzle and M. Perrin, Phys. Rev. D 31, 1841 (1985).
- [26] E. Prugovecki, Class. Quant. Grav. 4, 1659 (1987).
- [27] C. Duval, G. W. Gibbons and P. Horvathy, Phys. Rev. D 43, 3907 (1991) [hep-th/0512188].
- [28] P. A. Horvathy and P. -M. Zhang, Eur. Phys. J. C 65, 607 (2010) [arXiv:0906.3594 [physics.flu-dyn]].