

COMBINATORIAL RULES FOR THREE BASES OF POLYNOMIALS

COLLEEN ROSS AND ALEXANDER YONG

1. INTRODUCTION

We present combinatorial rules (one theorem and two conjectures) concerning three bases of $\text{Pol} = \mathbb{Z}[x_1, x_2, \dots]$.

Consider a basic question (studied for example in [L13+]):

How does one lift properties of the ring Λ of symmetric functions (and its Schur basis) to the entirety of Pol ?

The bases below lift the Schur polynomials. However, one wishes to analogize the relationship in Λ between rules for Schur polynomials and Littlewood-Richardson rules. For these bases, no rule has yet provided a parallel, explaining a desire for alternative forms.

First, we prove a “splitting” rule for the basis of *key polynomials* $\{\kappa_\alpha \mid \alpha \in \mathbb{Z}_{\geq 0}^\infty\}$, thereby establishing a new positivity theorem about them. This family was introduced by [D74] and first studied combinatorially in [LS89, LS90]. Combinatorial rules for their monomial expansion are known, see, e.g., [LS89, LS90, RS95, HHL09]. Our rule refines [RS95, Theorem 5(1)] and is compatible with the splitting rule [BKTY04, Corollary 3] for the basis of *Schubert polynomials* $\{\mathfrak{S}_w \mid w \in S_\infty\}$.

Second, we investigate a basis $\{\Omega_\alpha \mid \alpha \in \mathbb{Z}_{\geq 0}^\infty\}$ defined by [L01] that deforms the key basis. By extending the *Kohnert moves* of [K90] we conjecturally give the first combinatorial rule for the Ω -polynomials.

Third, in [K90], the Kohnert moves were used to conjecture the first combinatorial rule for Schubert polynomials (a proof was later presented in [W03]). Similarly, we use the extended Kohnert moves to give a conjecture for the basis of *Grothendieck polynomials* $\{\mathfrak{G}_w \mid w \in S_\infty\}$ [LS82]. This rule appears significantly different than earlier (proved) rules, such as those in [FK94, L01, BKTY05, LRS06].

1.1. Splitting key polynomials. Let S_∞ be the group of permutations of \mathbb{N} with finitely many non-fixed points. This acts on Pol by permuting the variables. Let s_i be the simple transposition interchanging x_i and x_{i+1} . The **divided difference operator** acts on Pol by

$$\partial_i = \frac{1 - s_i}{x_i - x_{i+1}}.$$

Define the **Demazure operator** by setting

$$\pi_i(f) = \partial_i(x_i \cdot f), \quad \text{for } f \in \text{Pol}.$$

For $\alpha = (\alpha_1, \alpha_2, \dots) \in \mathbb{Z}_{\geq 0}^\infty$, the **key polynomial** κ_α is

$$\kappa_\alpha = x_1^{\alpha_1} x_2^{\alpha_2} \cdots, \quad \text{if } \alpha \text{ is weakly decreasing.}$$

Date: February 1, 2013.

Otherwise,

$$\kappa_\alpha = \pi_i(\kappa_{\widehat{\alpha}}) \text{ where } \widehat{\alpha} = (\dots, \alpha_{i+1}, \alpha_i, \dots) \text{ and } \alpha_{i+1} > \alpha_i.$$

Since the leading term of κ_α is $x_1^{\alpha_1} x_2^{\alpha_2} \dots$, the key polynomials form a \mathbb{Z} -basis of Pol .

The key polynomials lift the Schur polynomials: when

$$(1) \quad \alpha = (\alpha_1, \alpha_2, \dots, \alpha_t, 0, 0, 0, \dots), \text{ where } \alpha_1 \leq \alpha_2 \leq \dots \leq \alpha_t, \text{ then}$$

$$(2) \quad \kappa_\alpha = s_{(\alpha_t, \dots, \alpha_2, \alpha_1)}(x_1, \dots, x_t).$$

A **descent** of α is an index i such that $\alpha_i \geq \alpha_{i+1}$; a **strict descent** is an index i such that $\alpha_i > \alpha_{i+1}$. Fix descents $d_1 < d_2 < \dots < d_k$ of α containing all strict descents of α . Since π_i symmetrizes $\{x_i, x_{i+1}\}$, κ_α is separately symmetric in each collection:

$$X_1 = \{x_1, x_2, \dots, x_{d_1}\}, X_2 = \{x_{d_1+1}, x_{d_1+2}, \dots, x_{d_2}\}, \dots, X_k = \{x_{d_{k-1}+1}, x_{d_{k-1}+2}, \dots, x_{d_k}\}.$$

(The variables $x_{d_k+1}, x_{d_k+2}, \dots$ do not appear in κ_α .) Therefore, uniquely:

$$(3) \quad \kappa_\alpha(X) = \sum_{\lambda^1, \dots, \lambda^k} \mathcal{E}_{\lambda^1, \dots, \lambda^k}^\alpha s_{\lambda^1}(X_1) \cdots s_{\lambda^k}(X_k),$$

where each λ^i is a partition. *A priori* one only knows $\mathcal{E}_{\lambda^1, \dots, \lambda^k}^\alpha \in \mathbb{Z}$.

Given $\alpha \in \mathbb{Z}_{\geq 0}^\infty$, there is a unique $w[\alpha] \in S_\infty$ such that $\text{code}(w[\alpha]) = \alpha$ (see, e.g., [M01, Proposition 2.1.2]). Here $\text{code}(w[\alpha]) \in \mathbb{Z}_{\geq 0}^\infty$ counts the number of boxes in columns of $\text{Rothe}(w[\alpha])$. We will need a special tableau coming from [S84, Section 4]:

The tableau $T[\alpha]$: Given $w[\alpha]$, $i_1 < i_2 < \dots < i_a$ in the first column of $T[\alpha]$ are given by having i_j be the largest descent position smaller than i_{j+1} in the permutation $ws_{i_a} s_{i_{a-1}} \cdots s_{i_1}$. The next column of $T[\alpha]$ is similarly determined, starting from $ws_{i_a} \cdots s_{i_1}$, etc.

An **increasing tableau** T of shape λ is a filling with strictly increasing rows and columns. (In fact, $T[\alpha]$ is an increasing tableau.) Let $\text{row}(T)$ be the reading word of T , obtained by reading the entries of T along rows, from right to left, and from top to bottom. Let $\min(T)$ be the smallest label in T . Finally, given a reduced word $\mathbf{a} = a_1 a_2 \dots a_m$, let $\text{EGLS}(\mathbf{a})$ be the output of the *Edelman-Greene correspondence* (see Section 2.1).

The following result shows $\mathcal{E}_{\lambda^1, \dots, \lambda^k}^\alpha \in \mathbb{Z}_{\geq 0}$. It is analogous to one on Schubert polynomials [BKTY04, Corollary 3] (which our proof uses).

Theorem 1.1. *The number $\mathcal{E}_{\lambda^1, \dots, \lambda^k}^\alpha$ counts sequences of increasing tableaux (T_1, T_2, \dots, T_k) where*

- T_i is of shape λ^i ;
- $\min T_1 > 0, \min T_2 > d_1, \min T_3 > d_2, \dots, \min T_k > d_{k-1}$; and
- $\text{row}(T_1) \cdot \text{row}(T_2) \cdots \text{row}(T_k)$ is a reduced word of $w[\alpha]$ such that $\text{EGLS}(\text{row}(T_1) \cdot \text{row}(T_2) \cdots \text{row}(T_k)) = T[\alpha]$.

When $d_j = j$ for all $j \geq 1$, Theorem 1.1 specializes to an instance of the monomial expansion formula [RS95, Theorem 5(1)] for κ_α (restated as Theorem 2.5 below). Also, when (1) holds, $k = 1, d_1 = t$ and thus Theorem 1.1 gives (2).

Example 1.2. The (strict) descents of $\alpha = (1, 3, 0, 2, 2, 1)$ are $d_1 = 2, d_2 = 5$, and

$$\begin{aligned} \kappa_{1,3,0,2,2,1} &= s_{3,2}(x_1, x_2) s_{2,1,1}(x_3, x_4, x_5) + s_{3,2}(x_1, x_2) s_{2,1}(x_3, x_4, x_5) s_1(x_6) \\ &\quad + s_{3,1}(x_1, x_2) s_{2,2}(x_3, x_4, x_5) s_1(x_6) + s_{3,1}(x_1, x_2) s_{2,2,1}(x_3, x_4, x_5). \end{aligned}$$

exhibits the claimed non-negativity of Theorem 1.1.

Also, $w[\alpha] = 2516743$ (one line notation) and $T[\alpha] = \begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & 5 & \\ \hline 4 & 6 & \\ \hline 5 & & \\ \hline 6 & & \\ \hline \end{array}$. Thus, $\mathcal{E}_{(3,2),(2,1,1),\emptyset}^{(1,3,0,2,2,1)} =$

$\mathcal{E}_{(3,2),(2,1),(1)}^{(1,3,0,2,2,1)} = \mathcal{E}_{(3,1),(2,2),(1)}^{(1,3,0,2,2,1)} = \mathcal{E}_{(3,1),(2,2,1),\emptyset}^{(1,3,0,2,2,1)} = 1$ are respectively witnessed by

$$\left(\begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & 5 & \\ \hline 4 & 6 & \\ \hline 5 & & \\ \hline 6 & & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 4 & 6 \\ \hline 5 & \\ \hline 6 & \\ \hline \end{array}, \emptyset \right), \left(\begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & 5 & \\ \hline 4 & 6 & \\ \hline 5 & & \\ \hline 6 & & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 4 & 6 \\ \hline 5 & \\ \hline 6 & \\ \hline \end{array}, \begin{array}{|c|} \hline 6 \\ \hline \end{array} \right), \left(\begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & & \\ \hline 4 & 5 & \\ \hline 5 & 6 & \\ \hline 6 & & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 4 & 5 \\ \hline 5 & 6 \\ \hline 6 & \\ \hline \end{array}, \begin{array}{|c|} \hline 6 \\ \hline \end{array} \right), \text{ and } \left(\begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & & \\ \hline 4 & 5 & \\ \hline 5 & 6 & \\ \hline 6 & & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 4 & 5 \\ \hline 5 & 6 \\ \hline 6 & \\ \hline \end{array}, \emptyset \right).$$

For example, for the leftmost sequence, $\text{EGLS}(43152 \cdot 6456 \cdot \emptyset) = T[\alpha]$ holds. \square

1.2. The Ω polynomials. A. Lascoux [L01] defines Ω_α for $\alpha = (\alpha_1, \alpha_2, \dots) \in \mathbb{Z}_{\geq 0}^\infty$ by replacing π_i in the definition of the key polynomials with the operator defined by

$$\tilde{\pi}_i(f) = \partial_i(x_i(1 - x_{i+1})f).$$

The initial condition is $\Omega_\alpha = x_1^{\alpha_1} x_2^{\alpha_2} x_3^{\alpha_3} \dots (= \kappa_\alpha)$, if α is weakly decreasing.

The **skyline diagram** is $\text{Skyline}(\alpha) = \{(i, y) : 1 \leq y \leq \alpha_i\} \subset \mathbb{N}^2$. Graphically, it is a collection of columns α_i high. For instance,

$$\text{Skyline}(1, 3, 0, 2, 2, 1) = \begin{pmatrix} \cdot & + & \cdot & \cdot & \cdot & \cdot \\ \cdot & + & \cdot & + & + & \cdot \\ + & + & \cdot & + & + & + \end{pmatrix}$$

Beginning with $\text{Skyline}(\alpha)$, **Kohnert's rule** [K90] generates diagrams D by sequentially moving any $+$ at the top of its column to the rightmost open position in its row and to its left. (The result of such a move need not be the skyline of any $\gamma \in \mathbb{Z}_{\geq 0}^\infty$.) Let $x^D = \prod_i x_i^{d_i}$ be the column weight where d_i is the number of $+$'s in column i of D . If the same D results from a different sequence of moves, it only counts once. Kohnert's theorem states $\kappa_\alpha = \sum x^D$, where the sum is over all such D . Extending this, we introduce:

The K -Kohnert rule: Each $+$ either moves as in Kohnert's rule, or stays in place *and* moves. In the latter case, mark the original position with a " g ". The g 's are unmovable, but a given $+$ treats g the same as other $+$'s when deciding if it can move, and to where. Diagrams with the same occupied positions but different arrangements of $+$'s and g 's are counted separately.

Example 1.3. Below, we give all K -Kohnert moves one step from D :

$$D = \begin{pmatrix} + & \cdot & g & + & \cdot \\ \cdot & + & + & + & + \end{pmatrix} \mapsto \begin{pmatrix} + & \cdot & g & + & \cdot \\ + & \cdot & + & + & + \end{pmatrix}, \begin{pmatrix} + & \cdot & g & + & \cdot \\ + & g & + & + & + \end{pmatrix}, \begin{pmatrix} + & + & g & \cdot & \cdot \\ \cdot & + & + & + & + \end{pmatrix},$$

$$\begin{pmatrix} + & + & g & g & \cdot \\ \cdot & + & + & + & + \end{pmatrix}, \begin{pmatrix} + & \cdot & g & + & \cdot \\ + & + & + & + & \cdot \end{pmatrix}, \begin{pmatrix} + & \cdot & g & + & \cdot \\ + & + & + & + & g \end{pmatrix}.$$

Let

$$J_\alpha^{(\beta)} = \sum \beta^{(\#g\text{'s appearing in } D)} x^D.$$

Conjecture 1.4. $J_\alpha^{(-1)} = \Omega_\alpha$.

Conjecture 1.4 has been checked by computer, for a wide range of cases up to α being of size 12, leaving us convinced. Clearly, $J_\alpha^{(0)} = \kappa_\alpha$, by Kohnert's theorem.

Example 1.5. Let $\alpha = (1, 0, 2)$. Then the diagrams contributing to $J_{(1,0,2)}$ are:

$$\begin{aligned} \text{Skyline}(1, 0, 2) = & \begin{pmatrix} \cdot & \cdot & + \\ + & \cdot & + \end{pmatrix}, \begin{pmatrix} \cdot & + & \cdot \\ + & \cdot & + \end{pmatrix}, \begin{pmatrix} + & \cdot & \cdot \\ + & \cdot & + \end{pmatrix}, \begin{pmatrix} + & \cdot & \cdot \\ + & + & \cdot \end{pmatrix}, \begin{pmatrix} \cdot & + & \cdot \\ + & + & \cdot \end{pmatrix}; \\ & \begin{pmatrix} + & g & \cdot \\ + & \cdot & + \end{pmatrix}, \begin{pmatrix} + & g & \cdot \\ + & + & \cdot \end{pmatrix}, \begin{pmatrix} + & \cdot & \cdot \\ + & + & g \end{pmatrix}, \begin{pmatrix} \cdot & + & \cdot \\ + & + & g \end{pmatrix}, \begin{pmatrix} \cdot & + & g \\ + & \cdot & + \end{pmatrix}, \begin{pmatrix} + & \cdot & g \\ + & \cdot & + \end{pmatrix}; \begin{pmatrix} + & g & \cdot \\ + & + & g \end{pmatrix}; \begin{pmatrix} + & g & g \\ + & \cdot & + \end{pmatrix}. \end{aligned}$$

Thus

$$\begin{aligned} J_{(1,0,2)} = & (x_1x_3^2 + x_1x_2x_3 + x_1^2x_3 + x_1^2x_2 + x_1x_2^2) \\ & - (x_1^2x_2x_3 + x_1^2x_2^2 + x_1^2x_2x_3 + x_1x_2^2x_3 + x_1x_2x_3^2 + x_1^2x_3^2) + (x_1^2x_2^2x_3 + x_1^2x_2x_3^2). \end{aligned}$$

The lowest degree homogeneous component of Ω_α is κ_α . Hence any $f \in \text{Pol}$ is a possibly infinite linear combination of the Ω_α 's. Finiteness is asserted in [L13+, Chapter 5]. We show in Section 4.2 that the J_α 's also form a (finite) basis.

1.3. Grothendieck polynomials. The **Grothendieck polynomial** [LS82] is defined using the **isobaric divided difference operator** whose action on $f \in \text{Pol}$ is given by:

$$\pi_i(f) = \partial_i((1 - x_{i+1})f).$$

Declare $\mathfrak{G}_{w_0}(X) = x_1^{n-1}x_2^{n-2}\cdots x_{n-1}$ where w_0 is the long element in S_n . Set $\mathfrak{G}_w(X) = \pi_i(\mathfrak{G}_{ws_i})$ if i is an ascent of w . The Grothendieck polynomials are known to lift $\{s_\lambda\}$ to Pol .

One has $\mathfrak{G}_w = \mathfrak{G}_w + (\text{higher degree terms})$. We now state the A. Kohnert's conjecture [K90] for \mathfrak{G}_w . The **Rothe diagram** is $\text{Rothe}(w) = \{(x, y) | y < w(x) \text{ and } x < w^{-1}(y)\} \subset [n] \times [n]$ (indexed so that the southwest corner is labeled $(1, 1)$). Starting with $\text{Rothe}(w)$, the Kohnert's rule generates diagrams D by applying the same rules as described for his rule for κ_α . Then $\mathfrak{G}_w = \sum x^D$; the sum is over all such D .

Analogously, we define

$$K_w^{(\beta)} = \sum_D \beta^{(\#g\text{'s appearing in } D)} \mathbf{x}^D$$

where the sum is over all diagrams D generated by the K -Kohnert rule. For example, if $w = 3142$ the diagrams contributing to $K_w^{(\beta)}$ are

$$\text{Rothe}(3142) = \begin{pmatrix} \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ + & \cdot & + & \cdot \\ + & \cdot & \cdot & \cdot \end{pmatrix}, \begin{pmatrix} \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ + & + & \cdot & \cdot \\ + & \cdot & \cdot & \cdot \end{pmatrix}, \begin{pmatrix} \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ + & + & g & \cdot \\ + & \cdot & \cdot & \cdot \end{pmatrix}.$$

and hence correspondingly, $K_{3142}^{(-1)} = (x_1^2x_3 + x_1^2x_2) - (x_1^2x_2x_3)$.

Conjecture 1.6. $K_w^{(-1)} = \mathfrak{G}_w$.

Note, $K_w^{(0)} = \mathfrak{G}_w$ is precisely Kohnert's conjecture. Conjecture 1.6 has been checked by computer for $n \leq 7$, and extensively for larger n . While Kohnert's rule for \mathfrak{G}_w is handy, it remains mysterious, even after [W03]. Conjectures 1.4 and 1.6 return to Kohnert's conjecture (albeit with a parameter β).

2. PROOF OF THEOREM 1.1

2.1. Reduced word combinatorics. Given $w \in S_n$, let

$$\mathbf{a} = (a_1, a_2, \dots, a_{\ell(w)}) \text{ and } \mathbf{i} = (i_1, i_2, \dots, i_{\ell(w)}).$$

In connection to [BJS93], we say the pair (\mathbf{a}, \mathbf{i}) is a **stable compatible pair for w** if $s_{a_1} \cdots s_{a_{\ell(w)}}$ is a reduced word for w and the following two conditions on \mathbf{i} hold:

- (cs.1) $1 \leq i_1 \leq i_2 \leq \cdots \leq i_{\ell(w)} < n$;
- (cs.2) $a_j < a_{j+1} \implies i_j < i_{j+1}$.

We will identify w with \mathbf{a} and the associated reduced word.

The **Edelman-Greene correspondence** [EG87] (the same basic construction is used in [LS82]) is a bijection

$$\text{EGLS} : (\mathbf{a}, \mathbf{i}) \mapsto (T, U)$$

where

- T is an increasing tableau such that $\text{row}(T)$ is a reduced word for \mathbf{a} ; and
- U is a semistandard tableau whose multiset of labels is precisely those in \mathbf{i} , and which has the same shape as T .

EGLS (column) insertion: Initially insert a_j into the leftmost column (of what will be T). If there are no labels strictly larger than a_j , we place a_j at the bottom of that column. If $a_j + t$ for $t > 2$ appears, we bump this $a_j + t$ to the next column to the right, replacing it with a_j . The same holds if $a_j + 1$ appears but not a_j . Finally, if both $a_j + 1$ and a_j already appear, we insert $a_j + 1$ into the next column to the right. Since \mathbf{a} is assumed to be reduced, the above enumerates all possibilities. Finally at step j a new box is created at a corner; in what will be U we place i_j .

Mildly abusing terminology, let $\text{EGLS}(\mathbf{a}) = T$.

2.2. Formulas for Schubert polynomials. A stable compatible pair (\mathbf{a}, \mathbf{i}) is a **compatible pair for w** if in addition to (cs.1) and (cs.2) the following holds:

- (cs.3) $i_j \leq a_j$.

Let $\text{Compatible}(w)$ be the set of compatible sequences for w . A rule of [BJS93] states:

$$(4) \quad \mathfrak{S}_w(X) = \sum_{(\mathbf{a}, \mathbf{i}) \in \text{Compatible}(w)} \mathbf{x}^{\mathbf{i}}.$$

A **descent** of w is an index j such that $w(j) > w(j+1)$. Let $\text{Descents}(w)$ be the set of descents of w . The following is [BKTY04, Corollary 3]:

Theorem 2.1. *Let $w \in S_n$ and suppose $\text{Descents}(w) \subseteq \{d_1 < d_2 < \dots < d_k\}$. Then*

$$(5) \quad \mathfrak{S}_w(X) = \sum_{\lambda^1, \dots, \lambda^k} c_{\lambda^1, \dots, \lambda^k}^w s_{\lambda^1}(X_1) \cdots s_{\lambda^k}(X_k)$$

where $c_{\lambda^1, \dots, \lambda^k}^w$ counts the number of tuples of increasing tableaux (T_1, \dots, T_k) where

- (i) T_i has shape λ^i ;
- (ii) $\min T_1 > 0, \min T_2 > d_1, \dots, \min T_k > d_{k-1}$; and
- (iii) $\text{row}(T_1) \cdots \text{row}(T_k)$ is a reduced word of w .

Assume for the remainder of the proof that

$$(6) \quad \text{Descents}(w) \subseteq \{d_1 < d_2 < \dots < d_k\}.$$

Let

$$\text{Tuples}(w) = \{(T_1, U_1), (T_2, U_2), \dots, (T_k, U_k)\}$$

where the T_i 's satisfy (i), (ii) and (iii) from Theorem 2.1, and each U_i is a semistandard tableau of shape λ^i using the labels $d_{i-1} + 1, d_{i-1} + 2, \dots, d_i$ ($d_0 = 0$).

2.3. "Splitting" the EGLS correspondence. Assuming (6) we define:

$$\Phi : \text{Compatible}(w) \rightarrow \text{Tuples}(w).$$

Description of Φ (using EGLS): Uniquely split $(\mathbf{a}, \mathbf{i}) \in \text{Compatible}$ as follows

$$(7) \quad ((\mathbf{a}^{(1)}, \mathbf{i}^{(1)}), (\mathbf{a}^{(2)}, \mathbf{i}^{(2)}), \dots, (\mathbf{a}^{(k)}, \mathbf{i}^{(k)}))$$

where

- $\mathbf{a} = \mathbf{a}^{(1)} \dots \mathbf{a}^{(k)}$ and $\mathbf{i} = \mathbf{i}^{(1)} \dots \mathbf{i}^{(k)}$ ("..." means concatenation); and
- the entries of $\mathbf{i}^{(j)}$ are contained in the set $\{d_{j-1} + 1, d_{j-1} + 2, \dots, d_j\}$.

Now define

$$\Phi((\mathbf{a}, \mathbf{i})) := (\text{EGLS}(\mathbf{a}^{(1)}, \mathbf{i}^{(1)}), \dots, \text{EGLS}(\mathbf{a}^{(k)}, \mathbf{i}^{(k)})).$$

Proposition 2.2. *The map $\Phi : \text{Compatible}(w) \rightarrow \text{Tuples}(w)$ is well-defined and a bijection.*

Proof. Φ is well-defined: The condition (i) is just says T_j and U_j have the same shape, which is true by EGLS's description. For (ii), the splitting says each label in $\mathbf{i}^{(j)}$ is strictly bigger than d_{j-1} . Now by (cs.3), each label in $\mathbf{a}^{(j)}$ is strictly bigger than d_{j-1} as well. By EGLS's definition, the set of labels appearing in T_j is the same as that of $\mathbf{a}^{(j)}$; hence (ii) holds. Lastly, $\text{row}(T_j)$ is a reduced word for $\mathbf{a}^{(j)}$. Then (iii) is clear.

Φ is a bijection: Since EGLS is a bijective correspondence, clearly Φ is an injection. Consider the weight function on $\text{Compatible}(w)$ that assigns (\mathbf{a}, \mathbf{i}) weight $\mathbf{x}^{\mathbf{i}}$ and assigns $[(T_1, U_1), \dots, (T_k, U_k)]$ the weight $\mathbf{x}^{U_1} \dots \mathbf{x}^{U_k}$, where \mathbf{x}^{U_i} is the usual monomial associated to the tableau U_i . Then clearly Φ is a weight-preserving map (since EGLS is similarly weight-preserving). Hence the surjectivity of Φ holds by (4) and Theorem 2.1. \square

See [L04, Section 5] for a proof of Theorem 2.1 which is close to the study of the split EGLS correspondence (the argument constructs certain crystal operators).

2.4. The tableau $T[\alpha]$. Recall $w[\alpha] \in S_\infty$ satisfies $\text{code}(w[\alpha]) = \alpha$. Let \prec be the pure reverse lexicographic total ordering on monomials. The Schubert polynomial $\mathfrak{S}_{w[\alpha]}$ has leading term \mathbf{x}^α (with respect to \prec). The same is true of κ_α (see [RS95, Corollary 7]) so

$$(8) \quad \mathfrak{S}_{w[\alpha]} = \kappa_\alpha + \text{linear combination of other key polynomials.}$$

Given an increasing tableau U , the **nil left key** $K_-^0(U)$ is defined by [LS89] (cf. [RS95, p.111–114]). Let $\text{sort}(\alpha)$ be the partition obtained by rearranging α into weakly decreasing order. Also let $\text{content}(T)$ the usual content vector of a semistandard tableau T . This is a result of A. Lascoux-M.-P. Schützenberger (cf. [RS95, Theorem 4]):

Theorem 2.3.

$$\mathfrak{S}_w(X) = \sum \kappa_{\text{content}(K_-^0(U))}$$

where the sum is over all increasing tableaux U of shape $\text{sort}(\alpha)$ with $\text{row}(U) = w$.

Thus, by (8) combined with Theorem 2.3 there exists a unique increasing tableau $U[\alpha]$ of shape $\text{sort}(\alpha)$ with $\text{row}(U[\alpha]) = w[\alpha]$ and such that $\alpha = \text{content}(K_-^0(U[\alpha]))$.

Let $F_w = \lim_{k \rightarrow \infty} \mathfrak{S}_{1^k \times w}$ be the **stable Schubert polynomial** associated to w . This is a symmetric polynomial in infinitely many variables. So therefore one has an expansion

$$(9) \quad F_w = \sum_{\lambda} a_{w,\lambda} s_{\lambda},$$

where the $a_{w,\lambda} \in \mathbb{Z}_{\geq 0}$ are counted by increasing tableaux A of shape λ with $\text{row}(A) = w$.

In [S84, Theorem 4.1], it is shown $a_{w,\mu(w)'} = 1$ for a certain explicitly described “maximal” $\mu'(w)$. Moreover a simple description of the witnessing tableau $A[\alpha]$ is given. Straightforwardly, $\mu'(w[\alpha]) = \text{sort}(\alpha)$. Then $T[\alpha]$ is precisely the witnessing tableau $A[\alpha]$ for $a_{w[\alpha],\lambda(w[\alpha])}$ (after accounting for the fact that [S84]’s conventions use $F_{w[\alpha]}$ for what we call $F_{w[\alpha]^{-1}}$). We leave the details to the reader.

Finally, the expansion of Theorem 2.3 refines (9); see, e.g., [RS95]. Hence, $T[\alpha] = A[\alpha] = U[\alpha]$. So, $T[\alpha]$ is an increasing tableau of shape $\text{sort}[\alpha]$ with $\text{row}(T[\alpha]) = w[\alpha]$ and $\text{content}(K_-(T[\alpha])) = \alpha$.

2.5. Conclusion of the proof of Theorem 1.1: From the definition of $\text{Rothe}(w[\alpha])$:

Lemma 2.4. *The descents of $w[\alpha]$ are contained in the set of descents $d_1 < d_2 < \dots < d_k$ of α .*

Thus,

$$(10) \quad \mathfrak{S}_{w[\alpha]}(X) = \sum_{(\mathbf{a}, \mathbf{i})} \mathbf{x}^{\mathbf{i}} = \sum_{\lambda^1, \dots, \lambda^k} c_{\lambda^1, \dots, \lambda^k}^{w[\alpha]} s_{\lambda^1}(X_1) \cdots s_{\lambda^k}(X_k).$$

We recall a formula [RS95, Theorem 5]:

Theorem 2.5. *Fix an increasing tableau T with $\text{content}(K_-^0(T)) = \alpha$. Then*

$$\kappa_{\alpha} = \sum_{(\mathbf{a}, \mathbf{i})} \mathbf{x}^{\mathbf{i}}$$

where the sum is over compatible sequences (\mathbf{a}, \mathbf{i}) satisfying (cs.1), (cs.2), (cs.3) and $\text{EGLS}(\mathbf{a}) = T$.

Two reduced words \mathbf{a} and \mathbf{a}' for the same permutation are in the same **Coxeter-Knuth class** if $\text{EGLS}(\mathbf{a}) = \text{EGLS}(\mathbf{a}') = T$. This T **represents** the class. This equivalence relation \sim on reduced words is defined by the symmetric and transitive closure of the relations:

$$(11) \quad \begin{aligned} \mathbf{A}i(i+1)i\mathbf{B} &\sim \mathbf{A}(i+1)i(i+1)\mathbf{B} \\ \mathbf{A}acb\mathbf{B} &\sim \mathbf{A}cab\mathbf{B} \\ \mathbf{A}bac\mathbf{B} &\sim \mathbf{A}bca\mathbf{B} \end{aligned}$$

where $a < b < c$. In particular, it is true that $\mathbf{a} \sim \text{row}(\text{EGLS}(\mathbf{a}))$.

Restrict Φ to those $(\mathbf{a}, \mathbf{i}) \in \text{Compatible}(w[\alpha])$ such that $\text{EGLS}(\mathbf{a}) = T[\alpha]$. Consider $\Phi(\mathbf{a}, \mathbf{i}) = [(T_1, U_1), \dots, (T_k, U_k)]$. Since $\text{EGLS}(\mathbf{a}^{(i)}) \sim \text{row}(T_i)$, by (11) we see

$$(12) \quad \text{row}(T_1) \cdots \text{row}(T_k) \sim \mathbf{a}^{(1)} \cdots \mathbf{a}^{(k)} = \mathbf{a}.$$

However, since we have assumed $\text{EGLS}(\mathbf{a}) = T[\alpha]$, therefore:

$$(13) \quad \text{EGLS}(\text{row}(T_1) \cdots \text{row}(T_k)) = T[\alpha],$$

The other two requirements on (T_1, \dots, T_k) hold since Φ is well-defined.

Conversely, suppose $[(T_1, U_1), \dots, (T_k, U_k)]$ has (T_1, \dots, T_k) satisfying Theorem 1.1's conditions. Since Φ is a bijection, $\Phi^{-1}([(T_1, U_1), \dots, (T_k, U_k)]) = (\mathbf{a}, \mathbf{i}) \in \text{Compatible}(w[\alpha])$. Also, by (12), $\mathbf{a} \sim \text{row}(T_1) \cdots \text{row}(T_k)$. Now, we assumed (13) holds. Hence, $\text{EGLS}(\mathbf{a}) = T[\alpha]$ as desired. This completes the proof of the Theorem 1.1. \square

3. ADDITIONAL REMARKS

3.1. Comments on Theorem 1.1. Since κ_α specialize non-symmetric Macdonald polynomials (see, e.g., [HHL09, Section 5.3]), can one extend Theorem 1.1 in that direction?

Theorem 1.1 implies that the key module of [RS95, Section 5] should have an action of $GL(d_1) \times GL(d_2 - d_1) \times \cdots \times GL(d_k - d_{k-1})$ such that the character is κ_α .

V. Reiner suggests a variation of Theorem 1.1 using the plactic theory. The derivation should be similar, using formulas from [RS94]. However we are missing the analogue of [BKTY04, Corollary 4]; cf. [KMS06, Sections 7, 8]. Theorem 1.1 naturally generalizes to Grothendieck polynomials, using [BKTY05, BKSTY08]; details may appear elsewhere.

3.2. J_α 's form a (finite) basis of Pol. Clearly, $J_\alpha(X) = \mathbf{x}^\alpha + \sum_{\beta \prec \alpha} c_\beta \mathbf{x}^\beta$. One decomposes $f \in \text{Pol}$ into a possibly infinite sum of J_α 's:

$$(14) \quad f = \sum_{\alpha} g_{\alpha} J_{\alpha}$$

That is, find the \prec largest monomial \mathbf{x}^{θ_0} appearing in $f^{(0)} := f$ (say with coefficient c_{θ_0}) and let $f^{(1)} := f - c_{\theta_0} \cdot J_{\theta_0}$. Thus $f^{(1)}$ only contains monomials strictly smaller in the \prec ordering. Now repeat, defining $f^{(t+1)} := f^{(t)} - c_{\theta_t} J_{\theta_t}$ where \mathbf{x}^{θ_t} is the \prec -largest monomial appearing in $f^{(t)}$ etc. Since J_α is not homogeneous, each step t potentially introduces \prec -smaller monomials but of higher degree. However, we claim:

Proposition 3.1. *The expansion (14) is finite.*

Proof. By the K -Kohnert rule, each β that appears in J_α is contained in the smallest rectangle R that contains α . So the above procedure only involves the finitely many diagrams contained in R for one of the finitely many initial $\alpha \in \mathbb{Z}_{\geq 0}^\infty$ such that \mathbf{x}^α is in f . \square

3.3. More on the interplay of Grothendieck and the Ω polynomials. M. Shimozono has suggested that the expansion of \mathfrak{G}_w into Ω_α should alternate in sign, by degree. An explicit rule exhibiting this has been conjectured by V. Reiner and the second author.

ACKNOWLEDGEMENTS

AY thanks Jim Haglund, Alain Lascoux, Mark Shimozono and Vic Reiner for inspiring discussions and correspondence. AY also thanks Oliver Pechenik and Luis Serrano for helpful comments. This project was initiated during a summer undergraduate research experience at UIUC supported by NSF grant DMS 0901331. AY also was supported by NSF grant DMS 1201595 and a Helen Corley Petit endowment at UIUC.

REFERENCES

- [BJS93] S. Billey, W. Jockusch and R. Stanley, *Some Combinatorial Properties of Schubert Polynomials*, J. of Algebraic Comb. Vol. 2 Num. 4, 1993, 345–374.
- [BKSTY08] A. Buch, A. Kresch, M. Shimozono, H. Tamvakis and A. Yong, *Stable Grothendieck polynomials and K-theoretic factor sequences*, Math. Ann. **340** (2008), no. 2, 359–382.
- [BKTY05] A. S. Buch, A. Kresch, H. Tamvakis and A. Yong, *Grothendieck polynomials and quiver formulas*, Amer. J. Math., **127** (2005), 551–567.
- [BKTY04] ———, *Schubert polynomials and quiver formulas*, Duke Math J., Volume 122 (2004), Issue 1, 125–143.
- [D74] M. Demazure, *Une nouvelle formule des caractères*, Bull. Sci. Math. **98**(1974), 163–172.
- [EG87] P. Edelman and C. Greene, *Balanced tableaux*, Adv. in Math. **63** (1987), no. 1, 42–99.
- [FK94] S. Fomin and A. N. Kirillov, *Grothendieck polynomials and the Yang-Baxter equation*, Proc. 6th Intern. Conf. on Formal Power Series and Algebraic Combinatorics, DIMACS, 1994, 183–190.
- [HHL09] J. Haglund, M. Haiman and N. Loehr, *A combinatorial formula for non-symmetric Macdonald polynomials*, Amer. J. of Math., 103 (2008), pp. 359–383.
- [KMS06] A. Knutson, E. Miller and M. Shimozono, *Four positive formulae for type A quiver polynomials*, Invent. Math. **166**(2006), 229–325.
- [K90] A. Kohnert, *Weintrauben, Polynome, Tableaux*, Bayreuth Math. Schrift. **38**(1990), 1–97.
- [L13+] A. Lascoux, *Polynomials*, in preparation, 2013.
- [L03] ———, *Schubert & Grothendieck: un bilan bidécennal*, Sémin. Lothar. Combin. **50** (2003/04), Art. B50i.
- [L01] ———, *Transition on Grothendieck polynomials*, Physics and Combinatorics, 2000 (Nagoya), pp. 164–179, World Scientific Publishing, River Edge (2001).
- [LS90] A. Lascoux and M.-P. Schützenberger, *Keys and standard bases*, in “Tableaux and Invariant Theory”, IMA Volumes in Math and its Applications (D. Stanton, Ed.), Vol. 19, pp. 125–144, Southend on Sea, UK, 1990.
- [LS89] ———, *Tableaux and non-commutative Schubert polynomials*, Funct. Anal. Appl. **23**(1989), 63–64.
- [LS82] ———, *Structure de Hopf de l’anneau de cohomologie et de l’anneau de Grothendieck d’une variété de drapeaux*, C. R. Acad. Sci. Paris Sér. I Math. **295** (1982), no. 11m 629–633.
- [L04] C. Lenart, *A unified approach to combinatorial formulas for Schubert polynomials*, J. Algebraic Combin., **20**(2004), 263–299.
- [LRS06] C. Lenart, S. Robinson and F. Sottile, *Grothendieck polynomials via permutation patterns and chains in the Bruhat order*, Amer. J. Math. **128** (2006), no. 4, 805–848.
- [M01] L. Manivel, *Symmetric functions, Schubert polynomials and degeneracy loci*, American Mathematical Society, Providence, RI, 2001.
- [RS95] V. Reiner and M. Shimozono, *Key polynomials and a flagged Littlewood-Richardson rule*, J. Comb. Theory. Ser. A., **70**(1995), 107–143.
- [RS94] ———, *Plactification*, J. Alg. Comb., **4**(1995), 331–351.
- [S84] R. P. Stanley, *On the number of reduced decompositions of elements of Coxeter groups*, Eur. J. Comb. **5**, 359–372 (1984).
- [W03] R. Winkel, *Diagram rules for the generation of Schubert polynomials*, J. Combin. Th. A., **86**(1999), 14–48.

DEPARTMENT OF INDUSTRIAL ENGINEERING AND MANAGEMENT SCIENCES, NORTHWESTERN UNIVERSITY, EVANSTON, IL 60208

E-mail address: ColleenRoss2012@u.northwestern.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL 61801

E-mail address: ayong@illinois.edu