

ON THE UNBOUNDED OF A CLASS OF FOURIER INTEGRAL OPERATOR ON $L^2(\mathbb{R}^n)$

A. SENOUSSAOUI

ABSTRACT. In this paper, we give an example of Fourier integral operator with a symbol belongs to $\bigcap_{0 < \rho < 1} S_{\rho,1}^0$ that cannot be extended as a bounded operator on $L^2(\mathbb{R}^n)$.

1. INTRODUCTION

A Fourier integral operator is a singular integral operator of the form

$$I(a, \phi)u(x) = \int \int e^{i\phi(x,y,\theta)} a(x, y, \theta) u(y) dy d\theta$$

defined under certain assumptions on the regularity and asymptotic properties of the phase function ϕ and the amplitude function a . Here θ plays the role of the covariable.

Fourier integral operators are more general than pseudodifferential operators, where the phase function is of the form $\langle x - y, \theta \rangle$.

Let us denote by $S_{\rho,\delta}^m(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \mathbb{R}^N)$ the space of $a(x, y, \theta) \in C^\infty(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \mathbb{R}^N)$, satisfying

$$|\partial_x^\alpha \partial_y^\beta \partial_\theta^\gamma a(x, y, \theta)| \leq C_{\alpha,\beta,\gamma} \lambda^{m-\rho|\gamma|+\delta(|\alpha|+|\beta|)}(\theta), \quad \forall (\alpha, \beta, \gamma) \in \mathbb{N}^{n_1} \times \mathbb{N}^{n_2} \times \mathbb{N}^N,$$

where $\lambda(\theta) = (1 + |\theta|)$.

The phase function $\phi(x, y, \theta)$ is assumed to be a $C^\infty(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2} \times \mathbb{R}^N, \mathbb{R})$ real function, homogeneous in θ of degree 1.

Since 1970, many efforts have been made by several authors in order to study this type of operators (see, e.g., [1, 3, 6, 7, 8]).

For the Fourier integral operators, an interesting question is under which conditions on a and ϕ these operators are bounded on L^2 or on the Sobolev spaces H^s .

It was proved in [10] that all pseudodifferential operators with symbol in $S_{\rho,\delta}^0$ are bounded on L^2 if $\delta < \rho$. When $0 < \delta = \rho < 1$, Calderon and Vaillancourt [2] have proved that all pseudodifferential operators with symbol in $S_{\rho,\rho}^0$ are bounded on L^2 . On the other hand, Kumano-Go [11] has given a pseudodifferential operator with symbol belonging to $\bigcap_{0 < \rho < 1} S_{\rho,1}^0$ which is not bounded on $L^2(\mathbb{R})$.

2000 *Mathematics Subject Classification.* Primary 35S30, 35S05 ; Secondary 47A10, 35P05.

Key words and phrases. Fourier integral operators, amplitude, symbol and phase, L^2 -boundedness.

The paper was realized during the stay of the author at Université Libre de Bruxelles. We would like to thank Professor J.-P. Gossez for several discussions.

For Fourier integral operators, it has been proved in [1] that the operator $I(a, \phi) : L^2 \rightarrow L^2$ is bounded if $\delta = m = \rho = 0$. Recently, M. Hasanov [6] constructed a class of unbounded Fourier integral operators on $L^2(\mathbb{R})$ with an amplitude in $S_{1,1}^0$.

For $u \in C_0^\infty(\mathbb{R}^n)$, the integral operators

$$(1.1) \quad I(a, S)\varphi(x) = \int e^{iS(x,\theta)} a(x, y, \theta) \mathcal{F}\varphi(\theta) d\theta$$

appear naturally in the expression of the solutions of hyperbolic partial differential equations (see [4, 5, 12]).

If we write formally the expression of the Fourier transformation $\mathcal{F}u(\theta)$ in (1.1), we obtain the following Fourier integral operators

$$(1.2) \quad I(a, S)u(x) = \iint e^{i(S(x,\theta)-y\theta)} a(x, y, \theta) u(y) dy d\theta$$

in which the phase function has the form $\phi(x, y, \theta) = S(x, \theta) - y\theta$. We note that in [13], we have studied the L^2 -boundedness and L^2 -compactness of a class of Fourier integral operator of the form (1.2).

In this article we give an example of a Fourier integral operator, in higher dimension, of the form (1.1) with symbol $a(x, \theta) \in \bigcap_{0 < \rho < 1} S_{\rho,1}^0$ independent on y , that cannot be extended to a bounded operator in $L^2(\mathbb{R}^n)$, $n \geq 1$. Here we take the phase function in the form of separate variable $S(x, \theta) = \varphi(x)\psi(\theta)$.

2. The boundedness on $C_0^\infty(\mathbb{R}^n)$ and on $D'(\mathbb{R}^n)$

If $\varphi \in C_0^\infty(\mathbb{R}^n)$, we consider the following integral transformations

$$(2.3) \quad \begin{aligned} (I(a, S)\varphi)(x) &= \int_{\mathbb{R}^N} e^{iS(x,\theta)} a(x, \theta) \mathcal{F}\varphi(\theta) d\theta \\ &= \int_{\mathbb{R}^N \times \mathbb{R}^n} e^{i(S(x,\theta)-y\theta)} a(x, \theta) \varphi(y) d\theta dy \end{aligned}$$

for $x \in \mathbb{R}^n$ and $N \in \mathbb{N}$.

In general the integral (2.3) is not absolutely convergent, so we use the technique of the oscillatory integral developed by L.Hörmander in [8]. The phase function S and the amplitude a are assumed to satisfy the hypothesis

- (H1) $S \in C^\infty(\mathbb{R}_x^n \times \mathbb{R}_\theta^N, \mathbb{R})$ (S real function)
- (H2) $\forall \beta \in \mathbb{N}^N, \exists C_\beta > 0$;

$$\left| \partial_\theta^\beta S(x, \theta) \right| \leq C_\beta(x) \lambda^{(1-|\beta|)_+}(\theta), \quad \forall (x, \theta) \in \mathbb{R}_x^n \times \mathbb{R}_\theta^N$$

where $\lambda(\theta) = (1 + |\theta|)$ and $(1 - |\beta|)_+ = \max(1 - |\beta|, 0)$.

- (H3) S satisfies

$$\left(\frac{\partial S}{\partial x}, \frac{\partial S}{\partial \theta} - y \right) \neq 0, \quad \forall (x, \theta) \in \mathbb{R}_x^n \times (\mathbb{R}_\theta^N \setminus \{0\}).$$

Remark 2.1. *If the phase function $S(x, \theta)$ is homogeneous in θ of degree 1, then it satisfies (H2).*

For any open Ω of $\mathbb{R}_x^n \times \mathbb{R}_\theta^N$, $m \in \mathbb{R}$, $\rho > 0$ and $\delta \geq 0$ we set

$$S_{\rho,\delta}^m(\Omega) = \left\{ \begin{array}{l} a \in C^\infty(\Omega); \quad \forall(\alpha, \beta) \in \mathbb{N}^n \times \mathbb{N}^N, \quad \exists C_{\alpha,\beta} > 0; \\ \left| \partial_x^\alpha \partial_\theta^\beta a(x, \theta) \right| \leq C_{\alpha,\beta} \lambda^{m-\rho|\beta|+\delta|\alpha|}(\theta). \end{array} \right\}$$

Theorem 2.2. *If S satisfies (H1), (H2), (H3) and if $a \in S_{\rho,\delta}^m(\mathbb{R}_x^n \times \mathbb{R}_\theta^N)$, then $I(a, \phi)$ is a continuous operator from $C_0^\infty(\mathbb{R}^n)$ to $C^\infty(\mathbb{R}^n)$ and from $\mathcal{E}'(\mathbb{R}^n)$ to $D'(\mathbb{R}^n)$, where $\rho > 0$ and $\delta < 1$.*

Proof. See [6], [4, pages 50-51]. \square

Corollary 2.3. *Let $\varphi(x), \psi(\theta) \in C^\infty(\mathbb{R}^n, \mathbb{R})$ two functions, $\psi(\theta)$ is homogeneous of degree 1 ($\psi(\theta) \neq 0$) and $\varphi(x)$ satisfies*

$$(2.4) \quad \varphi'(x) \neq 0, \quad \forall x \in \mathbb{R}^n.$$

Then the operator

$$(2.5) \quad (Fu)(x) = \int_{\mathbb{R}^n} e^{i\varphi(x)\psi(\theta)} a(x, \theta) \mathcal{F}u(\theta) d\theta, \quad u \in \mathcal{S}(\mathbb{R}^n)$$

maps continuously $C_0^\infty(\mathbb{R}^n)$ to $C^\infty(\mathbb{R}^n)$ and $\mathcal{E}'(\mathbb{R}^n)$ to $D'(\mathbb{R}^n)$ for every $a \in S_{\rho,\delta}^m(\mathbb{R}_x^n \times \mathbb{R}_\theta^N)$, where $\rho > 0$ and $\delta < 1$.

Proof. For the phase function $S(x, \theta) = \varphi(x)\psi(\theta)$, (H1), (H2) and (H3) are satisfied. \square

3. The unboundedness of the operator F on $L^2(\mathbb{R}^n)$

In this section we shall construct a symbol $a(x, \theta)$ in the Hörmander space $\bigcap_{0 < \rho < 1} S_{\rho,1}^0(\mathbb{R}_x^n \times \mathbb{R}_\theta^n)$, such that the Fourier integral operator F can not be extended as a bounded operator in $L^2(\mathbb{R}^n)$.

Lemma 3.1. *(Kumano-Go [11]). Let $f_0(t)$ be a continuous function on $[0, 1]$ such that*

$$(3.1) \quad f_0(0) = 0, \quad f_0(t) > 0 \text{ in }]0, 1].$$

Then, there exists a continuous function $b(t)$ on $[0, 1]$ such that $b(t)$ satisfies the conditions

$$(3.2) \quad \left\{ \begin{array}{l} f_0(t) \leq b(t) \text{ on } [0, 1], \\ b \in C^\infty(]0, 1]), \quad b(0) = 0, \quad b'(t) > 0 \text{ in }]0, 1], \\ |b^{(n)}(t)| \leq C_n t^{-n} \text{ in }]0, 1], \quad n \in \mathbb{N}^*, \quad C_n > 0. \end{array} \right.$$

Definition 3.2. *It is obvious that an operator A is extended as a bounded operator in $L^2(\mathbb{R}^n)$ if and only if there exists a constant $C > 0$ such that*

$$(3.3) \quad \|Au\|_{L^2} \leq C \|u\|_{L^2} \text{ for any } u \in \mathcal{S}(\mathbb{R}^n).$$

Theorem 3.3. *Let A be an operator, given at least for $x \in]0, \beta[^n$ ($\beta < 1$), by*

$$(Au)(x) = \int_{\mathbb{R}^n} u(g(x)z) \mathcal{F}\Psi(z) dz, \quad u \in \mathcal{S}(\mathbb{R}^n),$$

we denote here by $]0, \beta[^n = \prod_{j=1}^n]0, \beta[$.

If $\Psi \in \mathcal{S}(\mathbb{R}^n)$, $\Psi(0) = 1$ and the function $g \in C^0(]0, \beta[^n, \mathbb{R}_+)$ satisfies

$$(3.4) \quad \begin{cases} \lim_{|x| \rightarrow 0^+} \frac{g(x)}{|x|} = 0, \\ \forall i \in \{1, \dots, n\}; x_i \longrightarrow g(x_1, \dots, x_i, \dots, x_n) \text{ is increasing on }]0, \beta[. \end{cases}$$

Then the operator A cannot be extended to a bounded operator in $L^2(\mathbb{R}^n)$.

Proof. Using the Fourier inversion formula in $\mathcal{S}(\mathbb{R}^n)$, we have

$$(Au)(x) = \int_{\mathbb{R}^n} u(g(x)z) \mathcal{F}\Psi(z) dz, \quad u \in \mathcal{S}(\mathbb{R}^n).$$

Then there exists a constant $N_0 > 0$ such that

$$(3.5) \quad \left| (2\pi)^{-\frac{n}{2}} \int_{[-N, N]^n} \mathcal{F}\Psi(z) dz \right| \geq \beta \quad \text{for any } N \geq N_0.$$

Setting for $\varepsilon > 0$

$$u_\varepsilon(z) = \begin{cases} (2\pi)^{-\frac{n}{2}}, & \text{for } z \in [-\varepsilon, \varepsilon]^n \\ 0, & \text{for } z \notin [-\varepsilon, \varepsilon]^n \end{cases}.$$

Then, using the density of $\mathcal{S}(\mathbb{R}^n)$ in $L^2(\mathbb{R}^n)$, we see that $(Au_\varepsilon)(x)$ must be

$$(3.6) \quad (Au_\varepsilon)(x) = (2\pi)^{-\frac{n}{2}} \int_{[-\frac{\varepsilon}{g(x)}, \frac{\varepsilon}{g(x)}]^n} \mathcal{F}\Psi(z) dz \quad \text{for } x \in]0, \beta[^n.$$

By (3.4) for any $p \in \mathbb{N}^*$ there exists a small $\varepsilon_p \geq 0$ such that

$$\frac{\varepsilon_p}{g(p\varepsilon_p, \dots, p\varepsilon_p)} \geq N_0 \quad \text{and} \quad p\varepsilon_p \leq \beta.$$

It follows from the condition (3.4) that

$$\frac{\varepsilon_p}{g(x)} \geq \frac{\varepsilon_p}{g(p\varepsilon_p, \dots, p\varepsilon_p)} \geq N_0, \quad \text{holds for } x \in]0, p\varepsilon_p[^n,$$

so that, using (3.5) and (3.6), we have

$$(3.7) \quad \|Au_{\varepsilon_p}\|_{L^2}^2 = \int_{\mathbb{R}^n} |Au_{\varepsilon_p}(x)|^2 dx \geq \int_{[0, p\varepsilon_p]^n} |Au_{\varepsilon_p}(x)|^2 dx \geq \beta^2 (p\varepsilon_p)^n.$$

Assume that A is bounded on $L^2(\mathbb{R}^n)$. According to (3.3) there exists $C > 0$ such that:

$$\beta^2 (p\varepsilon_p)^n \leq \|Au_{\varepsilon_p}\|_{L^2}^2 \leq C^2 (2\varepsilon_p)^n \quad \text{for any } p.$$

Which is a contradiction. \square

Let $K(t)$ be a function from $\mathcal{S}(\mathbb{R})$ such that $K(t) = 1$ on $[-\delta, \delta]$ ($\delta < 1$), $b(t) \in C^0([0, 1])$ be a continuous function satisfying conditions (3.2) and $\varphi(x), \psi(\theta) \in C^\infty(\mathbb{R}^n, \mathbb{R})$ with $\psi(\theta)$ homogeneous of degree 1 ($\psi(\theta) \neq 0$). We assume that $\varphi(x)$ satisfies

$$(3.8) \quad |\varphi(x)| \leq C|x| \quad \text{for } |x| \leq 1.$$

We remark that if the function $\varphi(x)$ is homogeneous of degree 1, then it satisfies (3.8).

For $x = (x_1, \dots, x_n), \theta = (\theta_1, \dots, \theta_n) \in \mathbb{R}^n$, set

$$q(x, \theta) = e^{-i\varphi(x)\psi(\theta)} \prod_{j=1}^n K(b(|x|)|x|\theta_j)$$

Lemma 3.4. *The function $q \in C^\infty([-1, 1]^n \times \mathbb{R}_\theta^n)$ and the following estimate holds:*

$$(3.9) \quad \begin{aligned} & \forall (\alpha, \beta) \in \mathbb{N}^n \times \mathbb{N}^n, \exists C_{\alpha, \beta} > 0; \\ & \left| \partial_x^\alpha \partial_\theta^\beta q(x, \theta) \right| \leq C_{\alpha, \beta} \lambda^{|\alpha| - |\beta|}(\theta) \{b(\lambda^{-1}(\theta))\}^{-|\beta|} \text{ on } [-1, 1]^n \times \mathbb{R}_\theta^n. \end{aligned}$$

Proof. We adopt here the same strategy of Kumano-Go [11] lemma 2.

Since $\psi(\theta)$ is homogeneous of degree 1 ($\psi(\theta) \neq 0$), it will be sufficient to check the estimate for $n = 1$ on $[-1, 1] \times \mathbb{R}_\theta$, i.e.

$$(3.10) \quad \begin{aligned} & \forall (j, k) \in \mathbb{N} \times \mathbb{N}, \exists C_{j, k} > 0; \\ & \left| \partial_x^j \partial_\theta^k \left[e^{-i\varphi(x)\theta} K(b(|x|)x\theta) \right] \right| \leq C_{j, k} \lambda^{j-k}(\theta) \{b(\lambda^{-1}(\theta))\}^{-k}. \end{aligned}$$

Since $K(t) \in \mathcal{S}(\mathbb{R})$ and $K^{(n)}(t) = 0$ on $[-\delta, \delta]$, $n \in \mathbb{N}^*$, then

$$(3.11) \quad |t^l K(t)| \leq C_l, \quad \forall l \in \mathbb{N}$$

$$(3.12) \quad |t^l K^{(n)}(t)| \leq C_{l, n}, \quad \forall n \in \mathbb{N}^*, \forall l \in \mathbb{Z}.$$

By Leibnitz's formula we have

$$\begin{aligned} & \partial_x^j \partial_\theta^k \left[e^{-i\varphi(x)\theta} K(b(|x|)x\theta) \right] = \\ & \sum_{\substack{j_1 + j_2 = j \\ k_1 + k_2 = k}} C_{j_1, j_2, k_1, k_2}^{j, k} \partial_x^{j_1} \left[K^{(k_1)}(b(|x|)x\theta) (b(|x|)x)^{k_1} \right] \partial_x^{j_2} \left[(-i\varphi(x))^{k_2} e^{-i\varphi(x)\theta} \right], \end{aligned}$$

where $C_{j_1, j_2, k_1, k_2}^{j, k} = \frac{j!k!}{j_1!j_2!k_1!k_2!}$. Then, by means of (3.8) and (3.2), we have for constants $C'_{j, k}$

$$(3.13) \quad \begin{aligned} & \left| \partial_x^j \partial_\theta^k \left[e^{-i\varphi(x)\theta} K(b(|x|)x\theta) \right] \right| \leq C'_{j, k} K_{j+k}(b(|x|)x\theta) \\ & \sum_{\substack{j_1 + j_2 = j \\ k_1 + k_2 = k}} \max_{\substack{s_1 + s_2 + s_3 = j_1 \\ s_3 \leq k_1}} \left\{ |\theta|^{s_1} |x|^{-s_2} (b(|x|)x)^{k_1 - s_3} \right\} \max_{\substack{l_1 + l_2 = j_2 \\ l_2 \leq k_2}} \left\{ |\theta|^{l_1} |x|^{k_2 - l_2} \right\}, \end{aligned}$$

where $K_p(t)$, $p \in \mathbb{N}$ are defined by

$$K_0(t) = |K(t)|, \quad K_p(t) = \max_{1 \leq p' \leq p} |K^{(p')}(t)|, \quad p \in \mathbb{N}^*.$$

Writing $b(|x|)x = |b(|x|)x\theta| |\theta|^{-1}$ and $|x|^{-1} = |b(|x|)x\theta|^{-1} b(|x|) |\theta|$, then there exists a constant C ,

$$(3.14) \quad \begin{cases} |b(|x|)x| \leq C \lambda(b(|x|)x\theta) \lambda^{-1}(\theta), \\ |x|^{-1} \leq C |b(|x|)x\theta|^{-1} \lambda(\theta) \end{cases} \text{ on } [-1, 1] \times \mathbb{R}_\theta.$$

We have $b^{-1}(|x|) \leq b^{-1}(\lambda^{-1}(\theta))$ when $|x| \geq \lambda^{-1}(\theta)$ (because b is increasing). Then,

$$\begin{aligned} |x| &= |b(|x|)x\theta| (b(|x|) |\theta|)^{-1} \leq |b(|x|)x\theta| |\theta|^{-1} b^{-1}(\lambda^{-1}(\theta)) \\ &\leq C \lambda(b(|x|)x\theta) \lambda^{-1}(\theta) b^{-1}(\lambda^{-1}(\theta)). \end{aligned}$$

Bearing in mind the other case when $|x| \leq \lambda^{-1}(\theta)$, we obtain for a constant \tilde{C}

$$(3.15) \quad |x| \leq \tilde{C} \lambda(b(|x|)x\theta) \lambda^{-1}(\theta) b^{-1}(\lambda^{-1}(\theta)).$$

Finally from (3.11) to (3.15), we obtain (3.10). □

Lemma 3.5. *For any continuous function $b_0(t)$ on $[1, +\infty[$ such that*

$$(3.16) \quad b_0(t) > 0, \quad \lim_{t \rightarrow +\infty} b_0(t) = +\infty,$$

then there exists a continuous function $b(t)$ on $[0, 1]$ witch satisfies conditions (3.2) such that we have on $[-1, 1]^n \times \mathbb{R}_\theta^n$

$$(3.17) \quad \left| \partial_x^\alpha \partial_\theta^\beta q(x, \theta) \right| \leq C_{\alpha, \beta} \lambda^{|\alpha| - |\beta|}(\theta) \{b_0(\lambda(\theta))\}^{|\beta|}, \quad \alpha, \beta \in \mathbb{N}^n.$$

Proof. Setting

$$\begin{cases} f_0(t) = \{b_0(t^{-1})\}^{-1} & \text{on }]0, 1[\\ f_0(0) = 0, \end{cases}$$

then f_0 is a continuous function on $[0, 1]$ which verifies condition (3.1). Then, by lemma 3.1, there exists a continuous function $b(t)$ which satisfies (3.2). Noting that

$$\{b(\lambda^{-1}(\theta))\}^{-1} \leq \{f_0(\lambda^{-1}(\theta))\}^{-1} = b_0(\lambda(\theta))$$

this gives (3.17). □

Lemma 3.6. *Let $\{b_l(t)\}_{l \in \mathbb{N}^*}$ be a sequence of continuous functions on $[1, +\infty[$ which satisfy (3.16). Then, there exists a continuous function $b_0(t)$ verifying (3.16), such that, for any l_0 ,*

$$b_l(t) \geq b_0(t) \quad \text{on } [t_{l_0}, +\infty[, \quad l = 1, \dots, l_0$$

Finally, our but is to give an unbounded Fourier integral operator of the form (2.5) with symbol in $\bigcap_{0 < \rho < 1} S_{\rho, 1}^0(\mathbb{R}_x^n \times \mathbb{R}_\theta^n)$.

Theorem 3.7. *There exist a Fourier integral operator F of the form (2.5), with symbol $a \in \bigcap_{0 < \rho < 1} S_{\rho, 1}^0(\mathbb{R}_x^n \times \mathbb{R}_\theta^n)$, which cannot be extended to be a bounded operator on $L^2(\mathbb{R}^n)$.*

Proof. Let $\phi(s)$ be a $C_0^\infty(\mathbb{R})$ function such that

$$\begin{cases} \phi(s) = 1 & \text{on } [-\beta, \beta] \quad (\beta < 1) \\ \text{supp} \phi \subset [-1, 1]. \end{cases}$$

Define a C^∞ -symbol $a(x, \theta)$ by

$$a(x, \theta) = e^{-i\varphi(x)\psi(\theta)} \prod_{j=1}^n \phi(x_j) K(b(|x|)|x|\theta_j) \quad \text{in } \mathbb{R}_x^n \times \mathbb{R}_\theta^n$$

where $K(t)$ and $b(t)$ are the functions of lemma 3.4. Let $b_l(t) = \overbrace{\log \dots \log}^l (C_l + t)$ defined on $[1, +\infty[$ and C_l some large constant, then by lemmas 3.5 and 3.6 we have, $\forall (\alpha, \gamma) \in \mathbb{N}^n \times \mathbb{N}^n, \forall l \in \mathbb{N}^*$

$$(3.18) \quad \left| \partial_x^\alpha \partial_\theta^\gamma a(x, \theta) \right| \leq C_{\alpha, \gamma, l} \lambda^{|\alpha| - |\gamma|}(\theta) \{b_l(\lambda(\theta))\}^{|\gamma|},$$

ON THE UNBOUNDED OF A CLASS OF FOURIER INTEGRAL OPERATOR ON $L^2(\mathbb{R}^n)$

$C_{\alpha,\gamma,l}$ are constants, so that $a(x, \theta) \in \bigcap_{0 < \rho < 1} S_{\rho,1}^0(\mathbb{R}_x^n \times \mathbb{R}_\theta^n)$. Furthermore the corresponding Fourier integral F is

$$\begin{aligned} (Fu)(x) &= \int_{\mathbb{R}_\theta^n} e^{i\varphi(x)\psi(\theta)} a(x, \theta) \mathcal{F}u(\theta) d\theta \\ (3.19) \quad &= \prod_{j=1}^n \phi(x_j) \int_{\mathbb{R}_\theta^n} \prod_{j=1}^n K(b(|x|)|x|\theta_j) \mathcal{F}u(\theta) d\theta, \quad u \in \mathcal{S}(\mathbb{R}^n). \end{aligned}$$

We consider $(Fu)(x)$ in $]0, \beta]^n$. Then, using an adequate change of variable in the integral (3.19), we have

$$(Fu)(x) = \int_{\mathbb{R}_z^n} u(b(|x|)|x|z) \prod_{j=1}^n \mathcal{F}K(z_j) dz$$

which has the form of A in theorem 3.3. In addition the function $g(x) = b(|x|)|x|$ satisfies (3.4). Consequently the operator F cannot be extended as a bounded operator on $L^2(\mathbb{R}^n)$. \square

REFERENCES

- [1] Asada, K and Fujiwara, D.: *On some oscillatory transformation in $L^2(\mathbb{R}^n)$* . *Japan. J. Math.* vol 4 (2), 1978, p299-361.
- [2] Caldéron, A. P. and Vaillancourt, R.: *On the boundedness of pseudodifferential operators*. *J. Math. Soc. Japan* 23, 1972, p374-378.
- [3] Duistermaat, J.J.: *Fourier integral operators*. *Courant Institute Lecture Notes, New-York 1973*.
- [4] Egorov, Yu. V. and Shubin, M. A.: *Partial differential equations. Vol II, Springer-Verlag, Berlin, 1994*.
- [5] Egorov, Yu. V. and Shubin, M. A.: *Partial differential equations. Vol IV, Springer-Verlag, Berlin, 1994*.
- [6] Hasanov, M., *A class of unbounded Fourier integral operators*. *J. Math. Analysis and application* 225, 1998, p641-651.
- [7] Helffer, B.: *Théorie spectrale pour des opérateurs globalement elliptiques*. *Société Mathématiques de France, Astérisque 112, 1984*.
- [8] Hörmander, L.: *Fourier integral operators I*, *Acta Math.* vol 127, 1971, p33-57.
- [9] Hörmander, L.: *Pseudo-differential operators of type (1.1)*. *Comm. Pure Appl. Math.* 33 (1988), p1085-1111.
- [10] Hörmander, L.: *On the L^2 continuity of pseudodifferential operators*. *Comm. Pure Appl. Math.* 24 (1971), p529-535.
- [11] Kumano-Go, A.: *A problem of Nirenberg on pseudo-differential operators*. *Comm. Pure Appl. Math.* 23 (1970), p115-121.
- [12] Messirdi, B. and Senoussaoui, A.: *Parametrix du problème de Cauchy C^∞ pour un opérateur différentiel matriciel fortement hyperbolique muni des ordres de Leray-Volevič*. *Journal for Analysis and its Applications*. Vol 24, (3), 2005, 581-592.
- [13] Messirdi, B. and Senoussaoui, A.: *On the L^2 -boundedness and L^2 -compactness of a class of Fourier integral Operators*. *Electronic J. Diff. Equa.* Vol 2006, (26), 2006, 1-12.

UNIVERSITÉ D'ORAN, FACULTÉ DES SCIENCES, DÉPARTEMENT DE MATHÉMATIQUES. B.P. 1524 EL-MNAOUER, ORAN, ALGERIA.

E-mail address: senoussaoui_abdou@yahoo.fr, senoussaoui.abderahmane@univ-oran.dz