

REAL SOLUTIONS TO THE NONLINEAR HELMHOLTZ EQUATION WITH LOCAL NONLINEARITY

GILLES EVEQUOZ AND TOBIAS WETH

ABSTRACT. In this paper, we study real solutions of the nonlinear Helmholtz equation

$$-\Delta u - k^2 u = f(x, u), \quad x \in \mathbb{R}^N$$

satisfying the asymptotic conditions

$$u(x) = O(|x|^{\frac{1-N}{2}}) \quad \text{and} \quad \frac{\partial^2 u}{\partial r^2}(x) + k^2 u(x) = o(|x|^{\frac{1-N}{2}}) \quad \text{as } r = |x| \rightarrow \infty.$$

We develop the variational framework to prove the existence of nontrivial solutions for compactly supported nonlinearities without any symmetry assumptions. In addition, we consider the radial case in which, for a larger class of nonlinearities, infinitely many solutions are shown to exist. Our results give rise to the existence of standing wave solutions of corresponding nonlinear Klein–Gordon equations with arbitrarily large frequency.

1. INTRODUCTION

The study of the existence and qualitative properties of solutions to nonlinear wave equations

$$(1) \quad \frac{\partial^2 \psi}{\partial t^2}(t, x) - \Delta \psi(t, x) + V(x)\psi(t, x) = f(x, \psi(t, x)), \quad (t, x) \in \mathbb{R} \times \mathbb{R}^N,$$

goes back to the sixties, see e.g. the classical paper by Jörgens [13]. Since then, many authors have been investigating various aspects of this problem, including the question of existence and orbital stability of *standing wave (or solitary wave) solutions*. An ansatz for solutions of this type is given by

$$(2) \quad \psi(t, x) = e^{-i(\omega t + \varphi)} u(x), \quad \omega, \varphi \in \mathbb{R},$$

with a real-valued function u on \mathbb{R}^N . Taking for example a nonlinearity of the form $f(x, \psi) = g(x, |\psi|^2)\psi$ where g is a real-valued function, we see that such a ψ solves (1) if and only if u solves the reduced wave equation

$$(3) \quad -\Delta u + V(x)u - \omega^2 u = f(x, u), \quad x \in \mathbb{R}^N.$$

The requirement that u be real-valued guarantees that the corresponding energy density

$$E(t, x) = \frac{1}{2} \left(\left| \frac{\partial \psi}{\partial t}(t, x) \right|^2 + |\nabla \psi(t, x)|^2 + V(x)|\psi(t, x)|^2 - \int_0^{|\psi(t, x)|^2} g(x, \tau) d\tau \right)$$

is constant in t at every point $x \in \mathbb{R}^N$, a property which is characteristic of standing wave solutions of (1). Note that, due to the presence of the linear potential V , it is natural to consider nonlinearities f satisfying $\partial_u f(x, 0) = 0$ on \mathbb{R}^N . In this case, in almost all of the available literature it is assumed that ω^2 is not contained in the essential spectrum of

Key words and phrases. Helmholtz equation and entire solutions and local nonlinearity and Dirichlet to Neumann map and variational methods.

the Schrödinger operator $-\Delta + V$. We refer the reader to the surveys and monographs [3, 8, 20, 25, 26, 30] and the references therein for results in this case. In the special case where $V \equiv V_0$ is a constant, this restriction amounts to assuming $V_0 > \omega^2 \geq 0$. Some authors have also considered the limiting case where ω^2 coincides with the infimum or another boundary point of the essential spectrum of V , see e.g. [15] for a survey of classical results and [1, 2, 3, 6, 9, 18, 22, 31] for more recent work in this case. On the contrary, very little seems to be known if ω^2 is contained in the interior of the essential spectrum of V . In the present paper, we consider this situation in the special case $V \equiv V_0$ and $\omega^2 > V_0$, so by setting $k^2 = \omega^2 - V_0$ we arrive at the nonlinear Helmholtz equation

$$(4) \quad -\Delta u - k^2 u = f(x, u), \quad x \in \mathbb{R}^N,$$

with $k > 0$. It seems unclear a priori in which space one should approach this problem and whether variational methods can be used. In the present paper, we provide first results in the case where f is supported in a bounded subset of \mathbb{R}^N , i.e. f vanishes in $[\mathbb{R}^N \setminus B_R] \times \mathbb{R}$ for $R > 0$ sufficiently large. Here $B_R \subset \mathbb{R}^N$ denotes the open ball centered at 0 with radius R . Note that in this case no nontrivial solution of (4) exists in the space $L^2(\mathbb{R}^N)$, as follows immediately from a classical result of Rellich, see [21, Satz 1]. We will focus instead on solutions satisfying the asymptotic conditions

$$(5) \quad u(x) = O(|x|^{\frac{1-N}{2}}) \quad \text{and} \quad \frac{\partial^2 u}{\partial r^2}(x) + k^2 u(x) = o(|x|^{\frac{1-N}{2}}) \quad \text{as } r = |x| \rightarrow \infty.$$

Thus the solutions decay to zero as $|x| \rightarrow \infty$ if $N \geq 2$. We emphasize that in general not all solutions of (4) satisfy (5) if $N \geq 2$. In particular, in case $N \geq 2$ and $f \equiv 0$, (5) is not satisfied by one-dimensional standing wave solutions of (4) given by

$$(6) \quad x \mapsto \sin(kx \cdot \xi + \varphi) \quad \text{with a unit vector } \xi \in \mathbb{R}^N \text{ and } \varphi \in \mathbb{R}.$$

The above restriction on f allows us to work with the Dirichlet to Neumann map associated with the exterior problem for the linear Helmholtz equation $\Delta u + k^2 u = 0$ on $\mathbb{R}^N \setminus B_R$ together with a suitable asymptotic condition on u . To explain this in more detail, let us suppose for a moment that the nonlinearity $f(x, u)$ is replaced by an inhomogeneous source term $f(x)$ supported in B_R . In this case, a well-studied problem is to analyze the far field expansion of the (unique) complex solution of (4) satisfying the Sommerfeld (outgoing) radiation condition

$$(7) \quad \left| \frac{\partial u}{\partial r}(x) - iku(x) \right| = o(r^{\frac{1-N}{2}}) \quad \text{as } r = |x| \rightarrow \infty.$$

This condition has been introduced in Sommerfeld's classical work [23], and it corresponds to the study of outgoing waves excited by the source term $f(x)$. Moreover, by a well-known result (going back to Rellich [21, p.58]), for given and sufficiently regular Dirichlet boundary data on $S_R := \partial B_R$ there exists a unique complex solution of $\Delta u + k^2 u = 0$ in $\mathbb{R}^N \setminus B_R$ satisfying (7). Furthermore, the corresponding Dirichlet to Neumann map T_R on S_R , also called the *capacity operator* (see [19]), is well understood and can be computed explicitly in terms of spherical harmonics, see Section 6 below. This operator assigns to a given boundary datum on S_R the normal derivative of the corresponding unique solution $\Delta u + k^2 u = 0$ in $\mathbb{R}^N \setminus B_R$ satisfying (7). However, condition (7) rules out (real-valued) standing wave solutions u which are the subject of the present paper. Nevertheless, a standing wave solution u can — as explained in [23, pp. 328–329] — be realized as a superposition of an outgoing wave and an incoming wave having opposite frequency and thus as real part of a function satisfying (7). In particular, it still satisfies the weaker asymptotic condition (5). Technically, this amounts

to working with the real part K_R of the operator T_R restricted to a Sobolev space of real functions on S_R , see Section 2 below. With the help of the operator K_R , we will be able reduce the problem of finding real solutions of (4) satisfying (5) to a variational problem in $H^1(B_R)$. While such a reduction clearly requires that f vanishes outside of B_R , we shall also see that the vanishing of f leads to difficulties in the proof of Cerami's condition which is needed to show the existence of critical points of the corresponding functional.

To state our results, we need to introduce further notation and to state our assumptions. Let 2^* denote the critical Sobolev exponent, i.e. $2^* := 2N/(N-2)$ if $N \geq 3$ and $2^* := \infty$ if $N = 1, 2$. For our main result, we shall suppose that the nonlinearity $f : \mathbb{R}^N \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous, and that there exists a bounded set $\Omega \subset \mathbb{R}^N$ of positive measure such that the following holds:

- (f₀) $f(x, u) = 0$ for $x \in \mathbb{R}^N \setminus \Omega$, $u \in \mathbb{R}$.
- (f₁) There exists $a > 0$, $p \in (2, 2^*)$ such that $|f(x, u)| \leq a(1 + |u|^{p-1})$ for every $x \in \mathbb{R}^N$, $u \in \mathbb{R}$.
- (f₂) $f(x, u) = o(|u|)$ uniformly in x as $u \rightarrow 0$.
- (f₃) $F(x, u) \geq 0$ for every $x \in \mathbb{R}^N$, $u \in \mathbb{R}$, and $F(x, u)/u^2 \rightarrow \infty$ as $|u| \rightarrow \infty$ for every $x \in \Omega$.
- (f₄) There exists $s_0 > 0$ such that for every $x \in \Omega$ we have $f(x, -s_0) \leq 0 \leq f(x, s_0)$, and the map $u \mapsto f(x, u)/|u|$ is nondecreasing on (s_0, ∞) and on $(-\infty, -s_0)$.

Here we set $F(x, u) = \int_0^u f(x, s) ds$ for $x \in \mathbb{R}^N$, $u \in \mathbb{R}$. We point out that nonlinearities of the type

$$f(x, u) = q(x)|u|^{p-2}u \quad \text{or} \quad f(x, u) = q(x)u \log(1 + |u|^s), \quad s > 0,$$

satisfy these assumptions if q is continuous and $q > 0$ on Ω , $q \equiv 0$ on $\mathbb{R}^N \setminus \Omega$. Moreover, if f satisfies (f₀)–(f₄) and if $g : \mathbb{R}^N \times \mathbb{R} \rightarrow [0, \infty)$ is continuous, vanishes outside of a bounded subset of $\Omega \times [0, \infty)$ and satisfies (f₂), then the sum $f + g$ also satisfies (f₀)–(f₄).

We need one more definition related to the asymptotics of the solutions we consider. For $R > 0$, we let \mathcal{R}_R denote the set of all functions $u \in H_{loc}^1(\mathbb{R}^N, \mathbb{R})$ such that $u = \text{Re}(w)$ in $\mathbb{R}^N \setminus B_R$ for a function $w \in \mathcal{C}(\mathbb{R}^N \setminus B_R, \mathbb{C}) \cap \mathcal{C}^\infty(\mathbb{R}^N \setminus \overline{B}_R, \mathbb{C})$ satisfying

- (i) $w \equiv u$ on S_R ,
- (ii) $\Delta w + k^2 w = 0$ in $\mathbb{R}^N \setminus \overline{B}_R$,
- (iii) $w(x) = O(r^{\frac{1-N}{2}})$ and $|\frac{\partial w}{\partial r}(x) - ikw(x)| = o(r^{\frac{1-N}{2}})$ as $r = |x| \rightarrow \infty$.

Note that every function $u \in \mathcal{R}_R$ satisfies the asymptotic conditions (5). Our main result is the following.

Theorem 1.1. *There exists a countable set $\mathcal{D} \subset (0, \infty)$ such that for $R \in (0, \infty) \setminus \mathcal{D}$ with $\Omega \subset B_R$ and every nonlinearity f satisfying assumptions (f₀) – (f₄) we have:*

- (i) Equation (4) admits a solution in \mathcal{R}_R .
- (ii) If, in addition, $f(x, -t) = -f(x, t)$ for all $(x, t) \in \mathbb{R}^N \times \mathbb{R}$, then equation (4) admits a sequence of pairs of solutions $\{\pm u_n\}$, $n \in \mathbb{N}$ in \mathcal{R}_R with the property that

$$(8) \quad \|u_n\|_{H^1(B_R)} \rightarrow \infty \quad \text{as } n \rightarrow \infty.$$

Some remarks seem in order. Elliptic regularity theory implies that the solutions given by Theorem 1.1 belong to $W_{loc}^{2,q}(\mathbb{R}^N)$ for all $1 \leq q < \infty$, so there are strong solutions of (4) in $\mathcal{C}_{loc}^{1,\alpha}(\mathbb{R}^N)$ for all $0 < \alpha < 1$. The set \mathcal{D} in Theorem 1.1 is defined by the property that an associated R -dependent linear eigenvalue problem admits the eigenvalue zero iff $R \in \mathcal{D}$, see Lemma 2.3(iv) below for details. The restriction $R \in (0, \infty) \setminus \mathcal{D}$ in Theorem 1.1 is not

necessary, but the proof is more complicated in the case where $R \in \mathcal{D}$. Since the choice of R has no impact on the validity of the asymptotic conditions (5), we decided not to consider the case $R \in \mathcal{D}$ in the present paper. Nevertheless, it is natural to ask whether different choices of R in Theorem 1.1 give rise to different solutions of problem (4), (5). The following partial answer to this question can be derived from a careful study of the explicit representation of the capacity operator (and its real part) discussed in Section 6 below.

Theorem 1.2. *For fixed $R > 0$ and $u \in \mathcal{R}_R$ with $u|_{S_R} \not\equiv 0$, there exist at most countably many $R' > R$ such that $u \in \mathcal{R}_{R'}$.*

As a consequence, the problem (4), (5) admits infinitely many solutions if the nonlinearity f satisfies assumptions $(f_0) - (f_4)$.

It might be somewhat surprising at first glance that Theorem 1.2 also gives rise to infinitely many solutions u_n , $n \in \mathbb{N}$ of (4), (5) in the case where f is *not* odd in its second variable. However, a local unboundedness property as in (8) should not be expected for these solutions. It is natural to ask whether it is possible to relax the condition that f vanishes outside a compact set. We have no general answer to this question yet, but in the case where f is radial in x , a shooting argument yields radial solutions of (4), (5) under much less restrictive assumptions on f . For a precise result, see Theorem 5.2 below. We do not even need to assume that f tends to 0 as $r = |x| \rightarrow \infty$.

The paper is organized as follows. In Section 2, we set up the variational framework used to prove Theorem 1.1. Here we also state key properties of the capacity operator T_R and its real part K_R in the case $N \geq 2$, but we postpone the derivation of these properties to Section 6 since the underlying computations — relying on special properties of Hankel functions — are somewhat technical. In Section 3 we then complete the proof of Theorem 1.1 in the case $N \geq 2$. As already remarked above, a key difficulty in the proof is the validity of Cerami's condition (see Proposition 3.2), and the proof of this property is rather long. In Section 3, we also establish — under stronger assumptions on the nonlinearity — a rigid minimax principle with respect to families of half-spaces for the solution which minimizes the corresponding energy functional among all critical points, see Theorem 3.5. We believe that this minimax principle gives rise to additional properties of the corresponding (ground state) solutions. Section 4 contains a sketch of the proof of Theorem 1.1 in the one-dimensional case. This case is much easier than the case $N \geq 2$ but has to be treated slightly differently. Section 5 is devoted to the radial case. As pointed out already, we apply a shooting argument to prove the existence of infinitely many radial solutions of (4), (5) under much less restrictive assumptions on f , see Theorem 5.2. Finally, as noted above, in Section 6 we derive key properties of the capacity operator T_R and its real part K_R . We note that some of these properties are well known (see e.g. [19] for the case $N = 3$), but we could not find an appropriate reference for general $N \geq 2$. Moreover, it seems that the operator K_R has not been studied in the degree of detail which we need for our purposes. The final part of Section 6 is devoted to the proof of Theorem 1.2.

We close this introduction with a remark on some open questions which we plan to address in future work. First, one may ask for which compactly supported nonlinearities f satisfying (f_0) there exist solutions of (4) given as superposition of a function of type (6) and a function satisfying (5). Second, one may try to prove the existence of standing wave solutions without assumption (f_0) . This would require a completely different (variational) approach. Another natural question, more closely related to the scattering theory of (1), is the existence of complex solutions of (4) satisfying the radiation condition (7). It is not clear whether solutions

can be expected in the case where $f(\cdot, 0) \equiv 0$ as assumed in (f_2) . This problem is non-variational due to the lack of symmetry of the Dirichlet to Neumann map T_R associated with (7), so a completely different approach has to be used.

2. THE VARIATIONAL FRAMEWORK

We assume that $N \geq 2$ in the next two sections, referring to Section 4 for the case $N = 1$. In this section we will introduce the capacity operator and develop a variational framework for problem (4), (5). We start by fixing some notation. Let $R > 0$ be such that $\Omega \subset B_R$, where $B_R := B_R(0)$ is the open ball with radius R centered at zero. We also set $E_R := \mathbb{R}^N \setminus \overline{B_R}$ and consider the space

$$(9) \quad H_R^{\mathbb{C}} := \left\{ u \in H_{loc}^1(E_R, \mathbb{C}) : \frac{u}{(1+r^2)^{1/2}} \in L^2(E_R, \mathbb{C}), \frac{\nabla u}{(1+r^2)^{1/2}} \in L^2(E_R, \mathbb{C}^N), \frac{\partial u}{\partial r} - iku \in L^2(E_R, \mathbb{C}) \right\},$$

where — here and in the following — r always denotes the radial variable, i.e., $r = |x|$. It is known (see [19] for the case $N = 3$ and Section 6 below for general $N \geq 2$) that for every $u \in H^{\frac{1}{2}}(S_R, \mathbb{C})$, $S_R := \partial B_R$, there exists a unique weak solution $w \in H_R^{\mathbb{C}}$ of the problem

$$(10) \quad \begin{cases} \Delta w + k^2 w = 0 & \text{in } E_R, \\ w = u & \text{on } S_R. \end{cases}$$

Here weak solution means that $\text{trace}(w) = u$ and

$$\int_{E_R} (\nabla w \cdot \nabla \phi - k^2 w \phi) dx = 0 \quad \text{for all } \phi \in \mathcal{C}_c^1(E_R).$$

where $\mathcal{C}_c^1(E_R)$ denotes the space of \mathcal{C}^1 -functions with compact support in E_R . We then define the *capacity operator* (or *Dirichlet to Neumann map*)

$$(11) \quad T_R : H^{\frac{1}{2}}(S_R, \mathbb{C}) \rightarrow H^{-\frac{1}{2}}(S_R, \mathbb{C}), \quad T_R u = \frac{\partial w}{\partial \eta} \in H^{-\frac{1}{2}}(S_R, \mathbb{C})$$

where $w \in H_R^{\mathbb{C}}$ is the unique solution of (10) corresponding to $u \in H^{\frac{1}{2}}(S_R, \mathbb{C})$ and η denotes the normal unit vector field on S_R pointing outside B_R and inside E_R .

As shown in [19] for the case $N = 3$ and detailed in Section 6 below for general $N \geq 2$, the operator $T_R : H^{\frac{1}{2}}(S_R, \mathbb{C}) \rightarrow H^{-\frac{1}{2}}(S_R, \mathbb{C})$ is continuous, and suitably normalized spherical harmonics (when considered as functions of spherical angles) form an orthonormal basis of eigenfunctions of T_R . The operator T_R is not symmetric and therefore does not give rise to a variational formulation of our nonlinear problem given by (4) and (5). Therefore, we set $H^{\frac{1}{2}}(S_R) = H^{\frac{1}{2}}(S_R, \mathbb{R})$, $H^{-\frac{1}{2}}(S_R) = H^{-\frac{1}{2}}(S_R, \mathbb{R})$, and we let

$$(12) \quad K_R : H^{\frac{1}{2}}(S_R) \rightarrow H^{-\frac{1}{2}}(S_R), \quad K_R u = \text{Re}[T_R u]$$

denote the real part of the restriction of T_R to $H^{\frac{1}{2}}(S_R)$. This operator can be seen as the Dirichlet to Neumann map corresponding to the problem (10) with real data u on S_R and solutions w given as a real part of a function in $H_R^{\mathbb{C}}$. The explicit calculations of w in terms of u in Section 6 below immediately imply that w satisfies the asymptotic conditions (5). Moreover, the operator K_R has the following properties.

Lemma 2.1. *The operator K_R is bounded, symmetric and negative definite. More precisely, there are constants $\gamma_R > 0$ and $\Gamma_R > 0$ such that*

$$(13) \quad \|K_R u\|_{H^{-\frac{1}{2}}(S_R)} \leq \Gamma_R \|u\|_{H^{\frac{1}{2}}(S_R)} \quad \text{and} \quad \int_{S_R} u K_R u \, d\sigma \leq -\gamma_R \int_{S_R} u^2 \, d\sigma$$

for all $u \in H^{\frac{1}{2}}(S_R)$. Moreover,

$$(14) \quad \int_{S_R} v K_R u \, d\sigma = \int_{S_R} u K_R v \, d\sigma \quad \text{for } u, v \in H^{\frac{1}{2}}(S_R).$$

We postpone the proof of this lemma to Section 6 below. Setting $X := H^1(B_R)$, we now recall the standard estimate

$$(15) \quad \int_{B_R} u^2 \, dx \leq c \int_{B_R} |\nabla u|^2 \, dx + c \int_{S_R} u^2 \, d\sigma \quad \text{for } u \in X$$

with some constant $c = c(R) > 0$, see e.g. [25, Theorem A.9]. Moreover, we consider the bilinear form

$$(16) \quad \mathcal{B}_k : X \times X \rightarrow \mathbb{R}, \quad \mathcal{B}_k(u, v) = \int_{B_R} (\nabla u \cdot \nabla v - k^2 uv) \, dx - \int_{S_R} v K_R u \, d\sigma.$$

From (13) and (15), we easily deduce the following

Corollary 2.2. \mathcal{B}_0 defines a scalar product on $X = H^1(B_R)$ which is equivalent to the standard scalar product, i.e. the corresponding norms are equivalent.

We also set $\mathcal{B}_k(u) := \mathcal{B}_k(u, u)$ for $u \in X$ in the following. In the next lemma, we collect key facts concerning \mathcal{B}_k and the (nonlocal) eigenvalue problem

$$(17) \quad \begin{cases} -\Delta u = \lambda u, & \text{in } B_R, \\ \frac{\partial u}{\partial \eta} = K_R u & \text{on } S_R. \end{cases}$$

Lemma 2.3. (i) *The eigenvalue problem (17) admits an unbounded sequence of eigenvalues $0 < \lambda_1 \leq \lambda_2 \leq \dots$ and a corresponding system of eigenfunctions e_j , $j \in \mathbb{N}$ which is complete in X . Moreover, each eigenfunction e_j , $j \in \mathbb{N}$ is analytic in B_R .*

(ii) *There exists a scalar product $\langle \cdot, \cdot \rangle$, equivalent to the standard scalar product on $X = H^1(B_R)$, and an orthogonal splitting $X = X^- \oplus X^0 \oplus X^+$ such that*

$$\mathcal{B}_k(u) = \|u^+\|^2 - \|u^-\|^2 \quad \text{for all } u \in X,$$

where $\|\cdot\| = \langle \cdot, \cdot \rangle^{\frac{1}{2}}$ is the induced norm, and u^\pm, u^0 denote the corresponding orthogonal projections of u onto X^\pm, X^0 , respectively. More precisely,

$$X^- = \text{span}\{e_j : j \leq j_*\}, X^0 = \text{span}\{e_j : j_* < j < j^*\} \text{ and } X^+ = \overline{\text{span}\{e_j : j \geq j^*\}},$$

where $j_* := \max\{j \in \mathbb{N} : \lambda_j < k^2\}$ and $j^* := \min\{j \in \mathbb{N} : \lambda_j > k^2\}$. In particular, X^- and X^0 are finite dimensional.

(iii) *The family e_j , $j \in \mathbb{N}$ is orthogonal with respect to the scalar products $\langle \cdot, \cdot \rangle$, \mathcal{B}_0 and the scalar product of $L^2(B_R)$.*

(iv) *There exists a countable set $\mathcal{D} \subset (0, \infty)$ such that $X^0 \neq \{0\}$ if and only if $R \in \mathcal{D}$.*

Proof. (i) We first consider $H^1(B_R)$ with the equivalent scalar product \mathcal{B}_0 . Since K_R is negative definite, all eigenvalues of (17) must be positive. Hence $u \in H^1(B_R)$ is an eigenfunction

of (17) corresponding to $\lambda \in \mathbb{R}$ if and only if $\lambda > 0$ and u is an eigenfunction of the operator $K \in \mathcal{L}(X)$ defined by

$$(18) \quad \mathcal{B}_0(Ku, v) = \int_{B_R} uv \, dx \quad \text{for } u, v \in X$$

corresponding to the eigenvalue $\frac{1}{\lambda}$. The operator K is bounded, symmetric with respect to \mathcal{B}_0 , nonnegative and compact, since the embedding $X \hookrightarrow L^2(B_R)$ is compact. Moreover, 0 is not an eigenvalue of K . Hence K admits a sequence of positive eigenvalues $\mu_1 \geq \mu_2 \geq \dots$ such that $\mu_j \rightarrow 0$ as $j \rightarrow \infty$, and a corresponding complete system of \mathcal{B}_0 -orthogonal eigenfunctions e_j , $j \in \mathbb{N}$. Assertion (i) now follows with $\lambda_j = \frac{1}{\mu_j}$, $j \in \mathbb{N}$ and the family $\{e_j : j \in \mathbb{N}\}$ thus obtained.

(ii) For $u \in X$, let u^\pm, u^0 denote the $L^2(B_R)$ -orthogonal projections of u onto the subspaces X^\pm, X^0 , respectively, as defined in the assertion. For $u, v \in X$, we define

$$\langle u, v \rangle = \mathcal{B}_k(u^+, v^+) - \mathcal{B}_k(u^-, v^-) + \int_{B_R} u^0 v^0 \, dx.$$

It is easy to see that this scalar product has the desired properties, and by construction the splitting $X = X^- \oplus X^0 \oplus X^+$ is orthogonal with respect to this scalar product.

(iii) The \mathcal{B}_0 -orthogonality of the family $\{e_j : j \in \mathbb{N}\}$ has already been shown above, and the $L^2(B_R)$ -orthogonality then follows from (18). From this, the orthogonality with respect to $\langle \cdot, \cdot \rangle$ immediately follows by definition.

(iv) This part, which relies on special properties of the capacity operator and Hankel functions, will be proved in Section 6 below. \square

We now consider the functional

$$(19) \quad \Phi : X \rightarrow \mathbb{R}, \quad \Phi(u) = \frac{1}{2} \mathcal{B}_k(u) - \varphi(u) = \frac{1}{2} (\|u^+\|^2 - \|u^-\|^2) - \varphi(u),$$

where $\varphi(u) = \int_{B_R} F(x, u(x)) \, dx$ for $u \in X$ and $F(x, t) = \int_0^t f(x, s) \, ds$ for $t \in \mathbb{R}$. It is well known that $\varphi \in \mathcal{C}^1(X, \mathbb{R})$ as a consequence of assumption (f_1) , and

$$(20) \quad \varphi \text{ is nonnegative on } X$$

by (f_3) . Moreover, the critical points of Φ correspond to restrictions to B_R of solutions of (4). Indeed, if $u \in X$ is a critical point of Φ , then

$$0 = \int_{B_R} (\nabla u \cdot \nabla w - k^2 u w - f(x, u) w) \, dx - \int_{S_R} w K_R u \, d\sigma$$

for every $w \in X$, hence u is a weak solution of the problem

$$(21) \quad \begin{cases} -\Delta u - k^2 u = f(x, u) & \text{in } B_R, \\ \frac{\partial u}{\partial \eta} = K_R u & \text{on } S_R. \end{cases}$$

As explained in Section 6, extending u by the real part of the unique solution of (10) in $H_R^{\mathcal{C}}$ then yields a solution of (4), (5).

3. PROOF OF THEOREM 1.1

We fix $R \in (0, \infty) \setminus \mathcal{D}$ from now on, so that $X^0 = \{0\}$ by Lemma 2.3. As remarked in the introduction, this restriction is made only in order to simplify the proofs, whereas all results can still be proved — with additional effort — in the case where X^0 has positive dimension.

We first collect useful facts about the functional Φ defined in (19). For this we recall the following well-known consequence of (f_1) and (f_2) :

$$(22) \quad \forall \varepsilon > 0, \exists C_\varepsilon > 0 \text{ such that } |f(x, u)| \leq \varepsilon|u| + C_\varepsilon|u|^{p-1}, \forall (x, u) \in \mathbb{R}^N \times \mathbb{R}.$$

Lemma 3.1. (i) *There exists $\alpha_0 > 0$ such that $\inf_{\Sigma_\alpha} \Phi > 0$ for $\alpha \in (0, \alpha_0)$, where $\Sigma_\alpha := \{u \in X^+ : \|u\| = \alpha\}$.*

(ii) *Let \mathcal{Z} be a closed cone contained in a finite-dimensional subspace W of X and such that*

$$(23) \quad \begin{aligned} & \{x \in \Omega : w(x) \neq 0\} \text{ has positive measure for} \\ & \text{every } w \in \mathcal{Z} \setminus \{0\} \text{ with } \|w^+\| \geq \|w^-\|. \end{aligned}$$

Then there exists $\rho = \rho(\mathcal{Z}) > 0$ such that $\Phi(u) \leq 0$ for all $u \in \mathcal{Z}$ satisfying $\|u\| \geq \rho$.

Here and in the following, a set $\mathcal{Z} \subset X$ is called a cone if $\lambda x \in \mathcal{Z}$ for every $x \in \mathcal{Z}$, $\lambda \geq 0$. In particular, (ii) applies to $\mathcal{Z} = W$ if W is a finite-dimensional subspace of X .

Proof. (i) For $u \in X^+$ we have $\Phi(u) = \frac{1}{2}\|u\|^2 - \varphi(u)$ and $\varphi(u) = o(\|u\|^2)$ as $u \rightarrow 0$ by (22) and Sobolev embeddings. Hence the conclusion follows.

(ii) Suppose by contradiction that a sequence $(u_n)_n \subset \mathcal{Z}$ exists with $\Phi(u_n) > 0$ for all n and $\|u_n\| \rightarrow \infty$ as $n \rightarrow \infty$. Setting $w_n = \frac{u_n}{\|u_n\|}$, we may pass to a subsequence such that $w_n \rightarrow w \in W$ since W is finite-dimensional. Since $w_n \in \mathcal{Z}$ and $\|w_n\| = 1$ for all n , we have $w \in \mathcal{Z}$ and $\|w\| = 1$. Moreover, by (20) we have

$$0 \leq \liminf_{n \rightarrow \infty} \frac{\Phi(u_n)}{\|u_n\|^2} \leq \frac{1}{2} \lim_{n \rightarrow \infty} (\|w_n^+\|^2 - \|w_n^-\|^2) = \frac{1}{2} (\|w^+\|^2 - \|w^-\|^2)$$

and therefore $\|w^+\| \geq \|w^-\|$. Hence (23) implies that $\Omega_w := \{x \in \Omega : w(x) \neq 0\}$ has positive measure. Passing to a subsequence, we may also assume that $w_n \rightarrow w$ pointwise a.e. in B_R , which implies that

$$|u_n(x)| = \|u_n\| |w_n(x)| \rightarrow \infty \quad \text{as } n \rightarrow \infty \text{ for a.e. } x \in \Omega_w.$$

By (f_3) and Fatou's Lemma, we therefore deduce that

$$0 \leq \frac{\Phi(u_n)}{\|u_n\|^2} \leq \frac{1}{2} (\|w_n^+\|^2 - \|w_n^-\|^2) - \int_{\Omega_w} \frac{F(x, u_n)}{u_n^2} w_n^2 dx \rightarrow -\infty$$

as $n \rightarrow \infty$. This contradiction proves the claim. \square

To proceed with the proof of Theorem 1.1, we shall decompose the nonlinearity f as follows. We write $f = f_1 + f_2$, where $f_i : \mathbb{R}^N \times \mathbb{R} \rightarrow \mathbb{R}$, $i = 1, 2$ are defined by

$$f_1(x, u) = \begin{cases} f(x, u), & |u| \geq s_0, \\ \frac{u^2}{s_0^2} f(x, s_0), & 0 \leq u \leq s_0, \\ \frac{u^2}{s_0^2} f(x, -s_0), & -s_0 \leq u \leq 0, \end{cases}$$

and we put $f_2 = f - f_1$. Setting $F_i(x, u) = \int_0^u f_i(x, s) ds$, $i = 1, 2$ for $x \in \mathbb{R}^N$, $u \in \mathbb{R}$, we see that

$$(24) \quad \begin{aligned} f_2(x, u) &= 0 \text{ if } x \in \mathbb{R}^N \setminus \Omega \text{ or } |u| \geq s_0; \\ F_2(x, u) &= 0 \text{ if } x \in \mathbb{R}^N \setminus \Omega, u \in \mathbb{R}; \\ f_2 \text{ and } F_2 &\text{ are bounded on } \mathbb{R}^N \times \mathbb{R}. \end{aligned}$$

Moreover, f_1 satisfies condition (f_2) and the following stronger version of condition (f_4) . For every $x \in \mathbb{R}^N$,

$$(25) \quad u \mapsto \frac{f_1(x, u)}{|u|} \text{ is nondecreasing on } \mathbb{R} \setminus \{0\}.$$

We decompose the functional $\varphi : X \rightarrow \mathbb{R}$ accordingly and write $\varphi = \varphi_1 + \varphi_2$ with

$$\varphi_i(u) = \int_{B_R} F_i(x, u(x)) dx = \int_{\Omega} F_i(x, u(x)) dx, \quad i = 1, 2.$$

We note that φ_2 is bounded on X by (24). The following proposition will be a main technical step in the proof of Theorem 1.1.

Proposition 3.2. *Φ satisfies the Cerami condition in X , i.e., every sequence $(u_n)_n \subset X$ such that $\Phi(u_n) \rightarrow c$ for some $c \in \mathbb{R}$ and $(1 + \|u_n\|)\|\Phi'(u_n)\| \rightarrow 0$ as $n \rightarrow \infty$ has a subsequence which converges in X .*

The proof of this proposition is quite long and requires subtle estimates. Parts of the proof are inspired by [16] and [17], but we need new arguments to deal with the difficulty that the nonlinearity may vanish on a subset of B_R of positive measure. A key role in the proof is played by the useful inequality

$$(26) \quad \begin{aligned} f_1(x, u)[s(\frac{s}{2} + 1)u + (1 + s)v] + F_1(x, u) - F_1(x, [1 + s]u + v) &\leq 0 \\ \text{for } x \in \mathbb{R}^N, u, v \in \mathbb{R} \text{ and } s &\geq -1, \end{aligned}$$

which follows from (25). Indeed, as noted in [17], this inequality is a weak version of [27, Lemma 2.2]. As a consequence of (26) and of the properties of f_2 , we may derive the following

Lemma 3.3. *For every $K > 0$ there is a constant $C = C(K) > 0$ with the following property. If $u, s, v \in \mathbb{R}$ are numbers with $-1 \leq s \leq K$ and $|v| \leq K$, then*

$$f(x, u)[s(\frac{s}{2} + 1)u + (1 + s)v] + F(x, u) - F(x, [1 + s]u + v) \leq C \quad \text{for all } x \in \mathbb{R}^N.$$

Proof. By (24) there exists a constant $C_1 > 0$ (depending on K) such that

$$|f_2(x, u)| \leq C_1, \quad |F_2(x, u)| \leq C_1 \quad \text{and} \quad |f_2(x, u)s(\frac{s}{2} + 1)u| \leq C_1$$

for $u \in \mathbb{R}$, $x \in \mathbb{R}^N$, $|s| \leq K + 1$. Consequently, we have

$$\begin{aligned} f_2(x, u)[s(\frac{s}{2} + 1)u + (1 + s)v] + F_2(x, u) - F_2(x, [1 + s]u + v) \\ \leq C_1[1 + K(K + 1)] + 2C_1 \end{aligned}$$

for $x \in \mathbb{R}^N$ and $u, s, v \in \mathbb{R}$ with $-1 \leq s \leq K$ and $|v| \leq K$. Since $f = f_1 + f_2$, $F = F_1 + F_2$ and (26) holds, the claim follows with $C := C_1[1 + K(K + 1)] + 2C_1$. \square

The next step in the proof of Proposition 3.2 is the following relative energy estimate. In the following, a sequence $(u_n)_n$ in X is called a *Cerami sequence* for Φ if $\Phi(u_n) \rightarrow c$ for some $c \in \mathbb{R}$ and $(1 + \|u_n\|)\|\Phi'(u_n)\| \rightarrow 0$ as $n \rightarrow \infty$.

Lemma 3.4. *For every $\kappa > 0$ there exists $\tilde{C} = \tilde{C}(\kappa) > 0$ with the following property. If $(u_n)_n$ is a Cerami sequence for Φ , and $r_n \geq 0$, $v_n \in X^-$, $n \in \mathbb{N}$ are given with $r_n \leq \kappa$ and $\|v_n\| \leq \kappa$ for all $n \in \mathbb{N}$, then*

$$\Phi(r_n u_n + v_n) \leq \Phi(u_n) + \tilde{C} + o(1) \quad \text{as } n \rightarrow \infty.$$

Proof. We first note that, by a standard bootstrap argument using elliptic regularity theory, there exists a constant $K = K(\kappa) > \kappa$ such that

$$\|v\|_{L^\infty(B_R)} \leq K \quad \text{for every } v \in X^- \text{ with } \|v\| \leq \kappa.$$

We write $r_n = 1 + s_n$ with $-1 \leq s_n \leq \kappa - 1 \leq K$ and set $w_n = r_n u_n + v_n = (1 + s_n)u_n + v_n$ for $n \in \mathbb{N}$. Then

$$\begin{aligned} \Phi(w_n) - \Phi(u_n) &= \frac{1}{2}[\mathcal{B}_k(w_n) - \mathcal{B}_k(u_n)] + \int_{B_R} (F(x, u_n) - F(x, w_n)) dx \\ &= \frac{1}{2} \left([(1 + s_n)^2 - 1]\mathcal{B}_k(u_n) + 2(1 + s_n)\mathcal{B}_k(u_n, v_n) + \mathcal{B}_k(v_n) \right) \\ &\quad + \int_{B_R} (F(x, u_n) - F(x, w_n)) dx \\ &= -\frac{\|v_n\|^2}{2} + \mathcal{B}_k \left(u_n, s_n \left(\frac{s_n}{2} + 1 \right) u_n + (1 + s_n)v_n \right) + \int_{B_R} (F(x, u_n) - F(x, w_n)) dx \\ &\leq \mathcal{B}_k \left(u_n, s_n \left(\frac{s_n}{2} + 1 \right) u_n + (1 + s_n)v_n \right) + \int_{B_R} (F(x, u_n) - F(x, w_n)) dx \end{aligned}$$

Since $(u_n)_n$ is a Cerami sequence, $\|v_n\| \leq \kappa$ and $|s_n| \leq K + 1$ for all n , we have

$$\begin{aligned} &\left| \mathcal{B}_k \left(u_n, s_n \left(\frac{s_n}{2} + 1 \right) u_n + (1 + s_n)v_n \right) - \int_{\mathbb{R}^N} f(x, u_n) \left[s_n \left(\frac{s_n}{2} + 1 \right) u_n + (1 + s_n)v_n \right] dx \right| \\ &= \left| \Phi'(u_n) \left(s_n \left(\frac{s_n}{2} + 1 \right) u_n + (1 + s_n)v_n \right) \right| \\ &\leq c_1 \|\Phi'(u_n)\| \|u_n\| + c_2 \|\Phi'(u_n)\| \|v_n\| = o(1) \end{aligned}$$

as $n \rightarrow \infty$ with constants $c_1, c_2 > 0$ (depending on K). Consequently,

$$\begin{aligned} \Phi(w_n) - \Phi(u_n) &\leq \int_{B_R} \left(f(x, u_n) \left[s_n \left(\frac{s_n}{2} + 1 \right) u_n + (1 + s_n)v_n \right] \right. \\ &\quad \left. + F(x, u_n) - F(x, w_n) \right) dx + o(1). \end{aligned}$$

Choosing $C = C(K)$ as in Lemma 3.3, we conclude that

$$\Phi(w_n) - \Phi(u_n) \leq |B_R|C + o(1) \quad \text{as } n \rightarrow \infty.$$

Hence the assertion follows with $\tilde{C} = |B_R|C + 1$. \square

We may now complete the

Proof of Proposition 3.2. Let $(u_n)_n$ be a sequence with the assumed properties. We first show that $(u_n)_n$ is bounded in X . Assuming by contradiction that this is false, we may pass to a subsequence — still denoted by $(u_n)_n$ — such that $\|u_n\| \rightarrow \infty$ as $n \rightarrow \infty$. Setting

$w_n := \frac{u_n}{\|u_n\|}$, we may assume, passing again to a subsequence, that $w_n \rightharpoonup w$ weakly in X for some $w \in X$, $w_n \rightarrow w$ in $L^q(B_R)$ for all $1 \leq q < 2^*$, and $w_n \rightarrow w$ pointwise a.e. on B_R as $n \rightarrow \infty$. Moreover, we have $w_n^+ \rightharpoonup w^+$ weakly in X and $w_n^- \rightarrow w^-$ strongly in X as $n \rightarrow \infty$, since X^- is finite-dimensional. Passing to a further subsequence, we may also assume that $\|w_n^+\| \rightarrow s \geq 0$ as $n \rightarrow \infty$ for some $s \geq \|w^+\|$. Since

$$o(1) = \frac{\Phi(u_n)}{\|u_n\|^2} \leq \frac{1}{2} (\|w_n^+\|^2 - \|w_n^-\|^2),$$

by (20), we find that

$$(27) \quad \|w^-\| \leq s.$$

Hence, $1 = \|w_n\|^2 = \|w_n^+\|^2 + \|w_n^-\|^2 \rightarrow s^2 + \|w^-\|^2 \leq 2s^2$ as $n \rightarrow \infty$, indicating that $s > 0$. Next we suppose by contradiction that

$$(28) \quad w \equiv 0 \quad \text{a.e. on } \Omega$$

which implies that

$$(29) \quad \varphi(tw_n) \rightarrow 0 \quad \text{as } n \rightarrow \infty \text{ for all } t > 0.$$

We claim that there exist $t > 0$ and $v_n \in X^-$, $n \in \mathbb{N}$ with

$$(30) \quad \|v_n\| \leq 1 \quad \text{and} \quad \Phi(tw_n + v_n) > c + \tilde{C} + 1 \quad \text{for } n \text{ sufficiently large,}$$

where $\tilde{C} = \tilde{C}(1)$ is chosen as in Lemma 3.4 corresponding to $\kappa = 1$. To prove this, we have to distinguish different cases. We first note that

$$\Phi(tw_n) = \frac{t^2}{2} (s^2 - \|w^-\|^2) + o(1) \quad \text{as } n \rightarrow \infty$$

for every $t > 0$ by (29). Hence, if $\|w^-\| < s$, we can find $t > 0$ such that

$$\Phi(tw_n) > c + \tilde{C} + 1 \quad \text{for } n \text{ sufficiently large,}$$

and therefore (30) follows with $v_n = 0$ for every n .

Next we consider the remaining case $\|w^-\| = s$, and we note that for every $t > 0$ we have $tw_n + w_n^+ \rightarrow tw + w^+$ in $L^q(B_R)$ for all $1 \leq q < 2^*$ and also pointwise a.e. on B_R . Hence

$$(31) \quad \varphi(tw_n + w_n^+) \rightarrow \varphi(tw + w^+) = \varphi(w^+) \quad \text{as } n \rightarrow \infty$$

for all $t > 0$ by (28). Consequently,

$$\Phi(tw_n + w_n^+) = \frac{1}{2} [s^2(t+1)^2 - t^2s^2] - \varphi(w^+) + o(1) = s^2 \left(t + \frac{1}{2} \right) - \varphi(w^+) + o(1),$$

so that there exists $t > 0$ such that

$$\Phi(tw_n + w_n^+) > c + \tilde{C} + 1 \quad \text{for } n \text{ sufficiently large.}$$

Again, (30) follows with $t+1$ in place of t and $v_n = -w_n^-$, since $tw_n + w_n^+ = (t+1)w_n - w_n^-$ for every n . Next, fixing $t > 0$ and v_n , $n \in \mathbb{N}$ such that (30) holds, we write $tw_n + v_n = s_n u_n + v_n$ with $s_n = \frac{t}{\|u_n\|}$ for every n , so that $0 < s_n \leq 1$ for large n and $\|v_n\| \leq \|w_n\| = 1$. By Lemma 3.4, we therefore have

$$\Phi(tw_n + v_n) = \Phi(s_n u_n + v_n) \leq \Phi(u_n) + \tilde{C} + o(1)$$

as $n \rightarrow \infty$, which contradicts (30). The contradiction shows that (28) is false, and therefore the set $\Omega_w := \{x \in \Omega : w(x) \neq 0\}$ has positive measure. Moreover,

$$|u_n(x)| = \|u_n\| |w_n(x)| \rightarrow +\infty \text{ as } n \rightarrow \infty \text{ for almost every } x \in \Omega_w.$$

Hence, Fatou's Lemma, the L^2 -convergence $w_n \rightarrow w$ in B_R and (f_3) imply

$$o(1) = \frac{\Phi(u_n)}{\|u_n\|^2} \leq \frac{1}{2} - \int_{\Omega_w} \frac{F(x, u_n)}{|u_n|^2} |w_n|^2 dx \rightarrow -\infty,$$

as $n \rightarrow \infty$, a contradiction. The contradiction shows that $(u_n)_n$ is bounded in X . Therefore, we can find a subsequence — still denoted by $(u_n)_n$ — and some $u \in X$ such that $u_n \rightharpoonup u$ (weakly) in X , $u_n \rightarrow u$ in $L^q(B_R)$ for all $1 \leq q < 2^*$ and $u_n(x) \rightarrow u(x)$ for a.e. $x \in B_R$. As a consequence of (22), there holds

$$\int_{B_R} (f(x, u_n) - f(x, u))(u_n - u) dx \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Hence,

$$\begin{aligned} \mathcal{B}_0(u_n - u, u_n - u) &= (\Phi'(u_n) - \Phi'(u))(u_n - u) + k^2 \int_{B_R} (u_n - u)^2 dx \\ &\quad + \int_{B_R} (f(x, u_n) - f(x, u))(u_n - u) dx \rightarrow 0, \end{aligned}$$

as $n \rightarrow \infty$. It follows from Corollary 2.2 that $u_n \rightarrow u$ strongly in X . The proof is finished. \square

We may now complete the

Proof of Theorem 1.1 (Case $N \geq 2$). (i) The existence of a nontrivial solution follows from a variant of the classical linking theorem where the Palais–Smale condition is replaced by the Cerami condition [4, Theorem 2.3]. To see this, we proceed as follows. Considering the sequence of eigenfunctions $(e_j)_{j \in \mathbb{N}}$ of (17) given by Lemma 2.3, we set $u = e_{j^*} \in X^+$ and put

$$Q_\rho := \{tu + v : v \in X^-, \|v\| \leq \rho, 0 \leq t \leq \rho\} \quad \text{for } \rho > 0.$$

Note that the sets Q_ρ are contained in the finite dimensional subspace $W = X^- \oplus \mathbb{R}u \subset X$. Since every function in W is analytic in B_R , $\mathcal{Z} := W$ satisfies condition (23). Using Lemma 3.1 (ii) and the fact that Φ is nonpositive on X^- by (20), we thus find that

$$\sup_{\partial Q_\rho} \Phi = 0 \quad \text{for } \rho > 0 \text{ sufficiently large.}$$

According to Lemma 3.1 (i), we may further choose $\alpha > 0$ sufficiently small such that the sets Σ_α and ∂Q_ρ link and $\inf_{\Sigma_\alpha} \Phi > 0$ (see e.g. [25] or [4, Section 2] for the notion of linking of sets). Finally, we have $\sup_{Q_\rho} \Phi < +\infty$ by the compactness of Q_ρ . Taking Proposition 3.2 into account, we can apply the linking theorem and obtain that Φ has a nontrivial critical point $\hat{u} \in X$ such that

$$0 < \inf_{\Sigma_\alpha} \Phi \leq \Phi(\hat{u}) \leq \sup_{Q_\rho} \Phi.$$

(ii) Let us now assume that $f(x, -t) = -f(x, t)$ for all $(x, t) \in \mathbb{R}^N \times \mathbb{R}$. In this case, a variant of the Fountain Theorem (see [5, 7] and [30, Theorem 3.6]) yields the existence of a sequence of pairs $\{\pm u_n\}$, $n \in \mathbb{N}$ of critical points of Φ such that

$$(32) \quad \Phi(u_n) \rightarrow \infty \quad \text{as } n \rightarrow \infty.$$

More precisely, we use a version of the Fountain Theorem where the Cerami condition is used in place of the Palais–Smale condition. To see that such a variant exists, it suffices to note the validity of a deformation lemma giving rise to Cerami sequences instead of Palais–Smale sequences. Such a deformation lemma has already been established in [4, Theorem 1.3]. In

order to check the other assumptions of the Fountain Theorem, we remark that $X = \bigoplus_{j \in \mathbb{N}} \mathbb{R}e_j$ where $(e_j)_{j \in \mathbb{N}}$ is given by Lemma 2.3 (i). We set

$$X_j = \mathbb{R}e_j, \quad Y_j = \bigoplus_{\ell=1}^j X_\ell \quad \text{and} \quad Z_j = \overline{\bigoplus_{\ell=j}^{\infty} X_\ell}$$

for $j \in \mathbb{N}$. Since every function in Y_j is analytic, we see from Lemma 3.1 (ii), applied to $\mathcal{Z} = Y_j$, that for every $j \in \mathbb{N}$ there exists $\rho_j > 0$ such that $\Phi(u) \leq 0$ for $u \in Y_j$ with $\|u\| \geq \rho_j$. It only remains to check that for some sequence $(r_j)_j \subset (0, \infty)$

$$(33) \quad \inf\{\Phi(u) : u \in Z_j, \|u\| = r_j\} \rightarrow \infty \text{ as } j \rightarrow \infty.$$

This will be proved by similar arguments as in [30, Theorem 3.7]. Indeed, if $j \geq j^*$, then $Z_j \subset X^+$ and therefore, by (22),

$$(34) \quad \begin{aligned} \Phi(u) &= \frac{1}{2}\|u\|^2 - \int_{B_R} F(x, u) \, dx \geq \frac{1}{2}\|u\|^2 - \frac{\varepsilon}{2}\|u\|_{L^2(B_R)}^2 - \frac{C_\varepsilon}{p}\|u\|_{L^p(B_R)}^p \\ &\geq \frac{1}{4}\|u\|^2 - \frac{C_\varepsilon}{p}\beta_j^p\|u\|^p \quad \text{for } u \in Z_j, \end{aligned}$$

where

$$\varepsilon = \frac{1}{2} \inf_{u \in X \setminus \{0\}} \frac{\|u\|^2}{\|u\|_{L^2(B_R)}^2} > 0 \quad \text{and} \quad \beta_j := \sup\{\|u\|_{L^p(B_R)} : u \in Z_j, \|u\| = 1\}.$$

Since $(\beta_j)_j \subset [0, \infty)$ is a decreasing sequence, $\beta := \lim_{j \rightarrow \infty} \beta_j$ exists. Moreover, for each j we can find some $u_j \in Z_j$ such that $\|u_j\| = 1$ and $\beta_j \geq \|u_j\|_{L^p(B_R)} > \frac{\beta_j}{2}$. From the definition of Z_j we obtain that $u_j \rightarrow 0$ (weakly) in X and the compact Sobolev embedding $X \hookrightarrow L^p(B_R)$ then gives $u_j \rightarrow 0$ in $L^p(B_R)$. Thus, $\beta_j \rightarrow 0$ as $j \rightarrow \infty$. Choosing $r_j = (2C_\varepsilon\beta_j^p)^{\frac{1}{2-p}}$, we obtain from (34) that

$$\Phi(u) \geq r_j^2 \left(\frac{1}{4} - \frac{1}{2p} \right) \quad \text{for all } u \in Z_j \text{ with } \|u\| = r_j.$$

Since $r_j \rightarrow \infty$ as $j \rightarrow \infty$, the assertion follows. Moreover, since Φ is bounded on bounded subsets of $X = H^1(B_R)$, (32) implies that $\|u_n\|_{H^1(B_R)} \rightarrow \infty$ as $n \rightarrow \infty$, as claimed in (8).

To conclude the proof of Theorem 1.1, we remark that if u is a critical point of Φ , then the restriction of u to S_R belongs to $H^{\frac{1}{2}}(S_R)$ and there exists a unique weak solution w of (10). Therefore, extending u on \mathbb{R}^N by setting $u \equiv \text{Re}(w)$ on E_R , we see that $u \in \mathcal{R}_R$ is a solution of (4). \square

We close this section with an observation on a rigid minimax characterization of the ground state energy level (i.e., the least energy of a nontrivial critical point of Φ) in the case where (f_4) is replaced by a stronger condition.

Theorem 3.5. *Suppose that the nonlinearity f satisfies $(f_1) - (f_3)$ and the following stronger version of (f_4) : For every $x \in \mathbb{R}^N$,*

$$(35) \quad u \mapsto \frac{f(x, u)}{|u|} \text{ is nondecreasing on } \mathbb{R} \setminus \{0\}.$$

Then the ground state energy level

$$c := \inf\{\Phi(u) : u \text{ nontrivial critical point of } \Phi\}$$

is positive and equivalently given by

$$c = \inf_{u \in X^+ \setminus \{0\}} \sup_{t \geq 0, v \in X^-} \Phi(tu + v)$$

Moreover, c is attained within the set of nontrivial critical points of Φ , i.e., within the set of nontrivial solutions of (21).

Proof. Let u be a nontrivial critical point of Φ . Then $u \notin X^-$, since otherwise

$$\int_{B_R} f(x, u)u \, dx = \mathcal{B}_k(u) = -\|u^-\|^2 < 0,$$

contrary to (35) and (f_2) . Next we claim that

$$(36) \quad \Phi(u) \geq \Phi(w) \quad \text{for every } w \in X^- + \mathbb{R}^+u = X^- + \mathbb{R}^+u^+.$$

Indeed, similar to the proof of Lemma 3.4 we have, for $w = (1 + s)u + v$ with $s \geq -1$ and $v \in X^-$,

$$\begin{aligned} \Phi(w) - \Phi(u) &\leq \mathcal{B}_k(u, s(\frac{s}{2} + 1)u + (1 + s)v) + \int_{B_R} (F(x, u) - F(x, w)) \, dx \\ &= \int_{B_R} \left(f(x, u)[s(\frac{s}{2} + 1)u + (1 + s)v] + F(x, u) - F(x, w) \right) \, dx. \end{aligned}$$

Moreover, the last integral is nonpositive since, as a consequence of (35), the inequality (26) holds for f in place of f_1 . Hence (36) is true. Putting

$$c^* = \inf_{u \in X^+ \setminus \{0\}} \sup_{t \geq 0, v \in X^-} \Phi(tu + v),$$

we thus infer that $c \geq c^*$, whereas Lemma 3.1 (i) implies that $c^* \geq \inf_{\Sigma_\alpha} \Phi > 0$ for some $\alpha > 0$,

where $\Sigma_\alpha := \{u \in X^+ : \|u\| = \alpha\}$.

Furthermore, c is attained among nontrivial critical points of Φ . Indeed, if $(u_n)_n$ is a sequence of critical points of Φ with $\Phi(u_n) \rightarrow c$, then by Proposition 3.2 we have $u_n \rightarrow u_0$ in X for a subsequence, where u_0 is a critical point of Φ with $\Phi(u_0) = c > 0$, and therefore $u_0 \neq 0$.

It remains to show that $c \leq c^*$. For this, let $u \in X^+ \setminus \{0\}$ be such that

$$(37) \quad \sup_{t \geq 0, v \in X^-} \Phi(tu + v) < \infty.$$

Let $\mathcal{Z} := \{tu + v : v \in X^-, t \geq 0\}$. We claim that condition (23) holds for this set \mathcal{Z} . Suppose by contradiction that this is false, i.e., there exists $w \in \mathcal{Z} \setminus \{0\}$ with $\|w^+\| \geq \|w^-\|$ and such that $w \equiv 0$ a.e. on Ω . Since $\|w^+\| > 0$, we then find

$$\begin{aligned} \Phi(tw + w^+) &= \frac{1}{2} \left((t+1)^2 \|w^+\|^2 - t^2 \|w^-\|^2 \right) - \varphi(w^+) \\ &\geq \left(t + \frac{1}{2} \right) \|w^+\|^2 - \varphi(w^+) \rightarrow \infty, \end{aligned}$$

as $t \rightarrow \infty$. This contradicts (37), since $tw + w^+ = (t+1)w^+ + tw^- \in \mathcal{Z}$ for every $t > 0$. The contradiction shows that condition (23) holds for \mathcal{Z} . Since \mathcal{Z} is a closed cone contained in the finite dimensional space $X^- \oplus \mathbb{R}u$, Lemma 3.1 (ii) implies that there exists $\rho > 0$ with

$$(38) \quad \sup_{\partial Q_\rho} \Phi = 0 \quad \text{where} \quad Q_\rho := \{tu + v : v \in X^-, \|v\| \leq \rho, 0 \leq t \leq \rho\}.$$

Applying the linking theorem [4, Theorem 2.3] exactly as in the proof of Theorem 1.1 with the present choice of Q_ρ , we obtain a nontrivial critical point \hat{u} of Φ with

$$\Phi(\hat{u}) \leq \max_{Q_\rho} \Phi \leq \sup_{t \geq 0, v \in X^-} \Phi(tu + v).$$

We thus conclude that $c \leq c^*$, so equality holds and the proof is finished. \square

4. THE ONE-DIMENSIONAL CASE

In this section we sketch the proof of Theorem 1.1 in the case $N = 1$ which has to be treated slightly differently. Note that for $R > 0$ we have $B_R = (-R, R)$, $S_R := \{-R, R\}$ and $E_R = (-\infty, -R) \cup (R, +\infty)$, and consider the space $H_R^\mathbb{C}$ as defined in (9). For a given function $u : S_R \rightarrow \mathbb{C}$, it is easy to see that the unique solution $w \in H_R^\mathbb{C}$ of the problem

$$(39) \quad \begin{cases} w'' + k^2 w = 0 & \text{in } E_R, \\ w = u & \text{on } S_R \end{cases}$$

is then given by

$$w(x) = \begin{cases} u(-R)e^{-ik(x+R)} & \text{for } x \leq -R, \\ u(R)e^{ik(x-R)} & \text{for } x \geq R. \end{cases}$$

Hence the capacity operator T_R is simply given by $[T_R u](R) = ik u(R)$ and $[T_R u](-R) = ik u(-R)$ for any function $u : S_R \rightarrow \mathbb{C}$, and thus its real part K_R , as defined in (12), is zero. In particular, the second inequality in (13) is not true if $N = 1$, and also Corollary 2.2 does not hold in this case. On the other hand, the quadratic form \mathcal{B}_k defined in (16) is simply given by

$$\mathcal{B}_k(u, v) = \int_{-R}^R (u'v' - k^2 uv) dx, \quad u, v \in H^1(-R, R),$$

and (17) reduces to the homogeneous Neumann eigenvalue problem

$$(40) \quad -u'' = \lambda u \quad \text{in } (-R, R), \quad u'(\pm R) = 0.$$

As a consequence, the properties listed in Lemma 2.3 are well known and easy to check in the case $N = 1$. Furthermore, the inner problem (21) reduces to

$$(41) \quad \begin{cases} -u'' - k^2 u = f(x, u) & \text{in } (-R, R), \\ u'(\pm R) = 0, \end{cases}$$

and solutions of (41) correspond to critical points of the functional

$$\Phi : H^1(-R, R) \rightarrow \mathbb{R}, \quad \Phi(u) = \frac{1}{2} \int_{-R}^R \left((u')^2 - k^2 u^2 \right) dx - \int_{-R}^R F(x, u) dx.$$

Since the structure of Φ is the same as in the multidimensional case, the proof of Theorem 1.1 can be finished exactly as detailed in Section 3. In particular, the one-dimensional analogues of Lemma 3.1 and Proposition 3.2 are valid.

In addition, we remark that, by exactly the same proof, Theorem 3.5 is also true in the one-dimensional case.

5. EXISTENCE OF RADIAL SOLUTIONS

For $N \geq 2$, we look for radially symmetric solutions of

$$(42) \quad -\Delta u - k^2 u = f(|x|, u), \quad x \in \mathbb{R}^N,$$

where $k > 0$, and f satisfies the following assumptions:

(g_1) $f \in \mathcal{C}([0, \infty) \times \mathbb{R})$ with $f(\cdot, 0) = 0$, and f is locally Lipschitz continuous with respect to $u \in \mathbb{R}$.

(g_2) $0 \leq F(r, u) \leq \frac{1}{2} f(r, u) u$ for all $(r, u) \in [0, \infty) \times \mathbb{R}$, where $F(r, u) = \int_0^u f(r, s) ds$.

(g_3) For every $u \in \mathbb{R}$, $F(\cdot, u) \in \mathcal{C}^1(0, \infty)$ with $\partial_r F(r, u) \leq 0$ for all $r > 0$.

Setting $u(x) = u(r)$ with $r = |x|$, we can rewrite (42) as

$$(43) \quad -u'' - \frac{N-1}{r} u' - k^2 u = f(r, u), \quad r > 0$$

and we shall look for solutions of class \mathcal{C}^2 of this equation which satisfy $u'(0) = 0$.

The following local existence and uniqueness result (in a singular setting) is well known, see e.g. [29, Lemma 3.1] for a proof.

Lemma 5.1. *For every $\alpha \in \mathbb{R}$, there exists $\varepsilon > 0$ such that the initial value problem associated with (43), $u(0) = \alpha$ and $u'(0) = 0$ has a unique solution $u \in \mathcal{C}^2([0, \varepsilon), \mathbb{R})$.*

Using properties (g_2) and (g_3), we now continue the local solution given by Lemma 5.1 to a global solution on \mathbb{R}^+ and analyze its asymptotic behavior.

Theorem 5.2. *Let f satisfy (g_1) to (g_3). Then for every $\alpha \in \mathbb{R}$, there exists a unique radially symmetric solution of (42), $u = u(|x|)$, $u \in \mathcal{C}^2([0, \infty), \mathbb{R})$ satisfying $u(0) = \alpha$ and $u'(0) = 0$. Moreover,*

$$(44) \quad \sup_{r \geq 1} \{r^{N-1}(u'(r)^2 + u(r)^2)\} < \infty,$$

so that — identifying $u(x)$ with $u(|x|)$ for $x \in \mathbb{R}^N$ — we have $\frac{u}{\sqrt{1+|x|^2}}, \frac{|\nabla u|}{\sqrt{1+|x|^2}} \in L^2(\mathbb{R}^N)$.

If, in addition,

(g_4) $f(r, u) = o(|u|)$ as $u \rightarrow 0$, uniformly in $r > 0$,

then u satisfies the asymptotic condition (5).

Proof. Let $\alpha > 0$. By Lemma 5.1, there exists $\varepsilon > 0$ and a unique solution $u \in \mathcal{C}^2([0, 2\varepsilon), \mathbb{R})$ satisfying (43) and $u(0) = \alpha$, $u'(0) = 0$. Let $[0, r_0)$, $r_0 \in [2\varepsilon, \infty]$ denote the maximal existence interval of this solution. We claim that

$$(45) \quad \sup_{r \in [\varepsilon, r_0)} r^{N-1}(u'(r)^2 + u(r)^2) < \infty,$$

To see this, we use the change of variables

$$v(r) = r^{\frac{N-1}{2}} u(r), \quad r \in (0, r_0),$$

so that v solves

$$(46) \quad -v'' - \left\{ k^2 - \frac{(N-1)(N-3)}{4r^2} \right\} v = r^{\frac{N-1}{2}} f\left(r, r^{-\frac{N-1}{2}} v\right), \quad r \in (0, r_0).$$

We consider the \mathcal{C}^1 -function $r \mapsto \rho(r) := v'(r)^2 + k^2 v(r)^2$ on $(0, r_0)$ which satisfies

$$\frac{1}{2} \rho'(r) = \frac{(N-1)(N-3)}{4r^2} v(r) v'(r) - r^{\frac{N-1}{2}} f\left(r, r^{-\frac{N-1}{2}} v(r)\right) v'(r).$$

For $r \in [\varepsilon, r_0)$, we thus obtain

$$\begin{aligned}
 \rho(r) &= \rho(\varepsilon) + 2 \int_{\varepsilon}^r \frac{(N-1)(N-3)}{4s^2} v(s)v'(s) ds - 2 \int_{\varepsilon}^r s^{\frac{N-1}{2}} f(s, s^{-\frac{N-1}{2}} v(s)) v'(s) ds \\
 &= \rho(\varepsilon) + 2 \int_{\varepsilon}^r \frac{(N-1)(N-3)}{4s^2} v(s)v'(s) ds - 2s^{N-1} F(s, s^{-\frac{N-1}{2}} v(s)) \Big|_{\varepsilon}^r \\
 &\quad - (N-1) \int_{\varepsilon}^r s^{N-2} \{f(s, s^{-\frac{N-1}{2}} v(s)) s^{-\frac{N-1}{2}} v(s) - 2F(s, s^{-\frac{N-1}{2}} v(s))\} ds \\
 &\quad + 2 \int_{\varepsilon}^r s^{N-1} \partial_r F(s, s^{-\frac{N-1}{2}} v(s)) ds \\
 &\leq \rho(\varepsilon) + 2\varepsilon^{N-1} F(\varepsilon, \varepsilon^{\frac{1-N}{2}} v(\varepsilon)) + 2 \int_{\varepsilon}^r \frac{(N-1)(N-3)}{4s^2} v(s)v'(s) ds \\
 &\leq \rho(\varepsilon) + 2\varepsilon^{N-1} F(\varepsilon, \varepsilon^{\frac{1-N}{2}} v(\varepsilon)) + \int_{\varepsilon}^r \frac{(N-1)|N-3|}{4ks^2} \rho(s) ds,
 \end{aligned}$$

using (g_2) and (g_3) . Hence, Gronwall's inequality (see [12, Theorem III.1.1]) gives

$$\rho(r) \leq [\rho(\varepsilon) + 2\varepsilon^{N-1} F(\varepsilon, \varepsilon^{\frac{1-N}{2}} v(\varepsilon))] e^{\frac{(N-1)|N-3|}{4k\varepsilon^2}} \quad \text{for } r \in [\varepsilon, r_0).$$

From this we derive (45), since

$$r^{N-1} (u'(r)^2 + u(r)^2) \leq C_{\varepsilon} \rho(r) \quad \text{for } r \in [\varepsilon, r_0) \text{ with some constant } C_{\varepsilon} > 0.$$

As a consequence of (45), we find that $r_0 = \infty$ and that (44) holds.

We finally assume that (g_4) holds, and we check that the radiation condition in (5) is fulfilled.

For this we let $r \geq 1$ and write

$$\begin{aligned}
 r^{\frac{N-1}{2}} |u''(r) + k^2 u(r)| &= \left| \frac{(N-1)}{r} \left(\frac{(N-1)}{2r} v(r) - v'(r) \right) - \frac{f(r, r^{-\frac{(N-1)}{2}} v(r))}{r^{-\frac{(N-1)}{2}} v(r)} v(r) \right| \\
 &\leq \frac{(N-1)^2}{2r^2} \|v\|_{\infty} + \frac{(N-1)}{r} \|v'\|_{\infty} + \left| \frac{f(r, r^{-\frac{(N-1)}{2}} v(r))}{r^{-\frac{(N-1)}{2}} v(r)} \right| \|v\|_{\infty}.
 \end{aligned}$$

Since $\lim_{r \rightarrow \infty} r^{-\frac{(N-1)}{2}} v(r) = 0$, the assumption (g_4) gives $\lim_{r \rightarrow \infty} r^{\frac{N-1}{2}} |u''(r) + k^2 u(r)| = 0$, i.e., condition (5) holds. \square

We point out that if f satisfies $(g_1) - (g_4)$, every solution of (42) given by Theorem 5.2 is oscillatory. This follows from the fact that for every such solution u , 0 lies in the essential spectrum of any self-adjoint realization of the Sturm–Liouville differential expression

$$\tau w = -w'' - \left(k^2 - \frac{(N-1)(N-3)}{r^2} + \frac{f(r, u(r))}{u(r)} \right) w$$

associated to (46) (see [10, Corollary XIII.7.14 and Theorem XIII.7.40]).

6. PROPERTIES OF THE CAPACITY OPERATOR AND ITS REAL PART

In this section we derive key properties of the operator T_R introduced in Section 2 and its real part K_R in the case $N \geq 2$ (see Section 4 for the case $N = 1$). In particular, we will give the proofs of Lemma 2.1 and Lemma 2.3 (iv). Part of the material presented in this section should be well known to experts and appears in the standard literature in the special case $N = 3$ (see e.g. [19]). However we could not find the corresponding formulas for the general N -dimensional case. Moreover, as already remarked in the introduction, it seems that the

real part of the capacity operator has not been studied in the degree of detail which we need for our purposes.

Let $N \geq 2$, and consider the exterior Dirichlet problem for the Helmholtz equation

$$(47) \quad \begin{cases} \Delta w + k^2 w = 0 & \text{in } E_R := \mathbb{R}^N \setminus \overline{B_R}, \\ w = u & \text{on } S_R, \end{cases}$$

where $k > 0$, $R > 0$ and $u \in H^{\frac{1}{2}}(S_R, \mathbb{C})$ are given. Recall the space $H_R^{\mathbb{C}}$ defined in (9) which enforces a weak form of Sommerfeld's radiation condition. We wish to construct the unique solution $w \in H_R^{\mathbb{C}}$ explicitly in terms of a Fourier representation of the boundary datum u . For this we let Δ_S denote the Laplace–Beltrami operator on the unit sphere $S_1 \subset \mathbb{R}^N$, so that

$$(48) \quad \Delta w = \frac{1}{r^{N-1}} \frac{\partial}{\partial r} \left(r^{N-1} \frac{\partial w}{\partial r} \right) + \frac{1}{r^2} \Delta_S w.$$

We recall (see e.g. [24, Corollary 2.3]) that the linear span of the spherical harmonics (i.e. restrictions to S_1 of harmonic polynomials on \mathbb{R}^N with complex coefficients) is dense in $L^2(S_1, \mathbb{C})$. More precisely, denoting by \mathcal{H}_ℓ^N the space of spherical harmonics of degree $\ell \in \mathbb{N}_0$, then $d_\ell^N := \dim \mathcal{H}_\ell^N = \frac{(N+2\ell-2)}{(N+\ell-2)} \binom{N+\ell-2}{\ell}$ if $\ell \geq 1$, $d_0^N := \dim \mathcal{H}_0^N = 1$, and $\text{span} \bigcup_{\ell=0}^{\infty} \mathcal{H}_\ell^N$ is a dense subspace of $L^2(S_1, \mathbb{C})$. Furthermore, according to (48), every nonzero $\mathcal{Y}_\ell \in \mathcal{H}_\ell^N$ is an eigenfunction of Δ_S :

$$(49) \quad \Delta_S \mathcal{Y}_\ell + \ell(N + \ell - 2) \mathcal{Y}_\ell = 0 \quad \text{in } S_1$$

see [11, Theorem 3.2.11]. Moreover, starting from the orthogonal bases $\{1\}$ of \mathcal{H}_0^2 and $\{e^{i\ell\theta}, e^{-i\ell\theta}\}$ of \mathcal{H}_ℓ^2 , $\ell \geq 1$, ($\theta = \theta(x_1, x_2) = \text{sgn}(x_2) \arccos x_1$) we can inductively construct, according to [11, Lemma 3.5.3], orthogonal bases of \mathcal{H}_ℓ^N $\{\mathcal{Y}_1^\ell, \dots, \mathcal{Y}_{d_\ell^N}^\ell\}$, $\ell \in \mathbb{N}_0$ for all $N \geq 2$, with the property that for each element \mathcal{Y} in such a basis, the element $\overline{\mathcal{Y}}$ also belongs to this basis.

Proposition 6.1. *For $u \in H^{\frac{1}{2}}(S_R)$ given by the expansion $u(R\xi) = \sum_{\ell=0}^{\infty} \sum_{m=1}^{d_\ell^N} u_m^\ell \mathcal{Y}_m^\ell(\xi)$, $\xi \in S_1$, the problem (47) has a unique solution $w \in H_R^{\mathbb{C}}$ given by*

$$(50) \quad w(x) = \left(\frac{r}{R}\right)^{-\frac{N-2}{2}} \sum_{\ell=0}^{\infty} \frac{H_{\mu_\ell}^{(1)}(kr)}{H_{\mu_\ell}^{(1)}(kR)} \sum_{m=1}^{d_\ell^N} u_m^\ell \mathcal{Y}_m^\ell(\xi) \quad \text{for } r \geq R, \xi \in S_1 \text{ and } x = r\xi,$$

where, here and in the following, $\mu_\ell = \ell + \frac{(N-2)}{2}$ for $\ell \in \mathbb{N}_0$ and $H_\mu^{(1)}, H_\mu^{(2)}$ are the two Hankel functions (or Bessel functions of the third kind) of order μ (see e.g. [28, §3.6]).

Proof. We start by rescaling the space variable x to $\frac{x}{R}$. Setting $w(x) = v(\frac{x}{R})$, the problem (47) becomes

$$(51) \quad \begin{cases} \Delta v + (kR)^2 v = 0 & \text{in } E_1, \\ v = u_R & \text{on } S_1, \end{cases}$$

where $u_R(\xi) = u(R\xi)$, $\xi \in S_1$. We first remark that $w \in H_R^{\mathbb{C}}$ if and only if v belongs to the space

$$(52) \quad H_{1,kR}^{\mathbb{C}} := \left\{ u \in H_{loc}^1(E_1, \mathbb{C}) : \frac{u}{(1+r^2)^{1/2}} \in L^2(E_1, \mathbb{C}), \frac{\nabla u}{(1+r^2)^{1/2}} \in L^2(E_1, \mathbb{C}^N), \right. \\ \left. \frac{\partial u}{\partial r} - ikRu \in L^2(E_1, \mathbb{C}) \right\}.$$

We will therefore seek in this space, solutions of the form $v_m^\ell(x) = f_\ell(r)\mathcal{Y}_m^\ell(\xi)$ for some $\ell \in \mathbb{N}_0$, $1 \leq m \leq d_l^N$. Using (48) and (49), we may rewrite (51) for functions of this form to obtain

$$0 = \left\{ f_\ell''(r) + \frac{(N-1)}{r} f_\ell'(r) + \left((kR)^2 - \frac{\ell(N+\ell-2)}{r^2} \right) f_\ell(r) \right\} \mathcal{Y}_m^\ell(\xi).$$

Setting $g_\ell(s) = \left(\frac{s}{kR} \right)^{\frac{N-2}{2}} f_\ell\left(\frac{s}{kR} \right)$, we find

$$g_\ell''(s) + \frac{1}{s} g_\ell'(s) + \left\{ 1 - \frac{(\ell + \frac{N-2}{2})^2}{s^2} \right\} g_\ell(s) = 0, \quad s > 0,$$

which is Bessel's equation with parameter $\mu = \ell + \frac{N-2}{2}$. Its general solution is given by

$$g(s) = AH_\mu^{(1)}(s) + BH_\mu^{(2)}(s),$$

where $A, B \in \mathbb{C}$, and $H_\mu^{(1)}, H_\mu^{(2)}$ are the two Hankel functions of order μ . Now, we observe that the Sommerfeld radiation condition, given as the third condition in the definition of the space $H_{1,kR}^{\mathbb{C}}$ in (52), can be rewritten in the form

$$\int_{kR}^{\infty} s^{-1} \left| 2As[(H_\mu^{(1)})'(s) - iH_\mu^{(1)}(s)] - A(N-2)H_\mu^{(1)}(s) \right. \\ \left. + 2Bs[(H_\mu^{(2)})'(s) - iH_\mu^{(2)}(s)] - B(N-2)H_\mu^{(2)}(s) \right|^2 ds < \infty.$$

Using the recurrence formula $(H_\mu^{(p)})'(s) = \frac{\mu}{s} H_\mu^{(p)}(s) - H_{\mu+1}^{(p)}(s)$, $p = 1, 2$, we obtain

$$(53) \quad \int_{kR}^{\infty} s^{-1} \left| 2As[H_{\mu+1}^{(1)}(s) + iH_\mu^{(1)}(s)] + A(N-2-2\mu)H_\mu^{(1)}(s) \right. \\ \left. + 2Bs[H_{\mu+1}^{(2)}(s) + iH_\mu^{(2)}(s)] + B(N-2-2\mu)H_\mu^{(2)}(s) \right|^2 ds < \infty.$$

According to [28, Formulas 7.2 (1) and (2)], the asymptotic behavior of $H_\mu^{(p)}(s)$, $p = 1, 2$, is given by

$$H_\mu^{(1)}(s) = \sqrt{\frac{2}{\pi s}} e^{i(s - \frac{2\mu+1}{4}\pi)} [1 + O(s^{-1})], \quad H_\mu^{(2)}(s) = \sqrt{\frac{2}{\pi s}} e^{-i(s - \frac{2\mu+1}{4}\pi)} [1 + O(s^{-1})].$$

As a consequence,

$$|H_\mu^{(p)}(s)| = s^{-\frac{1}{2}} \left\{ \sqrt{\frac{2}{\pi}} + O(s^{-1}) \right\} \quad \text{for } p = 1, 2,$$

$$s|H_{\mu+1}^{(1)}(s) + iH_\mu^{(1)}(s)| = O(s^{-\frac{1}{2}}) \quad \text{and} \quad s|H_{\mu+1}^{(2)}(s) + iH_\mu^{(2)}(s)| = s^{\frac{1}{2}} \left[2\sqrt{\frac{2}{\pi}} + O(s^{-1}) \right].$$

Thus, (53) can only be satisfied, if $B = 0$. We conclude that a function in $H_{1,kR}^{\mathbb{C}}$ of the form $v_m^\ell(x) = f_\ell(r)Y_m^\ell(\xi)$ is a solution of the differential equation in (51) if and only if it can be written as

$$v_m^\ell(x) = Ar^{-\frac{(N-2)}{2}} H_{\mu_\ell}^{(1)}(kRr) \mathcal{Y}_m^\ell(\xi)$$

for some $A \in \mathbb{C}$ with $\mu_\ell = \ell + \frac{N-2}{2}$. Since $u_R(\xi) = \sum_{\ell=0}^{\infty} \sum_{m=1}^{d_\ell^N} u_m^\ell \mathcal{Y}_m^\ell(\xi)$, a solution of the boundary value problem (51) is thus given by

$$v(x) = r^{-\frac{(N-2)}{2}} \sum_{\ell=0}^{\infty} \frac{H_{\mu_\ell}^{(1)}(kRr)}{H_{\mu_\ell}^{(1)}(kR)} \sum_{m=1}^{d_\ell^N} u_m^\ell \mathcal{Y}_m^\ell(\xi) \quad \text{for } r \geq 1, \xi \in S_1 \text{ and } x = r\xi.$$

Rescaling back, we thus obtain the formula (50) for the (unique) solution $w \in H_R^{\mathbb{C}}$ of (47). \square

The formula (50) gives rise to the following expression for the capacity operator $T_R : H^{\frac{1}{2}}(S_R, \mathbb{C}) \rightarrow H^{-\frac{1}{2}}(S_R, \mathbb{C})$ (see [19] for the case $N = 3$):

$$(54) \quad [T_R u](R\xi) = \frac{1}{R} \sum_{\ell=0}^{\infty} z_\ell(kR) \sum_{m=1}^{d_\ell^N} u_m^\ell \mathcal{Y}_m^\ell(\xi), \quad \xi \in S_1$$

for $u \in H^{\frac{1}{2}}(S_R, \mathbb{C})$ given by $u(R\xi) = \sum_{\ell=0}^{\infty} \sum_{m=1}^{d_\ell^N} u_m^\ell \mathcal{Y}_m^\ell(\xi)$, $\xi \in S_1$, where

$$z_\ell(r) = \frac{r \frac{d}{dr} H_{\mu_\ell}^{(1)}(r)}{H_{\mu_\ell}^{(1)}(r)} - \frac{N-2}{2}, \quad r > 0.$$

We need some estimates for the coefficients $z_\ell(r)$. For this we recall that

$$H_\mu^{(1)} = J_\mu + iY_\mu, \quad \text{and} \quad H_\mu^{(2)} = J_\mu - iY_\mu = \overline{H_\mu^{(1)}} \quad \text{for } \mu \in \mathbb{R},$$

where J_μ and Y_μ denote the Bessel functions of the first and second kind. Setting $G_\mu(r) = \frac{r \frac{d}{dr} H_\mu^{(1)}(r)}{H_\mu^{(1)}(r)}$, for $r > 0$, $\mu \in \mathbb{R}$, we see that

$$(55) \quad \operatorname{Re} G_\mu(r) = \frac{r \operatorname{Re}(\overline{H_\mu^{(1)}(r)} \frac{d}{dr} H_\mu^{(1)}(r))}{|H_\mu^{(1)}(r)|^2} = \frac{r \frac{d}{dr} |H_\mu^{(1)}(r)|^2}{2 |H_\mu^{(1)}(r)|^2} = \frac{r \frac{d}{dr} (J_\mu^2(r) + Y_\mu^2(r))}{2 (J_\mu^2(r) + Y_\mu^2(r))}$$

$$(56) \quad \operatorname{Im} G_\mu(r) = \frac{r \operatorname{Im}(\overline{H_\mu^{(1)}(r)} \frac{d}{dr} H_\mu^{(1)}(r))}{|H_\mu^{(1)}(r)|^2} = \frac{r (J_\mu(r) Y_\mu'(r) - J_\mu'(r) Y_\mu(r))}{|H_\mu^{(1)}(r)|^2}.$$

We need the following estimates (see [19] for the case $N = 3$):

Lemma 6.2. *For $r > 0$ and $\ell \geq \max\{0, \frac{3-N}{2}\}$ we have*

$$(57) \quad \frac{(N-1)}{2} \leq -\operatorname{Re} z_\ell(r) \leq \ell + N - 2$$

$$(58) \quad 0 < \operatorname{Im} z_\ell(r) \leq r.$$

In the case $N = 2$, we also have $0 < -\operatorname{Re} z_0(r) \leq \frac{1}{2}$ and $\operatorname{Im} z_0(r) > 0$ for all $r > 0$.

Proof. We first prove (57). According to [28, §13.74], we have

$$(i) \quad \frac{d}{dr}(J_\mu^2(r) + Y_\mu^2(r)) < 0 \quad \text{for } r > 0, \mu \in \mathbb{R},$$

$$(ii) \quad \frac{d}{dr} r(J_\mu^2(r) + Y_\mu^2(r)) \begin{cases} \leq 0 & \text{if } \mu \geq \frac{1}{2} \\ \geq 0 & \text{if } \mu \leq \frac{1}{2}, \end{cases} \quad \text{for } r > 0.$$

From (i) we obtain $\operatorname{Re} G_\mu(r) < 0$ for all $r > 0$, $\mu \in \mathbb{R}$, and (ii) gives $\operatorname{Re} G_\mu(r) \leq -\frac{1}{2}$ for $r > 0$, $\mu \geq \frac{1}{2}$. Moreover, (ii) implies $\operatorname{Re} G_{\frac{1}{2}}(r) = -\frac{1}{2}$ for $r > 0$. Now, using the recurrence formula $\frac{d}{dr} H_\mu^{(1)}(r) = H_{\mu-1}^{(1)}(r) - \frac{\mu}{r} H_\mu^{(1)}(r)$, we find that

$$\begin{aligned} |H_\mu^{(1)}(r)|^2 (\operatorname{Re} G_\mu(r) + \mu) &= \operatorname{Re} \left(r \overline{H_\mu^{(1)}(r)} \frac{d}{dr} H_\mu^{(1)}(r) + \mu |H_\mu^{(1)}(r)|^2 \right) \\ &= r \operatorname{Re} \left(\overline{H_\mu^{(1)}(r)} H_{\mu-1}^{(1)}(r) \right) = r \operatorname{Re} \left(\overline{H_{\mu-1}^{(1)}(r)} H_\mu^{(1)}(r) \right) \\ &= -|H_{\mu-1}^{(1)}(r)|^2 (\operatorname{Re} G_{\mu-1}(r) - (\mu - 1)). \end{aligned}$$

Since $\operatorname{Re} G_\mu(r) < 0$ holds for all μ , the previous formula gives $\operatorname{Re} G_\mu(r) \geq -\mu$ for all $\mu \geq 1$. Summarizing, we have shown

$$-\mu \leq \operatorname{Re} G_\mu(r) \leq -\frac{1}{2}, \quad r > 0 \quad \text{for } \mu \in \left\{ \frac{1}{2} \right\} \cup [1, \infty).$$

This shows (57), since $z_\ell(r) = G_{\ell + \frac{N-2}{2}}(r) - \frac{N-2}{2}$. In the case $\mu = 0$, we have $-\frac{1}{2} \leq \operatorname{Re} G_0(r) < 0$ for all $r > 0$. Hence, when $N = 2$, we obtain $-\frac{1}{2} \leq z_0(r) < 0$ for all $r > 0$.

We now turn to the proof of (58). Using (56) and the fact that

$$(59) \quad \mathcal{W}(J_\mu, Y_\mu; r) := J_\mu(r) Y_\mu'(r) - J_\mu'(r) Y_\mu(r) = \frac{2}{\pi r} \quad \text{for } \mu \in \mathbb{R}, r > 0,$$

(see [28, §3.63 (1)]), we find that

$$\operatorname{Im} G_\mu(r) = \frac{r \mathcal{W}(J_\mu, Y_\mu; r)}{|H_\mu^{(1)}(r)|^2} = \frac{2}{\pi |H_\mu^{(1)}(r)|^2} > 0 \quad \text{for } \mu \in \mathbb{R}, r > 0.$$

Furthermore, using (ii) above, we see that

$$r |H_\mu^{(1)}(r)|^2 \geq \lim_{t \rightarrow \infty} t |H_\mu^{(1)}(t)|^2 = \frac{2}{\pi} \quad \text{for } r > 0, \mu \geq \frac{1}{2}.$$

We thus obtain $\operatorname{Im} G_\mu(r) \leq r$ for $r > 0$, $\mu \geq \frac{1}{2}$. As before, (58) follows from the identity $z_\ell(r) = G_{\ell + \frac{N-2}{2}}(r) - \frac{N-2}{2}$. \square

Recall that in Section 2 we have also introduced the operator

$$(60) \quad K_R : H^{\frac{1}{2}}(S_R) \rightarrow H^{-\frac{1}{2}}(S_R), \quad K_R u = \operatorname{Re}[T_R u],$$

where $H^{\frac{1}{2}}(S_R) = H^{\frac{1}{2}}(S_R, \mathbb{R})$, $H^{-\frac{1}{2}}(S_R) = H^{-\frac{1}{2}}(S_R, \mathbb{R})$. For $u \in H^{\frac{1}{2}}(S_R)$ given by $u(R\xi) =$

$\sum_{\ell=0}^{\infty} \sum_{m=1}^{d_\ell^N} u_m^\ell \mathcal{Y}_m^\ell(\xi)$ we then have

$$(61) \quad K_R u = \frac{1}{R} \sum_{\ell=0}^{\infty} \operatorname{Re}(z_\ell(kR)) \sum_{m=1}^{d_\ell^N} u_m^\ell \mathcal{Y}_m^\ell,$$

since $\sum_{m=1}^{d_\ell^N} u_m^\ell \mathcal{Y}_m^\ell$ is a real-valued function for every $\ell \in \mathbb{N}_0$. Using Lemma 6.2 we may therefore easily complete the

Proof of Lemma 2.1. We first note that, for $s \in \mathbb{R}$, one may define an equivalent norm on $H^s(S_R, \mathbb{C})$ by

$$\|u\|_s^2 = \sum_{\ell=0}^{\infty} (1 + \ell^2)^s \sum_{m=1}^{d_\ell^N} |u_m^\ell|^2$$

for $u \in H^{\frac{1}{2}}(S_R, \mathbb{C})$ given by $u(R\xi) = \sum_{\ell=0}^{\infty} \sum_{m=1}^{d_\ell^N} u_m^\ell \mathcal{Y}_m^\ell(\xi)$. Hence, by (57) and (58), the capacity operator $T_R : H^{\frac{1}{2}}(S_R, \mathbb{C}) \rightarrow H^{-\frac{1}{2}}(S_R, \mathbb{C})$ is bounded, and thus K_R defined in (60) is bounded as well. In particular, the integrals on both sides of (14) exist, and by (61) and the orthogonality of spherical harmonics (corresponding to different values of ℓ) they coincide. Finally, the second inequality in (13) also follows, by orthogonality, from (61) and the estimates in Lemma 6.2. \square

Next, we study the eigenspace X^0 of the eigenvalue problem (17) corresponding to the eigenvalue k^2 , and we complete the

Proof of Lemma 2.3(iv). Recall that $u \in X^0$ if and only if u solves

$$(62) \quad \begin{cases} \Delta u + k^2 u = 0 & \text{in } B_R, \\ \frac{\partial u}{\partial \eta} = v & \text{on } S_R \end{cases}$$

with $v = K_R u$ on S_R . Put, as before, $\mu_\ell = \ell + \frac{N-2}{2}$ for $\ell \in \mathbb{N}_0$. In the case where $\frac{d}{dr} \left(r^{\frac{2-N}{2}} J_{\mu_\ell}(kr) \right) \Big|_{r=R} \neq 0$ for all $\ell \in \mathbb{N}_0$, the inhomogeneous Neumann problem (62) has, for given $v \in H^{-\frac{1}{2}}(S_R)$, the unique solution

$$u(r\xi) = \left(\frac{r}{R} \right)^{-\frac{N-2}{2}} \sum_{\ell=0}^{\infty} \frac{J_{\mu_\ell}(kr)}{kJ'_{\mu_\ell}(kR) - \frac{(N-2)}{2R} J_{\mu_\ell}(kR)} \sum_{m=1}^{d_\ell^N} v_m^\ell \mathcal{Y}_m^\ell(\xi), \quad 0 < r < R,$$

where the coefficients v_m^ℓ are determined by $v(R\xi) = \sum_{\ell=0}^{\infty} \sum_{m=1}^{d_\ell^N} v_m^\ell \mathcal{Y}_m^\ell(\xi)$. In particular, the restriction of u to S_R satisfies

$$u(R\xi) = \sum_{\ell=0}^{\infty} \sum_{m=1}^{d_\ell^N} u_m^\ell \mathcal{Y}_m^\ell(\xi) \quad \text{with } u_m^\ell = \frac{J_{\mu_\ell}(kR)}{kJ'_{\mu_\ell}(kR) - \frac{(N-2)}{2R} J_{\mu_\ell}(kR)} v_m^\ell.$$

If, in addition, $v = K_R u$ on S_R , then by (61) we have

$$\begin{aligned} v_m^\ell &= \frac{1}{R} \operatorname{Re} \left(\frac{kR \frac{d}{dr} H_{\mu_\ell}^{(1)}(kR)}{H_{\mu_\ell}^{(1)}(kR)} - \frac{N-2}{2} \right) u_m^\ell \\ &= k \left(\frac{J'_{\mu_\ell}(kR) J_{\mu_\ell}(kR) + Y'_{\mu_\ell}(kR) Y_{\mu_\ell}(kR)}{J_{\mu_\ell}^2(kR) + Y_{\mu_\ell}^2(kR)} - \frac{N-2}{2kR} \right) u_m^\ell \end{aligned}$$

and therefore

$$u_m^\ell = \frac{J_{\mu_\ell}(kR)}{J'_{\mu_\ell}(kR) - \frac{(N-2)}{2kR}J_{\mu_\ell}(kR)} \left(\frac{J'_{\mu_\ell}(kR)J_{\mu_\ell}(kR) + Y'_{\mu_\ell}(kR)Y_{\mu_\ell}(kR)}{J_{\mu_\ell}^2(kR) + Y_{\mu_\ell}^2(kR)} - \frac{N-2}{2kR} \right) u_m^\ell$$

for $\ell \in \mathbb{N}_0$, $1 \leq m \leq d_\ell^N$. As a consequence of (59), this gives for each $\ell \in \mathbb{N}_0$ the alternative

$$u_m^\ell = 0 \quad \text{for all } 1 \leq m \leq d_\ell^N \quad \text{or} \quad Y_{\mu_\ell}(kR) = 0.$$

We may now finish the proof by setting

$$\mathcal{D} := \left\{ R > 0 : \frac{d}{dr} \left(r^{\frac{2-N}{2}} J_{\mu_\ell}(kr) \right) \Big|_{r=R} = 0 \text{ or } Y_{\mu_\ell}(kR) = 0 \text{ for some } \ell \in \mathbb{N}_0 \right\}$$

Indeed, since J_{μ_ℓ} and Y_{μ_ℓ} are analytic functions on $(0, \infty)$ for every $\ell \in \mathbb{N}_0$, the set \mathcal{D} is countable. Moreover, for $R \in (0, \infty) \setminus \mathcal{D}$ and $u \in X^0$ we conclude $u_m^\ell = 0$ for all $\ell \in \mathbb{N}_0$, $1 \leq m \leq d_\ell^N$ and therefore $u \equiv 0$. \square

We close this section with the

Proof of Theorem 1.2. Let $R > 0$ be fixed, and let $u \in \mathcal{R}_R$ with $u|_{S_R} \neq 0$. Furthermore, let $R' > R > 0$ be given such that $u \in \mathcal{R}_{R'}$. Then we have:

- (i) In E_R , u coincides with the real part of a solution w_1 of the linear Helmholtz equation satisfying (7);
- (ii) In $E_{R'}$, u coincides with the real part of a solution w_2 of the linear Helmholtz equation satisfying (7);
- (iii) $u|_{S_R} \equiv w_1|_{S_R}$ and $u|_{S_{R'}} \equiv w_2|_{S_{R'}}$.

Since $w := w_1 - w_2$ also satisfies the linear Helmholtz equation in $E_{R'}$ together with (7) and $\text{Re}(w) \equiv 0$ in $E_{R'}$, it follows that $w(x) = o(r^{\frac{1-N}{2}})$ as $r = |x| \rightarrow \infty$ and hence $w \equiv 0$ in $E_{R'}$ by a classical result of Rellich [21, Satz 1]. Hence $w_1 = w_2$ in $E_{R'}$, and by (iii) this implies that $\text{Im}(w_1|_{S_{R'}}) \equiv \text{Im}(w_2|_{S_{R'}}) \equiv 0$. Writing again

$$(63) \quad u(R\xi) = w_1(R\xi) = \sum_{\ell=0}^{\infty} u^\ell(\xi), \quad \xi \in S_1,$$

with $u^\ell(\xi) = \sum_{m=1}^{d_\ell^N} u_m^\ell \mathcal{Y}_m^\ell(\xi)$, we have, by Proposition 6.1,

$$w_1(R'\xi) = \left(\frac{R'}{R} \right)^{-\frac{(N-2)}{2}} \sum_{\ell=0}^{\infty} \frac{H_{\mu_\ell}^{(1)}(kR')}{H_{\mu_\ell}^{(1)}(kR)} u^\ell(\xi), \quad \xi \in S_1,$$

where the functions $\xi \mapsto u^\ell(\xi)$, $\ell \in \mathbb{N}_0$ are real-valued as a consequence of (63) and the fact that u is real-valued. Moreover, the assumption $u|_{S_R} \neq 0$ forces that $u^\ell \neq 0$ on S_1 for at least one $\ell \in \mathbb{N}_0$, and therefore

$$\text{Im} \left(\frac{H_{\mu_\ell}^{(1)}(kR')}{H_{\mu_\ell}^{(1)}(kR)} \right) = 0, \quad \text{i.e.,} \quad J_{\mu_\ell}(kR')Y_{\mu_\ell}(kR) = J_{\mu_\ell}(kR)Y_{\mu_\ell}(kR').$$

Since the functions J_{μ_ℓ} and Y_{μ_ℓ} are real analytic, the latter can only happen for countably many $R' > R$. We thus conclude that $u \in \mathcal{R}_{R'}$ for at most countably many $R' > R$.

It remains to be shown that problem (4), (5) admits infinitely many solutions if the nonlinearity f satisfies (f0)–(f4). For this we fix $R_1 \in (0, \infty) \setminus \mathcal{D}$ such that $\Omega \subset B_{R_1}$; then Theorem 1.1 yields a nontrivial solution $u_1 \in \mathcal{R}_{R_1}$ of (4), (5). The nontriviality of u implies

that $u_1|_{S_{R_1}} \neq 0$, since otherwise $u \equiv 0$ on E_{R_1} and therefore on all of \mathbb{R}^N as a consequence of unique continuation. Hence there exists $R_2 \in (R_1, \infty) \setminus \mathcal{D}$ such that $u_1 \notin \mathcal{R}_{R_2}$, whereas Theorem 1.1 yields a solution $u_2 \in \mathcal{R}_{R_2}$ of (4), (5). Moreover, there exists $R_3 \in (R_2, \infty) \setminus \mathcal{D}$ such that $u_1, u_2 \notin \mathcal{R}_{R_3}$, whereas Theorem 1.1 yields a solution $u_3 \in \mathcal{R}_{R_3}$ of (4), (5). Inductively, we now obtain an infinite sequence of pairwise different solutions of (4), (5). \square

REFERENCES

- [1] Alves, C.O., Souto, M.A.S., Montenegro, M.: Existence of solution for two classes of elliptic problems in \mathbb{R}^N with zero mass. *J. Differential Equations* **252**, 5735–5750 (2012)
- [2] Ambrosetti, A., Felli, V., Malchiodi, A.: Ground states of nonlinear Schrödinger equations with potentials vanishing at infinity. *J. Eur. Math. Soc. (JEMS)* **7**, 117–144 (2005)
- [3] Ambrosetti, A., Malchiodi, A.: Perturbation Methods and Semilinear Elliptic Problems on \mathbb{R}^n , *Progress in Mathematics*, vol. 240. Birkhäuser, Basel (2006)
- [4] Bartolo, P., Benci, V., Fortunato, D.: Abstract critical point theorems and applications to some nonlinear problems with “strong” resonance at infinity. *Nonlinear Anal.* **7**, 981–1012 (1983)
- [5] Bartsch, T.: Infinitely many solutions of a symmetric Dirichlet problem. *Nonlinear Anal.* **20**, 1205–1216 (1993)
- [6] Bartsch, T., Ding, Y.: On a nonlinear Schrödinger equation with periodic potential. *Math. Ann.* **313**, 15–37 (1999)
- [7] Bartsch, T., Willem, M.: Infinitely many nonradial solutions of a Euclidean scalar field equation. *J. Funct. Anal.* **177**, 447–460 (1993)
- [8] Cerami, G.: Some nonlinear elliptic problems in unbounded domains. *Milan J. Math.* **74**, 47–77 (2006)
- [9] Costa, D.G., Tehrani, H.: Existence of positive solutions for a class of indefinite elliptic problems in \mathbb{R}^N . *Calc. Var. Partial Differential Equations* **13**, 159–189 (2001)
- [10] Dunford, N., Schwartz, J.T.: *Linear Operators. Part II: Spectral Theory.* John Wiley & Sons, New York (1963)
- [11] Groemer, H.: Geometric Applications of Fourier Series and Spherical Harmonics, *Encyclopedia of Mathematics and its Applications*, vol. 61. Cambridge University Press, Cambridge (1996)
- [12] Hartman, P.: Ordinary Differential Equations, *Classics in Applied Mathematics*, vol. 38, second edn. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA (2002)
- [13] Jörgens, K.: Das Anfangswertproblem im Großen für eine Klasse nichtlinearer Wellengleichungen. *Math. Z.* **77**, 295–308 (1961)
- [14] Kato, T.: Growth properties of solutions of the reduced wave equation with a variable coefficient. *Comm. Pure Appl. Math.* **12**, 403–425 (1959)
- [15] Kuzin, I., Pohozaev, S.: Entire Solutions of Semilinear Elliptic Equations, *Progress in Nonlinear Differential Equations and their Applications*, vol. 33. Birkhäuser, Basel (1997)
- [16] Li, G., Wang, C.: The existence of a nontrivial solution to a nonlinear elliptic problem of linking type without the Ambrosetti-Rabinowitz condition. *Ann. Acad. Sc. Fenn. Math* **36**, 461–480 (2011)
- [17] Liu, S.: On superlinear Schrödinger equations with periodic potential. *Calc. Var. Partial Differential Equations* **45**, 1–9 (2012)
- [18] Moroz, V., Van Schaftingen, J.: Semiclassical stationary states for nonlinear Schrödinger equations with fast decaying potentials. *Calc. Var. Partial Differential Equations* **37**, 1–27 (2010)
- [19] Nédélec, J.C.: Acoustic and Electromagnetic Equations. Integral Representations for Harmonic Problems, *Applied Mathematical Sciences*, vol. 144. Springer-Verlag, New-York (2001)
- [20] Rabinowitz, P.H.: A note on a semilinear elliptic equation on \mathbb{R}^n , pp. 307–317. *Quaderni. Scuola Norm. Sup. Pisa* (1991)
- [21] Rellich, F.: Über das asymptotische Verhalten der Lösungen von $\Delta u + \lambda u = 0$ in unendlichen Gebieten. *Jber. Deutsch. Math. Verein.* **53**, 57–65 (1943)
- [22] Schneider, M.: Existence and nonexistence of positive solutions of indefinite elliptic problems in \mathbb{R}^N . *Adv. Nonlinear Stud.* **3**, 231–259 (2003)
- [23] Sommerfeld, A.: Die Greensche Funktion der Schwingungsgleichung. *Jber. Deutsch. Math. Verein.* **21**, 309–353 (1912)
- [24] Stein, E.M., Weiss, G.: Introduction to Fourier Analysis on Euclidean Spaces, *Princeton Mathematical Series*, vol. 32. Princeton University Press, Princeton NJ (1971)

- [25] Struwe, M.: Variational methods: Applications to Nonlinear Partial Differential Equations and Hamiltonian Systems, fourth edn. Springer-Verlag, Berlin (2008)
- [26] Stuart, C.A.: Bifurcation from the essential spectrum. In: Topological Nonlinear Analysis, II (Frascati, 1995), *Progress in Nonlinear Differential Equations and their Applications*, vol. 27, pp. 397–443. Birkhäuser, Boston (1997)
- [27] Szulkin, A., Weth, T.: Ground-state solutions for some indefinite variational problems. *J. Funct. Anal.* **257**, 3802–3822 (2009)
- [28] Watson, G.N.: A Treatise on the Theory of Bessel Functions. Reprint of the second (1944) edition. Cambridge Mathematical Library. Cambridge University Press, Cambridge (1996)
- [29] Weth, T.: Global bifurcation branches for radially symmetric Schrödinger equations. *Adv. Differential Equations* **10**, 721–746 (2005)
- [30] Willem, M.: Minimax Theorems, *Progress in Nonlinear Differential Equations and Their Application*, vol. 24. Birkhäuser, Basel (1996)
- [31] Willem, M., Zou, W.: On a Schrödinger equation with periodic potential and spectrum point zero. *Indiana Univ. Math. J.* **52**, 109–132 (2003)

INSTITUT FÜR MATHEMATIK - GOETHE-UNIVERSITÄT FRANKFURT, ROBERT-MAYER-STR. 10. - 60054
 FRANKFURT (GERMANY)
E-mail address: evequoz@math.uni-frankfurt.de

INSTITUT FÜR MATHEMATIK - GOETHE-UNIVERSITÄT FRANKFURT, ROBERT-MAYER-STR. 10. - 60054
 FRANKFURT (GERMANY)
E-mail address: weth@math.uni-frankfurt.de