

# Fractional Classical Mechanics

Nick Laskin\*

TopQuark Inc.  
Toronto, ON, M6P 2P2

## Abstract

Fractional classical mechanics has been introduced and developed as a classical counterpart of the fractional quantum mechanics. Lagrange, Hamilton and Hamilton-Jacobi frameworks have been implemented for the fractional classical mechanics. The Lagrangian of fractional classical mechanics has been introduced, and equation of motion has been obtained. Fractional oscillator model has been launched and solved in 1D case. A new equation for the period of oscillations of fractional classical oscillator has been found. The interplay between the energy dependency of the period of classical oscillations and the non-equidistant distribution of the energy levels for fractional quantum oscillator has been discussed.

We discuss as well, the relationships between new equations of fractional classical mechanics and the well-known fundamental equations of classical mechanics.

*PACS* numbers: 45.50.Dd; 45.05.+x; 45.20.Jj.

*Keywords:* Fractional classical mechanics, Lagrange, Hamilton and Hamilton-Jacobi approaches, Fractional oscillator model, fractional Kepler's third law.

## 1 Introduction

We introduce and elaborate fractional classical mechanics as a classical counterpart of the fractional quantum mechanics [1]-[4]. To begin with, let us

---

\*E-mail: nlaskin@rocketmail.com

consider the equation for canonical classical mechanics action  $S_\alpha(p, q)$  (see, Eq.(23) in [2]),

$$S_\alpha(p, q) = \int_{t_a}^{t_b} d\tau (p(\tau)\dot{q}(\tau) - H_\alpha(p(\tau), q(\tau))), \quad (1)$$

where the classical Hamiltonian  $H_\alpha(p(\tau), q(\tau))$  arrives from the Hamiltonian  $H_\alpha(p, q)$  [1]-[4]

$$H_\alpha(p, q) = D_\alpha |p|^\alpha + V(q), \quad 1 < \alpha \leq 2, \quad (2)$$

with  $V(q)$  as a potential energy and  $p \rightarrow p(\tau)$ ,  $q \rightarrow q(\tau)$ , and  $\{p(\tau), q(\tau)\}$  is the particle trajectory in phase space. The scaling factor  $D_\alpha$  has physical dimension  $[D_\alpha] = \text{erg}^{1-\alpha} \cdot \text{cm}^\alpha \cdot \text{sec}^{-\alpha}$ , see, [1]-[2]. Since the Hamiltonian  $H_\alpha(p, q)$  does not explicitly depend on the time, it represents a conserved quantity which is in fact the total energy of fractional classical mechanics system.

Here we consider 1D fractional classical mechanics. The 3D generalization is straightforward and it is based on the Hamiltonian [4]

$$H_\alpha(\mathbf{p}, \mathbf{q}) = D_\alpha |\mathbf{p}|^\alpha + V(\mathbf{q}), \quad 1 < \alpha \leq 2, \quad (3)$$

where  $\mathbf{p}$  and  $\mathbf{q}$  are 3D vectors.

The paper is organized as follows.

Fundamentals of fractional classical mechanics are covered in Sec.2, where Lagrange, Hamilton and Hamilton-Jacobi frameworks have been implemented.

In Sec.3 the scaling analysis of fractional classical motion equations has been developed based on the mechanical similarity. We discover and discuss *fractional Kepler's third law* which is a generalization of the well-known Kepler's third law.

The motion equations for the fractional classical mechanics have been integrated in Sec.4. Fractional classical oscillator model has been introduced. We found an equation for the period of oscillations of fractional classical oscillator. The map between the energy dependence of the period of classical oscillations and the non-equidistant distribution of quantum energy levels has been established.

In conclusion we outline the new developments.

## 2 Fundamentals of fractional classical mechanics

### 2.1 The Lagrange approach

The Lagrangian of fractional classical mechanics  $L_\alpha(\dot{q}, q)$  is defined as usual

$$L_\alpha(\dot{q}, q) = p\dot{q} - H_\alpha(p, q), \quad (4)$$

where the momentum  $p$  is

$$p = \frac{\partial L_\alpha(\dot{q}, q)}{\partial \dot{q}}, \quad (5)$$

and  $H_\alpha(p, q)$  is given by Eq.(2).

Hence, we obtain the Lagrangian of the fractional classical mechanics

$$L_\alpha(\dot{q}, q) = \left( \frac{1}{\alpha D_\alpha} \right)^{\frac{1}{\alpha-1}} \frac{\alpha-1}{\alpha} |\dot{q}|^{\frac{\alpha}{\alpha-1}} - V(q), \quad 1 < \alpha \leq 2. \quad (6)$$

For a free particle,  $V(q) = 0$ , the Lagrangian of the fractional classical mechanics is

$$L_\alpha^{(0)}(\dot{q}, q) = \left( \frac{1}{\alpha D_\alpha} \right)^{\frac{1}{\alpha-1}} \frac{\alpha-1}{\alpha} |\dot{q}|^{\frac{\alpha}{\alpha-1}}, \quad 1 < \alpha \leq 2. \quad (7)$$

Further, the Euler-Lagrange equation of motion has the standard form

$$\frac{d}{dt} \frac{\partial L_\alpha(\dot{q}, q)}{\partial \dot{q}} - \frac{\partial L_\alpha(\dot{q}, q)}{\partial q} = 0. \quad (8)$$

By substituting Eq.(6) into Eq.(8) we find the motion equation in Lagrangian form

$$\left( \frac{1}{\alpha D_\alpha} \right)^{\frac{1}{\alpha-1}} \frac{1}{\alpha-1} \ddot{q} |\dot{q}|^{\frac{2-\alpha}{\alpha-1}} + \frac{\partial V(q)}{\partial q} = 0. \quad (9)$$

This equation has to be accompanied by the initial conditions. We impose the following initial conditions. At  $t = 0$ , the initial displacement is denoted by  $q_0$  and the corresponding velocity is denoted by  $\dot{q}_0$ , that is,

$$q(t=0) = q_0, \quad \text{and} \quad \dot{q}(t=0) = \dot{q}_0. \quad (10)$$

As one can see, Eq.(9) has a non-linear kinematic term.

The new equation (9) is fractional generalization of the well known equation of motion of classical mechanics in the Lagrange form.

In the special case when  $\alpha = 2$ , Eq.(9) goes into

$$m\ddot{q} + \frac{\partial V(q)}{\partial q} = 0. \quad (11)$$

where  $m$  is a particle mass ( $D_2 = 1/2m$ ) and the initial conditions are given by Eq.(10).

## 2.2 The Hamilton approach

To obtain the Hamilton equations of motion for the fractional classical mechanics we apply variational principle,

$$\delta S_\alpha(p, q) = 0, \quad (12)$$

where the action  $S_\alpha(p, q)$  is given by Eq.(1).

Considering the momentum  $p$  and coordinate  $q$  as independent variables we can write

$$\delta S_\alpha(p, q) = \int_{t_a}^{t_b} d\tau (\delta p \dot{q} + p \delta \dot{q} - \frac{\partial H_\alpha(p, q)}{\partial p} \delta p - \frac{\partial H_\alpha(p, q)}{\partial q} \delta q) = 0. \quad (13)$$

Upon integration by parts of the second term  $p \delta \dot{q}$ , the variation  $\delta S_\alpha$  becomes

$$\delta S_\alpha(p, q) = \int_{t_a}^{t_b} d\tau \delta p (\dot{q} - \frac{\partial H_\alpha(p, q)}{\partial p}) + p \delta q|_{t_a}^{t_b} - \int_{t_a}^{t_b} d\tau \delta q (\dot{p} + \frac{\partial H_\alpha(p, q)}{\partial q}) = 0.$$

Since  $\delta q(t_a) = \delta q(t_b) = 0$  at the end points of the trajectory, the term  $p \delta q$  is 0. Between the end points  $\delta p$  and  $\delta q$  can take on any arbitrary value. Hence, the variation  $\delta S_\alpha$  can be 0 if the following conditions are satisfied

$$\dot{q} = \frac{\partial H_\alpha(p, q)}{\partial p}, \quad \dot{p} = -\frac{\partial H_\alpha(p, q)}{\partial q}. \quad (14)$$

This is, of course, the canonical Hamilton equations of motion. Thus, for the time-independent Hamiltonian given by Eq.(2) we obtain the equations

$$\dot{q} = \alpha D_\alpha |p|^{\alpha-1} \text{sgn} p, \quad \dot{p} = -\frac{V(q)}{\partial q}, \quad 1 < \alpha \leq 2, \quad (15)$$

where  $\text{sgn} p$  is the sign function which for nonzero values of  $p$  can be defined by the formula

$$\text{sgn} p = \frac{p}{|p|},$$

where  $|p|$  is the absolute value of  $p$ .

Equations (15) are the Hamilton equations for fractional classical mechanics system.

### 2.2.1 Poisson bracket approach to fractional classical mechanics

It is well-known that Hamiltonian classical mechanics could be reformulated in terms of the Poisson brackets. For the two arbitrary functions  $u(p, q)$  and  $v(p, q)$  of variables  $p$  and  $q$ , the Poisson bracket is defined as [5]

$$\{u(p, q), v(p, q)\} = \frac{\partial u}{\partial p} \frac{\partial v}{\partial q} - \frac{\partial u}{\partial q} \frac{\partial v}{\partial p}. \quad (16)$$

The Hamilton's equations of motion have an equivalent expression in terms of the Poisson bracket. Indeed, suppose that  $f(p, q, t)$  is a function of momentum  $p$ , coordinate  $q$  and time  $t$ . Then we have

$$\frac{df}{dt} = \frac{\partial f}{\partial t} + \left( \frac{\partial f}{\partial p} \dot{p} + \frac{\partial f}{\partial q} \dot{q} \right). \quad (17)$$

Substituting  $\dot{q}$  and  $\dot{p}$  given by Eq.(14) yields

$$\frac{df}{dt} = \frac{\partial f}{\partial t} + \{H_\alpha, f\}, \quad (18)$$

where  $\{H_\alpha, f\}$  is the Poisson bracket,

$$\{H_\alpha, f\} = \frac{\partial H_\alpha}{\partial p} \frac{\partial f}{\partial q} - \frac{\partial H_\alpha}{\partial q} \frac{\partial f}{\partial p}. \quad (19)$$

with  $H_\alpha$  given by Eq.(2).

### 2.3 The Hamilton-Jacobi approach

Having the Lagrangian (6), we can present classical mechanics action given by Eq.(1) as a function of coordinate  $q$ ,

$$S_\alpha(q) = \int_{t_a}^{t_b} d\tau L_\alpha(\dot{q}, q) = \int_{t_a}^{t_b} d\tau \left\{ \left( \frac{1}{\alpha D_\alpha} \right)^{\frac{1}{\alpha-1}} \frac{\alpha-1}{\alpha} |\dot{q}|^{\frac{\alpha}{\alpha-1}} - V(q) \right\}, \quad 1 < \alpha \leq 2, \quad (20)$$

Let's now treat the action  $S_\alpha(q, t_b)$  as a function of coordinate  $q$  and the upper limit of integration,  $t_b$ . Further, we suppose that motion goes along the actual path, that is we consider action as a functional of actual trajectory  $q(\tau)$  in coordinate space. To evaluate the variance  $\delta S_\alpha(q, t_b)$  we have to compare the values of  $S_\alpha(q, t_b)$  for trajectories having a common start point at  $q(t_a)$ , but passing through different end points at the  $t_b$ . Thus, for the variation of the action by the variation of the end point  $\delta q(t_b)$  of the trajectory we have

$$\delta S_\alpha(q, t_b) = \delta \int_{t_a}^{t_b} d\tau L_\alpha(\dot{q}, q) = \frac{\partial L_\alpha}{\partial \dot{q}} \delta q|_{t_a}^{t_b} + \int_{t_a}^{t_b} d\tau \delta q \left( \frac{\partial L_\alpha}{\partial q} - \frac{d}{dt} \frac{\partial L_\alpha}{\partial \dot{q}} \right) \quad (21)$$

Because of Eq.(8) and  $\delta q(t_a) = 0$  we obtain

$$\delta S_\alpha(q, t_b) = p \delta q(t_b),$$

or

$$p = \frac{\delta S_\alpha(q, t_b)}{\delta q(t_b)}. \quad (22)$$

For simplicity, letting  $\delta q(t_b) = \delta q$  and  $t_b = t$ , we can write

$$\frac{dS_\alpha(q, t)}{dt} = \frac{\partial S_\alpha(q, t)}{\partial t} + \frac{\partial S_\alpha(q, t)}{\partial q} \dot{q} = \frac{\partial S_\alpha(q, t)}{\partial t} + p\dot{q}. \quad (23)$$

According to the definition of the action its total time derivative along the trajectory is  $dS_\alpha/dt = L_\alpha$ , and Eq.(23) can be presented as

$$\frac{\partial S_\alpha(q, t)}{\partial t} = L_\alpha - p\dot{q} = -H_\alpha(p, q), \quad (24)$$

where Eq.(4) has been taken into account and  $p$  is given by Eq.(22).

Therefore, we come to the equation

$$\frac{\partial S_\alpha(q, t)}{\partial t} + H_\alpha\left(\frac{\partial S_\alpha(q, t)}{\partial q}, q\right) = 0. \quad (25)$$

With  $H_\alpha(p, q)$  given by Eq.(2) it takes the form

$$\frac{\partial S_\alpha(q, t)}{\partial t} + D_\alpha \left| \frac{\partial S_\alpha(q, t)}{\partial q} \right|^\alpha + V(q) = 0, \quad 1 < \alpha \leq 2, \quad (26)$$

where  $S_\alpha$  is called the Hamilton's principal function.

For time-independent Hamiltonian the variables  $q$  and  $t$  in this equation can be separated. That is, we can search for solution of Eq.(26) in the form

$$S_\alpha(q, t, E) = S_\alpha^{(0)}(q, E) - Et, \quad (27)$$

where  $S_\alpha^{(0)}(q, E)$  is time-independent Hamilton's principal function and  $E$ , is the constant of integration, which has been identified with the total energy. Substituting Eq.(27) into Eq.(26) yields

$$D_\alpha \left| \frac{\partial S_\alpha^{(0)}(q, E)}{\partial q} \right|^\alpha + V(q) = E, \quad 1 < \alpha \leq 2. \quad (28)$$

Equations (26) and (28) are the Hamilton-Jacobi equations of fractional classical mechanics.

As an example, let us consider a free particle,  $V(q) = 0$ . The Hamilton-Jacobi equation Eq.(26) for a free particle is

$$\frac{\partial S_\alpha(q, t)}{\partial t} + D_\alpha \left| \frac{\partial S_\alpha(q, t)}{\partial q} \right|^\alpha = 0, \quad 1 < \alpha \leq 2. \quad (29)$$

Therefore, from Eqs.(27) and (28) we obtain a solution to the Hamilton-Jacobi equation Eq.(29),

$$S_\alpha(q, t, E) = \left(\frac{E}{D_\alpha}\right)^{1/\alpha} q - Et. \quad (30)$$

Follow the Hamilton-Jacobi fundamentals (see, for instance, [5]) we differentiate Eq.(30) over the energy  $E$  and put the derivative equal to a new constant  $\delta$ ,

$$\delta = \frac{\partial S_\alpha(q, t, E)}{\partial E} = \frac{1}{\alpha D_\alpha} \left(\frac{E}{D_\alpha}\right)^{\frac{1}{\alpha}-1} q - t, \quad (31)$$

which yields

$$q = \alpha D_\alpha \left(\frac{E}{D_\alpha}\right)^{1-\frac{1}{\alpha}} (t + \delta), \quad (32)$$

and

$$p = \frac{\partial S_\alpha(q, t, E)}{\partial q} = \left(\frac{E}{D_\alpha}\right)^{1/\alpha}, \quad (33)$$

where  $E > 0$  and  $1 < \alpha \leq 2$ .

The equations (32) and (33) define the trajectory of a free particle in fractional classical mechanics framework.

In limit case when  $\alpha = 2$  and  $D_2 = 1/2m$ , Eqs.(9), (15), (26) and (28) are transformed into the well-known Hamilton, Lagrange and Hamilton-Jacobi equations of classical mechanics for a particle with mass  $m$  moving in the potential field  $V(q)$ .

### 3 Mechanical similarity

First of all, we intend to study the general properties of 1D fractional classical motion without integrating motion equations (9), (15), and (28).

When the potential energy  $V(q)$  is a homogeneous function of coordinate  $q$ , it is possible to find some general similarity relationships.

Let us carry out a transformation in which the coordinates are changed by a factor  $\rho$  and the time by a factor  $\tau$ :  $q \rightarrow q' = \rho q$ ,  $t \rightarrow t' = \tau t$ . Then all velocities  $\dot{q} = dq/dt$  are changed by a factor  $\rho/\tau$ , and the kinetic energy by a factor  $(\rho/\tau)^{\alpha/(\alpha-1)}$ . If  $V(q)$  is a homogeneous function of degree  $\beta$  then it satisfies,



$$V(\rho q) = \rho^\beta V(q). \quad (34)$$

It is obvious that if  $\rho$  and  $\tau$  are such that  $(\rho/\tau)^{\alpha/(\alpha-1)} = \rho^\beta$ , i.e.  $\tau = \rho^{1-\beta+\frac{\beta}{\alpha}}$ , then the transformation leaves the motion equation unaltered. This invariance is called mechanical similarity.

A change of all the coordinates of the particles by the same scale factor corresponds to replacement of the classical mechanical trajectories of the particles by other trajectories, geometrically similar but different in size. Thus, we conclude that, if the potential energy of the system is a homogeneous function of degree  $\beta$  in Cartesian coordinates, the fractional equations of motion permit a series of geometrically similar trajectories, and the times of the motion between corresponding points are in the ratio

$$\frac{t'}{t} = \left(\frac{l'}{l}\right)^{1-\beta+\frac{\beta}{\alpha}}, \quad (35)$$

where  $\frac{l'}{l}$  is the ratio of linear scales of the two paths. Not only the times but also any other mechanical quantities are in a ratio which is a power of  $\frac{l'}{l}$ . For example, the velocities and energies follow the scaling laws

$$\frac{\dot{q}'}{\dot{q}} = \left(\frac{l'}{l}\right)^{\beta-\frac{\beta}{\alpha}}, \quad \frac{E'}{E} = \left(\frac{l'}{l}\right)^\beta, \quad (36)$$

and

$$\frac{t'}{t} = \left(\frac{E'}{E}\right)^{\frac{1}{\alpha}+\frac{1}{\beta}-1}. \quad (37)$$

The following are some examples of the foregoing.

It follows from Eq.(37) that at  $\frac{1}{\alpha} + \frac{1}{\beta} = 1$  the period of oscillations does not depend on energy of oscillator. The condition  $\frac{1}{\alpha} + \frac{1}{\beta} = 1$  with  $1 < \alpha \leq 2$ ,  $1 < \beta \leq 2$  brings  $\alpha = 2$  and  $\beta = 2$  only. It means, for instance, that considering the fractional classical oscillator model with Hamiltonian (41), we can conclude that the standard classical harmonic oscillator only,

$$H_2(p, q) = p^2/2m + g^2 q^2, \quad (38)$$

has the period of oscillations which does not depend on energy of oscillator.

In the uniform field of force, the potential energy is a linear function of the coordinates, i.e.  $\beta = 1$ . From Eq.(37) we have  $\frac{t'}{t} = \left(\frac{l'}{l}\right)^{1/\alpha}$ . Therefore, the time of motion in a uniform field ( $\beta = 1$ ) is as the  $\alpha$ -root of the initial altitude.

If the potential energy is inversely proportional to the distance apart, i.e. it is a homogeneous function of degree  $\beta = -1$ , then Eq.(35) becomes

$$\frac{t'}{t} = \left(\frac{l'}{l}\right)^{2-\frac{1}{\alpha}}. \quad (39)$$

For instance, regarding the problem of orbital motion in 3D fractional classical mechanics, we can state that the orbital period to the power  $\alpha$  is proportional to the power  $2\alpha - 1$  of its orbit scale. This statement is in fact a generalization of the well-known Kepler's third law. We call Eq.(39) as *fractional Kepler's third law*. In special case  $\alpha = 2$ , Eq.(39) turns into the well-known Kepler's third law [5].

In general, for negative  $\beta$ ,  $\beta = -\gamma$ , in the space of the fractional Hamiltonians  $H_{\alpha,-\gamma}$  at

$$1 + \gamma - \frac{\gamma}{\alpha} = \frac{3}{2}, \quad (40)$$

there exists the subset of fractional dynamic systems, orbital motion of which follows the well-known Kepler's third law, that is, the square of the orbital period is proportional to the cube of the orbit size.

## 4 Integration of the motion equations

### 4.1 Fractional classical 1D oscillator

#### 4.1.1 The Lagrange approach

Fractional quantum oscillator model has been introduced in [1]. Fractional classical oscillator can be considered as a classical counterpart of the fractional quantum oscillator model.

We introduce fractional classical 1D oscillator as a mechanical system with the Hamiltonian

$$H(p, q) = D_{\alpha}|p|^{\alpha} + g^2|q|^{\beta}, \quad (41)$$

where  $g$  is a constant with physical dimension  $[g] = \text{erg}^{1/2} \cdot \text{cm}^{-\beta/2}$  and  $\alpha$  and  $\beta$  are parameters,  $1 < \alpha \leq 2$ ,  $1 < \beta \leq 2$ .

It follows from Eq.(6) that the Lagrangian of the fractional classical 1D oscillator is

$$L(\dot{q}, q) = \left( \frac{1}{\alpha D_\alpha} \right)^{\frac{1}{\alpha-1}} \frac{\alpha-1}{\alpha} |\dot{q}|^{\frac{\alpha}{\alpha-1}} - g^2 |q|^\beta, \quad 1 < \alpha \leq 2, \quad 1 < \beta \leq 2. \quad (42)$$

Hence, the motion equation of fractional classical 1D oscillator is

$$\left( \frac{1}{\alpha D_\alpha} \right)^{\frac{1}{\alpha-1}} \frac{1}{\alpha-1} \ddot{q} |\dot{q}|^{\frac{2-\alpha}{\alpha-1}} + \beta g^2 |q|^{\beta-1} \text{sgn} q = 0. \quad (43)$$

This is a new non-linear classical mechanics equation of motion. When  $\alpha = 2$  and  $\beta = 2$  it goes into the well-known linear equation of motion for classical mechanics oscillator which has the form,

$$m\ddot{q} + 2g^2 q = 0. \quad (44)$$

where  $m$  is a particle mass ( $D_2 = 1/2m$ ).

To integrate Eq.(43) we start from the law of energy conservation. For the Lagrangian Eq.(42) we have

$$D_\alpha \left( \frac{1}{\alpha D_\alpha} \right)^{\frac{\alpha}{\alpha-1}} |\dot{q}|^{\frac{\alpha}{\alpha-1}} + g^2 |q|^\beta = E, \quad (45)$$

where  $E$  is the total energy. Thus, we have

$$|\dot{q}| = \alpha D_\alpha^{1/\alpha} (E - g^2 |q|^\beta)^{\frac{\alpha-1}{\alpha}}, \quad (46)$$

which is a first-order differential equation, and it can be integrated.

Since the kinetic energy is positive, the total energy  $E$  always exceeds the potential energy, that is  $E > g^2 |q|^\beta$ . The points where the potential energy equals the total energy  $g^2 |q|^\beta = E$  are turning points of classical trajectory. The 1D motion bounded by two turning points is oscillatory, the particle moves repeatedly between those two points.

Substituting  $q = (E/g^2)^{1/\beta} y$  into Eq.(46) yields

$$|\dot{y}| = \alpha D_\alpha^{1/\alpha} g^{2/\beta} E^{1-(\frac{1}{\alpha} + \frac{1}{\beta})} (1 - |y|^\beta)^{1-\frac{1}{\alpha}}. \quad (47)$$

Hence, the period  $T(\alpha, \beta)$  of oscillations is

$$T(\alpha, \beta) = 4 \frac{E^{(\frac{1}{\alpha} + \frac{1}{\beta}) - 1}}{\alpha D_\alpha^{1/\alpha} g^{2/\beta}} \int_0^1 \frac{dy}{(1 - y^\beta)^{1 - \frac{1}{\alpha}}}.$$

By substituting  $z = y^\beta$  we rewrite the last equation in the form

$$T(\alpha, \beta) = 4 \frac{E^{(\frac{1}{\alpha} + \frac{1}{\beta}) - 1}}{\alpha \beta D_\alpha^{1/\alpha} g^{2/\beta}} \int_0^1 dz z^{\frac{1}{\beta} - 1} (1 - z)^{\frac{1}{\alpha} - 1}. \quad (48)$$

With the help of the  $B$ -function definition [6]

$$B\left(\frac{1}{\beta}, \frac{1}{\alpha}\right) = \int_0^1 dz z^{\frac{1}{\beta} - 1} (1 - z)^{\frac{1}{\alpha} - 1},$$

we finally find for the period of oscillations of fractional classical 1D oscillator

$$T(\alpha, \beta) = 4 \frac{E^{(\frac{1}{\alpha} + \frac{1}{\beta}) - 1}}{\alpha \beta D_\alpha^{1/\alpha} g^{2/\beta}} B\left(\frac{1}{\beta}, \frac{1}{\alpha}\right). \quad (49)$$

This new equation shows that the period depends on the energy of fractional classical 1D oscillator. The dependency on energy is in agreement with the scaling law given by Eq.(37).

It follows from Eq.(49) that period  $T(\alpha, \beta)$  doesn't depend on the energy of oscillator when  $(\frac{1}{\alpha} + \frac{1}{\beta}) = 1$ . If  $1 < \alpha \leq 2$  and  $1 < \beta \leq 2$  then condition  $(\frac{1}{\alpha} + \frac{1}{\beta}) = 1$  gives us that  $\alpha = 2$  and  $\beta = 2$ . Hence, we come to the standard classical mechanics harmonic oscillator with the Hamiltonian given by Eq.(38) and the energy independent oscillation period,  $T(2, 2) = \pi\sqrt{2m/g}$ .

Table 1 shows that the energy dependency of the period of fractional classical oscillator is a classical counterpart of the fractional quantum mechanics statement on non-equidistant energy levels of fractional quantum oscillator. For classical mechanics, independence on energy of the period of classical oscillator is a classical counterpart of the quantum mechanics statement on equidistant energy levels of quantum oscillator.

|                                     | Period of oscillations $T$   | Energy levels $\mathcal{E}_n$  |
|-------------------------------------|--|--|
| $1 < \alpha < 2$<br>$1 < \beta < 2$ | Fractional classical mechanics<br>$4 \frac{E^{(\frac{1}{\alpha} + \frac{1}{\beta}) - 1}}{\alpha \beta D_\alpha^{1/\alpha} g^{1/\beta}} B(\frac{1}{\beta}, \frac{1}{\alpha})$ | Fractional quantum mechanics<br>$\left( \frac{\pi \hbar \beta D_\alpha^{1/\alpha} g^{2/\beta}}{2B(\frac{1}{\beta}, \frac{1}{\alpha} + 1)} \right)^{\frac{\alpha \beta}{\alpha + \beta}} (n + \frac{1}{2})^{\frac{\alpha \beta}{\alpha + \beta}}$ |
| $\alpha = 2$<br>$\beta = 2$         | Classical mechanics<br>$\pi \sqrt{2m/g}$   | Quantum mechanics<br>$\hbar \sqrt{\frac{2}{m} g} (n + \frac{1}{2})$  |

Table 1. *Energy dependence/independence of the period of classical oscillator vs non-equidistant/equidistant energy levels of quantum oscillator*<sup>1</sup>.

#### 4.1.2 The Hamilton approach

For the fractional 1D oscillator the Hamilton equations of motion in accordance with Eqs.(14) and (41) are

$$\dot{q} = \frac{\partial H_{\alpha, \beta}}{\partial p} = \alpha D_\alpha |p|^{\alpha-1} \text{sgn} p, \quad (50)$$

$$\dot{p} = -\frac{\partial H_{\alpha, \beta}}{\partial q} = -\beta g^2 |q|^{\beta-1} \text{sgn} q, \quad (51)$$

where  $1 < \alpha \leq 2$ ,  $1 < \beta \leq 2$ .

Further, Hamilton equation (50) leads to

$$|p| = \left( \frac{1}{\alpha D_\alpha} \right)^{\frac{1}{\alpha-1}} |\dot{q}|^{\frac{1}{\alpha-1}}, \quad (52)$$

Substituting Eq.(52) into Eq.(41) yields exactly Eq.(45). Therefore, downstream integration can be done by the same way as it was done above in the framework of the Lagrange approach.

#### 4.1.3 The Hamilton-Jacobi approach

For fractional classical oscillator with the Hamilton function defined by Eq.(41) a complete integral of the Hamilton-Jacobi equation (27) is

---

<sup>1</sup>The period of oscillations  $T$  and frequency of oscillations  $\omega$  are related by  $T = 2\pi/\omega$ . Hence, the classical mechanics frequency is  $\omega = \sqrt{\frac{2}{m} g}$ , and in terms of the frequency, the quantum mechanic energy levels are  $\mathcal{E}_n = \hbar \omega (n + \frac{1}{2})$ .

The energy levels equation for the fractional quantum oscillator has been taken from [4].

$$S(q, t, E) = \int dq \left( \frac{1}{D_\alpha} (E - g^2 |q|^\beta)^{1/\alpha} - Et \right), \quad (53)$$

where the integration constant  $E$  can be identified with the total energy of the fractional classical oscillator.

It is known that in standard classical mechanics [5],  $\alpha = 2$ , the integral in Eq.(53) is considered as an indefinite integral. At this point, to move forward, we introduce the ansatz to treat the integral in Eq.(53) as

$$S(q, t, E) = \int_0^q dq \left( \frac{1}{D_\alpha} (E - g^2 |q|^\beta)^{1/\alpha} - Et \right). \quad (54)$$

Following the Hamilton-Jacobi fundamentals (see, for instance, [5]) we differentiate Eq.(53) over the energy  $E$  and put the derivative equal to a new constant  $\delta$ ,

$$\frac{\partial S(q, t, E)}{\partial E} = \frac{1}{\alpha D_\alpha^{1/\alpha}} \int_0^q dq' (E - g^2 |q'|^\beta)^{(1-\alpha)/\alpha} - t = \delta. \quad (55)$$

Substituting integration variable  $q'$  with new variable  $z$ ,  $q' = (E/g^2)^{1/\beta} z^{1/\beta}$  yields

$$t + \delta = \frac{E^{(\frac{1}{\alpha} + \frac{1}{\beta}) - 1}}{\alpha \beta D_\alpha^{1/\alpha} g^{2/\beta}} \int_0^{q^\beta g^2/E} dz z^{\frac{1}{\beta} - 1} (1 - z)^{\frac{1}{\alpha} - 1}. \quad (56)$$

The integral in the above equation can be expressed as the incomplete Beta function<sup>2</sup> and we obtain

$$t + \delta = \frac{E^{(\frac{1}{\alpha} + \frac{1}{\beta}) - 1}}{\alpha \beta D_\alpha^{1/\alpha} g^{2/\beta}} B_{q^\beta g^2/E} \left( \frac{1}{\beta}, \frac{1}{\alpha} \right). \quad (58)$$

This equation is the solution of the fractional 1D oscillator as function  $t(q)$  in terms of the incomplete Beta function.

---

<sup>2</sup>The incomplete Beta function is defined as [7]

$$B_x(\mu, \nu) = \int_0^x dy y^{\mu-1} (1-y)^{\nu-1}, \quad (57)$$

The incomplete Beta function  $B_x(\mu, \nu)$  has the hypergeometric representation

$$B_x(\mu, \nu) = \frac{x^\mu}{\mu} F(\mu, 1 - \nu; \mu + 1; x), \quad (59)$$

where  $F(\mu, 1 - \nu; \mu + 1; x)$  is the hypergeometric function [8]. Now we can write Eq.(58) in the form

$$t + \delta = \frac{E^{\frac{1}{\alpha}-1}}{\alpha D_\alpha^{1/\alpha}} q F\left(\frac{1}{\beta}, 1 - \frac{1}{\alpha}; \frac{1}{\beta} + 1; q^\beta g^2/E\right). \quad (60)$$

Thus, in terms of the hypergeometric function we found solution of the fractional 1D oscillator as function  $t(q)$ .

Let us show that the ansatz given by Eq.(54) and based on it solution given by Eq.(60) allow us to reproduce the well-known solution to standard classical harmonic oscillator,  $\alpha = 2$  and  $\beta = 2$ . Indeed, at  $\alpha = 2$  and  $\beta = 2$  we have

$$\omega(t + \delta) = x F\left(\frac{1}{2}, \frac{1}{2}; \frac{3}{2}; x^2\right), \quad (61)$$

where a new variable  $x$  has been introduced as,

$$x = g\sqrt{1/E}q, \quad (62)$$

and  $\omega = \sqrt{2/mg}$  is the frequency of classical 1D oscillator. Since  $x F(\frac{1}{2}, \frac{1}{2}; \frac{3}{2}; x^2) = \arcsin x$ , it follows from Eq.(61) that

$$x(t) = \sin \omega(t + \delta).$$

By restoring the original dynamic variable  $q(t)$  from Eq.(62) we obtain,

$$q(t) = \sqrt{\frac{2E}{m\omega^2}} \sin \omega(t + \delta), \quad (63)$$

which is the solution of the Hamilton-Jacobi equation for standard 1D harmonic oscillator with the Hamiltonian given by Eq.(38).

Thus, we proved that the ansatz given by Eq.(54) gives us the well-known solution to standard classical mechanics harmonic oscillator.

## 5 Conclusion

Fractional classical mechanics has been introduced as a classical counterpart of fractional quantum mechanics. The Lagrange, Hamilton and Hamilton-Jacobi frameworks have been developed for fractional classical mechanics. Scaling analysis of fractional classical motion equations has been implemented based on the mechanical similarity. We discover and discuss *fractional Kepler's third law* which is a generalization of the well-known Kepler's third law.

Fractional classical oscillator model has been introduced and motion equations for the fractional classical oscillator have been integrated. We found an equation for the period of oscillations of fractional classical oscillator. The map between the energy dependence of the period of classical oscillations and the non-equidistant distribution of the energy levels for fractional quantum oscillator has been established.

In the case when  $\alpha = 2$ , all new developments are turned into the well-known results of the classical mechanics.

## References

- [1] N. Laskin, Phys. Lett. A **268**, 298 (2000).
- [2] N. Laskin, Phys. Rev. E **62**, 3135 (2000) (also available online: <http://arxiv.org/abs/0811.1769>).
- [3] N. Laskin, Chaos **10**, 780 (2000).
- [4] N. Laskin, Phys. Rev. E **66**, 056108 (2002) (also available online: <http://arxiv.org/abs/quant-ph/0206098>)
- [5] L. D. Landau and E. M. Lifshitz, Mechanics, Vol. 1 of Course of Theoretical Physics, 3<sup>rd</sup> Edition (Pergamon Press, 1976).
- [6] <http://dlmf.nist.gov/5.12.E1>
- [7] <http://dlmf.nist.gov/8.17.E1>
- [8] <http://dlmf.nist.gov/8.17.E7>