A remark about polynomials with specified local minima and no other critical points

Eduardo D. Sontag

1 Introduction

This fact must surely be well-known, but it seems worth giving a simple and quite explicit proof:

Proposition 1.1 Take any finite subset X of \mathbb{R}^n , $n \ge 2$. Then, there is a polynomial function $P : \mathbb{R}^n \to \mathbb{R}$ which has local minima on the set X, and has no other critical points.

A weaker version, constructing one particular subset X of \mathbb{R}^2 with the stated property, was shown in a beautiful little note by an undergraduate, Ian Robertson, as part of an REU conducted by Alan Durfee [1]; references to this work, and the context from degree theory, can be found in [2]. Our construction generalizes Robertson's.

We were interested in this question because of the following consequence, obtained immediately by considering the gradient vector field $g(x) = -\nabla P(x)$:

Corollary 1.2 Take any finite subset X of \mathbb{R}^n , $n \ge 2$. Then, there is a polynomial vector field $g : \mathbb{R}^n \to \mathbb{R}^n$ which has asymptotically stable equilibria on the set X, and has no other equilibria.

Suppose that g is as in Corollary 1.2 and that X has more than one element. Let \mathcal{O}_x , $x \in X$, be the domains of attraction of the points in X. The union of the sets \mathcal{O}_x cannot equal all of \mathbb{R}^n , since these are disjoint open sets; pick any $\xi \in \mathbb{R}^n \setminus \bigcup_x \mathcal{O}_x$. The omega-limit set $\Omega^+(\xi)$ cannot intersect X (since points of X are asymptotically stable). Thus there are points that do not converge to any equilibria. (Alternatively, one could arrive at the same conclusion appealing to topological degree arguments.) Thus, the following is also of interest (and much easier to prove).

Proposition 1.3 Take any finite subset X of \mathbb{R}^n , $n \ge 2$. Then, there are a finite subset $X' \subset \mathbb{R}^n$ and a polynomial vector field $g : \mathbb{R}^n \to \mathbb{R}^n$ which has asymptotically stable equilibria on the set X, saddles on the set X', and no other equilibria, and moreover: (1) every solution of $\dot{x} = g(x)$ converges to $X \bigcup X'$ and (2) except for a measure-zero set of initial conditions, every solution converges to an equilibrium in X.

2 Proofs

We prove Proposition 1.1 by first treating the case of a set $X \subset \mathbb{R}^2$ of the special form $\mathcal{X} \times \{0\}$, and then using a coordinate change to reduce the general case to this one. The proof of Proposition 1.3, in contrast, only requires the coordinate change, plus a trivial construction.

2.1 A special case

Lemma 2.1 Let $\alpha : \mathbb{R} \to \mathbb{R}$ be a \mathcal{C}^2 function whose zeroes are all simple; that is, on the set

$$\mathcal{X} := \{ x \, | \, \alpha(x) = 0 \}$$

it holds that $\alpha'(x) \neq 0$. Introduce the following function $f : \mathbb{R}^2 \to \mathbb{R}$:

$$f(x,y) := \left(\alpha(x) - \left[\alpha(x) - \alpha'(x)\right]^2 y\right)^2 - \int \alpha(x) \left[\alpha(x) - \alpha'(x)\right] dx$$

(where the last term denotes an arbitrary anti-derivative). Consider the set of critical points of f,

$$\mathcal{C}(f) := \{(x,y) \mid f_x(x,y) = f_y(x,y) = 0\}$$

Then:

- 1. $\mathcal{C}(f) = \mathcal{X} \times \{0\}.$
- 2. At each $(x, y) \in C(f)$, the Hessian of f is positive definite.

As a consequence, f has local minima at the points in $\mathcal{X} \times \{0\}$, and no other critical points.

Proof. For convenience, we introduce $\beta(x) := \alpha(x) - \alpha'(x)$, so that

$$f(x,y) := \left(\alpha(x) - \beta(x)^2 y\right)^2 - \int \alpha(x)\beta(x) \, dx$$

We have:

$$f_x(x,y) = 2 \left(\alpha(x) - \beta(x)^2 y \right) \left(\alpha'(x) - 2\beta(x)\beta'(x)y \right) - \alpha(x)\beta(x)$$

$$f_y(x,y) = -2 \left(\alpha(x) - \beta(x)^2 y \right) \beta(x)^2.$$

Clearly, $\alpha(x) = 0$ and y = 0 imply $f_x(x, y) = f_y(x, y) = 0$, so we must only prove the converse. Pick any $(x, y) \in C(f)$.

From $f_y(x, y) = 0$, we have that one of these must hold:

$$\beta(x) = 0, \qquad (1)$$

$$\beta(x) \neq 0 \text{ and } \alpha(x) - \beta(x)^2 y = 0.$$
 (2)

If (1) holds, then $0 = f_x(x, y) = 2\alpha(x)\alpha'(x)$ implies that either $\alpha(x) = 0$ or $\alpha'(x) = 0$. On the other hand, the assumption of simple zeroes, " $\alpha(x) = 0 \Rightarrow \alpha'(x) \neq 0$ " can also be written in contrapositive form as " $\alpha'(x) = 0 \Rightarrow \alpha(x) \neq 0$," from which we have:

$$\begin{aligned} \alpha(x) &= 0 \quad \Rightarrow \quad \beta(x) = -\alpha'(x) \neq 0\\ \alpha'(x) &= 0 \quad \Rightarrow \quad \beta(x) = \alpha(x) \neq 0 \,. \end{aligned}$$

This rules out both $\alpha(x) = 0$ and $\alpha'(x) = 0$ when $\beta(x) = 0$, and thus case (1) cannot hold.

So case (2) holds, which means that $y = \alpha(x)/\beta(x)^2$. On the other hand, when $\alpha(x) - \beta(x)^2 y = 0$, we have that $0 = f_x(x, y) = -\alpha(x)\beta(x)$, and since $\beta(x) \neq 0$, it follows that $\alpha(x) = 0$, from which it also follows that y = 0. We conclude that $(x, y) \in \mathcal{X} \times \{0\}$, as desired.

To prove positive definiteness of the Hessian on C(f), we must show that on the set C(f), both $f_{yy}(x, y) > 0$ and $\Delta(x, y) > 0$, where $\Delta = f_{xx}f_{yy} - f_{xy}^2$. In general:

$$f_{xx} = 2(\alpha'(x) - 2\beta(x)\beta'(x)y)^{2} + 2(\alpha(x) - \beta(x)^{2}y)(\alpha''(x) - 2[\beta(x)\beta''(x) + (\beta'(x))^{2}]y) -\alpha'(x)\beta(x) - \alpha(x)\beta'(x)$$

$$f_{yy} = 2\beta(x)^{4}$$

$$f_{xy} = f_{yx} = -2(\alpha'(x) - 2\beta(x)\beta'(x)y)\beta(x)^{2} - 4(\alpha(x) - \beta(x)^{2}y)\beta(x)\beta'(x)$$

so in particular, on the set C(f), since $\alpha(x) = 0$ and y = 0:

$$f_{xx} = 2 (\alpha'(x))^2 - \alpha'(x)\beta(x)$$

$$f_{yy} = 2\beta(x)^4$$

$$f_{xy} = f_{yx} = -2 (\alpha'(x)) \beta(x)^2.$$

Since on this set $\beta = -\alpha'(x)$:

$$f_{xx} = 3 \left[\alpha'(x) \right]^2$$

$$f_{yy} = 2 \left[\alpha'(x) \right]^4$$

$$f_{xy} = f_{yx} = -2 \left[\alpha'(x) \right]^3$$

and thus

$$\Delta(x, y) = 2[\alpha'(x)]^6.$$

As $\alpha'(x) \neq 0$ on the set $\mathcal{C}(f)$, it follows that both $f_{yy} > 0$ and $\Delta > 0$, which completes the proof.

2.2 A coordinate change

Lemma 2.2 Let X be a finite subset of \mathbb{R}^n , $n \ge 2$. Then, there exists a polynomial map $F : \mathbb{R}^n \to \mathbb{R}^n$ such that:

$$F(X) \subseteq \mathbb{R} \times \{0\}^{n-1},\tag{3}$$

$$F$$
 has a polynomial inverse. (4)

Proof. For each pair of points $\xi \neq \eta$ in X, let $V_{\xi\eta} = \{p \in \mathbb{R}^n | p^T(\xi - \eta) = 0\}$. Pick a point p not belonging to the union of the finitely many hyperplanes $V_{\xi\eta}$. Choose any invertible mapping T which has p^T as its first row, and consider the change of variables z = Tx. Let $X = \{x^{(1)}, \ldots, x^{(k)}\}$ and $Z := TX = \{z^{(1)}, \ldots, z^{(k)}\}$, where $z^{(i)} = Tx^{(i)}$. Since $x \mapsto p^T x$ is one to one on the set X, the first coordinates of the $z^{(i)}$'s are all distinct, that is

 $z^{(i)} = (z_1^{(i)}, \dots, z_n^{(i)})$

and $z_1^{(i)} \neq z_1^{(j)}$ for each $i \neq j$. For each j = 2, ..., n, let p_j be the Lagrange interpolation polynomial that gives:

$$p_j(z_1^{(i)}) = z_j^{(i)}, \qquad i = 1, \dots, k$$

and define

$$\Pi: \mathbb{R}^n \to \mathbb{R}^n: \begin{pmatrix} z_1\\ z_2\\ \vdots\\ z_n \end{pmatrix} \mapsto \begin{pmatrix} z_1\\ z_2 - p_2(z_1)\\ \vdots\\ z_n - p_n(z_1) \end{pmatrix}.$$

This is invertible (with a polynomial inverse obtained by using instead " $z_j + p_j(z_1)$ " for each j > 1). Moreover, by definition, on the set Z we have that the *j*th coordinate of P(z), $P(z^{(i)})_j = z_j^{(i)} - p_j(z_1^{(i)}) = 0$, j > 1. In other words, P maps into $\mathbb{R} \times \{0\}^{n-1}$. The proof is completed by picking the composition $F = \Pi \circ T$.

2.3 **Proof of Proposition 1.1**

Let X be a finite subset of \mathbb{R}^n , $n \ge 2$, pick F as in Lemma 2.2, and let \mathcal{X} be such that $F(X) = \mathcal{X} \times \{0\}^{n-1}$. We will construct a polynomial $Q : \mathbb{R}^n \to \mathbb{R}$ whose only critical points are on the set $\mathcal{X} \times \{0\}^{n-1}$, and the Hessian is positive definite there. Then $P = Q \circ F$ will be as desired, because diffeomorphisms preserve critical points and their signature. (To be explicit: as $\nabla P(x) = \nabla Q(F(x)) \cdot JF(x)$ and the Jacobian JF is everywhere nonsingular, critical points map to critical points. Furthermore, at a critical point, the Hessian H transforms as $J^T H J$, so positive definiteness is preserved.)

Let
$$\alpha(x) := \prod_{i=1}^{k} (x - a_i)$$
, where $\mathcal{X} = \{a_1, \dots, a_k\}$. This α is as in Lemma 2.1; let f be as there. We define
 $Q(x_1, \dots, x_n) := f(x_1, x_2) + \frac{1}{2} \sum_{i>2} x_i^2$.

From $\partial Q/\partial x_i = 0$ for i > 2, the critical points of Q have $x_i = 0$ for all i > 2, so $\mathcal{C}(Q) = \mathcal{C}(f) \times \{0\}^{n-2} = \mathcal{X} \times \{0\}^{n-1}$. Moreover, at these points, the Hessian of Q is obtained by appending an identity matrix to the Hessian of f, and thus it is also positive definite, as required.

2.4 **Proof of Proposition 1.3**

We use the same change of variables, so that, up to a diffeomorphism, we can assume without loss of generality that $X = \mathcal{X} \times \{0\}^{n-1}$. Let

$$\gamma(x) := -\prod_{i=1}^{k} (x - a_i) \prod_{i=1}^{k-1} (x - b_i)$$

where $\mathcal{X} = \{a_1 < \ldots < a_k\}$ and the $b_i \in (a_i, a_{i+1})$ are arbitrary. The scalar differential equation $\dot{x} = \gamma(x)$ has stable equilibria at \mathcal{X} , unstable at $\mathcal{X}' = \{b_1, \ldots, b_{k-1}\}$, and no other equilibria. We then define

$$g(x) = (\gamma(x), -x_2, \dots, -x_n).$$

This satisfies the conclusions of the Lemma, with $X = \mathcal{X} \times \{0\}^{n-1}$ and $X' = \mathcal{X}' \times \{0\}^{n-1}$.

References

- [1] Ian Robertson. A polynomial with *n* maxima and no other critical points. Preprint; available at http://www.mtholyoke.edu/~adurfee/reu/92/reu92.htm, 1992.
- [2] Alan H. Durfee. The index of grad f(x, y). Topology **37**: 1339-1361, 1998