

Morse-Smale systems and horseshoes for three dimensional singular flows

Shaobo Gan

Dawei Yang*

Abstract

We prove that for every three-dimensional vector field, either it can be accumulated by Morse-Smale ones, or it can be accumulated by ones with a transverse homoclinic intersection of some hyperbolic periodic orbit in the C^1 topology.

1 Introduction

1.1 The main result

One of the main subjects in differentiable dynamical systems is to describe the dynamics of “most” dynamical systems. These theories were established in the last century. See [2] for instance. An important progress is due to Peixoto [47]:

Theorem (Peixoto). *Assume that M^2 is a closed surface. A C^1 vector field on M^2 is C^1 structurally stable vector field iff it is Morse-Smale and every vector field could be accumulated by a structurally stable one in the C^1 topology.*

Smale was interested in the generalization of Peixoto’s result and he asked whether Morse-Smale vector fields are dense in the space of vector fields. Soon, Levinson and Thom pointed out that Morse-Smale vector fields would not be dense (without a rigorous proof). See [2]. Smale noticed the point and he constructed his famous horseshoe (for two dimensional diffeomorphisms or three-dimensional vector fields) [55] which shows that the dynamics may be very complicated and Morse-Smale systems would not be dense in the space of diffeomorphisms or vector fields. As in [2, Page 16]: “At that moment the world turned upside down . . . , and a new life began”.

Actually, Poincaré found an important phenomenon in his famous work [49] on celestial mechanics, which was called “doubly asymptotical solution”. Nowadays mathematicians call it *transverse homoclinic intersection*. Smale found that his horseshoe is closely related to transverse homoclinic intersections. Three classical results are known:

- Poincaré showed that transverse homoclinic intersections can survive under small perturbations. Moreover, if a system has one transverse homoclinic intersection, then it has infinitely many transverse homoclinic intersections [49].
- Birkhoff showed that if a plane system has one transverse homoclinic intersection, then it has infinitely many hyperbolic periodic orbits [6].
- Smale proved that the existence of transverse homoclinic intersection is equivalent to the existence of horseshoe [55].

*D. Yang is the corresponding author. S. G. is supported by 973 project 2011CB808002, NSFC 11025101 and 11231001. D.Y. was partially supported by NSFC 11001101, Ministry of Education of P. R. China 20100061120098.

Hence there are two kinds of typical dynamical systems: Morse-Smale system or a system with a horseshoe. Their dynamical behavior is quite different:

- The dynamics of Morse-Smale system is very simple: the chain recurrent set of a Morse-Smale system is a set containing finitely many hyperbolic periodic orbits or singularities. The topological entropy is robustly zero.
- The dynamics of a system with a horseshoe is very complicated: its chain recurrent set contains a non-trivial basic set with dense periodic orbits. The topological entropy is robustly positive.

Is there other typical dynamics beyond the above two ones? Palis formulated the idea for diffeomorphisms, and he conjectured that

Conjecture (Palis [43, 44, 45]). *Every system can be approximated either by Morse-Smale systems or by systems exhibiting a horseshoe (non-trivial hyperbolic basic set).*

In this paper, we manage to prove such kind of results for three dimensional vector fields.

Theorem A (Main Theorem). *Every three dimensional vector field can be C^1 approximated by Morse-Smale ones or by ones exhibiting a Smale horseshoe. In other words, Morse-Smale vector fields and vector fields with a non-trivial hyperbolic set (Smale horseshoe) form a dense open set in the space of C^1 three dimensional vector fields.*

Important progress has been made for the conjecture of Palis for diffeomorphisms: in C^1 topology, Pujals-Sambarino [52] proved it for two-dimensional diffeomorphisms (as a corollary of a stronger result); Bonatti-Gan-Wen [11] gave a prove for three-dimensional diffeomorphisms; and finally Crovisier [16] proved the conjecture for *any*-dimensional diffeomorphisms.

Comparing with diffeomorphism case, singularities of vector fields bring more difficulties. This prevents one to use some techniques of diffeomorphisms to singular vector fields, such as Crovisier's central model. By considering the sectional Poincaré maps of the flows, sometimes one can get some (not all) similar properties between d -dimensional vector fields and $(d - 1)$ -dimensional diffeomorphisms. But singular vector field displays different dynamics, e.g., the famous *Lorenz attractor* [30]. In the spirit of Lorenz attractor, [1, 19, 20] constructed *geometric Lorenz attractor* in a theoretical way. Roughly, geometric Lorenz attractor is a robust attractor of three-dimensional vector field, and it contains a hyperbolic singularity which is accumulated by hyperbolic periodic orbits in a robust way.

Lorenz attractor is not hyperbolic because of the existence of singularity. On the other hand, Mañé [31] showed that a robust attractor of a surface diffeomorphism is hyperbolic. This implies that the dynamics of 3-dimensional singular flows are different from 2-dimensional diffeomorphisms. Morales-Pacifico-Pujals [34, 35, 36] studied geometric Lorenz attractor in an abstract way. They found the right concept, i.e., *singular hyperbolicity*, to describe the hyperbolicity of Lorenz attractor, and they proved that a robust transitive set of a three-dimensional vector field is singular hyperbolic. But the dynamics of singular hyperbolic set are not as clear as hyperbolic set. For instance, there is no shadowing lemma of Anosov-Bowen type.

The dynamics of general transitive sets with singularities for three-dimensional vector fields are even more unclear for us than singular hyperbolic sets, even if the transitive sets have some dominated splitting with respect to the linear Poincaré flow. These are main difficulties that we encounter. For a non-trivial transitive set *without singularities* of a generic any-dimensional vector field, one can adapt Crovisier's central model [16] to get a transverse homoclinic intersection of a hyperbolic periodic orbit.

Let us be more precise. Let M^d be a d -dimensional C^∞ compact Riemannian manifold without boundary. Denote by $\mathcal{X}^1(M^d)$ the space of C^1 vector fields on M^d . Given $X \in \mathcal{X}^1(M^d)$,

denote by $\phi_t = \phi_t^X$ the C^1 flow generated by X and $\Phi_t = d\phi_t : TM^d \rightarrow TM^d$ the tangent flow on the tangent bundle TM^d . If $X(\sigma) = 0$, then σ is called a *singularity* of X . Other points are called *regular*. Let $\text{Sing}(X)$ be the set of singularities of X . For a regular point p , if $\phi_t(p) = p$ for some $t > 0$, then p is called *periodic*. Let $\text{Per}(X)$ be the set of periodic points of X . If $x \in \text{Sing}(X) \cup \text{Per}(X)$, then x is called a *critical point* of X and $\text{Orb}(x)$ is called a *critical orbit* or *critical element* of X .

For an invariant set Λ and a Φ_t -invariant bundle $E \subset T_\Lambda M^d$, we say that E is *contracting* (w.r.t. the tangent flow Φ_t) if there are constants $C \geq 1, \lambda > 0$ such that $\|\Phi_t|_{E(x)}\| \leq Ce^{-\lambda t}$ for every $x \in \Lambda$ and $t \geq 0$; we say that E is *expanding* if it is contracting for $-X$.

An invariant set Λ of X is *hyperbolic* if TM^d has continuous Φ_t -invariant splitting

$$T_\Lambda M^d = E^s \oplus \langle X \rangle \oplus E^u$$

(where the fibre $\langle X(x) \rangle$ at x is 0-dimensional or 1-dimensional according to x is a singularity or not), such that E^s is contracting and E^u is expanding. If $\dim E^s$ is independent of $x \in \Lambda$, then $\dim E^s$ is called the *index* of Λ .

For a critical point x , if $\text{Orb}(x)$ is a hyperbolic set, then we say that x or $\text{Orb}(x)$ is hyperbolic. One can define its index as the index of the hyperbolic set $\text{Orb}(x)$.

Recall that a C^1 vector field X is *Morse-Smale* if the non-wandering set $\Omega(X)$ of X consists of only finitely many hyperbolic critical elements and their stable and unstable manifolds intersect transversely. We use \mathcal{MS} to denote the set of Morse-Smale vector fields in $\mathcal{X}^1(M^d)$. For a hyperbolic periodic orbit γ , define

$$\begin{aligned} W^s(\gamma) &= \{x \in M^d : \lim_{t \rightarrow +\infty} d(\phi_t(x), \gamma) = 0.\} \\ W^u(\gamma) &= \{x \in M^d : \lim_{t \rightarrow -\infty} d(\phi_t(x), \gamma) = 0.\} \end{aligned}$$

We know ([22]) that $W^s(\gamma)$ and $W^u(\gamma)$ are submanifolds, which are called the stable and unstable manifolds of γ . If $W^s(\gamma) \cap W^u(\gamma) \setminus \gamma \neq \emptyset$, then one says that γ has a *transverse homoclinic orbit*¹. One says that X has a *transverse homoclinic orbit* if for some hyperbolic periodic orbit γ of X , γ has a transverse homoclinic orbit. Recall that: Birkhoff-Smale theorem asserts that the existence of transverse homoclinic orbits is equivalent to the existence of Smale's horseshoe (non-trivial hyperbolic basic set). We denote:

$$\mathcal{HS} = \{X \in \mathcal{X}^1(M^d) : X \text{ has a transverse homoclinic orbit}\}.$$

One can restate Theorem A as:

$$\mathcal{MS} \cup \mathcal{HS} \text{ is open and dense in } \mathcal{X}^1(M^3).$$

1.2 More on three-dimensional flows

Given a vector field X , let ϕ_t be the flow generated by X . For any $\varepsilon > 0$, $\{x_0, x_1, \dots, x_n\}$ is called an ε -*chain* (or ε -pseudo-orbit) from x_0 to x_n if there are $t_i \geq 1$ such that $d(\phi_{t_i}(x_i), x_{i+1}) < \varepsilon$ for any $0 \leq i \leq n-1$. For $x, y \in M^d$, one says that y is *chain attainable* from x if for any $\varepsilon > 0$, there is an ε -chain from x and y . If x is chain attainable from itself, then x is called a *chain recurrent point*. The set of chain recurrent points is called *chain recurrent set* of X , and denoted by $\text{CR}(X)$. Chain bi-attainability is a closed equivalence relation in $\text{CR}(X)$. For each $x \in \text{CR}(X)$, the equivalent class containing x is called the chain recurrent class of x , and denoted by $C(x)$ or $C(\text{Orb}(x))$. These are standard by Conley's theory [14].

¹Singularities cannot have transverse homoclinic intersections.

Bonatti and Crovisier [7] extended the C^1 connecting lemma to pseudo-orbits. An application of their result gave a useful classification of chain recurrent classes for C^1 -generic vector fields²: if a chain recurrent class contains a periodic orbit, then it is the *homoclinic class* of this periodic orbit; otherwise, it is called an *aperiodic class*. Here, the *homoclinic class* of a hyperbolic periodic orbit is defined to be the closure of all transverse homoclinic orbits of this periodic orbit.

Hyperbolic periodic orbits may have non-transverse homoclinic intersections, which are called *homoclinic tangencies*. Newhouse [37, 38, 39] studied the bifurcations of homoclinic tangencies crucially, which generate rich dynamics. Newhouse phenomena give typical dynamics beyond uniformly hyperbolic dynamics. There are many results for diffeomorphisms far away from ones with a homoclinic tangency. One can see the introduction of [17].

In this work, we can prove that every *non-trivial* chain recurrent class is a homoclinic class for C^1 generic vector fields which are far away from homoclinic tangencies. Here, a chain recurrent class is called *non-trivial* if it is not reduced to be a critical orbit.

Theorem B. *There is a dense G_δ set $\mathcal{R} \subset \mathcal{X}^1(M^3)$ such that, for every $X \in \mathcal{R}$, if X cannot be accumulated by ones with a homoclinic tangency, then every non-trivial chain recurrent class of X is a homoclinic class.*

Theorem B is stronger than Theorem A. We will see this point in Section 4.

An important conjecture made by Palis for surface diffeomorphisms is: every two-dimensional diffeomorphism can be accumulated either by ones with a homoclinic tangency, or by uniformly hyperbolic ones. This was proved by Pujals-Sambarino [52] in the C^1 topology. For three-dimensional vector fields, as mentioned in [42], excluding homoclinic tangencies and uniform hyperbolic systems, the typical dynamics may include the homoclinic orbits of singularities or Lorenz-like attractors. Arroyo-Rodríguez [5] proved the conjecture of Palis if homoclinic orbits of singularities were involved for three-dimensional vector fields. It is still an open problem about the density of Lorenz-like attractors or repellers beyond uniform hyperbolicity and homoclinic bifurcations of periodic orbits (even in the C^1 topology).

Morales-Pacífico-Pujals [35, 36] defined what is “Lorenz-like” in a dynamical way. In [34], Morales-Pacífico gave the notion of singular Axiom A without cycle. Let’s be more precise.

We say that a continuous invariant splitting $T_\Lambda M^d = E \oplus F$ w.r.t. the tangent flow over a compact invariant set Λ is a *dominated splitting* with respect to the tangent flow Φ_t if there are constants $C \geq 1, \lambda > 0$, such that $\|\Phi_t|_{E(x)}\| \|\Phi_{-t}|_{F(\phi_t(x))}\| \leq Ce^{-\lambda t}$ for every $x \in \Lambda$ and $t \geq 0$. For a compact invariant set Λ , we say that Λ admits a *partially hyperbolic splitting* if there is a continuous invariant splitting $T_\Lambda M^d = E^s \oplus E^c \oplus E^u$ w.r.t. Φ_t , where E^s is contracting, E^u is expanding, and both $E^s \oplus (E^c \oplus E^u)$ and $(E^s \oplus E^c) \oplus E^u$ are dominated splittings. In the above definition, E^s or E^u is allowed to be trivial.

Definition 1.1. *A transitive set Λ of $X \in \mathcal{X}^1(M^3)$ is called a singular hyperbolic attractor if*

1. *There is a neighborhood U of Λ such that*

$$\Lambda = \bigcap_{t \geq 0} \phi_t(U).$$

2. *Λ contains a singularity of index 2, and every singularity in Λ has index 2.*
3. *Λ admits a partially hyperbolic splitting $T_\Lambda M^3 = E^s \oplus E^{cu}$, where $\dim E^s = 1$ and E^{cu} area-expanding: there are constants $C \geq 1, \lambda > 0$ such that for any $x \in \Lambda$ and for any $t \geq 0$, one has $|\det \Phi_{-t}|_{F(x)}| \leq Ce^{-\lambda t}$.*

²Their results are stated for diffeomorphisms. The proof can be adapted to the case of vector field by a parallel way.

Λ is called a singular hyperbolic repeller if it is a singular hyperbolic attractor for $-X$.

One knows that the geometric Lorenz attractors as in [19, 20, 1] are singular hyperbolic attractors. $X \in \mathcal{X}^1(M^3)$ is called *singular Axiom A without cycle* as in [34] if the chain recurrent set of X contains only finitely many chain recurrent classes; moreover each chain recurrent class is a hyperbolic basic set, or a singular hyperbolic attractor, or a singular hyperbolic repeller.

Singular Axiom A flows include Lorenz-like flows. [5, 34] asked whether singular Axiom A flows and flows with a homoclinic tangency are typical phenomena.

Conjecture. *Every $X \in \mathcal{X}^1(M^3)$ can be accumulated either by vector fields with a homoclinic tangency, or by singular Axiom A vector fields without cycle.*

We get some progress on this conjecture.

Theorem C. *There is a dense G_δ set $\mathcal{R} \subset \mathcal{X}^1(M^3)$ such that for any $X \in \mathcal{R}$ and σ , if the chain recurrent class $C(\sigma)$ is nontrivial and admits a dominated splitting $T_{C(\sigma)}M^3 = E \oplus F$ w.r.t. the tangent flow, then $C(\sigma)$ is a singular hyperbolic attractor or a singular hyperbolic repeller.*

Notice that in a joint work with C. Bonatti [12], we proved this result by adding an additional assumption that: $C(\sigma)$ contains a periodic orbit. So, according to this result, to prove the above theorem, we assume that $C(\sigma)$ contains no periodic orbits. Then we can consider the return map of (singular) cross-sections. After a sequence of perturbations, we will get a contradiction. This is one of the main points of this work.

1.3 Entropy of flows

The entropy of a flow is defined to be the entropy of the time-one map of the flow. The definition meets some problems: there are two topological equivalent flows, one has zero entropy and the other one has positive entropy. This pathology happens because of the existence of singularities. See [40, 57, 56] for references. But using the main theorem of the paper, we can prove:

Theorem 1.2. *There is a dense open set $\mathcal{U} \subset \mathcal{X}^1(M^3)$ such that for any $X \in \mathcal{U}$, for any Y topological equivalent to X , one has $h(X) = 0$ iff $h(Y) = 0$.*

Proof. By Theorem A, there is a dense open set $\mathcal{U} \subset \mathcal{X}^1(M^3)$ such that for any $X \in \mathcal{U}$, either X is Morse-Smale, or X has a non-trivial hyperbolic basic set. Thus for any $X \in \mathcal{U}$, one has

- either $h(X) = 0$, then X is Morse-Smale, thus for any Y which is topological equivalent to X , for any point x , the forward iteration and backward iteration of x with respect to ϕ_t^Y go to a critical element. This feature implies that $h(Y) = 0$.
- or $h(X) > 0$, then X has a non-trivial hyperbolic basic set. Since for non-singular equivalent flows X, Y , $h(X) > 0$ iff $h(Y) > 0$. We have that $h(Y) > 0$.

□

For the relationship between zero-entropy vector fields and Morse-Smale vector fields, one has

Theorem 1.3. *If a three-dimensional vector field $X \in \mathcal{X}^1(M^3)$ can be accumulated by C^1 robustly zero-entropy vector fields, then it can be C^1 accumulated by Morse-Smale vector fields.*

Proof. By the assumptions, for any C^1 neighborhood \mathcal{U} of X , there is an open set $\mathcal{V} \subset \mathcal{U}$ such that every vector field Y in \mathcal{V} has zero-entropy. By Theorem A, by reducing \mathcal{V} if necessary, one can assume that for every $Y \in \mathcal{V}$, either it is Morse-Smale, or it has a non-trivial hyperbolic basic set. Since $Y \in \mathcal{V}$ can only have zero-entropy, one has Y is Morse-Smale. This ends the proof.

□

1.4 Organization of this paper

The proof of the theorems is not short. Especially for vector fields, they involve more definitions and notations.

1. In section 2, we give various kinds of definitions of flows associated to a vector field X . Liao defined these flows in a very abstract way. In fact, all these flows have their geometric meanings. We will deal with dominated splittings for the tangent flow and the linear Poincaré flow. For the estimations stated in this section (which is crucial for singular flow), Liao had very original ideas by a sequence of papers. We restate some of them and give the proof ourself. The proof is more intuitive.
2. In Section 3, we will study generic properties by connecting lemmas, ergodic closing lemma. [7] gave a C^1 connecting lemma for pseudo-orbits which helps us to obtain generic results for chain recurrent classes. We notice that Lyapunov stable chain recurrent classes with a critical element are robust for generic vector fields. The proof is not difficult, but it opens a new door: Lyapunov stable chain recurrent class will survive under generic small perturbations.
3. In Section 4 we first give the proof of Theorem A by assuming Theorem B. Then we reduce the proofs of Theorem B and Theorem C to several sub-results.
4. In Section 5, we prove that the Lyapunov stable chain recurrent class admits a partially hyperbolic splitting $E^{ss} \oplus E^{cu}$ and every singularity in the chain recurrent class is Lorenz-like. In this section, the main novelty of this paper is that we use some uniform estimation on vector fields away from homoclinic tangencies and a suitable application of Liao's shadowing lemma.
5. In Section 6, we prove that every nontrivial partially hyperbolic chain recurrent class with singularities contains periodic orbits for generic three-dimensional vector fields. We notice that it contains a periodic orbit iff it is singular hyperbolic. When the chain recurrent class is not singular hyperbolic, it is not singular hyperbolic robustly. Then by a sequence of perturbations, the continuation of the chain recurrent class intersects the closure of the basin of some sink. This implies that the continuation of the chain recurrent class is not Lyapunov stable. We can get a contradiction by Lemma 3.14. The difficulty we encounter is similar to the case of one-dimensional endomorphisms with singularities, where "singularities" means that the points where the endomorphisms fail to be a local diffeomorphisms. For flows, for every central unstable curve in the cross-section, in principle we will know that its length will grow near the local stable manifold of the singularities. But when it is cut by local stable manifold of singularities, its image under return map will be disconnected. This facts make the dynamics unclear. We have a good control in this section for this phenomenon.

2 Flows associated to a vector field and dominated splittings

2.1 Tangent flow, linear Poincaré flow and their extensions

Given $X \in \mathcal{X}^1(M^d)$, X generates a C^1 flow $\phi_t : M^d \rightarrow M^d$, and the *tangent flow* $\Phi_t = d\phi_t : TM^d \rightarrow TM^d$. Denote by $\pi : TM^d \rightarrow M^d$ the bundle projection.

Denote the normal bundle of X by

$$\mathcal{N} = \mathcal{N}^X = \bigcup_{x \in M^d \setminus \text{Sing}(X)} \mathcal{N}_x,$$

where \mathcal{N}_x is the orthogonal complement of the flow direction $X(x)$, i.e.,

$$\mathcal{N}_x = \{v \in TM^d : v \perp X(x)\}.$$

Given $x \in M^d \setminus \text{Sing}(X)$ and $v \in \mathcal{N}_x$, $\psi_t(v)$ is the orthogonal projection of $\Phi_t(v)$ on $\mathcal{N}_{\phi_t(x)}$ along the flow direction, i.e.,

$$\psi_t(v) = \Phi_t(v) - \frac{\langle \Phi_t(v), X(\phi_t(x)) \rangle}{|X(\phi_t(x))|^2} X(\phi_t(x)),$$

where $\langle \cdot, \cdot \rangle$ is the inner product on $T_x M$ given by the Riemannian metric.

By the definition, $\|\psi_t\|$ is uniformly bounded for t in any bounded interval although it is just defined on the regular set which is not compact in general.

This flow could also be defined in a more general way by Liao [29]. [24] used the terminology of ‘‘extended linear Poincaré flow’’. For every point $x \in M^d$, one could define the sphere fiber at x by

$$S_x M^d = \{v : v \in T_x M^d, |v| = 1\}.$$

The sphere bundle $SM^d = \bigcup_{x \in M^d} S_x M^d$ is compact. One can define the *unit tangent flow*

$$\Phi_t^I : SM^d \rightarrow SM^d$$

as

$$\Phi_t^I(v) = \frac{\Phi_t(v)}{|\Phi_t(v)|}$$

for any $v \in SM^d$.

Given a compact invariant set Λ of X , denote

$$\tilde{\Lambda} = \text{Closure} \left(\bigcup_{x \in \Lambda \setminus \text{Sing}(X)} \frac{X(x)}{|X(x)|} \right).$$

in SM^d . Thus the essential difference between $\tilde{\Lambda}$ and Λ is on the singularities. We have more information on $\tilde{\Lambda}$: it tells us how regular points in Λ accumulate singularities.

For any $x \in M^d$, and any two orthogonal vectors $v_1, v_2 \in T_x M^d$, if $|v_1| \neq 0$, one can define

$$\chi_t(v_1, v_2) = (\Phi_t(v_1), \Phi_t(v_2) - \frac{\langle \Phi_t(v_1), \Phi_t(v_2) \rangle}{|\Phi_t(v_1)|^2} \Phi_t(v_1)).$$

By definition the two components of χ_t are still orthogonal. If one denotes

$$\chi_t = (\text{proj}_1(\chi_t), \text{proj}_2(\chi_t)),$$

for any regular point $x \in M^d$ and any vector $v \in \mathcal{N}_x$, one has

$$\psi_t(v) = \text{proj}_2 \chi_t(X(x), v).$$

One can normalize the first component of χ_t : for any $x \in M^d$, and any two orthogonal vectors $v_1, v_2 \in T_x M^d$, if $|v_1| = 1$, one can define

$$\chi_t^\#(v_1, v_2) = (\Phi_t^I(v_1), \Phi_t(v_2) - \frac{\langle \Phi_t(v_1), \Phi_t(v_2) \rangle}{|\Phi_t(v_1)|^2} \Phi_t(v_1)).$$

$\chi_t^\#$ is also a continuous flow, and for any regular point x and any $v \in \mathcal{N}_x$, one has

$$\chi_t^\# \left(\frac{X(x)}{|X(x)|}, v \right) = (\Phi_t^I \left(\frac{X(x)}{|X(x)|} \right), \psi_t(v)).$$

By the continuity of χ_t , one can extend the definition of ψ_t “to singularities”: for any $u \in \tilde{\Lambda}$, one defines $\tilde{\mathcal{N}}_u = \{v \in T_{\pi(u)}M : \langle u, v \rangle = 0\}$. $\tilde{\mathcal{N}}$ is a $(d-1)$ -dimensional vector bundle on the base space $\tilde{\Lambda}$ (for a formal discussion, see [24]). For $u \in \tilde{\Lambda}$ and $v \in \tilde{\mathcal{N}}_u$, one can define $\tilde{\psi}_t(v) = \text{proj}_2 \chi_t(u, v)$. By the definition we know that $\text{proj}_2 \chi_t$ is a continuous flow defined on $\tilde{\Lambda}$. Thus, $\tilde{\psi}_t$ can be viewed as a compactification of ψ_t .

2.2 Scaled linear Poincaré flow ψ_t^* and Liao’s estimation

For our purpose, we need another flow $\psi_t^* : \mathcal{N} \rightarrow \mathcal{N}$ (called *scaled linear Poincaré flow*). Given $x \in M^d \setminus \text{Sing}(X)$, and $v \in \mathcal{N}_x$,

$$\psi_t^*(v) = \frac{|X(x)|}{|X(\phi_t(x))|} \psi_t(v) = \frac{\psi_t(v)}{\|\Phi_t|_{\langle X(x) \rangle}\|}.$$

In our case, this scaled linear Poincaré flow will help us to overcome some difficulties produced by singularities. It gives uniform estimations on some non-compact sets.

Lemma 2.1. *For any $\tau > 0$, there is $C_\tau > 0$ such that for any $t \in [-\tau, \tau]$,*

$$\|\psi_t^*\| \leq C_\tau,$$

where $\|\psi_t^*\| = \sup\{|\psi_t^*(v)| : v \in \mathcal{N} \text{ and } |v| = 1\}$.

Proof. For $t \in [-\tau, \tau]$, first we know that Φ_t is uniformly bounded from 0 and ∞ ; from the definition of the linear Poincaré flow, we know that ψ_t is uniformly bounded. Thus, $\psi_t^*(x) = \psi_t(x)/\|\Phi_t|_{\langle X(x) \rangle}\|$ is uniformly bounded. \square

For each $\beta > 0$, one can define the normal manifold $N_x(\beta)$ of x as the following:

$$N_x(\beta) = \exp_x(\mathcal{N}_x(\beta)),$$

where $\mathcal{N}_x(\beta) = \{v \in \mathcal{N}_x : |v| \leq \beta\}$. Take $\beta_* > 0$ small enough such that for any $x \in M$, \exp_x is a diffeomorphism from $\mathcal{N}_x(\beta_*)$ to its image $N_x(\beta_*)$.

To study the dynamics in a small neighborhood of a periodic orbit of a vector field, Poincaré defined the sectional return map of a cross section of a periodic point. By generalizing this idea to every regular point, one can define the sectional Poincaré map for any two points in the same regular orbit. For our convenience, we define the sectional Poincaré map in the normal bundle.

Given $T > 0$ and $x \in M^d \setminus \text{Sing}(X)$, the flow ϕ_t defines a local holonomy map $P_{x, \phi_T(x)}$ from $N_x(\beta_*)$ to $N_{\phi_T(x)}(\beta_*)$ in a small neighborhood of x . Hence its lift map in the normal bundle gives a map $\mathcal{P}_{x, \phi_T(x)} : U \rightarrow \mathcal{N}_{\phi_T(x)}(\beta_*)$, where U is a small neighborhood of x in $\mathcal{N}_x(\beta_*)$ and $\mathcal{P}_{x, \phi_T(x)} = \exp_{\phi_T(x)}^{-1} \circ P_{x, \phi_T(x)} \circ \exp_x$. Note that when $T' > T > 0$, the domain of $\mathcal{P}_{x, \phi_{T'}(x)}$ is contained in the domain of $\mathcal{P}_{x, \phi_T(x)}$. Usually the size of U depends on the orbit of x : if x is very close to a singularity, then U should be very small. But after scaling, we have the following uniform estimation for the relative size of U .

Lemma 2.2. *Given $X \in \mathcal{X}^1(M^d)$ and $T > 0$, there is $\beta_T > 0$ such that for any regular point x , $\mathcal{P}_{x, \phi_T(x)}$ is well defined on $\mathcal{N}_x(\beta_T|X(x)|)$.*

Proof. After taking an orthonormal basis $\{e_1, e_2, \dots, e_d\}$ of $T_x M$, we get a coordinate system:

$$\text{Exp}_x : \mathbb{R}^d \rightarrow M^d,$$

such that

$$\text{Exp}_x(z) = \exp_x\left(\sum_{i=1}^d z_i e_i\right).$$

In the coordinate, the flow generated by the vector field X satisfies the following differential equation:

$$\frac{dz}{dt} = \hat{X}(z),$$

where

$$\hat{X}(z) = D\text{Exp}_x^{-1} \circ X(\text{Exp}_x(z)).$$

Note that \hat{X} depends on $x \in M^d$.

Assume that β_* is small enough such that for any $x \in M^d$, $|z| \leq \beta_*$, and any two unit vectors v and w ,

$$0.999 < |D\text{Exp}_x(z)v| < 1.001, \quad |\angle(D\text{Exp}_x(z)v, D\text{Exp}_x(z)w) - \angle(v, w)| < 0.001.$$

We will prove the lemma in the local coordinate. Note that $\text{Exp}_x(0) = x$ and $|X(x)| = |\hat{X}(0)|$.

Denote by

$$K = \sup_{x \in M^d, |z| \leq \beta_*} \{|\hat{X}(z)|, \|D\hat{X}(z)\|\}.$$

Since $D\text{Exp}_x$ and $D\text{Exp}_x^{-1}$ are uniformly bounded with respect to x , we have

$$K < \infty.$$

Assume $\beta_0 < \min\{1, \beta_*\}/(1000K)$. For any regular point x , take $e_1 = X(x)/|X(x)|$. Then $\hat{X}(0) = (|X(x)|, 0, \dots, 0)$. For $|z| \leq \beta_0|X(x)|$, according to the mean-value theorem, there exists $\xi \in \mathbb{R}^d$, $|\xi| \leq \beta_0|X(x)|$,

$$|\hat{X}(z)| = |\hat{X}(0) + D\hat{X}(\xi)z| \geq |X(x)| - K\beta_0|X(x)| \geq 0.999|X(x)| > 0.$$

This implies that $N_x(\beta_0|X(x)|) \cap \text{Sing}(X) = \emptyset$.

Denote by $\hat{\phi}_t(z) = (\hat{\phi}_t^1, \dots, \hat{\phi}_t^d)$ the solution of $\hat{X}(z)$ such that $\hat{\phi}_0(z) = z$. Then for $0 \leq t \leq \beta_0$, if $|z| \leq \frac{\beta_0}{3}|X(x)|$ and $|\hat{\phi}_s(z)| \leq \beta_0|X(x)|$ for $s \in [0, t]$, we have

$$|\hat{\phi}_t(z)| = |z + \int_0^t \hat{X}(\hat{\phi}_s(z))ds| \leq (\frac{\beta_0}{3} + 1.001t)|X(x)|.$$

Let t be the time such that $|\hat{\phi}_s(z)| \leq \beta_0|X(x)|$ for $s \in [0, t)$ and $|\hat{\phi}_t(z)| = \beta_0|X(x)|$. From the above estimation, we have that

$$t \geq \frac{2}{3}\beta_0/1.001 > \frac{1}{2}\beta_0.$$

By reducing β_0 if necessary, for $|z| \leq \beta_0|X(x)|$,

$$\sup_{t \in (-\beta_0, \beta_0)} \frac{|\hat{X}(z)|}{|\hat{X}(\hat{\phi}_t(z))|} < \frac{1}{1000K}, \quad \sup_{t \in (-\beta_0, \beta_0)} \angle(\hat{X}(z), \hat{X}(\hat{\phi}_t(z))) < \frac{1}{1000K}$$

Since $e_1 = X(x)/|X(x)|$, $\mathcal{N}_x = \{z : z_1 = 0\}$, and $\hat{X}(0) = (|X(x)|, 0, \dots, 0)$, we have that for $|z| \leq \beta_0|X(x)|$, $\hat{X}_1(z) \in [0.999|X(x)|, 1.001|X(x)|]$.

Claim. For $0 < r < \beta_0$, for any $z = (0, y)$, $y \in \mathbb{R}^{d-1}$, $|y| \leq r|X(x)|/2$, for any $t \in [\beta_0/3, 2\beta_0/3]$, there exists a unique $\tau = \tau(t, y) \in (0, \beta_0]$ such that $\hat{\phi}_\tau(0, y) \in \hat{N}_{\hat{\phi}_t(x)}(r)$, where

$$\hat{N}_x(r) = \text{Exp}^{-1}(N_x(r)).$$

Proof of Claim. Since $|z| < 1000\beta_0 K$, $t \in [0, \beta_0]$ and β_0 is small, Exp_x is almost an isometry, and $N_{\phi_t(x)}$ is the graph of a map $f = f_t : \mathbb{R}^{d-1} \rightarrow \mathbb{R}$ with $|\frac{\partial f}{\partial y}| < 0.001$. Let $z = (0, y)$ with $|y| \leq \beta_0 |X(x)|$. Then

$$\text{proj}_1 \hat{\phi}_{\beta_0}(z) = \int_0^{\beta_0} \hat{X}_1(\hat{\phi}_t(z)) dt \geq 0.999 |X(x)| \beta_0.$$

This means that z and $\hat{\phi}_{\beta_0}(z)$ are on the different sides of the graph of f_t for $t \in [\beta_0/3, 2\beta_0/3]$, from which one can define $\tau(t, y)$. Notice that f is C^1 and X is C^1 , we have that $\tau(t, y)$ is C^1 w.r.t. t and y . \square

For any $T \geq \beta_0$, let $n = \lceil 3T/\beta_0 \rceil$ and a partition

$$0 = t_0 < t_1 < \dots < t_{n-1} < t_n = T,$$

such that $t_i = i\beta_0/3$, $i = 0, 1, \dots, n-1$. Then we have $t_n - t_{n-1} \in [\beta_0/3, 2\beta_0/3]$. Then we can define β_T inductively. \square

Lemma 2.3. *Let $X \in \mathcal{X}^1(M^d)$ and $T > 0$. By reducing $\beta_T > 0$ as in Lemma 2.2 if necessary, for any $x \in M^d \setminus \text{Sing}(X)$, for the sectional Poincaré map*

$$\mathcal{P}_{x, \phi_T(x)} : \mathcal{N}_x(\beta_T |X(x)|) \rightarrow \mathcal{N}_{\phi_T(x)}(\beta_*),$$

$DP_{x, \phi_T(x)}(y)$ is uniformly continuous in the following sense: for any $\epsilon > 0$ there exists $\delta \in (0, \beta_T]$ such that for any $x \in M^d \setminus \text{Sing}(X)$ and $y, y' \in \mathcal{N}_x(\beta_T |X(x)|)$, if $|y - y'| \leq \delta |X(x)|$, then

$$|DP_{x, \phi_T(x)}(y) - DP_{x, \phi_T(x)}(y')| < \epsilon.$$

(Note that $DP_{x, \phi_T(x)}(0) = \psi_T|_{\mathcal{N}_x}$.) And hence there exists $K_T > 0$ (independent of x) such that

$$|DP_{x, \phi_T(x)}| \leq K_T.$$

Proof. We will still use the notations and terminologies as in the proof of Lemma 2.2. We first assume that $T < \beta_0$. Notice that we are in a local Euclidean coordinate, $N_z = \mathcal{N}_z$ for each regular point z .

In the local coordinate, we assume that the vector field \hat{X} has the following form $\hat{X}(x, y) = (f(x, y), g(x, y))$, where $x \in \mathbb{R}^1$, $y \in \mathbb{R}^{d-1}$, $f : \mathbb{R}^d \rightarrow \mathbb{R}^1$ and $g : \mathbb{R}^d \rightarrow \mathbb{R}^{d-1}$ are continuously C^1 maps such that

- $\hat{X}(0, 0) = (f(0, 0), 0)$, where $f(0, 0) > 0$.
- for any (x, y) in the local coordinate, one has $f(x, y) \in (0.999f(0, 0), 1.001f(0, 0))$ and $|g(x, y)| < f(0, 0)/1000$.

The flow of \hat{X} satisfies the following differential equations:

$$\begin{aligned} \frac{dx}{dt} &= f(x, y), \\ \frac{dy}{dt} &= g(x, y). \end{aligned}$$

We assume that the solution of these differential equations is $(\hat{\varphi}_t(x, y), \hat{\psi}_t(x, y))$, where $\hat{\varphi}_t(x, y) : \mathbb{R}^1 \times \mathbb{R}^d \rightarrow \mathbb{R}^1$ and $\hat{\psi}_t(x, y) : \mathbb{R}^1 \times \mathbb{R}^d \rightarrow \mathbb{R}^{d-1}$.

Now we consider the expression of the sectional Poincaré map $\mathcal{P}_T : \mathcal{N}_x(\beta_0|X(x)|) \rightarrow \mathcal{N}_{\phi_T(x)}(\beta_*)$ in this local coordinate. The local coordinate of x is $(0, 0)$. Thus $\mathcal{N}_{(0,0)}(\beta_*) \subset \{0\} \times \mathbb{R}^{d-1}$. Now we consider $(\widehat{\varphi}_T(0, 0), \widehat{\psi}_T(0, 0))$, whose normal manifold is contained in a graph of an affine map $h : \mathbb{R}^{d-1} \rightarrow \mathbb{R}$ such that $|Dh| < 0.001$.

For $y \in \mathbb{R}^{d-1}$,

$$\mathcal{P}(0, y) = \mathcal{P}(y) = (\widehat{\varphi}_\tau(0, y), \widehat{\psi}_\tau(0, y))^T,$$

where the time function $\tau : \mathbb{R}^{d-1} \rightarrow \mathbb{R}^1$ satisfies

$$\widehat{\varphi}_{\tau(y)}(0, y) = h(\widehat{\psi}_{\tau(y)}(0, y)).$$

By differentiating y in the above equality, one has

$$\left. \frac{\partial \widehat{\varphi}}{\partial t} \right|_{t=\tau(y)} \frac{\partial \tau}{\partial y} + \frac{\partial \widehat{\varphi}}{\partial y} = \frac{\partial h}{\partial y} \left(\frac{\partial \widehat{\psi}}{\partial t} \frac{\partial \tau}{\partial y} + \frac{\partial \widehat{\psi}}{\partial y} \right).$$

Notice that in the above equality, $\partial \tau / \partial y$, $\partial \widehat{\varphi} / \partial y$ and $\partial h / \partial y$ are row vectors with $d-1$ elements, $\partial \widehat{\psi} / \partial t$ is a column vector with $(d-1)$ elements, $\partial \widehat{\psi} / \partial y$ is a $(d-1) \times (d-1)$ matrix.

By solving the above equality, one has

$$\frac{\partial \tau}{\partial y} = \frac{\frac{\partial h}{\partial y} \frac{\partial \widehat{\psi}}{\partial y} - \frac{\partial \widehat{\varphi}}{\partial y}}{\frac{\partial \widehat{\psi}}{\partial t} - \frac{\partial h}{\partial y} \frac{\partial \widehat{\psi}}{\partial t}}.$$

Thus,

$$\frac{\partial \mathcal{P}}{\partial y} = \begin{pmatrix} \frac{\partial \widehat{\varphi}}{\partial t} \frac{\partial \tau}{\partial y} + \frac{\partial \widehat{\varphi}}{\partial y} \\ \frac{\partial \widehat{\psi}}{\partial t} \frac{\partial \tau}{\partial y} + \frac{\partial \widehat{\psi}}{\partial y} \end{pmatrix} = \begin{pmatrix} \frac{\partial \widehat{\varphi}}{\partial t} \\ \frac{\partial \widehat{\psi}}{\partial t} \end{pmatrix} \frac{\frac{\partial h}{\partial y} \frac{\partial \widehat{\psi}}{\partial y} - \frac{\partial \widehat{\varphi}}{\partial y}}{\frac{\partial \widehat{\psi}}{\partial t} - \frac{\partial h}{\partial y} \frac{\partial \widehat{\psi}}{\partial t}} + \begin{pmatrix} \frac{\partial \widehat{\varphi}}{\partial y} \\ \frac{\partial \widehat{\psi}}{\partial y} \end{pmatrix}.$$

By the expression of the differential equations, one has

$$\frac{\partial \mathcal{P}}{\partial y} = \widehat{X}(\mathcal{P}(y)) \frac{\frac{\partial h}{\partial y} \frac{\partial \widehat{\psi}}{\partial y} - \frac{\partial \widehat{\varphi}}{\partial y}}{\frac{\partial \widehat{\psi}}{\partial t} - \frac{\partial h}{\partial y} \frac{\partial \widehat{\psi}}{\partial t}} + \begin{pmatrix} \frac{\partial \widehat{\varphi}}{\partial y} \\ \frac{\partial \widehat{\psi}}{\partial y} \end{pmatrix}$$

In another form,

$$\frac{\partial \mathcal{P}}{\partial y} = \frac{\widehat{X} \circ \mathcal{P}(y)}{\widehat{f} \circ \mathcal{P}(y) - \frac{\partial h}{\partial y} \widehat{g} \circ \mathcal{P}(y)} \left(\frac{\partial h}{\partial y} \frac{\partial \widehat{\psi}}{\partial y} - \frac{\partial \widehat{\varphi}}{\partial y} \right) + \begin{pmatrix} \frac{\partial \widehat{\varphi}}{\partial y}(0, y) \\ \frac{\partial \widehat{\psi}}{\partial y}(0, y) \end{pmatrix}.$$

Since $0.999 < |f|/|\widehat{X}| < 1.001$ and $|g|/|X| < 0.001$, there is a uniform constant $\widehat{K} > 0$ such that

$$\left\| \frac{\partial \mathcal{P}}{\partial y} \right\| \leq \widehat{K}.$$

For any $\varepsilon > 0$, since the tangent flow Φ_t is uniformly continuous, there is $\delta > 0$ such that for any $y, y' \in N_x(\beta_T|X(x)|)$, if $|y - y'| \leq \delta|X(x)|$, one has

$$\left\| \frac{\partial \widehat{\varphi}}{\partial y}(0, y) - \frac{\partial \widehat{\varphi}}{\partial y}(0, y') \right\| < \varepsilon/4, \quad \left\| \frac{\partial \widehat{\psi}}{\partial y}(0, y) - \frac{\partial \widehat{\psi}}{\partial y}(0, y') \right\| < \varepsilon/4.$$

Let $\alpha(y) = \widehat{X} \circ \mathcal{P}(y)$ and $\beta(y) = \widehat{f} \circ \mathcal{P}(y) - \partial h / \partial y (\widehat{g} \circ \mathcal{P}(y))$, then we have

$$\frac{\alpha(y)}{\beta(y)} - \frac{\alpha(y')}{\beta(y')} = \frac{\alpha(y) - \alpha(y')}{\beta(y)} + \frac{\beta(y') - \beta(y)}{\beta(y)\beta(y')} \alpha(y').$$

We have the following estimation by the mean value theorem:

$$\|\alpha(y) - \alpha(y')\| \leq \|D\widehat{X}\| \|\mathcal{P}(y) - \mathcal{P}(y')\| \leq \widehat{K} \|D\widehat{X}\| \|y - y'\|.$$

By reducing δ if necessary, for any $y, y' \in N_x(\beta_0|X(x)|)$, if $|y - y'| \leq \delta|X(x)|$, one has

$$\left\| \frac{\alpha(y) - \alpha(y')}{\beta(y)} \right\| \leq 1.001 \frac{\widehat{K} \|D\widehat{X}\| \delta |X(x)|}{|X(x)|} \leq 2\delta \widehat{K}.$$

We just need to choose $\delta < \varepsilon/8\widehat{K}$

$$\left\| \frac{\beta(y) - \beta(y')}{\beta(y)\beta(y')} \alpha(y') \right\| \leq \frac{\|\alpha(y')\|}{|\beta(y')|} \frac{\widehat{K} \|D\widehat{f} + \widehat{K} D\widehat{g}\| \|y - y'\|}{|\widehat{X}|} < \varepsilon/4,$$

if we reduce δ again.

Let

$$F(y) = \frac{\widehat{X} \circ \mathcal{P}(y)}{\widehat{f} \circ \mathcal{P} - \partial h / \partial y} \widehat{g} \circ \mathcal{P}, \quad G(y) = \frac{\partial h}{\partial y} \frac{\partial \widehat{\psi}}{\partial y} - \frac{\partial \widehat{\varphi}}{\partial y}.$$

We know that $F(y)$ is uniformly continuous and $G(y)$ is uniformly continuous. Thus $F(y)G(y)$ is uniformly continuous.

Combining all above estimations, we can know that $\mathcal{P}_{x, \phi_T(x)}$ is uniformly continuous.

For any $T \geq \beta_0$, let $n = \lceil 3T/\beta_0 \rceil$ and a partition

$$0 = t_0 < t_1 < \cdots < t_{n-1} < t_n = T,$$

such that $t_i = i\beta_0/3, i = 0, 1, \dots, n-1$. Then we have $t_n - t_{n-1} \in [\beta_0/3, 2\beta_0/3]$. Then by using the prolongation, we know the result is true. \square

Sometimes one needs to consider the *scaled sectional Poincaré map* \mathcal{P}^* which is defined in the following way:

$$\mathcal{P}_{x, \phi_T(x)}^*(y) = \frac{\mathcal{P}_{x, \phi_T(x)}(y|X(x)|)}{|X(\phi_T(x))|}$$

for each $y \in \mathcal{N}_x(\beta_T)$. Thus $\mathcal{P}_{x, \phi_T(x)}^*$ is a map from $\mathcal{N}_x(\beta_T)$ to $\mathcal{N}_{\phi_T(x)}$.

Lemma 2.4. *Given $X \in \mathcal{X}^1(M^d)$ and $T > 0$, there are constants $\beta_T > 0$ and $K_T > 0$ such that for any $t \in (0, T)$ and any regular point $x \in M^d$,*

1. $\mathcal{P}_{x, \phi_t(x)}^*$ can be defined on $\mathcal{N}_x(\beta_T)$.
2. $D\mathcal{P}_{x, \phi_t(x)}^*$ is uniformly continuous: for any $\varepsilon > 0$, there is $\delta > 0$, such that for any $y, z \in \mathcal{N}_x(\beta_T)$, if $d(y, z) < \delta$, one has $\|D\mathcal{P}_{x, \phi_t(x)}^*(y) - D\mathcal{P}_{x, \phi_t(x)}^*(z)\| < \varepsilon$.
3. $D_y \mathcal{P}_{x, \phi_T(x)}^*(y)|_{y=0} = \psi_T^*(x)$.
4. $\|D\mathcal{P}_{x, \phi_t(x)}^*(y)\| \leq K_T$ for any $y \in \mathcal{N}_x(\beta_T)$.

Proof. Item 1 is true because of Lemma 2.2.

For any $y, z \in \mathcal{N}_x(\beta_T)$, one has

$$D\mathcal{P}_{x, \phi_T(x)}^*(y) - D\mathcal{P}_{x, \phi_T(x)}^*(z) = \frac{|X(x)|}{|X(\phi_T(x))|} (D\mathcal{P}_{x, \phi_T(x)}(y|X(x)|) - D\mathcal{P}_{x, \phi_T(x)}(z|X(x)|)).$$

Since $|X(x)|/|X(\phi_T(x))|$ is uniformly bounded, item 2 follows from Lemma 2.3.

For item 3, we have

$$D_y \mathcal{P}_{x, \phi_T(x)}^*(y) = D_y \left(\frac{\mathcal{P}_{x, \phi_T(x)}(y|X(x))}{|X(\phi_T(x))|} \right) = D_y \mathcal{P}_{x, \phi_T(x)}(y|X(x)) \frac{|X(x)|}{|X(\phi_T(x))|}.$$

Thus, when $y = 0$, one has

$$D_y \mathcal{P}_{x, \phi_T(x)}^*(0) = D_y \mathcal{P}_{x, \phi_T(x)} \frac{|X(x)|}{|X(\phi_T(x))|} = \frac{\psi_T(x)}{\|\Phi_T|_{\langle X(x) \rangle}\|} = \psi_T^*(x).$$

Item 4 holds because that ψ_T^* is uniformly bounded and item 2. □

2.3 Franks' Lemma and dominated splittings

As in the diffeomorphism case, one needs Franks' lemma to get some information on the derivative along periodic orbits. We state a version of Franks' lemma for flows which is taken from [13, Theorem A.1]. Liao also had a version by using his standard differential equations [25, Proposition 3.4].

Lemma 2.5. *Given $X \in \mathcal{X}^1(M^d)$ and a C^1 neighborhood $\mathcal{U} \subset \mathcal{X}^1(M)$ of X , there is a neighborhood $\mathcal{V} \subset \mathcal{U}$ of X and $\varepsilon > 0$ such that for any $Y \in \mathcal{U}$, for any periodic orbit $\text{Orb}(x)$ of Y with period $T \geq 1$, any neighborhood U of $\text{Orb}(x)$ and any partition of $[0, T]$:*

$$0 = t_0 < t_1 < \cdots < t_l = T, \quad 1 \leq t_{i+1} - t_i \leq 2, i = 0, 1, \dots, l-1,$$

and any linear isomorphisms $L_i : \mathcal{N}_{\phi_{t_i}^Y(x)} \rightarrow \mathcal{N}_{\phi_{t_{i+1}}^Y(x)}$, $i = 0, 1, \dots, l-1$ with $\|L_i - \psi_{t_{i+1}-t_i}^Y|_{\mathcal{N}_{\phi_{t_i}^Y(x)}}\| \leq \varepsilon$, there exists $Z \in \mathcal{U}$ such that $\psi_{t_{i+1}-t_i}^Z|_{\mathcal{N}_{\phi_{t_i}^Z(x)}} = L_i$ and $Z = Y$ on $(M^d \setminus U) \cup \text{Orb}(x)$.

Remark. *For simplicity, the time length of the partition is restricted to $[1, 2]$. We remark that this is not a serious restriction. Sometimes, the system may contain periodic orbits with period less than 1. But in our consideration, usually singularities are all hyperbolic. So there is a lower bound for the periods of periodic orbits. And after scaling, we may assume that the lower bound is 1.*

In this paper, we will use two estimations obtained by Franks' lemma: dominated splittings for vector fields away from homoclinic tangencies and uniform estimation along a sequence of periodic orbits restricted on the stable bundle.

Dichotomy for the stable bundle: For any hyperbolic periodic orbit γ , \mathcal{N}_γ admits a natural splitting $\mathcal{N}_\gamma = \mathcal{N}^s \oplus \mathcal{N}^u$ such that \mathcal{N}^s is contracting and \mathcal{N}^u is expanding w.r.t. the linear Poincaré flow ψ_t .

By using the methods of periodic linear systems as in [9, 13, 31, 59], one can have the following dichotomy result for a hyperbolic periodic orbit with large period:

Lemma 2.6. *Given $X \in \mathcal{X}^1(M^d)$, for any C^1 neighborhood \mathcal{U} of X , there are $\eta_{\mathcal{U}} > 0$ and $\iota_{\mathcal{U}} > 0$ and a neighborhood $\mathcal{V} \subset \mathcal{U}$ of X such that for any hyperbolic periodic orbit γ of index i of $Y \in \mathcal{V}$ with $\tau(\gamma) > \iota_{\mathcal{U}}$, then*

- either, there is $Z \in \mathcal{U}$ such that γ is a hyperbolic periodic orbit of index $i-1$;
- or, for any $x \in \gamma$, for any time partition

$$0 = t_0 < t_1 < \cdots < t_n = \tau(\gamma),$$

verifying $t_{i+1} - t_i \geq \iota_{\mathcal{U}}$ for $0 \leq i \leq n-1$, one has

$$\prod_{i=0}^{n-1} \|\psi_{t_{i+1}-t_i}|_{\mathcal{N}^s(\phi_{t_i}(x))}\| \leq C \exp\{-\eta_{\mathcal{U}}\tau(\gamma)\}.$$

Definition 2.7. Let $C > 0$, $\eta > 0$ and $T > 0$. For a hyperbolic periodic orbit γ , for a ψ_t -invariant bundle $E \subset \mathcal{N}_{\gamma}$, we say that γ is called (C, η, T, E) -contracting at the period (w.r.t. ψ_t) if there is $m \in \mathbb{N}$, and for any $x \in \gamma$, there is some time partition

$$0 = t_0 < t_1 < \dots < t_n = m\tau(\gamma),$$

with $t_{i+1} - t_i \leq T$ for $0 \leq i \leq n-1$, such that

$$\prod_{i=0}^{n-1} \|\psi_{t_{i+1}-t_i}|_{E(\phi_{t_i}(x))}\| \leq C \exp\{-\eta m\tau(\gamma)\},$$

For an ψ_t -invariant subbundle $F \subset \mathcal{N}_{\gamma}$, we say that γ is called (C, η, T, F) -expanding at the period if it is (C, η, T, F) -contracting at the period for $-X$.

Remark. Since for a periodic orbit γ , for any $x \in \gamma$, one has $\Phi_{\tau(\gamma)}X(x) = X(\phi_{\tau(\gamma)}(x)) = X(x)$, one can give the above definition by using ψ_t^* .

Corollary 2.8. Given $X \in \mathcal{X}^1(M^d)$, we assume that Λ is a compact invariant set and not reduced to a critical element. If there is a sequence of vector fields $\{X_n\}$ such that

- $\lim_{n \rightarrow \infty} X_n = X$,
- Each X_n has a sink γ_n such that $\lim_{n \rightarrow \infty} \gamma_n = \Lambda$.

then one has the following dichotomy:

- either, there is a sequence of vector fields $\{Y_n\}$ such that $\lim_{n \rightarrow \infty} d_{C^1}(X_n, Y_n) = 0$, and γ_n is a hyperbolic periodic orbit of Y_n of index $d-1$.
- or, there are $\eta > 0$ and $T > 0$ such that for n large enough, γ_n is a $(1, \eta, T, \mathcal{N})$ -contracting at the period w.r.t. $\psi_t^{Y_n}$.

Proof. Note that $\lim_{n \rightarrow \infty} \tau(\gamma_n) = \infty$. If the ‘‘either’’ case is not true, then by Lemma 2.6, one can get constants $\eta_{\mathcal{U}}$ and $\iota_{\mathcal{U}}$. Then the ‘‘or’’ case is true by taking $\eta = \eta_{\mathcal{U}}$ and $T = 2\iota_{\mathcal{U}}$. □

Vector fields away from homoclinic tangencies: For any invariant set Λ (maybe not compact) without singularities, if there are $\iota > 0$ and an invariant splitting $\mathcal{N}_{\Lambda} = \mathcal{N}^{cs} \oplus \mathcal{N}^{cu}$ w.r.t. ψ_t satisfying $\|\psi_{\iota}|_{\mathcal{N}^{cs}(x)}\| \|\psi_{-\iota}|_{\mathcal{N}^{cu}(\phi_{\iota}(x))}\| \leq 1/2$ for any $x \in \Lambda$, then we say that Λ admits a ι -dominated splitting w.r.t. ψ_t . If $\dim \mathcal{N}^{cs}(x)$ is independent of x , then it is called the *index* of this dominated splitting. Note that for any linear flow defined on some linear bundle, one can define the notion of dominated splitting for that linear flow. Recall that

$$\mathcal{HT} = \{X \in \mathcal{X}^1(M^d) : X \text{ has a homoclinic tangency.}\}.$$

By the similar arguments as diffeomorphisms and by using Franks’ lemma for flows, from [59, 60], we have

Lemma 2.9. For any $X \in \mathcal{X}^1(M^d) \setminus \overline{\mathcal{HT}}$, there is a C^1 neighborhood \mathcal{U} and constants $C > 0$, $\lambda > 0$, $\delta > 0$ and $\iota > 0$ such that for any periodic orbit γ of $Y \in \mathcal{U}$ with period $\pi(\gamma) > \iota$.

- $\Phi_{\pi(\gamma)}$ has at most two exponents in $(-\delta, \delta)$, and $\Phi_{\pi(\gamma)}$ has at least one zero exponent, and this exponent corresponds to the flow direction.
- There is an invariant splitting $\mathcal{N}_\gamma = G^s \oplus G^c \oplus G^u$ with respect to the linear Poincaré flow ψ_t^Y , where G^s is the invariant space corresponding to the exponents less than $-\delta$, G^c is the invariant space corresponding to the exponents in $(-\delta, \delta)$ and the dimension of G^c is zero or one, G^u is the invariant space corresponding to the exponents larger than δ ; moreover for any $x \in \gamma$, for any time partition

$$0 = t_0 < t_1 < \dots < t_n = \tau(\gamma),$$

verifying $t_{i+1} - t_i \geq \iota$ for $0 \leq i \leq n-1$, one has

$$\prod_{i=0}^{n-1} \|\psi_{t_{i+1}-t_i}|_{G^s(\phi_{t_i}(x))}\| \leq C \exp\{-\lambda\tau(\gamma)\},$$

$$\prod_{i=0}^{n-1} \|\psi_{t_i-t_{i+1}}|_{G^u(\phi_{t_{i+1}}(x))}\| \leq C \exp\{-\lambda\tau(\gamma)\}.$$

- If γ is hyperbolic, and $\mathcal{N}_\gamma = \mathcal{N}^s \oplus \mathcal{N}^u$ is the hyperbolic splitting with respect to ψ_t^Y , then for any $x \in \gamma$ and $T > \iota$, one has

$$\|\psi_T^Y|_{\mathcal{N}^s(x)}\| \|\psi_{-T}^Y|_{\mathcal{N}^u(\phi_T^Y(x))}\| \leq \frac{1}{2}.$$

Corollary 2.10. Let $X \in \mathcal{X}^1(M^d) \setminus \overline{\mathcal{HT}}$. We assume that

- There is a sequence of vector fields $\{X_n\}$ such that $\lim_{n \rightarrow \infty} X_n = X$.
- Each X_n has a hyperbolic periodic orbit γ_n of index i such that $\Lambda = \lim_{n \rightarrow \infty} \gamma_n$ in the Hausdorff topology.

Then

- $\mathcal{N}_{\Lambda \setminus \text{Sing}(X)}$ admits a dominated splitting of index i with respect to the linear Poincaré flow ψ_t .
- $\mathcal{N}_{\Lambda \setminus \text{Sing}(X)}$ admits a dominated splitting of index i with respect to the scaled linear Poincaré flow ψ_t^* .
- If one considers $\tilde{\Lambda}$, then $\tilde{\mathcal{N}}_{\tilde{\Lambda}}$ admits a dominated splitting of index i with respect to the flow $\tilde{\psi}_t$.

Proof. Since $X \in \mathcal{X}^1(M^d) \setminus \overline{\mathcal{HT}}$, $X_n \rightarrow X$ and γ_n is a hyperbolic periodic orbit of X_n of index i ,

$$\mathcal{N}_{\gamma_n} = \mathcal{N}^s(\gamma_n) \oplus \mathcal{N}^u(\gamma_n)$$

is an ι -dominated splitting of index i w.r.t. $\psi_t^{X_n}$ for some uniform $\iota > 0$.

For each $x \in \Lambda \setminus \text{Sing}(X)$, by taking a subsequence if necessary, one can assume that there is $x_n \in \gamma_n$ such that $\lim_{n \rightarrow \infty} x_n = x$. After taking another subsequence, one can assume that $\mathcal{N}^{cs}(x) = \lim_{n \rightarrow \infty} \mathcal{N}^s(x_n)$ and $\mathcal{N}^{cu}(x) = \lim_{n \rightarrow \infty} \mathcal{N}^u(x_n)$.

Thus $\mathcal{N}_{\Lambda \setminus \text{Sing}(X)} = \mathcal{N}^{cs} \oplus \mathcal{N}^{cu}$ is an ι -dominated splitting of index i . One can see [24] for more details.

Since

$$\psi_t^*(x) = \frac{\psi_t(x)}{\|\Phi_t|_{\langle X(x) \rangle}\|},$$

any dominated splitting of ψ_t is also a dominated splitting of ψ_t^* .

The dominated splitting of the linear Poincaré flow can be extended to the closure of its representation in the sphere bundle. See [24] for more details. \square

Lemma 2.11. *For every $X \in \mathcal{X}^1(M^d) \setminus \overline{\mathcal{HT}}$, there are $\iota > 0$, $C > 0$, $\eta > 0$ and a C^1 neighborhood \mathcal{U} of X such that for any $Y \in \mathcal{U}$, if γ is a periodic sink of Y with period $\tau(\gamma) > \iota$, then*

- either, \mathcal{N}_γ admits an ι -dominated splitting of index $d - 2$ with respect to ψ_t^Y .
- or γ is $(C, \eta, 2\iota, \mathcal{N})$ -contracting at the period w.r.t. ψ_t^Y .

Proof. Let C and ι be as in Lemma 2.9. If the conclusion is not true, there exist $\eta_n \rightarrow 0$, $X_n \rightarrow X$ and a periodic sink γ_n of X_n with $\tau(\gamma_n) > \iota$, neither item 1 nor item 2 is satisfied. Then according to Franks' Lemma, after a small perturbation of X_n of size η_n , we get a Y_n such that γ_n is a periodic orbit of index $d - 2$. Since $Y_n \rightarrow X$, for n large enough, $\psi_t^{Y_n}$ has an ι -dominated splitting over γ_n of index $d - 2$, and then we get a dominated splitting for the extended linear Poincaré flow over the limit. But the limits of X_n and Y_n are the same since $X_n|_{\gamma_n} = Y_n|_{\gamma_n}$. By the continuity of dominated splitting of the extended linear Poincaré flow, we know that for n large enough, X_n has also an ι -dominated splitting over γ_n of index $d - 2$, which gives a contradiction. \square

2.4 Mixed dominated splittings: from linear Poincaré flow to tangent flow

For two linear normed spaces E and F , and a linear operator $A : E \rightarrow F$, the mini-norm $m(A)$ is defined by

$$m(A) = \inf_{v \in E, |v|=1} |Av|.$$

We use $L(E, F)$ to denote the space of bounded linear maps from E to F .

The following lemma concerns how we can get the dominated splitting of the tangent flow from the dominated splitting of the linear Poincaré flow. [24, Lemma 5.5, Lemma 5.6] used this kind of ideas. Here we give a general version.

Lemma 2.12. *Let $\widehat{\Lambda} \subset SM^d$ be a compact invariant set of Φ_t^I . Suppose*

- $\widetilde{\mathcal{N}}_{\widehat{\Lambda}} = \Delta^{cs} \oplus \Delta^{cu}$ is a dominated splitting w.r.t. $\widetilde{\psi}_t$.
- There are $C > 0$ and $\lambda > 0$ such that for any $u \in \widehat{\Lambda}$, for any $t > 0$, one has

$$\frac{\|\widetilde{\psi}_t|_{\Delta^{cs}(u)}\|}{\|\Phi_t(u)\|} \leq Ce^{-\lambda t}.$$

Then the projection $\pi(\widehat{\Lambda})$ admits a dominated splitting $T_{\pi(\widehat{\Lambda})}M^d = E \oplus F$ w.r.t the tangent flow Φ_t , where $\dim E = \dim \Delta^{cs}$.

Proof. For each point $u \in \tilde{\Lambda} \subset SM^d$, the direct-sum splitting $T_{\pi(u)}M^d = \Delta^{cs} \oplus \langle u \rangle \oplus \Delta^{cu}$ is continuous w.r.t. u . With respect to this decomposition, the tangent flow Φ_T has the following form:

$$\begin{pmatrix} \tilde{\psi}_T|_{\Delta^{cs}(u)} & 0 & 0 \\ B(u) & \Phi_T|_{\langle u \rangle} & C(u) \\ 0 & 0 & \tilde{\psi}_T|_{\Delta^{cu}(u)} \end{pmatrix}$$

By the definitions of χ_t and $\tilde{\psi}_t$ (see Subsection 2.1), one has that $F(u) = \langle u \rangle \oplus \Delta^{cu}(u)$ is an invariant sub-bundle of Φ_t . Let's find another invariant sub-bundle of Φ_t .

Claim. *There is $C_1 > 0$ and $\lambda_1 > 0$ such that for any $u \in \hat{\Lambda}$ and any $t \geq 0$, one has*

$$\frac{\|\tilde{\psi}_t|_{\Delta^{cs}(u)}\|}{m(\Phi_t|_{F(u)})} \leq C_1 e^{-\lambda_1 t}.$$

Proof of the claim. By enlarging T if necessary, one can assume that for any $u \in \hat{\Lambda} \subset SM^d$, one has

$$\frac{\|\tilde{\psi}_T|_{\Delta^{cs}(u)}\|}{\|\Phi_T|_{\langle u \rangle}\|} \leq \frac{1}{2}, \quad \frac{\|\tilde{\psi}_T|_{\Delta^{cs}(u)}\|}{m(\tilde{\psi}_T|_{\Delta^{cu}(u)})} \leq \frac{1}{2}.$$

Since Φ_T is bounded, by the continuity of the splitting, there is $K > 0$ such that $\|\Phi_T\| \leq K$ and $m(\Phi_T) \geq 1/K$. Denote by

$$D(u) = \begin{pmatrix} \Phi_T|_{\langle u \rangle} & C(u) \\ 0 & \tilde{\psi}_T|_{\Delta^{cu}(u)} \end{pmatrix}.$$

For any $n \in \mathbb{N}$, one has

$$\Phi_{-nT}|_{F(\Phi_{nT}^I(u))} = \prod_{i=0}^{n-1} D^{-1}(\Phi_{iT}^I(\Phi_{nT}^I(u))) = D^{-n}(\Phi_{nT}^I(u)).$$

Since

$$D^n(u) = \begin{pmatrix} \Phi_{nT}|_{\langle u \rangle} & \sum_{i=0}^{n-1} \Phi_{(n-1-i)T}|_{\langle \Phi_{(i+1)T}^I(u) \rangle} C(\Phi_{iT}^I(u)) \tilde{\psi}_{iT}|_{\Delta^{cu}(u)} \\ 0 & \psi_{nT}|_{\Delta^{cu}(u)} \end{pmatrix},$$

we have

$$D^{-n}(\Phi_{nT}^I(u)) = \begin{pmatrix} \Phi_{-nT}|_{\langle \Phi_{nT}^I(u) \rangle} & \sum_{i=0}^{n-1} \Phi_{(-1-i)T}|_{\langle \Phi_{(i+1)T}^I(u) \rangle} C(\Phi_{iT}^I(u)) \tilde{\psi}_{(i-n)T}|_{\Delta^{cu}(\Phi_{nT}^I(u))} \\ 0 & \tilde{\psi}_{-nT}|_{\Delta^{cu}(\Phi_{nT}^I(u))} \end{pmatrix}.$$

This implies

$$\begin{aligned}
\|D^{-n}|_{\Phi_{nT}^I(u)}\| &\leq \|\Phi_{-nT}|_{\langle\Phi_{nT}^I(u)\rangle}\| + \|\tilde{\psi}_{-nT}|_{\Delta^{cu}(\Phi_{nT}^I(u))}\| \\
&+ \left\| \sum_{i=0}^{n-1} \Phi_{(-1-i)T}|_{\langle\Phi_{(i+1)T}^I(u)\rangle} C(\Phi_{iT}^I(u)) \tilde{\psi}_{(i-n)T}|_{\Delta^{cu}(\Phi_{nT}^I(u))} \right\| \\
&\leq 2 \cdot 2^{-n} \frac{1}{\|\tilde{\psi}_{nT}|_{\Delta^{cs}(u)}\|} + K \sum_{i=0}^{n-1} \|\Phi_{(-1-i)T}|_{\langle\Phi_{(i+1)T}^I(u)\rangle}\| \|\tilde{\psi}_{(i-n)T}|_{\Delta^{cu}(\Phi_{nT}^I(u))}\| \\
&\leq \frac{2^{-n+1}}{\|\tilde{\psi}_{nT}|_{\Delta^{cs}(u)}\|} + K \sum_{i=0}^{n-1} \frac{2^{-(i+1)}}{\|\tilde{\psi}_{(i+1)T}|_{\Delta^{cs}(u)}\|} \frac{2^{i-n}}{\|\tilde{\psi}_{(n-i)T}|_{\Delta^{cs}(\Phi_{iT}^I(u))}\|} \\
&\leq \frac{2^{-n+1}}{\|\tilde{\psi}_{nT}|_{\Delta^{cs}(u)}\|} + K^2 \sum_{i=0}^{n-1} \frac{2^{-n-1}}{\|\tilde{\psi}_{iT}|_{\Delta^{cs}(u)}\| \|\tilde{\psi}_{(n-i)T}|_{\Delta^{cs}(\Phi_{iT}^I(u))}\|} \\
&\leq \frac{2^{-n+1}}{\|\tilde{\psi}_{nT}|_{\Delta^{cs}(u)}\|} + K^2 \frac{n2^{-n-1}}{\|\tilde{\psi}_{nT}|_{\Delta^{cs}(u)}\|}.
\end{aligned}$$

Thus, when n large enough, one has

$$\frac{\|\tilde{\psi}_{nT}|_{\Delta^{cs}(u)}\|}{m(\Phi_{nT}|_{F(u)})} \leq \frac{1}{2}.$$

This inequality implies the claim. \square

Now we will start to find a Φ_t -invariant bundle $E(u)$ and $T' > 0$ such that for any $u \in \widehat{\Lambda}$,

$$\frac{\|\Phi_{T'}|_{E(u)}\|}{m(\Phi_{T'}|_{F(u)})} \leq \frac{1}{2}.$$

By the claim above, there is $T_0 > 0$ such that

$$\frac{\|\tilde{\psi}_{T_0}|_{\Delta^{cs}(u)}\|}{m(\Phi_{T_0}|_{F(u)})} \leq \frac{1}{2}.$$

Let $L(\widehat{\Lambda}) = \prod_{u \in \widehat{\Lambda}} L(\Delta^{cs}(u), F(u))$. For each $\Pi \in L(\widehat{\Lambda})$, one can define the norm $\|\Pi\| = \sup_{u \in \widehat{\Lambda}} \|\Pi(u)\|$. Under this norm, one knows that $L(\widehat{\Lambda})$ is a Banach space. For any $\Pi \in L(\widehat{\Lambda})$, one has

$$\begin{pmatrix} \tilde{\psi}_{T_0}|_{\Delta^{cs}(u)} & 0 \\ B(u) & \Phi_{T_0}|_{F(u)} \end{pmatrix} \begin{pmatrix} \Delta^{cs}(u) \\ \Pi(u)\Delta^{cs}(u) \end{pmatrix} = \begin{pmatrix} \tilde{\psi}_{T_0}|_{\Delta^{cs}(u)}\Delta^{cs}(u) \\ B(u)\Delta^{cs}(u) + \Phi_{T_0}|_{F(u)}\Pi(u)\Delta^{cs}(u) \end{pmatrix}.$$

Thus, if we want to find an invariant bundle w.r.t. Φ_t , we need to require that

$$B(u)\Delta^{cs}(u) + \Phi_{T_0}|_{F(u)}\Pi(u)\Delta^{cs}(u) = \Pi(\Phi_{T_0}^I(u))\tilde{\psi}_{T_0}|_{\Delta^{cs}(u)}\Delta^{cs}(u).$$

In the spirit of the above equality, one can define a map $\mathcal{F} : L(\widehat{\Lambda}) \rightarrow L(\widehat{\Lambda})$ by the following form:

$$\mathcal{F}\Pi(u) = (\Phi_{-T_0}|_{F(\Phi_{T_0}^I(u))})(\Pi(\Phi_{T_0}^I(u))\tilde{\psi}_{T_0}|_{\Delta^{cs}(u)} - B(u)).$$

Given $\Pi_1, \Pi_2 \in L(\widehat{\Lambda})$, one has

$$\mathcal{F}\Pi_1 - \mathcal{F}\Pi_2 = (\Phi_{-T_0}|_{F(\Phi_{T_0}^I u)})(\Pi_1 - \Pi_2)(\widetilde{\psi}_{T_0}|_{\Delta^{cs}(u)}).$$

Thus,

$$\|\mathcal{F}\Pi_1 - \mathcal{F}\Pi_2\| \leq \|\Phi_{-T_0}|_{F(\Phi_{T_0}^I u)}\| \|\Pi_1 - \Pi_2\| \|\widetilde{\psi}_{T_0}|_{\Delta^{cs}(u)}\| \leq \frac{1}{2} \|\Pi_1 - \Pi_2\|.$$

So, \mathcal{F} is a contracting map. By contraction mapping principle, \mathcal{F} has a unique fixed point $\Pi \in L(\widehat{\Lambda})$, i.e.,

$$B(u)\Delta^{cs}(u) + \Phi_{T_0}|_{F(u)}\Pi(u)\Delta^{cs}(u) = \Pi(\Phi_{T_0}^I(u))\widetilde{\psi}_{T_0}|_{\Delta^{cs}(u)}\Delta^{cs}(u).$$

As a corollary, $E = (\text{id}, \Pi)\Delta^{cs}$ is an invariant bundle of Φ_{T_0} .

Since E, F and Δ^{cs} are continuous bundles w.r.t. $u \in \widehat{\Lambda}$, there is $L > 0$, which depends on the angles between each two bundles, such that for any non-zero vector $v_E \in E, v^{cs} \in \Delta^{cs}$, if $v_E = (\text{id}, \Pi)v^{cs}$, then

$$|v_E| \leq \frac{1}{L}|v^{cs}|.$$

Thus, for each n , one has

$$\|\Phi_{nT_0}|_{E(u)}\| \leq \frac{1}{L} \|\widetilde{\psi}_{nT_0}|_{\Delta^{cs}(u)}\| \leq \frac{1}{L} \frac{1}{2^n} m(\Phi_{nT_0}|_{F(u)}).$$

From these, we get a Φ_t -invariant splitting $E \oplus F$ over Λ , which satisfies the condition of the dominated splitting. \square

2.5 The existence of invariant manifolds

We assume that Λ is a compact invariant set and $\mathcal{N}_{\Lambda \setminus \text{Sing}(X)}$ admits a dominated splitting with respect to the linear Poincaré flow. If $\Lambda \cap \text{Sing}(X) = \emptyset$, then Λ will have plaque family ([22]) as in the case of diffeomorphisms. If $\Lambda \cap \text{Sing}(X) \neq \emptyset$, then Λ won't have uniform size of plaque family: the plaque family is defined on a non-compact set and the size is scaled by the norm of the vector field.

The scaled Poincaré sectional map \mathcal{P}^* could be defined in a uniform neighborhood of the zero section of \mathcal{N} . Moreover, we have uniform estimations on $D\mathcal{P}_{x, \phi_T(x)}^*(y)$ by Lemma 2.4. For getting plaque families of dominated splittings, one needs the following abstract lemma.

Lemma 2.13. *For any $d \in \mathbb{N}, L > 0, r > 0$, and $\alpha > 0$, there is $\gamma_0 > 0, \varepsilon_0 > 0$ such that: for any $\gamma \leq \gamma_0$, there exists $\delta > 0$, if a sequence of C^1 diffeomorphisms*

$$f_i : \mathbb{R}^d(r) \rightarrow \mathbb{R}^d, \quad i \in \mathbb{Z}$$

satisfy the following properties:

1. $f_i(0) = 0$,
2. $\sup_{i \in \mathbb{Z}} \max\{|Df_i(0)|, |Df_i^{-1}(0)|\} \leq L$
3. *There is a sequence of invariant decompositions $\mathbb{R}^d = E_i \oplus F_i$ with the following properties:*
 - $Df_i(0)(E_i) = E_{i+1}, Df_i(0)(F_i) = F_{i+1}$,
 - $\angle(E_i, F_i) > \alpha$,
 -

$$\frac{\|Df_i(0)|_{E_i}\|}{m(Df_i(0)|_{F_i})} \leq \frac{1}{2}.$$

4. $\text{Lip}(f_i - Df_i(0)) < \varepsilon_0$.

Then there are two sequences of embedding maps $\phi_i^{cs} \in \text{Emb}(E_i^{cs}(\gamma), F_i^{cu})$ and $\phi_i^{cu} \in \text{Emb}(E_i^{cu}(\gamma), F_i^{cs})$ such that

- $\phi_i^{cs/cu}(0) = 0$, $D\phi_i^{cs/cu}(0) = 0$,
- $f_i W_i^{cs/cu}(\delta) \subset W_{i+1}^{cs/cu}(\gamma)$, where $W_i^{cs}(\gamma)$ is the graph of ϕ_i^{cs} restricted to $E_i^{cs}(\gamma)$ and $W_i^{cu}(\gamma)$ is the graph of ϕ_i^{cu} restricted to $E_i^{cu}(\gamma)$.

Moreover, the invariant manifolds are continuous with respect to the sequence of $f = (f_i)$. Precisely, for two sequences $f = (f_i), g = (g_i)$, define their metric as

$$|f - g|_{C^1} = \sum_{i=-\infty}^{\infty} \frac{|f_i - g_i|_{C^1}}{2^{|i|}}.$$

Then both $W_i^{cs}(\gamma, f)$ and $W_i^{cu}(\gamma, f)$ are continuous with respect to f , i.e., for every $i \in \mathbb{Z}$, if $f^{(n)} \rightarrow f$, $x_n \in W_i^{cs}(\gamma, f^{(n)})$, $x_n \rightarrow x$, then $x \in W_i^{cs}(\gamma, f)$, and $T_{x_n} W_i^{cs}(\gamma, f^{(n)}) \rightarrow T_x W_i^{cs}(\gamma, f)$.

The proof of Lemma 2.13 needs to adapt the argument of [22, Theorem 5.5]. We omit the proof here.

For diffeomorphisms, we know that plaque family of compact invariant set with dominated splittings exists. For vector fields, if a compact invariant singular set has a dominated splitting w.r.t. the linear Poincaré flow, we also have some similar results, but the form is changed: one should modify the size of the manifolds. Recall that $P_{x, \phi_t(x)}$ is the sectional Poincaré map between N_x and $N_{\phi_t(x)}$.

Lemma 2.14. *Let Λ be a compact invariant set of $X \in \mathcal{X}^1(M^d)$. We assume that $\Lambda \setminus \text{Sing}(X)$ admits a dominated splitting $\mathcal{N}_{\Lambda \setminus \text{Sing}(X)} = \Delta^{cs} \oplus \Delta^{cu}$ of index i with respect to the linear Poincaré flow ψ_t . Let $T > 0$. There are continuous maps: $\eta^{cs} : \Lambda \setminus \text{Sing}(X) \rightarrow \text{Emb}^1(B^i, TM^d)$ and $\eta^{cu} : \Lambda \setminus \text{Sing}(X) \rightarrow \text{Emb}^1(B^{d-1-i}, TM^d)$ verifying the following properties:*

1. $\eta^{cs}(x)(B^i(1)) \subset \mathcal{N}_x$ and $\eta^{cu}(x)(B^{d-2-i}(1)) \subset \mathcal{N}_x$ for any $x \in \Lambda \setminus \text{Sing}(X)$.
2. For any $\xi > 0$, we define two sub-manifolds by $W_\xi^{cs}(x) = \exp_x(\eta^{cs}(x)(B^i(\xi)))$ and $W_\xi^{cu}(x) = \exp_x(\eta^{cu}(x)(B^{d-2-i}(\xi)))$, then one has
 - $T_x W_1^{cs}(x) = \Delta^{cs}(x)$ and $T_x W_1^{cu}(x) = \Delta^{cu}(x)$.
 - For any $\varepsilon > 0$, there is $\delta > 0$ such that for any regular point $x \in \Lambda$ and any $t \in [0, T]$, one has $P_{x, \phi_t(x)}(W_{\delta|X(x)|}^{cs}) \subset W_{\varepsilon|X(\phi_t(x))|}^{cs}(\phi_t(x))$ and $P_{x, \phi_t(x)}(W_{\delta|X(x)|}^{cu}) \subset W_{\varepsilon|X(\phi_t(x))|}^{cu}(\phi_t(x))$.

Proof. We will mainly use Lemma 2.13 to prove this lemma. For each point x , \mathcal{N}_x is isomorphic to \mathbb{R}^{d-1} . Since $\Lambda \setminus \text{Sing}(X)$ admits a dominated splitting of index i w.r.t. the linear Poincaré flow, there is $T > 0$ such that

$$\frac{\|\psi_T^*|_{\Delta^{cs}(x)}\|}{m(\psi_T^*|_{\Delta^{cu}(x)})} \leq \frac{1}{2}, \quad \forall x \in \Lambda \setminus \text{Sing}(X).$$

For each $i \in \mathbb{Z}$, one take $f_i = \mathcal{P}_{\phi_{T^i}(x), \phi_{(T+1)^i}(x)}^*$ on $\mathcal{N}_{\phi_{T^i}(x)}$. By Lemma 2.4, all assumptions of Lemma 2.13 are satisfied. Then by Lemma 2.13, we get the existence of plaque family. \square

$W^{cs}(x)$ and $W^{cu}(x)$ are called *central stable plaques* and *central unstable plaques* respectively.

Corollary 2.15. *Let Λ be a compact invariant set of $X \in \mathcal{X}^1(M^d)$. We assume that $\Lambda \setminus \text{Sing}(X)$ admits a dominated splitting $\mathcal{N}_{\Lambda \setminus \text{Sing}(X)} = \Delta^{cs} \oplus \Delta^{cu}$ of index i with respect to the linear Poincaré flow ψ_t . Let $T > 0$. Then for any $\varepsilon > 0$, there is $\delta > 0$ such that for any $x, y \in \Lambda$, if $d(x, y) < \delta$, $d(x, \text{Sing}(X)) > \varepsilon$, $d(y, \text{Sing}(X)) > \varepsilon$, then $W^{cs}(x) \cap \phi_{[-T, T]}(W^{cu}(y)) \neq \emptyset$.*

The proof of this corollary bases on the uniform continuity of plaque families when points are far away from singularities.

2.6 Estimations on the size of stable/unstable manifolds

Definition 2.16. *Let Λ be an invariant set and $E \subset \mathcal{N}_{\Lambda \setminus \text{Sing}(X)}$ an invariant subbundle of the linear Poincaré flow ψ_t . For $C > 0$, $\eta > 0$ and $T > 0$, $x \in \Lambda \setminus \text{Sing}(X)$ is called (C, η, T, E) - ψ_t^* -contracting if for any partition of times $0 = t_0 < t_1 < \dots < t_n < \dots$ verifying*

- $t_{n+1} - t_n \geq T$ for any $n \in \mathbb{N}$ and $t_n \rightarrow \infty$ as $n \rightarrow \infty$,
- For any $n \in \mathbb{N}$,

$$\prod_{i=0}^{n-1} \|\psi_{t_{i+1}-t_i}^*|_{E(\phi_{t_i}(x))}\| \leq Ce^{-\eta t_n}.$$

$x \in \Lambda \setminus \text{Sing}(X)$ is called (C, η, T, E) - ψ_t^* -expanding if it's (C, η, T, E) - ψ_t^* -contracting for $-X$.

An increasing homeomorphism $\theta : \mathbb{R} \rightarrow \mathbb{R}$ is called a *reparametrization* if $\theta(0) = 0$. we meet the reparametrization problem for flows.

For any orbit $\text{Orb}(x)$, one defines

$$W^s(\text{Orb}(x)) = \{y \in M^d, \exists \text{ a reparametrization } \theta \text{ s.t., } \lim_{t \rightarrow \infty} d(\phi_{\theta(t)}(y), \phi_t(x)) = 0\},$$

$$W^s(\text{Orb}(x)) = \{y \in M^d, \exists \text{ a reparametrization } \theta \text{ s.t., } \lim_{t \rightarrow -\infty} d(\phi_{\theta(t)}(y), \phi_t(x)) = 0\}.$$

Lemma 2.17. *Let Λ be a compact invariant set of $X \in \mathcal{X}^1(M^d)$. Assume that $\Lambda \setminus \text{Sing}(X)$ admits a dominated splitting $\mathcal{N}_{\Lambda \setminus \text{Sing}(X)} = \Delta^{cs} \oplus \Delta^{cu}$ of index i with respect to the linear Poincaré flow ψ_t . For $C > 0$, $\eta > 0$ and $T > 0$, there is $\delta > 0$ such that*

- For any regular point $x \in \Lambda$, if x is $(C, \eta, T, \Delta^{cs})$ - ψ_t^* -contracting, then $W_{\delta|X(x)|}^{cs} \subset W^s(\text{Orb}(x))$;
- For any regular point $x \in \Lambda$, if x is $(C, \eta, T, \Delta^{cu})$ - ψ_t^* -expanding, then $W_{\delta|X(x)|}^{cu} \subset W^u(\text{Orb}(x))$.

Proof. We need to prove that there is $\delta > 0$ such that

$$\lim_{t \rightarrow +\infty} \text{diam} \left(\mathcal{P}_{x, \phi_t(x)}^* (\eta^{cs}(x)(B^i(\delta))) \right) = 0.$$

By the uniform continuity of $D\mathcal{P}_{x, \phi_t(x)}^*$ in Lemma 2.4, we have uniform linearized neighborhood of 0 in \mathcal{N}_z for each regular point z . Then the proof parallels to [52, Corollary 3.3]. \square

Corollary 2.18. *Under the assumption of Lemma 2.17 for any compact set $\Lambda_0 \subset \Lambda \setminus \text{Sing}(X)$, there is $\varepsilon > 0$ such that for any $x, y \in \Lambda_0$, if*

- $d(x, y) < \varepsilon$;
- x is $(C, \eta, T, \Delta^{cs})$ - ψ_t^* -contracting and y is $(C, \eta, T, \Delta^{cu})$ - ψ_t^* -expanding;

then $W^s(\text{Orb}(x)) \cap W^u(\text{Orb}(y)) \neq \emptyset$.

Similar to the proof of Lemma 2.17, we have

Lemma 2.19. *Let $X \in \mathcal{X}^1(M^d)$. For any $C > 0$, $\eta > 0$ and $T > 0$, there is $\delta = \delta(X, C, \eta, T) > 0$ such that if a regular point $x \in M^d$ is $(C, \eta, T, \mathcal{N})$ - ψ^* -contracting, then*

$$\lim_{t \rightarrow +\infty} \text{diam} (P_{x, \phi_t(x)}(N_x(\delta|X(x)|))) = 0.$$

In other words, $N_{x, \delta|X(x)|} \subset W^s(\text{Orb}(x))$.

Notice that not only x may be close to a singularity, but also $\omega(x)$ may contain singularities.

Theorem 2.20. *[29, Theorem 4.1, Proposition 6.2] Given $X \in \mathcal{X}^1(M^d)$ and a hyperbolic singularity σ of X , for any $C > 0$, $\eta > 0$ and $T > 1$, there exists a neighborhood U of σ such that there is no $(C, \eta, T, \mathcal{N})$ - ψ^* -contracting periodic point in U .*

The proof of Theorem 2.20 is not short and it is contained in [29]. Now we give some idea of Theorem 2.20: if Theorem 2.20 is not true, then there is a sequence of $(C, \eta, T, \mathcal{N})$ - ψ^* -contracting periodic points $\{p_n\}$ such that $\lim_{n \rightarrow \infty} p_n = \sigma$. This holds only if σ is a saddle. By the property of $\{p_n\}$, we have

- $T_\sigma M^d$ admits a dominated splitting $T_\sigma M^d = E^{cs} \oplus E^{uu}$ w.r.t. the tangent flow, where $\dim E^{uu} = 1$ and E^{uu} is strong unstable.
- There is $\delta > 0$, $N_{p_n}(\delta|X(p_n)|) \subset W^s(p_n)$.

Then by a careful estimation (which is not obvious), one has for n large enough, $W^{uu}(\sigma) \cap W^s(p_n) \neq \emptyset$, where $W^{uu}(\sigma)$ is the strong unstable manifold corresponding to E^{uu} . But $W^{uu}(\sigma) \setminus \{\sigma\}$ contains only two orbits, and p_n are distinct periodic orbits. This gives us a contradiction.

2.7 Pliss Lemma

We use the following lemma of Pliss type to get the points which can have uniform estimations to infinity.

Lemma 2.21. *Given $X \in \mathcal{X}^1(M^d)$, $C > 0$, $T > 0$ and $\eta > 0$, for any $\eta' \in (0, \eta)$, there is $N = N(C, T, \eta, \eta') > 0$, such that if γ is a periodic orbit with period $\tau(\gamma) > N$, $E \subset \mathcal{N}_\gamma$ is an invariant bundle w.r.t. ψ_t , and if γ is (C, η, T, E) -contracting at the period w.r.t. ψ_t , then there is $x \in \gamma$ such that x is $(1, \eta', T, E)$ - ψ_t^* -contracting.*

Proof. Since γ is a (C, η, T, E) -contracting at the period w.r.t. ψ_t , there is $m \in \mathbb{N}$ and a time partition

$$0 = t_0 < t_1 < \dots < t_n = m\tau(\gamma),$$

with $t_{i+1} - t_i \leq T$ for $0 \leq i \leq n - 1$, such that

$$\prod_{i=0}^{n-1} \|\psi_{t_{i+1}-t_i}|_{E(\phi_{t_i}(x))}\| \leq C \exp\{-\eta m\tau(\gamma)\}.$$

Since $\Phi_{\tau(\gamma)}(X(x)) = X(\phi_{\tau(\gamma)}(x)) = X(x)$, the above estimation is also true for ψ_t^* :

$$\prod_{i=0}^{n-1} \|\psi_{t_{i+1}-t_i}^*|_{E(\phi_{t_i}(x))}\| \leq C \exp\{-\eta m\tau(\gamma)\}.$$

When $\eta' < \eta$, if $\tau(\gamma)$ is large enough, one can cancel the constant C . Following [17, Lemma 2.14], one can get the $(1, \eta', T, E)$ - ψ_t^* -contracting point x . \square

Lemma 2.22. *Let $X \in \mathcal{X}^1(M^d)$ be Kupka-Smale. For any $C > 0$, $T > 0$ and $\eta > 0$, X can only have finitely many $(C, \eta, T, \mathcal{N})$ -contracting periodic orbits.*

Proof. If the conclusion is not true, then X has infinitely many distinct periodic orbits $\{\gamma_n\}$ such that

- $\lim_{n \rightarrow \infty} \tau(\gamma_n) = \infty$.
- Each γ_n is $(C, \eta, T, \mathcal{N})$ -contracting at the period w.r.t. ψ_t .

By Lemma 2.21, for each n large enough, there is $x_n \in \gamma_n$ such that x_n is a $(1, \eta/2, T, \mathcal{N})$ - ψ_t^* -contracting point. Thus, there is $\delta = \delta(X, \eta, T) > 0$ such that $N_{x_n}(\delta|X(x_n)|)$ is contained in the stable manifold of x_n . If x_n accumulates on singularities, then one can get a contradiction by Theorem 2.20. If x_n does not accumulate on singularities, then the basin of γ_n covers an open set with uniform size. This also gives a contradiction because the volume of M^d is finite and $\{\gamma_n\}$ are distinct periodic orbits. \square

3 Chain recurrence and genericity

3.1 Conley theory

A chain recurrent class is called *non-trivial* if it is not reduced to a critical element; otherwise, it is called *trivial*. For each hyperbolic critical element p of X , since $\text{Orb}(p)$ has a well-defined continuation $\text{Orb}(p_Y)$ for Y close to X , $C(p)$ also has a well-defined continuation $C(p_Y, Y)$.

A compact invariant set Λ of X (if it has a continuation) is called *lower semi-continuous* if for any sequence of vector fields $\{X_n\}$ verifying $\lim_{n \rightarrow \infty} X_n = X$, one has $\liminf_{n \rightarrow \infty} \Lambda_{X_n} \supset \Lambda$. A compact invariant set Λ of X (if it has a continuation) is called *upper semi-continuous* if for any sequence of vector fields $\{X_n\}$ verifying $\lim_{n \rightarrow \infty} X_n = X$, one has $\limsup_{n \rightarrow \infty} \Lambda_{X_n} \subset \Lambda$. It is well known that the closure of hyperbolic periodic orbits is lower semi-continuous and the chain recurrent set is upper semi-continuous. There is a classical result saying that lower semi-continuous sets and upper semi-continuous sets are continuous for generic vector fields.

Lemma 3.1. *For a hyperbolic critical element p , $C(p)$ is upper semi-continuous. As a corollary, if p_1 and p_2 are two critical elements of X with the property $C(p_1) \cap C(p_2) = \emptyset$, then there is a neighborhood \mathcal{U} of X such that for any $Y \in \mathcal{U}$, one has $C(p_{1,Y}) \cap C(p_{2,Y}) = \emptyset$.*

Proof. The fact that $C(p)$ is upper semi-continuous because of the continuity of the flows with respect to the vector fields.

By [14], if we have two chain recurrent classes $C(p_1)$ and $C(p_2)$ satisfying $C(p_1) \cap C(p_2) = \emptyset$, then there is an open set U such that

- $\phi_t(\bar{U}) \subset U$ for $t > 0$;
- $C(p_1) \subset U$ and $C(p_2) \subset \text{Int}(M \setminus \bar{U})$ or $C(p_2) \subset U$ and $C(p_1) \subset \text{Int}(M \setminus \bar{U})$

Then by the continuity of vector fields, there is a C^1 neighborhood \mathcal{U} of X such that for any $Y \in \mathcal{U}$, one has $C(p_{1,Y}) \subset U$ and $C(p_{2,Y}) \subset \text{Int}(M \setminus \bar{U})$ or $C(p_{2,Y}) \subset U$ and $C(p_{1,Y}) \subset \text{Int}(M \setminus \bar{U})$. As a corollary, one has $C(p_{1,Y}) \cap C(p_{2,Y}) = \emptyset$. \square

For each point $x \in M^d$, one can define the strong stable manifold $W^{ss}(x)$ and the strong unstable manifold $W^{uu}(x)$ as

$$W^{ss}(x) \triangleq \{y \in M^d : \lim_{t \rightarrow \infty} d(\phi_t(x), \phi_t(y)) = 0\},$$

$$W^{uu}(x) \triangleq \{y \in M^d : \lim_{t \rightarrow -\infty} d(\phi_t(x), \phi_t(y)) = 0\}.$$

But for flows, this definition is not enough in many cases. By the difference with diffeomorphisms, sometimes we need to reparameterize the time variable. This leads us to give the definition of $W^s(\text{Orb}(x))$ and $W^u(\text{Orb}(x))$ as in Section 2.

By the definitions, one has that $W^s(\text{Orb}(x))$ and $W^u(\text{Orb}(x))$ are invariant sets. The proof of the following lemma is fundamental, hence omitted.

Lemma 3.2. *For any hyperbolic critical point p , one has*

- *For any critical element p , one has $W^s(\text{Orb}(p)) = \cup_{t \geq 0} \phi_t(W^{ss}(p))$ and $W^u(\text{Orb}(p)) = \cup_{t \geq 0} \phi_t(W^{uu}(p))$.*
- *If $C(p)$ is non-trivial, then $C(p) \cap W^s(\text{Orb}(p)) \setminus \text{Orb}(p) \neq \emptyset$ and $C(p) \cap W^u(\text{Orb}(p)) \setminus \text{Orb}(p) \neq \emptyset$.*

For a compact invariant set Λ , one says that Λ is *Lyapunov stable* for X if for any neighborhood U of Λ , there is a neighborhood V of Λ such that $\phi_t(V) \subset U$ for any $t > 0$.

Lemma 3.3. *If Λ is Lyapunov stable, then $W^u(\text{Orb}(x)) \subset \Lambda$ for each $x \in \Lambda$.*

Proof. Given any $y \in W^u(\text{Orb}(x))$, for any neighborhood U of Λ , since Λ is Lyapunov stable, there is a neighborhood V of Λ such that $\phi_t(V) \subset U$ for any $t \geq 0$. For any $y \in W^u(\text{Orb}(x))$, there is an increasing homeomorphism $\theta : \mathbb{R} \rightarrow \mathbb{R}$ such that $d(\phi_t(x), \phi_{\theta(t)}(y)) \rightarrow 0$ as $t \rightarrow -\infty$. Thus there is $t_V > 0$ such that $\theta(-t_V) < 0$ and $\phi_{\theta(-t_V)} \in V$. By the Lyapunov stability one has $y = \phi_{-\theta(-t_V)}(\phi_{\theta(-t_V)}(y)) \in U$. By the arbitrary property of U , one has $y \in \Lambda$. \square

By the definition, if Λ is not Lyapunov stable, then there is a neighborhood U_0 of Λ such that there is a sequence of neighborhoods $\{V_n\}$ such that

- $\lim_{n \rightarrow \infty} \overline{V_n} = \Lambda$.
- $\phi_{t_n}(V_n) \not\subset U$ for some $t_n > 0$.

As a corollary, $\lim_{n \rightarrow \infty} t_n = \infty$ since Λ is an invariant set. Hence, if Λ is not Lyapunov stable, then there are $\{t_n\} \subset \mathbb{R}$ and a sequence of points $\{x_n\}$ such that

- $\lim_{n \rightarrow \infty} x_n \in \Lambda$.
- $\lim_{n \rightarrow \infty} t_n = \infty$.
- $\lim_{n \rightarrow \infty} \phi_{t_n}(x_n)$ exists and $\lim_{n \rightarrow \infty} \phi_{t_n}(x_n) \notin \Lambda$.

For $x \in M^d$, one can define the *chain unstable set* $W^{ch,u}(x)$ of x and the *chain stable set* $W^{ch,s}(x)$ of x in the following way:

$$\begin{aligned} W^{ch,u}(x) &= \{y \in M^d : \forall \varepsilon > 0, \exists \text{ an } \varepsilon\text{-pseudo orbit } \{x_i\}_{i=0}^n \text{ s.t. } x_0 = x, x_n = y\}, \\ W^{ch,s}(x) &= \{y \in M^d : \forall \varepsilon > 0, \exists \text{ an } \varepsilon\text{-pseudo orbit } \{x_i\}_{i=0}^n \text{ s.t. } x_0 = y, x_n = x\}. \end{aligned}$$

y is *chain attainable* from x iff $y \in W^{ch,u}(x)$. For $x, y \in M^d$, one says that they are *chain bi-attainable* if $y \in W^{ch,u}(x) \cap W^{ch,s}(x)$.

3.2 The C^1 connecting lemmas and ergodic closing lemma for flows

[3, 61] gave the following extension of Hayashi's C^1 connecting lemma [21]. We will use it in Section 6.

Lemma 3.4. *For any vector field $X \in \mathcal{X}^1(M^d)$, for any point $z \notin \text{Per}(X) \cup \text{Sing}(X)$, for any $\varepsilon > 0$, there are $L > 0$ and two neighborhoods $\widetilde{W}_z \subset W_z$ of z such that*

- *one can choose W_z and \widetilde{W}_z to be arbitrarily small neighborhoods of z ,*
- *for any p and q in M^d , if the positive orbit of p and the negative orbit of q enter \widetilde{W}_z , but the orbit segments $\{\phi_t(p) : 0 \leq t \leq L\}$ and $\{\phi_t(q) : -L \leq t \leq 0\}$ don't intersect W_z ,*

then there is Y ε - C^1 -close to X such that

- *q is in the positive orbit of p with respect to the flow ϕ_t^Y generated by Y .*
- *$Y(x) = X(x)$ for any $x \in M^d \setminus W_{L,z}$, where $W_{L,z} = \bigcup_{0 \leq t \leq L} \phi_t^X(W_z)$.*

Mañé's ergodic closing lemma [31] is also useful in this paper. First we state a flow version of Mañé's ergodic closing lemma taken from [58].

Definition 3.5. *Let $X \in \mathcal{X}^1(M^d)$. A regular point $x \in M^d$ is called strongly closable if for any C^1 neighborhood \mathcal{U} of X , and any $\varepsilon > 0$, there are $Y \in \mathcal{U}$ and a periodic point $y \in M^d$ of Y with period $\tau(y)$ such that*

- *$X(z) = Y(z)$ for any $x \in M^d \setminus \bigcup_{t \in \mathbb{R}} \phi_t(B(x, \varepsilon))$,*
- *$d(\phi_t^X(x), \phi_t^Y(y)) < \varepsilon$ for each $t \in [0, \tau(y)]$.*

Denote by $\Sigma(X)$ the set of strongly closable points of X .

The ergodic closing lemma [31, 58] states:

Lemma 3.6. *$\mu(\Sigma(X) \cup \text{Sing}(X)) = 1$ for every $T > 0$ and every ϕ_T^X -invariant probability Borel measure μ .*

One needs the following corollary which asserts that one can get a periodic orbit with some additional properties and preserve a compact invariant subset in a transitive set by small perturbations simultaneously. For the applications in this paper, one takes the compact invariant subset is the union of finitely homoclinic orbits of singularities. See Section 6.

Corollary 3.7. *Let $f : M^d \rightarrow \mathbb{R}$ be a continuous function. Let μ be an ergodic measure of a flow ϕ_t generated by $X \in \mathcal{X}^1(M^d)$, which is not supported on a singularity. Assume that Λ is a compact invariant set such that $\mu(\Lambda) = 0$. Then for any $\varepsilon > 0$, for any C^1 neighborhood \mathcal{U} of X , and any neighborhood U of $\text{supp}(\mu)$, there is a periodic orbit $\gamma \subset U$ of $Y \in \mathcal{U}$ such that*

- *Λ is a compact invariant set of Y .*
- *$|\int f d\delta_\gamma - \int f d\mu| < \varepsilon$, where δ_γ is the uniform distribution measure on γ , i.e., for any continuous function $g : M^d \rightarrow \mathbb{R}$,*

$$\int g(x) d\delta_\gamma(x) = \frac{1}{\tau(\gamma)} \int_0^{\tau(\gamma)} g(\phi_t(p)) dt,$$

where $p \in \gamma$.

Proof. Let $C = \max\{\sup_{x \in M^d} |f(x)|, 1\}$. Without loss of generality, one can assume that μ is not supported on a periodic orbit. Since μ is not supported on a singularity, one has $\mu(\Sigma(X)) = 0$. We choose $x \in \Sigma$ satisfying

$$\lim_{t \rightarrow \infty} \delta_{x,T} = \mu,$$

where $\delta_{x,T}$ is the uniform distribution measure supported on $\phi_{[0,T]}(x)$, i.e., for any continuous function $g : M^d \rightarrow \mathbb{R}$,

$$\int g(y) d\delta_{x,T}(y) = \frac{1}{T} \int_0^T g(\phi_t(x)) dt.$$

For any $\varepsilon > 0$, there is $T_0 > 0$ such that for any $T > T_0$, one has

$$\left| \int f d\delta_{x,T} - \int f d\mu \right| < \varepsilon/2.$$

By reducing ε if necessary, one can assume that $B(x, \varepsilon) \cap \Lambda = \emptyset$. Since Λ is invariant, one has that $\phi_t(B(x, \varepsilon)) \cap \Lambda = \emptyset$ for any $t \in \mathbb{R}$. By Lemma 3.6, there is Y which has a periodic point p with period $\tau(p)$ such that

- $d_{C^1}(X, Y) < \varepsilon$;
- Λ is a compact invariant set of Y ;
- $d(\phi_t^X(x), \phi_t^Y(p)) < \varepsilon/2C$ for each $t \in [0, \tau(p)]$.

x is not periodic because we assume that μ is not supported on a periodic orbit. Thus, one can assume that $\tau(p) > T_0$. Let $\gamma = \text{Orb}(p)$. Then

$$\begin{aligned} \left| \int f d\delta_\gamma - \int f d\mu \right| &\leq \left| \int f d\delta_\gamma - \int f d\delta_{x, \tau(\gamma)} \right| + \left| \int f d\delta_{x, \tau(\gamma)} - \int f d\mu \right| \\ &< C \frac{\varepsilon}{2C} + \frac{\varepsilon}{2} = \varepsilon. \end{aligned}$$

□

A vector field $X \in \mathcal{X}^r(M^d)$ is called *Kupka-Smale* if every critical element is hyperbolic, and the stable manifold of any critical element intersects the unstable manifold of any other critical element transversely. A classical generic result is: Kupka-Smale vector fields form a residual set in $\mathcal{X}^r(M^d)$. We need the following weak terminology:

Definition 3.8. *A vector field $X \in \mathcal{X}^r(M^d)$ is called weak Kupka-Smale if every critical element of X is hyperbolic.*

Since Kupka-Smale is C^r generic in $\mathcal{X}^r(M^d)$, we have that weak Kupka-Smale is also a C^r generic property in $\mathcal{X}^r(M^d)$.

We will state a connecting lemma for pseudo orbits, which was studied in [7]. [7] studied the connecting lemma for pseudo-orbits for weak Kupka-Smale diffeomorphisms. The assumption of weak Kupka-Smale is used since

- By using λ -lemma, for every non-periodic point x which is not in the stable/unstable manifold of a periodic point, the positive/negative iteration of x will be in a topological tower.

For flows, λ -lemma is true for both hyperbolic singularities and hyperbolic periodic orbits. The orbit structure is clear near hyperbolic critical elements. So the connecting lemma for pseudo-orbits is true for vector fields.

Lemma 3.9. ([7, Theorem 1.2])

Let $X \in \mathcal{X}^1(M^d)$ be a weak Kupka-Smale vector field. Given any finite set \hat{F} of periodic orbits of X , for any neighborhood U of \hat{F} , for any C^1 neighborhood \mathcal{U} of X , and for any $x, y \in M^d \setminus U$, if y is chain attainable from x , then there is $Y \in \mathcal{U}$ and $t > 0$ such that $\phi_t^Y(x) = y$ and $Y(z) = X(z)$ for any $z \in U$.

Moreover, if X is C^r , we can require that Y is also C^r .

The statement here is a little bit different from [7], but it is essentially contained there. See [7, Remarque 1.1].

Remark. Due to Lemma 3.9, we can consider a weak Kupka-Smale vector field close to a generic one. See Lemma 3.14 and Section 6 for more details.

Sometimes, one needs to perturb a vector field to be a weak Kupka-Smale vector field by preserving some non-transverse connection. This was used by Palis [41]. One can see R. Xi's master thesis [63] for a proof.

Theorem 3.10. [41, 63] For any vector field $X \in \mathcal{X}^r(M^d)$, any $n \in \mathbb{N}$ and any hyperbolic critical elements $\{P_1, Q_1, \dots, P_n, Q_n\}$, if $\text{Orb}(x_i) \subset W^s(P_i) \cap W^u(Q_i)$ is a non-transverse orbit for $1 \leq i \leq n$, then for any C^r neighborhood \mathcal{U} of X there exists $Y \in \mathcal{U}$,

- $\text{Orb}(x_i) \subset W^s(P_i) \cap W^u(Q_i)$ is still a non-transverse orbit of Y .
- Any other critical element of Y is hyperbolic, in other words, Y is weak Kupka-Smale.

3.3 Generic results

$\mathcal{R} \subset \mathcal{X}^1(M^d)$ is called *residual* if it contains a dense G_δ subsets of $\mathcal{X}^1(M^d)$. A property of vector fields is called C^1 *generic* if it holds on a residual set in $\mathcal{X}^1(M^d)$. Sometimes we use the terminology “for C^1 generic X ” which is equivalent to say that “there is a residual set $\mathcal{R} \subset \mathcal{X}^1(M^d)$ and $X \in \mathcal{R}$ ”. Being a Banach space, every residual set of $\mathcal{X}^1(M^d)$ is dense. Usually we can get a dense open property via a generic way.

One knows that lower semi-continuous maps and upper semi-continuous maps defined on a complete metric space are continuous on a residual set.

Lemma 3.11. For C^1 generic $X \in \mathcal{X}^1(M^d)$, for every critical element p , $C(p, X)$ is continuous at X . This means, if $\{X_n\}$ is a sequence of vector fields and $\lim_{n \rightarrow \infty} X_n = X$ in the C^1 topology, then $\lim_{n \rightarrow \infty} C(p_{X_n}, X_n) = C(p, X)$ in the Hausdorff topology.

Proof. The proof of this lemma just uses the upper semi-continuity of chain recurrent class. \square

Lemma 3.12. For C^1 generic $X \in \mathcal{X}^1(M^d)$, if p_1 and p_2 are two different critical elements, and if $C(p_1) = C(p_2)$, then there is a C^1 neighborhood \mathcal{U} of X such that for any $Y \in \mathcal{U}$, one has $C(p_{1,Y}, Y) = C(p_{2,Y}, Y)$.

Proof. Let \mathcal{C} be the metric space of all compact subsets of M^d , endowed with the Hausdorff metric. \mathcal{C} is a compact metric space. Let $\mathcal{B}_1, \mathcal{B}_2, \dots, \mathcal{B}_n, \dots$, be a countable basis of \mathcal{C} . Let $\mathcal{O}_1, \mathcal{O}_2, \dots, \mathcal{O}_n, \dots$, be the list of finite unions of elements of the countable basis. For each n and m , we define the sets $\mathcal{H}_{n,m}$ and $\mathcal{N}_{n,m}$ of vector fields as following:

- $X \in \mathcal{H}_{n,m}$ iff there is a neighborhood \mathcal{U} of X such that for any $Y \in \mathcal{U}$, for any hyperbolic critical element $p_n \in \mathcal{O}_n$ and any hyperbolic critical element $p_m \in \mathcal{O}_m$, one has $C(p_n) = C(p_m)$ for Y .

- $X \in \mathcal{N}_{n,m}$ iff there is a hyperbolic critical element $p_n \in \mathcal{O}_n$ and a hyperbolic critical element $p_m \in \mathcal{O}_m$ such that $C(p_n) \cap C(p_m) = \emptyset$. Since every chain recurrent class is upper semi continuous, one knows that $\mathcal{N}_{n,m}$ is open.

From the definitions, $\mathcal{H}_{n,m} \cup \mathcal{N}_{n,m}$ is open and dense in $\mathcal{X}^1(M^d)$. Let

$$\mathcal{R} = \bigcap_{n,m \in \mathbb{N}} (\mathcal{H}_{n,m} \cup \mathcal{N}_{n,m}).$$

\mathcal{R} is a residual subset. We will verify that every $X \in \mathcal{R}$ satisfies the the conclusion of the lemma. Let $X \in \mathcal{R}$. Thus, for each n and m one has $X \in \mathcal{H}_{n,m} \cup \mathcal{N}_{n,m}$. For any two hyperbolic critical elements p_1 and p_2 , there are $l \in \mathbb{N}$ and $k \in \mathbb{N}$ and a C^1 neighborhood \mathcal{U} of X such that

- For any $Y \in \mathcal{U}$, $\text{Orb}_Y(p_{1,Y})$ and $\text{Orb}_Y(p_{2,Y})$ are the maximal compact invariant sets in \mathcal{O}_l and \mathcal{O}_k respectively.

If $C(p_1) = C(p_2)$ for X , then $X \notin \mathcal{N}_{k,l}$. This implies that $X \in \mathcal{H}_{k,l}$. Let $\mathcal{U}_0 = \mathcal{U} \cap \mathcal{H}_{k,l}$. For each $Y \in \mathcal{U}_0$, one has

- Since $Y \in \mathcal{H}_{k,l}$, there is a critical point $p'_1 \in \mathcal{O}_l$ and a critical point $p'_2 \in \mathcal{O}_k$ of Y , one has $C(p'_1) = C(p'_2)$.
- Since $Y \in \mathcal{U}$, the unique critical point in \mathcal{O}_l is $p_{1,Y}$ and the unique critical point in \mathcal{O}_k is $p_{2,Y}$. As a corollary, $p_{1,Y} = p'_1$ and $p_{2,Y} = p'_2$.

Thus, one has $C(p_{1,Y}) = C(p_{2,Y})$ for any $Y \in \mathcal{U}_0$. □

Lemma 3.13. *For C^1 generic $X \in \mathcal{X}^1(M^d)$, and any hyperbolic critical element p of X , if $\overline{W^u(p)} \subset C(p)$, then there is a C^1 neighborhood \mathcal{U} of X such that $\overline{W^u(p_Y, Y)} \subset C(p_Y, Y)$.*

Proof. Let \mathcal{C} be the metric space of all compact subsets of M^d , endowed with the Hausdorff metric. \mathcal{C} is a compact metric space. Let $\mathcal{B}_1, \mathcal{B}_2, \dots, \mathcal{B}_n, \dots$, be a countable basis of \mathcal{C} . Let $\mathcal{O}_1, \mathcal{O}_2, \dots, \mathcal{O}_n, \dots$, be the list of finite unions of elements of the countable basis. For each n , one can define sets \mathcal{H}_n and \mathcal{N}_n as following:

- $X \in \mathcal{H}_n$ iff there is a C^1 neighborhood \mathcal{U} of X such that for any $Y \in \mathcal{U}$, every hyperbolic critical orbit $\text{Orb}_Y(p_Y) \in \mathcal{O}_n$ of Y satisfies $\overline{W^u(p_Y, Y)} \subset C(p_Y, Y)$. By definition, \mathcal{H}_n is open.
- $X \in \mathcal{N}_n$ iff X has a hyperbolic critical element $p \in \mathcal{O}_n$ such that $\overline{W^u(p, X)} \not\subseteq C(p, X)$. $\overline{W^u(p, X)}$ varies lower semi-continuously with respect to X and $C(p, X)$ varies upper semi-continuously with respect to X . So if $\overline{W^u(p, X)} \not\subseteq C(p, X)$, there is a neighborhood \mathcal{U} of X such that $\overline{W^u(p_Y, Y)} \not\subseteq C(p_Y, Y)$ for any $Y \in \mathcal{U}$. This would imply that \mathcal{N}_n is an open set in $\mathcal{X}^1(M^d)$.

It's clear that $\mathcal{H}_n \cup \mathcal{N}_n$ is open and dense in $\mathcal{X}^1(M^d)$. Let

$$\mathcal{R} = \bigcap_{n \in \mathbb{N}} (\mathcal{H}_n \cup \mathcal{N}_n).$$

\mathcal{R} is a residual subset of $\mathcal{X}^1(M^d)$. Take $X \in \mathcal{R}$. If p is a hyperbolic critical element of X , then there are n and a neighborhood \mathcal{U} of X such that for each $Y \in \mathcal{U}$, $\text{Orb}_Y(p_Y)$ is the unique hyperbolic critical orbit in \mathcal{O}_n . Since $\overline{W^u(p)} \subset C(p)$, one has $X \notin \mathcal{N}_n$. As a corollary, $X \in \mathcal{H}_n$. Let $\mathcal{U}_0 = \mathcal{U} \cap \mathcal{H}_n$. For any $Y \in \mathcal{U}_0$,

- Since $Y \in \mathcal{H}_n$, every hyperbolic critical element $q \in \mathcal{O}_n$ of Y , one has $\overline{W^u(q, Y)} \subset C(q, Y)$.
- Since $Y \in \mathcal{U}$, the unique critical element in \mathcal{O}_n is p_Y .

Thus, $\overline{W^u(p_Y, Y)} \subset C(p_Y, Y)$ for any $Y \in \mathcal{U}_0$. \square

The following lemma claims that for C^1 generic vector field, Lyapunov stability is a robust property under perturbations.

Lemma 3.14. *For C^1 generic $X \in \mathcal{X}^1(M^d)$, let p be a hyperbolic critical element of X . If $C(p, X)$ is a Lyapunov stable chain recurrent class, then there is a C^1 neighborhood \mathcal{U} of X such that for any weak Kupka-Smale vector field $Y \in \mathcal{U}$, $C(p_Y, Y)$ is a Lyapunov stable chain recurrent class of Y .*

Proof. Let $\mathcal{R} \subset \mathcal{X}^1(M^d)$ be the residual subset as in Lemma 3.13: $X \in \mathcal{R}$ iff for any hyperbolic critical element p of X , if $\overline{W^u(p, X)} \subset C(p, X)$, then there is a C^1 neighborhood $\mathcal{U} = \mathcal{U}_{X,p}$ such that $\overline{W^u(p_Y, Y)} \subset C(p_Y, Y)$ for any $Y \in \mathcal{U}$. We will prove that $C(p_Y, Y)$ is Lyapunov stable chain recurrent class for each weak Kupka-Smale $Y \in \mathcal{U}$. If not, there is a weak Kupka-Smale vector field $X_0 \in \mathcal{U}$ such that $C(p_{X_0}, X_0)$ is not Lyapunov stable. Thus we have

- $\overline{W^u(p_{X_0}, X_0)} \subset C(p_{X_0}, X_0)$.
- There is $y \notin C(p_{X_0}, X_0)$ such that $y \in W^{ch,u}(C(p_{X_0}, X_0))$.

Then, there is a C^1 neighborhood $\mathcal{U}_0 \subset \mathcal{U}$ of X_0 such that for any $Y \in \mathcal{U}_0$, $y \notin C(p_Y, Y)$ by the upper semi-continuity of chain recurrent classes. Choose $z \in W^u(p_{X_0}, X_0) \setminus \{p_{X_0}\}$. Since $z \in C(p_{X_0}, X_0)$, y is chain attainable from z . Choose a small neighborhood U of p_{X_0} .

One can assume that the negative orbit of z is contained in U . By Lemma 3.9, there is a vector field $Y \in \mathcal{U}_0$ such that

- $Y(x) = X_0(x)$ for any $x \in U$.
- y is in the positive orbit of z with respect to ϕ_t^Y .

As a corollary, y is in the unstable manifold of p_Y with respect to Y . This fact gives a contradiction. \square

The following lemma asserts that for a generic vector field, if the perturbed system has a periodic orbit which is $(C, \eta, d, \mathcal{N})$ -contracting at the period, then the original generic system already have itself by relaxing the constants.

Lemma 3.15. *There is a dense G_δ set $\mathcal{G} \subset \mathcal{X}^1(M^d)$ such that for any $X \in \mathcal{G}$, if for any two open sets $U, V \subset M^d$ with $\overline{U} \subset V$, for any C^1 neighborhood \mathcal{U} of X , and any three number $K \in \mathbb{N}$, $\eta > 0$ and $T > 0$, if some $Y \in \mathcal{U}$ has a hyperbolic periodic orbit γ_Y which is $(K, \eta, T, \mathcal{N})$ -contracting at the period w.r.t. ψ_t^Y such that $\gamma_Y \cap U \neq \emptyset$, then X has a periodic orbit γ which is $(K, \eta/2, 2T, \mathcal{N})$ -contracting at the period w.r.t. ψ_t such that $\gamma \cap V \neq \emptyset$.*

Proof. The proof of this lemma is an application of fundamental tricks for generic properties. Thus we just give a sketch. Let $O_1, O_2, \dots, O_n, \dots$ be a topological base of M^d . Let $\{\eta_m\}$ and $\{d_\ell\}$ be the sets of positive rational numbers. For each $n \in \mathbb{N}$, $K \in \mathbb{N}$, $m \in \mathbb{N}$ and $\ell \in \mathbb{N}$, one defines:

- $X \in \mathcal{H}_{n,K,m,\ell}$ iff there is a hyperbolic periodic orbit γ such that
 - $\gamma \cap O_n \neq \emptyset$.

- There is $x \in \gamma$ and a time partition $0 = t_0 < t_1 < \dots < t_q = \alpha\pi(\gamma)$ for some positive integer α satisfying $t_{i+1} - t_i \leq d_\ell$ for $0 \leq i \leq q - 1$ such that

$$\prod_{i=0}^{q-1} \|\psi_{t_{i+1}-t_i}|_{\mathcal{N}_{\phi_{t_i}(x)}}\| < Ke^{-\eta_m \alpha\pi(\gamma)}.$$

- $X \in \mathcal{N}_{n,K,m,\ell}$ iff there is a neighborhood \mathcal{U} of X such that for any $Y \in \mathcal{U}$, one has
 - either Y has no hyperbolic periodic orbit intersecting O_n ;
 - Or, Y has a hyperbolic periodic orbit γ such that $\gamma \cap O_n \neq \emptyset$ and for any $x \in \gamma$ and any time partition $0 = t_0 < t_1 < \dots < t_q = \alpha\pi(\gamma)$ for some positive integer α satisfying $t_{i+1} - t_i \leq d_\ell$ for $0 \leq i \leq q - 1$, one has

$$\prod_{i=0}^{q-1} \|\psi_{t_{i+1}-t_i}^Y|_{\mathcal{N}_{\phi_{t_i}(x)}}\| \geq Ke^{-\eta_m \alpha\pi(\gamma)}.$$

By the definitions, $\mathcal{H}_{n,K,m,\ell} \cup \mathcal{N}_{n,K,m,\ell}$ is a dense open set in $\mathcal{X}^1(M^d)$. Thus,

$$\mathcal{G} = \bigcap_{n,K,m,\ell \in \mathbb{N}} (\mathcal{H}_{n,K,m,\ell} \cup \mathcal{N}_{n,K,m,\ell})$$

is a residual subset of $\mathcal{X}^1(M^d)$. Now we check that every $X \in \mathcal{G}$ satisfies the properties of the lemma.

For any $\eta > 0$, $T > 0$, one can take a rational number $\eta_{m_0} \in (\eta/2, \eta)$ and $T_{\ell_0} \in (T, 2T)$. If there is a sequence of vector fields $\{X_n\}$ such that

- $\lim_{n \rightarrow \infty} X_n = X$.
- Each X_n has a hyperbolic periodic orbit γ_n which is $(K, \eta, T, \mathcal{N})$ -contracting at the period such that $\gamma_n \cap U \neq \emptyset$.

There is $x \in \overline{U}$ such that x is an accumulating point of γ_n . Thus, there is n_0 such that $x \in O_{n_0} \subset V$. Since $X \in \mathcal{G} \subset \mathcal{H}_{n_0,K,m_0,\ell_0} \cup \mathcal{N}_{n_0,K,m_0,\ell_0}$, one has $X \in \mathcal{H}_{n_0,K,m_0,\ell_0}$ by the definitions. Thus X has a periodic orbit in O_{n_0} itself which is $(K, \eta_{m_0}, T_{\ell_0}, \mathcal{N})$ -contracting at the period w.r.t. ψ_t . It's $(K, \eta/2, 2T, \mathcal{N})$ -contracting at the period w.r.t. ψ_t in V . \square

Lemma 3.16. *There is a dense G_δ set $\mathcal{G} \subset \mathcal{X}^1(M^d)$ such that for any $X \in \mathcal{G}$ and $x \in M^d$, for any $K \in \mathbb{N}$, $\eta > 0$ and $d > 0$, one has*

- either, x is contained in a periodic sink which is $(K, \eta/2, 2d, \mathcal{N})$ -contracting at the period w.r.t. ψ_t ;
- or, there is a C^1 neighborhood \mathcal{U} of X and a neighborhood U of x such that for any $Y \in \mathcal{U}$, Y has no periodic sink which is $(K, \eta, d, \mathcal{N})$ -contracting at the period w.r.t. ψ_t^Y , and intersects U .

Proof. Let \mathcal{G} be as in Lemma 3.15. If the conclusion of this lemma is not true, one has that

- x is not contained in a periodic sink which is $(K, \eta/2, 2d, \mathcal{N})$ -contracting at the period w.r.t. ψ_t .
- For any C^1 neighborhood \mathcal{U} of X and any neighborhood U of x , some $Y \in \mathcal{U}$ has a periodic sink, which is $(K, \eta, d, \mathcal{N})$ -contracting at the period w.r.t. ψ_t^Y , and intersects U for.

Thus, by Lemma 3.15, for any neighborhood of x , X itself has a periodic sink, which is $(K, \eta/2, 2d, \mathcal{N})$ -contracting at the period w.r.t. ψ_t . In other words, there is a sequence of periodic points x_n such that

- $\lim_{n \rightarrow \infty} x_n = x$.
- x_n is contained in a period sink γ_n , which is $(K, \eta, d, \mathcal{N})$ -contracting at the period for each n . Moreover, $\{\gamma_n\}$ are distinct.

We assert that $\tau(\gamma_n) \rightarrow \infty$ as $n \rightarrow \infty$. Otherwise, by taking a limit, x would be in a periodic sink which is $(K, \eta, d, \mathcal{N})$ -contracting at the period. Thus one can get a contradiction by Lemma 2.22. \square

Corollary 3.17. *Assume that $\dim M^3 = 3$. There is a residual set $\mathcal{G} \subset \mathcal{X}^1(M^3) \setminus \overline{\mathcal{HT}}$ and $\iota > 0$ such that for any $X \in \mathcal{G}$, any $\sigma \in \text{Sing}(X)$, there is a C^1 neighborhood \mathcal{U} of X and a neighborhood U of σ such that for any periodic orbit γ of Y , if $\gamma \cap U \neq \emptyset$, then \mathcal{N}_γ admits an ι -dominated splitting of index 1 w.r.t. the linear Poincaré flow.*

Proof. If $\text{ind}(\gamma) = 1$, then it is done by Lemma 2.9. Thus, one can assume that γ is a sink or source. Without loss of generality, we assume that it is a sink. By Lemma 2.11, if γ does not admit an ι -dominated splitting for some ι , then there are $C > 0$, $\eta > 0$ and $T > 0$ such that γ is $(C, \eta, T, \mathcal{N})$ -contracting at the period w.r.t. ψ_t . Since σ is a singularity, not a periodic point, by Lemma 3.16, one can get a contradiction. \square

We would like to list some other generic results we need in this paper.

Lemma 3.18. *There is a dense G_δ set $\mathcal{G} \subset \mathcal{X}^1(M^d)$ such that for each $X \in \mathcal{G}$, one has*

1. *For any non-trivial chain recurrent class $C(\sigma)$, where σ is a hyperbolic singularity of index $d - 1$, then every separatrix of $W^u(\sigma)$ is dense in $C(\sigma)$. Especially, $C(\sigma)$ is transitive.*
2. *Let $i \in [0, \dim M - 1]$. If there is a sequence of vector fields $\{X_n\}$ such that*
 - $\lim_{n \rightarrow \infty} X_n = X$,
 - *each X_n has a hyperbolic periodic orbits γ_{X_n} of index i such that $\lim_{n \rightarrow \infty} \gamma_{X_n} = \Lambda$,*

Then there is a sequence of hyperbolic periodic orbits γ_n of index i of X such that $\lim_{n \rightarrow \infty} \gamma_n = \Lambda$.

3. *There exists a neighborhood \mathcal{U} of X such that for any $Y \in \mathcal{U}$, Y has only finitely many singularities. Moreover, for every singularity σ of Y , the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_d$ of $DY(\sigma)$ satisfy:*

$$\text{Re}(\lambda_i) + \text{Re}(\lambda_j) \neq 0,$$

for any $1 \leq i, j \leq d$.

4. *For any hyperbolic periodic orbit P of X , then $C(P) = H(P)$, where $H(P)$ is the homoclinic class of P .*
5. *Every non-trivial chain transitive set is the limit of a sequence of periodic orbits in the Hausdorff topology.*
6. *X is Kupka-Smale.*

Remark. Item 1 is a corollary of the connecting lemma for pseudo-orbits [7]. There is no explicit version like this. [34, Section 4] gave some ideas about the proof of Item 1 without using of the terminology of chain recurrence. Item 2 is fundamental, one can see [60] for instance. Item 3 is fundamental. It is true because generic X can only have finitely many singularities. Moreover, the eigenvalues of the singularities have some continuous property. Item 4 is also a result in [7]. Item 5 is the main result in [15]. Item 6 is the classical Kupka-Smale theorem.

Let \mathcal{G}_0 be a dense G_δ set of $\mathcal{X}^1(M^d)$ such that $X \in \mathcal{G}_0$ iff X satisfies all generic properties in this subsection.

4 Reduce the proof into technical results

Now we proof Theorem A under the condition that Theorem B is true.

Proof of Theorem A. Now we give the proof of Theorem A by assuming the result of Theorem B. Suppose on the contrary that $\mathcal{X}^1(M^3) \setminus \overline{\mathcal{MS} \cup \mathcal{HS}}$ is not empty. Choose C^1 generic $X \in \mathcal{X}^1(M^3) \setminus \overline{\mathcal{MS} \cup \mathcal{HS}}$. Since every homoclinic tangency of a hyperbolic periodic orbit can be perturbed to be a transverse homoclinic intersection by an arbitrarily small perturbation, we have that X is far away from ones with a homoclinic tangency. Thus, by Theorem B, every chain recurrent class is a homoclinic class. Since we assume that Theorem A is not true, one has that every chain recurrent class is trivial: it is reduced to be a critical element. If there are finitely many chain recurrent classes, then we have that X is Morse-Smale. Thus, one can assume that X has infinitely many chain recurrent classes, and each chain recurrent class is a hyperbolic critical element. It is known that C^1 generic vector field can only have finitely many singularities since it is Kupka-Smale. Thus X has countably many distinct hyperbolic periodic orbits $\{\gamma_n\}$. By taking a subsequence if necessary, we can assume that $\{\gamma_n\}$ converges in the Hausdorff topology. Let Λ be the limit. Then, Λ is chain transitive, which implies that Λ is contained in a chain recurrent class. Because we know that every chain recurrent class of X is a hyperbolic critical element, Λ is a hyperbolic critical element. This cannot happen because hyperbolic critical elements are locally maximal. \square

To prove Theorem B, we need to prove the following Theorem 4.1 and 4.2.

Theorem 4.1. For C^1 generic $X \in \mathcal{X}^1(M^3) \setminus \overline{\mathcal{HT}}$, for the non-trivial chain recurrent class $C(\sigma)$ of some singularity σ , if $C(\sigma)$ is not a homoclinic class, then $C(\sigma)$ admits a dominated splitting $T_{C(\sigma)}M^3 = E \oplus F$ w.r.t. the tangent flow.

Theorem 4.2. For C^1 generic $X \in \mathcal{X}^1(M^3)$, for any non-trivial chain recurrent class $C(\sigma)$ of some singularity σ , if $C(\sigma)$ admits a dominated splitting $T_{C(\sigma)}M^3 = E \oplus F$ w.r.t. the tangent flow, then $C(\sigma)$ is a homoclinic class.

In general, we give the definition of singular hyperbolic sets as the following:³

Definition 4.3. Let Λ be a compact invariant set of $X \in \mathcal{X}^1(M^3)$, $E \subset T_\Lambda M^3$ a two dimensional invariant sub-bundle, we say that E is area-contracting, if there are $C > 0$, $\lambda > 0$ such that for any $x \in \Lambda$ and any $t > 0$, $|\det \Phi_t|_{E(x)}| \leq Ce^{-\lambda t}$; we say that E is area-expanding if it is area-contracting for $-X$.

A compact invariant set Λ of $X \in \mathcal{X}^1(M^3)$ is called singular hyperbolic, if

- either, Λ admits a partially hyperbolic splitting $T_\Lambda M^3 = E^{ss} \oplus E^{cu}$, where $\dim E^{ss} = 1$, E^{ss} is contracting and E^{cu} is area-expanding;

³We postpone to give the definition because we want the introduction to be easier to read.

- or, Λ admits a partially hyperbolic splitting $T_\Lambda M^3 = E^{cs} \oplus E^{uu}$, where $\dim E^{uu} = 1$, E^{uu} is expanding and E^{cs} is area-contracting.

Note that in the above definition, we don't require Λ contains a singularity or not. So every non-singular hyperbolic set is singular hyperbolic.

We also needs the following two results from [12].

Lemma 4.4. *For C^1 generic $X \in \mathcal{X}^1(M^3)$, if Λ is a non-singular chain transitive set and admits a dominated splitting $\mathcal{N}_\Lambda = \Delta^s \oplus \Delta^u$ with respect to ψ_t , then Λ is hyperbolic.*

Theorem 4.5. *For C^1 generic $X \in \mathcal{X}^1(M^3)$, if the chain recurrent class $C(\sigma)$ of a singularity σ contains a **periodic orbit** and admits a dominated splitting $T_{C(\sigma)}M = E \oplus F$ with respect to Φ_t , then $C(\sigma)$ is singular hyperbolic. Consequently, $C(\sigma)$ is an attractor or repeller according to the index of σ equals to 2 or 1.*

Proof of Theorem B. If Theorem B is not true, then there is a C^1 generic $X \in \mathcal{X}^1(M^3) \setminus \overline{\mathcal{HT}}$ and a chain recurrent class \mathcal{C} of X such that \mathcal{C} is not a homoclinic class. Now we have two cases:

\mathcal{C} contains no singularities. Since \mathcal{C} is chain transitive, there is a sequence of periodic orbits $\{\gamma_n\}$ such that $\gamma_n \rightarrow \mathcal{C}$ as $n \rightarrow \infty$ in the Hausdorff metric. Since \mathcal{C} is not reduced to a periodic orbit, one can assume that $\{\gamma_n\}$ are distinct periodic orbits and $\tau(\gamma_n) \rightarrow \infty$. By Corollary 2.8, if we cannot perturb γ_n of hyperbolic periodic orbit of index 1 for n large enough, then there are constants $C > 0$, $T > 0$, $\eta > 0$ such that γ_n are $(C, \eta, T, \mathcal{N})$ -contracting at the period w.r.t. ψ_t for X or $-X$. Then by Lemma 2.22, one can get a contradiction.

Thus, one can assume that the index of every γ_n is 1. From this, we have that \mathcal{C} admits a dominated splitting $\mathcal{N}_\mathcal{C} = \Delta^{cs} \oplus \Delta^{cu}$ of index 1 w.r.t. ψ_t . By Lemma 4.4, we have that \mathcal{C} is hyperbolic. This fact shows that \mathcal{C} is a homoclinic class, which gives a contradiction. \square

\mathcal{C} contains a singularity σ . We assume that \mathcal{C} is non-trivial. Since \mathcal{C} is not a homoclinic class, by Theorem 4.1, \mathcal{C} admits a dominated splitting $T_\mathcal{C}M^3 = E \oplus F$ w.r.t. the tangent flow Φ_t . Then by Theorem 4.2, \mathcal{C} is a homoclinic class, which gives a contradiction. \square

Proof of Theorem C. For C^1 generic $X \in \mathcal{X}(M^3)$, if $C(\sigma)$ admits a dominated splitting w.r.t. the tangent flow, then by Theorem 4.2, \mathcal{C} is a homoclinic class. By using Theorem 4.5, $C(\sigma)$ is singular hyperbolic. \square

4.1 Proof of Theorem 4.1

The proof of Theorem 4.1 can be divided into the following two propositions:

Proposition 4.6. *There is a dense G_δ set $\mathcal{G} \in \mathcal{X}^1(M^3) \setminus \overline{\mathcal{HT}}$ such that for any $X \in \mathcal{G}$, if σ is a hyperbolic saddle of X and $C(\sigma)$ is Lyapunov stable, then every singularity $\rho \in C(\sigma)$ is Lorenz-like, i.e., the eigenvalues $\lambda_1, \lambda_2, \lambda_3$ of $DX(\rho)$ satisfy:*

$$\lambda_1 < \lambda_2 < 0 < -\lambda_2 < \lambda_3.$$

Moreover, there is a C^1 neighborhood \mathcal{U}_X of X such that for any $Y \in \mathcal{U}_X$ and any $\rho \in C(\sigma) \cap \text{Sing}(X)$, one has $\rho_Y \in C(\sigma_Y, Y)$ and $W^{ss}(\rho_Y) \cap C(\sigma_Y, Y) = \{\rho_Y\}$.

Proposition 4.7. *Let Λ be a compact invariant set of $X \in \mathcal{X}^1(M^d)$ verifying the following properties:*

- $\Lambda \setminus \text{Sing}(X)$ admits an index i dominated splitting $\mathcal{N}_{\Lambda \setminus \text{Sing}(X)} = \Delta^{cs} \oplus \Delta^{cu}$ in the normal bundle w.r.t. the linear Poincaré flow ψ_t .
- Every singularity $\sigma \in \Lambda$ is hyperbolic and $\text{ind}(\sigma) > i$. Moreover, $T_\sigma M^d$ admits a partially hyperbolic splitting $T_\sigma M^d = E^{ss} \oplus E^{cu}$ with respect to the tangent flow, where $\dim E^{ss} = i$ and for the corresponding strong stable manifolds $W^{ss}(\sigma)$, one has $W^{ss}(\sigma) \cap \Lambda = \{\sigma\}$.
- For every $x \in \Lambda$, one has $\omega(x) \cap \text{Sing}(X) \neq \emptyset$.

Then one has

- either Λ admits a partially hyperbolic splitting $T_\Lambda M^d = E^{ss} \oplus F$ with respect to the tangent flow Φ_t , where $\dim E^{ss} = i$,
- or, there is a sequence of hyperbolic periodic orbits $\{\gamma_n\}_{n \in \mathbb{N}}$ of index i such that
 - $\tau(\gamma_n) \rightarrow \infty$,
 - $H(\gamma_n) \cap \Lambda \neq \emptyset$ for each $n \in \mathbb{N}$,
 - There is $T > 0$ such that for $x_n \in \gamma_n$,

$$\lim_{n \rightarrow \infty} \frac{1}{[\tau(\gamma_n)/T]} \sum_{i=0}^{[\tau(\gamma_n)/T]-1} \log \|\psi_T|_{\mathcal{N}^s(x_n)}\| = 0.$$

Now one can give a proof of Theorem 4.1 by Proposition 4.6 and Proposition 4.7.

Proof of Theorem 4.1. Since $\dim M^3 = 3$, without loss generality, one can assume that $\text{ind}(\sigma) = 2$. Otherwise, one considers $-X$. By Lemma 3.18, $C(\sigma)$ is Lyapunov stable. By Proposition 4.6, every singularity ρ in $C(\sigma)$ is Lorenz-like and $W^{ss}(\rho) \cap C(\sigma) = \{\rho\}$. Since $X \in \mathcal{X}^1(M^3) \setminus \overline{\mathcal{HT}}$, one has

- The normal bundle of $C(\sigma) \setminus \text{Sing}(X)$ admits a dominated splitting with respect to the linear Poincaré flow (See more details from Corollary 3.17).
- $C(\sigma)$ contains no periodic points. As a corollary, for every regular point $x \in C(\sigma)$, $\omega(x)$ contains a singularity. Otherwise, if $\omega(x)$ contains no singularities, then by Lemma 4.4, $\omega(x)$ is hyperbolic. Then one can get a periodic orbit in $C(\sigma)$ by the shadowing lemma, which is a contradiction.

By Proposition 4.7, either $C(\sigma)$ admits a partially hyperbolic splitting, or $C(\sigma) \cap H(\gamma) \neq \emptyset$ for some hyperbolic saddle γ . But the fact that $C(\sigma) \cap H(\gamma) \neq \emptyset$ gives a contradiction. \square

4.2 The proof of Theorem 4.2

The proof of Theorem 4.2 involves more notations and definitions. It's not easy to state the precise results here. But we can give some rough idea here.

Proposition 4.8. *For C^1 generic $X \in \mathcal{X}^1(M^3)$, for a Lyapunov stable chain recurrent class $C(\sigma)$, if $C(\sigma)$ contains no periodic orbits and $C(\sigma)$ admits a partially hyperbolic splitting $T_{C(\sigma)} M^3 = E^{ss} \oplus E^{cu}$ with $\dim E^{ss} = 1$, then $C(\sigma)$ admits a cross section system (Σ, F) . Moreover, this cross section system (Σ, F) has some continuous properties w.r.t. X .*

The precise definition of cross section system could be seen in Section 6.

Proposition 4.9. *For C^1 generic $X \in \mathcal{X}^1(M^3)$ and $\sigma \in \text{Sing}(X)$, if the chain recurrent class $C(\sigma)$ is Lyapunov stable, contains no periodic orbits and admits a partially hyperbolic splitting $T_{C(\sigma)}M^3 = E^{ss} \oplus E^{cu}$ with $\dim E^{ss} = 1$, then for any C^1 neighborhood \mathcal{U} of X , there is a weak Kupka-Smale vector field $Y \in \mathcal{U}$ such that Y has a periodic sink γ , and $C(\sigma_Y) \cap \overline{B(\gamma)} \neq \emptyset$, where $B(\gamma)$ is the basin of γ and $\overline{B(\gamma)}$ is the closure of $B(\gamma)$.*

The proof of Proposition 4.9 needs to use the cross section system constructed in Proposition 4.8

Proposition 4.9 could imply Theorem 4.2: Lemma 3.14 implies for every weak Kupka-Smale vector field Y close to X , $C(\sigma_Y)$ is Lyapunov stable for Y ; By the definitions, Lyapunov stable set cannot intersect the closure of the basin of some sink when it is not a sink itself.

5 Partial hyperbolicity: the proof of Theorem 4.1

5.1 Lorenz-like singularities

Lemma 5.1. *Let Λ be a non-trivial chain transitive set of $X \in \mathcal{X}^1(M^d)$. We assume*

- *Every singularity in Λ is hyperbolic.*
- *$\mathcal{N}_{\Lambda \setminus \text{Sing}(X)}$ admits a dominated splitting of index i w.r.t the linear Poincaré flow ψ_t .*

Then for every hyperbolic singularity $\sigma \in \Lambda$ with $\text{ind}(\sigma) > i$, $T_\sigma M^d$ admits a dominated splitting $T_\sigma M^d = E^{ss} \oplus E^{cu}$ with respect to the tangent flow Φ_t , where $\dim E^{ss} = i$.

Proof. For a hyperbolic singularity $\sigma \in \Lambda$, since Λ is a non-trivial chain transitive set, one has $W^s(\sigma) \cap \Lambda \setminus \{\sigma\} \neq \emptyset$ and $W^u(\sigma) \cap \Lambda \setminus \{\sigma\} \neq \emptyset$. One can see [12, Lemma 2.6] for more details about the proof. Recall the definition of Λ : the lift of Λ in the sphere bundle as in Subsection 2.1. One has that there is $v \in E^u(\sigma) \cap \tilde{\Lambda}$.

Since $\mathcal{N}_{\Lambda \setminus \text{Sing}(X)}$ admits a dominated splitting of index i with respect to the linear Poincaré flow, $\tilde{\mathcal{N}}_{\tilde{\Lambda}}$ admits a dominated splitting of index i with respect to the extended linear Poincaré flow $\tilde{\psi}_t$. We will consider the negative limit set $\alpha(v)$ with respect to the flow Φ_t^I .

By changing the Riemannian metric in a small neighborhood of σ , without loss of generality, one can assume $E^s(\sigma) \perp E^u(\sigma)$. Thus,

- $\tilde{\mathcal{N}}_{\alpha(v)}$ admits a dominated splitting $\Delta^{cs} \oplus \Delta^{cu}$ of index i w.r.t $\tilde{\psi}_t$ since $\mathcal{N}_{\Lambda \setminus \text{Sing}(X)}$ admits a dominated splitting of index i with respect to ψ_t .
- The hyperbolic splitting on $T_\sigma M^d$ implies that: $\tilde{\mathcal{N}}_{\alpha(v)}$ admits a dominated splitting of index $E^s \oplus F$ of index $\dim E^s$ for some F w.r.t. $\tilde{\psi}_t$, since σ is hyperbolic and $E^s(\sigma) \perp E^u(\sigma)$.

Thus by the properties of dominated splittings, one has $\Delta^{cs}(\alpha(v)) \subset E^s(\sigma)$. Thus, there are $C > 0$ and $\lambda > 0$ such that for any $u \in E^u \cap S_\sigma M^d$ and a splitting $\Delta^{cs} \oplus \Delta^{cu} = \tilde{\mathcal{N}}_{(\sigma, u)}$ with the following property:

- $$\frac{\|\tilde{\psi}_t|_{\Delta^{cs}(u)}\|}{\|\Phi_t|_{\langle u \rangle}\|} \leq Ce^{-\lambda t}, \quad \forall t \geq 0.$$
- $$\frac{\|\tilde{\psi}_t|_{\Delta^{cs}(u)}\|}{m(\tilde{\psi}_t|_{\Delta^{cu}(u)})} \leq Ce^{-\lambda t}, \quad \forall t \geq 0.$$

Then by Lemma 2.12, we know that σ admits a dominated splitting $T_\sigma M^d = E \oplus F$ w.r.t. the tangent flow Φ_t , where $\dim E = i$. Since $\text{ind} \sigma > i$, one can get the splitting as required. \square

Lemma 5.2. *There is a residual subset $\mathcal{G} \subset \mathcal{X}^1(M^3) \setminus \overline{\mathcal{HT}}$ such that for any hyperbolic singularity σ of index 2 of $X \in \mathcal{G}$, if $C(\sigma)$ is non-trivial, then $T_\sigma M^3$ admits a dominated splitting $T_\sigma M^3 = E^{ss} \oplus F$ w.r.t. the tangent flow Φ_t , where $\dim E^{ss} = 1$, E^{ss} is contracting, and $W^{ss}(\sigma) \cap C(\sigma) = \{\sigma\}$.*

Similarly, if $\text{ind}(\sigma) = 1$ and $C(\sigma)$ is non-trivial, then $W^{uu}(\sigma) \cap C(\sigma) = \{\sigma\}$.

Proof. Assume that X satisfies all generic properties in §3. We focus on the case of $\text{ind}(\sigma) = 2$. By Lemma 3.18, there is a sequence of periodic orbit γ_n such that $\lim_{n \rightarrow \infty} \gamma_n = C(\sigma)$ in the Hausdorff topology. By Corollary 3.17, there is $\iota > 0$ such that each γ_n admits an ι -dominated splitting in \mathcal{N}_{γ_n} w.r.t. the linear Poincaré flow. Then by Corollary 2.10, $\mathcal{N}_{C(\sigma) \setminus \text{Sing}(X)}$ admits a dominated splitting of index 1 with respect to the linear Poincaré flow ψ_t . As a corollary of Lemma 5.1, $T_\sigma M$ has dominated splitting $T_\sigma M = E^{ss} \oplus F$ w.r.t. the tangent flow Φ_t . Thus what we need to prove now is $W^{ss}(\sigma) \cap C(\sigma) = \{\sigma\}$. We will prove this by absurd. If this is not true, there is $x_0 \in W^{ss}(\sigma) \cap C(\sigma) \setminus \{\sigma\}$. One also notice that $C(\sigma)$ is Lyapunov stable since $\text{ind}(\sigma) = 2$ by Lemma 3.18. Hence $W^u(\sigma) \subset C(\sigma)$ and $W^u(\sigma)$ is dense in $C(\sigma)$. By changing the Riemannian metric in a small neighborhood, we may assume that $E^{ss}(\sigma)$, $E^{cs}(\sigma)$ and $E^u(\sigma)$ are mutually orthogonal, where $E^{cs}(\sigma) = E^s(\sigma) \cap F$.

Thus, by using the C^1 connecting lemma (Lemma 3.4), there is an arbitrarily small perturbation Y of X such that

- Y has a strong connection with respect to σ : there is $y \in M^3$ such that $y \in W^{ss}(\sigma) \cap W^u(\sigma) \setminus \{\sigma\}$.
- $Y(x) = X(x)$ if x is in a small neighborhood of σ .

By an extra small perturbation, one can assume that Y has the following extra properties:

- Y is linear in a small neighborhood of σ in some local chart;
- $E^{ss}(\sigma, Y)$, $E^{cs}(\sigma, Y)$ and $E^u(\sigma, Y)$ are still mutually orthogonal.

Let P be the plane spanned by E^{ss} and E^u in the local chart. One has that P is locally invariant: there is a neighborhood O_1 of σ such that for any $x \in P \cap O_1$, if $\phi_{[0,t]}^Y(x) \in O_1$, then $\phi_t^Y(x) \in P$. Now for Y , by an extra perturbation, there is a sequence of vector fields Y_n and a smaller neighborhood O_2 of σ such that

- $\lim_{n \rightarrow \infty} Y_n = Y$;
- For each n , $Y_n = Y$ in O_2 ;
- Y_n has a periodic orbit γ_n such that $\gamma_n \cap O_2 \subset P$ and $\lim_{n \rightarrow \infty} \gamma_n = \text{Orb}(y, Y)$.

By Corollary 3.17, one has

- There are $\iota = \iota(X)$ such that \mathcal{N}_{γ_n} has an ι -dominated splitting $\Delta^{cs} \oplus \Delta^{cu}$ of index 1 with respect to the linear Poincaré flow $\psi_t^{Y_n}$.

Thus, $\mathcal{N}_{\text{Orb}(y, Y)}$ admits an ι -dominated splitting w.r.t. ψ_t^Y . Let $\Gamma_Y = \text{Orb}(y, Y) \cup \sigma_Y$. It is a compact invariant set. Over the lift $\widetilde{\Gamma}_Y$ (see Subsection 2.1), there exists an dominated splitting $\widetilde{\mathcal{N}}_{\widetilde{\Gamma}_Y} = \Delta^{cs} \oplus \Delta^{cu}$ w.r.t. $\widetilde{\psi}_t$ such that

- For $v^u \in E^u(\sigma) \cap \tilde{\Gamma}_Y$, one has $\Delta^{cs}(v^u) = E^{ss}$ and $\Delta^{cu}(v^u) = E^{cs}$.
- For $v^{ss} \in E^{ss}(\sigma) \cap \tilde{\Gamma}_Y$, one has $\Delta^{cs}(v^{ss}) = E^{cs}$ and $\Delta^{cu}(v^{ss}) = E^u$.

Then by following the analysis in the proof of Lemma 4.3 in [24, page 255-256], we will get a contradiction. \square

Corollary 5.3. *There is a residual subset $\mathcal{G} \subset \mathcal{X}^1(M^3) \setminus \overline{\mathcal{HT}}$ such that if a chain recurrent class contains singularities, then all the singularities in the chain recurrent class have the same index.*

Proof. Let $\mathcal{G} = \mathcal{G}_0 \setminus \overline{\mathcal{HT}}$, where \mathcal{G}_0 is as in the end of Subsection 3.3. We will prove this corollary by absurd. If it's not true, then there is $X \in \mathcal{G}$ and a chain recurrent class C of X such that C contains singularities of different indices. Thus, one can assume that C contains two singularities σ_1 and σ_2 satisfying $\text{ind}(\sigma_1) = 1$ and $\text{ind}(\sigma_2) = 2$. Since by Lemma 5.2, $T_{\sigma_1}M^3 = E^{cs} \oplus E^{uu}$ is a dominated splitting w.r.t. Φ_t , where $\dim E^{uu} = 1$, and for the corresponding strong unstable manifold $W^{uu}(\sigma_1) \cap C = \{\sigma_1\}$. By Lemma 3.18, C is Lyapunov stable since C contains σ_2 . As a corollary, one has $W^{uu}(\sigma_1) \subset C$. This gives us a contradiction. \square

Corollary 5.4. *There is a residual subset $\mathcal{G} \subset \mathcal{X}^1(M^3) \setminus \overline{\mathcal{HT}}$ such that for any hyperbolic singularity σ of index 2 of $X \in \mathcal{G}$, there is a C^1 neighborhood \mathcal{U} of X such that for any $Y \in \mathcal{U}$ and for any singularity $\rho \in C(\sigma)$, one has*

- $\text{ind}(\rho) = 2$ and $\rho_Y \in C(\sigma_Y)$.
- $T_\rho M^3 = E^{ss} \oplus E^{cu}$ is a dominated splitting w.r.t. Φ_t , where $\dim E^{ss} = 1$.
- For the corresponding stable manifolds of E^{ss} , one has $W^{ss}(\rho_Y, Y) \cap C(\sigma_Y) = \{\rho_Y\}$

Proof. This is true just because that X is C^1 generic and the continuous property of the local strong stable manifolds. \square

Furthermore, we have

Theorem 5.5. *For a generic $X \in \mathcal{X}^1(M^3) \setminus \overline{\mathcal{HT}}$, and a hyperbolic singularity σ of index 2, if $C(\sigma)$ is non trivial, then σ is Lorenz-like for X , i.e., the eigenvalues $\lambda_1, \lambda_2, \lambda_3$ of $DX(\sigma)$ are all real and satisfy*

$$\lambda_1 < \lambda_2 < 0 < -\lambda_2 < \lambda_3. \quad (*)$$

Proof. First by Lemma 5.2, for the three eigenvalues $\lambda_1, \lambda_2, \lambda_3$ of $DX(\sigma)$, they are all real and

$$\lambda_1 < \lambda_2 < 0 < \lambda_3.$$

What's left is to prove that $\lambda_2 + \lambda_3 > 0$. The three corresponding eigenspaces are denoted by $E^{ss}(\sigma)$, $E^{cs}(\sigma)$ and $E^u(\sigma)$. By changing the Riemannian metric in a small neighborhood of σ , we can assume that they are mutually orthogonal. We will prove this by absurd, i.e., we assume that $\lambda_2 + \lambda_3 \leq 0$. Since X is C^1 generic, by Lemma 3.18 one has $\lambda_2 + \lambda_3 < 0$. Moreover,

- $W^u(\sigma) \subset C(\sigma)$ and $W^u(\sigma)$ is dense in $C(\sigma)$ by Lemma 3.18.
- $W^s(\sigma) \cap C(\sigma) \setminus \{\sigma\} \neq \emptyset$ since $C(\sigma)$ is non-trivial.

By using Lemma 3.4 (the C^1 connecting lemma), for any C^1 neighborhood \mathcal{U} of X , there is $Y \in \mathcal{U}$ such that

- Y has a homoclinic orbit Γ associated to σ_Y .
- For the three eigenvalues $\lambda_1^Y < \lambda_2^Y < 0 < \lambda_3^Y$ of $DY(\sigma_Y)$, one still has $\lambda_2^Y + \lambda_3^Y < 0$.
- $W^{ss}(\sigma_Y) \cap C(\sigma_Y) = \{\sigma_Y\}$ by Corollary 5.4.

By simple perturbations, there is a sequence of vector fields Y_n such that

- $\lim_{n \rightarrow \infty} Y_n = Y$.
- Each Y_n has a hyperbolic periodic orbit γ_n such that $\lim_{n \rightarrow \infty} \gamma_n = \Gamma \cup \sigma$.

Under our assumption, $\{\gamma_n\}$ could be sinks or saddles. Let $C > 0$, $\eta > 0$, $\iota > 0$ be as in Lemma 2.11. By Corollary 3.17 \mathcal{N}_{γ_n} admits an ι -dominated splitting of index 1 with respect to $\psi_t^{Y_n}$ for n large enough.

As a corollary, \mathcal{N}_Γ admits an ι -dominated splitting w.r.t. ψ_t^Y . Thus, $\widetilde{\mathcal{N}}_{\Gamma \cup \sigma}$ admits an ι -dominated splitting $\widetilde{\mathcal{N}}_{\Gamma \cup \sigma} = \Delta^{cs} \oplus \Delta^{cu}$ w.r.t. $\widetilde{\psi}_t^Y$.

Claim. For every $v \in T_\sigma M^3 \cap (\widetilde{\Gamma \cup \sigma})$, one has $\Delta^{cs}(v) = E^{ss}(\sigma)$.

Proof. For each $v \in T_\sigma M^3$, if $v \in \widetilde{\Gamma \cup \sigma}$, then $v \in E^{cs}(\sigma)$ or $v \in E^u(\sigma)$. Since $E^{ss}(\sigma)$, $E^{cs}(\sigma)$ and $E^u(\sigma)$ are mutually orthogonal, one has

- if $v \in E^{cs}(\sigma)$, since $\widetilde{\mathcal{N}}_v = E^{ss}(\sigma) \oplus E^u(\sigma)$ is a dominated splitting w.r.t. $\widetilde{\psi}_t$, one has $\Delta^{cs}(v) = E^{ss}(\sigma)$;
- if $v \in E^u(\sigma)$, since $\widetilde{\mathcal{N}}_v = E^{ss}(\sigma) \oplus E^{cs}(\sigma)$ is a dominated splitting w.r.t. $\widetilde{\psi}_t$, one has $\Delta^{cs}(v) = E^{ss}(\sigma)$.

□

Since the unique ergodic measure is supported on $\{\sigma\}$ for $\phi_t|_{\Gamma \cup \sigma}$, one has that there are $C > 0$ and $\lambda > 0$ such that for any $t > 0$ and any $(x, v) \in \widetilde{\Gamma \cup \sigma}$,

$$\frac{\|\widetilde{\psi}_t|_{\Delta^{cs}(x,v)}\|}{\|\Phi_t|_{\langle v \rangle}\|} \leq Ce^{-\lambda t}.$$

By Lemma 2.12, $\Gamma \cup \sigma$ admits a dominated splitting $T_{\Gamma \cup \sigma} M^3 = E \oplus F$ w.r.t. the tangent flow Φ_t , where $\dim E = 1$. Thus $E(\sigma) = E^{ss}(\sigma)$ by the uniqueness of dominated splittings. Since the unique ergodic measure is supported on $\{\sigma\}$, one has that E is uniformly contracting. Thus γ_n is also $(C, \eta, d, \mathcal{N})$ -contracting at the period w.r.t. $\psi_t^{Y_n}$ for some $C > 0$, $d > 0$ and $\eta \in (0, -(\lambda_2 + \lambda_3))$ which depends on X since γ_n stays in a small neighborhood of the singularity for most time. This also gives a contradiction by Lemma 3.16.

□

5.2 Liao's shadowing lemma for ψ_t^* and Liao's sifting lemma

Definition 5.6. Let $\eta > 0$ and $T > 0$. For any $x \in M^d \setminus \text{Sing}(X)$ and $T_0 > T$, the orbit arc $\phi_{[0, T_0]}(x)$ is called (η, T) - ψ_t^* -quasi hyperbolic with respect to a direct sum splitting $\mathcal{N}_x = E(x) \oplus F(x)$ if there is a partition

$$0 = t_0 < t_1 < \dots < t_l = T, \quad t_{i+1} - t_i \leq T,$$

such that for $k = 0, 1, \dots, l-1$, we have

$$\begin{aligned} \prod_{i=0}^{k-1} \|\psi_{t_{i+1}-t_i}^*|_{\psi_{t_i}(E(x))}\| &\leq e^{-\eta t_k}, \\ \prod_{i=k}^{l-1} m(\psi_{t_{i+1}-t_i}^*|_{\psi_{t_i}(F(x))}) &\geq e^{\eta(t_l-t_k)}, \\ \frac{\|\psi_{t_{k+1}-t_k}^*|_{\psi_{t_k}(E(x))}\|}{m(\psi_{t_{k+1}-t_k}^*|_{\psi_{t_k}(F(x))})} &\leq e^{-\eta(t_{k+1}-t_k)}. \end{aligned}$$

Remark. The third inequality is usually satisfied in an invariant set with a T^* -dominated splitting in the normal bundle with respect to the linear Poincaré flow.

Note that this definition is similar to the usual quasi hyperbolic orbit arc for linear Poincaré flow, while the only difference is that now we use the scaled linear Poincaré flow ψ_t^* instead of linear Poincaré flow ψ_t . Let d_T be the metric in TM^d induced by the Riemannian metric. For $x, y \in M^d$ and two linear subspaces $E(x)$ and $F(y)$, one defines

$$\tilde{d}(E(x), F(y)) = \sup_{\{u \in E(x), |u|=1\}} \inf_{\{v \in F(y), |v|=1\}} \{d_T(u, v)\}.$$

Theorem 5.7. Let $X \in \mathcal{X}^1(M^d)$, $\Lambda \subset M^d \setminus \text{Sing}(X)$ be a compact set, and $\eta > 0, T > 1$. For any $\varepsilon > 0$ there exists $\delta > 0$, such that for any (η, T) - ψ_t^* -quasi hyperbolic orbit arc $\phi_{[0, T]}(x)$ with respect to some direct sum splitting $\mathcal{N}_x = E(x) \oplus F(x)$ satisfying $x, \phi_T(x) \in \Lambda$ and $\tilde{d}(E(x), \psi_T(E(x))) \leq \delta$, there exists a point $p \in M^d$ and a C^1 strictly increasing function $\theta : [0, T] \rightarrow \mathbb{R}$ such that

- $\theta(0) = 0$ and $1 - \varepsilon < \theta'(t) < 1 + \varepsilon$;
- p is periodic: $\phi_{\theta(T)}(p) = p$;
- $d(\phi_t(x), \phi_{\theta(t)}(p)) \leq \varepsilon |X(\phi_t(x))|$, $t \in [0, T]$.
- There is a direct-sum splitting $\mathcal{N}_p = E(p) \oplus F(p)$ such that $\psi_{\theta(T)}^*(E(p)) = E(p)$, $\psi_{\theta(T)}^*(F(p)) = F(p)$, and for any $t \in [0, T]$,

$$\tilde{d}(\psi_t^*(E(x)), \psi_{\theta(t)}^*(E(p))) \leq \varepsilon,$$

$$\tilde{d}(\psi_t^*(F(x)), \psi_{\theta(t)}^*(F(p))) \leq \varepsilon.$$

Remark. If we consider an invariant set with a dominated splitting in the normal bundle w.r.t. the linear Poincaré flow, we can replace $\tilde{d}(E(x), \psi_T(E(x))) \leq \delta$ by $d(x, \phi_T(x)) < \delta$.

Note that in this version of shadowing lemma, we only need that the head and tail of orbit arc are far from singularities, while other part of the orbit arc can approximate singularities. This enables us to deal with some problems where regular orbits approximate singularities.

We also need Liao's sifting lemma [26, 27], whose aim is to find quasi-hyperbolic orbit segments. One can see [62] for a proof.

Lemma 5.8. Let $\phi_t : \Lambda \rightarrow \Lambda$ be a continuous flow on a compact metric space Λ and $f : \Lambda \rightarrow \mathbb{R}$ a continuous function. Let $\eta_2 > \eta_1 > 0$ and $T > 0$. Assume that

- there is $b \in \Lambda$ such that for any $n \in \mathbb{N}$,

$$\sum_{i=0}^{n-1} f(\phi_{iT}(b)) \geq 0;$$

- for any $c \in \Lambda$ verifying for any $n \in \mathbb{N}$,

$$\sum_{i=0}^{n-1} f(\phi_{iT}(c)) \geq -n\eta_1,$$

there is $g \in \omega(c)$ such that for any $n \in \mathbb{N}$,

$$\sum_{i=0}^{n-1} f(\phi_{iT}(g)) \leq -n\eta_2.$$

Then for any η_3, η_4 verifying $\eta_2 > \eta_3 > \eta_4 > \eta_1$, for any $k \in \mathbb{N}$, there is y in the positive orbit of x and integers $0 = n_0 < n_1 < \dots < n_k$ such that for each integer $i \in [0, k-1]$, for any integer $m \in [1, n_{i+1} - n_i]$ one has

$$\sum_{j=0}^{m-1} f(\phi_{jT}(\phi_{n_i T}(y))) \leq -m\eta_4,$$

$$\sum_{j=m-1}^{n_{i+1}-n_i-1} f(\phi_{jT}(\phi_{(n_i+m-1)T})) \geq -(n_{i+1} - n_i - m + 1)\eta_3.$$

We need the following fundamental lemma to prove hyperbolicity for compact sets.

Lemma 5.9. *Let $\phi_t : \Lambda \rightarrow \Lambda$ be a continuous flow on a compact metric space Λ and $f : \Lambda \rightarrow \mathbb{R}$ a continuous function. Given $T > 0$, if for any $x \in \Lambda$, there is $n_x \in \mathbb{N}$ such that*

$$\sum_{i=0}^{n_x-1} f(\phi_{iT}(x)) < 0,$$

then there are $C \geq 0$ and $\lambda < 0$ such that for any $x \in \Lambda$ and any $n \in \mathbb{N}$, one has

$$\sum_{i=0}^{n-1} f(\phi_{iT}(x)) \leq C + n\lambda.$$

5.3 Proof of Proposition 4.7

Now we can give the proof of Proposition 4.7.

Proof of Proposition 4.7. For proving this proposition, one assumes that the second case of the conclusion cannot occur.

By changing the Riemannian metric in a small neighborhood of singularities, one can assume that $E^{ss}(\sigma) \perp E^{cu}(\sigma)$ for any singularity $\sigma \in \Lambda$. There is $T^* > 0$ such that

- For any $\sigma \in \Lambda$ and any unit vector $v \in E^{cu}(\sigma)$, one has

$$\frac{\|\Phi_{T^*}|_{E^{ss}}\|}{|\Phi_{T^*}(v)|} \leq \frac{1}{4}.$$

- $\mathcal{N}_{\Lambda \setminus \text{Sing}(X)} = \Delta^{cs} \oplus \Delta^{cu}$ is a T^* -dominated splitting w.r.t. ψ_t .

We consider $\tilde{\Lambda}$: the lift of Λ in the sphere bundle as in Subsection 2.1. By considering the dynamics of χ_t , one has $\tilde{\mathcal{N}}_{\tilde{\Lambda}} = \tilde{\Delta}^{cs} \oplus \tilde{\Delta}^{cu}$ is a T^* -dominated splitting of index i w.r.t. $\tilde{\psi}_t = \text{proj}_2(\chi_t)$ verifying the following property: $\Delta^{cs}(x) = \tilde{\Delta}^{cs}(X(x)/|X(x)|)$ and $\Delta^{cu}(x) = \tilde{\Delta}^{cu}(X(x)/|X(x)|)$ for any regular point $x \in \Lambda$.

Since $W^{ss}(\sigma) \cap \Lambda = \{\sigma\}$, one has if $v \in \tilde{\Lambda} \cap T_\sigma M^d$, then $v \in E^{cu}(\sigma)$. On $\tilde{\Lambda}$, one defines the function $\tilde{\xi}$ by

$$\begin{aligned} \tilde{\xi} : \tilde{\Lambda} &\rightarrow \mathbb{R}, \\ v &\mapsto \log \|\tilde{\psi}_{T^*}|_{\tilde{\Delta}^{cs}(v)}\| - \log \|\Phi_{T^*}(v)\|. \end{aligned}$$

Since $\tilde{\Delta}^{cs}$ is a continuous bundle, $\tilde{\xi}$ is a continuous function.

On $\Lambda \setminus \text{Sing}(X)$, one can define the function ξ by

$$\begin{aligned} \xi : \Lambda \setminus \text{Sing}(X) &\rightarrow \mathbb{R}, \\ x &\mapsto \log \|\psi_{T^*}|_{\Delta^{cs}(x)}\| - \log \|\Phi_{T^*}|_{\langle X(x) \rangle}\|. \end{aligned}$$

By the definitions, for every regular point $x \in \Lambda$, $\xi(x) = \tilde{\xi}(X(x)/|X(x)|)$. $\tilde{\xi}$ is defined on a compact set and ξ is defined on a non-compact set.

Claim. *There is $C > 1$ and $0 < \lambda < 1$ such that for any $v \in \tilde{\Lambda}$ and $n \in \mathbb{N}$, one has*

$$\frac{\|\tilde{\psi}_{nT^*}|_{\tilde{\Delta}^{cs}(v)}\|}{|\Phi_{nT^*}(v)|} \leq C\lambda^n.$$

Proof of the Claim. The claim is equivalent to the following statement: There are $C > 1$ and $0 < \lambda < 1$ such that for any $v \in \tilde{\Lambda}$ and $n \in \mathbb{N}$,

$$\sum_{i=0}^{n-1} \tilde{\xi}(\Phi_{iT^*}^I(v)) \leq \log C + n \log \lambda.$$

If the claim is not true, by Lemma 5.9, for any $n \in \mathbb{N}$, there is $v_n \in \tilde{\Lambda}$ such that for any integer $\ell \in [1, n]$, one has

$$\sum_{j=0}^{\ell-1} \tilde{\xi}(\Phi_{iT^*}^I(v_n)) \geq 0.$$

Let $b \in \tilde{\Lambda}$ be an accumulation point of $\{v_n\}$. Then for any $n \in \mathbb{N}$,

$$\sum_{i=0}^{n-1} \tilde{\xi}(\Phi_{iT^*}^I(b)) \geq 0.$$

Since we assume that $E^{ss}(\sigma) \perp E^{cu}(\sigma)$, one has for any $v \in T_\sigma M^d \cap \tilde{\Lambda}$, $\tilde{\Delta}^{cs}(v) = E^{ss}(\sigma)$ and $v \in E^{cu}$. As a corollary, one has for any $n \in \mathbb{N}$,

$$\sum_{i=0}^{n-1} \tilde{\xi}(\Phi_{iT^*}^I(v)) \leq -n \log 4.$$

For every point $x \in \Lambda$, by assumptions, $\omega(x)$ contains a hyperbolic singularity. Thus for its lift $\tilde{\Lambda}$, for any point $c \in \tilde{\Lambda}$, there is a singularity $\sigma \in \Lambda$ and a unit vector $v \in E^{cu}(\sigma)$ such that

$v \in \omega(c)$. Thus for the function $\tilde{\xi}$ and the flow ϕ_t^I , the conditions of Lemma 5.8 (Liao's sifting lemma) are satisfied.

For any four numbers $\lambda_1, \lambda_2, \lambda_3$ and λ_4 satisfying $-\log 4/2 < \lambda_1 < \lambda_2 < \lambda_3 < \lambda_4 < 0$. Let

$$\tilde{\Lambda}_{\lambda_2} = \{v \in \tilde{\Lambda} : \tilde{\xi}(\Phi_{-T^*}^I(v)) \geq \lambda_2\}.$$

Since there are only finite many singularities, there is $\varepsilon_0 = \varepsilon_0(\lambda_2) > 0$, such that for any singularity $\sigma \in \Lambda$, and any unit vector $v \in T_\sigma M^d \cap \tilde{\Lambda}$, $d_T(v, \tilde{\Lambda}_{\lambda_2}) \geq \varepsilon_0$. By Lemma 5.8, for any $k \in \mathbb{N}$, there is u in the positive orbit of b and integers $0 = n_0 < n_1 < \dots < n_k$ such that for each integer $\ell \in [0, k-1]$, for any integer $m \in [1, n_{\ell+1} - n_\ell]$ one has

$$\begin{aligned} \sum_{j=0}^{m-1} \tilde{\xi}(\Phi_{jT^*}^I(\Phi_{n_i T^*}^I(u))) &\leq m\lambda_3, \\ \sum_{j=m-1}^{n_{\ell+1}-n_\ell-1} \tilde{\xi}(\Phi_{jT^*}^I(\Phi_{(n_\ell+m-1)T^*}^I(u))) &\geq (n_{\ell+1} - n_\ell - m + 1)\lambda_2. \end{aligned}$$

Thus $\Phi_{n_\ell T^*}^I(y) \in \tilde{\Lambda}_{\lambda_2}$ for $1 \leq \ell \leq k$. Assume that $b = X(x)/|X(x)|$ and $y = \Phi_{t_0}^I(b)$. Thus,

$$d(\phi_{n_i T^*}(\phi_{t_0}(x)), \text{Sing}(X)) \geq \varepsilon_0.$$

For any $\varepsilon > 0$ small enough such that: for any regular point $\beta \in \Lambda$, for any point $\theta \in B(\beta, \varepsilon|X(\beta)|)$, for any $T' \in [(1-\varepsilon)T^*, (1+\varepsilon)T^*]$, for any two subspaces $G(\beta) \subset \mathcal{N}_\beta$ and $G(\theta) \subset \mathcal{N}_\theta$ satisfying $\tilde{d}(G(\beta), G(\theta)) < \varepsilon$, one has

$$\lambda_1 - \lambda_2 \leq \log \|\psi_{T^*}^*|_{G(\beta)}\| - \log \|\psi_{T'}^*|_{G(\theta)}\| \leq \lambda_4 - \lambda_3.$$

$$\max\left\{\frac{\|\psi_{T'}^*|_{G(\theta)}\|}{\|\psi_{T^*}^*|_{G(\beta)}\|}, \frac{m(\psi_{T^*}^*|_{G(\beta)})}{m(\psi_{T'}^*|_{G(\theta)})}\right\} \leq \frac{\sqrt{2}}{2}.$$

For this $\varepsilon > 0$, there is $\delta = \delta(\varepsilon)$ as in Theorem 5.7 (Liao's shadowing). There is $k_\delta \in \mathbb{N}$ such that for any k_δ points $\{x_1, x_2, \dots, x_{k_\delta}\} \subset \tilde{\Lambda}_{\lambda_2}$, there are $1 \leq i_1 < i_2 \leq k_\delta$ such that $d(x_{i_1}, x_{i_2}) < \delta$. For this k_δ , there are $n_1 < n_2 < \dots < n_{k_\delta}$ and a point $y' \in \text{Orb}^+(x)$ such that for the function ξ and $0 \leq \ell \leq k_\delta - 1$, one has

$$\begin{aligned} \sum_{j=0}^{m-1} \xi(\phi_{jT^*}(\phi_{n_\ell T^*}(y'))) &\leq m\lambda_3, \\ \sum_{j=m-1}^{n_{\ell+1}-n_\ell-1} \xi(\phi_{jT^*}(\phi_{(n_\ell+m-1)T^*}(y'))) &\geq (n_{\ell+1} - n_\ell - m + 1)\lambda_2. \end{aligned}$$

Let $y_\ell = \phi_{n_\ell T^*}(y')$. By the dominated properties, one has for each y_ℓ , for any integer $m \in [1, n_{i+1} - n_i]$, one has

$$\begin{aligned} \prod_{j=0}^{m-1} \|\psi_{T^*}^*|_{\Delta^{cs}}(\phi_{jT^*}(y_\ell))\| &\leq e^{m\lambda_3}, \\ \prod_{j=m-1}^{n_{\ell+1}-n_\ell-1} \|\psi_{-T^*}^*|_{\Delta^{cu}}(\phi_{j+1}(y_\ell))\| &\leq \left(\frac{1}{4}\right)^{n_{\ell+1}-n_\ell-m} e^{-(n_{\ell+1}-n_\ell-m)\lambda_2}. \end{aligned}$$

Let $\eta = \min\{-\lambda_3, \log 4 + \lambda_2\}$. One has that

- $\phi_{[0, (n_{\ell+1} - n_{\ell})T^*]}(y_{\ell})$ is an (η, T^*) - ψ_t^* -quasi hyperbolic strings.
- $d(y_i, \text{Sing}(X)) > \varepsilon$.

By the choice of k_{δ} , there are y_{α} and y_{β} such that $d(y_{\alpha}, y_{\beta}) < \delta$. Thus by Theorem 5.7, the orbit segment from y_{α} to y_{β} can be shadowed: there is a periodic orbit P_{ε} with period $\tau(P_{\varepsilon})$ and $p_{\varepsilon} \in P_{\varepsilon}$ and a monotone increasing function $\theta_{\varepsilon}(t)$ such that

- $d(\phi_{\theta_{\varepsilon}(t)}(p_{\varepsilon}), \phi_t(y_{\alpha})) < \varepsilon |X(\phi_t(y_{\alpha}))|$ for any $0 \leq t \leq (n_{\beta} - n_{\alpha})T^*$.
- $1 - \varepsilon \leq \theta'_{\varepsilon}(t) \leq 1 + \varepsilon$ and $\theta_{\varepsilon}((n_{\beta} - n_{\alpha})T^*) = \tau(P_{\varepsilon})$.
- There is a direct-sum splitting $\mathcal{N}_{p_{\varepsilon}} = E(p_{\varepsilon}) \oplus F(p_{\varepsilon})$ such that $\psi_{\theta((n_{\beta} - n_{\alpha})T^*)}^*(E(p_{\varepsilon})) = E(p_{\varepsilon})$, $\psi_{\theta((n_{\beta} - n_{\alpha})T^*)}^*(F(p_{\varepsilon})) = F(p_{\varepsilon})$, and for any $t \in [0, (n_{\beta} - n_{\alpha})T^*]$,

$$\tilde{d}(\psi_t^*(E(y_{\alpha})), \psi_{\theta(t)}^*(E(p_{\varepsilon}))) \leq \varepsilon,$$

$$\tilde{d}(\psi_t^*(F(y_{\alpha})), \psi_{\theta(t)}^*(F(p_{\varepsilon}))) \leq \varepsilon.$$

Let $\eta' = \min\{-\lambda_4, \log 2 + \lambda_1\}$ and $T' = (1 + \varepsilon)T$. By the choosing of ε , one has that P_{ε} is an (η', T') - ψ_t^* -quasi hyperbolic string. Take a sequence $\{1/q\}_{q \in \mathbb{N}}$. For each q , there is a periodic orbit $P_{1/q}$ with period $\tau(P_{1/q})$ such that

- $\limsup_{q \rightarrow \infty} P_{1/q} \subset \Lambda$.
- There is $p_{1/q} \in P_{1/q}$ such that $d(p_{1/q}, \text{Sing}(X)) \geq \varepsilon_0$, $p_{1/q}$ is $(1, \eta', T', E)$ - ψ_t^* -contracting and $(1, \eta', T', F)$ - ψ_t^* -expanding.
- $P_{1/q}$ admits a T^* -dominated splitting w.r.t. ψ_t^* .

Without loss of generality, one can assume that $\{p_{1/q}\}_{q \in \mathbb{N}}$ converges. By Lemma 2.17⁴, for any two $w, m \in \mathbb{N}$ large enough, one has

$$W^s(P_{1/w}) \pitchfork W^u(P_{1/m}) \neq \emptyset, \quad W^u(P_{1/w}) \pitchfork W^s(P_{1/m}) \neq \emptyset.$$

In other words, they are homoclinically related. Thus for w large enough, one has $\Lambda \cap H(P_{1/w}) \neq \emptyset$. Moreover one has

$$\frac{1}{[\tau(P_{1/w})/T']^{\lfloor \tau(P_{1/w})/T' \rfloor - 1}} \sum_{i=0}^{\lfloor \tau(P_{1/w})/T' \rfloor - 1} \log \|\psi_{T'}|_{\mathcal{N}^s(p_{1/w})}\| \geq \lambda_1.$$

By the arbitrary property of λ_1 , we have the second case. □

By Lemma 2.12, and the claim above, one has Λ admits a dominated splitting $T_{\Lambda}M = E \oplus F$ w.r.t. the tangent flow, where $\dim E = i$ and $X(x) \subset F(x)$ for any regular point $x \in \Lambda$. We assume that it is a T -dominated splitting. We define the function $f : \Lambda \rightarrow \mathbb{R}$ by $f(x) = \log \|\Phi_T|_{E(x)}\|$.

Claim.

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{\ell=0}^{n-1} f(\phi_{\ell T}(x)) < 0, \quad \forall x \in \Lambda.$$

⁴This is the part that we need the existence of central plaques as in Section 2.

Proof. For every point $x \in \Lambda$, there are two cases: either $\omega(x) \subset \text{Sing}(X)$, or $\omega(x)$ contains a regular point $a \in \Lambda$. In the first case, since every singularity in Λ admits a partially hyperbolic splitting, one has that the claim is true. Now we consider the second case: $\omega(x)$ contains a regular point a . We fix a neighborhood U_a of a such that for any $z, y \in U_a$, one has

$$\frac{1}{2} \leq \frac{|X(z)|}{|X(y)|} \leq 2.$$

Let $f_0(z) = \log \|\Phi_T|_{E(z)}\| - \log \|\Phi_T|_{\langle X(z) \rangle}\|$ for every regular point $z \in \Lambda$. Since $T_\Lambda M^d = E \oplus F$ is a dominated splitting and $X(z) \subset F(z)$ for every regular point $z \in \Lambda$, one has $f_0(z) \leq -\log 2$ for any regular point z . There is $x_0 \in \text{Orb}^+(x)$ such that $x_0 \in U_a$. Since $a \in \omega(x)$, there is a sequence of times $\{t_n\}_{n \in \mathbb{N}}$ such that $\lim_{n \rightarrow \infty} \phi_{t_n}(x_0) = a$ and $\lim_{n \rightarrow \infty} t_n = \infty$.

For n large enough, we assume that $t_n = kT + t$, where $t \in [0, T]$. Thus we have

$$\frac{1}{k} \sum_{\ell=0}^{k-1} f(\phi_{\ell T}(x_0)) \leq -\log 2 + \frac{1}{k} (\log \|\Phi_T|_{\langle X(\phi_{kT}(x_0)) \rangle}\| - \log \|\Phi_T|_{X(x_0)}\|).$$

Since $k \rightarrow \infty$ as $n \rightarrow \infty$, one has

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{\ell=0}^{n-1} f(\phi_{\ell T}(x_0)) < 0.$$

Thus the same inequality holds for x . □

By the above claim, one has $\sum_{\ell=0}^{n-1} f(\phi_{\ell T}(x)) < 0$ for any $x \in \Lambda$. By Lemma 5.9, we have that E is uniformly contracting. This ends the proof of this proposition. □

6 Perturbations in partially hyperbolic quasi attractor

In this section, we will manage to prove Theorem 4.2. To prove Theorem 4.2, we notice that [12, Theorem B] proved that⁵

Theorem 6.1. *For C^1 generic $X \in \mathcal{X}^1(M^3)$, for the non-trivial chain recurrent class $C(\sigma)$ of some singularity σ , if $C(\sigma)$ admits a dominated splitting $T_{C(\sigma)}M^3 = E \oplus F$ w.r.t. the tangent flow, then $C(\sigma)$ admits a partially hyperbolic splitting; more precisely, $\text{ind}(\sigma) = 2$ iff $\dim E = 1$ and E is contracting.*

Thus, what we need to prove is

Theorem 6.2. *For C^1 generic $X \in \mathcal{X}^1(M^3)$, for the non-trivial chain recurrent class $C(\sigma)$ of some singularity σ , if $C(\sigma)$ admits a dominated splitting $T_{C(\sigma)}M^3 = E^s \oplus F$ w.r.t. the tangent flow, where E^s is one-dimensional and contracting, then $C(\sigma)$ is a homoclinic class.*

6.1 Cross sections of partially hyperbolic Lyapunov stable chain recurrent classes

Our aim in this subsection is to prove Proposition 4.8. We also need some other lemmas to get an open set of vector fields, which guarantee that we can do a perturbation in that open set.

For a singularity σ of $X \in \mathcal{X}^1(M^3)$, recall that σ is *Lorenz-like* if the eigenvalues $\lambda_1, \lambda_2, \lambda_3$ of $DX(\sigma)$ satisfy

$$\lambda_1 < \lambda_2 < 0 < -\lambda_2 < \lambda_3.$$

Since we not only need to consider $C(\sigma)$, but also its continuation, we give the notion of *pseudo-Lorenz* to characterize its properties.

⁵The statement is a little bit stronger than [12]. But the proof is contained there.

Definition 6.3. A compact invariant set Λ of $X \in \mathcal{X}^1(M^3)$ is pseudo-Lorenz if

- Λ is Lyapunov stable.
- Λ has a partially hyperbolic splitting $T_\Lambda M = E^{ss} \oplus E^{cu}$ with respect to Φ_t , where $\dim E^{ss} = 1$.
- Every singularity in Λ is Lorenz-like.
- For every singularity $\sigma \in \Lambda$, $W^{ss}(\sigma) \cap \Lambda = \{\sigma\}$.

This notion is a generalization of singular hyperbolic transitive attractor [36].

Proposition 6.4. For C^1 generic $X \in \mathcal{X}^1(M^3)$ and $\sigma \in \text{Sing}(X)$, if $C(\sigma)$ is pseudo-Lorenz, then there is a C^1 neighborhood \mathcal{U} of X such that for any $Y \in \mathcal{U}$,

- $C(\sigma_Y, Y)$ admits a partially hyperbolic splitting $T_{C(\sigma_Y, Y)} M^3 = E^{ss} \oplus E^{cu}$, where $\dim E^{ss} = 1$.
- Every singularity in $C(\sigma_Y, Y)$ is the continuation of some singularity in $C(\sigma)$, and Lorenz-like.
- For every singularity $\rho \in C(\sigma_Y, Y)$, one has $W^{ss}(\rho) \cap C(\sigma_Y, Y) = \{\rho\}$.

Moreover, if Y is weak Kupka-Smale, then $C(\sigma_Y, Y)$ is Lyapunov stable and hence it is pseudo-Lorenz.

Proof. Assume that X satisfies the generic properties in Subsection 3.3. We will prove that such an X satisfies the proposition.

Since X is Kupka-Smale, X has only finitely many hyperbolic singularities $\{\sigma_i\}_{i=1}^m$. Moreover, there is a neighborhood \mathcal{U}_0 such that for any $Y \in \mathcal{U}_0$, $\text{Sing}(Y) = \{\sigma_{i,Y}\}_{i=1}^m$, where $\sigma_{i,Y}$ is the continuations of σ_i . By Lemma 3.12, there is a C^1 neighborhood $\mathcal{U}_1 \subset \mathcal{U}_0$ such that for any $Y \in \mathcal{U}_1$, such that for any $1 \leq i, j \leq m$, we have $C(\sigma_{i,Y}, Y) = C(\sigma_{j,Y}, Y)$ iff $C(\sigma_i, X) = C(\sigma_j, X)$.

Given a pseudo-Lorenz set $C(\sigma)$ of X , we assume that $C(\sigma) \cap \text{Sing}(X) = \{\sigma_1, \sigma_2, \dots, \sigma_k\}$. Since the eigenvalues of $\sigma_{i,Y}$ vary continuously with Y ([46, 2.18 Proposition]) and $C(\sigma)$ is pseudo-Lorenz, there exists a neighborhood $\mathcal{U}_2 \subset \mathcal{U}_1$ such that for any $Y \in \mathcal{U}_2$, the index of every singularity in $C(\sigma_Y, Y)$ is 2, $C(\sigma_Y, Y) \cap \text{Sing}(Y) = \{\sigma_{1,Y}, \sigma_{2,Y}, \dots, \sigma_{k,Y}\}$ and every singularity in $C(\sigma_Y, Y)$ is Lorenz-like. Since $C(\sigma_Y, Y)$ is upper-continuous with Y , by shrinking \mathcal{U}_2 , we have that for every $Y \in \mathcal{U}_2$, $C(\sigma_Y, Y)$ has a partially hyperbolic splitting with respect to the tangent flow Φ_t^Y :

$$T_{C(\sigma_Y, Y)} M = E_Y^{ss} \oplus E_Y^{cu},$$

such that $\dim E_Y^{ss} = 1$ and E_Y^{ss} is uniformly contracting.

We have that, by shrinking \mathcal{U}_2 if necessary, for any $Y \in \mathcal{U}_2$, $W_{loc}^{ss}(\sigma_{i,Y}) \cap C(\sigma_Y, Y) = \{\sigma_{i,Y}\}$. In fact, suppose on the contrary that there exists a sequence $Y_n \rightarrow X$ such that $W_{loc}^{ss}(\sigma_{i,Y_n}) \cap C(\sigma_{Y_n}, Y_n) \neq \{\sigma_{i,Y_n}\}$. Then one branch of $W_{loc}^{ss}(\sigma_{i,Y_n})$ is contained in $\{\sigma_{i,Y_n}\}$. By the continuity of local strong stable manifolds of singularities, letting $n \rightarrow \infty$, we get that one branch of $W_{loc}^{ss}(\sigma_i)$ is also contained in $C(\sigma)$, which is a contradiction.

Finally, according to Lemma 3.14, $C(\sigma_Y, Y)$ is Lyapunov stable if Y is weak Kupka-Smale by shrinking \mathcal{U}_2 again if necessary. \square

For C^1 generic $X \in \mathcal{X}^1(M^3)$, it is proved that every Lyapunov stable chain recurrent class C is a quasi attractor by [7], i.e., there exists a sequence of neighborhoods U_n of C such that $\bigcap_{n \geq 1} U_n = C$ and $\phi_1(\overline{U}_n) \subset \text{Int}(U_n)$. Especially there is an arbitrarily small neighborhood U of $C(\sigma)$ such that $\phi_1(\overline{U}) \subset \text{Int}(U)$. As a consequence, there are $\varepsilon_0 > 0$, a C^1 neighborhood \mathcal{U} of X such that for any $Y \in \mathcal{U}$ and any $x \in U$, the strong stable manifold $W_{\varepsilon_0}^{ss}(x, Y)$ exists (e.g., see [10, page 289]).

Definition 6.5. For $Z \in \mathcal{X}^1(M^3)$, S is called a cross-section of Z if

- S is a C^1 surface which is homeomorphic to $(-1, 1)^2$.
- $\angle(T_x S, \langle Z(x) \rangle) > \pi/4, \forall x \in S$.

Definition 6.6. Let σ be a Lorenz-like singularity. S is called a singular cross-section associated to σ if S is a cross section and the following conditions are satisfied:

1. There is a homeomorphism $h = h(x, y) : [-1, 1]^2 \rightarrow \overline{S}$ such that $h((-1, 1)^2) = S$.
2. $\frac{\partial h}{\partial y}$ exists and is continuous on $[-1, 1]^2$.
3. $S \cap W_{loc}^s(\sigma) = h(\{0\} \times (-1, 1))$.

In fact, $(h, [-1, 1]^2)$ (or h for short) is a coordinate system of the surface S .

Denote by $\ell = S \cap W_{loc}^s(\sigma)$ and $S \setminus \ell = S^l \cup S^r$, where $S^l = h((-1, 0) \times (-1, 1))$ and $S^r = h((0, 1) \times (-1, 1))$ (see Figure 1). S^l is called *the left side* of S and S^r is called *the right side* of S .




Figure 1: Cross-section

For any $p, q \in S$, one can define the *horizontal distance* $L_S^h(p, q)$ and *vertical distance* $L_S^v(p, q)$ of p, q : $L_S^h(p, q) = |x_p - x_q|$ and $L_S^v(p, q) = |y_p - y_q|$, where $h(x_p, y_p) = p$ and $h(x_q, y_q) = q$.

Usually, for every Lorenz like singularity ρ , we will take two singular cross-sections S^+, S^- , which are on the opposite sides of $E^{ss}(\rho) \oplus E^u(\rho)$.

For $X \in \mathcal{X}^1(M^3)$, let Λ be a pseudo-Lorenz compact invariant set of X . Denote that $\Lambda \cap \text{Sing}(X) = \{\sigma_1, \dots, \sigma_k\}$ and

$$\Sigma = \bigcup_{1 \leq i \leq k} (S_i^+ \cup S_i^-),$$

where S_i^+ and S_i^- are two singular cross-sections associated to σ_i . For each S_i^\pm , let $h_i^\pm : [-1, 1]^2 \rightarrow \overline{S_i^\pm}$ be the homeomorphism in the definition of singular cross-section. For every $p \in \Sigma$, there exists a unique coordinate system h_i^\pm for some $i \in \{1, 2, \dots, k\}$ and $\pm \in \{+, -\}$. We denote h_i^\pm by h_p . Note that for every $p, p' \in \overline{S_i^\pm}$, $h_p = h_{p'}$.

For each $p \in \Sigma$, if $\{t > 0 : \phi_t(p) \in \Sigma\} \neq \emptyset$, we can define the first return time $t_p = \min\{t > 0 : \phi_t(p) \in \Sigma\}$ and define $F(p) = \phi_{t_p}(p)$. $F : \text{Dom}(F) \rightarrow \Sigma$ is called the *first return map* associated to Σ and Λ , where

$$\text{Dom}(F) = \{p \in \Sigma : \exists t > 0 \text{ s.t. } \phi_t(p) \in \Sigma\}.$$

If $h_p(x, y) = p$ and $F(p) = q$, denote by

$$\tilde{F}(x, y) = \tilde{F}_{p,q}(x, y) = h_q^{-1} \circ F \circ h_p(x, y).$$

Once $F^n(p) = q_n$ is well-defined, denote by

$$\begin{aligned} \tilde{F}^n(x, y) &= h_{q_n}^{-1} \circ F^n \circ h_p(x, y), \\ &= \tilde{F}_{q_{n-1}, q_n} \circ \tilde{F}_{q_{n-2}, q_{n-1}} \circ \cdots \circ \tilde{F}_{p, q_1}(x, y), \end{aligned}$$

where $q_j = F^j(p)$, for $j = 1, 2, \dots, n$ and $h_p(x, y) = p$.

Since S_i^\pm is transverse to X , $\text{Dom}(F)$ is open and t_x is upper semi-continuous with x .

Definition 6.7. For $X \in \mathcal{X}^1(M^3)$, and a pseudo-Lorenz compact invariant set Λ of X , assume that $\Lambda \cap \text{Sing}(X) = \{\sigma_1, \dots, \sigma_k\}$. In the notations above, (Σ, F) is a cross-section system of Λ if the following conditions are satisfied:

1. $\partial\Sigma \cap \Lambda = \emptyset$, where $\partial\Sigma$ is the boundary of Σ .
2. For each $p \in \Sigma$, there is $\varepsilon > 0$ such that $W_{loc}^{ss}(\phi_{(-\varepsilon, \varepsilon)}(p)) \cap \Sigma \subset h_p(\{x\} \times (-1, 1))$, where $h_p(x, y) = p$.
3. For each $1 \leq i \leq k$, and $x \in W_{loc}^s(\sigma_i) \cap \Lambda \setminus \{\sigma_i\}$, there exists $t \in \mathbb{R}$ such that $\phi_t(x) \in \ell_i^\pm$ and if $t > 0$, then $\phi_{[0, t]}(x) \subset W_{loc}^s(\sigma_i)$ and if $t < 0$, $\phi_{[t, 0]}(x) \subset W_{loc}^s(\sigma_i)$.
4. If $p \in \text{Dom}(F)$, then t_r is continuous for $r \in h_p(\{x_p\} \times (-1, 1))^6$.
5. For any $x \in \Lambda \setminus \bigcup_{1 \leq i \leq k} W_{loc}^s(\sigma_i)$, the positive orbit of x will intersect Σ . Especially,

$$\Lambda \cap \Sigma \setminus \bigcup_{1 \leq i \leq k} W_{loc}^s(\sigma_i) \subset \text{Dom}(F).$$

According to the uniform contracting of E^{ss} , we have

Lemma 6.8. Let (Σ, F) be a cross-section system of a pseudo-Lorenz set Λ . Then, there are $C \geq 1$ and $\lambda \in (0, 1)$ such that for any $p \in \Sigma$ with $h_p(x, y) = p$, for any $n \in \mathbb{N}$, if $F^n(p)$ is well-defined, then

$$\left| \frac{\partial \tilde{F}^n}{\partial y}(x, y) \right| \leq C\lambda^n.$$

□

Remark. In the above definition, the local stable manifolds of the singularities also have the exponential contracting property with respect to the flow ϕ_t . Thus,

$$\bigcup_{1 \leq i \leq k} \bigcup_{\pm \in \{+, -\}} \bigcup_{x \in (-1, 1)} h_i^\pm(\{x\} \times (-1, 1))$$

could be regarded as a stable foliation of the first return map.

⁶This will imply some adapted property: the iteration of each stable leaf under the first return map is totally contained in the cross section.

One can define the horizontal distance L^h and the vertical distance L^v of any two points x, y in Σ in the following way:

- If x, y are in a same singular cross-section $S \subset \Sigma$, then $L^h(x, y) = L_S^h(x, y)$ and $L^v(x, y) = L_S^v(x, y)$.
- If x, y are in different singular cross-sections, then $L^h(x, y) = \infty$ and $L^v(x, y) = \infty$.

For any set $\Gamma \subset \Sigma$, one can define the horizontal diameter $D^h(\Gamma)$ and the vertical diameter $D^v(\Gamma)$ by:

$$D^h(\Gamma) = \sup_{x, y \in \Gamma} \{L^h(x, y)\}, \quad D^v(\Gamma) = \sup_{x, y \in \Gamma} \{L^v(x, y)\}.$$

Morales-Pacífico [34] proved the following theorem:

Theorem 6.9. *For C^1 generic $X \in \mathcal{X}^1(M^3)$, if $C(\sigma)$ is a singular hyperbolic Lyapunov stable chain recurrent class, then $C(\sigma)$ is an attractor. As a corollary, $C(\sigma)$ contains periodic orbits.*

Lemma 6.10. *For C^1 generic $X \in \mathcal{X}^1(M^3)$ and $\sigma \in \text{Sing}(X)$, if $C(\sigma)$ is pseudo-Lorenz, then*

- either, $C(\sigma)$ is a homoclinic class;
- or, there is a C^1 neighborhood \mathcal{U}_0 of X such that $C(\sigma_Y)$ is not singular hyperbolic for any $Y \in \mathcal{U}_0$.

Proof. If for any neighborhood \mathcal{U} of X , there is $Y \in \mathcal{U}$ such that $C(\sigma_Y)$ is singular hyperbolic, then by the upper semi-continuity of chain recurrent classes, there is a C^1 neighborhood $\mathcal{U}_Y \subset \mathcal{U}$ of Y such that for every $Z \in \mathcal{U}_Y$, $C(\sigma_Z)$ is singular hyperbolic. This implies that for X , we have $C(\sigma)$ is already singular hyperbolic.

Thus, Theorem 6.9 implies the first case is true. The fact ends the proof. \square

For the study of non-hyperbolic set, Liao [27] and Mañé [32] gave the notion of minimally non-hyperbolic set: for any non-hyperbolic set Λ , there is a compact invariant set $\Lambda_0 \subset \Lambda$ such that every proper compact invariant subset of Λ_0 is hyperbolic. The proof is based on Zorn's Lemma. To understand the dynamics of a compact invariant set which is not singular hyperbolic, sometimes we need to use similar idea.

Definition 6.11. *A nonempty compact invariant set Λ is called a \mathcal{N} -set, if Λ is not singular hyperbolic, and every proper compact invariant set in Λ is singular hyperbolic.*

As in the case of minimally non-hyperbolic set, by Zorn's lemma, for any compact invariant set Λ , if Λ is not singular hyperbolic, then Λ contains a \mathcal{N} -set.

Lemma 6.12. *For C^1 generic $X \in \mathcal{X}^1(M^3)$ and $\sigma \in \text{Sing}(X)$, if $C(\sigma)$ is pseudo-Lorenz, and not a homoclinic class, then there is a C^1 neighborhood \mathcal{U}_0 of X such that for every $Y \in \mathcal{U}_0$, $C(\sigma_Y)$ contains a transitive \mathcal{N} -set Λ_Y such that $\Lambda_Y \not\subset \text{Sing}(Y)$.*

Proof. By Lemma 6.10 and Lemma 6.4, there is a neighborhood \mathcal{U}_0 of X such that for any $Y \in \mathcal{U}_0$, one has

- $C(\sigma_Y, Y)$ is not singular hyperbolic.
- $C(\sigma_Y, Y)$ admits a partially hyperbolic splitting $T_{C(\sigma_Y, Y)}M^3 = E^{ss} \oplus E^{cu}$, where $\dim E^{ss} = 1$, and every singularity in $C(\sigma_Y, Y)$ is Lorenz-like.

For each $Y \in \mathcal{U}_0$, $C(\sigma_Y)$ is not singular hyperbolic. Hence, there exists a \mathcal{N} -set $\Lambda_Y \subset C(\sigma_Y)$.

We will prove that Λ_Y is transitive. If not, for every $x \in \Lambda_Y$, $\alpha(x)$ is a proper subset of Λ_Y . As a consequence, $\alpha(x)$ is singular hyperbolic for every $x \in \Lambda_Y$. We have already known that Λ_Y admits a partially hyperbolic splitting $T_{\Lambda_Y}M = E^{ss} \oplus E^{cu}$ since $C(\sigma_Y)$ is pseudo-Lorenz. If $\alpha(x)$ is singular hyperbolic, then $\lim_{t \rightarrow \infty} |\text{Det}\Phi_{-t}^Y|_{E^{cu}(x)}| = 0$. Since every $x \in \Lambda_Y$ has this property, by a compact argument (e.g., see Lemma 5.9), one can prove that Λ_Y is singular hyperbolic. This contradicts to the assumption that Λ_Y is an \mathcal{N} -set. \square

Corollary 6.13. *Under the assumption of Lemma 6.12, for every \mathcal{N} -set Λ_Y , there is an ergodic measure μ_Y of ϕ_t^Y such that the support of μ_Y is Λ_Y , and for any $t > 0$, one has*

$$\int \log |\text{Det}\Phi_t|_{E^{cu}(x)}| d\mu_Y \leq 0.$$

Proof. By Lemma 5.9, there is a point $x \in \Lambda_Y$ such that $\log |\text{Det}\Phi_t|_{E^{cu}(x)}| \leq 0$ for any $t \geq 0$. By using a standard method, we can have an invariant measure ν such that for any $t > 0$, one has

$$\int \log |\text{Det}\Phi_t|_{E^{cu}(x)}| d\nu \leq 0.$$

By using the ergodic decomposition theorem, there is an ergodic component μ of ν such that

$$\int \log |\text{Det}\Phi_t|_{E^{cu}(x)}| d\mu \leq 0.$$

$\text{supp}(\mu) = \Lambda$: otherwise, $\text{supp}(\mu)$ is a proper compact invariant set of Λ_Y ; hence it is singular hyperbolic, which implies that the inequality above is false. \square

We know some structures about minimally non-hyperbolic *non-singular* set for C^2 vector fields when the set admits a dominated splitting w.r.t. the linear Poincaré flow, which is the main theorem in [5].

Theorem 6.14. *Let $X \in \mathcal{X}^2(M^3)$. If a compact invariant set Λ satisfies the following properties:*

- Λ is transitive.
- $\Lambda \cap \text{Sing}(X) = \emptyset$.
- The linear Poincaré flow ψ_t admits a dominated splitting on \mathcal{N}_Λ .
- Every periodic point in Λ is hyperbolic.

Then,

- either Λ is hyperbolic;
- or $\Lambda = \mathbb{T}^2$, Λ is a normally hyperbolic set with respect to Φ_t , and $\phi_t|_\Lambda$ is equivalent to an irrational flow.

Corollary 6.15. *Under the assumption of Lemma 6.12, every \mathcal{N} -set Λ_Y contains a singularity for every C^2 weak Kupka-Smale vector field $Y \in \mathcal{U}_0$*

Proof. This is just an application of Theorem 6.14 because $C(\sigma_Y)$ cannot contain a normally hyperbolic torus. \square

Remark. *This is another place that we need to use the assumption of “weak Kupka-Smale” besides the usage of the connecting lemma for pseudo orbits.*

Let us concentrate on the proof of Theorem 6.2. We need to fix a neighborhood such that we can do perturbations freely in the neighborhood. Under the assumptions of Theorem 6.2, if the conclusion is not true, then there is a C^1 neighborhood \mathcal{U}^* such that

- For every $Y \in \mathcal{U}^*$, $C(\sigma_Y)$ is not singular hyperbolic.
- Since chain recurrent classes is upper semi-continuous by Lemma 3.11, one has for any $Y \in \mathcal{U}^*$, $C(\sigma_Y)$ has a partially hyperbolic splitting $T_{C(\sigma_Y)}M^3 = E^{ss} \oplus E^{cu}$ with respect to the tangent flow.
- By the continuity of eigenvalues of singularity and Lemma 3.12, every singularity in $C(\sigma_Y)$ is a continuation of a singularity in $C(\sigma)$, and it is Lorenz-like.
- $C(\sigma_Y)$ is still Lyapunov stable by Lemma 3.14 if Y is weak Kupka-Smale.
- $C(\sigma_Y)$ is not singular hyperbolic by Lemma 6.10. As in Lemma 6.12, $C(\sigma_Y)$ contains a transitive \mathcal{N} -set inside, which is not reduced to a singularity.

Proposition 6.16. *For C^1 generic vector field $X \in \mathcal{U}^*$, and $\sigma \in \text{Sing}(X)$, if $C(\sigma)$ is pseudo-Lorenz and contains no periodic orbit, then $C(\sigma)$ admits a cross-section system (Σ, F) .*

Proof. The preparation: curves in the stable manifolds. Assume that

$$C(\sigma) \cap \text{Sing}(X) = \{\sigma_1, \sigma_2, \dots, \sigma_k\}.$$

For each singularity σ_i , one can choose a local chart $\phi_i^s : \mathbb{R}^2 \rightarrow W_{loc}^s(\sigma_i)$ such that

- $\phi_i^s(\{0\} \times \mathbb{R}) = W_{loc}^{ss}(\sigma_i)$,
- $\phi_i^s(\mathbb{R} \times \{0\})$ is an invariant central manifold.

Take two curves $\gamma_{i,1}, \gamma_{i,2} : \mathbb{R} \rightarrow W^s(\sigma_i)$ as the images of $y = x$ and $y = -x$ under ϕ_i^s such that $\gamma_{i,j}(0) = \sigma_i$ for $j = 1, 2$.

Claim. *There is $\rho_i > 0$ such that $C(\sigma) \cap \phi_i^s([- \rho_i, \rho_i]^2) \subset \phi_i^s(\{(x, y) : |x| < |y|\})$. As a corollary, $\gamma_{i,1}([- \rho_i, \rho_i]) \cap C(\sigma) = \{\sigma_i\}$ and $\gamma_{i,2}([- \rho_i, \rho_i]) \cap C(\sigma) = \{\sigma_i\}$.*

Proof of the claim. If the claim is not true, there are $x_n \in \phi_i^s(\{(x, y) : |x| \geq |y|\}) \cap C(\sigma)$ such that $\lim_{n \rightarrow \infty} x_n = \sigma_i$ and $x_n \neq \sigma_i$. The negative iterations of x_n are still in $C(\sigma)$. Choose a small neighborhood B_i of σ_i . Let

$$t_n = \sup\{t : \phi_{-s}(x_n) \in B_i, \forall 0 \leq s \leq t\}.$$

We have that $t_n \rightarrow \infty$ as $x_n \rightarrow \sigma_i$. Let a be an accumulation point of $\phi_{-t_n}(x_n)$. Then $\phi_t(a) \in \{(x, y) : |x| \geq |y|\}$ for $t \geq 0$. Hence $a \in C(\sigma) \cap W_{loc}^{ss}(\sigma_i) \cap \partial B_i$. This contradicts to the fact that $W_{loc}^{ss}(\sigma_i) \cap C(\sigma) = \{\sigma_i\}$. \square

The first step of the construction: Let $\rho = \min\{\rho_i : 1 \leq i \leq k\}$. For each σ_i , there are two connected components Θ_i^\pm of $W_{loc}^s(\sigma_i) \setminus W_{loc}^{ss}(\sigma_i)$. In the following, we will use Θ_i^+ to construct S_i^+ , while the other case can be constructed similarly.

There are two points $x_{i,1} \in \gamma_{i,1} \cap \Theta_i^+$ and $x_{i,2} \in \gamma_{i,2} \cap \Theta_i^+$ such that

- $x_{i,1}, x_{i,2} \notin C(\sigma)$.
- $x_{i,1} \in W_{loc}^{ss}(x_{i,2})$.

Thus there is a cross-section $\tilde{S}_i^+ = h_i^+((-1, 1)^2)$, where $h_i^+ : [-1, 1]^2 \rightarrow \overline{\tilde{S}_i^+}$ is a homeomorphism, such that

- $h_i^+((0, -1)) = x_{i,1}$, $h_i^+((0, 1)) = x_{i,2}$ and $h_i^+({0} \times (-1, 1))$ is a connected part of a strong stable manifold of ϕ_t .
- $h_i^+((-1, 1) \times \{-1, 1\}) \cap C(\sigma) = \emptyset$.
- \tilde{S}_i^+ is foliated by strong stable foliation in the following sense: for each $x \in \tilde{S}_i^+$, one define $\mathcal{F}^s(x)$ to be the connected component of $\bigcup_{t \geq 0} \phi_t(W_{\varepsilon_1}^{ss}(x)) \cap \tilde{S}_i^+$, \tilde{S}_i^+ can be foliated by \mathcal{F}^s . Moreover, $h_i^+({z} \times (-1, 1))$ is a leaf of the strong stable foliation.
- For any arbitrarily small number $\alpha > 0$, one can require that $D^h(\tilde{S}_i^+) < \alpha$.
- $\bigcup_{x \in (-1, 1)} h_i^+({x} \times (-1, 1))$ is a family of C^1 curves, and as a C^1 family, it varies continuously with respect to x .

One can construct \tilde{S}_i^- for Θ_i^- also.

Refine the construction: We take a neighborhood $U = B_\delta(C(\sigma))$ with δ small enough such that $B_\delta(\sigma_i)$ and $B_\delta(\sigma_j)$ are disjoint for $i \neq j$ and \overline{U} is disjoint from $h_i^\pm((-\beta_0, \beta_0) \times \{-1, 1\})$ for any i and any $\pm \in \{+, -\}$. Denote $S_i^\pm(\beta) = h_i^\pm((-\beta, \beta) \times (-1, 1))$ for $\beta \in (0, \beta_0]$. Denote by

$$\Sigma(\beta) = \bigcup_{1 \leq i \leq k, \pm \in \{+, -\}} S_i^\pm(\beta).$$

As before, consider the first return map F with respect to $\Sigma(\beta)$.

Claim. *If β is small enough, for any $x \in \text{Dom}(F)$, $t_x = \inf\{t > 0 : \phi_t(x) \in \Sigma(\beta)\}$ is continuous on $\mathcal{F}^s(x)$, the strong stable leaf of x with respect to F .*




Figure 2: Cross-section and return map: Λ does not intersect the shaded area and the image of any strong stable leaf under the return map does not intersect the shaded area.

Proof of the claim. Otherwise, $\phi_t(\mathcal{F}^s(x))$ will intersect some $h_i^\pm((-\beta_0, \beta_0) \times \{-1, 1\})$ for some $t > 0$. Since $C(\sigma)$ is Lyapunov stable, for $\epsilon > 0$ small enough, for any $t \geq 0$, $\phi_t(B_\epsilon(\sigma_i)) \subset U$. But for β small enough, for any $x \in \text{Dom}(F)$, there exists $T > 0$ such that $\phi_T(x) \in B_\epsilon(\sigma_i)$ for some i before it first returns to $\Sigma(\beta)$ since $S_i^\pm(\beta)$ is very close to the stable manifold of σ_i . So for $t > T$, $\phi_t(x) \in U$. Especially, $\phi_t(x) \cap h_i^\pm((-\beta_0, \beta_0) \times \{-1, 1\}) = \emptyset$ for any i and any $\pm \in \{+, -\}$. This contradiction proves the claim. \square

Since X is a C^1 generic vector field, $C(\sigma)$ can be accumulated by periodic orbits. Since $X \in \mathcal{U}_0$, $C(\sigma)$ contains no hyperbolic periodic orbit.

Claim. Given $\beta \in (0, \beta_0]$, for every $i = 1, 2, \dots, k$ and $\pm \in \{+, -\}$,

$$J_i^\pm = \{z \in (-\beta, \beta) : h_i^\pm(\{z\} \times (-1, 1)) \cap C(\sigma) = \emptyset\}$$

is open and dense in $(-\beta, \beta)$.

Proof of the claim. We only have to prove that: Given $a, b \in (-\beta, \beta)$, $a < b$, there exists $z \in (a, b)$ such that $h_i^\pm(\{z\} \times (-1, 1)) \cap C(\sigma) = \emptyset$. Otherwise, since $C(\sigma)$ can be accumulated by periodic orbits, there is a periodic point $p \in S_i^\pm$ close to $h_i^\pm(\{(a+b)/2\} \times (-1, 1))$. Thus $W_{\varepsilon_0}^{ss}(p) \cap C(\sigma) \neq \emptyset$. This contradicts to the fact that $C(\sigma) \cap \text{Per}(X) = \emptyset$. \square

For $\beta > 0$ small enough,

$$G = \bigcap_{1 \leq i \leq k, \pm \in \{+, -\}} (0, \beta) \cap J_i^\pm \cap (-J_i^\pm)$$

is open and dense in $(0, \beta)$.

So, take $\beta' \in G$ and let $\Sigma = \Sigma(\beta')$. After scaling, we may assume coordinate mappings h_i^\pm are defined on $(-1, 1)^2$. Then (Σ, F) satisfies item 1)-4) in the definition of cross-section system.

The domain of F : Since we assume that $C(\sigma)$ contains no periodic orbit, we have that $\omega(x)$ contains a singularity for any $x \in C(\sigma)$. This implies that for every $x \in C(\sigma)$, if x is not in the local stable manifold of some singularity, then the positive iterations of x will intersect Σ . This finishes the proof of the existence of cross-section system. \square

By summarizing the construction as in the above proof, we first find some ‘‘large’’ cross-section $\Sigma(1)$, then we just take some smaller part $\Sigma(\beta')$ which is modified by the strong stable foliation. Since the local strong stable manifolds are continuous with respect to the vector fields, for any $Y \in \mathcal{U}^*$ close to X , the intersection of the strong stable manifolds $W_{loc}^{ss}(z, Y)$ of $z \in \Sigma(\beta')$ and $\Sigma(1)$ is close to $\Sigma(\beta')$. Thus, the cross-section system has some continuous property from the proof of the above proposition.

Corollary 6.17. For C^1 generic vector field $X \in \mathcal{U}^*$, let $C(\sigma)$ be a chain recurrent class satisfying the assumption of Theorem 6.2. Then there is a C^1 neighborhood $\mathcal{U} \subset \mathcal{U}^*$ of X , such that for any $Y \in \mathcal{U}$, $C(\sigma_Y)$ admits a cross-section system (Σ_Y, F_Y) . Moreover, when $Y \rightarrow X$, we have $\Sigma_Y \rightarrow \Sigma$.

Proof. As explained above, we take

$$\Sigma_Y = \left(\bigcup_{z \in \Sigma_X} \phi_{[-\varepsilon(z), \varepsilon(z)]}(W_{loc}^{ss}(z, Y)) \right) \cap \Sigma(1).$$

Moreover, one can defined the first return map F_Y by using this cross section.

By the continuity of the local strong stable manifolds w.r.t. the vector fields, we have that Σ_Y is close to $\Sigma = \Sigma_X$ when Y is close to X . Since $C(\sigma_Y)$ is continuous w.r.t. Y , we have that

$C(\sigma_Y) \cap \partial\Sigma_Y = \emptyset$. By the definition of Σ_Y , Item 2 of the cross section system is satisfied. Item 3 is true because that $C(\sigma_Y) \cap W_{loc}^s(\sigma_Y)$ is continuous w.r.t. Y . Item 4 is true if the return time is long, which can be guaranteed if the cross section is thin. We have Item 5 because

- The return time is uniformly continuous when the point is not close to the local stable manifolds of the singularities.
- When the point is close to the local stable manifolds of the singularities, the return will follow the unstable manifolds of the singularities, which are stably contained in the new cross-section.

□

Now we take a C^1 neighborhood $\mathcal{U}^{**} \subset \mathcal{U}^*$ of X such that \mathcal{U}^{**} verifies Corollary 6.17.

6.2 Homoclinic orbits of singularities

Now we are going to prove Proposition 4.9: for generic $X \in \mathcal{X}^1(M^3)$, if $C(\sigma)$ is pseudo-Lorenz, then either $C(\sigma)$ is a singular hyperbolic attractor; or for any C^1 neighborhood \mathcal{U} of X , there is a weak Kupka-Smale vector field $Y \in \mathcal{U}$ such that Y has a periodic sink γ and $\text{Basin}(\gamma) \cap C(\sigma_Y) \neq \emptyset$. As the end of Section 4, once we know Proposition 4.9 is true, we will get the final result.

For each Lorenz-like singularity ρ and a small neighborhood U of this singularity, the local stable manifold of ρ divides U into two parts: the left part and the right part. Thus the unstable manifold $W^u(\rho)$ has two separatrices: the left separatrix and the right separatrix.

For each $Y \in \mathcal{U}^{**}$, one defines $n(Y)$ to be the number of homoclinic orbits of singularities contained in $C(\sigma_Y)$. Since there are only k singularities in $C(\sigma)$, $n(Y) \leq 2k$. Let

$$n = \max\{n(Y) : Y \in \mathcal{U}_1, Y \text{ is } C^2 \text{ and is weak Kupka - Smale}\}.$$

Let $\mathcal{M}_n \subset \mathcal{U}^{**}$ be the set of C^2 weak Kupka-Smale vector fields Y with $n(Y) = n$.

Lemma 6.18. *For $Y \in \mathcal{M}_n$ and for the transitive \mathcal{N} -set $\Lambda_Y \subset C(\sigma_Y)$ as in Lemma 6.12, if $\sigma_{i,Y} \in \Lambda_Y$, then $\sigma_{i,Y}$ has a homoclinic orbit Γ_i^\pm .*

Proof. Suppose on the contrary, Λ_Y contains a $\sigma_{i,Y}$, but $C(\sigma_Y)$ does not contain a homoclinic orbit of $\sigma_{i,Y}$. Since Λ_Y is transitive, $\Lambda_Y \cap W^s(\sigma_{i,Y}) \setminus \{\sigma_{i,Y}\} \neq \emptyset$ and $\Lambda_Y \cap W^u(\sigma_{i,Y}) \setminus \{\sigma_{i,Y}\} \neq \emptyset$. Take $x^s \in \Lambda_Y \cap W^s(\sigma_{i,Y}) \setminus \{\sigma_{i,Y}\}$ and $x^u \in \Lambda_Y \cap W^u(\sigma_{i,Y}) \setminus \{\sigma_{i,Y}\}$. By the assumptions, x^s and x^u are not in a homoclinic orbit.

Construction of perturbation boxes. For $Y \in \mathcal{M}_n$, one can choose $\varepsilon > 0$, such that $B(Y, \varepsilon) \subset \mathcal{U}^{**}$. By Lemma 3.4, one can choose $L > 0$ and neighborhoods $\widetilde{W}_{x^s} \subset W_{x^s}$ of x^s , $\widetilde{W}_{x^u} \subset W_{x^u}$ of x^u as in Lemma 3.4 such that

- $W_{L,x^s} \cap W_{L,x^u} = \emptyset$.
- $\Lambda_Y \setminus (W_{L,x^s} \cup W_{L,x^u}) \neq \emptyset$.
- $W_{L,x^s} \cup W_{L,x^u}$ is disjoint from any other homoclinic orbits of singularities.
- $W_{L,x^s} \cup W_{L,x^u}$ is disjoint from $\sigma_{i,Y}$.

Choosing the orbits. Since Λ_Y is transitive, there is $z \in \Lambda_Y \setminus (W_{L,x^s} \cup W_{L,x^u})$ such that $\alpha(z) = \omega(z) = \Lambda$. Choose $t_1, t_2 > 0$ such that $\phi_{-t_1}^Y(z) \in \widetilde{W}_{x^s}$ and $\phi_{t_2}^Y(z) \in \widetilde{W}_{x^u}$. Choose $t_s > 0$ and $t_u > 0$ such that $\phi_{t_s}^Y(x^s) \notin W_{L,x^s} \cup W_{L,x^u}$ and $\phi_{-t_u}^Y(x^u) \notin W_{L,x^s} \cup W_{L,x^u}$.

Connecting the orbit from $\phi_{-t_2}^Y(z)$ to $\phi_{t_s}^Y(x^s)$. Since the negative orbit of $\phi_{t_s}^Y(x^s)$ and the positive orbit of $\phi_{-t_2}^Y(z)$ both enter \widetilde{W}_{x^s} , by using Lemma 3.4, there is Y_1 which is ε -close to Y such that

- There is $T_1 > 0$ such that $\phi_{-T_1}^{Y_1}(\phi_{t_s}^Y(x^s)) = \phi_{-t_2}^Y(z)$.
- $Y_1(x) = Y(x)$ for any $x \in M^3 \setminus W_{L,x^s}$.

As a corollary, we have

- Any homoclinic orbit of Y is still a homoclinic orbit of Y_1 .
- $\sigma_{i,Y}$ is still a singularity of Y_1 , $\phi_{t_s}^Y(x^s)$ is still in the stable manifold of $\sigma_{i,Y}$ and $\phi_{-t_u}^Y(x^u)$ is still in the unstable manifold of $\sigma_{i,Y}$ with respect to Y_1 .

Connecting the orbit from $\phi_{-t_u}^Y(x^u)$ to $\phi_{t_s}^Y(x^s)$. Since $\phi_{-T_1}^{Y_1}(\phi_{t_s}^Y(x^s)) = \phi_{-t_2}^Y(z)$ is contained in \widetilde{W}_{x^u} , by using Lemma 3.4 again, there is Y_2 which is ε -close to Y_1 such that

- There is $T_2 > 0$ such that $\phi_{-T_2}^{Y_1}(\phi_{t_s}^Y(x^s)) = \phi_{-t_u}^Y(x^u)$.
- $Y_2(x) = Y_1(x)$ for any $x \in M^3 \setminus W_{L,x^u}$.

As a corollary, we have

- $Y(x) = Y_2(x)$ for any $x \in M \setminus (W_{L,x^s} \cup W_{L,x^u})$.
- Any homoclinic orbit of Y is still a homoclinic orbit of Y_1 .
- $\phi_{t_s}^Y(x^s)$ is in a homoclinic orbit of $\sigma_{i,Y}$.

Since $Y_2 \in \mathcal{U}_1$ and $Y \in \mathcal{M}_n$, we have that Y_2 has $n + 1$ homoclinic orbits of singularities in $C(\sigma_Y)$, which contradicts to the maximality of n . \square

Definition 6.19. A point $p \in M$ is called a typical point of a probability ergodic measure μ of a vector field Y , if the following conditions are satisfied:

1. p is strongly closable;
2. $\omega(p) = \text{supp}(\mu)$;
3. for every continuous function $f : M \rightarrow \mathbb{R}$,

$$\lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T f(\phi_t^Y(p)) dt = \int f(x) d\mu(x).$$

According to Ergodic Closing Lemma and Birkhoff Ergodic Theorem, the set of typical points of μ has μ -full measure.

Proposition 6.20. For any $Y \in \mathcal{M}_n$, and $\sigma_{i,Y} \in \Lambda_Y \cap \text{Sing}(Y)$, assume that

1. The right separatrix Γ of $\sigma_{i,Y}$ is a homoclinic orbit. Let $x \in W_{loc}^u(\sigma_{i,Y}) \cap \Gamma$ and denote

$$\mathcal{T}_x = \{t > 0 : \phi_t^Y(x) \in \Sigma\} = \{t_1 < t_2 < \cdots < t_{N(x)}\},$$

where $N(x) = \#\mathcal{T}_x$. We may assume that $\phi_{t_{N(x)}}^Y(x) \in S_i^+$.

2. There are⁷ $N \in \mathbb{N}$ and $\beta \in (0, 1)$ such that

$$F^N(h_i^+((0, \beta) \times (-1, 1))) \subset S_i^+.$$

3. Let μ_Y be the ergodic measure as in Corollary 6.13. If (Λ_Y, μ_Y) accumulates $\phi_{t_{N(x)}}^Y(x)$ on the right: there is a sequence of typical points $x_n \in h_i^+((0, 1) \times (-1, 1))$ of μ_Y such that $\lim_{n \rightarrow \infty} x_n = \phi_{t_{N(x)}}^Y(x)$.

Then for any C^1 neighborhood \mathcal{V} of Y , there is a weak Kupka-Smale vector field $Z \in \mathcal{V}$ such that Z has a periodic sink, whose basin accumulates on $C(\sigma_Z)$. In other words, Proposition 4.9 is true in this case.

Proof. Since every singularity in $C(\sigma_Y)$ is Lorenz-like, for β_0 small enough, we have $F^N(x)(h_i^+(\{\beta_0\} \times (-1, 1))) \subset h_i^+(\{\beta_1\} \times (-1, 1))$ for some $\beta_1 \in (\beta_0, 1)$.

Claim. *There is a weak Kupka-Smale vector field Z arbitrarily C^1 -close to Y such that Z has a periodic sink γ with the following properties:*

- γ intersects $S_i^{+,r}$, and one can choose the intersection points arbitrarily close to $\phi_{t_{N(x)}}^Y(x)$.
- Every homoclinic orbit of singularities of Y is still a homoclinic orbit of singularities of Z .

Proof of the claim. Given $T > 0$, define $f_T(x) = \log |\text{Det} \Phi_T^Y|_{E^{cu}(x)}|$ for any $x \in C(\sigma_Y)$. One knows that f_T is a continuous function on $C(\sigma)$. Since E^{cu} can be extended continuously in a small neighborhood of $C(\sigma)$, f_T can be also extended continuously. Denote by \widehat{E}^{cu} and \widehat{f}_T the extension of E^{cu} and f_T respectively. Note that we don't require that \widehat{E}^{cu} is invariant. By the property of μ_Y , one has $\int f_T d\mu_Y \leq 0$

Since μ_Y is ergodic and $\text{supp}(\mu_Y) = \Lambda$, one has that the set of homoclinic orbits of singularities has zero measure w.r.t. μ_Y . Choose a typical point x of μ_Y , which is very close to $\phi_{t_{N(x)}}^Y(x)$ and which is in $S_i^{+,r}$. Since x is typical, one can assume that x is a strong closable point by Lemma 3.6. Now by Corollary 3.7, for any $\varepsilon > 0$, there is Z ε -close to Y such that

- Every homoclinic orbit of singularities of Y is still a homoclinic orbit of singularities of Z .
- Z has a periodic γ such that

$$\left| \int \widehat{f}_T d\delta_\gamma - \int \widehat{f}_T d\mu_Y \right| < \varepsilon.$$

Since γ is close to Λ_Y , one knows that γ admits a partially hyperbolic splitting $T_\gamma M^3 = E^{ss,Z} \oplus E^{cu,Z}$. By the property of dominated splittings, for any $x \in \gamma$, one has that $\widetilde{d}(\widehat{E}^{cu}(x), E^{cu,Z}(x)) = O(\varepsilon)$, where $O(\varepsilon)$ means that there is a constant $C > 0$ such that $O(\varepsilon)/\varepsilon \leq C$ for ε small enough.

Thus, one has that $|\int \log |\text{Det} \Phi_T^Z| d\delta_\gamma - \int f_T d\mu_Y| = O(\varepsilon)$. As a consequence,

$$\int \log |\text{Det} \Phi_T^Z| d\delta_\gamma \leq O(\varepsilon).$$

Thus, by using the Franks Lemma (Lemma 2.5), by an extra small perturbation in an arbitrarily small neighborhood of γ , one gets that γ is a periodic sink. Finally, by Theorem 3.10, one can assume that Z is weak Kupka-Smale. \square

⁷Notice that here we use N but not $N(x)$, because we need to deal with some general case.

Since Z is close enough to Y , Z has a cross-section system close to Σ by Corollary 6.17. For simplicity, we still denote by Σ the cross-section system of Z .

Claim. *There is $\beta_Z > 0$ such that for any $\beta \in (0, \beta_Z]$, there is $N(\beta) \in \mathbb{N}$ such that $F_Z^{N(\beta)}(h_i^+((0, \beta) \times (-1, 1))) \supset h_i^+((0, \beta_Z] \times (-1, 1))$.*

Proof. This is true because the singularities is Lorenz-like: for the return map near the local stable manifold of the singularities, the horizontal direction will be expanded. Moreover, the local stable manifold of the singularities served as “fixed points” of the return map. \square

We need to consider the relation between the intersections of γ and Γ with S_j^\pm . For each j and $\pm \in \{+, -\}$, denote

$$\mathcal{T}_{j,\gamma}^\pm = \{t_\gamma : \gamma \cap h_j^\pm(\{t_\gamma\} \times [-1, 1]) \neq \emptyset\}.$$

$$\mathcal{T}_{j,\Gamma}^\pm = \{t_\Gamma : \Gamma \cap h_j^\pm(\{t_\Gamma\} \times [-1, 1]) \neq \emptyset\}.$$

$$a_j^{\pm,l} = \max \mathcal{T}_{j,\gamma}^\pm \cap [-1, 0), \quad a_j^{\pm,r} = \min \mathcal{T}_{j,\gamma}^\pm \cap (0, 1].$$

$$b_j^{\pm,l} = \max \mathcal{T}_{j,\Gamma}^\pm \cap [-1, 0), \quad b_j^{\pm,r} = \min \mathcal{T}_{j,\Gamma}^\pm \cap (0, 1].$$

$\mathcal{T}_{j,\Gamma}^\pm$ and $\mathcal{T}_{j,\gamma}^\pm$ may be empty for some j, \pm . Since γ can be chosen arbitrarily close to Γ and obtained by a typical orbit, we may assume that if $\mathcal{T}_{j,\Gamma}^\pm$ is nonempty, then $\mathcal{T}_{j,\gamma}^\pm$ is also nonempty. Moreover, consider the iterates of the right direction R attached to Γ in S_i^+ . We may assume that

1. If Γ intersects $S_j^{\pm,l}$, and the corresponding iterate of R is still the right direction in S_j^\pm , then $b_j^{\pm,l} < a_j^{\pm,l}$.
2. If Γ intersects $S_j^{\pm,r}$, and the corresponding iterate of R is the left direction in S_j^\pm , then $b_j^{\pm,r} > a_j^{\pm,r}$.

Since Z is weak Kupka-Smale, $C(\sigma_Z)$ is Lyapunov stable. This implies that for given $r \in \mathbb{N}$, $F_Z^r(h_i^+((0, \beta) \times (-1, 1)))$ is well defined for β small enough since the boundary of Σ_Z dose not intersect $C(\sigma_Z)$.

We will deal with the following two cases:

1. For any $\beta \in (0, \beta_Z]$, there exists some $r \geq N(\beta)$, such that $F_Z^r(h_i^+((0, \beta_Z) \times (-1, 1)))$ intersects l_j^\pm for some j, \pm .
2. For any $r > 0$, $F_Z^r(h_i^+((0, \beta_Z) \times (-1, 1)))$ will never intersect l_j^\pm for any j, \pm .

Case 1. We may assume that r is the smallest positive integer such that $F_Z^r(h_i^+((0, \beta_Z) \times (-1, 1)))$ intersects l_j^\pm for some j, \pm . Without loss of generality, we may assume that

$$\lim_{t \rightarrow 0^+} F_Z^r(h_i^+(\{t\} \times (-1, 1))) \in S_j^{\pm,l}.$$

Since $C(\sigma_Z)$ is Lyapunov stable, the corresponding iterate $F_Z^r(R)$ of R is still the right direction in S_j^\pm . And hence $F_Z^r(h_i^+((0, \beta) \times (-1, 1)))$ intersects $h_j^\pm(\{a_j^{\pm,l}\} \times (-1, 1))$, which are contained in the basin of the sink γ . The conclusion follows by letting $\beta \rightarrow 0$.

Case 2. For any $r > 0$, $F_Z^r(h_i^+((0, \beta_Z] \times (-1, 1)))$ will never intersect l_j^\pm for any j, \pm .

This will imply that for every $r > 0$, $F_Z^r(h_i^+((0, \beta_Z] \times (-1, 1)))$ is connected.

For any $\beta_0 \in (0, \beta_Z]$, there is an increasing sequence $\{\beta_r\}_{r \in \mathbb{N}} \subset (0, 1)$ such that $F_Z^r(h_i^+(\{\beta_0\} \times (-1, 1))) \subset h_i^+(\{\beta_r\} \times (-1, 1))$ for any $r \in \mathbb{N}$. We assume that $\beta' = \lim_{r \rightarrow \infty} \beta_r$. By the continuity of F_Z and h_i^+ , one has $F_Z(h_i^+(\{\beta'\} \times (-1, 1))) \subset h_i^+(\{\beta'\} \times (-1, 1))$. Since $h_i^+(\{\beta'\} \times (-1, 1))$ is a stable leaf of \tilde{F} , one has $h_i^+(\{\beta'\} \times (-1, 1))$ contains a periodic point of Z , which is a hyperbolic sink of Z . This implies that $h_i^+((0, \beta_Z] \times (-1, 1))$ is in a basin of the sink. \square

Remark. *In the above proof, since we are in the non-generic case, we need to do one extra perturbation to get a sink in a suitable position. This is one of the main difficulties in this paper.*

A direct application of Proposition 6.20 would be the following result:

Proposition 6.21. *For any $Y \in \mathcal{M}_n$, and $\sigma_{i,Y} \in \Lambda_Y \cap \text{Sing}(Y)$, assume that the right separatrix of $W^u(\sigma_{i,Y})$ is a homoclinic orbit Γ_i of $\sigma_{i,Y}$. Let $x \in W_{\text{loc}}^u(\sigma_{i,Y}) \cap \Gamma_i$ and denote by*

$$\mathcal{T}_x = \{t > 0 : \phi_t^Y(x) \in \Sigma\} = \{t_1 < t_2 < \dots < t_{N(x)}\},$$

where $N(x) = \#\mathcal{T}_x$. We may assume that $\phi_{t_{N(x)}}^Y(x) \in S_i^+$.

Let μ_Y be the ergodic measure as in Corollary 6.13. If (Λ_Y, μ_Y) accumulates $\phi_{t_{N(x)}}^Y(x)$ on the right: there is a sequence of typical points $x_n \in h_i^+((0, 1) \times (-1, 1))$ of μ_Y such that $\lim_{n \rightarrow \infty} x_n = \phi_{t_{N(x)}}^Y(x)$, then

- either, for $t > 0$ small enough we have that

$$F^{N(x)}(h_i^+((0, t) \times (-1, 1))) \subset h_i^+((-1, 0) \times (-1, 1)).$$

- or, for any neighborhood \mathcal{V} of Y , there is a weak Kupka-Smale vector field $Z \in \mathcal{V}$ such that Z has a periodic sink γ and $\overline{\text{Basin}(\gamma)} \cap C(\sigma_Z) \neq \emptyset$.




Figure 3:

Proof. First, notice that $N(x) = \#\mathcal{T}_x$ is finite. For $t_0 > 0$ small enough, $F^{N(x)}(h_i^+((0, t_0] \times (-1, 1)))$ is a connected set contained in S_i^+ . If $F^{N(x)}(h_i^+((0, t_0] \times (-1, 1))) \subset h_i^+((0, 1) \times (-1, 1))$, then it is true for any $t \in (0, t_0)$. Hence, by Proposition 6.20, we know that for any neighborhood \mathcal{V} of Y , there is a weak Kupka-Smale vector field $Z \in \mathcal{V}$ such that Z has a periodic sink γ and $\overline{\text{Basin}(\gamma)} \cap C(\sigma_Z) \neq \emptyset$. \square

Remark. For generic X , as in the proof of Proposition 6.16, the projection of $C(\sigma) \cap \Sigma$ to the horizontal direction along the strong stable foliation is nowhere dense. But $Y \in \mathcal{M}_n$ is not generic. This is why we need to consider a transitive subset Λ_Y and the ergodic measure supported on Λ .

Corollary 6.22. For any $Y \in \mathcal{M}_n$, and $\sigma_{i,Y} \in \Lambda_Y \cap \text{Sing}(Y)$, assume that the right separatrix of $W^u(\sigma_{i,Y})$ is a homoclinic orbit Γ_i of $\sigma_{i,Y}$. Let $x \in W_{loc}^u(\sigma_{i,Y}) \cap \Gamma_i$ and denote

$$\mathcal{T}_x = \{t > 0 : \phi_t^Y(x) \in \Sigma\} = \{t_1 < t_2 < \cdots < t_{N(x)}\},$$

where $N(x) = \#\mathcal{T}_x$. If $\phi_{t_{N(x)}}^Y(x)$ is contained in Λ_Y , then (Λ_Y, μ_Y) accumulates $\phi_{t_{N(x)}}^Y(x)$ on the left.

Proof. If (Λ_Y, μ_Y) accumulates $\phi_{t_{N(x)}}^Y(x)$ on the left, it is finished. Otherwise, (Λ_Y, μ_Y) will accumulate $\phi_{t_{N(x)}}^Y(x)$ on the right. Then according to Proposition 6.21, for $t > 0$ small enough we have that

$$F^{N(x)}(h_i^+((0, t) \times (-1, 1))) \subset h_i^+((-1, 0) \times (-1, 1)).$$

This implies that the typical points in Λ_Y also accumulate $\phi_{t_{N(x)}}^Y(x)$ on the left. \square

In the following, we will assume:

There is a neighborhood \mathcal{V}^* of Y such that for any $Z \in \mathcal{V}^*$, the basin of any sink of Z cannot accumulate on $C(\sigma_Z)$.

Corollary 6.23. For any $Y \in \mathcal{M}_n$, and $\sigma_{i,Y} \in \Lambda_Y \cap \text{Sing}(Y)$, if the right separatrix of $W^u(\sigma_{i,Y})$ is a homoclinic orbit Γ_i of $\sigma_{i,Y}$, then the left separatrix of $W^u(\sigma_{i,Y})$ is contained in Λ_Y .

Proof. If $\Gamma_i \subset \Lambda_Y$, then (Λ_Y, μ_Y) accumulates $\phi_{t_{N(x)}}^Y(x)$ on the left, which implies the left separatrix of $W^u(\sigma_{i,Y})$ is contained in Λ_Y by Corollary 6.22; otherwise, since $W^u(\sigma_{i,Y}) \setminus \sigma_{i,Y} \cap \Lambda_Y \neq \emptyset$, we have that the left separatrix of $W^u(\sigma_{i,Y})$ is contained in Λ_Y . \square

Corollary 6.24. For any $Y \in \mathcal{M}_n$, and the transitive \mathcal{N} -set $\Lambda_Y \subset C(\sigma_Y)$, if $\sigma_{i,Y} \in \Lambda_Y$ has two homoclinic orbits, then both the two homoclinic orbits are contained in Λ_Y .

Proof. Since Λ_Y is transitive and $\sigma_{i,Y} \in \Lambda_Y$, one separatrix of $W^u(\sigma_{i,Y})$ is contained in Λ_Y . We may assume that the right separatrix is contained in Λ_Y . According to Corollary 6.23, the left separatrix of $W^u(\sigma_{i,Y})$ is contained in Λ_Y , which means the left homoclinic orbit associated to $\sigma_{i,Y}$ is also contained in Λ_Y . \square

Proposition 6.25. Given $Y \in \mathcal{M}_n$, assume that Λ_Y contains a homoclinic orbit Γ_i of $\sigma_{i,Y}$. If $\Gamma_i \cap \ell_i^+ \neq \emptyset$, then $\Lambda_Y \cap \ell_i^- \neq \emptyset$, where $\ell_i^\pm = W_{loc}^s(\sigma_i) \cap S_i^\pm$, $\pm \in \{+, -\}$.

Proof. Suppose on the contrary, $\Lambda_Y \cap W_{loc}^s(\sigma_{i,Y}) \cap S_i^- = \emptyset$. Without loss of generality, we assume that Γ_i is the right separatrix of $W^u(\sigma_{i,Y})$. Since $\Lambda_Y \cap W_{loc}^s(\sigma_{i,Y}) \neq \emptyset$, we will have $\Lambda_Y \cap W_{loc}^s(\sigma_{i,Y}) \cap S_i^+ \neq \emptyset$. Since $\Gamma_i \subset \Lambda_Y$ and the ergodic measure μ_Y satisfies $\text{supp}(\mu_Y) = \Lambda_Y$, Γ_i can be approximated by typical points of μ_Y . As before, denote by

$$\mathcal{T}_x = \{t_i > 0 : \phi_{t_i}^Y(x) \in \Sigma\} = \{t_1 < t_2 < \cdots < t_{N(x)}\},$$

where $N(x) = \#\mathcal{T}_x$. We have that $\phi_{t_{N(x)}}^Y(x) \in S_i^+$. For the return map F , there are two cases:

1. The orientation preserving case: For $t > 0$ small enough, one has $F^{N(x)}(h_i^+((0, t) \times (-1, 1))) \subset S_i^{+,r}$.

2. The orientation reversing case: For $t > 0$ small enough, one has $F^{N(x)}(h_i^+((0, t) \times (-1, 1))) \subset S_i^{+,l}$.

In case 1, one will have

- For $t > 0$ small enough, one has $F^{N(x)}(h_i^-((0, t) \times (-1, 1))) \subset S_i^{+,l}$.
- Typical points of μ_Y in Λ_Y can not accumulate $\Gamma_i \cap W_{loc}^s(\sigma_{i,Y})$ on the right by Proposition 6.21.

Thus, typical points of μ_Y in Λ_Y accumulates $\Gamma_i \cap W_{loc}^s(\sigma_{i,Y})$ on the left. What we need to prove is that: for every point in Λ_Y which is close to x , then its backward iteration will intersect $S_i^+ \cup S_i^-$.

Claim. *Let $x_n \in \Lambda_Y \cap S_i^{+,l}$ such that $x_n \rightarrow \phi_{t_{N(x)}}^Y(x) \in l_i^+$. Then for n large enough $x_n \in F^{N(x)}(h_i^-((0, t) \times (-1, 1)))$.*




Figure 4: Illustration of the Claim

Proof of the claim: Let U be a small ball neighborhood of $\sigma_{i,Y}$. We may assume that $x \in U$ and $B = U \cap W_{loc}^s(\sigma_{i,Y})$ is a 2-disc, which is a neighborhood $\sigma_{i,Y}$ in $W_{loc}^s(\sigma_{i,Y})$. Moreover, we may assume that $l_i^\pm \subset B$. According to the construction of cross-section system, we know that

$$\left(B \setminus \overline{L_i^+ \cup L_i^-} \right) \cap C(\sigma_Y) = \emptyset,$$

where

$$L_i^\pm = \bigcup_{p \in l_i^\pm} \phi_{[t_p, +\infty)}^Y(p), \quad t_p = \min\{t : \phi_{[t, 0]}^Y(p) \subset B\}.$$

Denote by $y_n = \phi_{-t_{N(x)}}^Y(x_n)$. Then $\lim_{n \rightarrow \infty} y_n = x$. Let $\tau_n = \min\{t : \phi_{[t, 0]}^Y(y_n) \subset U\}$. Then $\lim_{n \rightarrow \infty} \tau_n = -\infty$. Let $z_n = \phi_{\tau_n}^Y(y_n)$. Then $z_n \in B \cap \partial U \cap \Lambda_Y$. Assume that $z = \lim_{n \rightarrow \infty} z_n$. Hence $z \in \partial B \cap \Lambda$. The positive orbit of z will intersect $l_i^+ \cup l_i^-$, where the claim follows. \square

According to the claim, we get a contradiction in Case 1.

In case 2, for $t > 0$ small enough, one has $F^{N(x)}(h_i^-((0, t) \times (-1, 1))) \subset S_i^{+,r}$. No matter typical points of μ_Y in Λ_Y accumulates $\Gamma_i \cap W_{loc}^s(\sigma_{i,Y})$ on the left or on the right, for any $t > 0$ small enough, $\Lambda_Y \cap h_i^-((0, t) \times (-1, 1)) \neq \emptyset$. Since Λ_Y is compact, one has $\Lambda_Y \cap \ell_i^- \neq \emptyset$. \square

Proposition 6.26. *For any $Y \in \mathcal{M}_n$, and the transitive set $\Lambda_Y \subset C(\sigma_Y)$, if $\sigma_{i,Y} \in \Lambda_Y$, then $\sigma_{i,Y}$ cannot have two homoclinic orbits.*

Proof. Suppose on the contrary, some $\sigma_{i,Y} \in \Lambda_Y$ has two homoclinic orbits. By Corollary 6.24, both homoclinic orbits are contained in Λ_Y . There are two cases:

- A. Two-side case: one homoclinic orbit intersects ℓ_i^+ , and the other intersects ℓ_i^- .
- B. One-side case: both homoclinic orbits intersect ℓ_i^+ (or ℓ_i^-).

We may have many subcases. For completeness, we will list all the cases here. We assume that $x_r \in W_{loc}^u(\sigma_{i,Y})$ is on the right separatrix and $x_l \in W_{loc}^u(\sigma_{i,Y})$ the left separatrix. For x_r and x_l , one can define

$$\begin{aligned} \mathcal{T}_r &= \{t > 0 : \phi_t^Y(x_r) \in \Sigma\} = \{t_1 < t_2 < \dots < t_{N_r}\}, \\ \mathcal{T}_l &= \{t > 0 : \phi_t^Y(x_l) \in \Sigma\} = \{\tau_1 < \tau_2 < \dots < t_{N_l}\}, \end{aligned}$$

where

$$N_r = \#\mathcal{T}_r, \quad N_l = \#\mathcal{T}_l.$$

1. The right separatrix of the unstable manifold intersect S_i^+ and the left separatrix of the unstable manifold intersect S_i^- (symmetrically, the right separatrix of the unstable manifold intersect S_i^- and the left separatrix of the unstable manifold intersect S_i^+).
 - Orientation-reversing: for small $s > 0$, $F^{N_r}(h_i^+((0, s) \times (-1, 1))) \subset S_i^{+,l}$ and $F^{N_l}(h_i^-((-s, 0) \times (-1, 1))) \subset S_i^{-,r}$.
 - Orientation-preserving: for small $s > 0$, $F^{N_r}(h_i^+((0, s) \times (-1, 1))) \subset S_i^{+,r}$ and $F^{N_l}(h_i^-((-s, 0) \times (-1, 1))) \subset S_i^{-,l}$.
 - Mixing: for small $s > 0$, $F^{N_r}(h_i^+((0, s) \times (-1, 1))) \subset S_i^{+,r}$ and $F^{N_l}(h_i^-((-s, 0) \times (-1, 1))) \subset S_i^{-,r}$; or $F^{N_r}(h_i^+((0, s) \times (-1, 1))) \subset S_i^{+,l}$ and $F^{N_l}(h_i^-((-s, 0) \times (-1, 1))) \subset S_i^{-,l}$.
2. Both the right separatrix and the left separatrix of the unstable manifolds intersect S_i^+ (symmetrically, both the right separatrix and the left separatrix of the unstable manifolds intersect S_i^-).
 - Orientation-reversing: for small $s > 0$, $F^{N_r}(h_i^+((0, s) \times (-1, 1))) \subset S_i^{+,l}$ and $F^{N_l}(h_i^+((-s, 0) \times (-1, 1))) \subset S_i^{+,r}$.
 - Orientation-preserving: for small $s > 0$, $F^{N_r}(h_i^+((0, s) \times (-1, 1))) \subset S_i^{+,r}$ and $F^{N_l}(h_i^+((-s, 0) \times (-1, 1))) \subset S_i^{+,l}$.

But in any case, the assumption of Proposition 6.20 will be satisfied. We only consider one case for instance. For the two-side case, without loss of generality, one can assume that the right separatrix of the unstable manifold intersect S_i^+ and the left separatrix of the unstable manifold intersect S_i^- .

We considers the following case:

- For any $s > 0$ small enough, $F^{N_r}(h_i^+((0, s) \times (-1, 1))) \subset S_i^{+,l}$ and $F^{N_l}(h_i^-((-s, 0) \times (-1, 1))) \subset S_i^{-,r}$.

Then for any $\pm \in \{+, -\}$, we have $F^{2(N_r+N_l)}(h_i^+((0, s) \times (-1, 1))) \subset h_i^+((0, 1) \times (-1, 1))$ and $F^{2(N_r+N_l)}(h_i^-((-s, 0) \times (-1, 1))) \subset h_i^-((-1, 0) \times (-1, 1))$. The typical point of μ_Y will accumulate local stable manifold of σ_i . Then we can apply Proposition 6.20 to get a contradiction. \square

Now we can give a contradiction and finish the proof of Theorem 6.2.

The contradiction. For any $Y \in \mathcal{M}_n$, and the transitive set $\Lambda_Y \subset C(\sigma_Y)$, according to Theorem 6.14, Λ_Y contains a singularity. Take a singularity $\rho = \rho_Y \in \Lambda_Y$. By Lemma 6.18, ρ has a homoclinic orbit Γ . We assume that Γ is the right separatrix of $W^u(\rho)$. By Proposition 6.26, the left separatrix of $W^u(\rho)$ is not a homoclinic orbit.

If $\Gamma \subset \Lambda_Y$, by Corollary 6.22, the left separatrix of $W^u(\rho)$ is contained in Λ_Y . We assume that $\Gamma \cap \ell^+ \neq \emptyset$. By Proposition 6.25, $\Lambda_Y \cap \ell^- \neq \emptyset$.

For any neighborhood \mathcal{U} of Y , by using Lemma 3.4 to a transitive orbit, we can get a C^2 vector field $Z \in \mathcal{U}$ without perturbing the homoclinic orbit of singularities of Y , which connects the orbit of the left separatrix of $W^u(\rho)$ and ℓ^- . Then Z has one more homoclinic orbit of ρ . And then by Theorem 3.10, there is a C^2 weak Kupka-Smale $Z' \in \mathcal{U}$ such that Z' has at least $(n + 1)$ homoclinic orbits of singularities. This fact contradicts the maximality of n . So, we proved that $\Gamma \not\subset \Lambda_Y$. Then by a similar argument, for any neighborhood \mathcal{U} of Y , there is a C^2 weak Kupka-Smale $Z \in \mathcal{U}$ which connects the orbit of the left separatrix of $W^u(\rho)$ and some point $a \in (\ell^+ \cup \ell^-) \cap \Lambda_Y$. Since $a \notin \Gamma$, Z also contains at least $(n + 1)$ homoclinic orbit of singularities. This finishes the proof of Theorem 6.2.

Acknowledgements.

We thank L. Wen for his comments and encouragements, S. Crovisier who listened the proof carefully. We also thank C. Bonatti, Y. Shi, Y. Zhang and X. Wang for useful discussion.

References

- [1] V. Afraïmovič, V. Bykov, and L. Silnikov, The origin and structure of the Lorenz attractor, *Dokl. Akad. Nauk SSSR*, **234** (1977), 336–339.
- [2] D. V. Anosov, Dynamical systems in the 1960s: The hyperbolic revolutions, *Mathematical events of the twentieth century*, Springer-Verlag, 2005.
- [3] M. Arnaud, Création de connexions en topologie C^1 , *Ergodic Theory Dynam. Systems*, **21**(2001), 339-381.
- [4] V. Araujo and M. Pacifico, Three-dimensional flows, Springer-Verlag, 2010.
- [5] A. Arroyo and F. Rodriguez Hertz, Homoclinic bifurcations and uniform hyperbolicity for threedimensional flows, *Ann. Inst. H. Poincaré-Anal. NonLinéaire*, **20**(2003), 805-841.
- [6] G. D. Birkhoff, Nouvelles recherches sur les systèmes dynamiques, *Mem. Pont. Acad. Sci. Nov. Lyncaei*, **53** (1935), 85-216.
- [7] C. Bonatti and S. Crovisier, Récurrence et généricité, *Invent. Math.*, **158** (2004), 33-104.

- [8] C. Bonatti and L. Díaz, On maximal transitive sets of generic diffeomorphisms, *Inst. Hautes études Sci. Publ. Math.*, **96**(2002), 171–197.
- [9] C. Bonatti, L. J. Díaz and E. R. Pujals, A C^1 -generic dichotomy for diffeomorphisms: Weak forms of hyperbolicity or infinitely many sinks or sources, *Annals of Math.* **158** (2003), 355–418.
- [10] C. Bonatti L. Díaz and M. Viana, Dynamics beyond uniform hyperbolicity. A global geometric and probabilistic perspective. *Encyclopaedia of Mathematical Sciences*, **102**. Mathematical Physics, III. Springer-Verlag, Berlin,(2005). xviii+384 pp.
- [11] C. Bonatti, S. Gan and L. Wen, On the existence of non-trivial homoclinic classes, *Ergodic Theory Dynam. Systems*, **27**(2007), 1473–1508.
- [12] C. Bonatti, S. Gan and D. Yang, Dominated chain recurrent classes with singularities, accepted by *Ann. Sc. Norm. Super. Pisa*, 2012.
- [13] C. Bonatti, N. Gourmelon and T. Vivier, Perturbations of the derivative along periodic orbits, *Ergodic Theory Dynam. Systems*, **26**(2006), 1307–1337.
- [14] C. Conley, *Isolated invariant sets and Morse index*, CBMS Regional Conference Series in Mathematics, **38**, AMS Providence, R.I., (1978).
- [15] S. Crovisier, Periodic orbits and chain transitive sets of C^1 -diffeomorphisms, *Publ. Math. Inst. Hautes Études Sci.*, **104**(2006), 87–141.
- [16] S. Crovisier, Birth of homoclinic intersections: a model for the central dynamics of partially hyperbolic systems, *Ann. Math.*, **172**(2010), 1641–1677.
- [17] S. Crovisier, M. Sambarino and D. Yang, Partial hyperbolicity and homoclinic tangencies, arXiv:1103.0869.
- [18] J. Franks, Necessary conditions for stability of diffeomorphisms, *Trans. Amer. Math. Soc.*, **158** (1971), 301–308.
- [19] J. Guckenheimer, A strange, strange attractor, *The Hopf bifurcation theorems and its applications (Applied Mathematical Series, 19)*, Springer-Verlag, 1976, 368–381.
- [20] J. Guckenheimer and R. Williams, Structural stability of Lorenz attractors, *Inst. Hautes Études Sci. Publ. Math.*, **50** (1979), 59–72.
- [21] S. Hayashi, Connecting invariant manifolds and the solution of the C^1 -stability and Ω -stability conjectures for flows, *Ann of Math.* **145**(1997), 81–137.
- [22] M. W. Hirsch; C. C. Pugh; M. Shub, Invariant manifolds, *Lecture Notes in Mathematics*, **583** Springer-Verlag 1977.
- [23] I. Kupka, Contribution à la théorie des champs génériques, *Contrib. Differ. Equ.*, **2**(1963), 457C484 and **3**(1964), 411–420.
- [24] M. Li, S. Gan and L. Wen, Robustly transitive singular sets via approach of extended linear Poincaré flow, *Discrete Contin. Dyn. Syst.* **13** (2005), 239–269.
- [25] S. Liao, A basic property of a certain class of differential systems, (Chinese) *Acta Math. Sinica*, **22**(1979), 316–343. (in chinese)
- [26] S. Liao, On the stability conjecture, *Chinese Annals of Math.*, **1**(1980), 9–30.

- [27] S. Liao, Obstruction sets II, *Acta Sci. Natur. Univ. Pekinensis*, **2** (1981), 1-36. (in chinese)
- [28] S. Liao, Certain uniformity properties of differential systems and a generalization of an existence theorem for periodic orbits, *Acta Sci. Natur. Univ. Pekinensis*, **2** (1981), 1-19. (in chinese)
- [29] S. Liao, On (η, d) -contractible orbits of vector fields, *Systems Science and Mathematical Sciences*, **2**(1989), 193-227.
- [30] E. N. Lorenz, Deterministic nonperiodic flow, *J. Atmosph. Sci.*, **20** (1963), 130–141.
- [31] R. Mañé, An ergodic closing lemma, *Ann. Math.*, **116**(1982), 503-540.
- [32] R. Mañé, Hyperbolicity, sinks and measure in one-dimensional dynamics, *Comm. Math. Phys.*, **100**(1985), 495-524.
- [33] C. Morales and M. Pacifico, Lyapunov stability of ω -limit sets, *Discrete Contin. Dyn. Syst.*, **8**(2002), 671-674.
- [34] C. Morales and M. Pacifico, A dichotomy for three-dimensional vector fields, *Ergodic Theory and Dynamical Systems*, **23**(2003), 1575-1600.
- [35] C. Morales, M. Pacifico, and E. Pujals, On C^1 robust singular transitive sets for three-dimensional flows, *C. R. Acad. Sci. Paris*, **326** (1998), 81–86.
- [36] C. Morales, M. Pacifico, and E. Pujals, Robust transitive singular sets for 3-flows are partially hyperbolic attractors or repellers, *Ann. of Math.*, **160** (2004), 375–432.
- [37] S. Newhouse, Nondensity of axiom A(a) on S^2 , *Global analysis I, Proc. Symp. Pure Math. AMS*, **14**(1970), 191-202.
- [38] S. Newhouse, Diffeomorphisms with infinitely many sinks, *Topology*, **13** (1974), 9–18.
- [39] S. Newhouse, The abundance of wild hyperbolic sets and nonsmooth stable sets for diffeomorphisms, *Inst. Hautes Études Sci. Publ. Math.*, **50** (1979), 101–151.
- [40] T. Ohno, A weak equivalence and topological entropy, *Publ. Res. Inst. Math. Sci.*, **16**(1980), 289-298.
- [41] J. Palis, On the C^1 Ω -stability conjecture. *Publ. Math. Inst. Hautes Études Sci.*, **66**(1988), 211-215.
- [42] J. Palis, Homoclinic bifurcations, sensitive-chaotic dynamics and strange attractors, *Dynamical systems and related topics (Nagoya, 1990)*, 466-472, Adv. Ser. Dynam. Systems, 9, *World Sci. Publ., River Edge, NJ*, 1991.
- [43] J. Palis, A global view of dynamics and a conjecture of the denseness of finitude of attractors, *Astérisque*, **261**(2000), 335–347.
- [44] J. Palis, A global perspective for non-conservative dynamics, *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **22**(2005), 485-507.
- [45] J. Palis, Open questions leading to a global perspective in dynamics, *Nonlinearity*, **21**(2008), 37-43.
- [46] J. Palis, and W. de Melo, Geometric theory of dynamical systems – an introduction, Springer-Verlag (1982).

- [47] M. M. Peixoto, Structural stability on two-dimensional manifolds, *Topology*, **1**(1962), 101-120.
- [48] V. Pliss, A hypothesis due to Smale, *Diff. Eq.* **8** (1972), 203-214.
- [49] H. Poincaré, Sur le problème des trois corps et les équations de la dynamique, *Acta Math.*, **13**(1890), 1-270.
- [50] H. Poincaré, *Les méthodes nouvelles de la mécanique céleste, Tome III*, Gauthier-Villars (1899).
- [51] C. Pugh, The closing lemma, *Amer. J. Math.*, **89**1967, 956–1009.
- [52] E. Pujals and M. Sambarino, Homoclinic tangencies and hyperbolicity for surface diffeomorphisms, *Annals of Math.*, **151** (2000), 961-1023.
- [53] L. P. Shil'nikov, Homoclinic Trajectories: From Poincaré to the present, *Mathematical events of the twentieth century*, Springer-Verlag, 2005.
- [54] S. Smale, Stable manifolds for differential equations and diffeomorphisms, *Ann. Sc. Norm. Super. Pisa*, **17**(1963), 97-116.
- [55] S. Smale, Diffeomorphisms with many periodic points. In: *Differential and Combinatorial Topology*. Princeton University Press, 1965, 63-80 (Princeton Math. Ser., 24).
- [56] W. Sun, T. Young and Y. Zhou, Topological entropies of equivalent smooth flows, *Trans. Amer. Math. Soc.*, **361**(2009), 3071-3082.
- [57] R. Thomas, Topological entropy of fixed-point free flows, *Trans. Amer. Math. Soc.*, **319**(1990), 601-618.
- [58] L. Wen, On the C^1 -stability conjecture for flows, *J. Differential Equations*, **129**(1996), 334-357.
- [59] L. Wen, Homoclinic tangencies and dominated splittings, *Nonlinearity* **15** (2002), 1445-1469.
- [60] L. Wen, Generic diffeomorphisms away from homoclinic tangencies and heterodimensional cycles, *Bull. Braz. Math. Soc. (N.S.)*, **35**(2004), 419-452.
- [61] L. Wen and Z. Xia, C^1 connecting lemmas, *Trans. Am. Math. Soc.*, **352**(2000), 5213-5230.
- [62] L. Wen, The selecting lemma of Liao, *Disc. Cont. Dynam. Syst.*, **20**(2008), 159-175.
- [63] R. Xi, Master thesis in Peking University, 2005.

Shaobo Gan

Laboratory of Mathematics and Applied Mathematics (LMAM),
 School of Mathematical Sciences
 Peking University, Beijing, 100871, P. R. China
 gansb@math.pku.edu.cn

Dawei Yang

School of Mathematics,
 Jilin University, Changchun, 130012, P.R. China
 yangdw1981@gmail.com