

DISCRETE SCHLESINGER TRANSFORMATIONS, THEIR HAMILTONIAN FORMULATION, AND DIFFERENCE PAINLEVÉ EQUATIONS

ANTON DZHAMAY, HIDETAKA SAKAI, AND TOMOYUKI TAKENAWA

ABSTRACT. Schlesinger transformations are algebraic transformations of a Fuchsian system that preserve its monodromy representation and act on the characteristic indices of the system by integral shifts. One of the important reasons to study such transformations is the relationship between Schlesinger transformations and discrete Painlevé equations; this is also the main theme behind our work. We derive *discrete Schlesinger evolution equations* describing discrete dynamical systems generated by elementary Schlesinger transformations and give their discrete Hamiltonian description w.r.t. the standard symplectic structure on the space of Fuchsian systems. As an application, we compute explicitly two examples of reduction from Schlesinger transformations to difference Painlevé equations. The first example, $d\text{-}P(D_4^{(1)})$ (or difference Painlevé V), corresponds to Bäcklund transformations for continuous P_{VI} . The second example, $d\text{-}P(A_2^{(1)*})$ (with the symmetry group $E_6^{(1)}$), is purely discrete. We also describe the role played by the geometry of the Okamoto space of initial conditions in comparing different equations of the same type.

1. INTRODUCTION

In the theory of ordinary linear differential equations on a complex domain, and in particular in the theory of Fuchsian systems, an important characteristic of the equation is its configuration of singularities and the characteristic indices at these singular points. Associated to this data is the notion of the monodromy representation of the equation. Roughly speaking, this representation describes how the fundamental solution matrix of the equation changes under analytic continuation along the closed paths around the singular points and it gives a significant insight into the global behavior of solutions of the equation.

The fruitful idea of deforming the equation by moving the location of critical points into different configurations, or by changing the characteristic indices, without changing its monodromy representation goes back to B. Riemann, but the actual foundations of this theory of *isomonodromic deformations* in the Fuchsian case were laid down in the works of R. Fuchs [Fuc07], L. Schlesinger [Sch12], and R. Garnier [Gar26]. An extension of the theory to the non-Fuchsian case was done much later in the series of papers by M. Jimbo, T. Miwa, and K. Ueno, [JMU81, JM82, JM81]. At present, the theory of isomonodromic deformations is a very active research field with deep connections to other areas such as the theory of integrable systems, classical theory of differential equations, and differential and algebraic geometry.

It is important to distinguish between continuous and discrete isomonodromic deformations. In the continuous case deformation parameters are locations of singular points of the system. The resulting isomonodromic flows on the space of coefficients of the Fuchsian system are given by Schlesinger *equations*. In [JMMS80] M. Jimbo, T. Miwa, Y. Mōri, and M. Sato showed that these equations, when considered as a dynamical system on the decomposition space of the Fuchsian system, can be written in the Hamiltonian form. For a 2×2 Fuchsian system with $n = 4$ poles, Schlesinger equations reduce to the most general of the Painlevé equations, Painlevé VI (this result goes back to the work of R. Fuchs, [Fuc07]). Painlevé equations are also Hamiltonian, due to the work of Okamoto, [Oka80], and in fact the reduction from Schlesinger equations to P_{VI} can be carried over on the level of Hamiltonians, see [Sak10]. Fuchsian systems with more than two accessory parameters give rise to higher order analogues of Painlevé equations, see the survey [KNS13] for the four-dimensional case.

It is also possible to deform the characteristic indices of the system, but in that case the isomonodromy condition requires that the indices change by integral shifts, and so the resulting dynamic is *discrete* and is

2010 *Mathematics Subject Classification*. 34M55, 34M56, 14E07.

Key words and phrases. Integrable systems, Painlevé equations, difference equations, isomonodromic transformations, birational transformations.

expressed in the form of *difference* equations called Schlesinger *transformations*. Similarly to the continuous case, these transformations reduce to discrete analogues of Painlevé-type equations and the study of this correspondence has been a major research topic in the field of discrete integrable systems over the last twenty years, [RGH91, PNGR92]. We need to remark here that Schlesinger transformations correspond to *difference* Painlevé equations, but the other two types of *discrete* Painlevé equations, *q-difference* and *elliptic-difference* Painlevé equations, can also be considered in a modification of this approach.

Discrete Painlevé equations share many properties with the differential Painlevé equations, e.g., the existence of special solutions such as algebraic solutions or solutions that can be expressed in terms of special functions, affine Weyl group symmetries, and the geometric classification of equations in terms of rational surfaces. Our goal for this paper is to describe, very explicitly, some aspects of such correspondence for the isomonodromic approach, following the outline above for the continuous case and adapting it to the so-called *elementary* Schlesinger transformations. We do this in three steps.

First, we give *explicit* evolution equations for the discrete dynamical system on the space of coefficient matrices that is generated by elementary Schlesinger transformations. We call such equations *difference Schlesinger evolution equations*. These equations make it easy to compute Schlesinger transformations and consequently to study reductions to difference Painlevé equations, since the usual computation of Schlesinger transformations from the compatibility condition between the Fuchsian equation and the deformation equation is often quite complicated. We should also mention here that in [Bor04] A. Borodin considered difference Schlesinger equations for isomonodromic deformations of *difference* linear systems. He showed that, under a certain continuous limit, such equations converge to implicit equations describing Schlesinger transformations of the corresponding continuous system.

Next, we use these equations to define the evolution dynamic on the decomposition space of the Fuchsian system and give the discrete Hamiltonian description of such flows. Since the Hamiltonian framework is not yet well-developed in the discrete case, having examples of such description is essential to better understanding of this formalism. Further, this is an essential first step towards developing the Hamiltonian theory of discrete Painlevé equations, as well as the Hamiltonian reduction procedure in the discrete case.

Finally, we consider explicit examples of reductions from Schlesinger transformations to difference Painlevé equations. In Sakai's geometric approach to Painlevé equations, difference equations are classified by the surface type (or symmetry type) as follows:

$$\begin{array}{rcccccccccccc}
 \text{Surface:} & A_0^{(1)**} & A_1^{(1)*} & A_2^{(1)*} & D_4^{(1)} & D_5^{(1)} & D_6^{(1)} & D_7^{(1)} & D_8^{(1)} & E_6^{(1)} & E_7^{(1)} & E_8^{(1)} \\
 \text{Symmetry:} & E_8^{(1)} & E_7^{(1)} & E_6^{(1)} & D_4^{(1)} & A_3^{(1)} & (A_1 + A_1)^{(1)} & A_1^{(1)} & - & A_2^{(1)} & A_1^{(1)} & -
 \end{array}$$

Difference Painlevé equations corresponding to surfaces of types D_l and E_l appear as Bäcklund transformations of differential Painlevé equations, but equations corresponding to surfaces of type A_l are purely discrete. Thus, we are particularly interested in the left side of the above list. In [Sak07] H. Sakai posed a problem of representing difference Painlevé equations with surface types $A_0^{(1)**}$, $A_1^{(1)*}$, and $A_2^{(1)*}$ as reductions of Schlesinger transformations of Fuchsian systems. In [Boa09] P. Boalch described the corresponding Fuchsian systems without computing the Schlesinger transformations and their reductions explicitly. Also, in [AB06] D. Arinkin and A. Borodin obtained difference Painlevé equation $d-P(A_2^{(1)*})$ from isomonodromy transformations of *difference* Fuchsian systems. In this paper we explicitly compute reductions from Schlesinger transformations to difference Painlevé equations $d-P(D_4^{(1)})$ (also known as difference Painlevé-V equation $d-P_V$) and $d-P(A_2^{(1)*})$. We start with a suitable parametrization for the space of Fuchsian system of the appropriate type (which for these examples is two-dimensional, and so we denote the parameters by x and y) and use the discrete Schlesinger evolution equations to explicitly compute the evolution $\bar{x} = \bar{x}(x, y)$ and $\bar{y} = \bar{y}(x, y)$. For this map we construct the Okamoto space of initial conditions that resolves the indeterminate points of the dynamic, verify that this surface is indeed of the required type, and then explicitly find the translation direction in the symmetry sub-lattice. We also compare the dynamic given by discrete Schlesinger transformations to other examples of difference Painlevé equations of the same type by first finding the isomorphism of blow-down structures of their Okamoto surfaces, and then comparing the corresponding translation direction.

The interesting feature of the $d-P(D_4^{(1)})$ example is that we can compute, for the same Fuchsian system and using the same coordinates, continuous deformations that are described by the Painlevé-VI equation P_{VI} ,

and discrete deformations that are described by the difference Painlevé-V equation $d-P_V$, which corresponds to the Bäcklund transformations of P_{VI} . We use this as a test case for our approach. The $d-P(A_2^{(1)*})$ example is new and is a partial answer to Sakai's question (we plan to consider reductions to difference Painlevé equations $d-P(A_1^{(1)*})$ and $d-P(A_0^{(1)**})$ in the subsequent publication). A particularly interesting feature of the second example, that we explain in detail in the main text, is that its translation direction is different from that of the standard form for this equation.

The text is organized as follows. In the next section we briefly set up the context of the problem and describe in more details some of the main objects. In Section 3 we derive the equations of the elementary Schlesinger transformations and in Section 4 we show how to write them in the Hamiltonian form. Section 5 is dedicated to the $d-P(D_4^{(1)})$ and $d-P(A_2^{(1)*})$ examples, and the final section is conclusions and discussion of the results.

Acknowledgements: This work started at the *Discrete Integrable Systems* program at the *Isaac Newton Institute for Mathematical Sciences* in Cambridge, UK. The authors are very grateful to the Institute for its hospitality and the stimulating working environment. H.S. was supported by Grant-in-Aid no. 24540205 of the Japan Society for the Promotion of Science. Part of this work was done when A.D. visited H.S. and T.T. in Tokyo, and he thanks the *Tokyo University of Marine Science and Technology* and the *University of Tokyo* for their hospitality. A.D. would also like to acknowledge the generous travel support by the *University of Northern Colorado* Provost Fund which was very important for the success of this collaboration.

2. PRELIMINARIES

2.1. Fuchsian Systems. Schlesinger transformations that we study originate from the deformation theory of Fuchsian systems. Recall that a *Fuchsian system* (or a *Fuchsian equation*) is a matrix linear differential equation on the Riemann sphere \mathbb{CP}^1 such that all of its singular points are *regular singular points*. We consider a generic case when a Fuchsian system can be written in the *Schlesinger normal form*, i.e., when its coefficient matrix is a rational function with at most simple poles at some (distinct) points z_1, \dots, z_n (and possibly at the point $z_0 = \infty$),

$$(2.1) \quad \frac{d\mathbf{Y}}{dz} = \mathbf{A}(z)\mathbf{Y} = \left(\sum_{i=1}^n \frac{\mathbf{A}_i}{z - z_i} \right) \mathbf{Y},$$

Here $\mathbf{A}_i = \text{res}_{z_i} \mathbf{A}(z) dz$ are constant $m \times m$ matrices. We also put $\mathbf{A}_\infty = \text{res}_\infty \mathbf{A}(z) dz = -\sum_{i=1}^n \mathbf{A}_i$ and so this equation is regular at infinity iff $\mathbf{A}_\infty = \mathbf{0}$.

Assumption 2.1. From now on we make an additional semi-simplicity assumption that the coefficient matrices \mathbf{A}_i are *diagonalizable*.

2.2. Spectral Type and Accessory Parameters. Geometrically, Schlesinger (or isomonodromic) dynamic takes place on the space of coefficients $\mathbf{A}(z)$ of the Fuchsian system (2.1), considered modulo gauge transformations. Here we briefly outline the description of this space following [Sak10].

First, given $\mathbf{A}(z)$, we separate the locations of singular points z_i (that we think of as parameters of the dynamic) from the residue matrices \mathbf{A}_i at those points. Thus, we define

$$\tilde{\mathcal{F}}(\mathbf{z}) = \tilde{\mathcal{F}}(z_1, \dots, z_n) = \{(\mathbf{A}_1, \dots, \mathbf{A}_n) \mid \mathbf{A}_i \in \text{End}(\mathbb{C}^m)\}$$

to be the set of all Fuchsian systems with possible singularities at the points z_i . Further, since Schlesinger dynamic either preserves (in the continuous case) or shifts (in the discrete case) the eigenvalues of \mathbf{A}_i , we treat the eigenvalues as the parameters of the dynamic as well. Thus, the appropriate phase space is a quotient of the fiber of the eigenvalue map by gauge transformations. Local coordinates on the phase space are called *accessory parameters*. Note that the space \mathcal{A} of accessory parameters is quite complicated and its dimension depends on the spectral type of $\mathbf{A}(z)$ that we define next.

Spectral type of $\mathbf{A}(z)$ encodes the degeneracy of eigenvalues θ_i^j of the coefficient matrices \mathbf{A}_i (θ_i^j are also called the *characteristic indices* of the system) via partitions $m = m_i^1 + \dots + m_i^{l_i}$, $m_i^1 \geq \dots \geq m_i^{l_i} \geq 1$, where m is the matrix size and m_i^j denotes the multiplicities of the eigenvalues of \mathbf{A}_i . The *spectral type* \mathbf{m} of $\mathbf{A}(z)$ is then defined to be the collection of these partitions for all indices (including ∞):

$$\mathbf{m} = m_1^1 \cdots m_1^{l_1}, m_2^1 \cdots m_2^{l_2}, \dots, m_n^1 \cdots m_n^{l_n}, m_\infty^1 \cdots m_\infty^{l_\infty}.$$

It can be used to classify Fuchsian systems up to isomorphisms and the operations of addition and middle convolution introduced by N. Katz [Kat96], as in the recent work by T. Oshima [Osh13].

Assumption 2.2. Since for a Fuchsian system we can use scalar local gauge transformation of the form $\tilde{\mathbf{Y}}(z) = w(z)^{-1}\mathbf{Y}(z)$, where $w(z)$ is a solution of the *scalar* equation

$$\frac{dw}{dz} = \sum_{i=1}^n \frac{\theta_i^j}{z - z_i} w,$$

to change the residue matrices by $\tilde{\mathbf{A}}_i = \mathbf{A}_i - \theta_i^j \mathbf{1}$, we can assume, without loss of generality, that one of the eigenvalues $\theta_i^j = 0$. Thus, we always assume that the eigenvalue θ_i^j of the highest multiplicity m_i^1 is zero and we denote the corresponding subset of $\tilde{\mathcal{F}}(\mathbf{z})$ of *reduced* Fuchsian equations by $\mathcal{F}(\mathbf{z})$. This amounts to choosing a representative in the quotient space of all Fuchsian equations by the group of local *scalar* gauge transformations.

In view of **Assumptions 2.1** and **2.2**, \mathbf{A}_i is similar to a diagonal matrix $\text{diag}\{\theta_i^1, \dots, \theta_i^{r_i}, 0, \dots, 0\}$, where $r_i = \text{rank}(\mathbf{A}_i)$. Omitting the zero eigenvalues, we put

$$\Theta_i = \text{diag}\{\theta_i^1, \dots, \theta_i^{r_i}\}.$$

Denote by $\tilde{\mathcal{T}}$ the set of all possible diagonal matrices of the spectral type \mathbf{m} with the highest degeneracy eigenvalues at *finite* points set to 0 and omitted, as above. Thus, $\tilde{\mathcal{T}} = \{\Theta = (\Theta_1, \dots, \Theta_n; \Theta_\infty)\}$ and $r_i = m - m_i^1$. Then we have the eigenvalue map $\text{Sp} : \mathcal{F}(\mathbf{z}) \rightarrow \tilde{\mathcal{T}}$ from the set of reduced Fuchsian equations to the set of characteristic indices, where $\mathcal{T} \subset \tilde{\mathcal{T}}$ is the subset of Θ satisfying the *Fuchs relation* (or the trace condition)

$$\sum_{i=1}^{n,\infty} \sum_{j=1}^{r_i} \theta_i^j = 0.$$

With these definitions, given $\Theta \in \mathcal{T}$, the fiber of the eigenvalue map is $\mathcal{F}(\mathbf{z})_\Theta = \text{Sp}^{-1}(\Theta)$.

We still need to take into account global similarity transformations. It is convenient to do so in two steps. First we *normalize* our equation at infinity by reducing \mathbf{A}_∞ to a particular form, e.g., making it diagonal, and second we take the quotient by the stabilizer subgroup $G_{\mathbf{A}_\infty}$ of the group $\mathbb{G}\mathbb{L}_m$ of global similarity transformations. Thus, for a given $\Theta \in \mathcal{T}$, we *fix* \mathbf{A}_∞ such that $\text{Sp}(\mathbf{A}_\infty) = \Theta_\infty$ and denote by $\mathcal{F}(\mathbf{z})_{(\Theta, \mathbf{A}_\infty)}$ the subset of all Fuchsian equations in $\mathcal{F}(\mathbf{z})_\Theta$ satisfying the condition $\mathbf{A}_1 + \dots + \mathbf{A}_n = -\mathbf{A}_\infty$. Then, finally, our phase space is

$$\mathcal{A} = \mathcal{A}_{(\Theta, \mathbf{A}_\infty)} = \mathcal{F}(\mathbf{z})_{(\Theta, \mathbf{A}_\infty)} / G_{\mathbf{A}_\infty}.$$

When \mathbf{A}_∞ is diagonal, which is the case that we are mainly interested in, the dimension of this space \mathcal{A} of accessory parameters of the spectral type \mathbf{m} is given by the formula

$$\dim \mathcal{A} = (n-1)m^2 - \sum_{i=1}^{n,\infty} \left(\sum_{j=1}^{l_i} (m_i^{l_i})^2 \right) + 2.$$

2.3. The Decomposition Space. To study the Hamiltonian structure of Schlesinger equations, and also of Schlesinger transformations, it is convenient to consider a larger space that we call the *decomposition space*. It is defined as follows.

In view of **Assumption 2.1**, there exist full sets of *right* eigenvectors $\mathbf{b}_{i,j}$, $\mathbf{A}_i \mathbf{b}_{i,j} = \theta_i^j \mathbf{b}_{i,j}$, and *left* eigenvectors $\mathbf{c}_i^{j\dagger}$, $\mathbf{c}_i^{j\dagger} \mathbf{A}_i = \theta_i^j \mathbf{c}_i^{j\dagger}$, (here we use the \dagger symbol to indicate a *row*-vector). In the matrix form, omitting vectors with indices $j > r_i$ that are in the kernel of \mathbf{A}_i , we can write

$$\mathbf{B}_i = [\mathbf{b}_{i,1} \cdots \mathbf{b}_{i,r_i}], \quad \mathbf{A}_i \mathbf{B}_i = \mathbf{B}_i \Theta_i, \quad \mathbf{C}_i^\dagger = \begin{bmatrix} \mathbf{c}_i^{1\dagger} \\ \vdots \\ \mathbf{c}_i^{r_i\dagger} \end{bmatrix}, \quad \mathbf{C}_i^\dagger \mathbf{A}_i = \Theta_i \mathbf{C}_i^\dagger,$$

with Θ_i defined by (2.2). Then we have a decomposition $\mathbf{A}_i = \mathbf{B}_i \mathbf{C}_i^\dagger$, provided that $\mathbf{C}_i^\dagger \mathbf{B}_i = \Theta_i$. This condition is related to the normalization ambiguity of the eigenvectors.

Thus, given \mathbf{A}_i , we can construct (in a non-unique way) a corresponding decomposition pair $(\mathbf{B}_i, \mathbf{C}_i^\dagger)$. The space of all such pairs for all finite indices $1 \leq i \leq n$, without any additional conditions, is our *decomposition space*. We denote it as

$$\mathcal{B} \times \mathcal{C} = (\mathbb{C}^{r_1} \times \cdots \times \mathbb{C}^{r_n}) \times ((\mathbb{C}^{r_1})^\dagger \times \cdots \times (\mathbb{C}^{r_n})^\dagger) \simeq (\mathbb{C}^{r_1} \times (\mathbb{C}^{r_1})^\dagger) \times \cdots \times (\mathbb{C}^{r_n} \times (\mathbb{C}^{r_n})^\dagger),$$

since it is convenient to write an element $(\mathbf{B}, \mathbf{C}^\dagger)$ of this space as a list of n pairs $(\mathbf{B}_1, \mathbf{C}_1^\dagger; \cdots; \mathbf{B}_n, \mathbf{C}_n^\dagger)$. The decomposition space can be equipped with the following natural symplectic structure. We define the symplectic form as

$$\omega = \sum_{i=1}^n \text{tr}(d\mathbf{C}_i^\dagger \wedge d\mathbf{B}_i),$$

and so the canonical coordinates are the matrix elements of \mathbf{B}_i and \mathbf{C}_i^\dagger .

Remark 2.3. To simplify the notation we use the same symbols, e.g., \mathbf{B}_i or $\mathbf{b}_{i,j}$, to denote both the canonical coordinate system on the decomposition space and an actual point in the space; the exact meaning of the symbol is always clear from the context.

The idea of using the decomposition space $\mathcal{B} \times \mathcal{C}$ to study the Hamiltonian structure of isomonodromic deformations goes back to the paper by M. Jimbo, T. Miwa, Y. Môri, and M. Sato [JMMS80]. Since then it has been used by many other researchers studying isomonodromic deformations, most notably by J. Harnad, see, e.g., [Har94].

Remark 2.4. There are two natural actions on the decomposition space $\mathcal{B} \times \mathcal{C}$. First, the group \mathbb{GL}_m of global gauge transformations of the Fuchsian system induces the following action. Given $\mathbf{P} \in \mathbb{GL}_m$, we have the action $\mathbf{A}_i \mapsto \mathbf{P}\mathbf{A}_i\mathbf{P}^{-1}$ which translates into the action $(\mathbf{B}_i, \mathbf{C}_i^\dagger) \mapsto (\mathbf{P}\mathbf{B}_i, \mathbf{C}_i^\dagger\mathbf{P}^{-1})$. We refer to such transformations as *similarity transformations*. It is often necessary to restrict this action to the subgroup $G_{\mathbf{A}_\infty}$ preserving the form of \mathbf{A}_∞ . Second, for any pair $(\mathbf{B}_i, \mathbf{C}_i^\dagger)$ the pair $(\mathbf{B}_i\mathbf{Q}_i, \mathbf{Q}_i^{-1}\mathbf{C}_i^\dagger)$ determines the same matrix \mathbf{A}_i for $\mathbf{Q}_i \in \mathbb{GL}_{r_i}$. The condition $\mathbf{Q}_i^{-1}\mathbf{C}_i^\dagger\mathbf{B}_i\mathbf{Q}_i = \mathbf{Q}_i^{-1}\Theta_i\mathbf{Q}_i = \Theta_i$ restricts \mathbf{Q}_i to the stabilizer subgroup G_{Θ_i} of \mathbb{GL}_{r_i} . In particular, when all θ_i^j are distinct, \mathbf{Q}_i has to be a diagonal matrix. We refer to such transformations as *trivial transformations*. These two actions obviously commute with each other.

We are now ready to give an alternative description of the space of accessory parameters $\mathcal{A}_{(\Theta, \mathbf{A}_\infty)}$. Given the pair $(\Theta, \mathbf{A}_\infty)$ as in the previous section, let

$$(\mathcal{B} \times \mathcal{C})_{(\Theta, \mathbf{A}_\infty)} = \{(\mathbf{B}_1, \mathbf{C}_1^\dagger; \cdots; \mathbf{B}_n, \mathbf{C}_n^\dagger) \in \mathcal{B} \times \mathcal{C} \mid \mathbf{C}_i^\dagger\mathbf{B}_i = \Theta_i, \sum_{i=1}^n \mathbf{B}_i\mathbf{C}_i^\dagger = -\mathbf{A}_\infty\} \subset \mathcal{B} \times \mathcal{C}.$$

Then

$$\mathcal{A} = \mathcal{A}_{(\Theta, \mathbf{A}_\infty)} = (\mathcal{B} \times \mathcal{C})_{(\Theta, \mathbf{A}_\infty)} / ((G_{\Theta_1} \times \cdots \times G_{\Theta_n}) \times G_{\mathbf{A}_\infty}).$$

The following diagram illustrates the relationship between different spaces defined in these two sections:




FIGURE 1. The decomposition space and the space of accessory parameters

2.4. Schlesinger Equations and Schlesinger Transformations. Consider now isomonodromic deformations of a Fuchsian system. For the continuous case we take the deformation parameters to be locations of critical points. Thus, we let $\mathbf{A} = \mathbf{A}(z; \mathbf{z}) = \mathbf{A}(z; z_1, \dots, z_n)$ (i.e., $\mathbf{A}_i = \mathbf{A}_i(\mathbf{z})$). In [Sch12] L. Schlesinger showed that isomonodromic deformations are described, if we use the modern terminology, by the following *differential-differential Lax pair*:

$$\begin{cases} \frac{d\mathbf{Y}}{dz} = \mathbf{A}(z; \mathbf{z})\mathbf{Y} = \left(\sum_{i=1}^n \frac{\mathbf{A}_i(\mathbf{z})}{z - z_i} \right) \mathbf{Y}. \\ \frac{\partial \mathbf{Y}}{\partial z_i} = -\frac{\mathbf{A}_i(\mathbf{z})}{z - z_i} \mathbf{Y}, \quad i = 1, \dots, n. \end{cases}$$

Compatibility conditions for these deformation equations are called the *Schlesinger equations*. These are partial differential equations on the coefficient matrices $\mathbf{A}_i(\mathbf{z})$ and they have the form

$$(2.2) \quad \frac{\partial \mathbf{A}_j}{\partial z_i} = \frac{[\mathbf{A}_j, \mathbf{A}_i]}{z_j - z_i} \quad (j \neq i), \quad \frac{\partial \mathbf{A}_i}{\partial z_i} = -\sum_{j \neq i} \frac{[\mathbf{A}_j, \mathbf{A}_i]}{z_j - z_i}.$$

In [JMMS80] Jimbo *et.al* showed that Schlesinger equations can be written as a Hamiltonian system on the decomposition space $\mathcal{B} \times \mathcal{C}$,

$$\frac{\partial \mathbf{B}_i}{\partial z_j} = \frac{\partial \mathcal{H}_j}{\partial \mathbf{C}_i^\dagger}, \quad \frac{\partial \mathbf{C}_i^\dagger}{\partial z_j} = -\frac{\partial \mathcal{H}_j}{\partial \mathbf{B}_i},$$

with the Hamiltonian

$$\mathcal{H}_j = \mathcal{H}_j(\mathbf{B}, \mathbf{C}^\dagger) = \sum_{i \neq j} \frac{\text{tr}(\mathbf{A}_j \mathbf{A}_i)}{z_j - z_i}, \quad \mathbf{A}_i = \mathbf{B}_i \mathbf{C}_i^\dagger.$$

Schlesinger equations admit reductions to Painlevé-type nonlinear differential equations, which remains valid on the level of the Hamiltonians as well; we review an example of P_{VI} in Section 5. The main question that we consider is what happens to this picture in the *discrete* case.

The discrete counterparts of Schlesinger equations are Schlesinger transformations. Those are rational transformations preserving the singularity structure and the monodromy data of the Fuchsian system (2.1), except for the integral shifts in the characteristic indices θ_i^j . The study of such transformations again goes back to Schlesinger [Sch12]. This, and the more general case of irregular singular points, was considered in great detail in [JM81]. Schlesinger transformations are given by the following *differential-difference Lax Pair*:

$$\begin{cases} \frac{d\mathbf{Y}}{dz} = \mathbf{A}(z; \Theta)\mathbf{Y} = \left(\sum_{i=1}^n \frac{\mathbf{A}_i(\Theta)}{z - z_i} \right) \mathbf{Y}, \\ \bar{\mathbf{Y}}(z) = \mathbf{R}(z)\mathbf{Y}(z). \end{cases},$$

where $\mathbf{R}(z)$ is a specially chosen rational matrix function called the *multiplier* of the transformation. The coefficient matrix $\mathbf{A}(z; \Theta)$ of our Fuchsian system then transforms to $\bar{\mathbf{A}}(z; \Theta)$ that is related to $\mathbf{A}(z; \Theta)$ by the equation

$$(2.3) \quad \bar{\mathbf{A}}(z; \Theta)\mathbf{R}(z) = \mathbf{R}(z)\mathbf{A}(z; \Theta) + \frac{d\mathbf{R}(z)}{dz}.$$

In this paper we focus on the *elementary* Schlesinger transformations that only change two of the characteristic indices by unit shifts, i.e., $\bar{\theta}_\alpha^\mu = \theta_\alpha^\mu - 1$ and $\bar{\theta}_\beta^\nu = \theta_\beta^\nu + 1$, $\alpha \neq \beta$. For the examples section, we also assume θ_α^μ and θ_β^ν have no multiplicities. In [JM81] such elementary transformation are denoted by $\left\{ \begin{smallmatrix} \alpha & \beta \\ \mu & \nu \end{smallmatrix} \right\}$. For such transformations we solve the equations (2.3) to obtain explicit discrete evolution equations for the coefficient matrices $\mathbf{A}_i(\Theta)$, we call such equations *discrete Schlesinger evolution equations* since they are the exact discrete analogue of equations (2.2), see Theorem 3.1. In Theorem 3.3 we lift these equations to get the dynamic on the decomposition space and in Theorem 4.2 we represent these equation as a discrete Hamiltonian system, in the sense explained in the next section, using the same canonical coordinates as [JMMS80]. We explicitly construct a discrete Hamiltonian function $\mathcal{H}^+(\mathbf{B}, \bar{\mathbf{C}}^\dagger)$ of the decomposition space

that generates our dynamic. Finally, we study reductions of this dynamic to the space of accessory parameters. The reduced dynamic is very complicated and is given by Painlevé-type equations. Schematically, the situation is as in Figure 2:




FIGURE 2. Schlesinger Transformations

Unfortunately, at present we do not have a discrete version of the Hamiltonian reduction procedure from the decomposition space to the space of accessory parameters. This is a very important problem for future research.

2.5. Discrete Lagrangian and Hamiltonian Formalism. In developing a discrete version of the Hamiltonian formalism, one approach is to discretize the continuous dynamic preserving its integrability properties, see the recent encyclopedic book by Y. Suris, [Sur03], but it requires underlying continuous dynamical system. An alternative and more direct procedure is to develop a discrete version based on the variational principles. In this case, the Lagrangian formalism is more natural, but it is possible to extend it to include the discrete version of the Hamiltonian formalism as well. This approach has its origin in the optimal control theory (see, e.g., B. Jordan and E. Polak [JP64], and J. Cadzow [Cad70]) and mechanics (see e.g., J. Logan [Log73] and S. Maeda [Mae82]). In the theory of integrable systems it was first used in the foundational works by A. Veselov [Ves91, Ves88] and A. Veselov and J. Moser [MV91]. A lot of recent work in this field is motivated by the development of very effective *symplectic integrators*, see an excellent review paper by J. Marsden and M. West [MW01] (and the references therein), as well as more recent works by S. Lall and M. West [LW06] and A. Bloch, M. Leok, and T. Oshawa [OBL11] that emphasize the Hamiltonian aspects of the theory. Below we give a *very* brief outline of this approach closely following [OBL11], we refer to this and other publications above for details.

Let \mathcal{Q} be our configuration space. Then, in the discrete case, the *Lagrangian* $\mathcal{L} \in \mathcal{F}(\mathcal{Q} \times \mathcal{Q})$ is a function on the *state space* $\mathcal{Q} \times \mathcal{Q}$. For the discrete time parameter $k \in \mathbb{Z}$, a trajectory of motion is a map $\mathbf{q} : \mathbb{Z} \rightarrow \mathcal{Q}$, $\mathbf{q}(k) =: \mathbf{q}_k$, or, equivalently, a sequence $\{\mathbf{q}_k \in \mathcal{Q}\}$. The action functional on the space of such sequences is given by $\mathfrak{S}(\{\mathbf{q}_k\}) = \sum_k \mathcal{L}(\mathbf{q}_k, \mathbf{q}_{k+1})$. Using the variational principle $\delta \mathfrak{S} = 0$ we obtain *discrete Euler-Lagrange* equations that have the form

$$D_2 \mathcal{L}(\mathbf{q}_{k-1}, \mathbf{q}_k) + D_1 \mathcal{L}(\mathbf{q}_k, \mathbf{q}_{k+1}) = \mathbf{0},$$

where D_1 (resp. D_2) denote vectors of partial derivatives w.r.t. the first (resp. second) sets of local coordinates on the state space $\mathcal{Q} \times \mathcal{Q}$. These equations then implicitly determine the map (or more precisely, the correspondence) $\mathbf{q}_{k+1} = \phi(\mathbf{q}_{k-1}, \mathbf{q}_k)$, which then defines the discrete Lagrangian flow $F_{\mathcal{L}} : \mathcal{Q} \times \mathcal{Q} \rightarrow \mathcal{Q} \times \mathcal{Q}$ on the state space via $F_{\mathcal{L}}(\mathbf{q}_{k-1}, \mathbf{q}_k) = (\mathbf{q}_k, \mathbf{q}_{k+1})$. This flow is symplectic w.r.t. the discrete Lagrangian symplectic form

$$\Omega_{\mathcal{L}}(\mathbf{q}_k, \mathbf{q}_{k+1}) := d\vartheta_{\mathcal{L}}^+ = d\vartheta_{\mathcal{L}}^- = D_1 D_2 \mathcal{L}(\mathbf{q}_k, \mathbf{q}_{k+1}) d\mathbf{q}_k \wedge d\mathbf{q}_{k+1},$$

$F_{\mathcal{L}}^* \Omega_{\mathcal{L}} = \Omega_{\mathcal{L}}$, where one-forms $\vartheta_{\mathcal{L}}^{\pm} : \mathcal{Q} \times \mathcal{Q} \rightarrow T^*(\mathcal{Q} \times \mathcal{Q})$ are defined by

$$\vartheta_{\mathcal{L}}^+(\mathbf{q}_k, \mathbf{q}_{k+1}) = D_2 \mathcal{L}(\mathbf{q}_k, \mathbf{q}_{k+1}) d\mathbf{q}_{k+1}, \quad \vartheta_{\mathcal{L}}^-(\mathbf{q}_k, \mathbf{q}_{k+1}) = -D_1 \mathcal{L}(\mathbf{q}_k, \mathbf{q}_{k+1}) d\mathbf{q}_k.$$

If we introduce *right* and *left discrete Legendre transforms* $\mathbb{F}\mathcal{L}^{\pm} : \mathcal{Q} \times \mathcal{Q} \rightarrow T^*\mathcal{Q}$ and the *momenta* variables $\mathbf{p}_k, \mathbf{p}_{k+1}$ by

$$\begin{aligned} \mathbb{F}\mathcal{L}^+(\mathbf{q}_k, \mathbf{q}_{k+1}) &= (\mathbf{q}_{k+1}, \mathbf{p}_{k+1}) = (\mathbf{q}_{k+1}, D_2 \mathcal{L}(\mathbf{q}_k, \mathbf{q}_{k+1})), \\ \mathbb{F}\mathcal{L}^-(\mathbf{q}_k, \mathbf{q}_{k+1}) &= (\mathbf{q}_k, \mathbf{p}_k) = (\mathbf{q}_k, -D_1 \mathcal{L}(\mathbf{q}_k, \mathbf{q}_{k+1})), \end{aligned}$$

we see that $\vartheta_{\mathcal{L}}^{\pm} = (\mathbb{F}\mathcal{L}^{\pm})^* \vartheta$ and $\Omega_{\mathcal{L}} = (\mathbb{F}\mathcal{L}^{\pm})^* \Omega$, where ϑ and Ω are the standard Liouville and symplectic forms on $T^*\mathcal{Q}$ respectively.

We can then define the discrete Hamiltonian flow $\tilde{F}_{\mathcal{L}} : T^*\mathcal{Q} \rightarrow T^*\mathcal{Q}$ by $\tilde{F}_{\mathcal{L}}(\mathbf{q}_k, \mathbf{p}_k) = (\mathbf{q}_{k+1}, \mathbf{p}_{k+1})$, i.e., $\tilde{F}_{\mathcal{L}} = \mathbb{F}\mathcal{L}^+ \circ (\mathbb{F}\mathcal{L}^-)^{-1}$. But then the equations

$$\mathbf{p}_k = -D_1 \mathcal{L}(\mathbf{q}_k, \mathbf{q}_{k+1}), \quad \mathbf{p}_{k+1} = D_2 \mathcal{L}(\mathbf{q}_k, \mathbf{q}_{k+1}),$$

mean that $\mathcal{L}(\mathbf{q}_k, \mathbf{q}_{k+1})$ is just the generating function (of type one, see [GPS00] for the terminology) of the canonical transformation $\tilde{F}_{\mathcal{L}}$. We can then define the *right* and *left discrete Hamiltonian functions* as generating functions of this canonical transformation of type two and three respectively. Namely, *right discrete Hamiltonian* is

$$\mathcal{H}^+(\mathbf{q}_k, \mathbf{p}_{k+1}) := \mathbf{p}_{k+1} \mathbf{q}_{k+1} - \mathcal{L}(\mathbf{q}_k, \mathbf{q}_{k+1}),$$

and then the map $\tilde{F}_{\mathcal{L}} : T^*\mathcal{Q} \rightarrow T^*\mathcal{Q}$ is given (implicitly) by the *right discrete Hamiltonian equations*

$$\mathbf{q}_{k+1} = D_2 \mathcal{H}^+(\mathbf{q}_k, \mathbf{p}_{k+1}), \quad \mathbf{p}_k = D_1 \mathcal{H}^+(\mathbf{q}_k, \mathbf{p}_{k+1}).$$

Similarly, *left discrete Hamiltonian*

$$\mathcal{H}^-(\mathbf{q}_{k+1}, \mathbf{p}_k) := -\mathbf{p}_k \mathbf{q}_k - \mathcal{L}(\mathbf{q}_k, \mathbf{q}_{k+1})$$

gives *left discrete Hamiltonian equations*

$$\mathbf{q}_k = -D_1 \mathcal{H}^-(\mathbf{q}_{k+1}, \mathbf{p}_k), \quad \mathbf{p}_{k+1} = -D_2 \mathcal{H}^-(\mathbf{q}_{k+1}, \mathbf{p}_k).$$

For completeness, we mention that the generating function of type four,

$$\begin{aligned} \mathcal{R}(\mathbf{p}_k, \mathbf{p}_{k+1}) &:= \mathbf{q}_k \mathbf{p}_k - \mathbf{q}_{k+1} \mathbf{p}_{k+1} + \mathcal{L}(\mathbf{q}_k, \mathbf{q}_{k+1}) = \mathbf{q}_k \mathbf{p}_k - \mathcal{H}^+(\mathbf{q}_k, \mathbf{p}_{k+1}) = -\mathbf{q}_{k+1} \mathbf{p}_{k+1} - \mathcal{H}^-(\mathbf{q}_{k+1}, \mathbf{p}_k), \\ \mathbf{q}_k &= D_1 \mathcal{R}(\mathbf{p}_k, \mathbf{p}_{k+1}), \quad \mathbf{q}_{k+1} = D_2 \mathcal{L}(\mathbf{q}_k, \mathbf{q}_{k+1}), \end{aligned}$$

is nothing but the Lagrangian function on the space of momenta satisfying the discrete Euler-Lagrange equations

$$D_2 \mathcal{R}(\mathbf{p}_{k-1}, \mathbf{p}_k) + D_1 \mathcal{R}(\mathbf{p}_k, \mathbf{p}_{k+1}) = \mathbf{0}.$$

2.6. Elementary Divisors. For elementary Schlesinger transformations $\left\{ \begin{smallmatrix} \alpha & \beta \\ \mu & \nu \end{smallmatrix} \right\}$ the multiplier matrix $\mathbf{R}(z)$ has a very special simple form. In this section we describe some of the properties of such matrices that we need for our computations. Consider a rational matrix function

$$\mathbf{R}(z) = \mathbf{I} + \frac{\mathbf{G}}{z - z_0}, \quad \text{where } \mathbf{G} = \mathbf{f} \mathbf{g}^\dagger \text{ is a matrix of rank 1.}$$

Because of their role in representing general matrix functions as a product of factors of this form, such matrices are sometimes called elementary divisors [Dzh09]. Assuming that $\text{tr}(\mathbf{G}) = \mathbf{g}^\dagger \mathbf{f} \neq 0$, which is the case we need, we can consider instead of \mathbf{G} a corresponding rank-one projector $\mathbf{P} = \mathbf{f}(\mathbf{g}^\dagger \mathbf{f})^{-1} \mathbf{g}^\dagger$, $\mathbf{P}^2 = \mathbf{P}$. Then $\mathbf{G} = (\mathbf{g}^\dagger \mathbf{f}) \mathbf{P} = (z_0 - \zeta_0) \mathbf{P}$, where z_0 and ζ_0 are the zero and the pole of $\det \mathbf{R}(z)$ respectively. We also put $\mathbf{Q} = \mathbf{I} - \mathbf{P}$ to be the complementary projector, $\mathbf{Q}^2 = \mathbf{Q}$ and $\mathbf{P}\mathbf{Q} = \mathbf{Q}\mathbf{P} = \mathbf{0}$.

Lemma 2.5. *Let*

$$\mathbf{R}(z) = \mathbf{I} + \frac{z_0 - \zeta_0}{z - z_0} \frac{\mathbf{f} \mathbf{g}^\dagger}{\mathbf{g}^\dagger \mathbf{f}} = \mathbf{I} + \frac{z_0 - \zeta_0}{z - z_0} \mathbf{P}.$$

Then we have the following basic properties:

$$\det \mathbf{R}(z) = \frac{z - \zeta_0}{z - z_0}, \quad \mathbf{R}^{-1}(z) = \mathbf{I} + \frac{\zeta_0 - z_0}{z - \zeta_0} \frac{\mathbf{f} \mathbf{g}^\dagger}{\mathbf{g}^\dagger \mathbf{f}}, \quad \mathbf{R}(z) \mathbf{f} = \left(\frac{z - \zeta_0}{z - z_0} \right) \mathbf{f}, \quad \mathbf{g}^\dagger \mathbf{R}(z) = \left(\frac{z - \zeta_0}{z - z_0} \right) \mathbf{g}^\dagger,$$

$$\mathbf{P}\mathbf{R}(z) = \mathbf{R}(z)\mathbf{P} = \frac{z - \zeta_0}{z - z_0}\mathbf{P}, \quad \mathbf{P}\mathbf{R}^{-1}(z) = \mathbf{R}^{-1}(z)\mathbf{P} = \frac{z - z_0}{z - \zeta_0}\mathbf{P}, \quad \mathbf{R}(z)\mathbf{Q} = \mathbf{Q}\mathbf{R}(z) = \mathbf{Q};$$

Proof. This is a simple direct computation, see also [Dzh09]. \square

3. SCHLESINGER TRANSFORMATIONS

In this section we derive evolution equations for an elementary Schlesinger transformation $\left\{ \begin{smallmatrix} \alpha & \beta \\ \mu & \nu \end{smallmatrix} \right\}$ in terms of the coefficient matrices \mathbf{A}_i of the Fuchsian system and then lift them to get the discrete Schlesinger dynamic on the decomposition space $\mathcal{B} \times \mathcal{C}$. Take $\mathbf{R}(z)$ to be of the form considered in Lemma 2.5. Then we have the following result.

Theorem 3.1. *Consider transformation (2.3) with*

$$(3.1) \quad \mathbf{R}(z) = \mathbf{I} + \frac{z_0 - \zeta_0}{z - z_0} \frac{\mathbf{f}\mathbf{g}^\dagger}{\mathbf{g}^\dagger\mathbf{f}} = \mathbf{I} + \frac{z_0 - \zeta_0}{z - z_0} \mathbf{P}, \quad z_0 \neq \zeta_0.$$

Then

(i) *the poles $\{\bar{z}_i\}$ of $\bar{\mathbf{A}}(z)$ coincide with the poles $\{z_i\}$ of $\mathbf{A}(z)$ if and only if, for some choice of indices α and β , $\beta \neq \alpha$, we have $z_0 = z_\alpha$, $\zeta_0 = z_\beta$, and*

$$(3.2) \quad \bar{\mathbf{A}}_\alpha \mathbf{P} = \mathbf{P}\mathbf{A}_\alpha - \mathbf{P}, \quad \mathbf{P}\bar{\mathbf{A}}_\beta = \mathbf{A}_\beta \mathbf{P} + \mathbf{P}.$$

Then, for some indices μ and ν ,

$$(3.3) \quad \mathbf{f} \sim \bar{\mathbf{b}}_{\alpha, \mu} \sim \mathbf{b}_{\beta, \nu}, \quad \mathbf{g}^\dagger \sim \mathbf{c}_\alpha^{\mu\dagger} \sim \bar{\mathbf{c}}_\beta^{\nu\dagger}, \quad \bar{\theta}_\alpha^\mu = \theta_\alpha^\mu - 1, \quad \bar{\theta}_\beta^\nu = \theta_\beta^\nu + 1,$$

where \sim means proportional, and so such $\mathbf{R}(z)$ defines the elementary transformation $\left\{ \begin{smallmatrix} \alpha & \beta \\ \mu & \nu \end{smallmatrix} \right\}$.

(ii) *For $\mathbf{R}(z)$ satisfying (3.3),*

$$(3.4) \quad \mathbf{R}(z) = \mathbf{I} + \frac{z_\alpha - z_\beta}{z - z_\alpha} \frac{\mathbf{b}_{\beta, \nu} \mathbf{c}_\alpha^{\mu\dagger}}{\mathbf{c}_\alpha^{\mu\dagger} \mathbf{b}_{\beta, \nu}},$$

the residue matrices of $\mathbf{A}(z)$ and $\bar{\mathbf{A}}(z)$ are connected by the equations

$$(3.5) \quad \bar{\mathbf{A}}_i \mathbf{R}(z_i) = \mathbf{R}(z_i) \mathbf{A}_i \quad \text{or equivalently} \quad \mathbf{A}_i \mathbf{R}^{-1}(z_i) = \mathbf{R}^{-1}(z_i) \bar{\mathbf{A}}_i, \quad i \neq \alpha, \beta,$$

$$(3.6) \quad \bar{\mathbf{A}}_\beta \mathbf{Q} = \mathbf{Q}\mathbf{A}_\beta \quad \text{and} \quad \mathbf{A}_\alpha \mathbf{Q} = \mathbf{Q}\bar{\mathbf{A}}_\alpha, \quad \text{where} \quad \mathbf{Q} = \mathbf{I} - \mathbf{P} = \mathbf{R}(z_\beta) = \mathbf{R}^{-1}(z_\alpha).$$

(iii) *Elementary Schlesinger transformation $\left\{ \begin{smallmatrix} \alpha & \beta \\ \mu & \nu \end{smallmatrix} \right\}$ corresponds to the following map on the coefficient matrices (discrete Schlesinger evolution equations):*

$$(3.7) \quad \bar{\mathbf{A}}_i = \mathbf{R}(z_i) \mathbf{A}_i \mathbf{R}^{-1}(z_i), \quad i \neq \alpha, \beta,$$

$$(3.8) \quad \bar{\mathbf{A}}_\alpha = \mathbf{A}_\alpha - [\mathbf{A}_\alpha, \mathbf{P}] - \mathbf{P} + \sum_{i \neq \alpha} \left(\frac{z_\beta - z_\alpha}{z_i - z_\alpha} \right) \mathbf{P}\mathbf{A}_i \mathbf{Q},$$

$$(3.9) \quad \bar{\mathbf{A}}_\beta = \mathbf{A}_\beta + [\mathbf{A}_\beta, \mathbf{P}] + \mathbf{P} + \sum_{i \neq \beta} \left(\frac{z_\alpha - z_\beta}{z_i - z_\beta} \right) \mathbf{Q}\mathbf{A}_i \mathbf{P}.$$

Proof. From the Schlesinger transformation equations

$$(3.10) \quad \bar{\mathbf{A}}(z)\mathbf{R}(z) = \mathbf{R}(z)\mathbf{A}(z) + \frac{d\mathbf{R}(z)}{dz} \quad \text{or} \quad \mathbf{R}^{-1}(z)\bar{\mathbf{A}}(z) = \mathbf{A}(z)\mathbf{R}^{-1}(z) - \frac{d\mathbf{R}^{-1}(z)}{dz},$$

and

$$\frac{d\mathbf{R}(z)}{dz} \mathbf{R}^{-1}(z) = -\mathbf{R}(z) \frac{d\mathbf{R}^{-1}(z)}{dz} = \left(\frac{1}{z - \zeta_0} - \frac{1}{z - z_0} \right) \frac{\mathbf{f}\mathbf{g}^\dagger}{\mathbf{g}^\dagger\mathbf{f}},$$

we see that

$$\text{tr} \frac{d\mathbf{R}(z)}{dz} \mathbf{R}^{-1}(z) = \left(\frac{1}{z - \zeta_0} - \frac{1}{z - z_0} \right) = \text{tr} \bar{\mathbf{A}}(z) - \text{tr} \mathbf{A}(z) = \sum_{i=1}^n \frac{\text{tr}(\bar{\mathbf{A}}_i) - \text{tr}(\mathbf{A}_i)}{z - z_i}.$$

Therefore, $z_0, \zeta_0 \in \{z_i\}_{i=1}^n$. Let $z_0 = z_\alpha$ and $\zeta_0 = z_\beta$. Then $\text{tr}(\bar{\mathbf{A}}_\alpha) - \text{tr}(\mathbf{A}_\alpha) = -1$ and $\text{tr}(\bar{\mathbf{A}}_\beta) - \text{tr}(\mathbf{A}_\beta) = 1$.

At z_α the first equation in (3.10) becomes

$$\left(\frac{\bar{\mathbf{A}}_\alpha}{z - z_\alpha} + \sum_{i \neq \alpha} \frac{\bar{\mathbf{A}}_i}{z - z_i} \right) \left(\mathbf{I} + \frac{z_\alpha - z_\beta}{z - z_\alpha} \mathbf{P} \right) = \left(\mathbf{I} + \frac{z_\alpha - z_\beta}{z - z_\alpha} \mathbf{P} \right) \left(\frac{\mathbf{A}_\alpha}{z - z_\alpha} + \sum_{i \neq \alpha} \frac{\mathbf{A}_i}{z - z_i} \right) - \frac{z_\alpha - z_\beta}{(z - z_\alpha)^2} \mathbf{P}.$$

The $(z - z_\alpha)^{-2}$ -terms give

$$\bar{\mathbf{A}}_\alpha \mathbf{P} = \mathbf{P} \mathbf{A}_\alpha - \mathbf{P},$$

and so we see that \mathbf{g}^\dagger has to be an eigenvector of \mathbf{A}_α and \mathbf{f} has to be an eigenvector of $\bar{\mathbf{A}}_\alpha$. Choose an index μ and let $\mathbf{g}^\dagger \sim \mathbf{c}_\alpha^{\mu\dagger}$ and $\mathbf{f} \sim \bar{\mathbf{b}}_{\alpha,\mu}$, then $\theta_\alpha^\mu = \bar{\theta}_\alpha^\mu + 1$. Repeating this argument for z_β and the second equation in (3.10) implies that $\mathbf{g}^\dagger \sim \bar{\mathbf{c}}_\beta^{\nu\dagger}$, $\mathbf{f} \sim \mathbf{b}_{\beta,\nu}$, and $\theta_\beta^\nu = \bar{\theta}_\beta^\nu - 1$, for some choice of the index ν , which completes the proof of (i).

Since $\mathbf{R}(z)$ is regular (as a complex function) and invertible (as a matrix) at z_i for $i \neq \alpha, \beta$, taking residues of (3.10) at z_i gives (3.5), which is also equivalent to (3.7). Taking the residue of $\mathbf{R}(z)$ at z_β of the first equation in (3.10) gives the first equation in (3.6) and taking the residue at z_α of the second equation in (3.10) gives the second equation in (3.6). This completes the proof of (ii).

Finally, taking the residue at infinity we get $-\bar{\mathbf{A}}_\infty = \sum_{i=1}^n \bar{\mathbf{A}}_i = \sum_{i=1}^n \mathbf{A}_i = -\mathbf{A}_\infty$, which becomes

$$(3.11) \quad \bar{\mathbf{A}}_\alpha + \bar{\mathbf{A}}_\beta = \mathbf{A}_\alpha + \mathbf{A}_\beta + \sum_{i \neq \alpha, \beta} (\mathbf{A}_i - \bar{\mathbf{A}}_i) = \mathbf{A}_\alpha + \mathbf{A}_\beta + \sum_{i \neq \alpha, \beta} (\mathbf{A}_i - \mathbf{R}(z_i) \mathbf{A}_i \mathbf{R}^{-1}(z_i)),$$

after using (3.5). It is easy to see that taking the residue at z_α of the first equation in (3.10) or the residue at z_β of the second equation in (3.10) results in equivalent equations.

Right-multiplying (3.11) by \mathbf{Q} and using (3.6)

$$\bar{\mathbf{A}}_\alpha \mathbf{Q} = \mathbf{A}_\alpha \mathbf{Q} + (\mathbf{A}_\beta \mathbf{Q} - \mathbf{Q} \mathbf{A}_\beta) + \sum_{i \neq \alpha, \beta} \left(-\frac{z_\alpha - z_\beta}{z_i - z_\alpha} \right) \mathbf{P} \mathbf{A}_i \mathbf{Q} = \mathbf{A}_\alpha - \mathbf{A}_\alpha \mathbf{P} + \sum_{i \neq \alpha} \left(\frac{z_\beta - z_\alpha}{z_i - z_\alpha} \right) \mathbf{P} \mathbf{A}_i \mathbf{Q},$$

since $\mathbf{Q} \mathbf{A}_\beta = \mathbf{Q} \mathbf{A}_\beta \mathbf{Q}$ and

$$(\mathbf{A}_i - \mathbf{R}(z_i) \mathbf{A}_i \mathbf{R}^{-1}(z_i)) \mathbf{Q} = (\mathbf{I} - \mathbf{R}(z_i)) \mathbf{A}_i \mathbf{Q} = -\frac{z_\alpha - z_\beta}{z_i - z_\alpha} \mathbf{P} \mathbf{A}_i \mathbf{Q}, \quad \text{since } \mathbf{R}^{-1}(z_i) \mathbf{Q} = \mathbf{Q}.$$

Finally, using $\bar{\mathbf{A}}_\alpha \mathbf{Q} = \bar{\mathbf{A}}_\alpha - \bar{\mathbf{A}}_\alpha \mathbf{P} = \bar{\mathbf{A}}_\alpha - \mathbf{P} \mathbf{A}_\alpha + \mathbf{P}$ and simplifying, we get (3.8). Equation (3.9) is obtained in the same way, only we start by left-multiplying (3.11) by \mathbf{Q} . \square

We now want to lift these equations to the decomposition space $\mathcal{B} \times \mathcal{C}$. Our strategy is the following. Imposing the *orthogonality conditions*

$$(3.12) \quad \mathbf{C}_i^\dagger \mathbf{B}_i = \Theta_i, \quad \text{and} \quad \bar{\mathbf{C}}_i^\dagger \bar{\mathbf{B}}_i = \Theta_i$$

allows us to “split” equations (3.7-3.9) to get the map from $(\mathcal{B} \times \mathcal{C})_{(\Theta, \mathbf{A}_\infty)}$ to $(\bar{\mathcal{B}} \times \bar{\mathcal{C}})_{(\bar{\Theta}, \bar{\mathbf{A}}_\infty)}$ that we then extend to the (dense open subset of the) whole of the decomposition space.

For convenience we list here some consequences of the orthogonality conditions that we often use:

$$\begin{aligned} \mathbf{P} \mathbf{b}_{\beta,\nu} &= \mathbf{b}_{\beta,\nu}, & \mathbf{Q} \mathbf{b}_{\beta,\nu} &= \mathbf{0}, & \mathbf{P} \mathbf{b}_{\alpha,j} &= \mathbf{0}, & \mathbf{Q} \mathbf{b}_{\alpha,j} &= \mathbf{b}_{\alpha,j} \quad (j \neq \mu), \\ \mathbf{c}_\alpha^{\mu\dagger} \mathbf{P} &= \mathbf{c}_\alpha^{\mu\dagger}, & \mathbf{c}_\alpha^{\mu\dagger} \mathbf{Q} &= \mathbf{0}, & \mathbf{c}_\beta^{j\dagger} \mathbf{P} &= \mathbf{0}, & \mathbf{c}_\beta^{j\dagger} \mathbf{Q} &= \mathbf{c}_\beta^{j\dagger} \quad (j \neq \nu). \end{aligned}$$

We begin with the following easy Lemma.

Lemma 3.2. *Let $\mathbf{A}(z)$ and $\bar{\mathbf{A}}(z)$ be connected by an elementary Schlesinger transformation $\left\{ \begin{smallmatrix} \alpha & \beta \\ \mu & \nu \end{smallmatrix} \right\}$ given by $\mathbf{R}(z)$ of the form (3.1) satisfying (3.3) and let $\mathbf{A}_i = \mathbf{B}_i \mathbf{C}_i^\dagger$ with $\mathbf{C}_i^\dagger \mathbf{B}_i = \Theta_i$, and similarly for $\bar{\mathbf{A}}_i$. Then the corresponding points $(\mathbf{B}, \mathbf{C}^\dagger)$ and $(\bar{\mathbf{B}}, \bar{\mathbf{C}}^\dagger)$ in the decomposition space are related as follows:*

(i) *If $i \neq \alpha, \beta$, $\mathbf{R}(z_i)$ is invertible and*

$$(3.13) \quad \bar{\mathbf{B}}_i \sim \mathbf{R}(z_i) \mathbf{B}_i \quad (\text{equivalently, } \mathbf{B}_i \sim \mathbf{R}^{-1}(z_i) \bar{\mathbf{B}}_i) \quad \text{and} \quad \mathbf{C}_i^\dagger \sim \bar{\mathbf{C}}_i^\dagger \mathbf{R}(z_i) \quad (\bar{\mathbf{C}}_i^\dagger \sim \mathbf{C}_i^\dagger \mathbf{R}^{-1}(z_i)).$$

Therefore

$$(3.14) \quad \bar{\mathbf{B}}_i \bar{\mathbf{D}}_i = \mathbf{R}(z_i) \mathbf{B}_i, \quad \bar{\Delta}_i \bar{\mathbf{C}}_i^\dagger = \mathbf{C}_i^\dagger \mathbf{R}^{-1}(z_i),$$

where $\bar{\mathbf{D}}_i$ and $\bar{\Delta}_i$ are diagonal normalization matrices, $\bar{\mathbf{D}}_i \bar{\Delta}_i = \mathbf{I}$.

(ii) For the special indices α and β we have

$$(3.15) \quad \mathbf{C}_\beta^\dagger \sim \bar{\mathbf{C}}_\beta^\dagger \mathbf{Q}, \quad \bar{\mathbf{B}}_\beta \sim \mathbf{Q} \mathbf{B}_\beta, \quad \text{and} \quad \bar{\mathbf{C}}_\alpha^\dagger \sim \mathbf{C}_\alpha^\dagger \mathbf{Q}, \quad \mathbf{B}_\alpha \sim \mathbf{Q} \bar{\mathbf{B}}_\alpha.$$

It is still possible to write

$$(3.16) \quad \bar{\mathbf{B}}_\beta \bar{\mathbf{D}}_\beta = \mathbf{Q} \mathbf{B}_\beta, \quad \bar{\Delta}_\alpha \bar{\mathbf{C}}_\alpha^\dagger = \mathbf{C}_\alpha^\dagger \mathbf{Q},$$

but since $\mathbf{Q} \mathbf{b}_{\beta,\nu} = \mathbf{c}_\alpha^{\mu\dagger} \mathbf{Q} = \mathbf{0}$, $(\bar{\mathbf{D}}_\beta)_\nu^\nu = (\bar{\Delta}_\alpha)_\mu^\mu = 0$.

Proof. Multiplying first equation in (3.5) on the right by \mathbf{B}_i gives $\bar{\mathbf{A}}_i \mathbf{R}(z_i) \mathbf{B}_i = \mathbf{R}(z_i) \mathbf{A}_i \mathbf{B}_i = \mathbf{R}(z_i) \mathbf{B}_i \Theta_i$, and so the matrix $\mathbf{R}(z_i) \mathbf{B}_i$ is the matrix of eigenvectors of $\bar{\mathbf{A}}_i$, $\bar{\mathbf{B}}_i \sim \mathbf{R}(z_i) \mathbf{B}_i$ or $\bar{\mathbf{B}}_i \bar{\mathbf{D}}_i = \mathbf{R}(z_i) \mathbf{B}_i$, where $\bar{\mathbf{D}}_i$ is a diagonal matrix of non-zero proportionality coefficients. The equations for $\bar{\mathbf{C}}_i^\dagger$ are proved similarly. From the normalization condition

$$\bar{\Delta}_i \bar{\mathbf{C}}_i^\dagger \bar{\mathbf{B}}_i \bar{\mathbf{D}}_i = \bar{\Delta}_i \bar{\Theta}_i \bar{\mathbf{D}}_i = \bar{\Delta}_i \bar{\mathbf{D}}_i \bar{\Theta}_i = \mathbf{C}_i^\dagger \mathbf{B}_i = \Theta_i = \bar{\Theta}_i,$$

where all matrices are diagonal, we immediately get $\bar{\mathbf{D}}_i \bar{\Delta}_i = \mathbf{I}$.

For the special indices α and β , away from the pairs (α, μ) and (β, ν) , the situation is exactly the same. \square

Theorem 3.3. The elementary Schlesinger transformation $\left\{ \begin{smallmatrix} \alpha & \beta \\ \mu & \nu \end{smallmatrix} \right\}$ defines the map

$$(\mathcal{B} \times \mathcal{C})_{(\Theta, \mathbf{A}_\infty)} \rightarrow (\bar{\mathcal{B}} \times \bar{\mathcal{C}})_{(\bar{\Theta}, \bar{\mathbf{A}}_\infty)}$$

given by the following evolution equations (grouped for convenience).

(i) Transformation vectors:

$$(3.17) \quad \bar{\mathbf{b}}_{\alpha,\mu} = \frac{1}{c_\alpha^\mu} \mathbf{b}_{\beta,\nu}, \quad \bar{\mathbf{c}}_\beta^{\nu\dagger} = c_\beta^\nu \mathbf{c}_\alpha^{\mu\dagger}.$$

(ii) Generic indices:

$$(3.18) \quad \bar{\mathbf{b}}_{i,j} = \frac{1}{c_i^j} \mathbf{R}(z_i) \mathbf{b}_{i,j}, \quad (i \neq \alpha \text{ and if } i = \beta, j \neq \nu); \quad \bar{\mathbf{c}}_i^{j\dagger} = c_i^j \mathbf{c}_i^{j\dagger} \mathbf{R}^{-1}(z_i), \quad (i \neq \beta \text{ and if } i = \alpha, j \neq \mu).$$

(iii) Special indices:

$$(3.19) \quad \bar{\mathbf{b}}_{\alpha,j} = \frac{1}{c_\alpha^j} \left(\mathbf{I} - \frac{\mathbf{P} \left(\sum_{i \neq \alpha} \frac{z_\beta - z_\alpha}{z_i - z_\alpha} \mathbf{A}_i \right)}{\theta_\alpha^\mu - \theta_\alpha^j - 1} \right) \mathbf{b}_{\alpha,j}, \quad j \neq \mu;$$

$$(3.20) \quad \bar{\mathbf{c}}_\beta^{j\dagger} = c_\beta^j \mathbf{c}_\beta^{j\dagger} \left(\mathbf{I} - \frac{\left(\sum_{i \neq \beta} \frac{z_\alpha - z_\beta}{z_i - z_\beta} \mathbf{A}_i \right) \mathbf{P}}{\theta_\beta^\nu - \theta_\beta^j + 1} \right), \quad j \neq \nu;$$

$$(3.21) \quad \bar{\mathbf{b}}_{\beta,\nu} = \frac{1}{c_\beta^\nu} \left((\theta_\beta^\nu + 1) \mathbf{I} + \mathbf{Q} \left(\mathbf{I} + \sum_{j \neq \nu} \frac{\mathbf{b}_{\beta,j} \mathbf{c}_\beta^{j\dagger}}{\theta_\beta^\nu - \theta_\beta^j + 1} \right) \left(\sum_{i \neq \beta} \frac{z_\alpha - z_\beta}{z_i - z_\beta} \mathbf{A}_i \right) \right) \mathbf{c}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu};$$

$$(3.22) \quad \bar{\mathbf{c}}_\alpha^{\mu\dagger} = c_\alpha^\mu \frac{\mathbf{c}_\alpha^{\mu\dagger}}{\mathbf{c}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu}} \left((\theta_\alpha^\mu - 1) \mathbf{I} + \left(\sum_{i \neq \alpha} \frac{z_\beta - z_\alpha}{z_i - z_\alpha} \mathbf{A}_i \right) \left(\mathbf{I} + \sum_{j \neq \mu} \frac{\mathbf{b}_{\alpha,j} \mathbf{c}_\alpha^{j\dagger}}{\theta_\alpha^\mu - \theta_\alpha^j - 1} \right) \mathbf{Q} \right).$$

Proof. Part (i) is just a restatement of (3.3) and part (ii) follows immediately from Lemma 3.2 if we put $\bar{\mathbf{D}}_i = \bar{\Delta}_i^{-1} = \text{diag}\{c_i^1, \dots, c_i^{r_i}\}$. Note, however, that for special indices α and β , since $(\bar{\mathbf{D}}_\beta)_\nu^\nu = (\bar{\Delta}_\alpha)_\mu^\mu = 0$, we have no information about $\bar{\mathbf{b}}_{\beta,\nu}$ and $\bar{\mathbf{c}}_\alpha^{\mu\dagger}$.

The difficult part of the Theorem is to establish equations in (iii). Right-multiplying (3.8) by $\mathbf{Q} \mathbf{b}_{\alpha,j} = \mathbf{b}_{\alpha,j}$, $j \neq \mu$, we get

$$\bar{\mathbf{A}}_\alpha \mathbf{Q} \mathbf{b}_{\alpha,j} = \theta_\alpha^j \mathbf{b}_{\alpha,j} + \sum_{i \neq \alpha} \left(\frac{z_\beta - z_\alpha}{z_i - z_\alpha} \right) \mathbf{P} \mathbf{A}_i \mathbf{b}_{\alpha,j}.$$

At the same time, in view of (3.18) and orthogonality

$$\bar{\mathbf{A}}_\alpha = \sum_{j \neq \mu} c_\alpha^j \bar{\mathbf{b}}_{\alpha,j} \mathbf{c}_\alpha^{j\dagger} \mathbf{Q} + \bar{\mathbf{b}}_{\alpha,\mu} \bar{\mathbf{c}}_\alpha^{\mu\dagger}, \quad \bar{\mathbf{A}}_\alpha \mathbf{Q} \mathbf{b}_{\alpha,j} = c_\alpha^j \bar{\mathbf{b}}_{\alpha,j} \theta_\alpha^j + \bar{\mathbf{b}}_{\alpha,\mu} \bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\alpha,j}.$$

Thus,

$$c_\alpha^j \bar{\mathbf{b}}_{\alpha,j} \theta_\alpha^j + \bar{\mathbf{b}}_{\alpha,\mu} \bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\alpha,j} = \theta_\alpha^j \mathbf{b}_{\alpha,j} + \sum_{i \neq \alpha} \frac{z_\beta - z_\alpha}{z_i - z_\alpha} \mathbf{P} \mathbf{A}_i \mathbf{b}_{\alpha,j}.$$

To evaluate $\bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\alpha,j}$ we now left-multiply by $\bar{\mathbf{c}}_\alpha^{\mu\dagger}$ and use the fact that $\bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{P} = (\bar{\theta}_\alpha^\mu \mathbf{c}_\alpha^{\mu\dagger}) / (\mathbf{c}_\alpha^{\mu\dagger} \bar{\mathbf{b}}_{\alpha,\mu})$:

$$\bar{\theta}_\alpha^\mu \bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\alpha,j} = \theta_\alpha^j \bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\alpha,j} + \sum_{i \neq \alpha} \frac{z_\beta - z_\alpha}{z_i - z_\alpha} \cdot \frac{\bar{\theta}_\alpha^\mu \mathbf{c}_\alpha^{\mu\dagger} \mathbf{A}_i \mathbf{b}_{\alpha,j}}{\mathbf{c}_\alpha^{\mu\dagger} \bar{\mathbf{b}}_{\alpha,\mu}}.$$

Hence

$$\bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\alpha,j} = \frac{\bar{\theta}_\alpha^\mu \mathbf{c}_\alpha^{\mu\dagger}}{\mathbf{c}_\alpha^{\mu\dagger} \bar{\mathbf{b}}_{\alpha,\mu}} \left(\sum_{i \neq \alpha} \frac{z_\beta - z_\alpha}{z_i - z_\alpha} \mathbf{A}_i \right) \frac{\mathbf{b}_{\alpha,j}}{\bar{\theta}_\alpha^\mu - \theta_\alpha^j}.$$

Then

$$\bar{\mathbf{b}}_{\alpha,j} c_\alpha^j \theta_\alpha^j = \theta_\alpha^j \mathbf{b}_{\alpha,j} - \bar{\theta}_{\alpha,\mu} \mathbf{P} \left(\sum_{i \neq \alpha} \frac{z_\beta - z_\alpha}{z_i - z_\alpha} \mathbf{A}_i \right) \frac{\mathbf{b}_{\alpha,j}}{\bar{\theta}_\alpha^\mu - \theta_\alpha^j} + \mathbf{P} \left(\sum_{i \neq \alpha} \frac{z_\beta - z_\alpha}{z_i - z_\alpha} \mathbf{A}_i \right) \mathbf{b}_{\alpha,j},$$

which, after easy simplification, becomes (3.19). Equation (3.20) is proved in the exactly same way.

Next, observe that in view of (3.17), (3.18), and (3.20) we can write

$$(3.23) \quad \bar{\mathbf{A}}_\beta = \sum_{j \neq \nu} \bar{\mathbf{b}}_{\beta,j} \bar{\mathbf{c}}_\beta^{j\dagger} + \bar{\mathbf{b}}_{\beta,\nu} \bar{\mathbf{c}}_\beta^{\nu\dagger} = \sum_{j \neq \nu} \mathbf{Q} \mathbf{b}_{\beta,j} \mathbf{c}_\beta^{j\dagger} \left(\mathbf{I} - \frac{\left(\sum_{i \neq \beta} \frac{z_\alpha - z_\beta}{z_i - z_\beta} \mathbf{A}_i \right) \mathbf{P}}{\theta_\beta^\nu - \theta_\beta^j + 1} \right) + c_\beta^\nu \bar{\mathbf{b}}_{\beta,\nu} \mathbf{c}_\alpha^{\mu\dagger}.$$

Thus, we can compute $\bar{\mathbf{A}}_\beta \mathbf{b}_{\beta,\nu}$ either using (3.23) or (3.9):

$$\begin{aligned} \bar{\mathbf{A}}_\beta \mathbf{b}_{\beta,\nu} &= c_\beta^\nu (\mathbf{c}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu}) \bar{\mathbf{b}}_{\beta,\nu} - \sum_{j \neq \nu} \frac{\mathbf{Q} \mathbf{b}_{\beta,j} \mathbf{c}_\beta^{j\dagger}}{\theta_\beta^\nu - \theta_\beta^j + 1} \left(\left(\sum_{i \neq \beta} \frac{z_\alpha - z_\beta}{z_i - z_\beta} \mathbf{A}_i \right) \mathbf{b}_{\beta,\nu} \right) \\ &= (\theta_\beta^\nu + 1) \mathbf{b}_{\beta,\nu} + \mathbf{Q} \sum_{i \neq \beta} \frac{z_\alpha - z_\beta}{z_i - z_\beta} \mathbf{A}_i \mathbf{b}_{\beta,\nu}. \end{aligned}$$

Solving for $\bar{\mathbf{b}}_{\beta,\nu}$ and simplifying gives (3.21). Equation (3.22) is proved similarly. \square

4. HAMILTONIAN FORMULATION

The main objective of this part is to give the Hamiltonian formulation of the discrete dynamic described in the previous section. First, pick points z_1, \dots, z_n , $z_i \neq z_j$, and characteristic indices θ_i^j , $1 \leq i \leq n$, $1 \leq j \leq r_i$, $\theta_i^j \neq \theta_i^k$ for $i \neq j$ (this is an important assumption that corresponds to Assumption 2.1 but is slightly more restrictive) and put $\bar{\theta}_\alpha^\mu = \theta_\alpha^\mu$, $\bar{\theta}_\beta^\nu = \theta_\beta^\nu + 1$ and $\bar{\theta}_i^j = \theta_i^j$ otherwise. These are parameters of our dynamic.

As before, put $\Theta_i = \text{diag}\{\theta_i^1, \dots, \theta_i^{r_i}\}$ and $\bar{\Theta}_i = \text{diag}\{\bar{\theta}_i^1, \dots, \bar{\theta}_i^{r_i}\}$ and let

$$(4.1) \quad \mathbf{P} = \frac{\mathbf{b}_{\beta,\nu} \bar{\mathbf{c}}_\beta^{\nu\dagger}}{\bar{\mathbf{c}}_\beta^{\nu\dagger} \mathbf{b}_{\beta,\nu}}, \quad \mathbf{R}(z) = \mathbf{I} + \frac{z_\alpha - z_\beta}{z - z_\alpha} \mathbf{P}, \quad \mathbf{Q} = \mathbf{I} - \mathbf{P}.$$

Our goal then is to find a function $\mathcal{H}^+ = \mathcal{H}^+(\mathbf{B}, \bar{\mathbf{C}}^\dagger)$ on $\mathcal{B} \times \bar{\mathcal{C}}$ so that, if we use equations (2.5) to define $\bar{\mathbf{b}}_{i,j}$ and $\mathbf{c}_i^{j\dagger}$ (recall Remark 2.3),

$$\bar{\mathbf{b}}_{i,j} = \frac{\partial \mathcal{H}^+}{\partial \bar{\mathbf{c}}_i^{j\dagger}}(\mathbf{B}, \bar{\mathbf{C}}^\dagger) \quad \text{and} \quad \mathbf{c}_i^{j\dagger} = \frac{\partial \mathcal{H}^+}{\partial \mathbf{b}_{i,j}}(\mathbf{B}, \bar{\mathbf{C}}^\dagger),$$

and construct corresponding matrices $\mathbf{B}_i, \bar{\mathbf{B}}_i, \mathbf{C}_i^\dagger, \bar{\mathbf{C}}_i^\dagger, \mathbf{A}_i = \mathbf{B}_i \mathbf{C}_i^\dagger$, and $\bar{\mathbf{A}}_i = \bar{\mathbf{B}}_i \bar{\mathbf{C}}_i^\dagger$, then the following is true, cf. equations (3.3–3.6) and (3.11):

$$(4.2) \quad \bar{\mathbf{b}}_{\alpha,\mu} \sim \mathbf{b}_{\beta,\nu} \quad \text{and} \quad \mathbf{c}_\alpha^{\mu\dagger} \sim \bar{\mathbf{c}}_\beta^{\nu\dagger};$$

$$(4.3) \quad \mathbf{C}_i^\dagger \mathbf{B}_i = \Theta_i \quad \text{and} \quad \bar{\mathbf{C}}_i^\dagger \bar{\mathbf{B}}_i = \bar{\Theta}_i;$$

$$(4.4) \quad \bar{\mathbf{A}}_i \mathbf{R}(z_i) = \mathbf{R}(z_i) \mathbf{A}_i \text{ for } i \neq \alpha, \beta, \quad \bar{\mathbf{A}}_\beta \mathbf{Q} = \mathbf{Q} \mathbf{A}_\beta, \quad \text{and} \quad \mathbf{Q} \bar{\mathbf{A}}_\alpha = \mathbf{A}_\alpha \mathbf{Q};$$

$$(4.5) \quad \sum_i \bar{\mathbf{A}}_i = \sum_i \mathbf{A}_i.$$

It is convenient to separate special indices (α, μ) and (β, ν) from generic indices (i, j) . Thus, we put

$$(4.6) \quad \mathbf{A}'_\alpha = \sum_{j \neq \mu} \mathbf{b}_{\alpha,j} \mathbf{c}_\alpha^{j\dagger} = \mathbf{B}'_\alpha (\mathbf{C}'_\alpha)^\dagger \quad \left(\text{thus, } \mathbf{A}_\alpha = \sum_j \mathbf{b}_{\alpha,j} \mathbf{c}_\alpha^{j\dagger} = \mathbf{A}'_\alpha + \mathbf{b}_{\alpha,\mu} \mathbf{c}_\alpha^{\mu\dagger} \right),$$

and so on. Then, since in view of (4.1) and (4.2)

$$(4.7) \quad \mathbf{Q} \mathbf{b}_{\beta,\nu} = \mathbf{Q} \bar{\mathbf{b}}_{\alpha,\mu} = \mathbf{0}, \quad \mathbf{c}_\alpha^{\mu\dagger} \mathbf{Q} = \bar{\mathbf{c}}_\beta^{\nu\dagger} \mathbf{Q} = \mathbf{0},$$

equations for $i = \alpha, \beta$ in (4.4) become $\bar{\mathbf{A}}'_\beta \mathbf{Q} = \mathbf{Q} \mathbf{A}'_\beta$, and $\mathbf{Q} \bar{\mathbf{A}}'_\alpha = \mathbf{A}'_\alpha \mathbf{Q}$.

Next, observe that the orthogonality conditions (4.3) impose some *a priori* conditions on the domain of \mathcal{H}^+ that cut out a certain locus $(\mathcal{B} \times \bar{\mathcal{C}})^{\text{ort}} \subset (\mathcal{B} \times \bar{\mathcal{C}})$. Indeed, from (4.2) and the condition $\bar{\mathbf{C}}_\alpha^\dagger \bar{\mathbf{B}}_\alpha = \bar{\Theta}_\alpha$ we get $(\bar{\mathbf{C}}_\alpha^\dagger)' \mathbf{b}_{\beta,\nu} = \mathbf{0}$. Similarly, $\bar{\mathbf{c}}_\beta^{\nu\dagger} \mathbf{B}'_\alpha = \mathbf{0}$. For $i \neq \alpha, \beta$ we get

$$\Theta_i (\bar{\mathbf{C}}_i^\dagger \mathbf{R}(z_i) \mathbf{B}_i) = \bar{\mathbf{C}}_i^\dagger \bar{\mathbf{A}}_i \mathbf{R}(z_i) \mathbf{B}_i = \bar{\mathbf{C}}_i^\dagger \mathbf{R}(z_i) \mathbf{A}_i \mathbf{B}_i = (\bar{\mathbf{C}}_i^\dagger \mathbf{R}(z_i) \mathbf{B}_i) \Theta_i,$$

and in view of our assumption $\theta_i^j \neq \theta_i^k$ for $j \neq k$, $\bar{\mathbf{C}}_i^\dagger \mathbf{R}(z_i) \mathbf{B}_i$ must be diagonal. We also need to require that it is invertible (which is an open condition). Similarly, $((\bar{\mathbf{C}}_\beta^\dagger)' \mathbf{Q} \mathbf{B}'_\beta)$ must be diagonal (and we also require it is invertible). We will later see that we need $(\bar{\mathbf{C}}_\alpha^\dagger)' \mathbf{B}'_\alpha$ to be diagonal (and invertible) as well. Thus, we put

$$(4.8) \quad (\mathcal{B} \times \bar{\mathcal{C}})^{\text{ort}} = \{ (\bar{\mathbf{C}}_\alpha^\dagger)' \mathbf{b}_{\beta,\nu} = \mathbf{0}, \bar{\mathbf{c}}_\beta^{\nu\dagger} \mathbf{B}'_\alpha = \mathbf{0}, (\bar{\mathbf{C}}_\alpha^\dagger)' \mathbf{B}'_\alpha, \bar{\mathbf{C}}_i^\dagger \mathbf{R}(z_i) \mathbf{B}_i \text{ are diagonal and invertible, } i \neq \alpha \}.$$

We can illustrate the relationship between evolution dynamic and discrete Hamiltonian by the diagram on Figure 3:




FIGURE 3. Discrete Hamiltonian Dynamic

Theorem 4.1. *If we restrict to $(\mathcal{B} \times \bar{\mathcal{C}})^{\text{ort}}$, equations (4.2–4.5) can then be solved for $\bar{\mathbf{b}}_{i,j} = \bar{\mathbf{b}}_{i,j}(\mathbf{B}, \bar{\mathbf{C}}^\dagger)$ and $\mathbf{c}_i^{j\dagger} = \mathbf{c}_i^{j\dagger}(\mathbf{B}, \bar{\mathbf{C}}^\dagger)$ to give the following expressions.*

(a) *Generic indices $i \neq \alpha$, $(i, j) \neq (\beta, \nu)$ (recall that $\mathbf{R}(z_\beta) = \mathbf{Q}$):*

$$(4.9) \quad \bar{\mathbf{b}}_{i,j} = \bar{\theta}_i^j \frac{\mathbf{R}(z_i) \mathbf{b}_{i,j}}{\bar{\mathbf{c}}_i^{j\dagger} \mathbf{R}(z_i) \mathbf{b}_{i,j}}, \quad \mathbf{c}_i^{j\dagger} = \theta_i^j \frac{\bar{\mathbf{c}}_i^{j\dagger} \mathbf{R}(z_i)}{\bar{\mathbf{c}}_i^{j\dagger} \mathbf{R}(z_i) \mathbf{b}_{i,j}}.$$

(b) Special index α :

$$(4.10) \quad \bar{\mathbf{b}}_{\alpha,\mu} = \bar{\theta}_\alpha^\mu \frac{\mathbf{b}_{\beta,\nu}}{\bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu}}, \quad \mathbf{c}_\alpha^{\mu\dagger} = \theta_\alpha^\mu \frac{\bar{\mathbf{c}}_\beta^{\nu\dagger}}{\bar{\mathbf{c}}_\beta^{\nu\dagger} \mathbf{b}_{\alpha,\mu}},$$

and for $j \neq \mu$,

$$(4.11) \quad \bar{\mathbf{b}}_{\alpha,j} = \bar{\theta}_\alpha^j \frac{\mathbf{R}_\alpha \mathbf{b}_{\alpha,j}}{\bar{\mathbf{c}}_\alpha^{j\dagger} \mathbf{R}_\alpha \mathbf{b}_{\alpha,j}}, \quad \mathbf{c}_\alpha^{j\dagger} = \theta_\alpha^j \frac{\bar{\mathbf{c}}_\alpha^{j\dagger} \mathbf{R}_\alpha}{\bar{\mathbf{c}}_\alpha^{j\dagger} \mathbf{R}_\alpha \mathbf{b}_{\alpha,j}}, \quad \text{where } \mathbf{R}_\alpha = \mathbf{I} - \frac{\mathbf{b}_{\alpha,\mu} \bar{\mathbf{c}}_\beta^{\nu\dagger}}{\bar{\mathbf{c}}_\beta^{\nu\dagger} \mathbf{b}_{\alpha,\mu}} - \frac{\mathbf{b}_{\beta,\nu} \bar{\mathbf{c}}_\alpha^{\mu\dagger}}{\bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu}}.$$

(c) Special index (β, ν) :

$$(4.12) \quad \bar{\mathbf{b}}_{\beta,\nu} = (\theta_\beta^\nu - \theta_\alpha^\mu + 1) \frac{\mathbf{b}_{\beta,\nu}}{\bar{\mathbf{c}}_\beta^{\nu\dagger} \mathbf{b}_{\beta,\nu}} + \theta_\alpha^\mu \frac{\mathbf{b}_{\alpha,\mu}}{\bar{\mathbf{c}}_\beta^{\nu\dagger} \mathbf{b}_{\alpha,\mu}} - \sum_{j \neq \mu} \theta_\alpha^j \frac{\bar{\mathbf{c}}_\alpha^{j\dagger} \mathbf{b}_{\alpha,\mu}}{\bar{\mathbf{c}}_\beta^{\nu\dagger} \mathbf{b}_{\alpha,\mu}} \cdot \frac{\mathbf{b}_{\alpha,j}}{\bar{\mathbf{c}}_\alpha^{j\dagger} \mathbf{b}_{\alpha,j}} \\ - \mathbf{Q} \left(\sum_{\substack{i \neq \alpha, j \\ (i,j) \neq (\beta,\nu)}} \left(\frac{z_\beta - z_\alpha}{z_i - z_\alpha} \right) \theta_i^j \frac{\mathbf{b}_{i,j} \bar{\mathbf{c}}_i^{j\dagger}}{\bar{\mathbf{c}}_i^{j\dagger} \mathbf{R}(z_i) \mathbf{b}_{i,j}} \right) \frac{\mathbf{b}_{\beta,\nu}}{\bar{\mathbf{c}}_\beta^{\nu\dagger} \mathbf{b}_{\beta,\nu}};$$

$$(4.13) \quad \mathbf{c}_\beta^{\nu\dagger} = (\theta_\beta^\nu - \theta_\alpha^\mu + 1) \frac{\bar{\mathbf{c}}_\beta^{\nu\dagger}}{\bar{\mathbf{c}}_\beta^{\nu\dagger} \mathbf{b}_{\beta,\nu}} + \bar{\theta}_\alpha^\mu \frac{\bar{\mathbf{c}}_\alpha^{\mu\dagger}}{\bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu}} - \sum_{j \neq \mu} \theta_\alpha^j \frac{\bar{\mathbf{c}}_\alpha^{j\dagger}}{\bar{\mathbf{c}}_\alpha^{j\dagger} \mathbf{b}_{\beta,\nu}} \cdot \frac{\mathbf{b}_{\alpha,j} \bar{\mathbf{c}}_\alpha^{j\dagger}}{\bar{\mathbf{c}}_\alpha^{j\dagger} \mathbf{b}_{\alpha,j}} \\ - \frac{\bar{\mathbf{c}}_\beta^{\nu\dagger}}{\bar{\mathbf{c}}_\beta^{\nu\dagger} \mathbf{b}_{\beta,\nu}} \left(\sum_{\substack{i \neq \alpha, j \\ (i,j) \neq (\beta,\nu)}} \left(\frac{z_\beta - z_\alpha}{z_i - z_\alpha} \right) \theta_i^j \frac{\mathbf{b}_{i,j} \bar{\mathbf{c}}_i^{j\dagger}}{\bar{\mathbf{c}}_i^{j\dagger} \mathbf{R}(z_i) \mathbf{b}_{i,j}} \right) \mathbf{Q}.$$

Proof. To prove (a) for $i \neq \alpha, \beta$, we right-multiply the equation $\bar{\mathbf{A}}_i \mathbf{R}(z_i) = \mathbf{R}(z_i) \mathbf{A}_i$ by \mathbf{B}_i to obtain $\bar{\mathbf{B}}_i (\bar{\mathbf{C}}_i^\dagger \mathbf{R}(z_i) \mathbf{B}_i) = \mathbf{R}(z_i) \mathbf{B}_i \Theta_i$ and then use (4.8) to get $\bar{\mathbf{B}}_i = \mathbf{R}(z_i) \mathbf{B}_i \Theta_i (\bar{\mathbf{C}}_i^\dagger \mathbf{R}(z_i) \mathbf{B}_i)^{-1}$, which is the first equation in (4.9). The second equation is obtained in the same way, only we start by-left multiplying by $\bar{\mathbf{C}}_i^\dagger$. The exact same argument works for indices (β, j) with $j \neq \nu$.

To prove (b), notice that equations (4.10) follow immediately from (4.2–4.3). Equations (4.11) are slightly less straightforward. First, given \mathbf{R}_α as in (4.11), it is easy to see

$$\mathbf{R}_\alpha \mathbf{Q} = \mathbf{I} - \frac{\mathbf{b}_{\beta,\nu} \bar{\mathbf{c}}_\alpha^{\mu\dagger}}{\bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu}} \quad \text{and} \quad \mathbf{Q} \mathbf{R}_\alpha = \mathbf{I} - \frac{\mathbf{b}_{\alpha,\mu} \bar{\mathbf{c}}_\beta^{\nu\dagger}}{\bar{\mathbf{c}}_\beta^{\nu\dagger} \mathbf{b}_{\alpha,\mu}}.$$

Then, since by (4.3) we require $\bar{\mathbf{c}}_\alpha^{\mu\dagger} \bar{\mathbf{b}}_{\alpha,j} = \mathbf{c}_\alpha^{j\dagger} \mathbf{b}_{\alpha,\mu} = 0$ for $j \neq \mu$, we get $\mathbf{R}_\alpha \mathbf{Q} \bar{\mathbf{A}}'_\alpha = \bar{\mathbf{A}}'_\alpha$, $\mathbf{A}'_\alpha \mathbf{Q} \mathbf{R}_\alpha = \mathbf{A}'_\alpha$ and so equation $\mathbf{Q} \bar{\mathbf{A}}'_\alpha = \mathbf{A}'_\alpha \mathbf{Q}$ in (4.4) becomes $\bar{\mathbf{A}}_\alpha \mathbf{R}_\alpha = \mathbf{R}_\alpha \mathbf{A}'_\alpha$. The rest of the argument proceeds exactly as in part (a), with one minor modification that $(\bar{\mathbf{C}}_\alpha^\dagger)' \mathbf{R}_\alpha \mathbf{B}'_\alpha$ becomes just $(\bar{\mathbf{C}}_\alpha^\dagger)' \mathbf{B}'_\alpha$ in view of orthogonality.

Given (a) and (b), equation (4.5) can be written as

$$(4.14) \quad \bar{\mathbf{b}}_{\beta,\nu} \bar{\mathbf{c}}_\beta^{\nu\dagger} - \mathbf{b}_{\beta,\nu} \mathbf{c}_\beta^{\nu\dagger} = \theta_\alpha^\mu \frac{\mathbf{b}_{\alpha,\mu} \bar{\mathbf{c}}_\beta^{\nu\dagger}}{\bar{\mathbf{c}}_\beta^{\nu\dagger} \mathbf{b}_{\alpha,\mu}} - \bar{\theta}_\alpha^\mu \frac{\mathbf{b}_{\beta,\nu} \bar{\mathbf{c}}_\alpha^{\mu\dagger}}{\bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu}} + \sum_{j \neq \mu} \theta_\alpha^j \frac{\mathbf{b}_{\alpha,j} \bar{\mathbf{c}}_\alpha^{j\dagger} \mathbf{R}_\alpha - \mathbf{R}_\alpha \mathbf{b}_{\alpha,j} \bar{\mathbf{c}}_\alpha^{j\dagger}}{\bar{\mathbf{c}}_\alpha^{j\dagger} \mathbf{b}_{\alpha,j}} \\ + \sum_{j \neq \nu} \theta_\beta^j \frac{\mathbf{b}_{\beta,j} \bar{\mathbf{c}}_\beta^{j\dagger} \mathbf{Q} - \mathbf{Q} \mathbf{b}_{\beta,j} \bar{\mathbf{c}}_\beta^{j\dagger}}{\bar{\mathbf{c}}_\beta^{j\dagger} \mathbf{Q} \mathbf{b}_{\beta,j}} + \sum_{i \neq \alpha, \beta} \sum_j \theta_i^j \frac{\mathbf{b}_{i,j} \bar{\mathbf{c}}_i^{j\dagger} \mathbf{R}(z_i) - \mathbf{R}(z_i) \mathbf{b}_{i,j} \bar{\mathbf{c}}_i^{j\dagger}}{\bar{\mathbf{c}}_i^{j\dagger} \mathbf{b}_{i,j}}.$$

To extract $\bar{\mathbf{b}}_{\beta,\nu}$ we multiply (4.14) on the left by \mathbf{Q} (recall that $\mathbf{Q}\mathbf{b}_{\beta,\nu} = \mathbf{0}$) and on the right by $\mathbf{b}_{\alpha,\mu}$ and use (4.3) and (4.8):

$$\begin{aligned} \mathbf{Q}(\bar{\mathbf{b}}_{\beta,\nu}\bar{\mathbf{c}}_{\beta}^{\nu\dagger})\mathbf{b}_{\alpha,\mu} &= \left(\bar{\mathbf{b}}_{\beta,\nu} - (\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\bar{\mathbf{b}}_{\beta,\nu})\frac{\mathbf{b}_{\beta,\nu}}{\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\beta,\nu}}\right)\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\alpha,\mu} = \left(\bar{\mathbf{b}}_{\beta,\nu} - \bar{\theta}_{\beta}^{\nu}\frac{\mathbf{b}_{\beta,\nu}}{\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\beta,\nu}}\right)\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\alpha,\mu} \\ &= \theta_{\alpha}^{\mu}\mathbf{Q}\mathbf{b}_{\alpha,\mu} + \sum_{j\neq\mu}\theta_{\alpha}^j\left(-\frac{\bar{\mathbf{c}}_{\alpha}^{j\dagger}\mathbf{b}_{\alpha,\mu}}{\bar{\mathbf{c}}_{\alpha}^{j\dagger}\mathbf{b}_{\alpha,j}}\right)\mathbf{b}_{\alpha,j} + \sum_{j\neq\nu}\theta_{\beta}^j\left(-\frac{(\bar{\mathbf{c}}_{\beta}^{j\dagger}\mathbf{b}_{\beta,\nu})(\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\alpha,\mu})}{(\bar{\mathbf{c}}_{\beta}^{j\dagger}\mathbf{Q}\mathbf{b}_{\beta,j})(\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\beta,\nu})}\right)\mathbf{Q}\mathbf{b}_{\beta,j} \\ &\quad + \sum_{i\neq\alpha,\beta}\sum_j\left(\frac{z_{\alpha}-z_{\beta}}{z_i-z_{\alpha}}\right)\frac{(\bar{\mathbf{c}}_i^{j\dagger}\mathbf{b}_{\beta,\nu})(\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\alpha,\mu})}{(\bar{\mathbf{c}}_i^{j\dagger}\mathbf{R}(z_i)\mathbf{b}_{i,j})(\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\beta,\nu})}\mathbf{Q}\mathbf{b}_{i,j}. \end{aligned}$$

After straightforward simplification this gives (4.12). Equation (4.13) is obtained in the same way, except that we start by multiplying (4.14) on the right by \mathbf{Q} and on the left by $\bar{\mathbf{c}}_{\alpha}^{\mu\dagger}$. \square

We now show that these equations are Hamiltonian.

Theorem 4.2. *Let*

$$(4.15) \quad \begin{aligned} \mathcal{H}^+(\mathbf{B}, \bar{\mathbf{C}}^{\dagger}; \Theta) &= (\theta_{\beta}^{\nu} - \theta_{\alpha}^{\mu} + 1)\log(\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\beta,\nu}) + \theta_{\alpha}^{\mu}\log(\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\alpha,\mu}) + (\theta_{\alpha}^{\mu} - 1)\log(\bar{\mathbf{c}}_{\alpha}^{\mu\dagger}\mathbf{b}_{\beta,\nu}) \\ &\quad + \sum_{j\neq\mu}\theta_{\alpha}^j\log(\bar{\mathbf{c}}_{\alpha}^{j\dagger}\mathbf{R}_{\alpha}\mathbf{b}_{\alpha,j}) + \sum_{j\neq\nu}\theta_{\beta}^j\log(\bar{\mathbf{c}}_{\beta}^{j\dagger}\mathbf{Q}\mathbf{b}_{\beta,j}) + \sum_{i\neq\alpha,\beta}\sum_j\theta_i^j\log(\bar{\mathbf{c}}_i^{j\dagger}\mathbf{R}(z_i)\mathbf{b}_{i,j}). \end{aligned}$$

Then, if we restrict to $(\mathcal{B} \times \bar{\mathcal{C}})^{\text{ort}}$, equations (4.9–4.13) are given by

$$\bar{\mathbf{b}}_{i,j} = \frac{\partial \mathcal{H}^+}{\partial \bar{\mathbf{c}}_i^{j\dagger}}(\mathbf{B}, \bar{\mathbf{C}}^{\dagger}) \quad \text{and} \quad \mathbf{c}_i^{j\dagger} = \frac{\partial \mathcal{H}^+}{\partial \mathbf{b}_{i,j}}(\mathbf{B}, \bar{\mathbf{C}}^{\dagger}).$$

Proof. This statement is immediate for all generic indices. The non-trivial part is to check it for derivatives of \mathcal{H}^+ w.r.t. the vectors $\mathbf{b}_{\alpha,\mu}$, $\mathbf{b}_{\beta,\nu}$, $\bar{\mathbf{c}}_{\alpha}^{\mu\dagger}$, and $\bar{\mathbf{c}}_{\beta}^{\nu\dagger}$ that appear in $\mathbf{R}(z_i)$, \mathbf{R}_{α} , and \mathbf{Q} . Using (4.8), we get

$$\begin{aligned} \frac{\partial \mathcal{H}^+}{\partial \bar{\mathbf{c}}_{\alpha}^{\mu\dagger}} &= (\theta_{\alpha}^{\mu} - 1)\frac{\mathbf{b}_{\beta,\nu}}{\bar{\mathbf{c}}_{\alpha}^{\mu\dagger}\mathbf{b}_{\beta,\nu}} + \sum_{j\neq\mu}\theta_{\alpha}^j\frac{(\bar{\mathbf{c}}_{\alpha}^{j\dagger}\mathbf{b}_{\beta,\nu})\partial_{\bar{\mathbf{c}}_{\alpha}^{j\dagger}}\left(-\frac{\bar{\mathbf{c}}_{\alpha}^{\mu\dagger}\mathbf{b}_{\alpha,j}}{\bar{\mathbf{c}}_{\alpha}^{\mu\dagger}\mathbf{b}_{\beta,j}}\right)}{\bar{\mathbf{c}}_{\alpha}^{j\dagger}\mathbf{R}_{\alpha}\mathbf{b}_{\alpha,j}} = (\theta_{\alpha}^{\mu} - 1)\frac{\mathbf{b}_{\beta,\nu}}{\bar{\mathbf{c}}_{\alpha}^{\mu\dagger}\mathbf{b}_{\beta,\nu}} = \bar{\mathbf{b}}_{\alpha,\mu}, \\ \frac{\partial \mathcal{H}^+}{\partial \bar{\mathbf{c}}_{\beta}^{\nu\dagger}} &= (\theta_{\beta}^{\nu} - \theta_{\alpha}^{\mu} + 1)\frac{\mathbf{b}_{\beta,\nu}}{\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\beta,\nu}} + \theta_{\alpha}^{\mu}\frac{\mathbf{b}_{\alpha,\mu}}{\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\alpha,\mu}} - \sum_{j\neq\mu}\theta_{\alpha}^j\frac{\bar{\mathbf{c}}_{\alpha}^{j\dagger}\mathbf{b}_{\beta,\nu}}{\bar{\mathbf{c}}_{\alpha}^{j\dagger}\mathbf{b}_{\alpha,j}}\partial_{\bar{\mathbf{c}}_{\beta}^{\nu\dagger}}\left(\frac{\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\alpha,j}}{\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\beta,\nu}}\right) \\ &\quad - \sum_{j\neq\nu}\theta_{\beta}^j\frac{\bar{\mathbf{c}}_{\beta}^{j\dagger}\mathbf{b}_{\beta,\nu}}{\bar{\mathbf{c}}_{\beta}^{j\dagger}\mathbf{Q}\mathbf{b}_{\beta,j}}\partial_{\bar{\mathbf{c}}_{\beta}^{\nu\dagger}}\left(\frac{\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\beta,j}}{\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\beta,\nu}}\right) + \sum_{i\neq\alpha,\beta}\sum_j\theta_i^j\frac{\left(\frac{z_{\alpha}-z_{\beta}}{z_i-z_{\alpha}}\right)\bar{\mathbf{c}}_i^{j\dagger}\mathbf{b}_{\beta,\nu}}{\bar{\mathbf{c}}_i^{j\dagger}\mathbf{R}(z_i)\mathbf{b}_{i,j}}\partial_{\bar{\mathbf{c}}_{\beta}^{\nu\dagger}}\left(\frac{\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{i,j}}{\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\beta,\nu}}\right) = \bar{\mathbf{b}}_{\beta,\nu}, \end{aligned}$$

since

$$\partial_{\bar{\mathbf{c}}_{\beta}^{\nu\dagger}}\left(\frac{\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{i,j}}{\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\beta,\nu}}\right) = \frac{(\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\beta,\nu})\mathbf{b}_{i,j} - (\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{i,j})\mathbf{b}_{\beta,\nu}}{(\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\beta,\nu})^2} = \frac{\mathbf{Q}\mathbf{b}_{i,j}}{(\bar{\mathbf{c}}_{\beta}^{\nu\dagger}\mathbf{b}_{\beta,\nu})}.$$

The other two equations are established in the same way. \square

Finally, we show that the evolution equations (3.17–3.22) can be obtained directly from the Hamiltonian.

Lemma 4.3. *Equations (4.9–4.13) give the same dynamic $(\mathcal{B} \times \bar{\mathcal{C}})_{(\Theta, \mathbf{A}_{\infty})} \rightarrow (\bar{\mathcal{B}} \times \bar{\mathcal{C}})_{(\bar{\Theta}, \bar{\mathbf{A}}_{\infty})}$ as in Theorem 3.3.*

Proof. Equation (3.17) is just a restatement of (4.10). It then follows that $\mathbf{R}(z)$ is given by (3.4). As usual, the statement is immediate for all indices $i \neq \alpha, \beta$ (note that the relationship between the normalization coefficients for $\bar{\mathbf{b}}_{i,j}$ and $\bar{\mathbf{c}}_i^{j\dagger}$ follows from the requirements (3.12) for the domain and the range of the map). Equation (4.9) gives the expression for $\bar{\mathbf{b}}_{\beta,j}$ in (3.18) for $j \neq \nu$, and right-multiplying $\mathbf{c}_{\alpha}^{j\dagger} \sim \bar{\mathbf{c}}_{\alpha}^{j\dagger}\mathbf{R}_{\alpha}$ in (4.11) by \mathbf{Q} gives $\bar{\mathbf{c}}_{\alpha}^{j\dagger} \sim \mathbf{c}_{\alpha}^{j\dagger}\mathbf{Q}$, which is just the expression for $\bar{\mathbf{c}}_{\alpha}^{j\dagger}$ in (3.18) for $j \neq \mu$. So it remains to establish equations (3.19–3.22). \square

First, we see that (4.12) now becomes

$$(4.16) \quad c_\beta^\nu \bar{\mathbf{b}}_{\beta,\nu} = (\theta_\beta^\nu + 1) \frac{\mathbf{b}_{\beta,\nu}}{\mathbf{c}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu}} + \mathbf{Q} \left(\sum_{i \neq \beta} \left(\frac{z_\alpha - z_\beta}{z_i - z_\beta} \mathbf{A}_i \right) \right) \frac{\mathbf{b}_{\beta,\nu}}{\mathbf{c}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu}} - \sum_{j \neq \nu} \left(\frac{\bar{\mathbf{c}}_\beta^{j\dagger} \mathbf{b}_{\beta,\nu}}{\mathbf{c}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu}} \right) \bar{\mathbf{b}}_{\beta,j}.$$

Left-multiplying it by $\bar{\mathbf{c}}_\beta^{k\dagger}$, $k \neq \mu$, gives

$$(4.17) \quad 0 = (\theta_\beta^\nu + 1 - \theta_\beta^k) \left(\frac{\bar{\mathbf{c}}_\beta^{k\dagger} \mathbf{b}_{\beta,\nu}}{\mathbf{c}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu}} \right) + c_\beta^k c_\beta^{k\dagger} \left(\sum_{i \neq \beta} \left(\frac{z_\alpha - z_\beta}{z_i - z_\beta} \mathbf{A}_i \right) \right) \frac{\mathbf{b}_{\beta,\nu}}{\mathbf{c}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu}}.$$

Solving (4.17) for $(\bar{\mathbf{c}}_\beta^{k\dagger} \mathbf{b}_{\beta,\nu}) / (\bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu})$ and substituting it into (4.16) gives, after simplifying, equation (3.21). Similarly, (4.13) becomes

$$(4.18) \quad \bar{\theta}_\alpha^\mu \frac{\bar{\mathbf{c}}_\alpha^{\mu\dagger}}{\mathbf{c}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu}} - \sum_{j \neq \mu} \frac{\bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\alpha,j}}{\bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu}} \cdot \mathbf{c}_\alpha^{j\dagger} \mathbf{Q} = \frac{\mathbf{c}_\alpha^{\mu\dagger}}{\mathbf{c}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu}} \left((\theta_\alpha^\mu - 1) \mathbf{I} + \sum_{i \neq \alpha} \left(\frac{z_\beta - z_\alpha}{z_i - z_\alpha} \right) \mathbf{A}_i \mathbf{Q} \right).$$

Right-multiplying it by $\mathbf{b}_{\beta,k}$, $k \neq \nu$ gives

$$(4.19) \quad (\bar{\theta}_\alpha^\mu - \theta_\alpha^k) \frac{\bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\alpha,k}}{\bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu}} = \frac{\mathbf{c}_\alpha^{\mu\dagger}}{\mathbf{c}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu}} \left(\sum_{i \neq \alpha} \left(\frac{z_\beta - z_\alpha}{z_i - z_\alpha} \right) \mathbf{A}_i \mathbf{b}_{\alpha,k} \right).$$

Solving (4.19) for $(\bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\alpha,k}) / (\bar{\mathbf{c}}_\alpha^{\mu\dagger} \mathbf{b}_{\beta,\nu})$, substituting it into (4.18), and simplifying gives (3.22). Substituting the expression for $\bar{\mathbf{c}}_\alpha^{\mu\dagger}$ into \mathbf{R}_α in $\mathbf{b}_{\alpha,j} \sim \mathbf{R}_\alpha \mathbf{b}_{\alpha,j}$ and simplifying gives (3.19), where we again use (3.12) to adjust the normalization coefficient. Finally, right-multiplying $\mathbf{c}_\beta^{j\dagger} \sim \bar{\mathbf{c}}_\beta^{j\dagger} \mathbf{Q}$ by $\mathbf{I} - \bar{\mathbf{b}}_{\beta,\nu} (\mathbf{c}_\alpha^{\mu\dagger} \bar{\mathbf{b}}_{\beta,\nu})^{-1} \mathbf{c}_\alpha^{\mu\dagger}$ gives

$$\bar{\mathbf{c}}_\beta^{j\dagger} \sim \mathbf{c}_\beta^{j\dagger} \left(\mathbf{I} - \frac{\bar{\mathbf{b}}_{\beta,\nu} \mathbf{c}_\alpha^{\mu\dagger}}{\mathbf{c}_\alpha^{\mu\dagger} \bar{\mathbf{b}}_{\beta,\nu}} \right).$$

Using the expression for $\bar{\mathbf{b}}_{\beta,\nu}$ and simplifying gives (3.20) and completes the proof of the Lemma. \square

5. EXAMPLES

In this section we consider two explicit examples of reductions from discrete Schlesinger transformations to difference Painlevé equations. We also emphasize the role played by the geometry of Painlevé equations in not only determining the type of the equation, but also in studying the relationship between different explicit forms of equations of the same type.

Recall that, according to the geometric theory of Painlevé equations developed in [Sak01], both continuous and discrete Painlevé equations can be classified by the type of their space of initial conditions, introduced by Okamoto [Oka79] in the continuous case. This space can be obtained by successively blowing up the projective plane \mathbb{P}^2 at nine (possibly infinitely-close) points that lie on a unique (possibly singular) cubic curve. The type of the resulting surface X is then given by the configurations of irreducible components D_i of its *unique* anti-canonical divisor D or, equivalently, by the root sub-lattice R spanned by D_i in the orthogonal complement D^\perp of D in the Picard lattice $\text{Pic}(X)$. The orthogonal complement R^\perp of R in D^\perp then corresponds to the symmetries of the equation — its Weyl group acts as automorphisms, called the *Cremona isometries*, of $\text{Pic}(X)$ preserving $R \subset D^\perp$ and the translational part of the Weyl group corresponds to discrete Painlevé equations.

We also want to stress the following point, see also [Sak07]. Given a discrete Painlevé equation, we can determine its type by constructing its space of initial conditions. However, there is no converse process, although there are some particularly nice forms (that we call *standard*) of equations of each type, see [Sak07, GRO03, Mur04]. Further, there can be many *non-equivalent equations* (i.e., equations that can not be transformed into each other by some change of coordinates) of the same type — such equations can correspond to different translation directions in the affine Weyl group of symmetries of X . And even when the two different equations are equivalent, finding the actual change of coordinates that will transform one equation into the other can be a very challenging problem. In each case, understanding the geometry and the

blowing-down structure of the space of initial conditions can be of great help in determining the relationship between two different equations of the same type, as we illustrate by two examples in this section.

For the first example, we consider the Fuchsian equation of matrix size $m = 2$ with $n = 3$ finite poles and rank-one condition $r_1 = r_2 = r_3 = 1$ for the residue matrices at those poles. It is well-known that continuous isomonodromic deformations of this system are described by the classical continuous Painlevé equation P_{VI} , the type of the space of initial conditions is $D_4^{(1)}$, and the discrete Painlevé V equation $d-P_V = d-P(D_4^{(1)})$ appears as Bäcklund transformations of P_{VI} . Thus, elementary Schlesinger transformations of this system should be described by $d-P_V$, and we explicitly show that this is indeed the case. We consider this example using two different parametrizations. One parametrization is given by the symplectic coordinates for P_{VI} . Unfortunately, in these coordinates the space of initial conditions is not minimal and we need to do blowing-downs in addition to blow-ups. Next, we find an isomorphism between the resulting surface X and the Okamoto surface \tilde{X} of the standard form of $d-P_V$ by finding an appropriate blowing-down structure in $\text{Pic}(X)$ that matches the blowing-down structure of \tilde{X} . This isomorphism can be used to explicitly find the change of coordinates that will transform one equation into the other, see the recent preprint [CT12] by A. Carstea and T. Takenawa for a more detailed introduction into this technique. We then outline how to compute the same example in a different parameterization that seems to be more natural for the discrete case.

The second example is new and is more interesting (and also more complicated). In [Sak07] one of the authors (H.S.) posed a problem of writing down discrete Painlevé equations of type $A_0^{(1)**}$, $A_1^{(1)*}$ and $A_2^{(1)*}$ as Schlesinger transformations of some Fuchsian equations. One difficulty here is that these equations do not correspond to any continuous Painlevé differential equations. This problem was solved theoretically by Boalch in [Boa09], where it was shown that the moduli spaces of monodromy groups of some Fuchsian equations are algebraically isomorphic to the spaces of initial conditions for these discrete Painlevé equations. In particular, Boalch showed that the Fuchsian system with matrix size $m = 3$ with $n = 2$ finite poles (and so there are no continuous deformation parameters) and rank-two residue matrices, $r_1 = r_2 = 2$, has as its symmetry the affine Weyl group of type $E_6^{(1)}$, and so the type of the corresponding surface is either $A_2^{(1)}$ (for the *multiplicative* or *q-difference* case) or $A_2^{(1)*}$ (for the *additive* or *difference* case). Since we consider the additive case, we expect that discrete Schlesinger transformations of this system should have type $d-P(A_2^{(1)*})$. However, an elementary discrete Schlesinger transformation, when written in coordinates, becomes very complicated and in fact it can not be reduced to a standard form of $d-P(A_2^{(1)*})$, as written in [Sak07]. Using the geometry of the space of initial conditions, we see that this is an example of a situation when the standard form of the equation corresponds to a direction that is a combination of two different directions given by different elementary Schlesinger transformations, and then we explicitly show how to combine them into a single Schlesinger transformation that corresponds to the standard form of $d-P(A_2^{(1)*})$, this is a new result.

5.1. Discrete Painlevé–V equation or $d-P(D_4^{(1)})$.

5.1.1. *Elementary Schlesinger transformations.* For this example we consider a Fuchsian system of the spectral type 11, 11, 11, 11, i.e., we have $n = 3$ (finite) poles and the matrix size is $m = 2$. In view of Assumption 2.2, it is possible to make the rank of the residue matrices at the finite poles equal to one. Further, using the Möbius transformations on \mathbb{P}^1 , we can map $(z_0 = \infty, z_1, z_3)$ to $(\infty, 1, 0)$. We put $z_2 = t$ and from now on we use $(1, t, 0)$ instead of the indices $(1, 2, 3)$, so $\theta_2 = \theta_t$, etc. Then

$$\mathbf{A}(z) = \frac{\mathbf{A}_1}{z-1} + \frac{\mathbf{A}_t}{z-t} + \frac{\mathbf{A}_0}{z}, \quad \mathbf{A}_\infty = -(\mathbf{A}_0 + \mathbf{A}_1 + \mathbf{A}_t) = \begin{bmatrix} \kappa_1 & 0 \\ 0 & \kappa_2 \end{bmatrix},$$

and we have the following *Riemann scheme* and the corresponding *Fuchs relation*:

$$\left\{ \begin{array}{cccc} z=0 & z=1 & z=t & z=\infty \\ \theta_0 & \theta_1 & \theta_t & \kappa_1 \\ 0 & 0 & 0 & \kappa_2 \end{array} \right\}, \quad \theta_0 + \theta_1 + \theta_t + \kappa_1 + \kappa_2 = 0.$$

In the rank-one case there is no need for the second index, and so we put $\mathbf{b}_i = \mathbf{b}_{i,1}$, $\mathbf{c}_i^\dagger = \mathbf{c}_i^{1\dagger}$. Then the spectral decomposition is simply $\mathbf{A}_i = \mathbf{b}_i \mathbf{c}_i^\dagger$ with $\text{tr}(\mathbf{A}_i) = \mathbf{c}_i^\dagger \mathbf{b}_i = \theta_i$.

Normalizing the eigenvectors \mathbf{b}_i and \mathbf{c}_i^\dagger , we can write

$$\mathbf{A}_i = a_i \begin{bmatrix} 1 \\ b_i \end{bmatrix} \begin{bmatrix} c_i & 1 \end{bmatrix} = a_i \begin{bmatrix} 1 \\ \beta_i - w \end{bmatrix} \begin{bmatrix} \gamma_i + w & 1 \end{bmatrix}, \quad \text{tr}(\mathbf{A}_i) = \theta_i = a_i(c_i + b_i) = a_i(\beta_i + \gamma_i).$$

Here $w = -b_0$ and so $\beta_0 = 0$ and $\gamma_0 = \theta_0/a_0$. The reason for introducing w is to simplify the relations between parameters a_i , b_i , and c_i that follow from the trace conditions and the equation $\mathbf{A}_\infty = -\text{diag}\{\kappa_1, \kappa_2\}$. These relations, written in terms of a_i , β_i , and γ_i , are

$$(5.1) \quad a_0 + a_1 + a_t = 0$$

$$(5.2) \quad a_1\beta_1\gamma_1 + a_t\beta_t\gamma_t + w(\kappa_1 - \kappa_2) = 0$$

$$(5.3) \quad a_1\beta_1 + a_t\beta_t + \kappa_2 = 0$$

$$(5.4) \quad \theta_0 + a_1\gamma_1 + a_t\gamma_t + \kappa_1 = 0,$$

where the last equation also follows from (5.3) and the Fuchs relation. Thus, everything is parameterized by a_1 , β_1 , a_t , and β_t subject to one additional constraint $a_1\beta_1 + a_t\beta_t + \kappa_2 = 0$. One additional degree of freedom is related to the global gauge action by constant non-degenerate diagonal matrices that allows us to normalize one of the a_i s, but it is convenient to keep this freedom for computations. Finally, note that the normalization parameter a_i should be considered as a part of either \mathbf{b}_i or \mathbf{c}_i^\dagger , where the actual choice depends on the context.

Same coordinates were used in [Sak10] to describe the isomonodromic dynamic in the continuous case and the reduction to P_{VI} . With a_i attached to \mathbf{c}_i^\dagger , the standard symplectic form on the decomposition space reduces as follows, see **Proposition 4** in [Sak10] for a more general statement:

$$\begin{aligned} \omega &= \sum_{i=1}^3 \text{tr} \left(d\mathbf{C}_i^\dagger \wedge d\mathbf{B}_i \right) = da_0 \wedge d(-w) + da_1 \wedge d(\beta_1 - w) + da_t \wedge d(\beta_t - w) \\ &= d(a_0 + a_1 + a_t) \wedge d(-w) + da_1 \wedge d\beta_1 + da_t \wedge d\beta_t = da_1 \wedge d\beta_1 + da_t \wedge d\beta_t \\ &= \frac{da_1}{a_1} \wedge d(a_1\beta_1) + \frac{da_t}{a_t} \wedge d(a_t\beta_t) = \left(\frac{da_t}{a_t} - \frac{da_1}{a_1} \right) \wedge d(a_t\beta_t) = d \left(\frac{a_t}{a_1} \right) \wedge d(a_1\beta_t), \end{aligned}$$

where we used (5.3) to eliminate $a_1\beta_1$. It is convenient to introduce new symplectic coordinates $p = (a_1\beta_t)/t$ and $q = -(ta_t)/a_1$. This gives, using (5.1–5.4),

$$\begin{aligned} a_1\beta_1 &= pq - \kappa_2, & a_1\beta_t &= tp, & a_1\gamma_1 &= \theta_1 + \kappa_2 - pq, \\ a_t\beta_t &= -pq, & a_t\beta_1 &= \frac{q}{t}(\kappa_2 - pq), & a_t\gamma_t &= \theta_t + pq. \end{aligned}$$

For the continuous deformations, the only deformation parameter is t and the corresponding Hamiltonian, given by equation (2.4), under this reduction becomes

$$\mathcal{H}_t = \frac{\text{tr}(\mathbf{A}_t\mathbf{A}_0)}{t} + \frac{\text{tr}(\mathbf{A}_t\mathbf{A}_1)}{t-1} = \frac{(\theta_t + pq)(\theta_0 + p(q-t))}{t} + \frac{(\theta_1 + \kappa_2 - p(q-t))(t\theta_t + \kappa_2q - pq(q-t))}{t(t-1)}.$$

Since

$$\begin{aligned} \omega &= d \left(-\frac{q}{t} \right) \wedge d(tp) = dp \wedge dq - d \left(\frac{pq}{t} \right) \wedge dt, \\ \omega - d\mathcal{H}_t \wedge dt &= dp \wedge dq - d \left(\mathcal{H}_t + \frac{pq}{t} \right) \wedge dt = dp \wedge dq - d\mathcal{H}_{VI} \wedge dt, \end{aligned}$$

where $\mathcal{H}_{VI} = \mathcal{H}_t + pq/t$ is the Hamiltonian function for P_{VI} .

Unfortunately, such a reduction procedure does not work in the discrete case and we need to work directly with the evolution equations. We give a brief outline of such computations and show the reduction to d- P_V .

From equations (3.7–3.9), that are more convenient here, we almost immediately get the following relations:

$$(5.5) \quad \bar{w} - w = \bar{\beta}_1 - \beta_t = \gamma_1 - \bar{\gamma}_t \quad (\text{and so } \bar{\beta}_1 + \bar{\gamma}_t = \beta_t + \gamma_1), \quad \beta_t\bar{\gamma}_t = t\bar{\beta}_1\gamma_1.$$

Multiplying the last equation in (5.5) by $a_1\bar{a}_t$, we get

$$(5.6) \quad (\theta_1 + \kappa_2 - pq)\bar{q}(\bar{\kappa}_2 - \bar{p}\bar{q}) = tp(\bar{\theta}_t + \bar{p}\bar{q}).$$

Combining the first equation in (5.5) with (5.2) gives

$$a_1\beta_1(\gamma_1 + \beta_t) + \beta_t(a_t\gamma_t - a_1\beta_1 + \kappa_1 - \kappa_2) = \bar{a}_1\bar{\beta}_t(\bar{\gamma}_t + \bar{\beta}_1) + \bar{\beta}_1(\bar{a}_1\bar{\gamma}_1 - \bar{a}_t\bar{\beta}_t + \kappa_1 - \kappa_2).$$

Dividing by $\gamma_1 + \beta_t = \bar{\gamma}_t + \bar{\beta}_1$ and doing some re-arranging (and using (5.6) in the process), we get

$$(5.7) \quad pq + \bar{p}\bar{q} - \kappa_2 = \frac{t(\theta_t + \kappa_1)}{q - t - \frac{\theta_1 + \kappa_2}{p}} + \frac{\bar{\theta}_1 + \kappa_1}{q - 1 - \frac{\theta_1 + \kappa_2}{p}}.$$

We now claim that after a certain change of variables equations (5.6) and (5.7) transform into the standard form for $d-P_V$. Although this can be seen directly without too much difficulty, we prefer to use the geometry of the Okamoto space of initial conditions for this equation to explain how to find this coordinate transformation, illustrating how such geometric machinery can be used.

5.1.2. *Geometry of $d-P(D_4^{(1)})$.* Let us briefly review the geometric description of the difference Painlevé V equation from the point of view of the theory of rational surfaces [Sak01]. We think about it as a birational map $\varphi : \mathbb{P}^1 \times \mathbb{P}^1 \dashrightarrow \mathbb{P}^1 \times \mathbb{P}^1$ with parameters a_0, \dots, a_4, s or, equivalently, as an automorphism of a field $\mathbb{C}(a_0, \dots, a_4; s; f, g)$,

$$\left(\begin{array}{ccc} a_1 & a_2 & a_3 \\ a_4 & a_0 & \end{array} ; s; f, g \right) \mapsto \left(\begin{array}{ccc} \bar{a}_1 & \bar{a}_2 & \bar{a}_3 \\ \bar{a}_4 & \bar{a}_0 & \end{array} ; \bar{s}; \bar{f}, \bar{g} \right), \quad \begin{array}{lll} \bar{a}_1 = a_1 + \delta, & \bar{a}_2 = a_2 - \delta, & \bar{a}_3 = a_3, \\ \bar{a}_4 = a_4, & \bar{a}_0 = a_0 + \delta, & \bar{s} = s, \end{array}$$

where $\delta = a_0 + a_1 + 2a_2 + a_3 + a_4$ and where \bar{f} and \bar{g} are given by the following system that is usually called the discrete Painlevé-V equation

$$(5.8) \quad \begin{cases} f + \bar{f} = a_3 + \frac{a_1}{g+1} + \frac{a_0}{sg+1} \\ g\bar{g} = \frac{(\bar{f} + \bar{a}_2)(\bar{f} + \bar{a}_2 + \bar{a}_4)}{s\bar{f}(\bar{f} - \bar{a}_3)} \end{cases};$$


we refer to it as the standard form of $d-P_V$. We lift this map to an *isomorphism* $\hat{\varphi}$ between rational surfaces $X_{\mathbf{a}} = X_{(a_0, \dots, a_4)}$ and $X_{\bar{\mathbf{a}}}$ obtained from $\mathbb{P}^1 \times \mathbb{P}^1$ via a sequence of blow-ups that resolve the indeterminacies of the map φ ,

$$\begin{array}{ccc} X_{\mathbf{a}} & \xrightarrow[\hat{\varphi}]{\cong} & X_{\bar{\mathbf{a}}} \\ \downarrow & & \downarrow \\ \mathbb{P}^1 \times \mathbb{P}^1 & \xrightarrow[\varphi]{} & \mathbb{P}^1 \times \mathbb{P}^1 \end{array}.$$

Decomposing φ as a sequence of two maps, $(f, g) \xrightarrow{\varphi_1} (\bar{f}, g) \xrightarrow{\varphi_2} (\bar{f}, \bar{g})$ it is easy to see that the indeterminate points of φ_1 are $(\infty, -1)$ and $(\infty, -1/s)$ and the indeterminate points of φ_2 are $(-\bar{a}_2, 0)$, $(-\bar{a}_2 - \bar{a}_4, 0)$, $(0, \infty)$, and (\bar{a}_3, ∞) . Applying φ_1^{-1} we obtain that the indeterminate points of φ are $p_1(-a_2 - a_4, 0)$, $p_2(-a_2, 0)$, $p_3(0, \infty)$, $p_4(a_3, \infty)$, $p_5(\infty, -1)$, and $p_6(\infty, -1/s)$. In exactly the same way we find that the indeterminate points of φ^{-1} are $\bar{p}_1(-\bar{a}_2 - \bar{a}_4, 0)$, \dots , $\bar{p}_6(\infty, -1/\bar{s})$.

As usual, we denote a local coordinate description of a blow-up at a point (x_0, y_0) by $(x, y) \leftarrow (u, v) \cup (U, V)$ where $x - x_0 = u = UV$ and $y - y_0 = uv = V$, then the exceptional divisor is given by $u = V = 0$. For example, extending the map φ at the blow-up at $p_1(-a_2 - a_4, 0)$, we use $u_1 = f + a_2 + a_4$ and $u_1v_1 = g$. Then $\bar{f} = \bar{f}(u_1, v_1)$ and $\bar{g} = g(u_1, v_1)$ become well-defined and we see that the exceptional divisor E_1 is mapped to the line $f = -\bar{a}_2$. On the level of divisor classes on $\text{Pic}(X)$ we get the map $E_1 \mapsto H_f - E_2$, where H_f is the class of the total transform of the line $f = \text{const}$.

Proceeding along the same lines and noticing that we need to blow-up at two additional points $p_7(0, a_1)$ on E_5 and $p_8(0, a_0/s)$ on E_7 (here we give the coordinates of the points in the (u_i, v_i) charts), we get the following blow-up diagram describing the construction of the surface $X_{\mathbf{a}}$ corresponding to our equation:



The Picard lattice of X is generated by the total transforms H_f and H_g of the coordinate lines and the classes of the exceptional divisors E_i ,

$$\text{Pic}(X) = \mathbb{Z}H_f \oplus \mathbb{Z}H_g \oplus \bigoplus_{i=1}^8 \mathbb{Z}E_i.$$


The anti-canonical divisor $-K_X$ uniquely decomposes as a positive linear combination of -2 -curves D_i ,

$$-K_X = 2H_f + 2H_g - \sum_{i=1}^8 E_i = D_0 + D_1 + 2D_2 + D_3 + D_4,$$

where


$$D_0 = H_g - E_1 - E_2, \quad D_1 = H_g - E_3 - E_4, \quad D_2 = H_f - E_5 - E_6, \quad D_3 = E_5 - E_7, \quad D_4 = E_6 - E_8.$$

The configuration of these -2 -curves (on which the blow-up points are located) is described by the affine Dynkin diagram of type $D_4^{(1)}$,



which is why we say that the difference Painlevé V equation has the type $d\text{-}P(D_4^{(1)})$. The corresponding divisors D_i generate a sub-lattice $R = \text{Span}_{\mathbb{Z}}\{D_0, \dots, D_4\} \subset (-K_X)^\perp \simeq E_8^{(1)}$ in $\text{Pic}(X)$. Symmetries of X are given by the affine Weyl group of *Cremona isometries* generated by the *complementary* root sub-lattice R^\perp in $(-K_X)^\perp$ acting as automorphisms of $\text{Pic}(X)$. In this case, $R^\perp = \text{Span}_{\mathbb{Z}}\{\alpha_0, \dots, \alpha_4\}$, where

$$\begin{aligned} \alpha_0 &= E_1 - E_2 \\ \alpha_1 &= E_3 - E_4 \\ \alpha_2 &= H_f - E_1 - E_3 \\ \alpha_3 &= H_g - E_5 - E_7 \\ \alpha_4 &= H_g - E_6 - E_8 \end{aligned}$$




which is again of the type $D_4^{(1)}$. Finally, we can compute the action of φ_* on $\text{Pic}(X)$ to be

$$\begin{aligned} H_f &\mapsto 5H_f + 2H_g - 2E_1 - 2E_2 - 2E_3 - 2E_4 - E_5 - E_6 - E_7 - E_8, & E_1 &\mapsto H_f - E_2, \\ H_g &\mapsto 2H_f + H_g - E_1 - E_2 - E_3 - E_4, & E_2 &\mapsto H_f - E_1, \\ E_5 &\mapsto 2H_f + H_g - E_1 - E_2 - E_3 - E_4 - E_8, & E_3 &\mapsto H_f - E_4, \\ E_6 &\mapsto 2H_f + H_g - E_1 - E_2 - E_3 - E_4 - E_7, & E_4 &\mapsto H_f - E_3, \\ E_7 &\mapsto 2H_f + H_g - E_1 - E_2 - E_3 - E_4 - E_6, \\ E_8 &\mapsto 2H_f + H_g - E_1 - E_2 - E_3 - E_4 - E_5, \end{aligned}$$

and so the induced action φ_* on the sub-lattice R^\perp is given by *translation*:

$$\varphi_* : (\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4) \mapsto (\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4) + (0, 0, 1, -1, -1)\delta, \quad \text{where } \delta = -K_X.$$


5.1.3. *Reduction to the standard form.* We now demonstrate how to reduce equations (5.6, 5.7) to equations (5.8). First, we resolve the indeterminacies of $\psi : (p, q) \rightarrow (\bar{p}, \bar{q})$ to get an isomorphism $\hat{\psi} : \tilde{X}_{\mathbf{a}} \rightarrow \tilde{X}_{\bar{\mathbf{a}}}$. The surface $\tilde{X} = \tilde{X}_{\mathbf{a}}$, with \mathbf{a} denoting the collection of parameters of the map, $\mathbf{a} = \{\theta_0, \theta_1, \theta_t, \kappa_1, \kappa_2\}$, is constructed by the following blow-up diagram:



Here the coordinates of the blow-up points are given w.r.t. the following local coordinate systems:

$$\begin{aligned} p_1 \left(\frac{1}{p} = 0, q = 0 \right) &\leftarrow p_2 \left(\frac{1}{p} = 0, pq = -\theta_t \right) \text{ and } p'_2 \left(\frac{1}{pq} = 0, q = 0 \right), \\ p_3 \left(p = 0, \frac{1}{q} = 0 \right) &\leftarrow p_4 \left(p = 0, \frac{1}{pq} = \frac{1}{\kappa_2} \right) \text{ and } p'_4 \left(p = 0, \frac{1}{pq} = \frac{1}{\theta_1 + \kappa_2} \right), \\ p_5 \left(\frac{1}{p} = 0, q = t \right) &\leftarrow p_7 \left(\frac{1}{p} = 0, p(q - t) = -\theta_0 \right), \\ p_6 \left(\frac{1}{p} = 0, q = 1 \right) &\leftarrow p_8 \left(\frac{1}{p} = 0, p(q - 1) = 1 - \kappa_1 + \kappa_2 \right). \end{aligned}$$

Note that the resulting configuration of exceptional divisors looks quite different from the previous case and moreover, $\text{rank}(\text{Pic}(\tilde{X})) = 12$. The reason for this is that the surface \tilde{X} is not *relatively minimal* for $\hat{\psi}$, see also [CT12] — there are two pairs of -1 -curves that are exchanged by $\hat{\psi}$: $F_2 \leftrightarrow F'_4$ and $F'_2 \leftrightarrow H_q - F_3$, and so blowing down any of these pairs will still result in an isomorphism between the corresponding surfaces. We blow down F'_2 and $H_q - F_3$, which eliminates -4 and -3 -curves and gives us the diagram on the left. This diagram now looks very similar to the configuration we had for the standard form of d - P_V (on the right),



and in particular we immediately see that \tilde{X} is again of type $D_4^{(1)}$. We now need to find the blowing-down structure for \tilde{X} to match the blowing-down structure for X . Thus, in $\text{Pic}(\tilde{X})$ we put $\mathcal{E}_9 = F'_2$, $\mathcal{E}_{10} = H_q - F_3$ and we are looking for *rational* 0 -curves $\mathcal{H}_f, \mathcal{H}_g$, and -1 -curves $\mathcal{E}_i, i = 1, \dots, 8$ such that

$$\mathcal{H}_f \bullet \mathcal{H}_g = 1, \quad \mathcal{E}_i^2 = -1, \quad \mathcal{H}_f^2 = \mathcal{H}_g^2 = \mathcal{H}_f \bullet \mathcal{E}_i = \mathcal{H}_g \bullet \mathcal{E}_i = \mathcal{E}_i \bullet \mathcal{E}_j = 0, \quad 1 \leq i \neq j \leq 10.$$

Using the above diagrams, the genus formula $2g(C) - 2 = C^2 + C \bullet K_{\bar{X}}$, and the distinguished -2 -curve D_2 , we see that it makes sense to take $\mathcal{H}_f = H_p + H_q - F_1 - F_3$ and $\mathcal{E}_i = F_i$, $i = 5, \dots, 8$. Then the possible choices for \mathcal{H}_g are H_q , $H_p + H_q - F_2 - F_3$, $H_p + H_q - F_4 - F_3$, and $H_p + H_q - F'_4 - F_3$. Attempting to represent $F_1 - F_2 = \mathcal{H}_g - \mathcal{E}_1 - \mathcal{E}_2$ eliminates the first two possibilities, and the remaining two are equivalent. To summarize, we have the following blowing-down structure:

$$\begin{aligned} \mathcal{H}_f &= H_p + H_q - F_1 - F_3, & \mathcal{E}_3 &= H_p - F_3, & \mathcal{E}_i &= F_i, & i &= 2, 4, \dots, 8, \\ \mathcal{H}_g &= H_p + H_q - F_3 - F'_4, & \mathcal{E}_9 &= F'_2, \\ \mathcal{E}_1 &= H_p + H_q - F_1 - F_3 - F'_4, & \mathcal{E}_{10} &= H_q - F_3. \end{aligned}$$

To find the expressions for coordinates f and g we proceed as follows. The linear system $|\mathcal{H}_f|$ consists of curves of bi-degree $(1, 1)$ passing through p_1 and p_3 , and so $|\mathcal{H}_f| = \{c_1 pq + c_2 = 0\}$. Thus, $[pq : 1]$ is a projective coordinate on $|\mathcal{H}_f|$, and so we can take $f = pq$. Similarly, $[pq - (\theta_1 + \kappa_2) : p]$ is a projective coordinate on $|\mathcal{H}_g|$. We take $g = -\frac{1}{t} \left(q - \frac{\theta_1 + \kappa_2}{p} \right)$, where the additional factor $-1/t$ is needed to ensure that the (f, g) -coordinates of p_5 are $(\infty, -1)$, as in the standard case.

We can now match the parameters of our elementary Schlesinger transformations with the parameters s and a_i in the standard form (5.8) of d - P_V . For example, \mathcal{E}_1 blows down to $(f, g) = (\theta_1 + \kappa_2, 0) = (-a_2 - a_4, 0)$. Doing this for other divisors \mathcal{E}_i results in the following

$$a_0 = \theta_1 - 1 + k_1, \quad a_1 = -\theta_t - k_1, \quad a_2 = \theta_t, \quad a_3 = \kappa_2, \quad a_4 = -\theta_1 - \theta_t - \kappa_2, \quad \delta = -1.$$

Thus, not only we can now immediately see that equations (5.6, 5.7) transform into the standard form (5.8) of d - P_V , but also that under this identification of parameters,

$$\bar{\theta}_1 = -\bar{a}_2 - \bar{a}_3 - \bar{a}_4 = -a_2 + \delta - a_3 - a_4 = \theta_1 - 1, \quad \bar{\theta}_t = -\bar{a}_0 - \bar{a}_1 - \bar{a}_2 - \bar{a}_3 - \bar{a}_4 - 1 = \theta_t + 1,$$

and the other parameters remain unchanged. Thus, we can *identify the action of d - P_V on the Riemann Scheme* of the Fuchsian system under this correspondence. We use the same observation in the next example.

5.1.4. Different Parameterization. A more direct derivation of d - P_V can be obtained if we make a different coordinate choice (which seems to be more natural for Schlesinger transformations and is used in the next example as well).

We use the global gauge group action to map vectors \mathbf{b}_t and \mathbf{b}_1 to the standard basis vectors and then use the trivial transformations to make components of \mathbf{b}_0 equal to 1 to get the following parameterization:

$$\mathbf{b}_t = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad \mathbf{c}_t = [\theta_t \quad \alpha], \quad \mathbf{b}_1 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad \mathbf{c}_1 = [\beta \quad \theta_1], \quad \mathbf{b}_0 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad \mathbf{c}_0 = [\gamma \quad \theta_0 - \gamma],$$

where three parameters α , β , and γ are constrained by the condition that the eigenvalues of \mathbf{A}_∞ are κ_1, κ_2 . Indeed, with this parameterization,

$$-\mathbf{A}_\infty = \begin{bmatrix} \theta_t + \gamma & \theta_0 + \alpha - \gamma \\ \beta + \gamma & \theta_0 + \theta_1 - \gamma \end{bmatrix}, \quad \begin{aligned} \text{tr } \mathbf{A}_\infty &= \kappa_1 + \kappa_2 = -(\theta_0 + \theta_1 + \theta_t), \\ \det \mathbf{A}_\infty &= \kappa_1 \kappa_2 = (\theta_0 + \theta_1 - \gamma)(\theta_t + \gamma) - (\theta_0 + \alpha - \gamma)(\beta + \gamma). \end{aligned}$$


The trace condition is just the Fuchs relation, and the determinant condition allow us to express, e.g., α in terms of β and γ . Thus, if we put $\beta = x$ and $\gamma = y$, $\alpha(x, y)$ is given by the expression

$$\alpha(x, y) = \frac{x(y - \theta_0) + y(\theta_1 - \theta_t) + (\theta_1 + \kappa_1 + \theta_0)(\theta_t + \kappa_1)}{x + y}.$$

Using the evolution equations (3.17–3.22) for the elementary Schlesinger transformation $\left\{ \begin{smallmatrix} 1 & t \\ & 1 \end{smallmatrix} \right\}$ then gives us the following map $\chi : (x, y) \rightarrow (\bar{x}, \bar{y})$:

$$(5.9) \quad \left\{ \begin{aligned} \bar{x} &= \frac{((x + \theta_1)y - \theta_0 x)((t - 1)((x + \theta_1)y + \theta_1 x) + x(tx - \theta_t))(t(x + y) - y)}{tx((t(x + y) - y)(\theta_1 + x) - x(1 + \theta_t))} \\ &\quad + \frac{tx(x + y)((\theta_1 - 1)(\theta_t + 1) - \kappa_1 \kappa_2) + xy\kappa_1 \kappa_2}{t((t(x + y) - y)(\theta_1 + x) - x(1 + \theta_t))} \\ \bar{y} &= \frac{(t(x + y) - y)(\theta_0 x - (x + \theta_1)y) + xy(\theta_t + 1)}{x(t(x + y) - y)} \end{aligned} \right.$$

The Okamoto surface X for this map is given by the blow-up diagram



where the coordinates of the blow-up points are

$$\begin{aligned}
p_1(x = \theta_t + \kappa_1, y = -\theta_t - \kappa_1), \quad p_2(x = \theta_t + \kappa_2, y = -\theta_t - \kappa_2), \quad p_3(x = 0, y = 0), \\
p_4\left(\frac{1}{x} = 0, y = \theta_0\right), \quad p_6\left(x = -\theta_1, \frac{1}{y} = 0\right), \\
p_5\left(\frac{1}{x} = 0, \frac{1}{y} = 0\right) \leftarrow p_7\left(\frac{1}{x} = 0, \frac{x}{y} = \frac{1-t}{t}\right) \leftarrow p_8\left(\frac{1}{x} = 0, t(x+y) - y = \theta_t + 1\right).
\end{aligned}$$

We compute the action of χ_* on $\text{Pic}(X)$ to be

$$\begin{aligned}
H_x &\mapsto 2H_x + 3H_y - 2F_1 - 2F_2 - F_4 - F_5 - F_6 - F_7, & F_1 &\mapsto H_y - F_2, \\
H_y &\mapsto 2H_x + 4H_y - 2F_1 - 2F_2 - 2F_4 - F_5 - F_6 - F_7 - F_8, & F_2 &\mapsto H_y - F_1, \\
F_3 &\mapsto 2H_x + 3H_y - 2F_1 - 2F_2 - F_4 - F_5 - F_6 - F_7 - F_8, & F_4 &\mapsto F_3, \\
F_5 &\mapsto H_x + 2H_y - F_1 - F_2 - F_4 - F_6 - F_7, & F_6 &\mapsto H_x + 2H_y - F_1 - F_2 - F_4 - F_5 - F_7, \\
F_7 &\mapsto H_x + 2H_y - F_1 - F_2 - F_4 - F_5 - F_6, & F_8 &\mapsto H_x + H_y - F_1 - F_2 - F_4.
\end{aligned}$$

To match this picture with the standard one, we look at the -2 -curves. The distinguished -2 -curve $D_2 = H_f - E_5 - E_6$ should correspond to $F_5 - F_7$. Next, if we let $D_0 = H_g - E_1 - E_2$ correspond to $H_x + H_y - F_1 - F_2 - F_3 - F_5$, $D_1 = H_g - E_3 - E_4$ should correspond to $H_x - F_4 - F_5$ to match the dynamic $D_0 \leftrightarrow D_1$. Let $D_3 = E_5 - E_7$ correspond to $H_y - F_5 - F_6$, then $D_4 = E_6 - E_7$ corresponds to the remaining -2 -curve $F_7 - F_8$. A change of basis giving such identification can be taken as

$$\mathcal{H}_f = H_y, \quad \mathcal{H}_g = H_x + H_y - F_3 - F_5, \quad \mathcal{E}_3 = H_y - F_3, \quad \mathcal{E}_5 = H_y - F_5, \quad \mathcal{E}_6 = F_7, \quad \mathcal{E}_7 = F_6,$$

and $\mathcal{E}_i = F_i$ for $i = 1, 2, 4, 8$. Under this identification the action of χ_* on the symmetry sub-lattice R^\perp coincides with the action of φ_* .

Let us now find the new coordinates f and g . Since $|\mathcal{H}_f| = |H_y|$, we can take $f = y$. The linear system $|\mathcal{H}_g|$ consists of curves of bi-degree $(1, 1)$ passing through $p_3(0, 0)$ and $p_5(\infty, \infty)$ and so $|\mathcal{H}_g| = \{c_1x + c_2y = 0\}$. Thus, $[x : y]$ is a projective coordinate on $|\mathcal{H}_g|$, and so we can provisionally put $g = x/y$, which gives $g(p_1) = g(p_2) = -1$, $g(p_3) = g(p_4) = \infty$, $g(p_5) = 0$, and we want them to be $0, \infty$, and -1 respectively. This is easy to fix by the Möbius transformation $g \mapsto -(g+1)$, and so our final change of coordinates is $f = y$ and $g = -(x+y)/y$. It is then easy to see that under this change of coordinates equations (5.9) transforms into the standard form (5.8). The matching of the standard parameters with the Fuchsian system parameters is given by

$$a_0 = -(\theta_t + 1), \quad a_1 = \theta_1, \quad a_2 = \theta_t + \kappa_2, \quad a_3 = \theta_0, \quad a_4 = \kappa_1 - \kappa_2, \quad s = t, \quad \delta = -1,$$

and under this identification of parameters,

$$\bar{a}_0 = -(\bar{\theta}_t + 1) = a_0 + \delta, \quad \bar{a}_1 = \bar{\theta}_1 = a_1 + \delta, \quad \bar{a}_2 = \bar{\theta}_t + \bar{\kappa}_2 = a_2 - \delta, \quad \bar{a}_3 = \theta_0 = a_3, \quad \bar{a}_4 = \bar{\kappa}_1 - \bar{\kappa}_2 = a_4,$$

as it should be.

5.2. Discrete Painlevé equation $d-P(A_2^{(1)*})$.

5.2.1. *Schlesinger Transformations.* We now consider Fuchsian systems of the spectral type $111, 111, 111$, which corresponds to $n = 2$ (finite) poles and the matrix size $m = 3$. As before, we make $\text{rank}(\mathbf{A}_i) = 2$ using scalar gauge transformations and map the finite poles to $z_0 = 0$ and $z_1 = 1$ by a Möbius transformation. Note that this example does not have any continuous deformation parameters but it still admits non-trivial Schlesinger transformations. So

$$\mathbf{A}(z) = \frac{\mathbf{A}_0}{z} + \frac{\mathbf{A}_1}{z-1}, \quad \mathbf{A}_i = \mathbf{B}_i \mathbf{C}_i^\dagger = [\mathbf{b}_{i,1} \quad \mathbf{b}_{i,2}] \begin{bmatrix} \mathbf{c}_i^{1\dagger} \\ \mathbf{c}_i^{2\dagger} \end{bmatrix},$$

and the corresponding *Riemann scheme* and the *Fuchs relation* are

$$\left\{ \begin{array}{ccc} z=0 & z=1 & z=\infty \\ \theta_0^1 & \theta_1^1 & \kappa_1 \\ \theta_0^2 & \theta_1^2 & \kappa_2 \\ 0 & 0 & \kappa_3 \end{array} \right\}, \quad \theta_0^1 + \theta_0^2 + \theta_1^1 + \theta_1^2 + \sum_{j=1}^3 \kappa_j = 0.$$

As in the previous example, we use the global similarity transformations to map the vectors $\mathbf{b}_{0,1}$, $\mathbf{b}_{0,2}$, and $\mathbf{b}_{1,1}$ to the standard basis vectors and then use the trivial transformations to adjust the scale on the coordinate axes so that all components of $\mathbf{b}_{1,2}$ are equal to 1. Then, using the condition $\mathbf{C}_i^\dagger \mathbf{B}_i = \Theta_i$, we get the following parameterization:

$$\mathbf{B}_0 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad \mathbf{C}_0^\dagger = \begin{bmatrix} \theta_0^1 & 0 & \alpha \\ 0 & \theta_0^2 & \beta \end{bmatrix}, \quad \mathbf{B}_1 = \begin{bmatrix} 0 & 1 \\ 0 & 1 \\ 1 & 1 \end{bmatrix}, \quad \mathbf{C}_1^\dagger = \begin{bmatrix} -\gamma - \theta_1^1 & \gamma & \theta_1^1 \\ \theta_1^2 - \delta & \delta & 0 \end{bmatrix}.$$

Requiring that the eigenvalues of \mathbf{A}_∞ are κ_1 , κ_2 , and κ_3 results in the equations $\text{tr}(\mathbf{A}_\infty) = \kappa_1 + \kappa_2 + \kappa_3$ (which is just the Fuchs relation), $|\mathbf{A}_\infty|_{11} + |\mathbf{A}_\infty|_{22} + |\mathbf{A}_\infty|_{33} = \kappa_2 \kappa_3 + \kappa_3 \kappa_1 + \kappa_1 \kappa_2$ (where $|\mathbf{A}|_{ij}$ denotes the (ij) -minor of \mathbf{A}) and $\det(\mathbf{A}_\infty) = \kappa_1 \kappa_2 \kappa_3$. The last two equations are linear in α , β , γ , and δ so we can write them as a linear system on α and β :

$$(5.10) \quad (\gamma + \delta + \theta_1^1 - \theta_1^2)\alpha - (\gamma + \delta)\beta = \kappa_2 \kappa_3 + \kappa_3 \kappa_1 + \kappa_1 \kappa_2 + (\theta_0^2 - \theta_0^1)\delta - (\theta_0^2 + \theta_1^1)(\theta_0^1 + \theta_1^2) - \theta_0^2 \theta_1^1, \\ -(\theta_0^2(\gamma + \delta + \theta_1^1 - \theta_1^2) + \theta_1^2 \gamma + \theta_1^1 \delta)\alpha + (\theta_0^1(\gamma + \delta) + \theta_1^2 \gamma + \theta_1^1 \delta)\beta = \kappa_1 \kappa_2 \kappa_3 + \theta_1^1((\theta_0^1 - \theta_0^2)\delta + \theta_0^2(\theta_0^1 + \theta_1^2)).$$

At this point we can choose parameterization variables x and y to simplify the structure of the substitution rule. Noticing that the coefficients of the matrix of the above linear system are written in terms of the expressions $\gamma + \delta$, $\gamma + \delta + \theta_1^1 - \theta_1^2$, and $\theta_1^2 \gamma + \theta_1^1 \delta$, we make the following choice of coordinates:

$$(5.11) \quad x = \frac{(\gamma + \delta)(\theta_0^1 - \theta_0^2)}{\theta_1^1 - \theta_1^2}, \quad y = \frac{\theta_1^2 \gamma + \theta_1^1 \delta}{\gamma + \delta + \theta_1^1 - \theta_1^2}.$$

The reasons for this particular choice are first to simplify the entries of the matrix of the above linear system, and second to simplify its determinant, which is the denominator for the expressions $\alpha(x, y)$ and $\beta(x, y)$ — its zeroes contribute to the indeterminate points for $\alpha(x, y)$ and $\beta(x, y)$ and through that to the indeterminate points of the dynamic as well. We then get

$$\alpha(x, y) = \frac{1}{(x+y)(\theta_1^1 - \theta_1^2)} \left(yr_1 + \frac{x(\theta_0^2 r_1 + r_2)}{x + \theta_0^1 - \theta_0^2} \right), \quad \beta(x, y) = \frac{1}{(x+y)(\theta_1^1 - \theta_1^2)} ((y + \theta_0^2)r_1 + r_2),$$

where


$$r_1 = r_1(x, y) = \kappa_1 \kappa_2 + \kappa_2 \kappa_3 + \kappa_3 \kappa_1 - (y - \theta_1^2)(x - \theta_0^2) - \theta_0^1(y + \theta_0^2) - \theta_1^1(\theta_0^1 + \theta_0^2 + \theta_1^2), \\ r_2 = r_2(x, y) = \kappa_1 \kappa_2 \kappa_3 + \theta_1^1((y - \theta_1^2)(x - \theta_0^2) + \theta_0^1(y + \theta_0^2))$$

are the right-hand-sides of the linear system (5.10) (written now in variables x and y).

Using the evolution equations (3.17–3.22) for the elementary Schlesinger transformation $\begin{Bmatrix} 0 & 1 \\ 1 & 1 \end{Bmatrix}$ then gives us the following map $\varphi : (x, y) \rightarrow (\bar{x}, \bar{y})$:

$$\begin{cases} \bar{x} = \frac{(\alpha - \beta)(\alpha x(\theta_1^1 - \theta_1^2) + (1 + \theta_0^2)(x(y - \theta_1^2) + y(\theta_0^1 - \theta_0^2)))}{(\alpha - \beta)(x(y - \theta_1^2) + (\theta_0^1 - \theta_0^2)y) - \alpha(\theta_1^1 + 1)(\theta_0^1 - \theta_0^2)} \\ \bar{y} = \frac{(\alpha - \beta)(y(x + \theta_0^1 - \theta_0^2) - \theta_1^2 x)}{\alpha(\theta_0^1 - \theta_0^2)} \end{cases},$$

where we still need to substitute $\alpha = \alpha(x, y)$ and $\beta = \beta(x, y)$, but this can be easily done using computer algebra. We claim that the resulting dynamical system has the type $A_2^{(1)*}$, as expected. Indeed, the Okamoto surface for the map $\psi : (x, y) \rightarrow (\bar{x}, \bar{y})$ is given by the blow-up diagram




where the coordinates of the blow-up points are

$$\begin{aligned} p_1(\theta_0^2 + \kappa_1, -\theta_0^2 - \kappa_1), \quad p_3(\theta_0^2 + \kappa_3, -\theta_0^2 - \kappa_3), \quad p_5(\infty, \theta_1^1), \quad p_7(\theta_0^2 - \theta_0^1, \infty), \\ p_2(\theta_0^2 + \kappa_2, -\theta_0^2 - \kappa_2), \quad p_4(0, 0), \quad p_6(\infty, \theta_1^2), \quad p_8(\theta_0^2 + 1, \infty). \end{aligned}$$

The anti-canonical divisor $-K_X = 2H_f + 2H_g - \sum_{i=1}^8 E_i$ uniquely decomposes as a positive linear combination of -2 -curves D_i ,

$$-K_X = D_0 + D_1 + D_2, \quad \text{where } D_0 = H_x + H_y - F_1 - F_2 - F_3 - F_4, \quad D_1 = H_x - F_5 - F_6, \quad D_2 = H_y - F_7 - F_8.$$

So we see that the configuration of components D_i is indeed described by the Dynkin diagram of type $A_2^{(1)}$. To this diagram correspond two different types of surfaces, the generic one corresponds to the multiplicative system of type $A_2^{(1)}$, and the degenerate configuration, where all three components D_i intersect at one point, corresponds to the additive system denoted by $A_2^{(1)*}$, this is our case (on the picture below we use the equivalent description of \mathbf{X} as a blow-up of \mathbb{P}^2 instead of $\mathbb{P}^1 \times \mathbb{P}^1$, since all lines are then equivalent):



Thus, we have showed that an elementary Schlesinger transformation of this Fuchsian system reduces to a discrete Painlevé equation of type d- $P(A_2^{(1)*})$. We now want to compare the Schlesinger dynamic with the standard Painlevé dynamic of this type.


5.2.2. *Comparison with the standard case.* We take as the model equation for the $d-P(A_2^{(1)*})$ dynamic the following equation written by Grammaticos, Ramani, and Ohta, [GRO03], see also Murata [Mur04] and Sakai [Sak07]. We consider $d-P(A_2^{(1)*})$ to be a birational map $\varphi : \mathbb{P}^1 \times \mathbb{P}^1 \dashrightarrow \mathbb{P}^1 \times \mathbb{P}^1$ with parameters b_1, \dots, b_8

$$(5.12) \quad \left(\begin{matrix} b_1 & b_2 & b_3 & b_4 \\ b_5 & b_6 & b_7 & b_8 \end{matrix}; f, g \right) \mapsto \left(\begin{matrix} \bar{b}_1 & \bar{b}_2 & \bar{b}_3 & \bar{b}_4 \\ \bar{b}_5 & \bar{b}_6 & \bar{b}_7 & \bar{b}_8 \end{matrix}; \bar{f}, \bar{g} \right), \quad \begin{matrix} \bar{b}_1 = b_1, & \bar{b}_3 = b_3, & \bar{b}_5 = b_5 + \delta, & \bar{b}_7 = b_7 - \delta, \\ \bar{b}_2 = b_2, & \bar{b}_4 = b_4, & \bar{b}_6 = b_6 + \delta, & \bar{b}_8 = b_8 - \delta, \end{matrix}$$

$\delta = b_1 + \dots + b_8$, and \bar{f} and \bar{g} are given by the equation

$$(5.13) \quad \begin{cases} (f+g)(\bar{f}+g) = \frac{(g+b_1)(g+b_2)(g+b_3)(g+b_4)}{(g-b_5)(g-b_6)} \\ (\bar{f}+g)(\bar{f}+\bar{g}) = \frac{(\bar{f}-b_1)(\bar{f}-b_2)(\bar{f}-b_3)(\bar{f}-b_4)}{(\bar{f}+b_7-\delta)(\bar{f}+b_8-\delta)} \end{cases}$$

Note that equation (5.13) is the asymmetric version of equation (1.3) in [GRO03]. We see that the resulting blow-up diagram matches the one that we obtained for the Schlesinger transformation dynamic.



Thus, we can put $\mathcal{H}_f = H_x$ and $\mathcal{H}_g = H_y$ and let $f = x$, $g = y$. We also put $\mathcal{E}_i = F_i$ and use this to identify Schlesinger parameters with parameters b_i :

$$b_1 = \theta_0^2 + \kappa_1, \quad b_2 = \theta_0^2 + \kappa_2, \quad b_3 = \theta_0^2 + \kappa_3, \quad b_4 = 0, \quad b_5 = \theta_1^1, \quad b_6 = \theta_1^2, \quad b_7 = \theta_0^1 - \theta_0^2, \quad b_8 = -\theta_0^2 - 1.$$

The dynamic itself, however, is different. There are two ways to see that. First, in view of the above identification of parameters we see that the dynamic on the Riemann Schemes is different. Indeed, since $\delta = b_1 + \dots + b_8 = -1$, the induced action of the standard Painlevé dynamic on the Riemann scheme is

$$\left\{ \begin{matrix} z=0 & z=1 & z=\infty \\ \theta_0^1 & \theta_1^1 & \kappa_1 \\ \theta_0^2 & \theta_1^2 & \kappa_2 \\ 0 & 0 & \kappa_3 \end{matrix} \right\} \xrightarrow{d-P(A_2^{(1)*})} \left\{ \begin{matrix} z=0 & z=1 & z=\infty \\ \theta_0^1 & \theta_1^1 - 1 & \kappa_1 + 1 \\ \theta_0^2 - 1 & \theta_1^2 - 1 & \kappa_2 + 1 \\ 0 & 0 & \kappa_3 + 1 \end{matrix} \right\},$$


whereas our elementary Schlesinger transformation acts as

$$\left\{ \begin{matrix} z=0 & z=1 & z=\infty \\ \theta_0^1 & \theta_1^1 & \kappa_1 \\ \theta_0^2 & \theta_1^2 & \kappa_2 \\ 0 & 0 & \kappa_3 \end{matrix} \right\} \xrightarrow{\begin{Bmatrix} 0 & 1 \\ 1 & 1 \end{Bmatrix}} \left\{ \begin{matrix} z=0 & z=1 & z=\infty \\ \theta_0^1 - 1 & \theta_1^1 + 1 & \kappa_1 \\ \theta_0^2 & \theta_1^2 & \kappa_2 \\ 0 & 0 & \kappa_3 \end{matrix} \right\}.$$

Second, we can explicitly compute the induced dynamic for both φ_* and ψ_* on the Picard lattice. Then we find out that the action on the components of the anti-canonical divisor is the same and is a permutation $(D_0 D_1 D_2)$, but the resulting translations on the symmetry sub-lattice are different. Indeed, take the

symmetry sub-lattice $R^\perp = \text{Span}_{\mathbb{Z}}\{\alpha_0, \dots, \alpha_6\}$, where

$$\begin{aligned} \alpha_0 &= E_3 - E_4, & \alpha_1 &= E_2 - E_3, \\ \alpha_2 &= E_1 - E_2, & \alpha_3 &= H_f - E_1 - E_7, \\ \alpha_4 &= E_7 - E_8, & \alpha_5 &= H_g - E_1 - E_5, \\ \alpha_6 &= E_5 - E_6 \end{aligned}$$



which is of the type $E_6^{(1)}$. Then φ_* and ψ_* act on R^\perp as translations in two *different* directions:

$$\begin{aligned} \psi_* : (\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6) &\mapsto (\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6) + (0, 0, 0, -1, 1, 1, -1)\delta, \\ \varphi_* : (\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6) &\mapsto (\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6) + (0, 0, 0, 1, 0, -1, 0)\delta, \end{aligned}$$

where in both cases $\delta = -K_X$. Hence, these two dynamics are *not equivalent* and we can not transform the elementary Schlesinger transformation dynamic into the standard form by a change of variables.

In fact, we can represent the standard Painlevé dynamic as a composition of two different Schlesinger transformations, combined with some automorphisms of our Fuchsian system, as follows.

Let $\sigma_i(j, k)$ be the map that exchanges the j -th and the k -th eigenvectors of \mathbf{A}_i . If the eigenvalues θ_i^j and θ_i^k are non-zero, this map is just a permutation on the decomposition space $\mathcal{B} \times \mathcal{C}$. For this example we need $\sigma_0(1, 3)$, but it also defines a mapping on $\mathcal{B} \times \mathcal{C}$, since $\mathbf{b}_0^3 \in \text{Ker}(\mathbf{C}_0^\dagger)$ and $\mathbf{c}_0^{3\dagger} \in \text{Ker}(\mathbf{B}_0)$. Combining $\sigma_0(1, 3)$ with the scalar gauge transformation $\rho_0(\theta_0^1)$, where $\rho_i(s) : \mathbf{A}_i \mapsto \mathbf{A}_i - s\mathbf{I}$, $\mathbf{A}_\infty \mapsto \mathbf{A}_\infty + s\mathbf{I}$, see **Assumption 2.2**, we get the action $\Sigma_0(1, 3) = \rho_0(\theta_0^1) \circ \sigma_0(1, 3)$ on the spectral data (as encoded in the Riemann scheme) and the decomposition space:

$$\begin{aligned} \Sigma_0(1, 3) : \left\{ \begin{array}{ccc} z_0 & z_1 & \infty \\ \theta_0^1 & \theta_1^1 & \kappa_1 \\ \theta_0^2 & \theta_1^2 & \kappa_2 \\ 0 & 0 & \kappa_3 \end{array} \right\} &\xrightarrow{\sigma_0(1,3)} \left\{ \begin{array}{ccc} z_0 & z_1 & \infty \\ 0 & \theta_1^1 & \kappa_1 \\ \theta_0^2 & \theta_1^2 & \kappa_2 \\ \theta_0^1 & 0 & \kappa_3 \end{array} \right\} \xrightarrow{\rho_0(\theta_0^1)} \left\{ \begin{array}{ccc} z_0 & z_1 & \infty \\ -\theta_0^1 & \theta_1^1 & \kappa_1 + \theta_0^1 \\ \theta_0^2 - \theta_0^1 & \theta_1^2 & \kappa_2 + \theta_0^1 \\ 0 & 0 & \kappa_3 + \theta_0^1 \end{array} \right\} \\ \Sigma_0(1, 3) : (\mathbf{b}_0^1, \mathbf{b}_0^2; \mathbf{c}_0^{1\dagger}, \mathbf{c}_0^{2\dagger}; \mathbf{b}_1^1, \mathbf{b}_1^2; \mathbf{c}_1^{1\dagger}, \mathbf{c}_1^{2\dagger}) &\mapsto \left((\mathbf{c}_0^{1\dagger} \times \mathbf{c}_0^{2\dagger})^t, \mathbf{b}_0^2; \frac{-\theta_0^1(\mathbf{b}_0^1 \times \mathbf{b}_0^2)^t}{(\mathbf{c}_0^{1\dagger} \times \mathbf{c}_0^{2\dagger})(\mathbf{b}_0^1 \times \mathbf{b}_0^2)}, \mathbf{c}_0^{2\dagger}; \mathbf{b}_1^1, \mathbf{b}_1^2; \mathbf{c}_1^{1\dagger}, \mathbf{c}_1^{2\dagger} \right), \end{aligned}$$

where we had to take into account the normalization condition $\mathbf{C}_0^\dagger \mathbf{B}_0 = \Theta_0$. Then, looking at the action on the Riemann scheme, we see that

$$d-P(A_2^{(1)*}) = \Sigma_0(1, 3) \circ \left\{ \begin{array}{cc} 1 & 0 \\ 2 & 1 \end{array} \right\} \circ \Sigma_0(1, 3) \circ \left\{ \begin{array}{cc} 1 & 0 \\ 1 & 1 \end{array} \right\}.$$

Using computer algebra, we can verify directly that this composition dynamic satisfies equation (5.13).

6. CONCLUSIONS

To summarize, we developed the theory of elementary Schlesinger transformations of Fuchsian systems. We found explicit *discrete Schlesinger evolution equations* describing these transformations and showed that these equations can be written in discrete Hamiltonian form, when considered as a dynamical system on the decomposition space for Fuchsian systems. We found the explicit formula for the discrete Hamiltonian for such systems. We also stressed the parallel between the discrete and continuous cases. Finally, we used the machinery developed in the first part of this paper to compute two explicit examples of reductions from discrete Schlesinger transformations to difference Painlevé equations. In doing so we emphasized the role played by the geometry of the Okamoto spaces of initial conditions in doing such reductions, as well as in comparing different equations of the same type or difference forms of the same equation. We also gave a partial answer to the earlier question of H. Sakai by explicitly representing the difference Painlevé equation $d-P(A_2^{(1)*})$ as such reduction.

REFERENCES

- [AB06] D. Arinkin and A. Borodin, *Moduli spaces of d -connections and difference Painlevé equations*, Duke Math. J. **134** (2006), no. 3, 515–556, [arXiv:math/0411584](#) [[math.AG](#)]. MR MR2254625 (2007h:14047)
- [Boa09] Philip Boalch, *Quivers and difference Painlevé equations*, Groups and symmetries, CRM Proc. Lecture Notes, vol. 47, Amer. Math. Soc., Providence, RI, 2009, pp. 25–51, [arXiv:0706.2634](#) [[math.AG](#)]. MR 2500553 (2011g:39028)
- [Bor04] Alexei Borodin, *Isomonodromy transformations of linear systems of difference equations*, Ann. of Math. (2) **160** (2004), no. 3, 1141–1182, [arXiv:math/0209144](#) [[math.CA](#)]. MR MR2144976
- [Cad70] J.A. Cadzow, *Discrete calculus of variations.*, Int. J. Control, I. Ser. **11** (1970), 393–407 (English).
- [CT12] Adrian Stefan Carstea and Tomoyuki Takenawa, *A note on minimization of rational surfaces obtained from birational dynamical systems*, preprint, [arXiv:1211.5393](#) [[math.DS](#)].
- [Dzh09] Anton Dzhamay, *Factorizations of rational matrix functions with application to discrete isomonodromic transformations and difference Painlevé equations*, J. Phys. A: Math. Theor. **42** (2009), no. 45, 10pp., [arXiv:0904.1971](#) [[math-ph](#)].
- [Fuc07] R. Fuchs, *Über lineare homogene Differentialgleichungen zweiter Ordnung mit drei im Endlichen gelegenen wesentlich singulären Stellen*, Math. Ann. **63** (1907), 301–321.
- [Gar12] R. Garnier, *Sur les équations différentielles du troisième ordre dont l'intégrale générale est uniforme et sur une classe d'équations nouvelles d'ordre supérieur dont l'intégrale générale a ses points critiques fixes*, Ann. Sci. École Norm. Sup. **29** (1912), 1–126.
- [Gar26] ———, *Solution du problème de riemann pour les systemes différentiels linéaires du second ordre*, Annales scientifiques de l'É.N.S. 3e série **43** (1926), 177–307.
- [GPS00] Herbert Goldstein, Charles Poole, and John Safko, *Classical mechanics*, Addison-Wesley, 2000.
- [GRO03] B. Grammaticos, A. Ramani, and Y. Ohta, *A unified description of the asymmetric q - P_V and d - P_{IV} equations and their Schlesinger transformations*, J. Nonlinear Math. Phys. **10** (2003), no. 2, 215–228, [arXiv:nlin/0310050](#) [[nlin.SI](#)]. MR 1976382 (2004c:34273)
- [Har94] J. Harnad, *Dual isomonodromic deformations and moment maps to loop algebras*, Comm. Math. Phys. **166** (1994), no. 2, 337–365, [arXiv:hep-th/9301076](#). MR MR1309553 (96e:58073)
- [JMMS80] Michio Jimbo, Tetsuji Miwa, Yasuko Môri, and Mikio Sato, *Density matrix of an impenetrable Bose gas and the fifth Painlevé transcendent*, Phys. D **1** (1980), no. 1, 80–158. MR MR573370 (84k:82037)
- [JM81] Michio Jimbo and Tetsuji Miwa, *Monodromy preserving deformation of linear ordinary differential equations with rational coefficients. II*, Phys. D **2** (1981), no. 3, 407–448. MR MR625446 (83k:34010b)
- [JM82] ———, *Monodromy preserving deformation of linear ordinary differential equations with rational coefficients. III*, Phys. D **4** (1981/82), no. 1, 26–46. MR MR636469 (83k:34010c)
- [JMU81] Michio Jimbo, Tetsuji Miwa, and Kimio Ueno, *Monodromy preserving deformation of linear ordinary differential equations with rational coefficients. I. General theory and τ -function*, Phys. D **2** (1981), no. 2, 306–352. MR MR630674 (83k:34010a)
- [JS96] Michio Jimbo and Hidetaka Sakai, *A q -analog of the sixth Painlevé equation*, Lett. Math. Phys. **38** (1996), no. 2, 145–154, [arXiv:chao-dyn/9507010](#). MR MR1403067 (97m:39002)
- [JP64] B. W. Jordan and E. Polak, *Theory of a class of discrete optimal control systems*, Journal of Electronics and Control **17** (1964), no. 6.
- [Kat96] Nicholas M. Katz, *Rigid local systems*, Annals of Mathematics Studies, vol. 139, Princeton University Press, Princeton, NJ, 1996. MR 1366651 (97e:14027)
- [KNS13] Hiroshi Kawakami, Akane Nakamura, and Hidetaka Sakai, *Toward a classification of four-dimensional Painlevé-type equations*, Algebraic and geometric aspects of integrable systems and random matrices, Contemp. Math., vol. 593, Amer. Math. Soc., Providence, RI, 2013, pp. 143–161. MR 3087954
- [Log73] John David Logan, *First integrals in the discrete variational calculus*, Aequationes Math. **9** (1973), 210–220. MR 0328397 (48 #6739)
- [LW06] S. Lall and M. West, *Discrete variational Hamiltonian mechanics*, J. Phys. A **39** (2006), no. 19, 5509–5519. MR 2220773 (2007h:37080)
- [Mae82] Shigeru Maeda, *Lagrangian formulation of discrete systems and concept of difference space*, Math. Japon. **27** (1982), no. 3, 345–356. MR 663162 (83k:39003)
- [MV91] Jürgen Moser and Alexander P. Veselov, *Discrete versions of some classical integrable systems and factorization of matrix polynomials*, Comm. Math. Phys. **139** (1991), no. 2, 217–243. MR MR1120138 (92g:58054)
- [MW01] J. E. Marsden and M. West, *Discrete mechanics and variational integrators*, Acta Numer. **10** (2001), 357–514. MR MR2009697 (2004h:37130)
- [Mur04] Mikio Murata, *New expressions for discrete Painlevé equations*, Funkcial. Ekvac. **47** (2004), no. 2, 291–305, [arXiv:nlin/0304001](#) [[nlin.SI](#)]. MR 2108677 (2005j:39014)
- [OBL11] Tomoki Ohsawa, Anthony M. Bloch, and Melvin Leok, *Discrete Hamilton-Jacobi theory*, SIAM J. Control Optim. **49** (2011), no. 4, 1829–1856, [arXiv:0911.2258](#) [[math.OC](#)]. MR 2837576 (2012g:70028)
- [Oka79] Kazuo Okamoto, *Sur les feuilletages associés aux équations du second ordre à points critiques fixes de P. Painlevé*, Japan. J. Math. (N.S.) **5** (1979), no. 1, 1–79. MR 614694 (83m:58005)
- [Oka80] Kazuo Okamoto, *Polynomial Hamiltonians associated with Painlevé equations.*, Proc. Japan Acad. Ser. A Math. Sci. **56** (1980), no. 6, 264–268. MR 596006 (82j:34010b)

- [Osh13] Toshio Oshima, *Classification of Fuchsian systems and their connection problem*, Exact WKB Analysis and Microlocal Analysis, RIMS Kôkyûroku Bessatsu, B37, Res. Inst. Math. Sci. (RIMS), Kyoto, 2013, pp. 163–192. [arXiv:0811.2916](#) [[math.CA](#)].
- [PNGR92] V. G. Papageorgiou, F. W. Nijhoff, B. Grammaticos, and A. Ramani, *Isomonodromic deformation problems for discrete analogues of Painlevé equations*, Phys. Lett. A **164** (1992), no. 1, 57–64. MR MR1162062 (93c:58092)
- [RGH91] A. Ramani, B. Grammaticos, and J. Hietarinta, *Discrete versions of the Painlevé equations*, Phys. Rev. Lett. **67** (1991), no. 14, 1829–1832. MR 1125951 (92j:39011)
- [Sak01] Hidetaka Sakai, *Rational surfaces associated with affine root systems and geometry of the Painlevé equations*, Comm. Math. Phys. **220** (2001), no. 1, 165–229. MR MR1882403 (2003c:14030)
- [Sak07] ———, *Problem: discrete Painlevé equations and their Lax forms*, Algebraic, analytic and geometric aspects of complex differential equations and their deformations. Painlevé hierarchies, RIMS Kôkyûroku Bessatsu, B2, Res. Inst. Math. Sci. (RIMS), Kyoto, 2007, pp. 195–208. MR 2310030 (2008c:33020)
- [Sak10] ———, *Isomonodromic deformation and 4-dimensional Painlevé type equations*, Tech. Report 17, Graduate School of Mathematical Sciences, The University of Tokyo, 2010.
- [Sch12] L. Schlesinger, *Über eine Klasse von Differentialsystemen beliebiger Ordnung mit festen kritischen Punkten*, J. Reine Angew. Math. **141** (1912), 96–145.
- [Sur03] Yuri B. Suris, *The problem of integrable discretization: Hamiltonian approach*, Progress in Mathematics, vol. 219, Birkhäuser Verlag, Basel, 2003. MR MR1993935 (2004h:37086)
- [Ves88] A. P. Veselov, *Integrable systems with discrete time, and difference operators*, Funktsional. Anal. i Prilozhen. **22** (1988), no. 2, 1–13, 96. MR MR947601 (90a:58081)
- [Ves91] ———, *Integrable Lagrangian correspondences and factorization of matrix polynomials*, Funktsional. Anal. i Prilozhen. **25** (1991), no. 2, 38–49, 96. MR MR1142207 (93b:58081)

SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF NORTHERN COLORADO, CAMPUS BOX 122, 501 20TH STREET, GREELEY, CO 80639, USA

E-mail address: adzham@unco.edu

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3–8–1 KOMABA MEGURO-KU, TOKYO 153-8914, JAPAN

E-mail address: sakai@ms.u-tokyo.ac.jp

FACULTY OF MARINE TECHNOLOGY, TOKYO UNIVERSITY OF MARINE SCIENCE AND TECHNOLOGY, 2-1-6 ETCHU-JIMA, KOTO-KU, TOKYO, 135-8533, JAPAN

E-mail address: takenawa@kaiyodai.ac.jp