

Exponentially growing finite energy solutions for the Klein-Gordon equation on sub-extremal Kerr spacetimes

Yakov Shlapentokh-Rothman*

September 20, 2018

Abstract

For any sub-extremal Kerr spacetime with non-zero angular momentum, we find an open family of non-zero masses for which there exist smooth, finite energy, and exponentially growing solutions to the corresponding Klein-Gordon equation. If desired, for any non-zero integer m , an exponentially growing solution can be found with mass arbitrarily close to $\frac{|am|}{2Mr_+}$. In addition to its direct relevance for the stability of Kerr as a solution to the Einstein-Klein-Gordon system, our result provides the first rigorous construction of a superradiant instability. Finally, we note that this linear instability for the Klein-Gordon equation contrasts strongly with recent work establishing linear stability for the wave equation.

Contents

1	Introduction	2
1.1	Linear Stability of The Wave Equation	5
1.2	The Geometry of Kerr: The Ergoregion and Superradiance	6
1.2.1	Coordinate Systems	6
1.2.2	The Ergoregion and Superradiance	7
1.3	Precise Statement of Results	8
1.4	Previous Works on the Klein-Gordon Equation	9
1.5	Black-Hole Bombs and The Physics Literature	10
2	Mode Solutions	10

*Department of Mathematics, MIT, Cambridge, MA 02139, USA

3	Proof of Theorem 1.3: Restrictions on Mode Solutions Corresponding to Real ω	12
3.1	Part 1	12
3.2	Part 2	13
3.3	Part 3	14
4	Proof of Theorem 1.2: Construction of Mode Solutions	14
4.1	Outline of Proof	14
4.2	Existence of Real Mode Solutions	15
4.2.1	A Variational Interpretation of Real Mode Solutions	15
4.2.2	Analysis of the Variational Problem	16
4.3	Construction of the Exponentially Growing Modes	20
4.4	Modes Crossing the Real Axis	22
5	Following the Unstable Modes in the Upper Half Plane	26
6	Acknowledgements	26
A	Linear ODE's with Regular Singularities	27
B	The Angular ODE	29
C	Local Theory for the Radial ODE	31
C.1	The Horizon	31
C.2	Infinity	32
C.3	Connection Coefficients	35
D	Distant Bound Keplerian Orbits and Unstable Modes	35

1 Introduction

The Kerr spacetime $(\mathcal{M}, g_{a,M})$ is a two parameter family of asymptotically flat, stationary, and axisymmetric solutions to the vacuum Einstein equations $\text{Ric}(g) = 0$. As a precursor to establishing the conjectured non-linear stability of Kerr, there has been much study of the linear stability problem for various equations on a fixed Kerr background. In this paper we will study the Klein-Gordon equation:

$$(\square_g - \mu^2) \psi = 0.$$

Here $\mu \geq 0$ is the mass of the scalar field ψ . In contrast to previous works on the wave equation ($\mu = 0$) showing linear stability, we will produce an

open family of masses for which the Klein-Gordon equation exhibits linear *instability*, i.e. for these masses there exists smooth, finite energy solutions which grow exponentially in time:

Theorem 1.1. *Fix a Kerr spacetime $(\mathcal{M}, g_{a,M})$ with $a \neq 0$ (see section 1.2). Then there exists an open family of masses μ with $\epsilon_\mu > 0$ and a non-zero, smooth, and finite energy solution ψ to the corresponding Klein-Gordon equation*

$$(\square_g - \mu^2) \psi = 0$$

such that

$$e^{\epsilon_\mu t} |\partial^\alpha \psi(0, r, \theta, \phi)| \lesssim_\alpha |\partial^\alpha \psi(t, r, \theta, \phi)| \text{ for all multi-indices } \alpha.$$

These statements should be understood with respect to Boyer-Lindquist coordinates (see section 1.2). For every non-zero integer m , μ can be chosen arbitrarily close to $\frac{|am|}{2Mr_+}$. In particular, μ can be made arbitrarily small as $a \rightarrow 0$.

This may suggest that the Kerr spacetime is non-linearly *unstable* as a solution to the Einstein-Klein-Gordon system.¹ Additionally, our result provides the first rigorous construction of a superradiant instability. Informally put, superradiance occurs in a black hole spacetime when there does not exist a globally defined Killing vector field which is both timelike or null at infinity and timlike or null on the horizon. For such spacetimes “energy” may radiate out of the black hole and, depending on the particular dynamics under consideration, lead to a superradiant instability.

To make these ideas more concrete, let’s focus on the Klein-Gordon equation and begin by briefly recalling the energy-momentum tensor formalism (see [1] for a proper introduction). Let g denote the (Lorentzian) metric on our spacetime and ∇ denote covariant differentiation. For any function ψ we define the energy-momentum tensor

$$\mathbf{T}_{\alpha\beta} := \text{Re} (\nabla_\alpha \psi \overline{\nabla_\beta \psi}) - \frac{1}{2} g_{\alpha\beta} (|\nabla \psi|^2 + \mu^2 |\psi|^2).$$

¹The Einstein-Klein-Gordon system for a spacetime (\mathcal{M}, g) and massive scalar field ψ is

$$\begin{aligned} \text{Ric}_{\alpha\beta}(g) - \frac{1}{2} R(g) g_{\alpha\beta} &= 8\pi \mathbf{T}_{\alpha\beta}(g, \psi), \\ (\square_g - \mu^2) \psi &= 0. \end{aligned}$$

Here $R(g)$ is the scalar curvature and $\mathbf{T}_{\alpha\beta}$ is the energy-momentum tensor,

$$\mathbf{T}_{\alpha\beta} := \text{Re} (\nabla_\alpha \psi \overline{\nabla_\beta \psi}) - \frac{1}{2} g_{\alpha\beta} (|\nabla \psi|^2 + \mu^2 |\psi|^2).$$

For any vector field X we define a corresponding 1-form, called a “current,” by

$$\mathbf{J}_\alpha^X := \mathbf{T}_{\alpha\beta} X^\beta. \quad (1.1)$$

The key identity is

$$\nabla^\alpha \mathbf{J}_\alpha^X = \text{Re} \left((\nabla^\alpha \nabla_\alpha \psi - \mu^2 \psi) \overline{(X\psi)} \right) + \frac{1}{2} \mathbf{T}_{\alpha\beta} \pi^{\alpha\beta}. \quad (1.2)$$

Here π denotes the deformation tensor of X . This vanishes if and only if X is Killing. In particular, if ψ solves the Klein-Gordon equation and X is Killing, we find that \mathbf{J}_α^X is divergence free. In this case, for any two homologous hypersurfaces Σ_1 and Σ_2 , the divergence theorem gives a conservation law:

$$\int_{\Sigma_1} \mathbf{J}_\alpha^X n_{\Sigma_1}^\alpha = \int_{\Sigma_2} \mathbf{J}_\alpha^X n_{\Sigma_2}^\alpha. \quad (1.3)$$

Here n_{Σ_i} denotes the (future oriented) normal to the hypersurface Σ_i , and the integrals are with respect to the natural volume forms (the ones that make the divergence theorem true). For the identity (1.3) to be useful, we need some positivity of $\mathbf{J}_\alpha^X n_{\Sigma_i}^\alpha$. One may show (see [1]) that $\mathbf{J}_\alpha^X n_{\Sigma_i}^\alpha$ is a positive definite non-degenerate quadratic form in ψ and its derivatives at a point x_0 if and only if X and n_{Σ_i} are timelike at x_0 . If we allow the quadratic form to be degenerate, then we may allow X and n_{Σ_i} to also be null. The significance of superradiance for stability problems should now be clear.

Of course, the canonical example of a spacetime admitting superradiance is Kerr.²³ The geometry of Kerr is reviewed in section 1.2; for now, let us simply recall that there is a unique Killing vector field which is timelike at infinity and that this vector field is spacelike on (almost all of) the horizon. If we let T denote this vector field and Φ denote the unique Killing vector field which vanishes along the axis of symmetry, then the energy density along the horizon \mathcal{H}^+ for a solution ψ to the Klein-Gordon equation is

$$\text{Re} \left(T\psi \overline{\left(T\psi + \frac{a}{2Mr_+} \Phi\psi \right)} \right). \quad (1.4)$$

²Note that any spacetime of the form $(\mathbb{R} \times M, -dt^2 + g_M)$, with (M, g_M) a Riemannian manifold, admits a globally defined timelike Killing vector field corresponding to time translation. Thus, superradiance is a truly Lorentzian phenomenon.

³The next most interesting examples from the instability point of view are perhaps “small” Kerr-Ads black holes.

When the black hole possesses non-zero angular momentum ($a \neq 0$) it is clearly possible for this quantity to be negative, and thus, in principle, energy can radiate out of the black hole.

Our exponentially growing solutions will be superradiant bound states,⁴ i.e. the energy flux will be negative along the horizon, and the solution will spatially decay exponentially fast so that no energy is radiated away to infinity. For such solutions the energy coming out of the black hole cannot escape; this is what provides the mechanism for the exponential growth. For technical reasons we will first construct bound state solutions which have exactly zero energy flux on the horizon. As one expects, these solutions will neither grow nor decay. Then, a perturbation argument will produce the exponentially growing solutions.

1.1 Linear Stability of The Wave Equation

The wave equation is simply the Klein-Gordon equation with $\mu = 0$. It is instructive to contrast our instability result for the Klein-Gordon equation with previous work showing that the wave equation is linearly stable. Due to the difficulties of superradiance and the complicated trapping structure of Kerr, the most basic boundedness and decay statements for the wave equation on the Kerr spacetime remained unresolved until the quite recent [10] (we should also mention the previous [24] and [16]). This breakthrough followed a lot of earlier work restricted to Kerr spacetimes with ($|a| \ll M$), e.g. [7], [9], [8], [2], [18], [26], [27], and [19].

The instability result of this paper serves to emphasize the subtle effects superradiance can have on the linear stability problem and helps to “explain” why even establishing boundedness for the wave equation on Kerr is difficult. In particular, since the Klein-Gordon equation decays faster than the wave equation on Minkowski space, one may have expected that the Klein-Gordon equation would be easier to control. However, as $|a| \rightarrow 0$ (where superradiance is *weaker* and one expects the problem to get *easier*) we have produced exponentially growing and finite energy solutions with arbitrarily small mass. Thus, any argument used for the wave equation must break down for Klein-Gordon equations with arbitrarily small mass. On a more conceptual level, we see that as one passes into the relativistic world, new obstructions to *boundedness*, not just decay, arise in the superradiant bounded-frequency regime.

⁴Recall that it only makes sense to consider bound state solutions when the mass of the scalar field is non-zero.

1.2 The Geometry of Kerr: The Ergoregion and Superradiance

In this section we shall briefly review the relevant aspects of the geometry of Kerr. For a true introduction to the Kerr spacetime we recommend [28] and [21]. The lecture notes [9] provide a good introduction to the interaction between the geometry of Kerr and the behavior of linear waves.

1.2.1 Coordinate Systems

Outside the black hole, the Kerr metric in Boyer-Lindquist coordinates $(t, r, \theta, \phi) \in \mathbb{R} \times (r_+, \infty) \times \mathbb{S}^2$ is given by

$$g = - \left(1 - \frac{2Mr}{\rho^2} \right) dt^2 - \frac{4Mar \sin^2 \theta}{\rho^2} dt d\phi + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2 + \sin^2 \theta \frac{\Pi}{\rho^2} d\phi^2,$$

$$r_{\pm} := M \pm \sqrt{M^2 - a^2},$$

$$\Delta := r^2 - 2Mr + a^2 = (r - r_+)(r - r_-),$$

$$\rho^2 := r^2 + a^2 \cos^2 \theta,$$

$$\Pi := (r^2 + a^2)^2 - a^2 \sin^2 \theta \Delta.$$

There are two free parameters M and a . The first parameter M is the mass of the black hole, and aM is the angular momentum. The 1-parameter family where $a = 0$ is known as the ‘‘Schwarzschild’’ spacetime. Some non-zero mass M of the spacetime will be fixed throughout the paper, so the term ‘‘mass’’ will always refer to μ . For various reasons relating to the global geometry of Kerr, physically relevant Kerr spacetimes must satisfy $|a| \leq M$ [28]. Furthermore, we shall assume that we are sub-extremal, i.e. $|a| < M$. This assumption guarantees that r_{\pm} both exist and are distinct.

Though Boyer-Lindquist coordinates are often convenient, they break down when $r = r_+$. We shall thus introduce another coordinate system. Let us define two functions $\bar{t}(r)$ and $\bar{\phi}(r)$ on (r_+, ∞) up to a constant by

$$\frac{d\bar{t}}{dr} := \frac{r^2 + a^2}{\Delta}$$

$$\frac{d\bar{\phi}}{dr} := \frac{a}{\Delta}.$$

Then we define Kerr-star coordinates (t^*, r, θ, ϕ^*) by

$$t^*(t, r) := t + \bar{t}(r)$$

$$\phi^*(\phi, r) := \phi + \bar{\phi}(r).$$

In these coordinates the metric becomes

$$g = - \left(1 - \frac{2Mr}{\rho^2} \right) (dt^*)^2 - \frac{4Mar \sin^2 \theta}{\rho^2} dt^* d\phi^* + 2dt^* dr + \rho^2 d\theta^2 + \sin^2 \theta \frac{\Pi}{\rho^2} (d\phi^*)^2 - 2a \sin^2 \theta dr d\phi^*.$$

Note that we can now allow $(t^*, r, \theta, \phi^*) \in \mathbb{R} \times (0, \infty) \times \mathbb{S}^2$. The horizon is defined to be the null hypersurface $\{r = r_+\}$. This is the boundary of the black hole. We call the region $\{r > r_+\}$ the “domain of outer communication.” Lastly, we note that in their common domain, ∂_t in Boyer-Lindquist coordinates is equal to ∂_{t^*} in Kerr-star coordinates. A similar statement applies to ∂_ϕ and ∂_{ϕ^*} .

1.2.2 The Ergoregion and Superradiance

On Minkowski space, the Killing vector field $T := \partial_t$ is everywhere timelike. Combining this with the energy-momentum formalism [1] immediately implies that $\|\psi(t)\|_{\dot{H}_x^1}^2 + \mu^2 \|\psi(t)\|_{L_x^2}^2$ is constant in time. As in Minkowski space, in Kerr we set $T := \partial_t$. One finds that T is the unique Killing vector field which is timelike for all large r . Unfortunately, when

$$\Delta - a^2 \sin^2 \theta < 0$$

then T is spacelike. This region is known as the ergoregion. Due to the ergoregion, the conserved quantity associated to T has no definite sign. Even more disturbing than this loss of a “free” boundedness statement is the Penrose process [28],[6],[22]. This is a thought experiment where a test particle exploits the ergoregion to extract energy from the black hole. In the introduction we have already explained how to see this energy extraction at the level of the Klein-Gordon equation (1.4).

It is important to keep in mind that superradiance cannot occur if either a or $\Phi\psi$ vanishes. Lastly, we would like to emphasize that the problems of the ergoregion and superradiance occur for the wave equation. Hence, the presence of these two features alone certainly does not imply linear instability.

1.3 Precise Statement of Results

We will rigorously construct finite energy solutions to the Klein-Gordon equation

$$(\square_g - \mu^2) \psi = 0$$

on sub-extremal Kerr which grow exponentially. These growing solutions will be “mode solutions” of the form

$$\psi(t, r, \theta, \phi) := e^{-i\omega t} e^{im\phi} S_{ml}(\theta) R(r) \text{ with } \omega \in \mathbb{C} \text{ and } m \in \mathbb{Z}. \quad (1.5)$$

Here (t, r, θ, ϕ) are Boyer-Lindquist coordinates where, as is well known, the Klein-Gordon equation separates. The functions S_{ml} and R must satisfy appropriate ordinary differential equations and boundary conditions (see section 2) so that ψ solves the Klein-Gordon equation, extends smoothly to the horizon (where Boyer-Lindquist coordinates break down), and has finite energy. For our mode solution to grow with time, we must have $\text{Im}(\omega) > 0$. Such solutions are called “unstable modes.” We say that these modes “lie in the upper half-plane.” Mode solutions with $\omega \in \mathbb{R}$ will be called “real modes.” We say that these modes “lie on the real axis.” It will be convenient to refer to the tuple (ω, m, l, μ) as the “parameters” of the mode.

We can now state our main result:

Theorem 1.2. *Fix a sub-extremal Kerr spacetime with mass M and angular momentum aM . Let $m \in \mathbb{Z}$ and $\omega_0 \in \mathbb{R}$ satisfy $am - 2Mr_+\omega_0 = 0$ and $am \neq 0$. Then, for each l and sufficiently small $\delta > 0$, there exists $\mu(0) > |\omega_R(0)|$, real analytic $\omega_R(\epsilon)$, and real analytic $\mu(\epsilon)$ such that for every $-\delta < \epsilon < \delta$, there exists a mode solution with parameters $(\omega_R(\epsilon) + i\epsilon, m, l, \mu(\epsilon))$. These unstable modes must all be superradiant*

$$|am| - 2Mr_+ \sqrt{\omega_R^2(\epsilon) + \epsilon^2} > 0.$$

Lastly, the modes lose mass as they become unstable

$$\frac{\partial \mu}{\partial \epsilon}(0) < 0.$$

The construction of the exponentially growing modes is achieved by perturbing modes corresponding to real ω . Thus, before proving Theorem 1.2, we will undertake an analysis of modes corresponding to real ω . For these modes we have two main results. The first is an existence result (already contained in Theorem 1.2). The second shows that the assumptions on the frequency parameters from Theorem 1.2 are necessary.

Theorem 1.3. *Suppose there exists a mode solution with parameters (ω, m, l, μ) such that $\omega \in \mathbb{R}$ and $\mu^2 > \omega^2$. Then the following statements are true.*

1. *We have $am - 2Mr_+\omega = 0$.*
2. *We have $am \neq 0$.*
3. *There exists a function $C(m, l)$ such that $\omega^2 < \mu^2 < \omega^2 + C(m, l)$ and*

$$\lim_{l \rightarrow \infty} C(m, l) = 0.$$

We will close the section with two remarks. First, we note that we can rephrase the condition $am - 2Mr_+\omega = 0$ more geometrically. Let V be a null generator of the horizon, e.g. $T + \frac{a}{2Mr_+}\Phi$. Then

$$am - 2Mr_+\omega = 0 \Leftrightarrow V\psi = 0 \Leftrightarrow \text{No energy flux along the horizon}$$

Thus, our real mode solutions are simply solutions to the Klein-Gordon equation with exactly vanishing energy flux along the horizon.

Second, an appropriately modified version of Theorem 1.3 also holds if $\omega^2 > \mu^2$. Instead of requiring finite energy, one should enforce the outgoing condition

$$R \sim \frac{e^{i\sqrt{\omega^2 - \mu^2}r^*}}{r} \text{ at } \infty.$$

Here r^* is defined up to a constant by

$$\frac{dr^*}{dr} = \frac{r^2 + a^2}{\Delta}.$$

Though we will not pursue this here, one can rule out such solutions by modifying the arguments of [25].

1.4 Previous Works on the Klein-Gordon Equation

To the best of the author's knowledge, there are no previous rigorous constructions of growing solutions for the Klein-Gordon equation. However, there are a few results which rule out growing modes in certain parameter ranges. In [3] Beyer showed that no unstable modes can exist if

$$\mu \geq \frac{|am|}{2Mr_+} \sqrt{1 + \frac{2M}{r_+}}.$$

In [9] it is noted that if a is small, and μ is small relative to a and m , then the techniques developed by Dafermos and Rodnianski in [7] can be used to show that no unstable modes exist. Finally, unstable modes for the wave equation were ruled out in the ground-breaking mode stability work of Whiting [29].

1.5 Black-Hole Bombs and The Physics Literature

Soon after the discovery of superradiant wave scattering [30], the authors of [23] speculated about placing a mirror around a black hole which would reflect superradiant frequencies. They argued that this would create a positive feedback loop and result in a “black-hole bomb.” Naturally, one is led to wonder if this superradiant instability can arise in a more physically natural fashion. A key breakthrough came in 1976 when Damour, Deruelle, and Ruffini observed that a good candidate is the Klein-Gordon equation with non-zero mass [11]. A few years later, Zouros and Eardley [31] and Detweiler [12] developed more involved heuristics, all leading to the same conclusion. In particular, a connection was drawn between unstable modes for the Klein-Gordon equation and the existence of bound Keplerian orbits outside the ergoregion (see appendix D). Furthermore, they gave some approximations for the instability rates. Various extensions/refinements, numerical and otherwise, of these results continue to appear in the physics literature, e.g. [17], [13], and the references therein. Many of the studies of unstable modes in the physics literature rely on the WKB approximation ([12] is an exception). Even if these WKB arguments were made rigorous, they would only become accurate as $l \rightarrow \infty$. Since our techniques are variational, no large parameter is necessary, and we produce a much more complete picture. We also remark that it is expected that small Kerr-Ads black holes should exhibit superradiant instabilities [4], [5].

2 Mode Solutions

Before discussing the proofs of the theorems, we will provide a brief review of mode solutions and the corresponding boundary conditions. As mentioned above, we work in Boyer-Lindquist coordinates and consider solutions of the form

$$\psi(t, r, \theta, \phi) := e^{-i\omega t} e^{im\phi} S_{ml}(\theta) R(r) \text{ with } \omega \in \mathbb{C}.$$

In order for ψ to satisfy the Klein-Gordon equation, S_{ml} and R must satisfy appropriate equations:

$$\frac{1}{\sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{dS_{ml}}{d\theta} \right) - \left(\frac{m^2}{\sin^2 \theta} - a^2 (\omega^2 - \mu^2) \cos^2 \theta \right) S_{ml} + \lambda_{ml} S_{ml} = 0, \quad (2.1)$$

$$\Delta \frac{d}{dr} \left(\Delta \frac{dR}{dr} \right) - V_\mu R = 0, \quad (2.2)$$

$$V_\mu := -(r^2 + a^2)^2 \omega^2 + 4Mamr\omega - a^2 m^2 + \Delta (\lambda_{ml} + a^2 \omega^2 + \mu^2 r^2).$$

We will refer to (2.1) as the angular ODE and (2.2) as the radial ODE. We would like to impose boundary conditions so that ψ extends smoothly to the whole spacetime and has finite energy. First of all, we must have $m \in \mathbb{Z}$. Next, when ω is real, one can show that imposing the condition that $e^{im\phi} S_{ml}(\theta)$ extends to \mathbb{S}^2 leads to a regular Sturm-Liouville problem for S_{ml} with simple eigenvalues. In appendix B we will show that one may embed these eigenvalues into holomorphic curves $\lambda_{ml}(\kappa)$ such that for each $\kappa \in \mathbb{C}$ with sufficiently small imaginary part, the λ_{ml} is an eigenvalue for an angular ODE with $\kappa = a^2(\omega^2 - \mu^2)$. We thus get a discrete set of eigenvalues $\{\lambda_{ml}\}$ for which equation (2.1) has a solution. For real ω we will have $\lim_{l \rightarrow \infty} \lambda_{ml} = \infty$. We will index the S_{ml} in such a way that in the $a = 0$ case, the S_{ml} are simply spherical harmonics with eigenvalues $\lambda_{ml} = l(l+1)$ ($l \geq |m|$). From this it is not hard to see that when ω is real,

$$\lambda_{ml} + a^2 \omega^2 \geq |m|(|m| + 1).$$

We are left with the question of boundary conditions for R . Since Boyer-Lindquist coordinates break down on the horizon, we will need to change to Kerr-star coordinates (section 1.2.1). We now ask if

$$\psi(t^*, r, \phi^*, \theta) = e^{-i\omega(t^* - \bar{t}(r))} e^{im(\phi^* - \bar{\phi}(r))} S_{ml}(\theta) R(r)$$

extends smoothly to $r = r_+$. This will happen if we can write

$$R(r) = e^{-i(\omega \bar{t}(r) - m \bar{\phi}(r))} f(r) \quad (2.3)$$

where f extends smoothly to r_+ . Let's define

$$\xi := \frac{i(am - 2Mr_+ \omega)}{r_+ - r_-}.$$

It is easy to see that (2.3) is equivalent to

$$R(r) = (r - r_+)^{\xi} \rho(r) \tag{2.4}$$

for some smooth function ρ . The local theory in appendix C.1 will show that this gives a one dimensional space of solutions to the radial ODE. Recall that we defined an r^* coordinate up to a constant by

$$\frac{dr^*}{dr} := \frac{r^2 + a^2}{\Delta}.$$

Then, assuming ω is real and equation (2.4) holds, we will have

$$\begin{aligned} \frac{dR}{dr^*} &= \frac{\xi(r_+ - r_-)}{2Mr_+} R + O(r - r_+) \Rightarrow \\ \frac{dR}{dr^*} + i \left(\omega - \frac{am}{2Mr_+} \right) R &= O(r - r_+). \end{aligned} \tag{2.5}$$

Lastly, we will also need ψ to have finite energy. Hence, for large r we must require

$$\int_{r_++1}^{\infty} \left(|R|^2 + \left| \frac{dR}{dr} \right|^2 \right) r^2 dr < \infty. \tag{2.6}$$

3 Proof of Theorem 1.3: Restrictions on Mode Solutions Corresponding to Real ω

We will start with the proof of Theorem 1.3 since it is simpler than and motivates the hypotheses of Theorem 1.2. We have placed some more technical aspects of the argument in the appendix.

3.1 Part 1

Let R be a mode solution with parameters (ω, m, l, μ) such that $\omega \in \mathbb{R}$ and $\mu^2 > \omega^2$. We wish to show that

$$am - 2Mr_+\omega = 0.$$

As reviewed in appendix C, an asymptotic analysis of the radial ODE shows that all solutions are either exponentially growing or exponentially decaying at infinity. Since our mode solution must have finite energy (2.6), we conclude that R is exponentially decaying at infinity.

Next, let's define the energy current,

$$Q_T := \text{Im} \left(\Delta \frac{dR}{dr} \overline{R} \right).$$

The radial ODE implies that

$$\frac{dQ_T}{dr} = 0.$$

Since R must decay exponentially at infinity, we have $Q_T(\infty) = 0$. Hence, using the horizon boundary condition (2.5), we get

$$0 = Q_T(r_+) = (2Mr_+) \text{Im} \left(\frac{dR}{dr^*}(r_+) \overline{R(r_+)} \right) = (am - 2Mr_+\omega) |R(r_+)|^2.$$

Thus, either $am - 2Mr_+\omega = 0$ or $R(r_+) = 0$. However, $R(r_+) = 0$ implies that R is identically 0 (appendix C). We conclude that $am - 2Mr_+\omega = 0$.

3.2 Part 2

Again we let R be a mode solution with parameters (ω, m, l, μ) such that $\omega \in \mathbb{R}$ and $\mu^2 > \omega^2$. From the previous section we know that we must have

$$am - 2Mr_+\omega = 0.$$

We now wish to show that

$$am \neq 0.$$

Using $am - 2Mr_+\omega = 0$, we may write

$$V_\mu = -(r^2 + a^2)^2 \omega^2 + 4M^2 \omega^2 r_+ (2r - r_+) + \Delta (\lambda_{ml} + a^2 \omega^2 + r^2 \mu^2). \quad (3.1)$$

We now argue by contradiction. If $2Mr_+\omega = am = 0$, then

$$V_\mu = \Delta (\lambda_{ml} + r^2 \mu^2) \geq 0.$$

Now consider the function

$$f(r) := \text{Re} \left(\Delta \frac{dR}{dr} \overline{R} \right).$$

Since our mode solution must be exponentially decreasing at infinity, we see that $f(\infty) = 0$. The boundary conditions at the horizon imply that $f(r_+) = 0$. Hence,

$$0 = \int_{r_+}^{\infty} \frac{df}{dr} dr = \int_{r_+}^{\infty} \left(\Delta \left| \frac{dR}{dr} \right|^2 + \frac{V_\mu}{\Delta} |R|^2 \right) dr.$$

This contradicts the non-triviality of R .

3.3 Part 3

We still let R be a mode solution with parameters (ω, m, l, μ) such that $\omega \in \mathbb{R}$ and $\mu^2 > \omega^2$. From the previous two sections we know that

$$am - 2Mr_+\omega = 0,$$

$$am \neq 0.$$

We wish to show that there exists a function $C(m, l)$ such that

$$\omega^2 < \mu^2 < \omega^2 + C(m, l).$$

It is immediately clear from (3.1) that there exists $C(m, l)$ such that

$$\mu^2 > \omega^2 + C(m, l) \Rightarrow V_\mu \geq 0.^5$$

Now the proof concludes exactly as in section 3.2.

4 Proof of Theorem 1.2: Construction of Mode Solutions

Now we will prove Theorem 1.2. As in the previous section, we have placed some more technical aspects of the argument in the appendix.

4.1 Outline of Proof

Before beginning the proof we will give a brief outline. As mentioned in the introduction, we start by constructing real mode solutions. The key technical insight is a variational interpretation of real mode solutions. The variational problem will possess a degeneracy, but this will turn out to be a minor technical problem. Next, we will perturb our real mode solution into the upper complex half-plane by slightly varying ω and μ . This argument relies on observing that mode solutions are in a one to one correspondence with zeros of a certain holomorphic function of ω and μ . Given this, an appropriate application of the implicit function will conclude the argument. Lastly, we analyze how a mode in the upper half-plane can cross the real axis. The upshot will be that as long as we are in a bound state regime ($\mu^2 > \omega^2$), a mode must become superradiant (proposition 4.6) and lose mass (proposition 4.8) as it enters the upper half-plane. Putting everything together will conclude the proof of Theorem 1.2.

⁵The key point being that $am - 2Mr_+\omega = 0 \Rightarrow V_\mu(r_+) = 0$.

4.2 Existence of Real Mode Solutions

We begin with construction of modes corresponding to real ω . In light of Theorem 1.3 we shall set $am - 2Mr_+\omega = 0$, assume $\omega \neq 0$, and be ready to take μ^2 sufficiently close to ω^2 .

4.2.1 A Variational Interpretation of Real Mode Solutions

First, we shall need to review the local theory for the radial ODE. As recalled in appendix C, when $am - 2Mr_+\omega = 0$, a local basis around r_+ of solutions to the radial ODE is given by

$$\{\varphi_1, \log(r - r_+)\varphi_2 + \varphi_3\}$$

where the φ_i are all analytic near r_+ , $\varphi_1(r_+) = 1$, $\varphi_2(r_+) = 1$, and $\varphi_3(r_+) = 0$. Our to be constructed mode solution R should be a non-zero multiple of φ_1 . As we already observed during the proof of Theorem 1.3, the finite energy requirement (2.6) and the local theory from appendix C implies that near infinity R must be exponentially decreasing. It will be useful to further observe that the local analysis shows that if a solution of the radial ODE is not exponentially decreasing, then it is exponentially increasing.

Next, we explore the graph of the potential V_μ . It is not hard to see that for μ^2 sufficiently close to ω^2 , there exists $r_+ < r_A(\mu^2) < r_B(\mu^2) < \infty$ such that V_μ is positive on (r_+, r_A) , negative on (r_A, r_B) , and then positive on (r_B, ∞) . This suggests that we could look for bound states of the radial ODE by minimizing the functional

$$\mathcal{L}_\mu(f) := \int_{r_+}^{\infty} \left(\Delta \left| \frac{df}{dr} \right|^2 + \frac{V_\mu}{\Delta} |f|^2 \right) dr$$

over functions of unit L^2 norm. Note that any solution f of the radial ODE with $\mathcal{L}_\mu(f) < \infty$ will automatically satisfy the correct boundary conditions. This is the crucial way that the $am - 2Mr_+\omega = 0$ assumption enters the construction. The degeneration of the radial ODE at r_+ poses some difficulties for a direct variational analysis of \mathcal{L}_μ . Nevertheless, we will be able to overcome this by working with regularized versions of \mathcal{L}_μ . In section 4.2.2 we will prove the following two propositions.

Proposition 4.1. *For every μ sufficiently close to but larger than ω , there exists a non-zero f_μ satisfying the boundary conditions of a mode solution and $\nu_\mu \leq 0$ such that*

$$\Delta \frac{d}{dr} \left(\Delta \frac{df_\mu}{dr} \right) - V_\mu f_\mu + \nu_\mu \Delta f_\mu = 0.$$

Furthermore, ν_μ can be taken to be increasing in μ^2 .

Proposition 4.2. *There exists μ_0 and corresponding f_{μ_0} such that $\nu_{\mu_0} = 0$.*

The f_{μ_0} is the real mode solution we seek.

We will close the section with a preparatory lemma. Let's fix ω , m , and l which are assumed to satisfy $am - 2Mr_+\omega = 0$ and $\omega \in \mathbb{R} \setminus \{0\}$. Define

$$\mathcal{A} := \{\mu > 0 : \mu^2 > \omega^2 \text{ and } \exists f \in C_0^\infty \text{ with } \mathcal{L}_\mu(f) < 0\}.$$

Lemma 4.3. *Let μ be sufficiently close to but larger than ω . Then we will have*

$$\mu \in \mathcal{A}.$$

Proof. For every fixed f , $\mathcal{L}_\mu(f)$ is continuous in μ . Thus, it is sufficient to find a smooth f with compact support such that

$$\mathcal{L}_\omega(f) < 0.$$

First, we note that near infinity

$$\frac{V_\omega}{\Delta} = -2M\omega^2 r + O(1).$$

Hence, for f supported in (A, ∞) with A large, we write

$$\mathcal{L}_\omega(f) = \int_A^\infty \left((r^2 + O(r)) \left| \frac{df}{dr} \right|^2 - (2M\omega^2 r + O(1)) |f|^2 \right) dr.$$

Since $\omega \neq 0$, if we set f to be equal to $r^{-3/4}$ on a sufficiently large compact set K and 0 outside a slight enlargement of K , it is clear that we can make $\mathcal{L}_\omega(f)$ as negative as we please. \square

We remark that this lemma is the only place where we shall use the $\omega \neq 0$ hypothesis.

4.2.2 Analysis of the Variational Problem

It will be useful to consider the following regularization of \mathcal{L}_μ :

$$\mathcal{L}_\mu^{(\epsilon)}(f) := \int_{r_++\epsilon}^\infty \left(\Delta \left| \frac{df}{dr} \right|^2 + \frac{V_\mu}{\Delta} |f|^2 \right) dr.$$

Lemma 4.4. *If $\mu^2 > \omega^2$, then there exists $f_\mu^{(\epsilon)} \in H_0^1(r_+ + \epsilon, \infty)$ with unit $L^2(r_+ + \epsilon, \infty)$ norm such that $\mathcal{L}_\mu^{(\epsilon)}$ achieves its infimum over $H_0^1(r_+ + \epsilon, \infty)$ functions of unit $L^2(r_+ + \epsilon, \infty)$ norm on $f_\mu^{(\epsilon)}$.*

Proof. If omitted, all integration ranges are over $(r_+ + \epsilon, \infty)$. Since V_μ is increasing in μ^2 , is non-negative near r_+ , and goes to infinity at $r = \infty$, we can find $r_+ < B_0 < B_1$, $C_0 > 0$, and $C_1 > 0$ only depending on a lower bound for μ^2 such that

$$\int \left(\Delta \left| \frac{df}{dr} \right|^2 + C_0 r^2 1_{[B_0, B_1]^c} (\mu^2 - \omega^2) |f|^2 \right) dr \leq C_1 \int_{B_0}^{B_1} |f|^2 dr + L_\mu^{(\epsilon)}(f). \quad (4.1)$$

From this it is clear that

$$\nu_\mu^{(\epsilon)} := \inf \left\{ \mathcal{L}_\mu^{(\epsilon)}(f) : f \in C_c^\infty \text{ and } \|f\|_{L^2} = 1 \right\} > -\infty.$$

Let $\{f_{n,\mu}^{(\epsilon)}\}_{n=1}^\infty$ be a sequence of smooth functions, compactly supported in $(r_+ + \epsilon, \infty)$, with $\|f_{n,\mu}^{(\epsilon)}\|_{L^2} = 1$, such that

$$\mathcal{L}_\mu^{(\epsilon)}(f_{n,\mu}^{(\epsilon)}) \rightarrow \nu_\mu^{(\epsilon)}.$$

The bound (4.1) implies that $\|f_{n,\mu}^{(\epsilon)}\|_{H^1}$ is uniformly bounded. We now apply Rellich compactness to produce a $f_\mu^{(\epsilon)} \in H_0^1$ such that a re-labeled subsequence of $\{f_{n,\mu}^{(\epsilon)}\}$ converges to $f_\mu^{(\epsilon)}$ weakly in H^1 and strongly in L^2 on compact subsets of (r_+, ∞) .

We claim that no mass is lost in the limit, i.e. $\|f_\mu^{(\epsilon)}\|_{L^2} = 1$. Suppose not. Then, for any compact set K , there will exist infinitely many of the $f_{n,\mu}^{(\epsilon)}$'s such that

$$\|f_{n,\mu}^{(\epsilon)}\|_{L^2([r_+, \infty) \setminus K)} \geq \alpha > 0.$$

It is easy to see from (4.1) that this will give a contradiction if K is sufficiently large.

Using the boundedness of weak limits and the strong L^2 convergence, we then get

$$\nu_\mu^{(\epsilon)} \leq \mathcal{L}_\mu^{(\epsilon)}(f_\mu^{(\epsilon)}) \leq \liminf_{n \rightarrow \infty} \mathcal{L}_\mu^{(\epsilon)}(f_{n,\mu}^{(\epsilon)}) = \nu_\mu^{(\epsilon)}.$$

This implies that $\mathcal{L}_\mu^{(\epsilon)}$ achieves its minimum on $f_\mu^{(\epsilon)}$. \square

Now we are ready to prove proposition 4.1.

Proof. First we observe that $\left\{ \nu_\mu^{(\epsilon)} \right\}_{\epsilon > 0}$ is bounded and decreasing in ϵ . Set $\nu_\mu = \lim_{\epsilon \rightarrow 0} \nu_\mu^{(\epsilon)}$. Lemma 4.3 implies that $\mu \in \mathcal{A}$ which in turn implies that $\nu_\mu < 0$. For any interval $K = (r_+ + \frac{1}{n}, n)$ with n large, (4.1) implies that

$$\begin{aligned} \sup_{\epsilon > 0} \left\| f_\mu^{(\epsilon)} \right\|_{H^1(K)} &< \infty, \\ \inf_{\epsilon > 0} \left\| f_\mu^{(\epsilon)} \right\|_{L^2(K)} &> 0. \end{aligned}$$

After an application of Rellich compactness and passing to a subsequence, we may find a non-zero $f_\mu \in H^1$ that is a weak H^1 and strong L^2_{loc} limit of $f_\mu^{(\epsilon)}$.

Using the Euler-Lagrange equations associated to $\mathcal{L}_\mu^{(\epsilon)}$, we find

$$\Delta \frac{d}{dr} \left(\Delta \frac{df_\mu}{dr} \right) - V_\mu f_\mu + \nu_\mu \Delta f_\mu = 0.$$

On any compact subset K of (r_+, ∞) , boundedness of weak limits and the L^2_{loc} convergence of the $f_\mu^{(\epsilon)}$ imply that

$$\begin{aligned} \int_K \left(\Delta \left| \frac{df_\mu}{dr} \right|^2 + \frac{V_\mu}{\Delta} |f_\mu|^2 \right) &\leq \nu_\mu \Rightarrow \\ \int_{r_+}^\infty \Delta \left| \frac{df_\mu}{dr} \right|^2 dr &< \infty. \end{aligned} \tag{4.2}$$

Near r_+ the local theory from appendix C implies that

$$f_\mu = A\varphi_1 + B(\log(r - r_+)\varphi_2 + \varphi_3)$$

for some constants A and B and non-zero analytic functions φ_i . However, if $B \neq 0$, then

$$\int_{r_+}^\infty \Delta \left| \frac{df_\mu}{dr} \right|^2 dr = \infty.$$

Hence, $B = 0$. Near infinity the local theory from appendix C implies that that f_μ is asymptotic to a linear combination of an exponentially growing solution and an exponentially decaying solution. The bound (4.2) clearly implies that f_μ is in fact exponentially decaying. Thus, f_μ satisfies the boundary conditions of a mode solution. \square

Finally, we can prove proposition 4.2.

Proof. First we will show that ν_μ is continuous for $\mu \in \mathcal{A}$. Let us normalize each $f_\mu^{(\epsilon)}$ so that $\|f_\mu^{(\epsilon)}\|_{L^2} = 1$. We have

$$\begin{aligned} \nu_{\mu_1}^{(\epsilon)} &= \mathcal{L}_{\mu_1}^{(\epsilon)}(f_{\mu_1}^{(\epsilon)}) = \mathcal{L}_{\mu_2}^{(\epsilon)}(f_{\mu_1}^{(\epsilon)}) + \int_{r_++\epsilon}^{\infty} \frac{V_{\mu_1} - V_{\mu_2}}{\Delta} |f_{\mu_1}^{(\epsilon)}|^2 dr \geq \\ &\nu_{\mu_2}^{(\epsilon)} - |\mu_1^2 - \mu_2^2| \int_{r_++\epsilon}^{\infty} r^2 |f_{\mu_1}^{(\epsilon)}|^2 dr. \end{aligned}$$

Reversing the roles of μ_1 and μ_2 gives

$$\begin{aligned} |\nu_{\mu_1}^{(\epsilon)} - \nu_{\mu_2}^{(\epsilon)}| &\leq |\mu_1^2 - \mu_2^2| \int_{r_++\epsilon}^{\infty} r^2 \left(|f_{\mu_1}^{(\epsilon)}|^2 + |f_{\mu_2}^{(\epsilon)}|^2 \right) dr \leq \\ &C |\mu_1^2 - \mu_2^2|. \end{aligned}$$

We have used (4.1) in the last inequality. Since the constant C is independent of ϵ , we may take ϵ to 0.

Since $\mathcal{A} \neq \emptyset$, by lemma 4.3 we may set

$$\mu_0 := \sup \mathcal{A}.$$

It is clear that for any $\mu \in \mathcal{A}$, we cannot have V_μ strictly positive on (r_+, ∞) . Thus $\mu_0 < \infty$. Since ν_μ is increasing in μ , we may extend ν_μ continuously so that ν_{μ_0} exists. We will of course have $\nu_{\mu_0} \leq 0$. Suppose that $\nu_{\mu_0} < 0$. Then, by continuity we could slightly increase μ_0 to μ'_0 with $\nu_{\mu'_0} < 0$. This of course contradicts the definition of μ_0 . We conclude that $\nu_{\mu_0} = 0$. It remains to show that there exists a corresponding f_{μ_0} . From the local theory in appendix C, for every μ and ν , we have a unique solution $\tilde{R}(r, \mu, \nu)$ to

$$\Delta \frac{d}{dr} \left(\Delta \frac{d\tilde{R}}{dr} \right) - V_\mu \tilde{R} + \nu \Delta \tilde{R} = 0$$

which satisfies $\tilde{R}(r_+, \mu, \nu) = 1$. At infinity there will be a local basis of solutions spanned by $\tilde{\rho}_1(r, \mu, \nu)$ and $\tilde{\rho}_2(r, \mu, \nu)$ where $\tilde{\rho}_1$ is exponentially increasing, $\tilde{\rho}_2$ is exponentially decreasing, and both depend analytically on r , μ , and ν . Lastly, we have analytic connection coefficients $\tilde{A}(\mu, \nu)$ and $\tilde{B}(\mu, \nu)$ defined by

$$\tilde{R}(r, \mu, \nu) = \tilde{A}(\mu, \nu) \tilde{\rho}_1(r, \mu, \nu) + \tilde{B}(\mu, \nu) \tilde{\rho}_2(r, \mu, \nu).$$

We may pick $\mu_i \uparrow \mu_0$ such that $\tilde{A}(\mu_i, \nu_{\mu_i}) = 0$. It follows that $\tilde{A}(\mu_0, 0) = 0$. We may then set

$$f_{\mu_0}(r) := \tilde{R}(r, \mu_0, 0).$$

□

4.3 Construction of the Exponentially Growing Modes

In this section our goal is to perturb the real modes into the complex upper half-plane with an appropriate application of the implicit function theorem. Using the previous section we may start with a real mode R with frequency parameters $(\omega_R(0), m, l, \mu(0))$ such that $\omega_R(0) \in \mathbb{R}$ and $\mu^2(0) > \omega_R^2(0)$. For any $\omega = \omega_R + i\omega_I$ and μ sufficiently close to $\omega_R(0)$ and $\mu(0)$ respectively, the local theory from appendix C will give us two linearly independent solutions to the radial ODE $\rho_1(r, \omega, \mu)$ and $\rho_2(r, \omega, \mu)$ such that ρ_1 is exponentially increasing at infinity, ρ_2 is exponentially decreasing at infinity, and both depend holomorphically on ω and analytically on μ . Furthermore, the local theory around r_+ tells us that, up to normalizing properly, for each $\omega = \omega_R + i\omega_I$ and μ we have a unique local solution $R(r, \omega, \mu)$ around r_+ satisfying the boundary conditions of a mode solution. We have

$$R(r, \omega, \mu) = A(\omega, \mu)\rho_1(r, \omega, \mu) + B(\omega, \mu)\rho_2(r, \omega, \mu). \quad (4.3)$$

As shown in appendix C, A and B are holomorphic in ω and μ . Finding a mode solution is equivalent to finding a zero of A . We have picked our parameters so that $A(\omega_R(0), \mu(0)) = 0$. Let's write $A = A_R + iA_I$. Next, we note that an application of the implicit function theorem will produce our unstable modes if we can establish

$$\det \begin{pmatrix} \frac{\partial A_R}{\partial \omega_R} & \frac{\partial A_R}{\partial \mu} \\ \frac{\partial A_I}{\partial \omega_R} & \frac{\partial A_I}{\partial \mu} \end{pmatrix} (\omega_R(0), \mu(0)) \neq 0.$$

In order to do this, we shall return to the the energy current

$$Q_T = \text{Im} \left(\Delta \frac{dR}{dr} \overline{R} \right).$$

Recall that in section 3 we saw

$$\frac{dQ_T}{dr} = 0,$$

$$Q_T(r_+) = am - 2Mr_+\omega_R.$$

We have used the normalization $|R(r_+)|^2 = 1$ in the second statement. Next, let's write Q_T in terms of ρ_1 and ρ_2 .

$$Q_T = |A|^2 \Delta \text{Im} \left(\frac{d\rho_1}{dr} \overline{\rho_1} \right) + \Delta \text{Im} \left(A \frac{d\rho_1}{dr} \overline{B\rho_2} \right) + \Delta \text{Im} \left(B \frac{d\rho_2}{dr} \overline{A\rho_1} \right) + |B|^2 \Delta \text{Im} \left(\frac{d\rho_2}{dr} \overline{\rho_2} \right).$$

Before examining this at infinity, let us note the precise asymptotics of the ρ_i as recalled in appendix C.

$$\rho_1 \sim e^{\sqrt{\mu^2 - \omega_R^2} r}^{-1 + \frac{M(2\omega_R^2 - \mu^2)}{\sqrt{\mu^2 - \omega_R^2}}},$$

$$\rho_2 \sim e^{-\sqrt{\mu^2 - \omega_R^2} r}^{-1 - \frac{M(2\omega_R^2 - \mu^2)}{\sqrt{\mu^2 - \omega_R^2}}}.$$

Furthermore, it's easy to see from the construction of the ρ_i that they are both real valued. Now let's compute $Q_T(\infty)$. Since the ρ_i are real, the first and last terms clearly vanish. The exponential powers cancel in the middle terms, and we find

$$Q_T(\infty) = \sqrt{\mu^2 - \omega_R^2} \text{Im} (A\overline{B}) - \sqrt{\mu^2 - \omega_R^2} \text{Im} (B\overline{A}) = 2\sqrt{\mu^2 - \omega_R^2} \text{Im} (A\overline{B}).$$

We conclude that

$$am - 2Mr_+ \omega_R = 2\sqrt{\mu^2 - \omega_R^2} \text{Im} (A\overline{B}). \quad (4.4)$$

Since $A(\omega_R(0), \mu(0)) = 0$, taking derivatives of (4.4) implies that

$$-2Mr_+ = 2\sqrt{\mu^2(0) - \omega_R^2(0)} \text{Im} \left(\frac{\partial A}{\partial \omega_R} (\omega_R(0), \mu(0)) \overline{B} (\omega_R(0), \mu(0)) \right),$$

$$0 = 2\sqrt{\mu^2(0) - \omega_R^2(0)} \text{Im} \left(\frac{\partial A}{\partial \mu} (\omega_R(0), \mu(0)) \overline{B} (\omega_R(0), \mu(0)) \right).$$

Since $B(\omega_R(0), \mu(0)) \neq 0$, these two equations imply that $\frac{\partial A}{\partial \omega_R}$ and $\frac{\partial A}{\partial \mu}$ are linearly independent at $(\omega_R(0), \mu(0))$ if and only if $\frac{\partial A}{\partial \mu} (\omega_R(0), \mu(0)) \neq 0$, i.e.

$$\det \begin{pmatrix} \frac{\partial A_R}{\partial \omega_R} & \frac{\partial A_R}{\partial \mu} \\ \frac{\partial A_I}{\partial \omega_R} & \frac{\partial A_I}{\partial \mu} \end{pmatrix} (\omega_R(0), \mu(0)) \neq 0 \Leftrightarrow \frac{\partial A}{\partial \mu} (\omega_R(0), \mu(0)) \neq 0.$$

It remains to establish

Lemma 4.5.

$$\frac{\partial A}{\partial \mu} \neq 0.$$

Proof. For the sake of contradiction, suppose that $\frac{\partial A}{\partial \mu}(\omega_R(0), \mu(0)) = 0$. Differentiating (4.3) gives

$$\begin{aligned} \frac{\partial R}{\partial \mu}(r, \omega_R(0), \mu(0)) &= \frac{\partial B}{\partial \mu}(\omega_R(0), \mu(0)) \rho_2(r, \omega_R(0), \mu(0)) + \\ &B(\omega_R(0), \mu(0)) \frac{\partial \rho_2}{\partial \mu}(r, \omega_R(0), \mu(0)). \end{aligned}$$

This implies that $\frac{\partial R}{\partial \mu}$ is exponentially decreasing at infinity. Differentiating the radial ODE with respect to μ , multiplying by \bar{R} , and integrating gives

$$\int_{r_+}^{\infty} \left(\frac{\partial}{\partial r} \left(\Delta \frac{\partial^2 R}{\partial r \partial \mu} \right) - \frac{V_\mu}{\Delta} \frac{\partial R}{\partial \mu} \right) \bar{R} dr = \int_{r_+}^{\infty} \left(2\mu r^2 + \frac{\partial \lambda}{\partial \mu} \right) |R|^2 dr.$$

Integrating by parts twice on the left hand side will produce no boundary terms since both R and $\frac{\partial R}{\partial \mu}$ are exponentially decreasing at infinity. Taking real parts then gives

$$\int_{r_+}^{\infty} \left(2\mu r^2 + \frac{\partial \lambda}{\partial \mu} \right) |R|^2 dr = 0 \Rightarrow R = 0.$$

Proposition B.3 from the appendix says that

$$\frac{\partial \lambda}{\partial \mu} > 0.$$

This contradicts the non-triviality of R . □

4.4 Modes Crossing the Real Axis

In this section we shall investigate how a mode can “cross” the real axis. Let’s introduce a little more notation. From the analysis of the previous section we have a family of mode solutions $R(r, \epsilon)$ with parameters $(\omega(\epsilon), m, l, \mu(\epsilon))$ where $\omega(\epsilon) = \omega_R(\epsilon) + i\epsilon$. Implicitly we have also been using the existence of a family $\lambda_{ml}(\epsilon)$ of eigenvalues to the angular ODE (see proposition B.1). These functions are all defined for $|\epsilon| \ll 1$. In what follows we will often omit the ϵ ’s and we shall assume $0 < \omega_R(\epsilon) < \mu(\epsilon)$. Using the

symmetry of the equations under $(\omega, m) \mapsto (-\omega, -m)$ one may check that this assumption implies no loss of generality. The function R will satisfy

$$\frac{\partial}{\partial r} \left(\Delta \frac{\partial R}{\partial r} \right) - \frac{V_\mu}{\Delta} R = 0,$$

$$R \sim (r - r_+)^{\frac{i(am - 2Mr_+ \omega)}{r_+ - r_-}} \text{ at } r_+,$$

$$R \sim e^{-r\sqrt{\mu^2 - \omega^2}} r^{-1 - \frac{M(2\omega^2 - \mu^2)}{\sqrt{\mu^2 - \omega^2}}} \text{ at } r = \infty,$$

$$V_\mu := -(r^2 + a^2)^2 \omega^2 + 4Mamr\omega - a^2 m^2 + \Delta (\lambda + a^2 \omega^2 + \mu^2 r^2).$$

We also have

$$\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial S}{\partial \theta} \right) - \left(\frac{m^2}{\sin^2 \theta} - a^2 (\omega^2 - \mu^2) \cos^2 \theta \right) S + \lambda S = 0$$

where $S(\cdot, \epsilon) : \theta \in (0, \pi) \rightarrow \mathbb{C}$ is given boundary conditions which make it regular at $\theta = 0, \pi$ (see appendix B). Note that we have suppressed the m and l indices from S_{ml} and λ_{ml} .

From Theorem 1.3 we know that

$$\omega_R(0) = \frac{am}{2Mr_+}. \quad (4.5)$$

We wish to investigate the signs of $\frac{\partial \omega_R}{\partial \epsilon}(0)$ and $\frac{\partial \mu}{\partial \epsilon}(0)$. The condition 4.5 corresponds to our mode solution being exactly on the threshold of superradiance. This makes sense because when $\epsilon = 0$ the solution neither grows nor decays with time. For $\epsilon > 0$ the mode solution will grow with time. Hence, we expect the mode to become superradiant:

$$\omega_R(\epsilon) < \frac{am}{2Mr_+}. \quad (4.6)$$

This leads us to

Proposition 4.6. *If $\epsilon > 0$ we must have*

$$\omega_R^2(\epsilon) + \epsilon^2 < \left(\frac{am}{2Mr_+} \right)^2.$$

In particular

$$\frac{\partial \omega_R}{\partial \epsilon}(0) \leq 0.$$

Proof. Once again we return to (a slight variant of) the microlocal energy current:

$$\tilde{Q}_T := \text{Im} \left(\Delta \frac{\partial R}{\partial r} \overline{\omega R} \right).$$

For $\epsilon > 0$ we have

$$\begin{aligned} \tilde{Q}_T(\infty) &= \tilde{Q}_T(r_+) = 0 \\ \frac{\partial \tilde{Q}_T}{\partial r} &= -\epsilon \Delta \left| \frac{\partial R}{\partial r} \right|^2 + \text{Im} \left(\frac{V_\mu \overline{\omega}}{\Delta} \right) |R|^2 \Rightarrow \\ \int_{r_+}^{\infty} \left(\epsilon \Delta \left| \frac{\partial R}{\partial r} \right|^2 - \text{Im} \left(\frac{V_\mu \overline{\omega}}{\Delta} \right) |R|^2 \right) dr &= 0. \end{aligned}$$

We have

$$\text{Im}(-V_\mu \overline{\omega}) = \epsilon \left((r^2 + a^2)^2 |\omega|^2 - a^2 m^2 + \Delta r^2 \mu^2 \right) - \Delta \text{Im}((\lambda + a^2 \omega^2) \overline{\omega}).$$

Proposition B.2 from the appendix gives

$$-\text{Im}((\lambda + a^2 \omega^2) \overline{\omega}) > 0.$$

Furthermore, $\text{Im}(-V_\mu \overline{\omega})$ is increasing in r . Thus, $R \neq 0$ implies that

$$\begin{aligned} \text{Im}(-V_\mu \overline{\omega})(r_+) &< 0 \Leftrightarrow \\ \omega_R^2(\epsilon) + \epsilon^2 &< \left(\frac{am}{2Mr_+} \right)^2. \end{aligned}$$

□

Proposition 4.7.

$$\frac{\partial \mu}{\partial \epsilon}(0) \geq 0 \Rightarrow \text{Re} \left(\frac{\partial \lambda}{\partial \epsilon} \right)(0) \geq 0.$$

Proof. Let's set

$$S_\epsilon := \frac{\partial S}{\partial \epsilon}.$$

At $\epsilon = 0$ we have

$$\begin{aligned} \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial S_\epsilon}{\partial \theta} \right) - \left(\frac{m^2}{\sin^2 \theta} - a^2 (\omega^2 - \mu^2) \cos^2 \theta \right) S_\epsilon + \lambda S_\epsilon = \\ - \left(2a^2 \cos^2 \theta \left(\omega_R \left(\frac{\partial \omega_R}{\partial \epsilon} + i \right) - \mu \frac{\partial \mu}{\partial \epsilon} \right) + \frac{\partial \lambda}{\partial \epsilon} \right) S. \end{aligned}$$

One may check that S_ϵ is regular at $\theta = 0, \pi$. Multiplying by \overline{S} and integrating by parts implies

$$\int_0^\pi \left(2a^2 \cos^2 \theta \left(\omega_R \left(\frac{\partial \omega_R}{\partial \epsilon} + i \right) - \mu \frac{\partial \mu}{\partial \epsilon} \right) + \frac{\partial \lambda}{\partial \epsilon} \right) |S|^2 \sin \theta d\theta = 0.$$

Using proposition 4.6 we conclude that

$$\frac{\partial \mu}{\partial \epsilon}(0) \geq 0 \Rightarrow \operatorname{Re} \left(\frac{\partial \lambda}{\partial \epsilon} \right)(0) \geq 0.$$

□

Finally, we examine $\frac{\partial \mu}{\partial \epsilon}(0)$.

Proposition 4.8.

$$\frac{\partial \mu}{\partial \epsilon}(0) < 0.$$

Proof. Let's set

$$R_\epsilon := \frac{\partial R}{\partial \epsilon}.$$

We have

$$\frac{\partial}{\partial r} \left(\Delta \frac{\partial R_\epsilon}{\partial r} \right) - \frac{V_\mu}{\Delta} R_\epsilon = \frac{\partial}{\partial \epsilon} \left(\frac{V_\mu}{\Delta} \right) R. \quad (4.7)$$

Now we want to multiply by \overline{R} and integrate by parts. However, we have to be careful with regards to R'_ϵ 's boundary conditions. At infinity R_ϵ may easily be seen to be exponentially decreasing, but at r_+ the proper condition is more subtle. By construction

$$(r - r_+) \frac{-i(am - 2Mr_+ \omega(\epsilon))}{r_+ - r_-} R(r, \epsilon) =: G(r, \epsilon)$$

is analytic in r and ϵ near $(r_+, 0)$. At $\epsilon = 0$ we have

$$\begin{aligned} & (r - r_+) \frac{-i(am - 2Mr_+ \omega)}{r_+ - r_-} R_\epsilon - \\ & \frac{2Mr_+}{r_+ - r_-} \left(1 - i \frac{\partial \omega_R}{\partial \epsilon} \right) (r - r_+) \frac{-i(am - 2Mr_+ \omega)}{r_+ - r_-} \log(r - r_+) R = \frac{\partial G}{\partial \epsilon} \Rightarrow \\ & R_\epsilon(r, 0) = \frac{2Mr_+}{r_+ - r_-} \left(1 - i \frac{\partial \omega_R}{\partial \epsilon} \right) \log(r - r_+) R(r, 0) + \frac{\partial G}{\partial \epsilon}(r, 0). \end{aligned}$$

Now we multiply (4.7) by \overline{R} , take the real part, and integrate by parts. We end up with

$$\begin{aligned}
-2Mr_+ |R(r_+)|^2 &= \int_{r_+}^{\infty} \operatorname{Re} \left(\frac{\partial}{\partial \epsilon} \left(\frac{V_\mu}{\Delta} \right) \right) |R|^2 dr = \\
\int_{r_+}^{\infty} \Delta^{-1} \left(2\omega_R \frac{\partial \omega_R}{\partial \epsilon} (-\Delta^2 + (a^2 - 4Mr)\Delta - 4M^2 r(r - r_+)) \right) |R|^2 dr + \\
\int_{r_+}^{\infty} \left(\operatorname{Re} \left(\frac{\partial \lambda}{\partial \epsilon} \right) + 2r^2 \mu \frac{\partial \mu}{\partial \epsilon} \right) |R|^2 dr.
\end{aligned}$$

Now proposition 4.7 finishes the proof. □

5 Following the Unstable Modes in the Upper Half Plane

Following our construction of unstable modes near the real axis, it is natural to ask if one can continue to decrease μ and produce more unstable modes. We will not explore this in detail in this paper, but we will briefly describe the expected behavior. One believes that one can vary μ and produce a 1-parameter (at least continuous) family of modes with frequency parameters $(\omega(\mu), \mu, m, l)$.⁶ As long as these modes are in the upper half plane, proposition 4.6 shows that they will remain superradiant. Hence, if and when they cross the real axis, they will satisfy $|\omega| \leq \frac{|am|}{2Mr_+}$. Now, note that proposition 4.8 implies that in the bound state regime ($\mu^2 > \omega^2$), an unstable mode can cross the real axis only by increasing the mass. Hence, as long as we decrease μ and maintain $\mu > \frac{|am|}{2Mr_+}$, the curve of modes cannot cross the real axis, and, by continuity, we conclude that these modes would have to remain unstable.

6 Acknowledgements

I would like to thank Igor Rodnianski, Mihalis Dafermos, Gustav Holzegel, and Jonathan Luk for useful conversations and for advice on the exposition.

⁶A potential approach is to more directly exploit the underlying analyticity, see [24] and [16] for ideas along these lines.

A Linear ODE's with Regular Singularities

Let's recall some facts about linear ODEs in the complex plane.

Lemma A.1. *Consider the complex ODE*

$$\frac{d^2 H}{dz^2} + f(z, \lambda) \frac{dH}{dz} + g(z, \lambda) H = 0. \quad (\text{A.1})$$

We will assume that there exists $\{f_j(\lambda)\}$, $\{g_j(\lambda)\}$, $\{F_j^{(K)}\}$, $\{G_j^{(K)}\}$, r , and open $U \subset \mathbb{C}$ such that in $B_r(z_0)$ and for every compact $K \subset U$

$$|f_j(\lambda)| \leq F_j^{(K)} \quad \text{and} \quad |g_j(\lambda)| \leq G_j^{(K)} \quad \text{when } \lambda \in K,$$

$$\sum_{j=0}^{\infty} G_j^{(K)} (z - z_0)^j \quad \text{and} \quad \sum_{j=0}^{\infty} F_j^{(K)} (z - z_0)^j \quad \text{converge absolutely,}$$

$$\{f_j(\lambda)\} \quad \text{and} \quad \{g_j(\lambda)\} \quad \text{are holomorphic in } \lambda \in U,$$

$$(z - z_0)f(z, \lambda) = \sum_{j=0}^{\infty} f_j(\lambda)(z - z_0)^j \quad \text{and} \quad (z - z_0)^2 g(z) = \sum_{j=0}^{\infty} g_j(\lambda)(z - z_0)^j.$$

If these hypotheses hold we say that z_0 is a regular singularity. Set

$$Q(\alpha, \lambda) := \alpha(\alpha - 1) + f_0(\lambda)\alpha + g_0(\lambda).$$

The indicial equation is

$$Q(\alpha, \lambda) = 0.$$

We suppose that a holomorphic $\alpha(\lambda)$ has been chosen such that

$$Q(\alpha(\lambda), \lambda) = 0 \quad \text{and} \quad \min_{j \in \mathbb{Z}^+} |Q(\alpha(\lambda) + j)| = A(\lambda) > 0.$$

Then there exists a unique solution to (A.1) of the form

$$h(z, \lambda) = (z - z_0)^{\alpha(\lambda)} \rho(z, \lambda)$$

such that $\rho(z_0, \lambda) = 1$. Furthermore, ρ is holomorphic for $z \in B_r(z_0)$ and $\lambda \in U$.

Proof. One can extract a proof of this from the discussion of regular singularities in [20]. For the sake of completeness we will give the needed slight

extension here. Without loss of generality we may set $z_0 = 0$. We begin by looking for a formal solution of the form

$$h(z, \lambda) = z^{\alpha(\lambda)} \sum_{j=0}^{\infty} \rho_j(\lambda) z^j$$

where we set $\rho_0(\lambda) = 1$. Formally plugging this into (A.1) we find (see [20])

$$Q((\alpha(\lambda), \lambda)) = 0,$$

$$Q(\alpha(\lambda) + j, \lambda) \rho_j(\lambda) = - \sum_{k=0}^{j-1} ((\alpha(\lambda) + k) f_{j-k}(\lambda) + g_{j-k}(\lambda)) \rho_k(\lambda) \text{ for } j \geq 1.$$

Since $Q(\alpha(\lambda), \lambda) = 0$ by hypothesis, the first equation is satisfied. Furthermore, by assumption $Q(\alpha(\lambda) + j) \neq 0$ for any j . Hence, the second equation determines $\rho_j(\lambda)$ recursively. This establishes the uniqueness of ρ . It remains to check that the series converges appropriately. We will do this by majorizing the series. Let us pick an arbitrary compact set $K \subset U$ and $r_0 < r$. After applying Cauchy's estimate to the holomorphic functions $\sum_j F_j^{(K)} z^j$ and $\sum_j G_j^{(K)} z^j$, we may find a constant C_K so that

$$|f_j(\lambda)| \leq C_K r_0^{-j} \text{ and } |g_j(\lambda)| \leq C_K r_0^{-j} \text{ for } \lambda \in K.$$

Let $\beta(\lambda)$ be the other root of $Q(\cdot, \lambda)$, and set $n(\lambda) := |\alpha(\lambda) - \beta(\lambda)|$. Since $Q(\alpha(\lambda) + k, \lambda) = k(k + \alpha(\lambda) - \beta(\lambda))$, our hypotheses imply that $\alpha(\lambda) - \beta(\lambda) \notin \mathbb{Z}_{\leq 0}$. Next, define $b_j(\lambda)$ by

$$b_j(\lambda) = |\rho_j(\lambda)| \text{ for } j \leq n,$$

$$j(j - |\alpha(\lambda) - \beta(\lambda)|) b_j(\lambda) = C_K \sum_{k=0}^{j-1} (|\alpha(\lambda)| + k + 1) b_k(\lambda) r_0^{k-j} \text{ for } j > n.$$

It is easy to check by induction that $|\rho_j(\lambda)| \leq b_j(\lambda)$ for all j . For sufficiently large j , one finds that

$$r_0 j(j - |\alpha(\lambda) - \beta(\lambda)|) b_j(\lambda) - (j - 1)(j - 1 - |\alpha(\lambda) - \beta(\lambda)|) b_{j-1}(\lambda) =$$

$$C_K (|\alpha(\lambda)| + j) b_{j-1}(\lambda).$$

Now the ratio test implies that the series $\sum_{j \geq 0} b_j(\lambda) z^j$ converges in the ball of radius r_0 . Hence, by the comparison test, $\sum_{j=0}^{\infty} \rho_j(\lambda) z^j$ converges in the same ball. Since r_0 was arbitrary, we find that for every $\lambda \in K$, $\sum_{j=0}^{\infty} \rho_j(\lambda) z^j$ converges and is holomorphic in $z \in B_r(0)$. Next we may freeze $z \in B_r(0)$ and consider $\rho(z, \lambda) = \sum_{j=0}^{\infty} \rho_j(\lambda) z^j$ as a function of λ . For every compact $K \subset U$, our proof has shown that $\rho(z, \cdot)$ is a uniform limit of holomorphic functions. Hence, $\rho(z, \lambda)$ is holomorphic for $\lambda \in U$. \square

B The Angular ODE

In this section we will establish the needed facts about the eigenvalues of the angular ODE. We assume throughout this section that $m \neq 0$.

Set $\kappa := a^2(\omega^2 - \mu^2)$. Then the angular ODE is

$$\frac{1}{\sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{dS}{d\theta} \right) - \left(\frac{m^2}{\sin^2 \theta} - \kappa \cos^2 \theta \right) S + \lambda S = 0.$$

We have suppressed the m and l indices.

Proposition B.1. *Suppose that for some fixed $\kappa_0 \in \mathbb{R}$ we have an eigenvalue λ_0 . Then, for κ sufficiently close to κ_0 , we can uniquely find a holomorphic curve $\lambda(\kappa)$ of eigenvalues for the angular ODE with parameter κ such that $\lambda_0 = \lambda(\kappa_0)$.*

Proof. Let's change variables to $x := \cos \theta$. Then the angular ODE becomes

$$\frac{d}{dx} \left((1-x^2) \frac{dS}{dx} \right) - \left(\frac{m^2}{1-x^2} - \kappa x^2 \right) S + \lambda S = 0 \text{ with } x \in (-1, 1).$$

An asymptotic analysis (appendix A) at $x = \pm 1$ shows that any solution must be asymptotic to a linear combination of $(1 \mp x)^{|m|}$ and $(1 \mp x)^{-|m|}$ as $x \rightarrow \pm 1$. If S is an eigenfunction we clearly must have

$$S \sim (1 \pm x)^{|m|} \text{ as } x \rightarrow \pm 1.$$

For any κ and λ we can uniquely define a solution $S(\theta, \kappa, \lambda)$ by requiring that

$$\begin{aligned} S(\theta, \kappa, \lambda) &\sim (1+x)^{|m|} \text{ as } x \rightarrow -1 & \text{(B.1)} \\ \left(S(\cdot, \kappa, \lambda)(1+\cdot)^{-|m|} \right) (x=-1) &= 1. \end{aligned}$$

We then have holomorphic functions $F(\kappa, \lambda)$ and $G(\kappa, \lambda)$ such that

$$S(\theta, \kappa, \lambda) \sim F(\kappa, \lambda)(1-x)^{-|m|} + G(\kappa, \lambda)(1-x)^{|m|} \text{ as } x \rightarrow 1.$$

Since λ_0 is an eigenvalue, we have $F(\kappa_0, \lambda_0) = 0$. We will be able to uniquely define our curve $\lambda(\kappa)$ for κ near κ_0 via an application of the implicit function theorem if we can verify that

$$\frac{\partial F}{\partial \lambda}(\kappa_0, \lambda_0) \neq 0.$$

For the sake of contradiction, assume that

$$\frac{\partial F}{\partial \lambda}(\kappa_0, \lambda_0) = 0.$$

Set

$$S_\lambda := \frac{\partial S}{\partial \lambda}.$$

By differentiating (B.1) and using that F and $\frac{\partial F}{\partial \lambda}$ vanish at (κ_0, λ_0) , one may easily check that S_λ still satisfies the boundary conditions of an eigenfunction. It will also satisfy

$$\frac{d}{dx} \left((1-x^2) \frac{dS_\lambda}{dx} \right) - \left(\frac{m^2}{1-x^2} - \kappa_0 x^2 \right) S_\lambda + \lambda_0 S_\lambda = -S.$$

Multiplying both sides of this equation by \overline{S} , integrating over $(0, \pi)$, and then integrating by parts will imply that

$$\int_0^\pi |S|^2 \sin \theta d\theta = 0.$$

This is clearly a contradiction. \square

Proposition B.2.

$$\omega_I > 0 \Rightarrow \operatorname{Im}((\lambda + a^2 \omega^2) \overline{\omega}) < 0.$$

Proof. Multiplying the angular ODE by $\overline{\omega S}$, integrating by parts, and taking imaginary parts gives

$$\begin{aligned} \omega_I \int_0^\pi \left(\left| \frac{dS}{d\theta} \right|^2 + \left(\frac{m^2}{\sin^2 \theta} + a^2 |\omega|^2 \sin^2 \theta + a^2 \mu^2 \cos^2 \theta \right) |S|^2 \right) \sin \theta d\theta = \\ - \int_0^\pi \operatorname{Im}((\lambda + a^2 \omega^2) \overline{\omega}) |S|^2 \sin \theta d\theta. \end{aligned}$$

\square

Proposition B.3. *When ω is real, we have*

$$\frac{\partial \lambda}{\partial \mu} > 0.$$

Proof. Let

$$S_\mu := \frac{\partial S}{\partial \mu}.$$

We have

$$\begin{aligned} \frac{1}{\sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{dS_\mu}{d\theta} \right) - \left(\frac{m^2}{\sin^2 \theta} - a^2(\omega^2 - \mu^2) \cos^2 \theta \right) S_\mu + \lambda S_\mu = \\ \left(2a^2 \mu \cos^2 \theta - \frac{\partial \lambda}{\partial \mu} \right) S. \end{aligned}$$

Multiplying the equation by \bar{S} , integrating by parts, and taking the real part gives

$$\int_0^\pi \left(2a^2 \mu \cos^2 \theta - \frac{\partial \lambda}{\partial \mu} \right) |S|^2 \sin \theta d\theta = 0.$$

□

C Local Theory for the Radial ODE

C.1 The Horizon

Let's apply the theory from appendix A to the radial ODE. Recall that we earlier set

$$\xi = \frac{i(am - 2Mr_+\omega)}{r_+ - r_-}.$$

First we consider the case where $am - 2Mr_+\omega \neq 0$. In this case the indicial equation has two distinct roots which do not differ by an integer. Hence a local basis of solutions to the radial ODE around r_+ will be given by

$$\left\{ (r - r_+)^\xi \rho_1(r), (r - r_+)^{-\xi} \rho_2(r) \right\}$$

where each $\rho_i(r)$ is holomorphic near r_+ and is normalized to have $\rho_i(r_+) = 1$. Our mode analysis showed that a mode solution must be of the form $A(r - r_+)^\xi \rho_1(r)$ for some $A \in \mathbb{C}$. Hence, for every ω and μ so that λ is defined, we have a unique solution to the radial ODE of the form

$$(r - r_+)^\xi \rho(r, \omega, \mu) \tag{C.1}$$

where $\rho(r, \omega, \mu)$ is analytic in r , holomorphic in ω , analytic in μ , and $\rho(r_+, \omega, \mu) = 1$. Let us remark that if a mode solution with real ω and $am - 2Mr_+\omega \neq 0$ vanishes at r_+ , it must vanish identically.

Now let's consider what happens if $am - 2Mr_+\omega = 0$. In this case the indicial equation has a double root at $\alpha = 0$ and lemma A.1 only produces one solution near r_+ . One must then consider solutions which have a logarithmic singularity at r_+ . The standard theory (see [20]) then implies that a local basis of solutions is given by

$$\{\varphi_1(r), \log(r - r_+)\varphi_2(r) + \varphi_3(r)\}$$

where the ν_i are all holomorphic near r_+ , $\varphi_1(r_+) = 1$, $\varphi_2(r_+) = 1$, and $\varphi_3(r_+) = 0$. It will be important to note that lemma A.1 implies that φ_1 is embedded in the family of local solutions C.1.

Lastly, it will be useful for the bound state analysis to note that everything said in this section so far applies verbatim to the equation

$$\Delta \frac{d}{dr} \left(\Delta \frac{dR}{dr} \right) - V_\mu R - \nu \Delta R = 0 \text{ for } \nu \in \mathbb{R}.$$

C.2 Infinity

The local existence theorems quoted in this section can be found in chapter 7 of [20]. Let us note that the radial ODE can be written as

$$\begin{aligned} \frac{dR^2}{dr^2} + \frac{\partial_r \Delta}{\Delta} \frac{dR}{dr} - \frac{V_\mu}{\Delta^2} R &= 0 \Rightarrow \\ \frac{dR^2}{dr^2} + \left(\frac{2}{r} + O(r^{-2}) \right) \frac{dR}{dr} + \left((\omega^2 - \mu^2) + \frac{2M(2\omega^2 - \mu^2)}{r} + O(r^{-2}) \right) R &= 0. \end{aligned}$$

Let's write $\omega = \omega_R + i\omega_I$. We will need to construct a local basis at infinity that depends holomorphically on ω and analytically on μ .

Lemma C.1. *For all ω and μ with $\mu^2 - \omega^2 \notin (-\infty, 0]$ there is a unique $\rho_1(r, \omega, \mu)$ which solves the radial ODE and satisfies*

$$\rho_1(r, \omega, \mu) = e^{-\sqrt{\mu^2 - \omega^2} r} r^{-1 - \frac{M(2\omega^2 - \mu^2)}{\sqrt{\mu^2 - \omega^2}}} + O \left(e^{-\sqrt{\mu^2 - \omega^2} r} r^{-2 - \frac{M(2\omega^2 - \mu^2)}{\sqrt{\mu^2 - \omega^2}}} \right).$$

Furthermore, ρ_1 depends holomorphically on ω and μ . The square root is defined by making a branch cut along the negative real numbers.

Proof. One can more or less extract a proof of this from the discussion of irregular singularities in Chapter 7 section 2 of [20]. For the sake of completeness we will give the needed slight extension. We let C denote a

sufficiently large constant which can be taken holomorphic in μ and ω . One may find a formal solution to the radial ODE of the form

$$L(r, \omega, \mu) := e^{-\sqrt{\mu^2 - \omega^2} r} r^{-1 - \frac{M(2\omega^2 - \mu^2)}{\sqrt{\mu^2 - \omega^2}}} \sum_{j=0}^{\infty} \frac{a_j(\omega, \mu)}{z^j}$$

where $a_0 = 1$ and the a_j are holomorphic in ω and μ . See Chapter 7 section 1 of [20] for the computations behind this. Let's set

$$L_n(r, \omega, \mu) := e^{-\sqrt{\mu^2 - \omega^2} r} r^{-1 - \frac{M(2\omega^2 - \mu^2)}{\sqrt{\mu^2 - \omega^2}}} \sum_{j=0}^{n-1} \frac{a_j(\omega, \mu)}{z^j}.$$

Then

$$\frac{d^2 L_n}{dr^2} + \frac{\partial_r \Delta}{\Delta} \frac{dL_n}{dr} - \frac{V_\mu}{\Delta^2} L_n = e^{-\sqrt{\mu^2 - \omega^2} r} r^{-1 - \frac{M(2\omega^2 - \mu^2)}{\sqrt{\mu^2 - \omega^2}}} B_n(r, \omega, \mu)$$

where $B_n(r, \omega, \mu) \leq Cr^{-n-1}$. Let's look for a solution ρ_1 of the form

$$\rho_1(r, \omega, \mu) = L_n(r, \omega, \mu) + \epsilon(r, \omega, \mu).$$

We must have

$$\begin{aligned} \frac{d^2 \epsilon}{dr^2} + \frac{\partial_r \Delta}{\Delta} \frac{d\epsilon}{dr} - \frac{V_\mu}{\Delta^2} \epsilon &= -e^{-\sqrt{\mu^2 - \omega^2} r} r^{-1 - \frac{M(2\omega^2 - \mu^2)}{\sqrt{\mu^2 - \omega^2}}} B_n \Leftrightarrow \\ &\frac{d^2 \epsilon}{dr^2} + (\omega^2 - \mu^2) \epsilon = \\ &-e^{-\sqrt{\mu^2 - \omega^2} r} r^{-1 - \frac{M(2\omega^2 - \mu^2)}{\sqrt{\mu^2 - \omega^2}}} B_n - \frac{\partial_r \Delta}{\Delta} \frac{d\epsilon}{dr} + \left(\frac{V_\mu}{\Delta^2} + (\omega^2 - \mu^2) \right) \epsilon. \end{aligned}$$

Let's set

$$K(r, t) := \frac{e^{\sqrt{\mu^2 - \omega^2}(r-t)} - e^{-\sqrt{\mu^2 - \omega^2}(r-t)}}{2\sqrt{\mu^2 - \omega^2}}.$$

Variation of parameters gives

$$\begin{aligned} \epsilon(r, \omega, \mu) &= \\ &\int_r^\infty K(r, t) \left(e^{-\sqrt{\mu^2 - \omega^2} t} t^{-1 - \frac{M(2\omega^2 - \mu^2)}{\sqrt{\mu^2 - \omega^2}}} B_n(t) - \left(\frac{V_\mu(t)}{\Delta^2(t)} + \omega^2 - \mu^2 \right) \epsilon(t) + \frac{\partial_t \Delta(t)}{\Delta(t)} \frac{d\epsilon}{dr}(t) \right) dt. \end{aligned}$$

We may solve this by iterating in the usual fashion. Set $h_0(r, \omega, \mu) = 0$ and

$$h_{j+1}(r, \omega, \mu) = \int_r^\infty K(r, t) \left(e^{-\sqrt{\mu^2 - \omega^2} t} t^{-1 - \frac{M(2\omega^2 - \mu^2)}{\sqrt{\mu^2 - \omega^2}}} B_n - \left(\frac{V_\mu(t)}{\Delta^2(t)} + \omega^2 - \mu^2 \right) h_j(t) + \frac{\partial_t \Delta(t)}{\Delta(t)} \frac{dh_j}{dr}(t) \right) dt.$$

It is easy to see that

$$|h_1(r, \omega, \mu)| + \left| \frac{dh_1}{dr}(r, \omega, \mu) \right| \leq \frac{C e^{-\sqrt{\mu^2 - \omega^2} r} r^{-1 - \frac{M(2\omega^2 - \mu^2)}{\sqrt{\mu^2 - \omega^2}}}}{r^n} \left(n + \frac{M(2\omega^2 - \mu^2)}{\sqrt{\mu^2 - \omega^2}} \right)^{-1}.$$

Then, with induction one can show that

$$|h_{j+1} - h_j|(r, \omega, \mu) + \left| \frac{dh_{j+1}}{dr} - \frac{dh_j}{dr} \right|(r, \omega, \mu) \leq \frac{C^j e^{-\sqrt{\mu^2 - \omega^2} r} r^{-1 - \frac{M(2\omega^2 - \mu^2)}{\sqrt{\mu^2 - \omega^2}}}}{r^n} \left(n + \frac{M(2\omega^2 - \mu^2)}{\sqrt{\mu^2 - \omega^2}} \right)^{-j}.$$

For ω and μ in a sufficiently small compact set and sufficiently large n , the $h_j(r, \omega, \mu)$ will converge uniformly in r , ω , and μ . \square

It is of course easy to pick a second holomorphic family of solutions $\rho_2(r, \omega, \mu)$ that is linearly independent of ρ_1 . One can show (Chapter 7 of [20]) that we must then have

$$\rho_1(r, \omega, \mu) \sim e^{-\sqrt{\mu^2 - \omega^2} r} r^{-1 - \frac{M(2\omega^2 - \mu^2)}{\sqrt{\mu^2 - \omega^2}}},$$

$$\rho_2(r, \omega, \mu) \sim e^{\sqrt{\mu^2 - \omega^2} r} r^{-1 + \frac{M(2\omega^2 - \mu^2)}{\sqrt{\mu^2 - \omega^2}}}.$$

Lastly, we note that a similar discussion can be carried out for the equation

$$\Delta \frac{d}{dr} \left(\Delta \frac{dR}{dr} \right) - V_\mu R + \nu \Delta R = 0.$$

C.3 Connection Coefficients

Let's fix some set of frequency parameters with $\mu^2 - \omega_R^2 \notin (-\infty, 0]$. Above we constructed $\rho(r, \omega, \mu)$ holomorphic in ω and μ so that $(r - r_+)^{\xi} \rho(r, \omega, \mu)$ gives a solution to the radial ODE with the correct boundary condition at r_+ . We can then introduce connection coefficients $A(\omega, \mu)$ and $B(\omega, \mu)$:

$$R(r, \omega, \mu) := (r - r_+)^{\xi} \rho(r, \omega, \mu) = A(\omega, \mu) \rho_1(r, \omega, \mu) + B(\omega, \mu) \rho_2(r, \omega, \mu).$$

Let $W(\cdot, \cdot)$ denote the Wronskian. Then

$$A = \frac{W(R, \rho_2)}{W(\rho_1, \rho_2)}.$$

Thus A is holomorphic in ω and analytic in μ . Similarly, B is holomorphic in ω and analytic in μ .

D Distant Bound Keplerian Orbits and Unstable Modes

Since it presents a convenient and intuitive, if overly simplified, picture, we will informally present the heuristic argument connecting distant bound Keplerian orbits and unstable modes. We refer the reader to [31] for more details. We start by positing the existence of timelike geodesics which are “very far away” from the black hole yet are also bounded away from infinity. Additionally, we assume that that our geodesics are “slowly prograde” in the sense that

$$0 < \frac{d\phi}{dt} \ll \frac{a}{2Mr_+}. \quad (\text{D.1})$$

On physical grounds, one expects to be able to construct approximate wave packet solutions to the Klein-Gordon equation which are localized near these geodesics. It is reasonable to expect such a wave packet to be given by a superposition of modes

$$\varphi(t, r, \theta, \phi) := e^{-i\omega t} e^{im\phi} S_{ml}(\theta) R(r)$$

where we allow complex $\omega = \omega_R + i\omega_I$, but assume that $|\omega_I| \ll \omega_R$. It is easy to see that (D.1) implies that the modes must be superradiant:

$$am - 2Mr_+\omega_R > 0.$$

Since the geodesics are bounded away from infinity, we shall want R to fall off at a sufficiently fast exponential rate near spatial infinity. An analysis of R 's ODE (see section 2 and appendix C) shows that this can be achieved if

$$0 < \omega_R < \mu.$$

Let χ be a bump function supported near r_+ . Then, using Kerr-star coordinates, set

$$\tilde{t}(t^*, r) := \chi(r)t^* + (1 - \chi(r))t(t^*, r).$$

Now, for every $\tau > 0$, let Σ_τ be the hypersurface defined by $\tilde{t} = \tau$. This is a uniformly spacelike hypersurface which intersects the horizon transversally. Since φ is superradiant, one easily finds that $\tau_2 > \tau_1$ implies

$$\int_{\Sigma_{\tau_2}} \mathbf{J}_\alpha^T n_{\Sigma_{\tau_2}}^\alpha > \int_{\Sigma_{\tau_1}} \mathbf{J}_\alpha^T n_{\Sigma_{\tau_1}}^\alpha. \quad ^7$$

That is, the energy along Σ_τ must be growing with time. Note that the exponential decay of R at infinity implies that the energy along Σ_τ is finite. Since our geodesic lives far outside the ergoregion, we expect that R peaks far away from the ergoregion, and hence that the energy along Σ_τ can increase if and only if φ is increasing with time. Of course, this is only possible if $\omega_I > 0$.

References

- [1] S. Alinhac *Geometric Analysis of Hyperbolic Differential Equations: An Introduction* Cambridge University Press, 2010
- [2] L. Andersson and P. Blue *Hidden symmetries and decay for the wave equation on the Kerr spacetime* (2009) arXiv:0908.2265 [math.AP]
- [3] H. Beyer *On the stability of the massive scalar field in Kerr space-time* J. Math. Phys. **52** (2011), Issue 10, 102502
- [4] V. Cardoso, O. Dias, J. Lemos, S. Yoshida *Black-hole bomb and super-radiant instabilities* Phys. Rev. D **70** (2004), Issue 4, 044039
- [5] V. Cardoso, O. Dias, S. Yoshida *Classical instability of Kerr-AdS black holes and the issue of final state* Phys. Rev. D **74** (2006), Issue 4, 044008

⁷Recall that \mathbf{J}_α^T is defined by (1.1).

- [6] D. Christodoulou *Reversible and irreversible transformations in black-hole physics* Phys. Rev. Lett. **25** (1970), 15961597
- [7] M. Dafermos and I. Rodnianski *A proof of the uniform boundedness of solutions to the wave equation on slowly rotating Kerr backgrounds* Inventiones Math. **185** (2011), no. 3, 467-559
- [8] M. Dafermos and I. Rodnianski *Decay for solutions of the wave equation on Kerr exterior spacetimes I-II: The cases $|a| \ll M$ or axisymmetry* (2010) arXiv:1010.5132 [gr-qc]
- [9] M. Dafermos and I. Rodnianski *Lectures on black holes and linear waves* Institut Mittag-Leffler Report no. 14, 2008/2009 (2008)
- [10] M. Dafermos and I. Rodnianski *The black hole stability problem for linear scalar perturbations* Proceedings of the 12 Marcel Grossmann Meeting (2012)
- [11] T. Damour, N. Deruelle, R. Ruffini *On Quantum Resonances in Stationary Geometries* Lett. Al Nuovo Cimento **15** (1976), no. 8, 257-262
- [12] S. Detweiler *Klein-Gordon equation and rotating black holes* Phys. Rev. D **22** (1980), no. 10, 2323-2326
- [13] S. Dolan *Instability of the massive Klein-Gordon field on the Kerr spacetime* Phys. Rev. D **76** (2007), Issue 8, 084001
- [14] F. Finster, N. Kamran, J. Smoller, S.-T. Yau *Decay of solutions of the wave equation in the Kerr geometry* Comm. Math. Phys. **264** (2006), no. 2, 465-503
- [15] F. Finster, N. Kamran, J. Smoller, S.-T. Yau *Erratum: Decay of solutions of the wave equation in the Kerr geometry* Comm. Math. Phys. **280** (2008), no. 2, 563-573
- [16] J. Hartle and D. C. Wilkins *Analytic Properties of the Teukolsky Equation* Comm. Math. Phys. **38** (1974), no. 1, 47-63
- [17] S. Hod and O. Hod *Analytic treatment of the black-hole bomb* Phys. Rev. D **81** (2010), Issue 6, 061502(R)
- [18] J. Luk *A vector field method approach to improved decay for solutions to the wave equation on a slowly rotating Kerr black hole* Analysis and PDE **5** (2012), no. 3, 553-623

- [19] J. Metcalfe, D. Tataru, and M. Tohaneanu *Price's law on non-stationary spacetimes* Adv. in Math. **230** (2012), no. 3, 995-1028
- [20] F. Olver *Asymptotics and Special Functions* A K Peters, 1997
- [21] B. O'Neill *The Geometry of Kerr Black Holes* A K Peters, 1995
- [22] R. Penrose *Gravitational collapse: the role of general relativity* Rev. del Nuovo Cimento **1**, (1969) 272-276
- [23] W. Press and S. Teukolsky *Floating orbits, superradiant scattering and the black-hole bomb* Nature **238** (1972), 211-212
- [24] W. Press and S. Teukolsky *Perturbations of a rotating black hole. II. Dynamical stability of the Kerr metric* Astrophysical Journal, **185** (1973), 649-673
- [25] Y. Shlapentokh-Rothman *Quantitative Mode Stability on the Real Axis for the Wave Equation on the Kerr Spacetime* preprint. (2013)
- [26] D. Tataru and M. Tohaneanu *Local energy estimates on Kerr black hole backgrounds* IMRN **2011** (2011), no. 2, 248-292
- [27] M. Tohaneanu *Strichartz estimates on Kerr black hole backgrounds* Transactions of the AMS **364** (2012), n0. 2, 689-702
- [28] R. Wald *General Relativity* The University of Chicago Press, 1984
- [29] B. Whiting *Mode stability of the Kerr black hole* J. Math. Phys. **30** (1989), no. 6, 1301-1305
- [30] Y. Zeldovich *The Generation of Waves by a Rotating Body* Pisma v Redakstiyu Zhurnal Eksperimentalnoi i Teoreticheskoi Fiziki **14** (1971), no. 270, English translation: JETP Letters **14**, 180 (1971).
- [31] T. Zouros and D. Eardley *Instabilities of massive scalar perturbations of a rotating black hole* Ann. of Phys **118** (1976), Issue 1, 139-155