

Clonal interference and Muller's ratchet in spatial habitats

Jakub Otwinowski*[†], Joachim Krug[‡]

October 4, 2022

Abstract

Competition between independently arising beneficial mutations is enhanced in spatial populations due to the linear rather than exponential growth of clones. Recent theoretical studies have pointed out that the resulting fitness dynamics is analogous to a surface growth process, where new layers nucleate and spread stochastically, leading to the build up of scale-invariant roughness. This scenario differs qualitatively from the standard view of adaptation in that the speed of adaptation becomes independent of population size while the fitness variance does not. Here we exploit recent progress in the understanding of surface growth processes to obtain precise predictions for the universal, non-Gaussian shape of the fitness distribution for one-dimensional habitats, which are verified by simulations. When the mutations are deleterious rather than beneficial the problem becomes a spatial version of Muller's ratchet. In contrast to the case of well-mixed populations, the rate of fitness decline remains finite even in the limit of an infinite habitat, provided the ratio U_d/s^2 between the deleterious mutation rate and the square of the (negative) selection coefficient is sufficiently large. Using again an analogy to surface growth models we show that the transition between the stationary and the moving state of the ratchet is governed by directed percolation.

1 Introduction

The appearance of a mutation in a population and its fixation or loss is the most basic process of adaptation. This process determines the rate of evolution, or how quickly populations adapt to new environments. One approach in quantitative models of adaptation is to assume that non-neutral mutations are rare. In this regime, generally referred to as periodic selection [1, 2], the population has no genetic variation except for brief periods when a mutation sweeps through and fixates, and therefore the rate of fixation is mutation limited.

When mutations are more common, genetic variation builds up, and when there is little or no recombination, mutations must compete with each other to fixate. This is relevant in the case of

*Emory University, Physics Department Atlanta, Georgia, United States

[†]University of Pennsylvania, Biology Department, Philadelphia, Pennsylvania, United States

[‡]University of Cologne, Institute for Theoretical Physics, Köln, Germany

beneficial mutations, since recent microbial experiments suggest they are more common than previously thought [3, 4, 5]. Microbial evolution experiments have observed reduced rates of evolution due to the competition between beneficial mutations [6, 7]. Fisher’s fundamental theorem equates the rate of evolution with the variance of the fitness distribution [8], which can be approximated analytically in simplified population genetic models. These recent theoretical analyses have found the rate of evolution in large populations of asexuals is not proportional to the total supply rate of beneficial mutations, but depends much more weakly (logarithmically) on population size and mutation rate [9, 10, 11, 12, 13, 14].

Deleterious mutations are more common than beneficial ones, but their chance of fixation is much smaller and vanishes for infinite populations. However, in finite populations and in the absence of beneficial mutations and recombination, deleterious mutations will eventually fix by genetic drift, leading to a fitness decline known as Muller’s ratchet [15, 16]. Determining the rate of the ratchet as a function of population size, mutation rate and selection strength is a long-standing problem that continues to attract considerable interest [17, 18, 19, 20, 21, 22, 23, 24]. Recombination can prevent Muller’s ratchet and also mitigates the slowdown in the rate of evolution from clonal interference, which is why Muller’s ratchet and clonal interference are often argued as reasons for an evolutionary advantage of sex [25, 26, 27].

These analyses were limited to well-mixed populations, where each individual competes with the whole population, such as microbes in liquid culture. However, many populations are not well-mixed, but are confined in space such that they only compete with a limited neighborhood population on timescales of a generation. Spatially structured population genetics have been studied with finitely subdivided, and continuous populations [28, 29, 30, 31, 32, 33]. Interestingly, when mutations are rare a single beneficial mutation can effectively compete with the whole population, and the fixation probability is the same in well-mixed and spatially structured populations [34, 35].

Recently, models incorporating large amounts of beneficial mutations and one or two dimensional spatial structure have found the rate of evolution to be even slower than in well-mixed populations, as the slower than exponential growth of clones increases the likelihood of competition [36, 37, 38, 39, 40, 41]. In fact, the rate of evolution becomes independent of system size, while the variance (in the steady state) scales as a power of population size, violating Fisher’s theorem [38, 39]. This also implies that there is a long transient regime during which the stationary variance builds up, while the speed of adaptation is constant.

Here, we study the transient regime of Wright-Fisher models of evolution on a one-dimensional lattice for both adaptation with beneficial mutations, and Muller’s ratchet. In the former case the fitness variance grows as a power law in time, and saturates at a value determined by the system size (also as a power law) [38, 39]. The fitness may be pictured as a surface in space, which advances over

time. This behavior is analogous to surface growth models in physics, where particles are deposited on an initially flat surface, which develops roughness over time [42, 43].

In section 2, we introduce the model for adaptation on a one dimensional lattice, and review previous results. In section 3, we show the model of adaptation belongs to a class of surface growth models called the Kardar-Parisi-Zhang (KPZ) universality class [44, 45, 46, 47]. By exploiting the equivalence to models of surface growth, this scenario can be described in great detail, including in particular the non-Gaussian shape of the fitness distribution. In section 4, we modify the model to study Muller's ratchet. We find that for certain parameters the rate of fitness decline does not go to zero as the population size becomes large, and we characterize the transition between fitness decline and no decline. The model with deleterious mutations is similar to a different class of models in surface growth physics, and we use this analogy to find other asymptotic properties.

2 Model

The spatial constraints are realized as a one dimensional lattice of size L with periodic boundary conditions, where each point represents a single organism that occupies a space [38]. The evolution follows standard Wright-Fisher dynamics in discrete generations, where the next fitness of each site is chosen randomly from one of the parents in the neighborhood, weighted according to their fitness. The smallest possible neighborhood in one dimension is such that the child in the next generation inherits the fitness from only two possible parents, that is, the fitness $f_i(t+1)$ of site i at generation $t+1$ is chosen from either $f_i(t)$ or $f_{i+1}(t)$.

In the case of a homogeneous system of fitness 1, where a single mutant appears with fitness $1+s$, the fixation probability for a beneficial mutation is the same as in the well-mixed case, $\pi = 2s$ for $s \ll 1$ [34, 35]. Intuitively, the fixation probability is unaffected because a single mutation has ample time to compete with the entire system, regardless of spatial structure. Since the fixation probability is the same, the speed of evolution in the periodic selection regime is the same as in the well-mixed case. What is different is the timescale of fixation.

The boundary between two domains with different fitnesses is a biased random walker, and the speed of this walker is the expected value of its displacement after one time step, $c = s/2$ for small s . In the continuum limit, this model corresponds to a special case of the more general stochastic Fisher equation (or SFKPP equation) [48, 49, 50], where it is possible to have traveling waves with speed $c \sim s$ in the strong noise regime, or $c \sim \sqrt{s}$ in the weak noise regime. However, the dependence of the wave speed on s does not change the essential features.

Importantly, the time for fixation may be much longer in the presence of spatial structure compared

to well-mixed populations. A wave spreading with finite speed c will take time $t_{\text{fix}} \sim L/c$ to cover the whole system (and total population size $N \sim L$), as opposed to a well-mixed population where $t_{\text{fix}} \sim \log(N)$. The slow spread of mutations make it more likely that many clones exist simultaneously in large systems. A site may also contain more than one organism, in which case c is different, but it does not change the overall results [39] (unless interference happens within one site).

Since we are interested in the rate of evolution during competition, a steady rate of beneficial mutations is supplied, akin to a population adapting to a new environment. Beneficial mutations appear randomly at rate U_b per site per generation (deleterious mutations are studied in section 4). We assume that mutations have independent effects, with no epistasis, and therefore increase the fitness according to $\log f' = \log f + s$, where s is a constant on the order of 1%.

An important quantity is the rate of fitness changes $V = \lim_{t \rightarrow \infty} \langle \log f \rangle / t$, where the average is over the population. When the time between mutations to appear and become established, $t_{\text{mut}} = (\pi U_b L)^{-1}$, is much longer than t_{fix} , V is mutation limited: $V = s \pi U_b L = 2s^2 U_b L$. However, when $t_{\text{mut}} \sim t_{\text{fix}}$, multiple unfixed mutations in the population compete with each other, slowing down V . In well mixed populations the condition for mutation limited adaptation is that there should be less than one new beneficial mutation per generation. In contrast, with spatial structure $t_{\text{mut}} \sim t_{\text{fix}}$ defines a characteristic interference length scale $L_c \sim (c/U_b)^{1/2}$, above which mutation competition sets in. In this competitive regime, the rate of evolution no longer depends on the supply of beneficial mutations, but V becomes independent of L for $L > L_c$ [38, 39]. Using this observation and dimensional analysis, one may deduce that this maximum speed grows as $U_b^{1/2}$ in one dimension, and $U_b^{1/3}$ in two dimensions.

In the following section we describe the fitness distribution in the transient regime of the evolution, before reaching the steady state, by exploiting an analogy to surface growth physics.

3 Adaptation with many beneficial mutations

3.1 Analogy to surface growth

The rough spatial profile of the fitness resembles a typical surface seen in surface growth models [38, 42]. In surface growth, particles are deposited on an initially smooth surface randomly, and they may diffuse or stick to each other, gradually forming a rough surface. Many simple models of surface growth were studied by statistical physicists interested in non-equilibrium systems [42, 43]. They discovered that a large number of models share the same properties in the continuum, long-time limit, where many of the microscopic details of the model do not matter, and these classes of models, or universality classes, share the same symmetries.

The evolutionary model defined here is equivalent to a surface growth model called polynuclear growth

[51, 52, 53, 54] (PNG), in the limit $s \rightarrow \infty$. In PNG, the process of surface growth may be divided into two parts, nucleation (mutation), and spreading (selection). Nucleation occurs with low probability at any point, at a certain rate, U_b , which corresponds to adding a small block of height to the surface (log fitness). The nucleated block then grows laterally forming a new layer. Depending on the size of the lattice, the surface grows layer by layer (corresponding to the periodic selection regime) or the surface roughens due to multiple simultaneous nucleation events (corresponding to clonal interference) [51, 52]. In the rough regime the PNG model belongs to the universality class of growth processes described on large length and time scales by the KPZ equation, a nonlinear stochastic partial differential equation [44, 53, 54].

While in PNG the spreading is fast and deterministic, in the evolutionary model it is stochastic, and the new layer may even disappear [52]. The boundaries may collide with each other, and they either annihilate or stack up creating differences in log fitness greater than s . From the point of view of surface growth it is natural to hypothesize that the universal features of the PNG model are robust with respect to these differences, but this has to be verified by explicit simulations. The test of the universality hypothesis proceeds in two steps. First, one estimates the scaling exponents governing the power law dependence of the standard deviation of the surface height (or log fitness) distribution on time and system size. Second, the shape of the full distribution of height fluctuations is considered.

In surface growth, starting from flat initial conditions, the standard deviation of the surface height distribution grows in time as $\sigma(t) \sim t^\beta$, where β is the growth exponent, then reaches a steady state when the correlation length reaches the size of the system [42, 55]. In the steady state, $\sigma(t \rightarrow \infty) \sim L^\alpha$ where α is the saturation exponent. Figure 1a confirms this scenario for the evolution model. The crossover time is where saturation sets in (the elbow), and it scales as $L^{\alpha/\beta}$. One may try to measure the exponents from the simulations, but based on the similarity to the PNG model one expects that the scaling exponents are those of the one-dimensional KPZ-equation, $\alpha = 1/2$, $\beta = 1/3$ and $\alpha/\beta = 3/2$. Figure 1b shows that the data indeed collapses when plotted as σ^2/L versus $t/L^{3/2}$. In the evolutionary context the saturation time scale $\sim L^{3/2}$ is proportional to the fixation time of beneficial mutations [39]. Note that these values of the exponents characterize the asymptotic, long time and large scale behavior of the model, and the behavior in the pre-asymptotic regime may be somewhat different [38].

3.2 Transient fitness distributions

Over the past decade, a much more refined characterization of the KPZ universality class has been developed that extends beyond the values of the scaling exponents α and β to the full probability distribution of surface height fluctuations [45, 46, 47]. The essence of this refined universality hypothesis




Figure 1: (a) Variance σ^2 of the log fitness distribution as a function of time for different system sizes, $L = 2^{13}$ (green circles), $L = 2^{14}$ (blue squares), and $L = 2^{15}$ (red crosses), with $s = 0.05$ and $U_b = 10^{-5}$. After a transient regime, σ^2 saturates at a value that depends on L . (b) When the data is rescaled as σ^2/L and $t/L^{3/2}$ it collapses onto a single curve, indicating that in fact $\sigma(t) \sim t^{1/3}$ and $\sigma(t \rightarrow \infty) \sim L^{1/2}$ as predicted by KPZ theory. Data points are averages over 50 simulations.

is that the log fitnesses (or surface heights) can be written as

$$\log f_i(t) = Vt + (\Gamma t)^{1/3} \chi, \quad (1)$$

where χ is a random variable from one of the *Tracy-Widom (TW) distributions*, V is the long-time growth rate, and Γ is a constant related to the parameters of the KPZ equation [45]. From eq. (1) we find the width of the distribution:

$$\sigma^2(t) = \text{var}(\log f_i(t)) = (\Gamma t)^{2/3} \text{var}(\chi). \quad (2)$$

The TW distributions were first discovered in fluctuations of the largest eigenvalues of random matrices [56]. The relation to the PNG model was established by mapping the PNG surface height to the length of the longest increasing subsequence of random permutations [54, 57], and subsequently TW universality was derived directly from the KPZ equation [47, 58]. Remarkably, the distributions were found to be geometry dependent, with the flat (monomorphic) initial condition leading to the TW distribution characteristic of random matrices from the Gaussian orthogonal ensemble (GOE).

Here we show numerically that, despite the additional randomness of the stochastic spreading, the distribution of fitnesses in the non-stationary regime of the spatial evolution model is a TW distribution




Figure 2: Skewness and kurtosis of the fitness distributions from 200 simulations compared to the known values for the GOE Tracy-Widom distribution. $L = 2^{14}$, $s = 0.05$, $U_b = 10^{-5}$.

characteristic of the KPZ universality class. One signature of the TW distributions can be seen by measuring higher moments, such as skewness, $\langle \left(\frac{\log f - \langle \log f \rangle}{\sigma} \right)^3 \rangle$ and excess kurtosis, $\langle \left(\frac{\log f - \langle \log f \rangle}{\sigma} \right)^4 \rangle - 3$, which do not depend on the parameters V and Γ . Figure 2 shows that the skewness and kurtosis of the fitness distributions are non-zero, indicating non-Gaussianity, and they approach the known values of the GOE TW distribution.

It is also possible to compare the fitness distribution directly to the TW distribution. The parameters V and Γ can be found from the simulation data by applying linear regression to the means of equations (1) and (2). The fitnesses from the simulation are then rescaled as

$$\chi_{sim} = \frac{\log f_i - Vt}{(\Gamma t)^{1/3}}. \quad (3)$$

Figure 3 shows that in the non-stationary regime, the fitnesses fall onto the universal GOE TW distribution, which is skewed towards higher fitnesses, with tail behaviors $-\ln P(\chi)_{\chi \rightarrow \infty} \sim \chi^{3/2}$ and $-\ln P(\chi)_{\chi \rightarrow -\infty} \sim |\chi|^3$. To demonstrate the robustness of this result, we simulated a variant of the model where the selective advantage of beneficial mutations, s , is a random variable generated from an exponential distribution, a common choice in this field [11, 13]. The two data sets can be seen to be indistinguishable.

In addition, two other initial conditions were simulated. The droplet geometry in the PNG model is when the initial condition is a single nucleation site, with no additional nucleations (or mutations) allowed outside. The boundary of the initial seed grows over time, making the fitness profile curved. The deviations from this curved profile converge to the TW distribution of the Gaussian unitary ensemble (GUE) [45, 46, 54]. The droplet geometry has an interesting evolutionary analogy: It corresponds to a mutation that raises the mutation rate significantly (a mutator strain), and competes with a population




Figure 3: Scaled fitness distributions for three different initial conditions: Flat (blue squares and circles), droplet (orange crosses), and rough (green diamonds). Lines indicate the Tracy-Widom GOE (blue solid), GUE (orange dotted), and the F_0 (green dashed) distributions respectively (calculated using [59]). The scaled fitness distributions were taken from 200 simulations after 10^6 generations, with $L = 2^{18}$, $U_b = 10^{-5}$ and $s = 0.05$, except the blue squares which had exponentially distributed selection coefficients, with mean $\langle s \rangle = 0.05$. For rough initial conditions, the simulation was first run to the steady state ($L^{3/2}$ generations), and deviations from the initial condition were calculated. For the droplet geometry, a single mutation was first allowed to establish, and mutations were only allowed in that lineage. The exact shape of the droplet is unknown, so only fitnesses from the position of the initial mutation (the peak of the droplet) were used in the distribution.

that has essentially no mutations.

The third initial condition corresponds to a system with fully developed, stationary diversity (surface roughness). In this case the distribution of the deviations from the initial fitness profile is predicted to converge to a universal distribution F_0 , which does not appear directly in random matrix theory but is closely related to the TW distributions [54]. Again, the data fall nicely onto the predicted distribution.

4 Deleterious mutations

Our model may be modified to include only deleterious mutations by having a single negative selection coefficient $s < 0$. Once a mutation appears, its expansion is unfavorable, and happens only due to genetic drift. For a single mutation, the probability of fixation vanishes exponentially in N as $\pi \sim e^{Ns}$ for large negative Ns [34, 35]. When many deleterious mutations are present simultaneously in the population, the rate of fitness decline is governed by the time scale on which the subpopulation of individuals with the smallest number of mutations (the ‘least loaded class’) goes extinct by genetic drift. In the well-mixed case the number of individuals in the least loaded class is on the order of $n_0 \sim N e^{-U_d/|s|}$, where

U_d is the deleterious mutation rate [16]. Correspondingly for

$$n_0|s| \sim N|s|e^{-U_d/|s|} \gg 1. \quad (4)$$

the probability of fixation of an additional deleterious mutation in this class is exponentially small. Detailed analysis shows that under condition (4) the rate of Muller’s ratchet is also exponentially small in N [20, 22, 24], whereas for $n_0|s| < 1$ the fitness of the population declines continuously, and a description in terms of a traveling wave in fitness space, similar to that used in the context of adaptation ($s > 0$), is applicable [10]. Importantly, for a given set of mutation parameters (U_d, s) the slow ratchet condition (4) is always attained for large populations, which implies that the fitness decline effectively ceases for $N \rightarrow \infty$.

4.1 Muller’s ratchet in spatial populations




Figure 4: Deleterious mutations accumulate at a constant rate in large spatial habitats. The figure shows the number of accumulated deleterious mutations per site, or F/s after 10^6 generations for $s = -0.01$, $U_d = 10^{-3}$ (blue circles), $U_d = 10^{-4}$ (green squares), and $U_d = 10^{-5}$ (red crosses). For $U_d \leq 10^{-4}$ the number of mutations per site falls below unity for large L , while for $U_d = 10^{-3}$ about 20% of all 10^6 U_d mutations that occurred have been fixed. Results were averaged over 10 instances.

Simulations of the one-dimensional spatial model show a fundamentally different behavior in the rate of fitness decline, which we measure with $F = \langle \log f \rangle$, so that V is approximately F/t after some long time. Fig. 4 shows that for sufficiently large U_d deleterious mutations accumulate at a constant rate, which becomes independent of the habitat size, L , for large L . Exploration of the parameter space reveals that this transition in the fitness decline is sharp for certain values of U_d and s (Fig 5). The rate of fitness decline is non-monotonic in $|s|$. Initially the larger mutation effects lead to a higher rate of fitness decline with increasing $|s|$, but at the same time selection becomes more effective in eliminating the deleterious mutations, which eventually halts the fitness decline. Rescaling the fitness

by $U_d s$ collapses the curves in the region of large $|s|$, where $F \sim U_d/s$, while for very small $|s|$, mutations accumulate at close to the maximal possible rate, $F \approx U_d s t$.




Figure 5: (a) The rate of fitness decline, as indicated by the mean log fitness F , changes sharply for certain parameters U_d and s . (b) Rescaling by $U_d s$ collapses parts of the curves. Dashed line indicates s^{-2} . F was measured after 10^7 generations with $L = 2^{18}$, and mutation rates were $U_d = 10^{-6}$ (blue diamonds), 10^{-5} (green pluses), 10^{-4} (red crosses), 10^{-3} (cyan squares), and 10^{-2} (magenta circles). Results were averaged over 30 simulations, except for $U_d = 10^{-4}$ and $U_d = 10^{-3}$, which were only run once.




Figure 6: (a) A sharp transition in the density of sites with no mutations, ρ_0 , as a function of mutation effect size. When ρ_0 is large the fitness does not decline, while when $\rho_0 = 0$ the fitness declines indefinitely. (b) Scaling the x -axis by $U_d^{-1/2}$ reveals that the critical parameters are $U_d/s^2 \approx 1$. Simulations are described in Fig. 5.

To further elucidate the nature of the transition we examine the density of sites with no mutations, ρ_0 . Figure 6 shows a sharp transition in ρ_0 , between regimes where the fitness is steadily declining (the moving ratchet) and where the fitness is not declining (the stationary ratchet). The collapse of curves in Fig. 6b indicates that the transition occurs when

$$\frac{U_d}{s^2} \approx 1. \quad (5)$$

To explain this relation, consider a patch of deleterious mutants created in a single mutational event. Because $|s| \ll 1$, the boundaries of the patch perform almost symmetric random walks that are weakly biased inwards by selection. The patch disappears when the two boundaries meet. The life time τ of such an isolated patch is therefore equal to the first passage time of a random walk on the half-line with a bias $\sim |s|$ towards the origin, which has a distribution of the form [60].

$$P(\tau) \sim \tau^{-3/2} e^{-s^2\tau}. \quad (6)$$

When U_d is small, deleterious patches are created and disappear independently of each other (Fig. 7).




Figure 7: Schematic space-time view of the creation and extinction of patches of deleterious mutations in the spatial Muller’s ratchet problem. The boundaries of a patch are weakly biased random walks and the patch disappears when the two walks meet. A patch of life time τ reaches a maximal width $\sim \tau^{1/2}$. The distribution of life times is heavy-tailed for small $|s|$ (see Eq. 6).

To estimate the density ρ_0 of sites without deleterious mutations, we note that $1 - \rho_0$ is the probability that a deleterious mutation is found at a randomly chosen point in space, x , at a random time t . In other words, $1 - \rho_0$ is the fraction of the space-time area in Fig. 7 that is covered by deleterious patches. A patch with life time τ reaches a spatial extension of order $\tau^{1/2}$, and hence its area is $a \sim \tau^{3/2}$. Using (6) the distribution of a is found to be $P(a) \sim a^{-4/3} e^{-s^2 a^{2/3}}$, from which the average area of a patch is deduced as $\langle a \rangle \sim s^{-2}$. Since patches are created with probability U_d per unit time and space, it follows finally that

$$1 - \rho_0 \sim \frac{U_d}{s^2} \quad (7)$$

at least when $U_d/s^2 \ll 1$ so that the patches remain isolated. Assuming that the dependence of ρ_0 on the parameter combination U_d/s^2 continues to hold up to the point where the merging of patches leads to the global extinction of the least loaded class ($\rho_0 = 0$), we conclude that the transition from the stationary to the moving ratchet is indeed determined by a condition of the form (5). Support for

this assumption is provided in Fig. 8a, which shows that simulation results for ρ_0 obtained for different values of s and U_d collapse onto a single curve when plotted against U_d/s^2 .




Figure 8: (a) The density of sites with no mutations, ρ_0 , is a function of U_d/s^2 that vanishes at $U_d/s^2 \approx 1$. The behavior of $\rho_0^{1/\beta}$ is approximately linear at the critical point, which is consistent with the DP prediction (8). (b) Similarly, the scaled rate of fitness decline $V/U_d s$ in the moving phase is a function of U_d/s^2 and vanishes at the transition in accordance with (8). Dashed line indicates $U_d/s^2 = 1$. Simulations are described in Fig. 5. Here V was estimated more accurately to exclude the genetic load, by finding the slope of $F(t)$ for the second half of simulation. Points with the lowest mutation rate, $U_d = 10^{-6}$, were omitted, because they did not have enough time to reach the asymptotic velocity.

The relation (7) also explains the behavior of the fitness in the regime of large $|s|$ in Fig. 5. In the stationary phase of the ratchet the fitness is independent of time and given by $F = s(1 - \rho_0) \sim U_d/s$, hence $F/U_d s \sim 1/s^2$ independent of U_d . Note that the behavior of ρ_0 in (7) is different from the well-mixed case, where $\rho_0 = e^{U_d/s} \approx 1 - \frac{U_d}{|s|}$. For a given selection strength $|s|$, the deleterious mutation rate required to set the ratchet into motion is $U_d \sim s^2$ in the spatial case, much smaller than the corresponding value $U_d \sim |s| \ln(N|s|)$ obtained from (4) in the well-mixed setting.

4.2 Nonequilibrium wetting and critical exponents

For a detailed characterization of the transition between the stationary and the moving spatial ratchet we exploit the similarity of our model to a class of surface growth models that are referred to as non-equilibrium wetting models [61, 62, 63, 64]. In a wetting transition a macroscopic layer of one phase (typically a liquid) forms on top of another phase (typically a solid substrate). Non-equilibrium wetting describes the transition between a layer that is bound to the substrate (the stationary ratchet), and one that grows indefinitely (the moving ratchet). Specifically, in the limit $s \rightarrow \infty$ our model becomes equivalent to an unrestricted solid-on-solid model with no evaporation inside plateaus [62], for which the wetting transition has been shown to be governed by directed percolation (DP) [65]. Directed percolation is a broad universality class of nonequilibrium phase transitions that occur between an ‘active’ and an

‘extinct’ state, for example, an infectious disease spreading in a population. In the present context the active phase is the population in the least loaded class that persists indefinitely in the stationary ratchet state and goes extinct at the transition.




Figure 9: Decay of the density of sites with no mutations, ρ_0 , with time approximately follows a power law associated with the DP class at the critical point. High mutation rates cause ρ_0 to approach zero at long times, while small amounts of deleterious mutations lead to a non-zero value of ρ_0 at long times. In between, around $U_d/s^2 \approx 1$, $\rho_0(t) \sim t^{-\theta}$, with DP-exponent $\theta \approx 0.159$ (black line). Mutation rates varied from $U_d = 9 \times 10^{-5}$ (top line), to $U_d = 1 \times 10^{-4}$ (bottom line), with increments of 10^{-6} . $L = 2^{18}$, $s = -0.01$, and data were averaged over 24 runs.

The association with DP predicts power law behaviors of various quantities near the transition. In particular, at the transition, $\rho_0(t) \sim t^{-\theta}$, where $\theta = 0.159464\dots$ [65]. Figure 9 shows that ρ_0 decays slowly for parameters close to $U_d/s^2 = 1$, consistent with the power law predicted by DP. Moreover, the stationary density of sites with no mutations and the rate of fitness decline are predicted to behave as

$$\rho_0 \sim (u_c - u)^{\beta_{DP}} \text{ for } u < u_c \text{ and } |V| \sim (u - u_c)^{\nu_{\parallel}} \text{ for } u > u_c, \quad (8)$$

at the transition, where $u = U_d/s^2$ is the scaled mutation rate with critical value $u_c \approx 1$, and $\beta_{DP} = 0.276486\dots$ and $\nu_{\parallel} = 1.733847\dots$ are the order parameter exponent and the temporal correlation length exponent of DP, respectively [65]. While the additional stochasticity associated with the smallness of the selection coefficient in our model makes it difficult to find these exponents numerically with any accuracy, the data shown in Fig. 8 are clearly consistent with the power laws (8). At the transition the (negative) mean fitness and the variance of the fitness are predicted to grow logarithmically with time [61, 63], which is also borne out by the simulations (Fig. 10).




Figure 10: Logarithmic dependence of fitness and fitness variance near the critical point. Simulations are described in figure 9.

5 Discussion

In this paper we have explored the effects of spatial structure on two common evolutionary scenarios characterized by a large and constant supply of beneficial or deleterious mutations. In both cases the fact that selection acts through local, rather than global competition leads to profound modifications of the familiar well-mixed dynamics. For the case of adaptation the most conspicuous effect is the existence of a limiting rate of adaptation that becomes independent of the population size for large populations. At the same time the lack of communication between different parts of the habitat implies that the fitness variance grows without bound, invalidating the proportionality between these two quantities expected from Fisher’s fundamental theorem [38, 39]. Similarly, our results for Muller’s ratchet show that selection in spatial habitats is weakened to the extent that the fitness declines at a finite rate even for infinitely large populations, provided the condition $U_d/s^2 > 1$ is satisfied. Figure 11 summarizes the behavior of the rate of fitness change in the different regimes considered in this paper.

By exploiting analogies with models of surface growth, we have arrived at a detailed statistical characterization of the fitness evolution in one-dimensional spatial habitats. The model with beneficial mutations has the scaling exponents and universal distribution that belong to the KPZ universality class, and we provide evidence that the model with only deleterious mutations is in the directed percolation class. While our model becomes similar to the PNG and non-equilibrium wetting models in the limit of strong selection, it was not a priori evident that the additional stochasticity associated with genetic drift would leave the asymptotic behavior unchanged.

Knowing the universality class has implications for generalizations of the model. For example, based on our understanding of KPZ-type surface growth processes, we may conclude that the saturation of the speed of adaptation holds in any habitat dimension and for a broad class of distributions of selection




Figure 11: Summary of evolutionary regimes as a function of mutation rate U and selection coefficient s , where the latter is assumed to be the same for all mutations. In all regimes the rate of fitness change V remains finite in the limit of infinite population size or habitat size. It is positive for adaptation ($s > 0$), negative in the moving Muller’s ratchet regime where $s < 0$ and $U > s^2$, and zero in the stationary ratchet regime.

coefficients, including those that are fatter than exponential. Also the association between the spatial Muller’s ratchet and DP is expected to extend to two-dimensional (planar) habitats, including the dependence of the transition on the parameter combination U_d/s^2 .

A recent simulation study has investigated a range of two-dimensional KPZ-models and identified a set of geometry-dependent universal distributions that are qualitatively similar to those found in the one-dimensional case [66]. Spatial evolution models in planar habitats have been considered in the context of cancer progression, where the distribution of waiting times t_k until the occurrence of a given number k of mutations is of central interest [67]. In the surface growth analogy, this corresponds to the time when the surface reaches a given height. Using the probabilistic concept of first passage percolation, it can be shown that such waiting times in KPZ-type growth processes again follow KPZ statistics [45]. This implies that the distribution of ‘waiting times to cancer’, which was argued in [67] to be Gaussian for small k , should asymptotically approach the two-dimensional analogue of the TW distribution found in [66].

A natural open question concerns the behavior of spatial populations that can acquire both deleterious and beneficial mutations. In the well-mixed case it is known that beneficial mutations dominate the behavior of large populations, in the sense that the fitness increases at a positive rate provided that a finite fraction of mutations are beneficial [12, 14]. In preliminary simulations we have explored a one-dimensional model where both types of mutations occur at rates U_b and U_d , respectively, with a single selection magnitude $|s| = 0.01$. When U_d is small the deleterious mutations do not accumulate, but do provide a genetic load. The genetic load does not affect the adaptation of beneficial mutations, and the associated growth exponents and fitness distribution are the same as for the model without any deleterious mutations. When U_d is larger, there is a competition between the accumulation of

deleterious and beneficial mutations, and the fitness may go either up or down. Nevertheless, also in this situation the growth exponents are close to their KPZ values, even if the fitness is declining. A detailed investigation of this model in the light of the analogy to nonequilibrium wetting processes appears to be an interesting problem for further research.

Acknowledgements

This work was supported by DFG within SFB 680 and SPP 1590. J.O. acknowledges support from The Halle Foundation. We thank Oskar Hallatschek and Stefan Boettcher for useful discussions.

References

- [1] KC Atwood, LK Schneider, and FJ Ryan. Periodic selection in *Escherichia coli*. *Proceedings of the National Academy of Sciences of the United States of America*, 37(3):146–155, 1951.
- [2] J Arjan G M de Visser and Daniel E Rozen. Clonal interference and the periodic selection of new beneficial mutations in *Escherichia coli*. *Genetics*, 172(4):2093–100, April 2006.
- [3] Michael M Desai, Daniel S Fisher, and Andrew W Murray. The speed of evolution and maintenance of variation in asexual populations. *Current biology : CB*, 17(5):385–94, March 2007.
- [4] Lília Perfeito, Lisete Fernandes, Catarina Mota, and Isabel Gordo. Adaptive mutations in bacteria: high rate and small effects. *Science (New York, N.Y.)*, 317(5839):813–5, August 2007.
- [5] Paul D Sniegowski and Philip J Gerrish. Beneficial mutations and the dynamics of adaptation in asexual populations. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 365(1544):1255–63, April 2010.
- [6] J. Arjan G M de Visser, Clifford W Zeyl, Philip J Gerrish, Jeffrey L Blanchard, and Richard E Lenski. Diminishing Returns from Mutation Supply Rate in Asexual Populations. *Science*, 283(5400):404–406, January 1999.
- [7] Aaron C Shaver, Peter G Dombrowski, Joseph Y Sweeney, Tania Treis, Renata M Zappala, and Paul D Sniegowski. Fitness evolution and the rise of mutator alleles in experimental *Escherichia coli* populations. *Genetics*, 162(2):557–566, 2002.
- [8] R.A. Fisher. *The Genetical Theory of Natural Selection*. Clarendon Press, Oxford, 1930.
- [9] Michael M Desai and Daniel S Fisher. Beneficial mutation selection balance and the effect of linkage on positive selection. *Genetics*, 176(3):1759–98, July 2007.

- [10] Igor M Rouzine, Eric Brunet, and Claus O Wilke. The traveling-wave approach to asexual evolution: Muller’s ratchet and speed of adaptation. *Theoretical population biology*, 73(1):24–46, February 2008.
- [11] Su-Chan Park, Damien Simon, and Joachim Krug. The Speed of Evolution in Large Asexual Populations. *Journal of Statistical Physics*, 138(1-3):381–410, January 2010.
- [12] Feng Yu, Alison Etheridge and Charles Cuthbertson. Asymptotic behavior of the rate of adaptation. *Annals of Applied Probability*, 20:978–1004 (2010).
- [13] Benjamin H Good, Igor M Rouzine, Daniel J Balick, Oskar Hallatschek, and Michael M Desai. Distribution of fixed beneficial mutations and the rate of adaptation in asexual populations. *Proceedings of the National Academy of Sciences of the United States of America*, 109(13):4950–5, March 2012.
- [14] Michael Kelly. Upper bound on the rate of adaptation of an asexual population. *Annals of Applied Probability*, 23:1377–1408 (2013).
- [15] H.J. Muller. The relation of recombination to mutational advance. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 1(1):2–9, May 1964.
- [16] John Haigh. The accumulation of deleterious genes in a population: Muller’s Ratchet. *Theoretical Population Biology*, 14(2):251–267, October 1978.
- [17] W Stephan, L Chao, and J G Smale. The advance of Muller’s ratchet in a haploid asexual population: approximate solutions based on diffusion theory. *Genetical research*, 61:225–231, 1993.
- [18] I Gordo and B Charlesworth. The degeneration of asexual haploid populations and the speed of Muller’s ratchet. *Genetics*, 154:1379–1387, 2000.
- [19] Isabel Gordo and Brian Charlesworth. On the Speed of Muller’s Ratchet. *Genetics*, 156(4):2137–2140, December 2000.
- [20] Kavita Jain. Loss of least-loaded class in asexual populations due to drift and epistasis. *Genetics*, 179(4):2125–34, August 2008.
- [21] Etheridge A, Pfaffelhuber P, and Wakolbinger A. How often does the ratchet click? Facts, heuristics, asymptotics. In Blath J, Mörters P, and Scheutzow M, editors, *Trends in Stochastic Analysis*, chapter 13, pages 365–390. Cambridge University Press, Cambridge, 2009.
- [22] D. Waxman and L. Loewe. A stochastic model for a single click of Muller’s ratchet. *Journal of Theoretical Biology*, 264(4):1120–1132, 2010.
- [23] Richard A Neher and Boris I Shraiman. Fluctuations of fitness distributions and the rate of Muller’s ratchet. *Genetics*, 191(4):1283–93, August 2012.

- [24] Jakob J Metzger and Stephan Eule. Distribution of the Fittest Individuals and the Rate of Muller’s Ratchet in a Model with Overlapping Generations. *PLoS computational biology*, 9(11):e1003303, November 2013.
- [25] H.J Muller. Some genetic aspects of sex. *Am. Nat.*, 66:118–138, 1932.
- [26] Brian Charlesworth. The effects of deleterious mutations on evolution at linked sites. *Genetics*, 190(1):5–22, January 2012.
- [27] S.-C. Park and J. Krug. Rate of adaptation in sexual and asexual populations: A solvable model of the Fisher-Muller effect. *Genetics*, 195:941–955, 2013.
- [28] R A Fisher. The wave of advance of advantageous genes. *Annals of Eugenics*, 7(Part 4):355–369, June 1937.
- [29] A N Kolmogorov, I G Petrovskii, and N S Piskunov. Study of the diffusion equation with growth of the quantity of matter and its application to a biological problem. *Moscow Univ Bull Math*, 1(1):1–25, 1937.
- [30] S Wright. Isolation by distance. *Genetics*, 28:114–138 (1943).
- [31] Motoo Kimura and GH Weiss. The stepping stone model of population structure and the decrease of genetic correlation with distance. *Genetics*, (480):561–576, 1964.
- [32] Gustave Malécot. Heterozygosity and relationship in regularly subdivided populations. *Theoretical Population Biology*, 8(2):212–241, 1975.
- [33] K. S. Korolev, Oskar Hallatschek, and David R. Nelson. Genetic demixing and evolution in linear stepping stone models. *Reviews of Modern Physics*, 82(2):1691–1718, May 2010.
- [34] Takeo Maruyama. On the fixation probability of mutant genes in a subdivided population. *Genetical Research*, 15(02):221, April 1970.
- [35] Takeo Maruyama. A simple proof that certain quantities are independent of the geographical structure of population. *Theoretical Population Biology*, 5(2):148–154, April 1974.
- [36] Isabel Gordo and Paulo R A Campos. Adaptive evolution in a spatially structured asexual population. *Genetica*, 127(1-3):217–29, May 2006.
- [37] L. Perfeito, I. Gordo, and P. R.a. Campos. The effect of spatial structure in adaptive evolution. *The European Physical Journal B*, 51(2):301–306, June 2006.
- [38] Jakub Otwinowski and Stefan Boettcher. Accumulation of beneficial mutations in one dimension. *Physical Review E*, 84(1):011925, July 2011.
- [39] Erik A Martens and Oskar Hallatschek. Interfering waves of adaptation promote spatial mixing. *Genetics*, 189(3):1045–60, November 2011.

- [40] Elder S. Claudino, M. L. Lyra, Iram Gleria, and Paulo R. A. Campos. Adaptive evolution on a continuous lattice model. *Physical Review E*, 87(3):032711, March 2013.
- [41] Cornelia Pokalyuk, Lisha A Mathew, Dirk Metzler, and Peter Pfaffelhuber. Competing islands limit the rate of adaptation in structured populations. *Theoretical population biology*, 90:1–11 (2013).
- [42] Albert-László Barabási and Harry Eugene Stanley. *Fractal concepts in surface growth*. Cambridge University Press, 1995.
- [43] Joachim Krug. Origins of scale invariance in growth processes. *Advances in Physics*, 46(2):139–282, April 1997.
- [44] Mehran Kardar, Giorgio Parisi, and Yi-Cheng Zhang. Dynamic Scaling of Growing Interfaces. *Physical Review Letters*, 56(9):889–892, 1986.
- [45] Thomas Kriecherbauer and Joachim Krug. A pedestrian’s view on interacting particle systems, KPZ universality and random matrices. *Journal of Physics A: Mathematical and Theoretical*, 43(40):403001, October 2010.
- [46] Kazumasa A Takeuchi, Masaki Sano, Tomohiro Sasamoto, and Herbert Spohn. Growing interfaces uncover universal fluctuations behind scale invariance. *Scientific reports*, 1:34, January 2011.
- [47] Ivan Corwin. The Kardar-Parisi-Zhang equation and universality class. *Random Matrices: Theory and Applications*, 01(01):1130001, January 2012.
- [48] C Doering, C Mueller, and P Smereka. Interacting particles, the stochastic Fisher-Kolmogorov-Petrovsky-Piscounov equation, and duality. *Physica A: Statistical Mechanics and its Applications*, 325(1-2):243–259, July 2003.
- [49] Oskar Hallatschek and K. Korolev. Fisher Waves in the Strong Noise Limit. *Physical Review Letters*, 103(10):108103, September 2009.
- [50] Oskar Hallatschek and David R Nelson. Life at the front of an expanding population. *Evolution*, 64(1):193–206, January 2010.
- [51] F.C. Frank. Nucleation-controlled growth on a one-dimensional growth of finite length. *Journal of Crystal Growth*, 22(3):233–236, 1974.
- [52] Nigel Goldenfeld. Kinetics of a model for nucleation-controlled polymer crystal growth. *Journal of Physics A: Mathematical and General*, 2807, 1984.
- [53] J Krug and H Spohn. Anomalous Fluctuations in the Driven and Damped Sine-Gordon Chain. *Europhysics Letters (EPL)*, 8(3):219–224, February 1989.
- [54] Michael Prähofer and Herbert Spohn. Universal Distributions for Growth Processes in 1+1 Dimensions and Random Matrices. *Physical Review Letters*, 84(21):4882–4885, May 2000.

- [55] F Family and T Vicsek. Scaling of the active zone in the Eden process on percolation networks and the ballistic deposition model. *Journal of Physics A: Mathematical and General*, 18(2):L75–L81, February 1985.
- [56] Craig A. Tracy and Harold Widom. Fredholm determinants, differential equations and matrix models. *Communications in Mathematical Physics*, 163(1):33–72, June 1994.
- [57] Jinho Baik, Percy Deift, and Kurt Johansson. On the distribution of the length of the longest increasing subsequence of random permutations. *Journal of the American Mathematical Society*, 12(4):1119–1178, 1999.
- [58] Tomohiro Sasamoto and Herbert Spohn. One-Dimensional Kardar-Parisi-Zhang Equation: An Exact Solution and its Universality. *Physical Review Letters*, 104(23):230602, June 2010.
- [59] Momar Dieng. RMLab, 2006.
- [60] Sidney Redner. *A Guide to First Passage Processes*. Cambridge University Press, 2001.
- [61] János Kertész and Dietrich E Wolf. Anomalous roughening in growth processes. *Physical Review Letters*, 62:2571–2574 (1989).
- [62] U. Alon, M. Evans, H. Hinrichsen, and D. Mukamel. Smooth phases, roughening transitions, and novel exponents in one-dimensional growth models. *Physical Review E*, 57(5):4997–5012, May 1998.
- [63] Haye Hinrichsen. Logarithmic roughening in a growth process with edge evaporation. *Physical Review E*, 67(1):016110, January 2003.
- [64] Andre Cardoso Barato. Nonequilibrium Wetting. *Journal of Statistical Physics*, 138(4-5):728–766, December 2009.
- [65] Malte Henkel, Haye Hinrichsen and Sven Lübeck. *Nonequilibrium Phase Transitions. Volume I: Absorbing Phase Transitions*. Springer, Dordrecht 2008.
- [66] Timothy Halpin-Healy. (2+1)-Dimensional Directed Polymer in a Random Medium: Scaling Phenomena and Universal Distributions. *Physical Review Letters*, 109(17):170602, October 2012.
- [67] Erik A Martens, Rumen Kostadinov, Carlo C Maley, and Oskar Hallatschek. Spatial structure increases the waiting time for cancer. *New journal of physics*, 13(11):115014, November 2011.