

A NEW CRITICAL CURVE FOR THE LANE-EMDEN SYSTEM

WENJING CHEN, LOUIS DUPAIGNE, AND MARIUS GHERGU

ABSTRACT. We study stable positive radially symmetric solutions for the Lane-Emden system $-\Delta u = v^p$ in \mathbb{R}^N , $-\Delta v = u^q$ in \mathbb{R}^N , where $p, q \geq 1$. We obtain a new critical curve that optimally describes the existence of such solutions.

1. INTRODUCTION

We consider the Lane-Emden system

$$(1.1) \quad \begin{cases} -\Delta u = v^p, u > 0 & \text{in } \mathbb{R}^N, \\ -\Delta v = u^q, v > 0 & \text{in } \mathbb{R}^N, \end{cases}$$

where $N \geq 1$ and $p \geq q > 0$. Introduced independently by Mitidieri [14] and Van der Vorst [21], the Sobolev critical hyperbola plays a crucial role in the analysis of (1.1). In particular, Mitidieri [15] (see also Serrin and Zou [18]) proved that (1.1) has a nontrivial radially symmetric solution if and only if (p, q) lies on or above the hyperbola i.e. when

$$(1.2) \quad \frac{1}{p+1} + \frac{1}{q+1} \leq 1 - \frac{2}{N}.$$

The Lane-Emden conjecture states that such a result should continue to hold for any positive solution (not necessarily radially symmetric). See Souplet [19] and the references therein for the progress on this conjecture.

In this paper we characterize the stability of radially symmetric solutions of the Lane-Emden system (1.1), the definition of which we recall now.

Definition 1.1. *A solution (u, v) to (1.1) is stable if there exists a positive supersolution of the linearized system i.e. if there exists $(\phi, \psi) \in C^2(\mathbb{R}^N)^2$ such that*

$$\begin{cases} -\Delta \phi \geq pv^{p-1}\psi & \text{in } \mathbb{R}^N, \\ -\Delta \psi \geq qu^{q-1}\phi & \text{in } \mathbb{R}^N, \\ \phi, \psi > 0 & \text{in } \mathbb{R}^N. \end{cases}$$

Let us also recall that if (1.2) holds, then

$$(1.3) \quad (u_s, v_s) = (a|x|^{-\alpha}, b|x|^{-\beta}), \quad x \in \mathbb{R}^N \setminus \{0\}$$

is a weak solution of (1.1) provided

$$(1.4) \quad \alpha = \frac{2(p+1)}{pq-1}, \quad \beta = \frac{2(q+1)}{pq-1}$$

and $a = (ST^p)^{\frac{1}{pq-1}}$, $b = (S^qT)^{\frac{1}{pq-1}}$, $S = \alpha(N-2-\alpha)$, $T = \beta(N-2-\beta)$.

Our main result states that the stability of a radial solution of the Lane-Emden system is determined by the position of the exponents (p, q) with respect to a new critical curve, which we christen ‘‘Joseph and Lundgren’’, since the exponent introduced by these authors in [11] is the intersection of the curve with the diagonal $p = q$.

Theorem 1.2. *Assume $p \geq q \geq 1$.*

- (i) *If $N \geq 11$ and (p, q) lies on or above the Joseph-Lundgren critical curve i.e.*

$$(1.5) \quad \left[\frac{(N-2)^2 - (\alpha - \beta)^2}{4} \right]^2 \geq pq\alpha\beta(N-2-\alpha)(N-2-\beta),$$

then any radially symmetric solution (u, v) of (1.1) is stable and satisfies

$$u < u_s \quad \text{and} \quad v < v_s \quad \text{in } \mathbb{R}^N \setminus \{0\},$$

where (u_s, v_s) is the singular solution given by (1.3) and α, β are the scaling exponents given by (1.4).

- (ii) *If $N \leq 10$ or if $N \geq 11$ and (1.5) fails, then there is no stable radially symmetric solution of (1.1).*

Remark 1.3. Equation (1.5) is derived by studying the stability of the singular solution (u_s, v_s) given by (1.3).

Remark 1.4.

- The above theorem was first proved by Cowan for $1 \leq N \leq 10$, $p \geq q \geq 2$ and (u, v) not necessarily radial. See [4].
- In the case $p = q$, using Remarks 1.1(a) and 2.1(a) of Souplet [19] and Farina’s seminal work for the case of a single equation [9], part (ii) of the theorem readily follows. The result continues to hold for possibly nonradial solutions, assumed to be stable only outside a compact set.
- In the biharmonic case $q = 1$, the theorem was first proved by Karageorgis [13] using the asymptotics found by Gazzola and Grunau in [12].
- In all the other cases, only partial results were known. To the authors knowledge, the state of the art for nonradial solutions is contained in the following references: Wei and D. Ye [23], Wei, Xu and Yang [22], Hajlaoui, A. Harrabi and D. Ye [10] for the biharmonic case, and Cowan [4] for the general case. We believe that the methods of the paper [8] by Goubet, Warnault and two of the authors should




FIGURE 1. The stable region (shaded) for radially symmetric solutions of the Lane-Emden system (1.1).

slightly improve the known results (and coincide with [10] in the biharmonic case).

- Our result does not cover the case where one of the exponents is less than 1.
- The left hand-side in (1.5) is related to the following Hardy-Rellich inequality :

$$(1.6) \quad \int_{\mathbb{R}^N} |x|^{2-\gamma} |\Delta \varphi|^2 dx \geq C_\gamma \int_{\mathbb{R}^N} |x|^{-2-\gamma} \varphi^2 dx.$$

The optimal constant C_γ in the class of radially symmetric functions $\varphi = \varphi(|x|)$ is given by

$$(1.7) \quad C_\gamma = \inf_{\substack{\varphi \in C_c^\infty(\mathbb{R}^N \setminus \{0\}) \\ 0 \neq \varphi = \varphi(|x|)}} \frac{\int_{\Omega} |x|^{2-\gamma} |\Delta \varphi|^2 dx}{\int_{\Omega} |x|^{-2-\gamma} \varphi^2 dx} = \left[\frac{(N-2)^2 - \gamma^2}{4} \right]^2,$$

and the above infimum is never achieved. See Caldiroli and Musina [2]. We remark that the optimal constant C_γ in (1.7) corresponds to the left hand-side in (1.5) with $\gamma = \alpha - \beta \in [0, 2)$.

As an immediate corollary of Theorem 1.2 and standard blow-up analysis, we obtain the following regularity result.

Corollary 1.5. *Let B denote the unit ball of \mathbb{R}^N , $N \geq 1$, $\lambda, \mu > 0$. Let $f, g \in C^1(\mathbb{R})$ be two nondecreasing functions such that $f(0) \geq 0$, $g(0) > 0$, $f'(0)g'(0) > 0$ and*

$$\lim_{t \rightarrow +\infty} \frac{f'(t)}{t^{p-1}} = a, \quad \lim_{t \rightarrow +\infty} \frac{g'(t)}{t^{q-1}} = b$$

for some $a, b > 0$, $p \geq q \geq 1$, $pq > 1$. Then, any extremal solution to the system

$$(1.8) \quad \begin{cases} -\Delta u = \lambda f(v), & u > 0 & \text{in } B, \\ -\Delta v = \mu g(u), & v > 0 & \text{in } B, \\ u = v = 0 & & \text{on } \partial B \end{cases}$$

is bounded if either $N \leq 10$ or if $N \geq 11$ and (p, q) lies below the Joseph-Lundgren critical curve i.e.(1.5) fails.

For the notion of extremal solution for systems, we refer to Montenegro [16]. See also Cowan [3] for partial results on general domains. The proof is a straightforward adaptation of Theorem 1.8 in [5], using the version of the blow-up technique introduced by Polacik, Quittner and Souplet [17], so we skip it.

2. PRELIMINARY RESULTS

The following three results will serve for the purpose of comparing solutions. In the lemma below, we say that a solution is strictly stable in a bounded region $\Omega \subset \mathbb{R}^N$ if the principle eigenvalue of the linearized equation in Ω is strictly positive.

Lemma 2.1. *Let $(u, v) \in C^2(\mathbb{R}^N)^2$ be a stable solution of (1.1). Then, given any bounded domain $\Omega \subset \mathbb{R}^N$, (u, v) is strictly stable in Ω . In particular, the linearized operator satisfies the maximum principle, that is, any pair $(\phi, \psi) \in C^2(\overline{\Omega})^2$ such that*

$$\begin{cases} -\Delta \phi \geq p v^{p-1} \psi & \text{in } \Omega, \\ -\Delta \psi \geq q u^{q-1} \phi & \text{in } \Omega, \\ \phi, \psi \geq 0 & \text{on } \partial \Omega, \end{cases}$$

satisfies $\phi, \psi \geq 0$ in Ω .

Proof. Since (u, v) is stable in \mathbb{R}^N , the linearized equation has a strict supersolution in Ω . As observed by Sweers [20] and Busca-Sirakov [1], this implies in turn that the principle eigenvalue of the linearized operator in Ω is strictly positive and equivalently that the maximum principle holds. \square

In the next lemma, we say that a solution is minimal if it lies below any (local) supersolution of the same equation. See e.g. [7] for the notion of minimal solution.

Lemma 2.2. *Assume $p \geq q \geq 1$. Let $a, b, R > 0$. If $(u, v) \in C^2(\overline{B_R})^2$ is a strictly stable solution of*

$$(2.1) \quad \begin{cases} -\Delta u = v^p & \text{in } B_R, \\ -\Delta v = u^q & \text{in } B_R, \\ u = a, v = b & \text{on } \partial B_R, \end{cases}$$

then (u, v) is minimal.

Proof. Assume that (u, v) is a strictly stable solution of (2.1). By the maximum principle, $u \geq a, v \geq b$ in B_R . In particular, there exists the minimal solution (u_m, v_m) and $u \geq u_m \geq a, v \geq v_m \geq b$ in B_R .

Set $\phi = u - u_m, \psi = v - v_m$. Then, $\phi, \psi \geq 0$ in B_R and, since $p \geq q \geq 1$,

$$\begin{cases} -\Delta \phi = v^p - v_m^p \leq p v^{p-1} \psi & \text{in } B_R, \\ -\Delta \psi = u^q - u_m^q \leq q u^{q-1} \phi & \text{in } B_R, \\ \phi = \psi = 0 & \text{on } \partial B_R. \end{cases}$$

Since (u, v) is strictly stable, the maximum principle holds and implies that $\phi, \psi \leq 0$ in B_R . It follows that $\phi \equiv \psi \equiv 0$, that is, $u = u_m$ and $v = v_m$. \square

As an immediate consequence of the two previous lemmas, we obtain

Corollary 2.3. *Let (u, v) be a stable solution of (1.1) in \mathbb{R}^N and let (u_s, v_s) be the singular solution defined by (1.3). If there exists $R > 0$ such that $u(R) \leq u_s(R)$ and $v(R) \leq v_s(R)$, then*

$$u < u_s \quad \text{and} \quad v < v_s \quad \text{in } B_R \setminus \{0\}.$$

2.1. Stability of the singular solution. In this part we investigate the stability of the singular solution (u_s, v_s) given by (1.3).

Proposition 2.4. *The following are equivalent:*

- (i) *The singular solution (u_s, v_s) is stable in $\mathbb{R}^N \setminus \{0\}$;*
- (ii) *The singular solution (u_s, v_s) is stable outside of some compact set;*
- (iii) *(p, q) satisfies (1.5).*

Proof. Since the implication (i) \Rightarrow (ii) is trivial, we only need to prove the implications

$$(ii) \Rightarrow (iii) \Rightarrow (i)$$

Assume first that (ii) holds, that is, the singular solution (u_s, v_s) is stable outside of a compact set. Thus, (u_s, v_s) is stable in $\mathbb{R}^N \setminus \overline{B_r}$ for some $r > 0$. By scale invariance, (u_s, v_s) is stable in $\mathbb{R}^N \setminus \overline{B_\rho}$ for all $\rho > 0$.

Set $\gamma = \alpha - \beta$, where α, β are the scaling exponents given by (1.4) and let K_1, K_2 be the constants such that

$$p v_s^{p-1} = K_1 |x|^{-2+\gamma} \quad \text{and} \quad q u_s^{q-1} = K_2 |x|^{-2-\gamma}.$$

Then, (p, q) satisfies (1.5) if and only if

$$C_\gamma \geq K_1 K_2,$$

where C_γ is given by (1.7). Assume by contradiction that (p, q) does not satisfy (1.5). Then, we may find an open annular region $\Omega = B_{R_1} \setminus \overline{B_{R_2}}$ such that

$$(2.2) \quad \lambda := \min_{\varphi \in H \setminus \{0\}} \frac{\int_{\Omega} |x|^{2-\gamma} |\Delta \varphi|^2 dx}{\int_{\Omega} |x|^{-2-\gamma} \varphi^2 dx} < K_1 K_2,$$

where H is the space of radial functions φ such that $\int_{\Omega} |x|^{2-\gamma} |\Delta \varphi|^2 dx < +\infty$ and $\varphi = 0$ on $\partial\Omega$. Let $\varphi > 0$ be a minimizer of (2.2), so that letting $\psi = |x|^{2-\gamma} (-\Delta \varphi)$, we have

$$\begin{cases} -\Delta \varphi = |x|^{-2+\gamma} \psi, \varphi > 0 & \text{in } \Omega, \\ -\Delta \psi = \lambda |x|^{-2-\gamma} \varphi, \psi > 0 & \text{in } \Omega, \\ \varphi = \psi = 0 & \text{on } \partial\Omega. \end{cases}$$

Since (u_s, v_s) is strictly stable in Ω , thanks to [20, Theorem 1.1], there also exists $(\tilde{\varphi}, \tilde{\psi}) \in C^2(\overline{\Omega})^2$ such that

$$\begin{cases} -\Delta \tilde{\varphi} = K_1 |x|^{-2+\gamma} \tilde{\psi}, \tilde{\varphi} > 0 & \text{in } \Omega, \\ -\Delta \tilde{\psi} = K_2 |x|^{-2-\gamma} \tilde{\varphi} + 1, \tilde{\psi} > 0 & \text{in } \Omega, \\ \tilde{\varphi} = \tilde{\psi} = 0 & \text{on } \partial\Omega. \end{cases}$$

A straightforward integration by part shows that φ and $\tilde{\varphi}$ satisfy

$$\langle \varphi, \tilde{\varphi} \rangle := \int_{\Omega} |x|^{2-\gamma} \Delta \varphi \Delta \tilde{\varphi} dx \leq 0$$

which is impossible, since both ψ and $\tilde{\psi}$ are positive. Hence (p, q) satisfies (1.5) and we have proved that (ii) implies (iii).

Assume now (iii). It is easy to see that

$$(2.3) \quad \phi(x) = \frac{4K_1}{(N-2-\gamma)(N-2+\gamma)} |x|^{-\frac{N-2-\gamma}{2}}, \quad \psi(x) = |x|^{-\frac{N-2+\gamma}{2}}$$

satisfy

$$(2.4) \quad \begin{aligned} -\Delta \phi &= p v_s^{p-1} \psi \\ -\Delta \psi &\geq q u_s^{q-1} \phi \end{aligned}$$

in $\mathbb{R}^N \setminus \{0\}$, which means that (u_s, v_s) is stable in $\mathbb{R}^N \setminus \{0\}$. \square

3. PROOF OF THEOREM 1.2

Assume (p, q) satisfies (1.5). Then by Proposition 2.4, the singular solution (u_s, v_s) is stable in $\mathbb{R}^N \setminus \{0\}$. Theorem 1.2(i) follows from the proposition below.

Proposition 3.1. *Assume (p, q) satisfies (1.5). Then, for any radially symmetric solution (u, v) of (1.1) we have*

$$(3.1) \quad u < u_s \quad \text{and} \quad v < v_s \quad \text{in } \mathbb{R}^N \setminus \{0\}.$$

Proof. Assume by contradiction that there exists a radially symmetric solution (u, v) of (1.1) for which (3.1) fails to hold and set

$$U = u_s - u, \quad V = v_s - v.$$

Since (3.1) is not fulfilled, U' and V' must change sign in $(0, \infty)$. Indeed, otherwise $U' < 0$ or $V' < 0$ in $(0, \infty)$ which implies (since $U(\infty) = V(\infty) = 0$) that $u_s \geq u$ or $v_s \geq v$ in $(0, \infty)$. Now, the maximum principle yields $u_s \geq u$ and $v_s \geq v$ in $(0, \infty)$ and this contradicts our assumption.

Let $r_1 > 0$ (resp. $r_2 > 0$) be the first zero of U' (resp. V'). Thus

$$U' < 0 \text{ in } (0, r_1), \quad U'(r_1) = 0, \quad V' < 0 \text{ in } (0, r_2), \quad V'(r_2) = 0.$$

Without losing the generality, we may assume $r_2 \geq r_1$. Set next

$$r_3 := \inf\{r > 0 : V(r) < 0\} \in (0, \infty]$$

and we claim that $r_3 < r_1$. If $r_3 \geq r_1$ then $V > 0$ in $(0, r_1)$ which means

$$(3.2) \quad v < v_s \quad \text{in } (0, r_1).$$

Integrating in (1.1) and using (3.2) we find

$$(r^{N-1}u')' = -r^{N-1}v^p > -r^{N-1}v_s^p = (r^{N-1}u_s')' \quad \text{in } (0, r_1).$$

Integrating the above inequality over $[0, r_1]$ we find $u'(r_1) > u_s'(r_1)$ which contradicts $U'(r_1) = 0$. Hence $r_3 \in (0, r_1)$. Similarly we define

$$r_4 := \inf\{r > 0 : U(r) < 0\} \in (0, \infty]$$

and as before we deduce $r_4 \in (0, r_2)$. In fact, we show that $r_4 \leq r_1$. Assuming the contrary, that is, $r_4 > r_1$, we find $r_1 < r_4 < r_2$. Further, since $V' < 0$ in $(0, r_2)$ we deduce $V(r) < V(r_3) = 0$ for all $r \in (r_3, r_2)$ so $v_s < v$ in (r_3, r_2) . Therefore,

$$(r^{N-1}u')' = -r^{N-1}v^p < -r^{N-1}v_s^p = (r^{N-1}u_s')' \quad \text{in } (r_3, r_2).$$

Integrating over $[r_1, r]$, $r_1 < r < r_2$, and using $U'(r_1) = 0$ we obtain $u'(r) < u_s'(r)$ for all $r \in (r_1, r_2)$. This means that U is increasing in (r_1, r_2) . In particular, $U(r_1) < U(r_4) = 0$. On the other hand, from the definition of r_4 we have $U(r_1) > 0$, contradiction. We have thus obtained $r_3 < r_1$, $r_4 \leq r_1 \leq r_2$ which yield

$$(3.3) \quad U(r_1) \leq 0, \quad U'(r_1) = 0, \quad V(r_1) < 0, \quad V'(r_1) \leq 0.$$

Next, let (ϕ, ψ) be defined by (2.3) and recall that (ϕ, ψ) solves the linearized equation (2.4) in $\mathbb{R}^N \setminus \{0\}$. Also, since $p \geq q \geq 1$, (U, V) satisfies

$$(3.4) \quad \begin{cases} -\Delta U \leq p v_s^{p-1} V \\ -\Delta V \leq q u_s^{q-1} U \end{cases} \quad \text{in } \mathbb{R}^N \setminus \{0\}.$$

We multiply the equations in (2.4) by V and U , and the two equations in (3.4) by ψ and ϕ respectively. Integrating over B_r , $r > 0$, we find

$$\int_{B_r} (-\Delta U)\psi \leq \int_{B_r} (-\Delta\phi)V \quad \text{and} \quad \int_{B_r} (-\Delta V)\phi \leq \int_{B_r} (-\Delta\psi)U.$$

Adding the above inequalities we deduce

$$\int_{B_r} (V\Delta\phi - \phi\Delta V) + \int_{B_r} (U\Delta\psi - \psi\Delta U) \leq 0 \quad \text{for all } r > 0,$$

that is,

$$\int_{\partial B_r} \left(V \frac{\partial\phi}{\partial\nu} - \phi \frac{\partial V}{\partial\nu} \right) + \int_{\partial B_r} \left(U \frac{\partial\psi}{\partial\nu} - \psi \frac{\partial U}{\partial\nu} \right) \leq 0 \quad \text{for all } r > 0.$$

Since U, V, ϕ, ψ are radially symmetric, this yields

$$(3.5) \quad V\phi' - \phi V' + U\psi' - \psi U' \leq 0 \quad \text{in } (0, \infty).$$

Now, let us remark that $\phi, \psi > 0$ and $\phi', \psi' < 0$ in $(0, \infty)$. Combining this fact with (3.3) we deduce that (3.5) does not hold at $r = r_1$, a contradiction. Hence $u < u_s$ and $v < v_s$ in $\mathbb{R}^N \setminus \{0\}$. \square

Assume next that (1.5) fails to hold. We establish first the following result.

Proposition 3.2. *Assume (p, q) does not satisfy (1.5). Then, for any stable solution (u, v) of (1.1) we have*

$$u < u_s \quad \text{and} \quad v < v_s \quad \text{in } \mathbb{R}^N \setminus \{0\}.$$

Proof. Assume by contradiction that $u - u_s$ changes sign in $\mathbb{R}^N \setminus \{0\}$. Then $v - v_s$ also changes sign in $\mathbb{R}^N \setminus \{0\}$ for otherwise $v - v_s \leq 0$ in $\mathbb{R}^N \setminus \{0\}$ implies

$$-\Delta(u - u_s) = v^p - v_s^p \leq 0 \quad \text{in } \mathbb{R}^N \setminus \{0\}.$$

Also $u - u_s < 0$ in a neighborhood of the origin and $u(x) - u_s(x) \rightarrow 0$ as $|x| \rightarrow \infty$. By the maximum principle, we deduce $u - u_s \leq 0$ in $\mathbb{R}^N \setminus \{0\}$ which contradicts our assumption.

Hence $u - u_s$ and $v - v_s$ change sign on $(0, \infty)$. Denote by r_1 (resp. r_2) the first sign-changing zero of $u - u_s$ (resp. $v - v_s$). From Corollary 2.3, $u - u_s$ (resp. $v - v_s$) cannot be zero in a whole neighborhood of r_1 (resp. r_2). Without losing generality, we may assume that $r_1 \leq r_2$.

We claim that $u - u_s$ has a second sign-changing point $r_3 > r_1$. Indeed, otherwise $u - u_s \geq 0$ in $\mathbb{R}^N \setminus B_{r_1}$ which by the maximum principle implies that $v - v_s \geq 0$ in $\mathbb{R}^N \setminus B_{r_2}$. Therefore, $u \geq u_s, v \geq v_s$ in $\mathbb{R}^N \setminus B_{r_2}$ which implies that (u_s, v_s) is a stable solution of (1.1) in $\mathbb{R}^N \setminus B_{r_2}$ and thus, contradicts Proposition 2.4. Hence, there exists $r_3 > r_1$ a second sign-changing point of $u - u_s$. Further, we must have $r_3 \geq r_2$ for otherwise $r_1 < r_3 < r_2$. Then $u(r_3) = u_s(r_3)$ and $v(r_3) < v_s(r_3)$ which by Corollary 2.3 yields $u < u_s, v < v_s$ in $B_{r_3} \setminus \{0\}$. But this is impossible since $u(r_1) = u_s(r_1)$. Thus, $r_3 \geq r_2$.

We next claim that $v - v_s$ has a second sign-changing point $r_4 > r_2$. As before, if this is not true, then $v - v_s \geq 0$ in $\mathbb{R}^N \setminus B_{r_2}$ and by the maximum principle we find $u - u_s \geq 0$ in $\mathbb{R}^N \setminus B_{r_3}$. Then $u \geq u_s, v \geq v_s$ in $\mathbb{R}^N \setminus B_{r_3}$, so (u_s, v_s) is stable in $\mathbb{R}^N \setminus B_{r_3}$ which contradicts Proposition 2.4.

We show next that $r_4 \geq r_3$. Assuming the contrary we have $r_2 < r_4 < r_3$. At this stage, two cases may occur:

CASE 1: $v \leq v_s$ in (r_4, r_3) . Remark that $u(r_3) = u_s(r_3)$ and $v(r_3) \leq v_s(r_3)$. By Corollary 2.3 we deduce $u < u_s$ in B_{r_3} which is impossible since $u(r_1) = u_s(r_1)$.

CASE 2: $v - v_s$ has a third sign-changing point $\rho \in (r_4, r_3)$. Then $v - v_s > 0$ on (r_2, r_4) and $v - v_s < 0$ on (r_4, ρ) . On the other hand,

$$-\Delta(v - v_s) = u^q - u_s^q \geq 0 \quad \text{in } B_\rho \setminus \overline{B}_{r_4}$$

and $v - v_s = 0$ on $\partial(B_\rho \setminus B_{r_4})$. The maximum principle yields $v - v_s > 0$ on (r_4, ρ) , a contradiction. We have proved that $r_4 \geq r_3$.

We claim that $u - u_s$ has a third sign-changing point $r_5 > r_3$. Indeed, if this is not true, then $u - u_s \leq 0$ in $\mathbb{R}^N \setminus B_{r_3}$ and by the maximum principle we have $v - v_s \leq 0$ in $\mathbb{R}^N \setminus B_{r_4}$. Hence $u \leq u_s, v \leq v_s$ in $\mathbb{R}^N \setminus B_{r_4}$ which combined with Corollary 2.3 produces $u < u_s, v < v_s$ in B_{r_4} . This is clearly impossible since $u(r_1) = u_s(r_1)$. Hence, $u - u_s$ has a third sign-changing point $r_5 > r_3$.

If $r_5 \leq r_4$ then

$$-\Delta(u - u_s) = v^p - v_s^p \geq 0 \quad \text{in } B_{r_5} \setminus \overline{B}_{r_3}$$

and $u - u_s = 0$ on $\partial(B_{r_5} \setminus B_{r_3})$. By the maximum principle we infer that $u - u_s \geq 0$ in $B_{r_5} \setminus B_{r_3}$ which implies $u - u_s \geq 0$ in $B_{r_5} \setminus B_{r_1}$. This contradicts the fact that $r_3 \in (r_1, r_5)$ is a sign-changing point of $u - u_s$.

If $r_5 > r_4$ then $u(r_4) \leq u_s(r_4)$ and $v(r_4) = v_s(r_4)$. By Corollary 2.3 we deduce $u < u_s, v < v_s$ in B_{r_4} which is again a contradiction. \square

We are now ready to complete the proof of Theorem 1.2(ii). We adapt an idea introduced in [6]. Assume there exists a positive stable radially symmetric solution (u, v) of (1.1) and set

$$M_1 = \sup_{r \in (0, \infty)} \frac{u(r)}{u_s(r)}, \quad M_2 = \sup_{r \in (0, \infty)} \frac{v(r)}{v_s(r)}.$$

By Proposition 3.2 we have $M_1, M_2 \leq 1$. Since $\lim_{r \rightarrow \infty} u(r) = 0$, u coincides with the Newtonian potential of v^p . Hence

$$\begin{aligned} u(x) &= c_N \int_{\mathbb{R}^N} |x - y|^{2-N} v^p(y) dy \\ &\leq M_2^p \left\{ c_N \int_{\mathbb{R}^N} |x - y|^{2-N} v_s^p(y) dy \right\} = M_2^p u_s(x). \end{aligned}$$

Thus, $M_1 \leq M_2^p$ and similarly $M_2 \leq M_1^q$. It follows that $M_1 \leq M_1^{pq}$. So, since $pq > 1$ we have either $M_1 = 0$ or $M_1 = 1$. If $M_1 = 0$ then $u \equiv 0$

and this yields $v \equiv 0$ which is impossible. Therefore $M_1 = 1$ and similarly $M_2 = 1$, i.e.

$$\sup_{r \in (0, \infty)} \frac{u(r)}{u_s(r)} = \sup_{r \in (0, \infty)} \frac{v(r)}{v_s(r)} = 1.$$

By the strong maximum principle, (u, v) cannot touch (u_s, v_s) , so there exists a sequence $\{R_k\}$ converging to $+\infty$ such that

$$(3.6) \quad \lim_{k \rightarrow \infty} \frac{u(R_k)}{u_s(R_k)} = 1.$$

Define

$$u_k(r) = R_k^\alpha u(R_k r), \quad v_k(r) = R_k^\beta v(R_k r) \quad r \geq 0.$$

By scale invariance we have

$$(3.7) \quad 0 < u_k < u_s, \quad 0 < v_k < v_s \quad \text{in } \mathbb{R}^N \setminus \{0\}$$

and (u_k, v_k) solves the Lane-Emden system (1.1). By elliptic regularity, $\{(u_k, v_k)\}$ converges uniformly in $C_{loc}^2(\mathbb{R}^N)$ to a solution (\tilde{u}, \tilde{v}) of (1.1) which, in view of (3.7), also satisfies

$$0 \leq \tilde{u} \leq u_s, \quad 0 \leq \tilde{v} \leq v_s \quad \text{in } \mathbb{R}^N \setminus \{0\}.$$

Let us remark that by (3.6) we have

$$\tilde{u}(1) = \lim_{k \rightarrow \infty} u_k(1) = \lim_{k \rightarrow \infty} R_k^\alpha u(R_k) = \lim_{k \rightarrow \infty} R_k^\alpha u_s(R_k) = u_s(1).$$

On the other hand,

$$\begin{cases} -\Delta(\tilde{u} - u_s) = \tilde{v}^p - v_s^p \leq 0 & \text{in } \mathbb{R}^N \setminus \{0\}, \\ \lim_{|x| \rightarrow 0} (\tilde{u} - u_s) \leq 0, \quad \lim_{|x| \rightarrow \infty} (\tilde{u} - u_s) \leq 0. \end{cases}$$

By the strong maximum principle we deduce that $\tilde{u} \equiv u_s$. This is impossible, since \tilde{u} is bounded while u_s is singular.

ACKNOWLEDGEMENT

The first named author was partially supported by the Ecos-Conycit grant C09E06 between France and Chile. The second named author wishes to thank J. Wei and the math department at the Chinese University of Hong Kong, where part of this work was completed. The research of the third named author has been supported by a Ulysses project between France and Ireland.

REFERENCES

- [1] Jérôme Busca and Boyan Sirakov, *Harnack type estimates for nonlinear elliptic systems and applications*, Ann. Inst. H. Poincaré Anal. Non Linéaire **21** (2004), no. 5, 543–590, DOI 10.1016/j.anihpc.2003.06.001 (English, with English and French summaries). MR2086750 (2005g:35071)
- [2] Paolo Caldiroli and Roberta Musina, *Rellich inequalities with weights*, Calc. Var. Partial Differential Equations **45** (2012), no. 1-2, 147–164, DOI 10.1007/s00526-011-0454-3. MR2957654

- [3] Craig Cowan, *Regularity of stable solutions of a Lane-Emden type system*, <http://arxiv.org/abs/1206.4273> (19 june 2012).
- [4] ———, *Liouville theorems for stable Lane-Emden systems and biharmonic problems*, <http://arxiv.org/abs/1207.1081> (4 july 2012).
- [5] Juan Dávila, Louis Dupaigne, and Alberto Farina, *Partial regularity of finite Morse index solutions to the Lane-Emden equation*, *J. Funct. Anal.* **261** (2011), no. 1, 218–232, DOI 10.1016/j.jfa.2010.12.028. MR2785899
- [6] Juan Dávila, Louis Dupaigne, and Marcelo Montenegro, *The extremal solution of a boundary reaction problem*, *Commun. Pure Appl. Anal.* **7** (2008), no. 4, 795–817, DOI 10.3934/cpaa.2008.7.795.
- [7] Serge Dumont, Louis Dupaigne, Olivier Goubet, and Vicentiu Rădulescu, *Back to the Keller-Osserman condition for boundary blow-up solutions*, *Adv. Nonlinear Stud.* **7** (2007), no. 2, 271–298. MR2308040 (2008e:35062)
- [8] Louis Dupaigne, Marius Ghergu, Olivier Goubet, and Guillaume Warnault, *The Gelfand problem for the biharmonic operator*, to appear in *Arch. Ration. Mech. Anal.*
- [9] Alberto Farina, *On the classification of solutions of the Lane-Emden equation on unbounded domains of \mathbb{R}^N* , *J. Math. Pures Appl.* (9) **87** (2007), no. 5, 537–561, DOI 10.1016/j.matpur.2007.03.001 (English, with English and French summaries). MR2322150 (2008c:35070)
- [10] H. Hajlaoui, A. Harrabi, and D. Ye, *On stable solutions of biharmonic problem with polynomial growth*, arXiv:1211.2223.
- [11] D. D. Joseph and T. S. Lundgren, *Quasilinear Dirichlet problems driven by positive sources*, *Arch. Rational Mech. Anal.* **49** (1972/73), 241–269. MR0340701 (49 #5452)
- [12] Filippo Gazzola and Hans-Christoph Grunau, *Radial entire solutions for supercritical biharmonic equations*, *Math. Ann.* **334** (2006), no. 4, 905–936, DOI 10.1007/s00208-005-0748-x. MR2209261 (2007b:35114)
- [13] Paschalis Karageorgis, *Stability and intersection properties of solutions to the nonlinear biharmonic equation*, *Nonlinearity* **22** (2009), no. 7, 1653–1661, DOI 10.1088/0951-7715/22/7/009. MR2519683 (2011a:35240)
- [14] Enzo Mitidieri, *A Rellich type identity and applications*, *Comm. Partial Differential Equations* **18** (1993), no. 1-2, 125–151, DOI 10.1080/03605309308820923. MR1211727 (94c:26016)
- [15] ———, *Nonexistence of positive solutions of semilinear elliptic systems in \mathbf{R}^N* , *Differential Integral Equations* **9** (1996), no. 3, 465–479. MR1371702 (96m:35087)
- [16] Marcelo Montenegro, *Minimal solutions for a class of elliptic systems*, *Bull. London Math. Soc.* **37** (2005), no. 3, 405–416, DOI 10.1112/S0024609305004248. MR2131395 (2005k:35099)
- [17] Peter Poláčik, Pavol Quittner, and Philippe Souplet, *Singularity and decay estimates in superlinear problems via Liouville-type theorems. I. Elliptic equations and systems*, *Duke Math. J.* **139** (2007), no. 3, 555–579, DOI 10.1215/S0012-7094-07-13935-8. MR2350853 (2009b:35131)
- [18] James Serrin and Henghui Zou, *Non-existence of positive solutions of semilinear elliptic systems*, A tribute to Ilya Bakelman (College Station, TX, 1993), *Discourses Math. Appl.*, vol. 3, Texas A & M Univ., College Station, TX, 1994, pp. 55–68. MR1423369 (97k:35065)
- [19] Philippe Souplet, *The proof of the Lane-Emden conjecture in four space dimensions*, *Adv. Math.* **221** (2009), no. 5, 1409–1427, DOI 10.1016/j.aim.2009.02.014. MR2522424 (2010h:35088)
- [20] Guido Sweers, *Strong positivity in $C(\overline{\Omega})$ for elliptic systems*, *Math. Z.* **209** (1992), no. 2, 251–271, DOI 10.1007/BF02570833. MR1147817 (92j:35046)
- [21] R. C. A. M. Van der Vorst, *Variational identities and applications to differential systems*, *Arch. Rational Mech. Anal.* **116** (1992), no. 4, 375–398, DOI 10.1007/BF00375674. MR1132768 (93d:35043)

- [22] Juncheng Wei, Xingwang Xu, and Wen Yang, *Classification of stable solutions to biharmonic problems in large dimensions*. <http://www.math.cuhk.edu.hk/~wei/publicationpreprint.html>
- [23] Juncheng Wei and Dong Ye, *Liouville Theorems for finite Morse index solutions of Biharmonic problem*. <http://www.math.cuhk.edu.hk/~wei/publicationpreprint.html>

DEPARTAMENTO DE INGENIERÍA MATEMÁTICA AND CMM, UNIVERSIDAD DE CHILE,
CASILLA 170 CORREO 3, SANTIAGO, CHILE
E-mail address: wjchen1102@yahoo.com.cn

LAMFA, UMR CNRS 7352, UNIVERSITÉ DE PICARDIE JULES VERNE, 33 RUE ST
LEU, 80039, AMIENS CEDEX, FRANCE
E-mail address: louis.dupaigne@math.cnrs.fr

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY COLLEGE DUBLIN, BELFIELD,
DUBLIN 4, IRELAND
E-mail address: maris.ghergu@ucd.ie