

Exotic topological order in fractal spin liquids

Beni Yoshida

*Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA and
Institute for Quantum Information and Matter, California Institute of Technology, Pasadena, California 91125, USA*

(Dated: April 6, 2022)

We present a large class of three-dimensional spin models that possess topological order with stability against local perturbations, but are beyond description of topological quantum field theory. Conventional topological spin liquids, on a formal level, may be viewed as condensation of string-like extended objects with discrete gauge symmetries, being at fixed points with continuous scale symmetries. In contrast, ground states of fractal spin liquids are condensation of highly-fluctuating fractal objects with certain algebraic symmetries, corresponding to limit cycles under real-space renormalization group transformations which naturally arise from discrete scale symmetries of underlying fractal geometries. A theoretical framework for unified analysis on physical properties of fractal spin liquids is presented, which is applicable to all the previously known examples.

I. INTRODUCTION

In a quantum many-body system at zero temperature, topological order may arise when a gapped ground state possesses long-range entanglement that cannot be detected by any local measurement or local order parameter [1]. Ground state properties of topologically ordered systems are stable against any types of local perturbations, regardless of symmetries of perturbations, and depend only on global properties of geometric manifolds on which the whole systems are supported [2]. The discovery of topological order, such as fractional quantum hall systems [3, 4], came as a great surprise as it was beyond description of the Landau-Ginzburg theory which was once believed to be *the* ultimate theory of a quantum many-body system (see [5] for a review). It is now widely believed that the notion of topological order is essential in understanding the emergence of *quantum phases with no local order* in gapped quantum spin liquids, as seen in some resonating valence-bond states of frustrated anti-ferromagnets [6–13]. In addition to fundamental importance in condensed matter physics, the study of topologically ordered systems is of practical importance since they are physically natural platforms for realizations of fault-tolerant quantum information processing [1]. Searches for novel topological phases and their classification in a variety of theoretical and experimental systems have been an active area of research.

For topologically ordered spin liquids with discrete gauge symmetries, their low energy behavior is relatively well understood on a formal level as they can be effectively described by topological quantum field theory (TQFT) [2, 7], a field theory with an invariance under continuous deformations (diffeomorphism) [14, 15]. This is due to an observation that physical properties of topologically ordered systems do not depend on local structures of the systems, and topological properties of geometric manifolds play a crucial role in classifying topological phases. Indeed, a fairly complete class of two-dimensional TQFT-based spin systems with non-chiral topological order has been proposed by Levin and Wen where condensation of highly-fluctuating extended objects, called “string-nets”, are responsible for emergence of topological order [16]. A different, but closely related approach based on renormalization group (RG) ansatz of ground state wave-functions at fixed points has further verified the validity of TQFT as effective theory of topological phases in the presence of continuous scale invariance [17].

Yet, in some cases, quantum spin liquids may exhibit topological order that is beyond description of TQFT. For example, in three spatial dimensions, the Cubic code, recently proposed by Haah [18], possesses topological order with the stability against local perturbations, but seems to exhibit physical properties that are completely different from those of conventional topological spin liquids. For one thing, the number of degenerate ground states is exponential in the linear length of the lattice. Furthermore, unlike string-net condensates, the model is free from string-like extended objects, and the mobility of quasi-particle excitations is highly constrained via some algebraic rules. It has been argued that these exotic features are particularly useful for storing quantum bits (qubits) securely in thermal environment, but with some subtlety pointed out due to its thermal instability at non-zero temperature [19]. The discovery of the Cubic code and relevant models [20, 21] clearly indicates that classification of topological phases via TQFT is incomplete; TQFT is just a subset of some universal theory of topological order which is yet to be found. The necessary first step is to find a family of topological spin liquids that are beyond TQFT, and to develop a general theoretical framework to discuss their physical properties in a unified way.

The goal of this paper is to present a large class of exactly solvable topological spin liquids on a three-dimensional lattice which possess exotic topological order that cannot be described by TQFT. Instead of string-like (one-dimensional) or membrane-like (two-dimensional) objects with continuous geometries, ground states of the model are condensation of extended objects with non-integer dimensionality, namely *fractal objects*. In this paper, we discuss physical properties of such *quantum fractal liquids*, including; (a) the number of ground states and its dependence on geometric shapes of the manifolds, (b) the presence of topological order that is beyond TQFT, (c) propagations of quasi-particle excitations and their algebraic properties, and (d) topological entanglement entropies as an order parameter. Theoretical tools for analyses on quantum fractal liquids are also developed.

Emergence of fractal objects in correlated spin systems is not a completely new idea. In [22], Newman and Moore proposed a toy model of translation symmetric *classical* spin liquid on a two-dimensional square lattice. It has degenerate ground states whose spin configurations resemble the Sierpinski triangle with the famous “ $\log 3 / \log 2$ ” fractal dimension. Further studies revealed that a family of such fractal systems, referred to as *classical fractal liquids*, are particularly useful for storing a large amount of classical information reliably as they saturate a theoretical limit on classical information storage capacity of local Hamiltonians with mass gap [23]. Ground states of classical fractal liquids do not have continuous scale symmetries, but have discrete scale symmetries only, exhibiting *limit cycle* behaviors under real-space RG transformations as demonstrated in this paper. Physical properties of classical fractal liquids are intrinsically akin to those of a black hole since inner states are completely determined by degrees of freedom on a one-dimensional surface, and their coding properties obey an “area law” asymptotically [23]. Such exotic features of classical fractal liquids indicate a possibility of novel quantum phases that cannot be described by a field theory with continuous scale invariance. Construction of quantum fractal liquids can be viewed as natural generalization of classical fractal liquids to a quantum setting.

We first present a physical picture of quantum fractal liquids in section II by reviewing how condensation of extended objects emerge in topological spin liquids. In conventional topological spin liquids described by TQFT, extended objects with continuous geometries and integer dimensionality may emerge from underlying discrete gauge symmetries [7, 8]. In contrast, quantum fractal liquids are condensation of fractal objects with discrete geometries and non-integer dimensionality which emerge from certain algebraic symmetries. Geometric properties of extended objects can be characterized by topological classes of non-trivial symmetry operators where fractal symmetry operators are associated with quantum fractal liquids. In particular, geometric shapes of symmetry operators can be used as “order parameters” for classifying quantum phases where discontinuity in quantum phase transition is triggered by non-analytic changes of topologically distinct logical operators.

We then present a general framework to construct a family of classical fractal liquids with a variety of fractal geometries in section III. Quasi-particle excitations in classical fractal liquids may violate charge conservation since they do not possess gauge symmetries, but obey more general *algebraic symmetries* which are associated with polynomials that generate fractal geometries. Classical fractal liquids differ from conventional classical spin liquids as correlation functions have peculiar oscillatory behaviors associated with power-law decay with imaginary scaling dimensions. We demonstrate that ground states of classical fractal liquids correspond to limit cycles of real-space RG transformations. In section IV, we present a general framework to construct a family of three-dimensional quantum fractal liquids by weaving a pair of two-dimensional classical fractal liquids coherently such that their two-dimensional holographic images are two-dimensional classical fractal liquids. It is shown that quantum fractal liquids are topologically ordered with stability against local perturbations, but are beyond TQFT. We then discuss algebraic properties of quasi-particle excitations and derive a necessary and sufficient condition for quantum fractal liquids to be free from string-like logical operators. Several examples of quantum fractal liquids are also studied. In particular, the Cubic code is shown to be unitarily equivalent to a certain model of quantum fractal liquids, which we call a second-order model.

Finally, in section V, we compute topological entanglement entropies of quantum fractal liquids to establish connection between ground state and excitation properties. We demonstrate that topological entanglement entropies defined for tube-like regions can distinguish quantum fractal liquids from conventional topologically ordered systems. We give a physical interpretation based on propagations of quasi-particle excitations and their quantum dimension. Section VI is devoted to conclusions and open questions. Technical tools to analyze a family of translation symmetric stabilizer Hamiltonians are presented in appendix A which may be of independent interest for quantum information scientists.

Some comments on the paper follow. We adopt the stability against local perturbations as the definition of topological order. By TQFT, we mean an axiomatic formulation by Atiyah which admits only a finite number of degenerate ground states [15]. For a review of TQFT, see [24]. By topological spin liquids, we mean gapped spin

systems without local symmetries, i.e. topologically ordered spin systems. A connection between topological order and discrete gauge theory was pointed out by Read and Sachdev [7] and Wen [8]. Discussion on symmetry-protected topological order is beyond the scope of this paper. Discussion on gapless quantum spin liquids is beyond the scope of this paper. Our construction of quantum fractal liquids is theoretically motivated, and its relevance to experimental realization may not be immediately clear. Some technical tools are borrowed from a recent work by Haah [25] which aims to classify translation symmetric stabilizer Hamiltonians via commutative algebra. See [20] for a few other examples of string-free models proposed by Kim. See [21] for a model with pairs of fractal and string-like logical operators proposed by Castelnovo and Chamon. Limit cycles in RG transformations are not new phenomena as observed in the Efimov effect in non-relativistic conformal field theory with imaginary scaling dimensions [26–28]. As for relativistic field theory, it remains open for $D > 2$ [29, 30]. See [31] and subsequent papers by others for a recent proposal and debate. Discussion on coding properties is beyond the scope of this paper, and is briefly presented in appendix A.

II. TOPOLOGICAL SPIN LIQUID

A. Topological spin liquid and string-nets

In this section, we review how condensation of extended objects emerge in topological phases and present a physical picture of quantum fractal liquids, *condensation of fractal objects*. We begin with the simplest string-net model, known as $\mathbb{Z}_2$ spin liquid (or the Toric code) [1, 32] (Fig 1(a)). Consider a square lattice where qubits live on edges of the lattice with periodic boundary conditions. The Hamiltonian is

$$H = - \sum_s A_s - \sum_p B_p, \quad A_s = \prod_{r \in s} X_r, \quad B_p = \prod_{r \in p} Z_r$$

where s represents a star and p represents a plaquette. Pauli X and Z operators act on each qubit as follows: $Z|0\rangle = |0\rangle$, $Z|1\rangle = -|1\rangle$, $X|0\rangle = |1\rangle$ and $X|1\rangle = |0\rangle$. The model is exactly solvable as interaction terms A_s and B_p commute with each other, and ground states satisfy

$$A_s|\psi\rangle = |\psi\rangle, \quad B_p|\psi\rangle = |\psi\rangle, \quad \forall s, p.$$

A ground state can be viewed as condensation of string-like extended objects. Consider a trivial product state $|0\rangle^{\otimes N}$ over the entire lattice (N is the total number of qubits) and observe that $B_p|0\rangle^{\otimes N} = |0\rangle^{\otimes N}$. The following is a ground state:

$$|\psi_{loop}\rangle = \prod_s (1 + A_s) |0\rangle^{\otimes N} \quad (1)$$

since $A_s(1 + A_s) = 1 + A_s$. The normalization factor is omitted. It is a superposition of terms $A_{s_1} A_{s_2} A_{s_3} \cdots |0\rangle^{\otimes N}$. Since A_s is a product of Pauli- X operators, it flips the signs of qubits: $|0\rangle \rightarrow |1\rangle$. Then a term $A_s |0\rangle^{\otimes N}$ can be viewed as a state with one small loop on a “dual lattice”, and a term $A_{s_1} A_{s_2} |0\rangle^{\otimes N}$ with neighboring stars s_1 and s_2 is a state with a larger loop (Fig. 1(b)). In general, $A_{s_1} A_{s_2} A_{s_3} \cdots |0\rangle^{\otimes N}$ is a state with loops of various sizes and shapes. A ground state is a superposition of all the loop states (Fig. 1(c)):

$$|\psi_{loop}\rangle = \sum_{\forall \gamma} |\gamma\rangle \quad (2)$$

where γ represents an arbitrary loop configuration. Therefore a ground state is condensation of *fluctuating string-like objects* with $\mathbb{Z}_2$ gauge symmetry. Note the model has four-fold degeneracy on a torus.

One can construct a general quantum many-body system with several types of strings constrained by discrete gauge symmetry. Levin and Wen derived the most general form of wave-functions that are represented as condensation of string-like extended objects on a two-dimensional lattice by further assuming that wave-functions possess *scale invariance* and correspond to fixed-points of RG transformations [16]. Indeed, a ground state of $\mathbb{Z}_2$ spin liquid has scale invariance as it is a superposition of loops of all the different sizes and shapes. Note that scale invariance is




FIG. 1: $\mathbb{Z}_2$ spin liquid (the Toric code). (a) The Hamiltonian. (b) Loop states on a dual lattice. (c) Condensation of loops. (d) Logical operators.

required for systems described by TQFT since they must be invariant under continuous deformations. Yet, scale invariance is not a necessary condition for the presence of topological order. As we will see, quantum fractal liquids do not have full continuous scale symmetries. Instead, they have *discrete scale symmetries* where systems are invariant only under a limited set of scale transformations and ground states correspond to limit cycles of RG transformations.

Geometric properties of extended-objects can be characterized by topological properties of global symmetry operators. Formally, symmetries of the Hamiltonian can be captured by unitary transformations that leave the Hamiltonian invariant:

$$U^\dagger H U = H \quad \Rightarrow \quad [H, U] = 0. \quad (3)$$

Interaction terms A_s and B_p are gauge symmetry operators for $\mathbb{Z}_2$ spin liquid since $[A_s, H] = [B_p, H] = 0$ where the ground state space is an invariant subspace under actions of interaction terms. There also exist *non-trivial symmetry operators* which act non-trivially inside the ground state space (see Fig 1(d)):

$$[H, \ell_0^{(Z)}] = [H, \ell_1^{(Z)}] = [H, \ell_0^{(X)}] = [H, \ell_1^{(X)}] = 0.$$

which have non-trivial winding on a torus.

Since non-trivial symmetry operators commute with the Hamiltonian, they do not change the energy of the system. Yet, they cannot be written as products of A_s or B_p and transform degenerate ground states into each other. Recall that $|\psi_{loop}\rangle$ is a condensation of loops that can be shrunk into a vacuum under $\mathbb{Z}_2$ gauge symmetry. An application of $\ell_0^{(X)}$ to $|\psi_{loop}\rangle$ creates a non-trivial loop winding around the lattice, and $\ell_0^{(X)}|\psi_{loop}\rangle$ is condensation of loops with non-trivial winding in the $\hat{x}$ direction. Similarly, $\ell_1^{(X)}|\psi_{loop}\rangle$ is condensation of loops with non-trivial winding in the $\hat{y}$ direction. Four degenerate ground states may be indexed by winding numbers as $|\tilde{\gamma}_x\rangle \otimes |\tilde{\gamma}_y\rangle$ with $\gamma_x, \gamma_y = 0, 1$ where γ_x and γ_y represent the presence or absence of windings in the $\hat{x}$ and $\hat{y}$ directions respectively. Then non-trivial symmetry operators $\ell_0^{(X)}$ and $\ell_1^{(X)}$ act like Pauli-X operators on a pair of *logical qubits* $|\tilde{\gamma}_x\rangle \otimes |\tilde{\gamma}_y\rangle$. It is convenient to represent their commutation relation as follows

$$\left\{ \begin{array}{cc} \ell_0^{(Z)}, & \ell_1^{(Z)} \\ \ell_0^{(X)}, & \ell_1^{(X)} \end{array} \right\}$$

where operators in the same column anti-commute with each other while logical operators in different columns commute with each other. Anti-commuting pairs of non-trivial symmetry operators can be viewed as logical Pauli-Z and Pauli-X acting on logical qubits $|\tilde{\gamma}_x\rangle \otimes |\tilde{\gamma}_y\rangle$. These non-trivial symmetry operators are often called *logical operators* in quantum information community as they can rewrite encoded logical qubits in the ground state space.

Note $\mathbb{Z}_2$ spin liquid is a good quantum error-correcting code as only global operators with non-trivial winding can change encoded logical qubits, and no local errors can destroy stored quantum information. This is the origin of the stability against local perturbations in topological phases from quantum information perspective [1, 33].

Another useful way of revealing topological properties of $\mathbb{Z}_2$ spin liquid is to consider its *one-dimensional limit*. Let us shrink the $\hat{y}$ axis of the lattice and consider a one-dimensional limit. Then a system consists of a pair of non-interacting ferromagnets in the $\hat{x}$ direction. Similarly, one obtains another pair of non-interacting ferromagnets by shrinking the $\hat{x}$ axis. Therefore, two-dimensional $\mathbb{Z}_2$ spin liquid can be constructed by weaving one-dimensional ferromagnets coherently such that its one-dimensional holographic images are ferromagnets.

One can generalize $\mathbb{Z}_2$ spin liquid to higher-dimensional systems. For instance, the D -dimensional Toric code has anti-commuting pairs of m -dimensional and $(D - m)$ -dimensional logical operators:

$$m\text{-dim} \leftrightarrow (D - m)\text{-dim} \quad m : \text{integer}$$

Its ground states can be viewed as condensation of m -dimensional extended-objects, or $(D - m)$ -dimensional extended objects in a dual description. In general, for a quantum many-body system described by TQFT, extended objects (Wilson loops or surfaces) have continuous geometries with integer dimensionality. In fact, the dimensional duality of non-trivial symmetry operators is a consequence of the Poincaré duality for systems described by TQFT and can be derived analytically from continuous deformability of logical operators as shown in [34].

B. Emergence of fractal geometry

In conventional topologically ordered systems, extended objects have continuous geometries with integer dimensionality. Yet, one may construct a quantum many-body system with fractal objects which have discrete geometries with non-integer dimensionality. In this subsection, we give a physical picture of quantum fractal liquids by reviewing how fractal geometries arise in classical spin systems.

Consider a square lattice where $L \times L$ spins live on vertices of the lattice. The Hamiltonian is

$$H = - \sum_{i=0}^{L-2} \sum_{j=0}^{L-1} \Pi_{ij}, \quad \Pi_{ij} = Z_{i,j} Z_{i+1,j} Z_{i+1,j+1} = \begin{pmatrix} Z_{i,j}, & Z_{i+1,j} \\ I, & Z_{i+1,j+1} \end{pmatrix}$$

where we represented interaction terms *graphically* as a matrix. We denote spin values at (i, j) for $i, j = 0, \dots, L - 1$ as $s_{i,j} = 0, 1$. Ground states must satisfy $\Pi_{ij}\psi = \psi$, and

$$s_{i,j} + s_{i+1,j} = s_{i+1,j+1} \pmod{2} \quad (4)$$

for all i and for $j = 0, \dots, L - 2$. The following is a ground state:

$$\begin{bmatrix} 1, & 0, & 0, & 0, & 0, & 0, & 0, & 0, & \dots \\ 1, & 1, & 0, & 0, & 0, & 0, & 0, & 0, & \dots \\ 1, & 0, & 1, & 0, & 0, & 0, & 0, & 0, & \dots \\ 1, & 1, & 1, & 1, & 0, & 0, & 0, & 0, & \dots \\ 1, & 0, & 0, & 0, & 1, & 0, & 0, & 0, & \dots \\ 1, & 1, & 0, & 0, & 1, & 1, & 0, & 0, & \dots \\ 1, & 0, & 1, & 0, & 1, & 0, & 1, & 0, & \dots \\ 1, & 1, & 1, & 1, & 1, & 1, & 1, & 1, & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

where the upper-left corner corresponds to $s_{0,0}$. Configuration of sites with spin value 1 forms the *Sierpinski triangle* whose fractal dimension is $\log 3 / \log 2$ (see Fig. 4(a)). It is interesting to observe that a ground state does not have translation symmetries while the parent Hamiltonian is translation symmetric. We call this spontaneous breaking of translation symmetries *strong-breaking of translation symmetries* in contrast to weak-breaking of translation symmetries introduced in [13].

The model has a large number of degenerate ground states. Let us pick up arbitrary spin values on the first row of the lattice $\vec{s} = (s_{0,0}, s_{1,0}, s_{2,0}, \dots, s_{L-1,0})$. Then, according to Eq. (4), spin values on other rows of the lattice are determined. Since there are 2^L possible choices for $\vec{s}$, there are 2^L degenerate ground states. It is convenient to view the model as a time-evolution of one-dimensional cellular automaton where spin values on lower rows are computed via an update rule in Eq. (4). It is well known that one-dimensional cellular automata with linear update rules generate a variety of fractal geometries (see [35] for a review). One may consider a general class of classical spin models with fractal ground states by designing interaction terms which imitate update rules of one-dimensional cellular automata. Such *classical fractal liquid* is particularly useful for storing classical bits of information securely as it asymptotically saturate a theoretical limit of classical information storage capacity of gapped spin systems [23].

The Sierpinski triangle model has liquid-like order, but is different from conventional classical spin liquids such as anti-ferromagnetic Ising models on geometrically frustrated lattices [36–40]. Due to unconventional three-body interactions, the model does not have magnetic order at any temperature including $T = 0$. A zero temperature thermodynamic entropy is large, but not extensive: $S = \sqrt{N}$. In the Sierpinski triangle model, whether two spins have ferromagnetic or anti-ferromagnetic interactions depends on a spin value of the third spin, which leads to emergence of peculiarly regulated fractal geometries. Whether the model may select an ordered ground state via order by disorder mechanism is not known. In this paper, we refer to a family of the Sierpinski triangle model as *classical fractal liquids* despite technical subtleties mentioned above.

The model does not have power law decay of two-point correlation functions as observed in conventional classical spin liquids. Instead, we point out that it has an oscillatory power-law behavior with imaginary scaling dimensions, exhibiting *discrete scale symmetries*. Consider the following three-point correlation function:

$$C(r) = \langle Z_{i,j} Z_{i+r,j} Z_{i+r,j+r} \rangle. \quad (5)$$

In the ground space manifold, $C(r)$ is

$$\begin{aligned} C(r) &= 1, & r &= 2^m \\ &= 0, & r &\neq 2^m \end{aligned}$$

with oscillatory behaviors in $\log r$, instead of r . At finite temperature, the three-point correlation function reads

$$C(r) = (1 - 2p)^{r^{\log 3 / \log 2}}, \quad r = 2^m$$

with $p = e^{-\beta} / (e^{-\beta} + e^{\beta})$ where the exponent has a peculiar dependence on a fractal dimension. The correlation function for all r is

$$C(r) \propto \exp(-A \cdot r^{\log 3 / \log 2}) \cdot \sum_{k=-\infty}^{\infty} r^{i \frac{2\pi k}{\log 2}}$$

where oscillatory behaviors in $\log r$ are represented by power law with imaginary scaling dimensions, which is characteristic of systems with discrete scale symmetries [27].




FIG. 2: Discrete scale symmetries and imaginary scaling dimensions in three-point correlation function.

Geometric properties of degenerate ground states in the Sierpinski triangle model can be captured by geometric

shapes of logical operators:

$$\ell = \begin{bmatrix} Z, & I, & I, & I, & I, & I, & I, & I, & \dots \\ I, & I, & I, & I, & I, & I, & I, & I, & \dots \\ I, & I, & I, & I, & I, & I, & I, & I, & \dots \\ I, & I, & I, & I, & I, & I, & I, & I, & \dots \\ I, & I, & I, & I, & I, & I, & I, & I, & \dots \\ I, & I, & I, & I, & I, & I, & I, & I, & \dots \\ I, & I, & I, & I, & I, & I, & I, & I, & \dots \\ I, & I, & I, & I, & I, & I, & I, & I, & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}, \quad r = \begin{bmatrix} X, & I, & I, & I, & I, & I, & I, & I, & \dots \\ X, & X, & I, & I, & I, & I, & I, & I, & \dots \\ X, & I, & X, & I, & I, & I, & I, & I, & \dots \\ X, & X, & X, & X, & I, & I, & I, & I, & \dots \\ X, & I, & I, & I, & X, & I, & I, & I, & \dots \\ X, & X, & I, & I, & X, & X, & I, & I, & \dots \\ X, & I, & X, & I, & X, & I, & X, & I, & \dots \\ X, & X, & X, & X, & X, & X, & X, & X, & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

where they commute with all the interaction terms, but cannot be represented as products of interaction terms. So, it has the following pairs of logical operators:

$$0\text{-dim} \quad \leftrightarrow \quad \frac{\log 3}{\log 2}\text{-dim}.$$

While the model has a fractal logical operator, a ground state does not have any quantum fluctuation, and is not topologically ordered since its partner is a trivial logical operator with zero-dimensional geometry. To have topological order, both logical operators must have topologically non-trivial geometries (i.e. they must be finite-dimensional).

Given the large scope of theory of fractal geometries, it is natural to wonder if fluctuations of fractal objects can describe general topologically ordered quantum systems that are beyond descriptions of TQFT. In the reminder of the paper, we present a large class of quantum spin systems with fractal liquid order and analyze its physical properties. In particular, we construct a model of quantum spin systems which has pairs of anti-commuting fractal logical operators:

$$\text{fractal-dim} \quad \leftrightarrow \quad \text{fractal-dim}$$

by combining features of both $\mathbb{Z}_2$ spin liquid and classical fractal liquids.

C. Topological phase transition

A quantum many-body system with topological order can be viewed as condensation of extended objects with a variety of geometric shapes. It is natural to expect that two ground states with different types of extended objects exhibit totally different physical properties, and thus belong to distinct topological phases. In this subsection, we make this intuition more precise by arguing that two spin systems with topologically different classes of logical operators are always separated by quantum phase transitions.

We begin by reviewing a basic idea of classification of quantum phases by following [17, 41]. The notion of quantum phases characterizes long-range ground state properties of a many-body quantum system with mass gap at zero temperature. Ground states in different quantum phases cannot be connected smoothly at the thermodynamic limit. Let us consider two ground states $|\psi_A\rangle$ and $|\psi_B\rangle$ of two different gapped Hamiltonian H_A and H_B and ask if they are separated by quantum phase transitions (non-analytic changes of ground state properties) or not. Two ground states $|\psi_A\rangle$ and $|\psi_B\rangle$ are said to be in different quantum phases when there always exist some quantum phase transitions between H_A and H_B regardless of choices of paths connecting H_A and H_B . Conversely, if there exists a continuous change from H_A to H_B without crossing quantum phase transitions, two ground states $|\psi_A\rangle$ and $|\psi_B\rangle$ are in the same quantum phase. In summary, one has the following classification principle:

$$\text{Being always separated by QPTs} \quad \Leftrightarrow \quad \text{Different quantum phases}$$

An equivalent, but more convenient way of classifying quantum phases uses local unitary transformations. Two ground states $|\psi_A\rangle$ and $|\psi_B\rangle$ are considered to be in the same quantum phase when there exists some local unitary transformation connecting $|\psi_A\rangle$ and $|\psi_B\rangle$. By local unitary transformations, we mean transformations generated by a set of geometrically local quantum operations, applied for a finite duration. On the other hand, when there is no




FIG. 3: Classification of quantum phases. (a) Continuous deformability of ground states. (b) Continuous deformability of logical operators.

local unitary transformation connecting $|\psi_A\rangle$ and $|\psi_B\rangle$, they are in different quantum phases:

$$\text{Being connected only by global unitary} \Leftrightarrow \text{Different quantum phases}$$

Note that only global unitary transformations can change long-range properties of ground states.

These two classification principles of quantum phases, namely quantum phase transitions and local unitary transformations, are equivalent under appropriate assumptions. If two gapped Hamiltonians H_A and H_B can be transformed into each other continuously without closing the energy gap, correlation lengths of ground states remain finite, and ground states at each stage of transformation can be approximated by applying some quasi-local unitary transformations to original ground states [33]. Conversely, if $|\psi_A\rangle$ and $|\psi_B\rangle$ are connected by some local unitary transformations, one can always continuously transform H_A into H_B . Discussion so far is summarized in Fig. 3(a).

The classification of quantum phases, based on continuous deformability of ground state wave-functions, reminds us of the study of *topology* in mathematics, which aims to classify geometric shapes of object based on continuous deformability. Roughly speaking, two objects are considered to be equivalent when they can be transformed into each other via continuous deformations (diffeomorphism). Yet, if one cannot continuously deform an object to the other, they are considered to be topologically different. The similarity between classifications of quantum phases, based on continuous deformability of wave functions, and classifications of geometric shapes, based on continuous deformability of geometric objects, allows us to use the notion of topology in classifying quantum phases. Indeed, the following relation holds [42, 43]:

$$\text{Logical operators are topologically different} \Rightarrow \text{Two systems belong to different quantum phases.}$$

The argument roughly goes as follows. Consider two systems with topologically distinct logical operators ℓ and ℓ' . Let us suppose that they belong to the same quantum phase. Then, there must be some local unitary transformation U such that $U\ell U^\dagger = \ell'$. Yet, this is not possible since local unitary transformation can change geometric shapes of logical operators only continuously at the thermodynamic limit. Therefore, models with topologically different types of logical operators belong to different quantum phases and are always separated by quantum phase transitions. Note that there is no local unitary that transforms a string-like logical operator to a fractal logical operator, and thus, fractal models are different from conventional topologically ordered systems. “ $\Leftarrow$ ” of the above relation is proven only for stabilizer Hamiltonians with continuous scale symmetries (i.e. within TQFT) [43].

In summary, we expect that there will be four classes of quantum phases arising in stabilizer Hamiltonians.

- **Ferromagnetic phase** has zero-dimensional logical operators and D -dimensional logical operators as in fer-

romagnet. It possesses continuous scale symmetries with finite ground state degeneracy, and its ground states correspond to fixed points of RG transformations. Translation symmetries are not strongly broken.

- **Classical fractal phase** has zero-dimensional logical operators and fractal logical operators as in classical fractal liquids. It possesses discrete scale symmetries with increasing ground state degeneracy, and its ground states correspond to limit cycles of RG transformations. Translation symmetries are strongly broken.
- **Topological phase** has m -dimensional logical operators and $(D - m)$ -dimensional logical operators where $m > 0$ is an integer as in $\mathbb{Z}_2$ spin liquid. It is topologically ordered with stability against local perturbations. It possesses continuous scale symmetries with finite ground state degeneracy, and its ground states correspond to fixed-points of RG transformations. Translation symmetries are not strongly broken in the ground state space.
- **Quantum fractal phase** has pairs of fractal logical operators as in quantum fractal liquids. It is topologically ordered with stability against local perturbations. It possesses discrete scale symmetries with increasing ground state degeneracy, and its ground states correspond to limit cycles of RG transformations. Translation symmetries are strongly broken.

III. CLASSICAL FRACTAL LIQUID

A. Fractal and algebraic symmetry




FIG. 4: Examples of fractal geometries generated by polynomials. Unfilled elements represent zero, gray elements represent one, and red elements represent two. (a) The Sierpinski triangle from $f = 1 + x$ over $\mathbb{F}_2$ with the fractal dimension $\mathcal{D} = \log 3 / \log 2$. (b) The Fibonacci model from $f = 1 + x + x^2$ over $\mathbb{F}_2$ with the fractal dimension $\mathcal{D} = \log(1 + \sqrt{5}) / \log 2$. (c) The generalized Sierpinski triangle from $f = 1 + x$ over $\mathbb{F}_3$ with the fractal dimension $\mathcal{D} = \log 6 / \log 3$.

In this section, we present a theoretical framework to construct a family of classical fractal liquids. It is well known that a variety of fractal geometries can be generated by polynomials over finite fields. We begin with polynomial representation of the Sierpinski triangle (Fig. 4(a)). Consider a polynomial $f = 1 + x$ over $\mathbb{F}_2$ and its j -th powers:

$$\begin{aligned}
 f^0 &= 1 \\
 f^1 &= 1 + x \\
 f^2 &= 1 + x^2 \\
 f^3 &= 1 + x + x^2 + x^3 \\
 f^4 &= 1 + x^4 \\
 f^5 &= 1 + x + x^4 + x^5
 \end{aligned}$$

where coefficients are computed modulo 2. More graphically, one has

$$\begin{bmatrix} f^0 \\ f^1 \\ f^2 \\ f^3 \\ f^4 \\ f^5 \end{bmatrix} = \begin{bmatrix} 1 & & & & & \\ 1 & x & & & & \\ 1 & & x^2 & & & \\ 1 & x & x^2 & x^3 & & \\ 1 & & & & x^4 & \\ 1 & x & & & x^4 & x^5 \end{bmatrix}$$

where “+” signs are omitted. One will notice that the Sierpinski triangle emerges in a geometric pattern of non-zero coefficients in f^j ($j = 0, 1, \dots$).

The entire Sierpinski triangle can be represented as a single polynomial with x and y :

$$\mathbf{f}(x, y) = 1 + fy + f^2y^2 + f^3y^3 + \dots$$

where j -th row is indexed by y^j . More graphically, one has

$$\mathbf{f}(x, y) = \begin{bmatrix} 1 & & & & & \\ y & xy & & & & \\ y^2 & & x^2y^2 & & & \\ y^3 & xy^3 & x^2y^3 & x^3y^3 & & \\ y^4 & & & & x^4y^4 & \\ y^5 & xy^5 & & & x^4y^5 & x^5y^5 \end{bmatrix}$$

where non-zero coefficients of $x^i y^j$ correspond to filled elements of the Sierpinski triangle at (i, j) . Note that row vectors of the Sierpinski triangle evolves in the $\hat{y}$ direction via applications of generating polynomial f .

Another interesting example of fractal geometries is generated by $f = 1 + x + x^2$ over $\mathbb{F}_p$:

$$\begin{bmatrix} f^0 \\ f^1 \\ f^2 \\ f^3 \\ f^4 \end{bmatrix} = \begin{bmatrix} 1 & & & & & \\ 1 & x & x^2 & & & \\ 1 & & x^2 & x^4 & & \\ 1 & x & & x^3 & x^5 & x^6 \\ 1 & & & x^4 & & x^8 \end{bmatrix}.$$

Again, the entire fractal geometry can be represented as $\mathbf{f}(x, y) = 1 + fy + f^2y^2 + f^3y^3 + \dots$. The model is often called the Fibonacci model since its fractal dimension is given by $\frac{\log 1 + \sqrt{5}}{\log 2}$ (Fig. 4(b)). The above construction of fractal geometries can be generalized to polynomials over $\mathbb{F}_p$ ($p > 2$) with an arbitrary prime p . For instance, consider $f = 1 + x$ over $\mathbb{F}_3$ ($p = 3$):

$$\begin{bmatrix} f^0 \\ f^1 \\ f^2 \\ f^3 \\ f^4 \\ f^5 \end{bmatrix} = \begin{bmatrix} 1 & & & & & \\ 1 & x & & & & \\ 1 & 2x & x^2 & & & \\ 1 & & & x^3 & & \\ 1 & x & & x^3 & x^4 & \\ 1 & 2x & x^2 & x^3 & 2x^4 & x^5 \end{bmatrix}$$

which is a generalization of the Sierpinski triangle for three-dimensional spins (Fig. 4(c)).

The self-similarity in fractal geometries arises from *discrete scale symmetries* of generating polynomials. Consider an arbitrary polynomial f over $\mathbb{F}_p$ with prime p :

$$f = c_0 + c_1x + c_2x^2 + c_3x^3 + \dots \quad (6)$$

where $c_j = 0, \dots, p-1$. Then its p -th power is

$$f^p = c_0 + c_1 x^p + c_2 x^{2p} + c_3 x^{3p} + \dots \quad (7)$$

For instance, with $f = 1 + 2x + x^2$ over $\mathbb{F}_3$, one finds

$$f = 1 + 2x + x^2 \quad f^3 = 1 + 2x^3 + x^6 \quad f^9 = 1 + 2x^9 + x^{18} \quad f^{27} = 1 + 2x^{27} + x^{54}.$$

Due to discrete scale symmetries, the generated fractal geometry $\mathbf{f}(x, y) = 1 + fy + f^2 y^2 + f^3 y^3 + \dots$ has a self-similarity where the same pattern appear repeatedly at different length scales.

Fractal geometries do not possess gauge symmetries since growth of filled elements violates *charge conservation* where a single element may evolve into multiple elements of the same type in the $\hat{y}$ direction. This is in strong contrast with the fact that continuous geometries often have physical interpretations based on conservation laws associated with underlying gauge symmetries as in the case of TQFT [16]. Charge conservation in scale invariant spin models originate from group theoretical constraints imposed on the parent Hamiltonian. Fractal geometries obey a more general form of symmetries, which will be referred to as *algebraic symmetries*, for a possible relation to theory of algebraic geometry which concerns geometric structures of solutions of polynomial equations.

B. Polynomial representation of Pauli operators

To construct parent Hamiltonians of classical fractal liquids, it is convenient to represent interaction terms by polynomials too. We begin with construction of Pauli operators for a system of qubits on a one-dimensional infinite chain, following [25]. Consider a polynomial f over $\mathbb{F}_2$:

$$f = \sum_{j=-\infty}^{\infty} c_j x^j, \quad c_j = 0, 1. \quad (8)$$

We define the corresponding Pauli operators as follows

$$Z(f) = \prod_{j=-\infty}^{\infty} Z_j^{c_j}, \quad X(f) = \prod_{j=-\infty}^{\infty} X_j^{c_j} \quad (9)$$

where Z_j and X_j are Pauli operators acting on j -th qubit. So, a polynomial f encodes positions of qubits where Pauli operators Z_j or X_j may act. For instance, $f = 1 + x + x^2$ and $Z(f) = Z_0 Z_1 Z_2$.

The polynomial representation of Pauli operators is particularly useful for studying spin systems with translation symmetries since translations can be concisely described in terms of polynomials. For instance, consider a Pauli operator $Z(f) = Z_0 Z_1 Z_2$ for $f = 1 + x + x^2$. Then, its translation in the $\hat{x}_+$ direction is given by $Z_1 Z_2 Z_3$, whose polynomial representation is $Z(xf)$:

$$\begin{aligned} f = 1 + x + x^2 &\rightarrow xf = x + x^2 + x^3 \\ Z(f) = Z_0 Z_1 Z_2 &\rightarrow Z(xf) = Z_1 Z_2 Z_3. \end{aligned}$$

In general, $Z(xf)$ is a translation of $Z(f)$ in the $\hat{x}_+$ direction. Similarly, a translation in the $\hat{x}_-$ direction is given by $Z(x^{-1}f)$. One may generalize this formalism to higher-dimensional systems by adding extra variables $y, z, \dots$.

To gain more insights, let us represent a one-dimensional ferromagnet by polynomials over $\mathbb{F}_2$:

$$H = - \sum_j Z(x^j(1+x))$$

where $Z(x^j(1+x)) = Z_j Z_{j+1}$. The Sierpinski triangle model is

$$H = - \sum_{ij} Z(x^i y^j (1+x+xy))$$

where interaction terms are translations of $Z(1+x+xy)$. In general, one may consider a classical translation symmetric Hamiltonian

$$H = - \sum_{i,j,\dots} Z(x^i y^j \dots \alpha) \quad (10)$$

with an arbitrary polynomial $\alpha(x, y, \dots)$ where interaction terms are translations of $Z(\alpha)$. Ground states of a Hamiltonian obey

$$Z(x^i y^j \dots \alpha) |\psi\rangle = |\psi\rangle \quad \forall i, j, \dots \quad (11)$$

The polynomial representation of Pauli operators becomes particularly powerful in analyzing commutation relations between Pauli operators $Z(f)$ and $X(g)$. Since we are interested in translation symmetric systems, we would like to obtain commutation relations between $Z(f)$ and translations of $X(g)$. Let us imagine that we check commutation relations between $Z(f)$ and $X(x^j g)$ for all j and assign integers $d_j = 0, 1$ as follows

$$\begin{aligned} d_j = 0 & \quad \text{for} \quad [Z(f), X(x^j g)] = 0 \\ d_j = 1 & \quad \text{for} \quad \{Z(f), X(x^j g)\} = 0. \end{aligned}$$

Based on d_j , we define the *commutation polynomial* $P(f, g)$ as follows:

$$P(f, g) = \sum_j d_j x^j \quad \text{s.t} \quad Z(f) X(x^j g) = (-1)^{d_j} X(x^j g) Z(f). \quad (12)$$

Thus, the commutation polynomial $P(f, g)$ is a collection of commutation relations between $Z(f)$ and $X(x^j g)$. For instance, with $f = 1 + x + x^2$ and $g = 1 + x$, $Z(f)$ anti-commutes only with $X(x^{-1}g)$ and $X(x^2g)$. So, the commutation polynomial is $P(f, g) = x^{-1} + x^2$.

The commutation polynomial $P(f, g)$ can be concisely written by introducing the notion of *dual*:

$$f = \sum_{j=-\infty}^{\infty} c_j x^j \quad \rightarrow \quad \bar{f} = \sum_{j=-\infty}^{\infty} c_j x^{-j} \quad (13)$$

where the dual $\bar{f}$ is obtained by taking $x \rightarrow x^{-1}$. Then, the commutation polynomial is given by the *convolution*

$$P(f, g) = f \bar{g} \quad (14)$$

For instance, one has $f \bar{g} = (1 + x + x^2)(1 + x^{-1}) = x^{-1} + 2 + 2x + x^2 = x^{-1} + x^2$ for the above example. The proof of Eq. (14) is straightforward. Generalization to polynomials over $\mathbb{F}_p$ is straightforward.

C. Classical fractal liquid

We present general construction of classical fractal liquids. Consider a two-dimensional square lattice with $L \times L$ spins ($L = 2^m$). The Hamiltonian is

$$H = - \sum_{ij} Z(x^i y^j \bar{\alpha}), \quad \alpha = 1 + f(x)y \quad (15)$$

where $f(x)$ is an arbitrary polynomial over $\mathbb{F}_2$ with x only. We put periodic boundary conditions both in the $\hat{x}$ and $\hat{y}$ directions, and assume that $f(x)$ is reversible. (See appendix A for discussion on reversibility).

In finding ground states of the Hamiltonian, it is convenient to find its logical operators. Z -type logical operators are trivial single Pauli operators $\ell_j^{(Z)} = Z_{j0} = Z(x^j)$ for $j = 0, \dots, L-1$ while X -type logical operators have fractal

geometries:

$$\ell_j^{(X)} = X(x^j \mathbf{f}(x, y)), \quad \mathbf{f}(x, y) = 1 + fy + \cdots + (fy)^{L-1}$$

where $\mathbf{f}(x, y)$ is a fractal geometry generated by $f(x)$. One can see that $\ell_j^{(X)}$ commute with all the stabilizer generators since the commutation polynomial between $Z(1 + \bar{f}\bar{y})$ and $X(\mathbf{f}(x, y))$ is

$$P(1 + \bar{f}\bar{y}, 1 + fy + \cdots + (fy)^{L-1}) = (1 + \bar{f}\bar{y})(1 + \bar{f}\bar{y} + \cdots + (\bar{f}\bar{y})^{L-1}) = 0$$

when $f(x)$ is reversible. We list all the logical operators as follows:

$$\left\{ \begin{array}{l} \ell_0^{(X)}, \dots, \ell_{L-1}^{(X)} \\ \ell_0^{(Z)}, \dots, \ell_{L-1}^{(Z)} \end{array} \right\}$$

where pairs of logical operators in the same column anti-commute with each other while pairs of logical operators in different columns commute with each other. So, there are $k = L$ logical bits in total.

By using X-type logical operators $\ell_j^{(X)}$, one can find all the ground states of a classical fractal liquid. Let us denote spin values at (i, j) as $s_{ij} = 0, 1$ for $i, j = 0, \dots, L-1$, and represent a ground state ψ as

$$\psi = \sum_{ij} s_{ij} x^i y^j. \quad (16)$$

Since the Hamiltonian consists only of Z-type Pauli operators, $\psi = 0$ with $s_{ij} = 0$ is a ground state of the Hamiltonian. (Recall $Z|0\rangle = |0\rangle$ and $Z|1\rangle = -|1\rangle$ in our notation). To find another ground state, one applies $\ell_0^{(X)}$ to $\psi = 0$ and obtains a fractal ground state:

$$\psi(1) = 1 + fy + \cdots + (fy)^{L-1} = \mathbf{f}(x, y) \quad (17)$$

One can find all the other ground states by applying fractal logical operators $\ell_j^{(X)}$. There are 2^L degenerate ground states, represented by

$$\psi(\gamma) = \gamma(x)(1 + fy + \cdots + (fy)^{L-1}) = \gamma(x)\mathbf{f}(x, y) \quad (18)$$

where $\gamma(x)$ is an arbitrary polynomial with x only. Noting $\dim \gamma = L$, one finds $k = L$.

Classical fractal liquids discussed so far are based on *first-order* cellular automata whose present states at $t = \tau$ depend on states at $t = \tau - 1$. In higher-order cellular automata, the present states at $t = \tau$ may depend on states at $t = \tau - q, \dots, \tau - 1$ for $q > 1$. One can construct classical fractal liquids based on higher-order cellular automata by taking

$$\alpha = 1 + f_1(x)y + f_2(x)y^2 + \cdots + f_q(x)y^q.$$

However, it is generally difficult to write down spin configurations of higher-order classical fractal liquids explicitly.

Since the model does not have gauge symmetries, its quasi-particle excitations violate charge conservation and propagate according to algebraic symmetries imposed by generating polynomial $f(x)$. Recall that ground states ψ satisfy $Z(x^i y^j \bar{\alpha})\psi = \psi$ for all i, j , and quasi-particle excitations may be viewed as violations of these algebraic constraints. It is convenient to represent positions of excitations by an *excitation polynomial*:

$$E(x, y) = \sum_{i,j,\ell} c_{ij} x^i y^j$$

where an excited state ψ' has

$$\begin{array}{ll} c_{ij} = 0 & Z(x^i y^j \bar{\alpha})\psi' = +\psi' \\ c_{ij} = 1 & Z(x^i y^j \bar{\alpha})\psi' = -\psi' \end{array}$$

such that a quasi-particle is present at (i, j) if and only if $c_{ij} = 1$.

Excitations in classical spin liquids are caused by Pauli- X spin flips. Consider quasi-particle excitations caused by $X(e(x, y))$ where $e(x, y)$ are polynomials representing positions of spin flips. Since anti-commutations between $e(x, y)$ and $Z(x^i y^j \bar{\alpha})$ create quasi-particles at (i, j) , the excitation polynomial is

$$E(x, y) = e(x, y)\alpha. \quad (19)$$

For instance, if $X_{0,0}$ with $e = 1$ is applied, one has multiple excitations $E(x, y) = \alpha$. Consider an isolated excitation at $(0, 0)$. An application of $X_{0,0}$ makes it propagate in the $\hat{y}$ direction to multiple excitations represented by $f(x)y$. So, quasi-particle excitations propagate via applications of $f(x)$ just like a one-dimensional cellular automaton. Since the model does not have gauge symmetries, one cannot associate conserved charge to quasi-particle excitations. Indeed, a single quasi-particle may split into multiple quasi-particles of the same type. The energy barrier between degenerate ground states is $O(\log(L))$ which leads to quasi-glassy thermodynamic relaxation as shown by Newman and Moore [22].




FIG. 5: Propagation of quasi-particle excitations and violation of charge conservation.

D. Limit cycle under RG transformation

Classical fractal liquids have an interesting symmetry property concerning how their ground states behave under RG transformations. Let us consider the Sierpinski triangle model for $L = 8$. A ground state is

$$\psi = \begin{bmatrix} 1, & 0, & 0, & 0, & 0, & 0, & 0, & 0 \\ 1, & 1, & 0, & 0, & 0, & 0, & 0, & 0 \\ 1, & 0, & 1, & 0, & 0, & 0, & 0, & 0 \\ 1, & 1, & 1, & 1, & 0, & 0, & 0, & 0 \\ 1, & 0, & 0, & 0, & 1, & 0, & 0, & 0 \\ 1, & 1, & 0, & 0, & 1, & 1, & 0, & 0 \\ 1, & 0, & 1, & 0, & 1, & 0, & 1, & 0 \\ 1, & 1, & 1, & 1, & 1, & 1, & 1, & 1 \end{bmatrix}.$$

RG transformations, denoted by RG_{ij} ($i, j = 0, 1$), pick up spins at (x, y) with $x = i \pmod{2}$ and $y = j \pmod{2}$:

$$\begin{aligned} \text{RG}_{0,0}(\psi) &= \begin{bmatrix} 1, & 0, & 0, & 0 \\ 1, & 1, & 0, & 0 \\ 1, & 0, & 1, & 0 \\ 1, & 1, & 1, & 1 \end{bmatrix}, & \text{RG}_{0,1}(\psi) &= \begin{bmatrix} 0, & 0, & 0, & 0 \\ 0, & 0, & 0, & 0 \\ 0, & 0, & 0, & 0 \\ 0, & 0, & 0, & 0 \end{bmatrix} \\ \text{RG}_{1,0}(\psi) &= \begin{bmatrix} 1, & 0, & 0, & 0 \\ 1, & 1, & 0, & 0 \\ 1, & 0, & 1, & 0 \\ 1, & 1, & 1, & 1 \end{bmatrix}, & \text{RG}_{1,1}(\psi) &= \begin{bmatrix} 1, & 0, & 0, & 0 \\ 1, & 1, & 0, & 0 \\ 1, & 0, & 1, & 0 \\ 1, & 1, & 1, & 1 \end{bmatrix}. \end{aligned}$$

and

$$\text{RG}_{0,0}(\psi(1)) = \begin{bmatrix} 1, & 0, & 0, & 0, & 0, & 0, & 0, & 0, & 0, & \dots \\ 1, & 1, & 0, & 0, & 0, & 0, & 0, & 0, & 0, & \dots \\ 1, & 2, & 1, & 0, & 0, & 0, & 0, & 0, & 0, & \dots \\ 1, & 0, & 0, & 1, & 0, & 0, & 0, & 0, & 0, & \dots \\ 1, & 1, & 0, & 1, & 1, & 0, & 0, & 0, & 0, & \dots \\ 1, & 2, & 1, & 1, & 2, & 1, & 0, & 0, & 0, & \dots \\ 1, & 0, & 0, & 2, & 0, & 0, & 1, & 0, & 0, & \dots \\ 1, & 1, & 0, & 2, & 2, & 0, & 1, & 1, & 0, & \dots \\ 1, & 2, & 1, & 2, & 1, & 2, & 1, & 2, & 1, & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}, \quad \text{RG}_{1,2}(\psi(1)) = \begin{bmatrix} 2, & 0, & 0, & 0, & 0, & 0, & 0, & 0, & 0, & \dots \\ 2, & 2, & 0, & 0, & 0, & 0, & 0, & 0, & 0, & \dots \\ 2, & 1, & 2, & 0, & 0, & 0, & 0, & 0, & 0, & \dots \\ 2, & 0, & 0, & 2, & 0, & 0, & 0, & 0, & 0, & \dots \\ 2, & 2, & 0, & 2, & 2, & 0, & 0, & 0, & 0, & \dots \\ 2, & 1, & 2, & 2, & 1, & 2, & 0, & 0, & 0, & \dots \\ 2, & 0, & 0, & 1, & 0, & 0, & 2, & 0, & 0, & \dots \\ 2, & 2, & 0, & 1, & 1, & 0, & 2, & 2, & 0, & \dots \\ 2, & 1, & 2, & 1, & 2, & 1, & 2, & 1, & 2, & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

For the Fibonacci model $f = 1 + x + x^2$ over $\mathbb{F}_2$, one has:

$$\psi(1) = \begin{bmatrix} 1, & 0, & 0, & 0, & 0, & 0, & 0, & 0, & 0, & \dots \\ 1, & 1, & 1, & 0, & 0, & 0, & 0, & 0, & 0, & \dots \\ 1, & 0, & 1, & 0, & 1, & 0, & 0, & 0, & 0, & \dots \\ 1, & 1, & 0, & 1, & 0, & 1, & 1, & 0, & 0, & \dots \\ 1, & 0, & 0, & 0, & 1, & 0, & 0, & 0, & 1, & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

and

$$\text{RG}_{00}(\psi(1)) = \begin{bmatrix} 1, & 0, & 0, & 0, & 0, & 0, & 0, & 0, & 0, & \dots \\ 1, & 1, & 1, & 0, & 0, & 0, & 0, & 0, & 0, & \dots \\ 1, & 0, & 1, & 0, & 1, & 0, & 0, & 0, & 0, & \dots \\ 1, & 1, & 0, & 1, & 0, & 1, & 1, & 0, & 0, & \dots \\ 1, & 0, & 0, & 0, & 1, & 0, & 0, & 0, & 1, & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}, \quad \text{RG}_{01}(\psi(1)) = \begin{bmatrix} 1, & 1, & 0, & 0, & 0, & 0, & 0, & 0, & 0, & \dots \\ 1, & 0, & 0, & 1, & 0, & 0, & 0, & 0, & 0, & \dots \\ 1, & 1, & 1, & 1, & 1, & 1, & 0, & 0, & 0, & \dots \\ 1, & 0, & 1, & 1, & 1, & 1, & 0, & 1, & 0, & \dots \\ 1, & 1, & 0, & 0, & 1, & 1, & 0, & 0, & 1, & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

Discrete scale symmetries provide an useful algorithm to compute the fractal dimension of $\mathbf{f}(x, y)$ [44]. We illustrate the algorithm for the Fibonacci model: $f = 1 + x + x^2$ over $\mathbb{F}_2$. We denote a ground state with an initial condition γ as $\psi(\gamma)$. Then, renormalization of ground states $\psi_m(1)$ and $\psi_m(1+x)$ gives the following ground states for $L = 2^{m-1}$:

$$\begin{aligned} \text{RG}_{00}(\psi_m(1)) &= \psi_{m-1}(1) & \text{RG}_{00}(\psi_m(1+x)) &= \psi_{m-1}(1) \\ \text{RG}_{10}(\psi_m(1)) &= \psi_{m-1}(0) & \text{RG}_{10}(\psi_m(1+x)) &= \psi_{m-1}(1) \\ \text{RG}_{01}(\psi_m(1)) &= \psi_{m-1}(1+x) & \text{RG}_{01}(\psi_m(1+x)) &= \psi_{m-1}(1) \\ \text{RG}_{11}(\psi_m(1)) &= \psi_{m-1}(1) & \text{RG}_{11}(\psi_m(1+x)) &= \psi_{m-1}(x) \end{aligned}$$

Let us denote the weights of $\psi_m(1)$ and $\psi_m(1+x)$ as A_m and B_m . Then, one has

$$\begin{pmatrix} A_m \\ B_m \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 4 & 0 \end{pmatrix} \begin{pmatrix} A_{m-1} \\ B_{m-1} \end{pmatrix}. \quad (21)$$

This matrix has eigenvalues $1 \pm \sqrt{5}$, and thus, A_m and B_m scale as $O(L^{\frac{\log 1 + \sqrt{5}}{\log 2}})$ for large L .

The above RG transformations concern classical fractal liquids on a finite lattice. If one performs RG transformations on an infinite lattice, $\text{RG}_{ij}(\psi)$ becomes a group operation where $\text{RG}_{ij}(\psi)$ is a linear map inside the ground state space. In the case of a ferromagnet, the RG functions are always trivial; $\text{RG}_{ij}(\psi) = \psi$ since ψ is spatially uniform. Yet, for classical fractal liquids, translation symmetries of ground states are strongly broken, and as a result, $\text{RG}_{ij}(\psi)$ may be different from ψ in general.

This gives an interesting possibility of limit cycle behaviors under RG transformations. Consider $f = 1 + x$ over $\mathbb{F}_3$.

Let us apply an RG transformation for a ground state $\psi(1) = \mathbf{f}(x, y)$ where $\mathbf{f}(x, y) = 1 + fy + f^2y^2 + \dots$ is defined on a infinite lattice. Then, $\text{RG}_{12}(\psi)$ gives the following sequence

$$\psi(1) \rightarrow \psi(2) \rightarrow \psi(1) \rightarrow \psi(2) \rightarrow \dots$$

where a ground state $\psi(1)$ jumps to a different ground state $\psi(2)$, and the RG sequence exhibits a limit-cycle behavior. Next, for $f = 1 + x + x^2$ over $\mathbb{F}_2$, consider a ground state $\psi(1) = \mathbf{f}(x, y)$. Then, one has the following sequence under RG_{01} :

$$\psi(1) \rightarrow \psi(1+x) \rightarrow \psi(1) \rightarrow \psi(1+x) \rightarrow \dots$$

which is also a limit cycle.

Finally, consider $f = 1 + x + x^2$ over $\mathbb{F}_5$. We list some of its ground states as follows:

$$\begin{bmatrix} 1 \\ 1 \ 1 \ 1 \\ 1 \ 2 \ 3 \ 2 \ 1 \\ 1 \ 3 \ 1 \ 2 \ 1 \ 3 \ 1 \\ 1 \ 4 \ 0 \ 1 \ 4 \ 1 \ 0 \ 4 \ 1 \end{bmatrix} \begin{bmatrix} 1 \ 1 \\ 1 \ 2 \ 2 \ 1 \\ 1 \ 3 \ 0 \ 0 \ 3 \ 1 \\ 1 \ 4 \ 4 \ 3 \ 3 \ 4 \ 4 \ 1 \\ 1 \ 0 \ 4 \ 1 \ 0 \ 0 \ 1 \ 4 \ 0 \ 1 \end{bmatrix} \begin{bmatrix} 1 \ 2 \\ 1 \ 3 \ 3 \ 2 \\ 1 \ 4 \ 2 \ 3 \ 0 \ 2 \\ 1 \ 0 \ 2 \ 4 \ 0 \ 0 \ 2 \ 2 \\ 1 \ 1 \ 3 \ 1 \ 1 \ 4 \ 2 \ 4 \ 4 \ 2 \end{bmatrix} \\ \begin{bmatrix} 1 \ 3 \\ 1 \ 4 \ 4 \ 3 \\ 1 \ 0 \ 4 \ 1 \ 2 \ 3 \\ 1 \ 1 \ 0 \ 0 \ 2 \ 1 \ 0 \ 3 \\ 1 \ 2 \ 2 \ 1 \ 2 \ 3 \ 3 \ 4 \ 3 \ 3 \end{bmatrix} \begin{bmatrix} 1 \ 4 \\ 1 \ 0 \ 0 \ 4 \\ 1 \ 1 \ 1 \ 4 \ 4 \ 4 \\ 1 \ 2 \ 3 \ 2 \ 4 \ 2 \ 3 \ 4 \\ 1 \ 3 \ 1 \ 2 \ 4 \ 3 \ 4 \ 4 \ 2 \ 4 \end{bmatrix}.$$

A transformation RG_{02} generates the following limit cycle and fixed point:

$$1 \rightarrow 1+x \rightarrow 1+3x \rightarrow 1+2x \rightarrow 1, \quad 1+4x \rightarrow 1+4x$$

where ground states are represented by their initial conditions. A transformation RG_{04} leads to

$$1 \rightarrow 1+x \rightarrow 1, \quad 1+2x \rightarrow 1+4x \rightarrow 1+3x \rightarrow 1+3x.$$

These sequences are shown in Fig. 6.




FIG. 6: Limit cycles in RG transformations for a classical fractal liquid with $f = 1 + x + x^2$ over $\mathbb{F}_5$. (a) RG_{02} . (b) RG_{04} .

How ground states transform under RG transformations can be formulated as a transition rule of polynomials. Let us define RG functions for one-dimensional polynomials, denoted by RG_j as follows; for $\gamma = \sum_i c_i x^i$, one has

$$\text{RG}_j(\gamma) = \sum_i c_{pi+j} x^i. \quad (22)$$

In other words, we pick up coefficients of x^{pi+j} . Let us consider a ground state $\psi(\gamma)$. Then, its renormalization is

given by:

$$\text{RG}_{ij}(\psi(\gamma)) = \psi(\gamma'), \quad \gamma' = \text{RG}_j(\gamma f^i). \quad (23)$$

We do not know whether limit cycle behaviors lead to some interesting physical phenomena that are observable. Perhaps quantum criticality of classical fractal liquids (if exists) may have some reminiscent of limit cycles. We have considered wave-function renormalization so far. If one performs Hamiltonian renormalization, one obtains p copies of the original Hamiltonian after each transformation.

IV. QUANTUM FRACTAL LIQUID

A. $\mathbb{Z}_2$ spin liquid and polynomial

In this section, we present a general framework to construct a family of quantum fractal liquids which are condensation of fractal objects. We begin by representing $\mathbb{Z}_2$ spin liquid by polynomials. Following [43], we group two qubits into a single composite particle (Fig. 7) such that composite particles live on vertices of a square lattice and obtain:

$$S_{i,j}^{(Z)} = \begin{bmatrix} Z_A Z_B & Z_A \\ Z_B & I \end{bmatrix} \quad S_{i,j}^{(X)} = \begin{bmatrix} I & X_A \\ X_B & X_A X_B \end{bmatrix}$$

where each qubit inside a composite particle is labelled by A and B . In polynomial representation, the parent Hamiltonian is

$$H = - \sum_{ij} Z \begin{pmatrix} x^i y^j (1+x) \\ x^i y^j (1+y) \end{pmatrix} - \sum_{ij} X \begin{pmatrix} x^i y^j (1+y^{-1}) \\ x^i y^j (1+x^{-1}) \end{pmatrix}$$

where the upper (lower) row represents Pauli operators acting on A (B). Interaction terms are translations of

$$Z \begin{pmatrix} 1+x \\ 1+y \end{pmatrix}, \quad X \begin{pmatrix} 1+y^{-1} \\ 1+x^{-1} \end{pmatrix}$$

In this form, it is immediate to see that $\mathbb{Z}_2$ spin liquid consists of one-dimensional ferromagnets at its one-dimensional limit since classical Hamiltonians generated by $Z(1+x)$ and $Z(1+y)$ correspond to one-dimensional ferromagnets in the $\hat{x}$ and $\hat{y}$ directions respectively.




FIG. 7: Reduction of $\mathbb{Z}_2$ spin liquid. Two qubits are grouped into a composite particle.

Logical operators are

$$\begin{aligned}\ell_0^{(Z)} &= Z \begin{pmatrix} 0 \\ 1+x+x^2+\dots \end{pmatrix}, & \ell_1^{(Z)} &= Z \begin{pmatrix} 1+y+y^2+\dots \\ 0 \end{pmatrix} \\ \ell_0^{(X)} &= X \begin{pmatrix} 0 \\ 1+y+y^2+\dots \end{pmatrix}, & \ell_1^{(X)} &= X \begin{pmatrix} 1+x+x^2+\dots \\ 0 \end{pmatrix}\end{aligned}$$

It is worth representing them graphically as follows:

$$\ell_0^{(Z)} = \begin{bmatrix} Z_B & Z_B & \cdots & Z_B & Z_B \\ I & I & \cdots & I & I \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ I & I & \cdots & I & I \\ I & I & \cdots & I & I \end{bmatrix}, \quad \ell_0^{(X)} = \begin{bmatrix} Z_A & I & I & \cdots & I \\ Z_A & I & I & \cdots & I \\ \vdots & \vdots & \vdots & \ddots & I \\ Z_A & I & I & \cdots & I \\ Z_A & I & I & \cdots & I \end{bmatrix}$$

and

$$\ell_0^{(X)} = \begin{bmatrix} X_B & I & I & \cdots & I \\ X_B & I & I & \cdots & I \\ \vdots & \vdots & \vdots & \ddots & I \\ X_B & I & I & \cdots & I \\ X_B & I & I & \cdots & I \end{bmatrix}, \quad \ell_1^{(X)} = \begin{bmatrix} X_A & X_A & \cdots & X_A & X_A \\ I & I & \cdots & I & I \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ I & I & \cdots & I & I \\ I & I & \cdots & I & I \end{bmatrix}.$$

One can see that logical operators commute with interaction terms by computing commutation polynomials. For instance, $\ell_0^{(Z)} = Z(0, 1+x+x^2+\dots)^T$ commutes with $X(1+y^{-1}, 1+x^{-1})$ since $(1+x+x^2+\dots)(1+x) = 0$.

One can generalize construction of $\mathbb{Z}_2$ spin liquid with arbitrary polynomials $\alpha(x, y, \dots)$ and $\beta(x, y, \dots)$:

$$Z \begin{pmatrix} \alpha \\ \beta \end{pmatrix}, \quad X \begin{pmatrix} \bar{\beta} \\ \bar{\alpha} \end{pmatrix} \quad (24)$$

where $\bar{\alpha}$ and $\bar{\beta}$ are duals of α and β obtained by taking $x \rightarrow x^{-1}$, $y \rightarrow y^{-1}$, $\dots$. Note that interaction terms commute with each other as their commutation polynomial is $\alpha\beta + \beta\alpha = 0$ over $\mathbb{F}_2$. A parent Hamiltonian is

$$H = - \sum_{ij\dots} Z \begin{pmatrix} x^i y^j \cdots \alpha \\ x^i y^j \cdots \beta \end{pmatrix} - \sum_{ij\dots} X \begin{pmatrix} x^i y^j \cdots \bar{\beta} \\ x^i y^j \cdots \bar{\alpha} \end{pmatrix}.$$

Various physical properties of above Hamiltonians can be discussed from algebraic arguments on α and β . See appendix A for further discussion.

B. Quantum fractal liquid

Consider a three-dimensional $L \times L \times L$ square lattice where two qubits live on each site with $L = 2^m$ and periodic boundary conditions. Quantum fractal liquids have

$$\alpha = 1 + f(x)y, \quad \beta = 1 + g(x)z \quad (25)$$

where $f(x)$ and $g(x)$ are reversible polynomials over $\mathbb{F}_2$. More explicitly, interaction terms are translations of

$$Z \begin{pmatrix} 1 + f(x)y \\ 1 + g(x)z \end{pmatrix}, \quad X \begin{pmatrix} 1 + \bar{g}(x)\bar{z} \\ 1 + \bar{f}(x)\bar{y} \end{pmatrix}. \quad (26)$$

Therefore, interaction terms are characterized by a pair of fractal generators $f(x)$ and $g(x)$. In this sense, quantum fractal liquids can be viewed as a coherent combination of a pair of classical fractal liquids living on $(\hat{x}, \hat{y})$ -plane and

$(\hat{x}, \hat{z})$ -plane respectively. Generalization to polynomials over $\mathbb{F}_p$ is straightforward.

Logical operators of quantum fractal liquids have fractal shapes which are generated by polynomials $f(x)$ and $g(x)$:

$$\begin{aligned} \mathbf{f}(x, y) &= 1 + fy + f^2y^2 + \dots & \bar{\mathbf{f}}(x, y) &= 1 + \bar{f}\bar{y} + \bar{f}^2\bar{y}^2 + \dots \\ \mathbf{g}(x, z) &= 1 + gz + g^2z^2 + \dots & \bar{\mathbf{g}}(x, z) &= 1 + \bar{g}\bar{z} + \bar{g}^2\bar{z}^2 + \dots \end{aligned}$$

Note $\mathbf{f}(x, y)$ lies on a $(\hat{x}, \hat{y})$ -plane while $\mathbf{g}(x, z)$ lies on a $(\hat{x}, \hat{z})$ -plane. Quantum fractal liquids have $k = 2L$, and there are $2L$ of Z -type logical operators and $2L$ of X -type logical operators:

$$\begin{aligned} \ell_i^{(Z)} &= Z \begin{pmatrix} 0 \\ x^i \mathbf{f}(x, y) \end{pmatrix}, & r_i^{(Z)} &= Z \begin{pmatrix} x^i \mathbf{g}(x, z) \\ 0 \end{pmatrix} \\ \ell_i^{(X)} &= X \begin{pmatrix} x^i \bar{\mathbf{f}}(x, y) \\ 0 \end{pmatrix}, & r_i^{(X)} &= X \begin{pmatrix} 0 \\ x^i \bar{\mathbf{g}}(x, z) \end{pmatrix} \end{aligned}$$

where $i = 0, \dots, L-1$. Therefore, Z -type logical operators have geometric shapes of $\mathbf{f}(x, y)$ and $\mathbf{g}(x, y)$ while X -type logical operators have geometric shapes of $\bar{\mathbf{f}}(x, y)$ and $\bar{\mathbf{g}}(x, y)$.

To show that above operators are logical operators, we need to verify the following two things; a) they commute with interaction terms, b) they can be grouped into pairs of anti-commuting logical operators. One may see that logical operators commute with all the interaction terms by computing commutation polynomials. For instance, a commutation polynomial between $\ell_0^{(Z)} = Z(0, \mathbf{f}(x, y))^T$ and stabilizers $X(1 + \bar{g}\bar{y}, 1 + \bar{f}\bar{x})^T$ is given by $(1 + fy)\mathbf{f}(x, y) = 0$. Logical operators obey the following commutation relations:

$$\left\{ \ell_0^{(Z)}, \dots, \ell_{L-1}^{(Z)}, r_0^{(Z)}, \dots, r_{L-1}^{(Z)}, \right. \\ \left. r_0^{(X)}, \dots, r_{L-1}^{(X)}, \ell_0^{(X)}, \dots, \ell_{L-1}^{(X)} \right\}.$$

To see that quantum fractal liquids are topologically ordered, we begin by showing that they are good quantum error-correcting code with $d \rightarrow \infty$ for $L \rightarrow \infty$ where d is the quantum code distance of the ground state space. A standard way to prove this considers a bi-partition of the system into two complementary subsets A and B and uses the following bi-partition formula which holds for arbitrary stabilizer codes [45]:

$$g_A + g_B = 2k \tag{27}$$

where g_A and g_B represent the number of independent logical operator supported inside A and B respectively. Let us assume A to be a connected region with finite support. Then, its complementary subset B accommodates some $(\hat{x}, \hat{y})$ -plane and $(\hat{x}, \hat{z})$ -plane where all the $2k$ independent logical operators can be supported. So, one has $g_B = 2k$. This leads to $g_A = 0$. Therefore, weights of logical operators are not finite (i.e. unbounded), and $d \rightarrow \infty$ for $L \rightarrow \infty$. Another way uses a criteria presented in appendix A (see [20, 25] also). Quantum fractal liquids are characterized by a pair of polynomials $\alpha = 1 + f(x)y$ and $\beta = 1 + g(x)z$ which do not have common factors. Therefore, the system always has $d \rightarrow \infty$ for $L \rightarrow \infty$ (as long as $k \neq 0$).

For stabilizer Hamiltonians, being a quantum code ($d \rightarrow \infty$ for $L \rightarrow \infty$) automatically implies the presence of topological order with stability against local perturbations. In [33], Bravyi, Hastings and Michalakis proved that frustration-free Hamiltonians with an ability of quantum error-correcting code have stability against local perturbations when Hamiltonians satisfy a certain condition, called TQO-2. Roughly speaking, TQO-2 states that locally computed density matrices are consistent with ground states which are computed globally. One can check that quantum fractal liquids satisfy TQO-2, and thus have stability against local perturbations. Recall that quantum fractal liquids have 2^L ground states. Under a small but finite local perturbations, the energy splitting among these ground states is always exponentially suppressed, and the energy gap between the ground states and excited states remains finite. Note quantum codes with TQO-2 condition have efficient decoding algorithm based on RG-like transformations [25].

We then discuss the number of degenerate ground states and its dependence on the system size. A key feature of quantum fractal liquids is that the number of logical qubits k has a fairly sensitive dependence on the system size $\vec{L} = (L_1, L_2, L_3)$. The number of logical qubits k is given by counting the dimension of solutions γ satisfying the

following equation:

$$f(x)^{L_2}\gamma(x) = g(x)^{L_3}\gamma(x) = \gamma(x), \quad x^{L_1} = 1. \quad (28)$$

with $k = 2 \dim \gamma$. For instance, with $f = x^i$ and $g = x^j$, one has $k = 2 \gcd(L_1, iL_2, jL_3)$ where k depends crucially on the system size $\vec{L} = (L_1, L_2, L_3)$. In general, it is a very challenging task to write down an explicit form of $k(L_1, L_2, L_3)$ for a given pair of $f(x)$ and $g(x)$. Yet, $k(L_1, L_2, L_3)$ has a nice symmetry property under scale transformations:

$$k(pL_1, pL_2, pL_3) = pk(L_1, L_2, L_3). \quad (29)$$

The proof is straightforward from discrete scale symmetries of polynomials over $\mathbb{F}_p$. See appendix A for quantum fractal liquids with open boundary conditions.

Ground states of quantum fractal liquids correspond to limit cycles under real-space RG transformations on an infinite lattice. To obtain RG transformations with meaningful attractors that do not flow to disordered states or trivial product states, one needs to apply some appropriate projection operators on sites that are to be coarse-grained. Consider a pair of qubits at site (i, j, ℓ) , denoted as $|\phi\rangle_{ij\ell} = |\phi_A\rangle_{ij\ell} \otimes |\phi_B\rangle_{ij\ell}$, and apply the following projections to a ground state:

$$(I + Z_A^\ell \otimes Z_B^j)(I + X_A^j \otimes X_B^\ell)|\phi_A\rangle_{ij\ell} \otimes |\phi_B\rangle_{ij\ell}. \quad (30)$$

Note that projection operators commute with each other, and projections are applied only to sites (i, j, ℓ) with $j \neq 0$ or $\ell \neq 0$ modulo 2. As a result, pairs of qubits on sites (i, j, ℓ) with $j = \ell = 0$ modulo 2 are completely decoupled from the rest where projectors are applied. With some speculation, one notices that stabilizer generators for remaining sites (i, j, ℓ) with $j = \ell = 0$ modulo 2 are given by

$$Z \begin{pmatrix} \alpha^2 \\ \beta^2 \end{pmatrix}, \quad Z \begin{pmatrix} x\alpha^2 \\ x\beta^2 \end{pmatrix}, \quad X \begin{pmatrix} \bar{\beta}^2 \\ \bar{\alpha}^2 \end{pmatrix}, \quad X \begin{pmatrix} x\bar{\beta}^2 \\ x\bar{\alpha}^2 \end{pmatrix}$$

and their translations that are generated by applications of $x^{2i'}y^{2j'}z^{2\ell'}$. (See [46] for transformations of stabilizer generators under projections). This corresponds to two copies of original quantum fractal liquids. Let us pick up sites with $(i, j, \ell) = (0, 0, 0)$ modulo 2 and throw away sites with $(i, j, \ell) = (1, 0, 0)$ modulo 2 via some arbitrary projections. Taking $x^2 \rightarrow x$, $y^2 \rightarrow y$ and $z^2 \rightarrow z$, stabilizer generators are $Z(\alpha, \beta)^T$ and $X(\bar{\beta}, \bar{\alpha})^T$. So, this RG transformation maps a ground state of a quantum fractal liquid into some ground state which may be different from the original. One can keep track of how ground states flow in RG transformations by looking at polynomial representation of fractal logical operators which can be analyzed in a way similar to ground states of classical fractal liquids. If one performs Hamiltonian renormalization, instead of wave-function renormalization, one obtains two copies of the original Hamiltonian after each transformation. For quantum fractal liquids with higher-order cellular automata rules, RG flows become more complicated.

C. Quasi-particles and no-string rule

We discuss properties of quasi-particle excitations in quantum fractal liquids. Without loss of generality, one can concentrate on excitations caused by Pauli- Z type operators which flips X -type interaction terms. Following a treatment of classical fractal liquids, we represent positions of excitations as a polynomial:

$$E(x, y, z) = \sum_{i,j,\ell} c_{ij\ell} x^i y^j z^\ell$$

where $c_{ij\ell} = 1$ means an excitation is present at (i, j, ℓ) . We consider excitations caused by a Pauli operator $Z(e_A, e_B)^T$ which is given by a commutation polynomial between $(e_A, e_B)^T$ and $(\bar{\beta}, \bar{\alpha})^T$:

$$E(x, y, z) = P \left[\begin{pmatrix} e_A \\ e_B \end{pmatrix}, \begin{pmatrix} 1 + \bar{g}\bar{z} \\ 1 + \bar{f}\bar{y} \end{pmatrix} \right] = e_A(1 + gz) + e_B(1 + fy). \quad (31)$$

Excitations caused by a Pauli operator $Z_{0,0,0}^{(A)}$ ($e_A = 1, e_B = 0$) are given by

$$E(x, y, z) = (1 + gz) \quad (32)$$

while excitations caused by a Pauli operator $Z_{0,0,0}^{(B)}$ ($e_A = 0, e_B = 1$) are given by

$$E(x, y, z) = (1 + fy). \quad (33)$$

So, if an isolated excitation is present at $(i, j, \ell) = (0, 0, 0)$, it will propagate to multiple excitations represented by gz via an application of $Z_{0,0,0}^{(A)}$ (Fig. 8(a)). Similarly, it will propagate to multiple excitations represented by fy via an application of $Z_{0,0,0}^{(B)}$ (Fig. 8(a)). In general, for a single isolated excitation, $f(x)$ is applied when propagating in the $\hat{y}$ direction, and $g(x)$ is applied when propagating in the $\hat{z}$ direction. Let us consider a one-dimensional excitation pattern, represented by $e(x)$, located at $j = \ell = 0$, and think of moving it to $j = j'$ and $\ell = \ell'$ while keeping the pattern of excitations one-dimensional. Then, the resulting excitation pattern is given by

$$E(x, y, z) = e(x)f(x)^{j'}g(x)^{\ell'}y^{j'}z^{\ell'}. \quad (34)$$

This may be viewed as time evolution of an initial condition $e(x)$, updated j' times by $f(x)$ and ℓ' times by $g(x)$ respectively.




FIG. 8: (a) Propagation of quasi-particles by $f(x)$ and $g(x)$ (b) A pair of localized excitations e_1 and e_2 with elongated excitations e'_1 and e'_2

Let us consider simple cases with $f(x) = g(x)$ and consider propagations of an excitation originally at $(0, 0, 0)$ to $j = j'$ and $\ell = \ell'$. Then, the resulting excitation pattern is given by

$$E(x, y, z) = f(x)^{j'}g(x)^{\ell'}y^{j'}z^{\ell'} = f(x)^{j'+\ell'}y^{j'}z^{\ell'}. \quad (35)$$

So, if $j' + \ell' = 0$, excitations are single quasi-particles, and excitation energy remains finite. Therefore, quasi-particle excitations can propagate freely (without costing much energy) in the $\hat{y} - \hat{z}$ direction which implies the presence of a string-like logical operator. Indeed, the following string-like operators are logical operators:

$$Z \begin{pmatrix} 0 \\ (y+z)^{L-1} \end{pmatrix}, \quad X \begin{pmatrix} (\bar{y} + \bar{z})^{L-1} \\ 0 \end{pmatrix}.$$

Note that $(y+z)^{L-1}$ is a string-like polynomial extending in the $\hat{y} - \hat{z}$ direction.

A naturally arising question is when quasi-particle excitations propagate freely and quantum fractal liquids have string-like logical operators. Without loss of generality, one can assume that $f(x)$ and $g(x)$ start from the origin, meaning that $f(x)$ and $g(x)$ have non-zero constant terms and have only positive powers. We say that $f(x)$ and $g(x)$

are *algebraically related* when there exists some finite integers c_f and c_g such that

$$f(x)^{c_f} = g(x)^{c_g} \quad (36)$$

without considering periodic boundary conditions. Then, one notices that quasi-particles can propagate in the $c_a \hat{y} - c_b \hat{z}$ direction. Therefore, there exist string-like logical operators.

It turns out that Eq. (36) is a necessary and sufficient condition for the presence of string-like logical operators:

$$\text{“No string-like logical operator”} \Leftrightarrow \text{“Polynomials } f(x) \text{ and } g(x) \text{ are not algebraically related”}. \quad (37)$$

To verify this equivalence relation, let us consider a pair of localized excitations $e_1(x, y, x)$ and $e_2(x, y, x)$ which are contained in a cubic box of $w \times w \times w$ sites with $w \ll L$, which are separated by L_δ . We assume L_δ is sufficiently larger than w , and sufficiently smaller than L . We ask whether e_1 and e_2 are connected by a string-like object or not. One can make quasi-particle excitations in $e_1(x, y, z)$ and $e_2(x, y, z)$ propagate by applying $f(x)$ and $g(x)$ such that they become some elongated line-like excitations whose lengths are $O(w)$ (see Fig. 8(b)). We denote polynomials corresponding to these elongated excitations as e'_1 and e'_2 , and assume that e'_1 is at $j = \ell = 0$ and e'_2 is at $j = j'$ and $\ell = \ell'$ where $|j'| + |\ell'| \sim O(L_\delta)$. One may write $e'_1 = e_1^*(x)$ and $e'_2 = e_2^*(x)y^{j'}z^{\ell'}$. Then, one must have $e_2^* = e_1^* f(x)^{j'} g(x)^{\ell'}$ in order to connect these excitations by some object. If a pair of excitations is connected by a string-like object, $f^{j'} g^{\ell'}$ must remain finite for large j' and ℓ' . This requires f and g to be algebraically related; otherwise, one can prove that the size of $f^{j'} g^{\ell'}$ grows at least linearly as $|j'|$ and $|\ell'|$ grow. So, a pair of isolated excitations is not connected by a string-like object, and there is no string-like logical operators. One may construct a formal proof by following an argument in [47].

When there is no string-like logical operators, energy barrier separating different ground states is lower bounded by $O(\log L)$ as shown in [22, 47]. The presence of large energy barrier is favorable for storing logical qubits securely in thermal environment. For sufficiently small system size L where entropic contributions are negligible, the memory time of encoded logical qubits can be estimated by an Arrhenius law $\tau \sim \exp(-\Delta/T)$ where Δ is an energy barrier. For models without string-like logical operators, one has $\tau \sim \text{POLY}(L)$ which is an improvement over $\tau \sim O(1)$ of conventional models with string-like logical operators. For large L , however, one does not see any significant improvement since entropic contributions dominate over energetic constraints as the number of ground states and local minima increases. With this subtlety, fractal models without string-like logical operators are said to be marginally self-correcting [19]. This behavior is closely related to the thermal instability of fractal models as they undergo thermal phase transition at $T_c = 0$. So, the transition to quantum fractal glassy phase is a continuous one (i.e. a crossover), and is not sharp. See [48, 49] for proposals to overcome this challenge in quantum information community.

D. Several examples

Finally, we study several examples of quantum fractal liquids.

(a) We begin with a trivial case:

$$Z \left(\begin{array}{c} 1 + y \\ 1 + z \end{array} \right) \quad \text{over } \mathbb{F}_2$$

This is a stack of slices of two-dimensional $\mathbb{Z}_2$ model where each copy lives on a $(\hat{y}, \hat{z})$ -plane. It has pairs of string-like logical operators since $f = 1$ and $g = 1$ are generators of strings. Similarly, for $f = 1$ and $g = 1$ over $\mathbb{F}_p$ ($p > 2$), the model is a stack of $\mathbb{Z}_p$ spin liquids with p -dimensional qudits.

(b) For $f = x$ and $g = 1$, stabilizer generators are given by

$$Z \left(\begin{array}{c} 1 + xy \\ 1 + z \end{array} \right) \quad \text{over } \mathbb{F}_2.$$

This is a stack of slices of two-dimensional $\mathbb{Z}_2$ model, but each copy lives on a $(\hat{x} + \hat{y}, \hat{z})$ -plane. It has string-like logical operators, running in the $\hat{x} + \hat{y}$ direction. One can reduce this model to the model in (a) by a *modular transformation* $x \rightarrow x$, $xy \rightarrow y$ and $z \rightarrow z$ which corresponds to a lattice distortion. In general, two models are considered to be

equivalent when they are connected by modular transformations.

(c) Let us look at an example with pairs of fractal logical operators and string-like logical operators:

$$Z \left(\begin{array}{c} 1 + (1 + x + x^2)y \\ 1 + z \end{array} \right) \quad \text{over } \mathbb{F}_2,$$

the model has pairs of fractal logical operators and string-like logical operators. In particular, geometric shapes of fractal logical operators are generated by $f = 1 + x + x^2$ (the Fibonacci model). Note that fractal logical operators live on a two-dimensional plane and string-like logical operators are perpendicular to the two-dimensional plane.

(d) Some models do not have any logical qubits under periodic boundary conditions:

$$Z \left(\begin{array}{c} 1 + (1 + x)y \\ 1 + z \end{array} \right) \quad \text{over } \mathbb{F}_2$$

since $f = 1 + x$ is not reversible over $\mathbb{F}_2$. When the model is defined with open boundary conditions (as in appendix A), it has pairs of fractal logical operators and string-like logical operators.

(e) Consider

$$Z \left(\begin{array}{c} 1 + (1 + x)y \\ 1 + (1 + x)z \end{array} \right), \quad \text{over } \mathbb{F}_2.$$

One might think that the model has pairs of fractal logical operators, and is free from string-like logical operators. Yet, the model has hidden string-like logical operators running in the $\hat{y} - \hat{z}$ direction. In fact, this model is unitarily equivalent to the following model

$$Z \left(\begin{array}{c} 1 + (1 + x)y \\ 1 + y^{-1}z \end{array} \right), \quad \text{over } \mathbb{F}_2.$$

See appendix A for further discussion on local unitary transformations.

(f) Let us consider the cases without any string-like logical operators for $\mathbb{F}_2$:

$$Z \left(\begin{array}{c} 1 + (1 + x + x^2)y \\ 1 + (1 + x + x^3)z \end{array} \right) \quad \text{over } \mathbb{F}_2$$

The model is free from string-like logical operators as $f = 1 + x + x^2$ and $g = 1 + x + x^3$ are algebraically unrelated. Interaction terms are seven-body.

(g) Next, let us consider a model over $\mathbb{F}_p$ ($p > 2$):

$$Z \left(\begin{array}{c} 1 + (1 + x)y \\ 1 + (1 + x^2)z \end{array} \right) \quad \text{over } \mathbb{F}_3$$

The model is also free from string-like logical operators as $f = 1 + x$ and $g = 1 + x^2$ are algebraically unrelated. Interaction terms are five-body.

(h) The Cubic code can be viewed as a second-order quantum fractal liquid. In polynomial representation, one has

$$Z \left(\begin{array}{c} 1 + x + y + z \\ 1 + xy + yz + zx \end{array} \right) \quad \text{over } \mathbb{F}_2.$$

The model can be mapped to the following second-order quantum fractal liquid through local unitary transformations and modular transformations:

$$Z \left(\begin{array}{c} 1 + f(x)y \\ 1 + g_1(x)z + g_2(x)z^2 \end{array} \right) \quad \text{over } \mathbb{F}_2$$

where

$$f(x) = 1 + x + x^2, \quad g_1(x) = 1 + x, \quad g_2(x) = 1 + x + x^2.$$

See appendix A for further discussion.

V. TOPOLOGICAL ENTANGLEMENT ENTROPY

In this section, we compute entanglement entropies of quantum fractal liquids in order to establish a connection between ground state and excitation properties. We demonstrate that topological entanglement entropies, defined for tube-like regions with holes, can distinguish quantum fractal liquids from conventional topologically ordered systems.

We begin with a review of computational technique of entanglement entropies. For stabilizer Hamiltonians with topological order, calculation of entanglement entropies can be reduced to a simple counting argument. Consider an arbitrary stabilizer Hamiltonian, $H = -\sum_j S_j$ and a bi-partition into complementary subsets A and $B = \bar{A}$. We denote a group of Pauli operators generated by S_j as $\mathcal{S} = \langle \{S_j : \forall j\} \rangle$ and its restriction into A as $\mathcal{S}_A$:

$$\mathcal{S}_A = \langle \{U \in \mathcal{S} : U|_B = I\} \rangle \quad (38)$$

where $U|_B$ is a restriction of U on B . Note that $\mathcal{S}_A$ consists of all the stabilizer operators supported inside A . If the stabilizer Hamiltonian possesses topological order, there is no logical operator supported inside A , and one has

$$E_A = |A| - \dim \mathcal{S}_A \quad (39)$$

where E_A is an entanglement entropy, $|A|$ is the total number of qubits inside A , and $\dim \mathcal{S}_A$ is the number of independent generators for $\mathcal{S}_A$.

Let us now compute an entanglement entropy of $\mathbb{Z}_2$ spin liquid, represented by $Z(1+x, 1+y)^T$, for a region A with $w_1 \times w_2$ sites. One has $|A| = 2w_1w_2$ and $\dim \mathcal{S} = 2(w_1 - 1)(w_2 - 1)$, so $E_A = 2(w_1 + w_2) - 2$. There is a constant correction term to the area-law of entanglement entropies. Yet, this correction is not universal since it is dependent on shapes of boundaries. To obtain a universal correction term correctly, one needs to cancel out non-universal contributions by adding or subtracting entanglement entropies for several regions.




FIG. 9: Computations of topological entanglement entropies.

There are two standard methods for this [50, 51]. The first method considers the mutual information among three neighboring regions (Fig. 9(a)):

$$I(A, B, C) = E_A + E_B + E_C - E_{AB} - E_{BC} - E_{CA} + E_{ABC}. \quad (40)$$

Since $E_R = |R| - \dim \mathcal{S}_R$ for regions R , contribution from volumes $|R|$ exactly cancels. As for stabilizer generators, $U \in \mathcal{S}_A$ implies $U \in \mathcal{S}_{AB}, \mathcal{S}_{CA}, \mathcal{S}_{ABC}$. So, contribution from $U \in \mathcal{S}_A$ cancels. Only stabilizers $U \in \mathcal{S}_{ABC}$, but with $U \notin \mathcal{S}_A, \mathcal{S}_B, \mathcal{S}_C, \mathcal{S}_{AB}, \mathcal{S}_{BC}, \mathcal{S}_{CA}$ contribute to $I(A, B, C)$. Therefore, one needs to count the number of independent stabilizer generators with supports on all A , B and C . One may see that there is only one such stabilizer, so $I(A, B, C) = -1$. However, we note that there exist models of stabilizer Hamiltonians which are connected to product

states, but possess non-zero $I(A, B, C)$. For instance, a model with $\alpha = 1$ and $\beta = 1 + x + y + xy$ has $I(A, B, C) = -2$ though it is connected to a product state through local unitary transformations. This discrepancy is perhaps because boundaries of topologically ordered systems are not necessarily described by conformal field theory on a precise level.

The other method considers the following combinations of entanglement entropies (Fig. 9(b))

$$E^* = E_A - E_{B_1} - E_{B_2} + E_{C_1 \cup C_2} \quad (41)$$

where A is a loop-like region. One needs to find stabilizer generators with $U \in \mathcal{S}_A$ and $U \notin \mathcal{S}_{B_1}, \mathcal{S}_{B_2}, \mathcal{S}_{C_1 \cup C_2}$. To find such stabilizers, it is convenient to find identity generators R (see appendix A for detailed discussion):

$$R = \sum_{i,j} x^i y^j, \quad R\alpha = R\beta = 0$$

which creates an identity operator as $Z(R\alpha, R\beta)^T = I$. Consider a subpart of R , given by $R^* = \sum_{i,j=0}^{w-2} x^i y^j$. This creates the following loop-like stabilizer operator

$$Z \begin{pmatrix} R^* \alpha \\ R^* \beta \end{pmatrix} = \begin{bmatrix} Z_A Z_B & Z_B & \cdots & Z_B & Z_A \\ Z_A & I & \cdots & I & Z_A \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ Z_A & I & \cdots & I & Z_A \\ Z_B & Z_B & \cdots & Z_B & I \end{bmatrix} \quad (42)$$

which has supports on all the four regions B_1, B_2, C_1, C_2 . One can construct a similar operator from X -type stabilizers. So, one has $E^* = 2$. These stabilizer operators may be viewed as string-like extended objects. Since endpoints of loops correspond to quasi-particle excitations, these loop-like stabilizer operators characterize propagations of quasi-particles along the loop A . There are two independent quasi-particles circling around the loop in $\mathbb{Z}_2$ spin liquid.

Let us proceed to discussion on topological entanglement entropies of quantum fractal liquids in three-dimensional systems. There are four possible types of topological phases; (a) the three-dimensional Toric code, (b) a stack of slices of two-dimensional Toric code, (c) quantum fractal liquids with string-like logical operators, and (d) quantum fractal liquids without string-like logical operators. The goal is to distinguish these four phases. For this purpose, we consider topological entanglement entropies defined for a tube-like region with a hole inside it (Fig. 10). We assume that the width of the tube w is sufficiently large, but is small compared to L . We add and subtract entanglement entropies in a way similar to Fig. 9(b). We argue that, one can, in principle, distinguish four phases by measuring topological entanglement entropies for tube-like regions in various directions.

(a) The three-dimensional Toric code is

$$Z \begin{pmatrix} 1+y \\ 1+x \\ 0 \end{pmatrix} \quad Z \begin{pmatrix} 0 \\ 1+z \\ 1+y \end{pmatrix} \quad Z \begin{pmatrix} 1+z \\ 0 \\ 1+x \end{pmatrix} \quad X \begin{pmatrix} 1+\bar{x} \\ 1+\bar{y} \\ 1+\bar{z} \end{pmatrix}$$

with three qubits per site. (See appendix A for higher-dimensional Toric code). One can form a loop-like Z -type stabilizer by multiplying $Z(0, 1+z, 1+y)^T$ with $R^* = \sum_{j,\ell=0}^{w-2} y^j z^\ell$ and a tube-like X -type stabilizer by multiplying $X(1+\bar{x}, 1+\bar{y}, 1+\bar{z})^T$ with $R^* = \sum_{i=0}^{L-1} \sum_{j,\ell=0}^{w-2} x^i y^j z^\ell$. So, one has $E^* = 2$ since there are only two independent quasi-particles. One obtains the same result for tube-like regions extending in the $\hat{y}$ or $\hat{z}$ directions. So, topological entanglement entropies remain finite for tubes in any directions. This is related to the fact that propagations of quasi-particles are rotationally invariant at the thermodynamic limit and translation symmetries are not broken in any directions.

(b) We then compute topological entanglement entropies for a stack of slices of two-dimensional Toric code. To be specific, we assume that each copy of the Toric code lies on a $(\hat{y}, \hat{z})$ -plane. For a tube-like region extending in the $\hat{x}$ direction, one has $E^* = 2L$ which corresponds to the number of quasi-particles that can circle around the tube. Yet, for a tube-like region extending in the $\hat{y}$ direction, one has $E^* = 0$ since quasi-particles are constrained to $(\hat{y}, \hat{z})$ -planes and they can propagate “along” the tube, without circling around the tube. This is related to the fact that translation symmetries are strongly broken in the $\hat{x}$ direction.

(c) Next, we compute topological entanglement entropies for quantum fractal liquids with strings. To be specific, we consider the case with $f(x) \neq x^i$ and $g(x) = 1$ where string-like logical operators are along the $\hat{z}$ direction. Identity generators are

$$R = \gamma(x)\mathbf{f}(x, y)(1 + z + z^2 + \cdots)$$

where $\gamma(x)$ is an arbitrary polynomial with x . For a tube-like region extending in the $\hat{x}$ direction, one can form tube-like Z -type stabilizers with $R^* = \gamma(x) \sum_{i=0}^{L-1} \sum_{j,\ell=0}^{w-2} x^i (fy)^j z^\ell$. There are L independent tube-like Z -type stabilizers, so one has $E^* = 2L$. This corresponds to the total number of independent quasi-particles that can circle around the tube. In fact, one obtains the same result for a tube-like region extending in the $\hat{y}$ directions: $E^* = 2L$. Yet, for a tube extending in the $\hat{z}$ direction, with some speculations, one has $E^* = 0$ since quasi-particles can propagate only along the $\hat{z}$ direction, without circling around the tube. Note that translation symmetries are strongly broken in the $\hat{x}$ and $\hat{y}$ directions.

(d) Finally, we discuss quantum fractal liquids without strings. Identity generators are

$$R = \gamma(x)\mathbf{f}(x, y)\mathbf{g}(x, z).$$

For a tube-like region extending in the $\hat{x}$ direction, one can form L independent tube-like Z -types stabilizers, so $E^* = 2L$. In fact, one obtains the same result for tubes in the $\hat{y}$ and $\hat{z}$ directions: $E^* = 2L$. This is due to an observation that quasi-particles cannot propagate along the tube, regardless of its direction, and need to circle around the tube. Note that translation symmetries are broken in all the directions, yet, rotational invariance of topological entanglement entropies are persevered.

Below, we summarize the results:

	Tube in $\hat{x}$	Tube in $\hat{y}$	Tube in $\hat{z}$
3-dim Toric code	$O(1)$	$O(1)$	$O(1)$
2-dim Toric code	$O(L)$	0	0
Quantum fractal liquid (with string)	$O(L)$	$O(L)$	0
Quantum fractal liquid (no string)	$O(L)$	$O(L)$	$O(L)$

One may notice that topological entanglement entropies become zero or finite when string-like logical operators are present along the axis of the tube. So, by measuring topological entanglement entropies for tube-like regions in various directions, one may distinguish quantum fractal liquids from conventional topologically ordered systems in principle. (There are some artificial counterexamples to our claim. For instance, three copies of stacks of two-dimensional Toric codes, put on three different directions, have $O(L)$ topological entanglement entropies in all the three directions. Stability of these fractal order parameter is an important problem, but is beyond the scope of this paper). Another useful method of detecting quantum fractal order is to see if translation symmetries are strongly broken in the ground state space or not. One may be able to deduce fractal generators $f(x)$ and $g(x)$ by analyzing behaviors of $k(L_1, L_2, L_3)$.




FIG. 10: A tube-like region extending in the $\hat{x}$ direction.

VI. CONCLUSIONS

In this paper, we have presented general construction of classical and quantum fractal liquids and discussed their physical properties. We have found that correlation functions of classical fractal liquids have peculiar oscillatory behaviors which correspond to power-law decay with imaginary scaling dimensions. We have demonstrated that degenerate ground states of classical and quantum fractal liquids have discrete scale symmetries and correspond to limit cycles under real-space RG transformations. We have argued that quasi-particle excitations in classical and quantum fractal liquids violate charge conservation as they possess algebraic symmetries which are more general than gauge symmetries. We have derived a necessary and sufficient condition for quantum fractal liquids to be free from string-like logical operators by using an analogy to cellular automata. We have demonstrated that topological entanglement entropies can detect quantum fractal order in principle. We hope that our construction and theory of fractal spin liquids will provide a stepping stone toward complete understanding and classification of quantum phases of matter.

Quasi-particle excitations arising in quantum fractal liquids are all Abelian. Topological order arising in quantum fractal liquids is non-chiral. It is unclear to what extent our results may generalize to chiral topological phases and non-Abelian topological phases. Effective field theoretical descriptions of classical and quantum fractal liquids are currently not known. It may be interesting to analyze how classical and quantum fractal liquids behave under RG transformations in the language of matrix and tensor product state representations [52–55]. Connection to limit cycles in non-relativistic field theory has not been established. Classification of quantum phases arising in translation symmetric stabilizer Hamiltonians is an important problem. In [43], we proved that topological phases arising in two-dimensional translation symmetric Hamiltonians are equivalent to copies of the Toric code. Yet, classification of three-dimensional topological phases remains as an open problem. An underlying difficulty in physically realizing the Sierpinski triangle model lies in the fact that the model has three-body interaction terms. Yet, one may simulate three-body interactions easily by using hopping particles as proposed in [56]. Note that fractal geometries appear in various biological systems, such as broccoli and sea shells in nature. So, we think that physical realization of fractal geometries is possible in principle, and is within the reach of current experimental techniques.

Conventional classical spin liquids are systems that violate the third law of thermodynamics with extensive thermodynamic entropy at zero temperature. Whether one can have topological spin liquids that violate the third law of thermodynamics is an interesting open problem. According to an analytical result obtained in quantum information community [57], for commuting frustration-free quantum spin systems, one cannot have topological spin liquid that violate the third law of thermodynamics. Yet, the problem is open for frustrated systems. Indeed, frustration is the very origin that leads to all the interesting phenomena in both gapped and gapless spin liquids.

Another important open question concerns coding properties of quantum fractal liquids. In [23], we showed that a family of classical fractal liquids asymptotically saturates a theoretical limit on the classical information storage capacity of discrete spin systems. Yet, a quantum code which saturate the quantum information storage capacity has not been found. A mathematical method of finding lower bounds on code distances of quantum fractal liquids is presented in appendix A.

Acknowledgment

I thank Peter Shor, John Preskill, Xiao-Gang Wen and Shintaro Takayoshi for helpful discussion and comments. I acknowledge fruitful discussion with Oliver Buerschaper on RG transformations. At MIT, I was supported by DOE Grant No. DE-FG02-05ER41360. At Caltech, I am supported by David and Ellen Lee Postdoctoral fellowship. I acknowledge funding provided by the Institute for Quantum Information and Matter, an NSF Physics Frontiers Center with support of the Gordon and Betty Moore Foundation (Grants No. PHY-0803371 and PHY-1125565).

Appendix A: Tool-box for translation symmetric stabilizer Hamiltonian

Translation symmetric stabilizer Hamiltonian is a particularly interesting class of exactly solvable quantum spin systems which may possess rich varieties of quantum phases. Systematic studies of translation symmetric stabilizer Hamiltonians were initiated in [34, 43] where topological phases of two-dimensional stabilizer Hamiltonians are com-

pletely classified and are shown to be equivalent to copies of the Toric code. It has been also pointed out that translation symmetric stabilizer Hamiltonians can be studied via commutative algebra [25].

In this appendix, we develop theoretical tools for analyzing physical properties of translation symmetric stabilizer Hamiltonians. For simplicity of discussion, we concentrate mainly on Hamiltonians with the following interaction terms:

$$Z\left(\begin{smallmatrix}\alpha\\ \beta\end{smallmatrix}\right), \quad X\left(\begin{smallmatrix}-\bar{\beta}\\\bar{\alpha}\end{smallmatrix}\right) \quad (\text{A1})$$

where α and β are arbitrary polynomials over $\mathbb{F}_p$. We refer to the model as *canonical model*.

Logical qubits: Consider a polynomial R over $\mathbb{F}_p$ which satisfies

$$R\alpha = R\beta = 0. \quad (\text{A2})$$

The number of logical qubits corresponds to the number of solutions to the above equation since R generates an identity operator: $Z(R\alpha, R\beta)^T = I$. We call such polynomials R the *identity generators* and call Eq. (A2) the *self-consistency equation*. The total number of logical qubits is

$$k = 2 \dim R \quad (\text{A3})$$

by counting contributions from Z -type and X -type stabilizer generators. Note there are p^k ground states in total. For instance, with $\alpha = 1 + x^3$ and $\beta = 1 + x^2$ over $\mathbb{F}_2$, one needs to find solutions to $R(1 + x^3) = R(1 + x) = 0$. Note $1 + x^3 = (1 + x)(1 + x + x^2)$. For $L = 2^m$, one finds that $(1 + x + x^2)^L = 1$, and $(1 + x + x^2)$ has an inverse under periodic boundary conditions. Then the self-consistency equation is further reduced to $R(1 + x) = 0$. Note that $(1 + x)$ is not reversible since $(1 + x)^L = 0$. The only possible non-trivial solution to the self-consistency equation is $R = (1 + x)^{L-1} = 1 + x + x^2 + \dots + x^{L-1}$, and one has $\dim R = 1$ and $k = 2$.

The irreversibility of α and β is important in finding identity generators R . Let us assume that α^{-1} exists. Then the self-consistency equation for R is given by $R\alpha = 0$ which has $R = 0$ only as a solution. So, in order for $k \neq 0$, α and β need to be irreversible. It turns out that irreversibility of α and β is a necessary and sufficient condition for $k \neq 0$ when $L = p^m$. Recall that, for a polynomial $f = \sum_i c_i x^i$, one has $f^p = \sum_i c_i x^{p^i}$. So, for $L = p^m$, one has

$$f^L = \sum_j c_j x^{Lj} = \sum_j c_j = f(x = 1). \quad (\text{A4})$$

One may notice the following:

$$f \text{ is irreversible} \quad \Leftrightarrow \quad f(x = 1) = 0. \quad (\text{A5})$$

Similarly, for higher-dimensional systems, one has

$$f \text{ is irreversible} \quad \Leftrightarrow \quad f(x = 1, y = 1, z = 1, \dots) = 0. \quad (\text{A6})$$

Then, $R = \sum_{i,j,\ell} x^i y^j z^\ell \dots$ is an identity generator since $R\alpha = R\beta = 0$ for irreversible α and β . Therefore, one has the following criteria for logical qubits:

$$\text{A model has a logical qubit } (k \neq 0). \quad \Leftrightarrow \quad \alpha \text{ and } \beta \text{ are irreversible.}$$

Topological order: A necessary and sufficient condition for a canonical model to have topological order can be concisely described by algebraic relations between α and β . Without loss of generality, one may assume that α and β have terms with positive finite powers of $x, y, z, \dots$ only. (Otherwise, we shift the lattice positions). We also assume $k \neq 0$. For now, let us *lift periodic boundary conditions* and consider α and β on a semi-infinite lattice $(x, y, z, \dots = 0, \dots, \infty)$. Let us assume that α and β have a common factor γ :

$$\alpha = \gamma\alpha_0, \quad \beta = \gamma\beta_0$$

on a semi-infinite lattice. Here, we consider the largest common factor γ between α and β . Note that α_0 , β_0 and γ are finite polynomials with local supports since α and β are finite polynomials due to the locality of interaction terms.

Now, let us return to a system under periodic boundary conditions ($x^L = 1$). Then, one has the following criteria for topological order (assuming $k \neq 0$):

$$\text{A canonical model has topological order.} \quad \Leftrightarrow \quad \text{The largest common factor } \gamma \text{ of } \alpha \text{ and } \beta \text{ is reversible.}$$

We emphasize that the reversibility of a common factor γ is defined under periodic boundary conditions while a common factor γ itself is derived on a semi-infinite lattice.

It is easy to prove “ $\Rightarrow$ ” of the above claim. When α and β have an irreversible common factor γ , one can show that the following operator is a logical operator:

$$\ell^{(Z)} = Z \begin{pmatrix} \alpha_0 \\ \beta_0 \end{pmatrix}$$

which leads to $d \sim O(1)$. To prove “ $\Leftarrow$ ” of the claim, we assume that a canonical model does not have topological order and there exists a finite logical operator $\ell^{(Z)} = Z(a, b)^T$ with finite polynomials a and b with $a\beta - b\alpha = 0$. Suppose that γ is reversible. Then $Z(\alpha_0, \beta_0)^T$ is a stabilizer, and $a\beta_0 - b\alpha_0 = 0$. As α_0 and β_0 do not have common factor, one must have $a = \delta\alpha_0$ and $b = \delta\beta_0$ for some finite polynomial δ . Then $\ell^{(Z)}$ is a stabilizer, which leads to a contradiction.

Local unitary transformation: Here, we discuss how a canonical model is transformed into the other via a sequence of two-qubit control gates. We then show that the Cubic code is unitarily equivalent to a certain second-order quantum fractal liquid.

Two-qubit gates can be characterized by a two-qubit Pauli operator $V = V_1 \otimes V_2$. Consider an arbitrary two-qubit Pauli operator $U = U_1 \otimes U_2$. A two-qubit gate generated by V transforms U as follows:

$$U_1 \otimes U_2 \rightarrow U_1 V_1^{c_2} \otimes U_2 V_2^{c_1} \quad (\text{A7})$$

where

$$U_1 V_1 = (-1)^{c_1} V_1 U_1, \quad U_2 V_2 = (-1)^{c_2} V_2 U_2.$$

For instance, with $V = X_1 \otimes X_2$, one has the following transformations:

$$\left\{ \begin{matrix} Z_1, & Z_2 \\ X_1, & X_2 \end{matrix} \right\} \rightarrow \left\{ \begin{matrix} Z_1 X_2, & X_1 Z_2 \\ X_1, & X_2 \end{matrix} \right\}.$$

These two-qubit gates may be viewed as generalizations of control- Z operation. One may see that transformations in Eq. (A7) preserve commutation relations by direct calculations, and thus are indeed unitary transformations.

Let us apply these two-qubit gates to a canonical model with $Z(\alpha, \beta)^T$ and $X(\bar{\beta}, \bar{\alpha})^T$. We think of applying a sequence of two-qubit gates, characterized by $X_A \otimes Z_B$, on neighboring sites in the $\hat{x}$ direction. Then, one has the following transformations:

$$Z \begin{pmatrix} \alpha \\ \beta \end{pmatrix} \rightarrow Z \begin{pmatrix} \alpha \\ \beta + \alpha x \end{pmatrix}, \quad X \begin{pmatrix} \bar{\beta} \\ \bar{\alpha} \end{pmatrix} \rightarrow X \begin{pmatrix} \bar{\beta} + \bar{\alpha} \bar{x} \\ \bar{\alpha} \end{pmatrix} \quad (\text{A8})$$

with $\alpha \rightarrow \alpha$ and $\beta \rightarrow \beta + \alpha x$. Note that these two-qubit gates can be applied simultaneously as they commute with each other. By generalizing this transformation, the following transformations are allowed:

$$\alpha \rightarrow \alpha, \quad \beta \rightarrow \beta + x^i y^j z^\ell \alpha \quad (\text{A9})$$

where i, j, ℓ are some finite integers.

For the Cubic code ($\alpha = 1 + x + y + z$, $\beta = 1 + xy + yz + zx$), we apply two-qubit gates $(\alpha, \beta) \rightarrow (\alpha, \beta + \alpha x)$, modular

transformations $(x, y, z) \rightarrow (x, yz^{-1}, z)$, shifting of lattice sites in $\hat{z}$ directions and two-qubit gates $(\alpha, \beta) \rightarrow (\alpha + \beta, \beta)$:

$$\begin{aligned} \begin{pmatrix} 1+x+y+z \\ 1+xy+yz+zx \end{pmatrix} &\rightarrow \begin{pmatrix} 1+x+y+z \\ 1+x+x^2+yz \end{pmatrix} \rightarrow \begin{pmatrix} 1+x+yz^{-1}+z \\ 1+x+x^2+y \end{pmatrix} \\ &\rightarrow \begin{pmatrix} y+(1+x)z+z^2 \\ 1+x+x^2+y \end{pmatrix} \rightarrow \begin{pmatrix} (1+x+x^2)+(1+x)z+z^2 \\ 1+x+x^2+y \end{pmatrix}. \end{aligned}$$

This is a second-order quantum fractal liquid.

Open boundary condition: One may also consider open boundary conditions by defining stabilizer generators nicely so that they commute with each other at boundaries. Without loss of generality, one can assume that α and β have the following forms:

$$\alpha = 1 + c_1x + c_2x^2 + \cdots, \quad \beta = 1 + d_1x + d_2x^2 + \cdots.$$

For Z -type stabilizer generators for $j = 0, \dots, L-1$, we impose the following constraints:

$$x^j = 0 \quad \text{for } j \geq L.$$

In other words, Z -type stabilizers are truncated at the boundary on the right hand side (Fig. 11). X -type stabilizer generators are defined by taking the duals of Z -type stabilizer generators. They are truncated at the boundary on the left hand side. One may see that stabilizer generators commute with each other at the boundaries. A similar construction works for higher-dimensional systems too. We note that this construction of canonical models with open boundary conditions reproduces a model of the Toric code with boundaries, originally proposed by Bravyi and Kitaev where boundaries are classified into two types; rough and smooth boundaries [58].




FIG. 11: A construction of stabilizer Hamiltonians with open boundary conditions. An example with $\alpha = 1 + x + x^2$ and $\beta = 1 + x$ is shown. For Z -type stabilizers, Pauli operators are truncated at the right edge, and for X -type stabilizers, Pauli operators are truncated at the left edge.

Higher-dimensional model: The D -dimensional Toric code with m -dimensional and $(D-m)$ -dimensional logical operators can be described by a model with ${}_DC_m$ qubits per site with variables $x_1, \dots, x_D$. We label sublattices by a set of m integers $(A_1, A_2, \dots, A_m)$ for $A_j \in \mathbb{Z}_D$. For Z -type stabilizer generators, we choose an arbitrary set of

$m + 1$ integers A^* from $\mathbb{Z}_D$. If $(A_1, A_2, \dots, A_m) \cup A_0 = A^*$, we put $1 + x_{A_0}$ for a sublattice $(A_1, A_2, \dots, A_m)$. As for X -type stabilizer generators, we consider a dual of $(A_1, A_2, \dots, A_m)$ (a maximal set of integers which do not overlap with $(A_1, A_2, \dots, A_m)$) and choose an arbitrary set of $D - m + 1$ integers.

For instance, with $D = 3$ and $m = 1$, three sublattices are labeled by $(1), (2), (3)$. One may consider $A^* = (1, 2), (2, 3), (3, 1)$ and obtain Z -type stabilizers:

$$\begin{pmatrix} 1 + x_2 \\ 1 + x_1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 + x_3 \\ 1 + x_2 \end{pmatrix}, \begin{pmatrix} 1 + x_3 \\ 0 \\ 1 + x_1 \end{pmatrix}.$$

As for X -type stabilizers, sublattices are labelled by duals $(2, 3), (3, 1), (1, 2)$. Taking $A^* = (1, 2, 3)$, one has

$$\begin{pmatrix} 1 + \bar{x}_1 \\ 1 + \bar{x}_2 \\ 1 + \bar{x}_3 \end{pmatrix}.$$

With $D = 4$ and $m = 2$, sublattices are labelled by $(1, 2), (1, 3), (1, 4), (2, 3), (2, 4), (3, 4)$, and for $A^* = (1, 2, 3), (1, 2, 4), (1, 3, 4), (2, 3, 4)$, Z -type stabilizers are

$$\begin{pmatrix} 1 + x_3 \\ 1 + x_2 \\ 0 \\ 1 + x_1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 + x_4 \\ 0 \\ 1 + x_2 \\ 0 \\ 1 + x_1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 + x_4 \\ 1 + x_3 \\ 0 \\ 0 \\ 1 + x_1 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 + x_4 \\ 1 + x_3 \\ 1 + x_2 \end{pmatrix}.$$

As for X -type stabilizers, sublattices are labelled by duals $(3, 4), (2, 4), (2, 3), (1, 4), (1, 3), (1, 2)$, and one has

$$\begin{pmatrix} 0 \\ 0 \\ 1 + \bar{x}_1 \\ 0 \\ 1 + \bar{x}_2 \\ 1 + \bar{x}_3 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 + \bar{x}_1 \\ 0 \\ 1 + \bar{x}_2 \\ 0 \\ 1 + \bar{x}_4 \end{pmatrix}, \begin{pmatrix} 1 + \bar{x}_1 \\ 0 \\ 0 \\ 1 + \bar{x}_3 \\ 1 + \bar{x}_4 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 + \bar{x}_2 \\ 1 + \bar{x}_3 \\ 1 + \bar{x}_4 \\ 0 \\ 0 \\ 0 \end{pmatrix}.$$

To construct fractal versions of these models in $(D + 1)$ -dimensional systems, we add an extra variable x_0 , and replace

$$x_j \rightarrow f_j(x_0)x_j$$

where $f_j(x_0)$ is a polynomial with x_0 only.

Lower bound on code distance: Below, we reduce a problem of finding lower bounds on code distances of quantum fractal liquids to a certain problem of decomposing polynomials $f(x)$ and $g(x)$. Consider a quantum fractal liquid for $L = p^m$ with reversible $f(x)$ and $g(x)$ with $f(x = 1) = 1$ and $g(x = 1) = 1$. Then, one has the following holds.

- Consider all the tensors $C_{ij}^{(\ell)}$ satisfying the following equation:

$$\gamma(x)g(x)^\ell = \sum_{ij} x^i f^j(x) C_{ij}^{(\ell)}, \quad \text{for all } \ell. \quad (\text{A10})$$

and consider all the tensors $D_{ij}^{(\ell)}$ satisfying the following equation:

$$\gamma(x)f(x)^\ell = \sum_{ij} x^i g(x)^j D_{ij}^{(\ell)}, \quad \text{for all } \ell. \quad (\text{A11})$$

We denote minimal weights of tensors $C_{ij}^{(\ell)}$ and $D_{ij}^{(\ell)}$ as $W_m^{(C)}$ and $W_m^{(D)}$ for $\gamma(x) \neq 0$. We denote the minimum of $W_m^{(C)}$ and $W_m^{(D)}$ as W_m . Then the code distance d_m is lower bounded by

$$d_m \geq W_m. \quad (\text{A12})$$

Below, we present the proof of the above claim. Consider a quantum fractal liquid with $L \times L \times L$ sites with $L = p^m$, and denote the ℓ -th layer of $L \times L \times 1$ sites as Q_ℓ for $\ell = 0, \dots, L-1$. We think of removing sites in a subset $E = E_0 \cup E_1 \cup \dots \cup E_{L-1}$ with E being a set of site indices and $E_\ell \subseteq Q_\ell$. We ask if $\ell^{(Z)}$ -type logical operators can be supported inside the remaining sites $\bar{E}$ or not. $\ell^{(Z)}$ -type logical operators can be represented as:

$$\ell^{(Z)}(\gamma) = Z \begin{pmatrix} 0 \\ \gamma(x)\mathbf{f}(x, y) \end{pmatrix}, \quad \gamma(x) = c_0 + c_1x + \dots + c_{L-1}x^{L-1}$$

where $c_j = 0, \dots, p-1$. Let us denote coefficients of $x^i y^j$ in $\gamma(x)\mathbf{f}(x, y)$ as $S_{ij0}(\gamma)$:

$$\gamma(x)\mathbf{f}(x, y) = \sum_{ij} S_{ij0}(\gamma) x^i y^j.$$

Observe that a logical operator $\ell^{(Z)}(\gamma)$ can be supported inside $Q_0/(i, j, 0)$ if and only if a polynomial $\gamma(x)\mathbf{f}(x, y)$ has a zero coefficient for $x^i y^j$. Then $\ell^{(Z)}(\gamma)$ is supported inside Q_0/E_0 if and only if $S_{ij0}(\gamma) = 0$ for all $(i, j, 0) \in E_0$.

It is convenient to introduce polynomials R_{ij0} which satisfy the following relations:

$$R_{ij0} = r_0^{(ij0)} + r_1^{(ij0)}x + \dots + r_{L-1}^{(ij0)}x^{L-1}, \quad \text{where} \quad S_{ij0}(\gamma) = \sum_{a=0}^{L-1} r_a^{(ij0)} c_a. \quad (\text{A13})$$

For instance, one has $R_{000} = 1$, $R_{100} = x$ and $R_{200} = x^2$ since $S_{000}(\gamma) = c_0$, $S_{100}(\gamma) = c_1$ and $S_{200}(\gamma) = c_2$. With some speculation, one can prove that

$$R_{ij0} = x^i \bar{f}(x)^j \quad (\text{A14})$$

where $\bar{f}(x)$ is the dual of $f(x)$. R_{ij0} may be viewed as a polynomial representation of coefficients $S_{ij0}(\gamma)$. In order for $\ell^{(Z)}(\gamma)$ to be supported inside Q_0/E_0 , $\gamma(x)$ needs to satisfy a set of constraints that are characterized by polynomials inside $S(Q_0) \equiv \langle R_{ij0} : (i, j, 0) \in E_0 \rangle$.

Next, let us find logical operators which have equivalent representations inside Q_1/E_1 . Note the following equivalence relation:

$$\ell^{(Z)}(\gamma) = Z \begin{pmatrix} 0 \\ \gamma(x)\mathbf{f}(x, y) \end{pmatrix} \sim Z \begin{pmatrix} 0 \\ \gamma(x)g(x)\mathbf{f}(x, y)z \end{pmatrix}.$$

So, $\ell^{(Z)}(\gamma)$ can be supported inside Q_1/E_1 if and only if $S_{ij1}(\gamma) = 0$ for all $(i, j, 1) \in E_1$ where S_{ij1} corresponds to coefficients of $x^i y^j z$ in $z\gamma(x)\mathbf{f}(x, y)g(x)$. These constraints can be represented as

$$R_{ij1} = r_0^{(ij1)} + r_1^{(ij1)}x + \dots + r_{L-1}^{(ij1)}x^{L-1} = x^i \bar{f}(x)^j \bar{g}(x), \quad S_{ij1}(\gamma) = \sum_{a=0}^{L-1} r_a^{(ij1)} c_a. \quad (\text{A15})$$

In general, for polynomials

$$R_{ij\ell} = \sum_a r_a^{(ij\ell)} x^a = x^i \bar{f}(x)^j \bar{g}(x)^\ell, \quad (\text{A16})$$

one has

$$S_{ij\ell}(\gamma) = \sum_a r_a^{(ij\ell)} c_a$$

where

$$\gamma(x)\mathbf{f}(x,y)\mathbf{g}(x,z) = \sum_{ij\ell} S_{ij\ell} x^i y^j z^\ell.$$

Consider the following spaces of constraints on $\gamma(x)$ for Q_ℓ/E_ℓ :

$$S(Q_\ell) = \langle R_{ij\ell} : (i,j,\ell) \in E_\ell \rangle. \quad (\text{A17})$$

Then, all the logical operators can be supported inside $\bar{E}$ if

$$S(Q_0) \cap S(Q_1) \cap \cdots \cap S(Q_{L-1}) = \emptyset.$$

In other words, if there is no common constraint for $\gamma(x)$ to satisfy among $S(Q_\ell)$, logical operators $\ell^{(Z)}(\gamma)$ with arbitrary $\gamma(x)$ can be supported inside $\bar{E}$.

To analyze properties of E which satisfy Eq. (A17), let us consider the case where $S(Q_\ell)$ have some common element for all ℓ . Then, a tensor $C_{ij}^{(\ell)}$ satisfying the following equation must exist:

$$\gamma(x) = \sum_{ij} x^i \bar{f}(x)^j \bar{g}(x)^\ell C_{ij}^{(\ell)}, \quad \text{for all } \ell. \quad (\text{A18})$$

Consider all the possible solutions $C_{ij}^{(\ell)}$ to Eq. (A18) and their weights (the number of non-zero entries), and denote the minimal weight as $W_m^{(C)}(\gamma)$. Then $\ell^{(Z)}(\gamma)$ can be supported inside $\bar{E}$ if $|E| < W_m^{(C)}(\gamma)$. Let us further define the minimum of $W_m^{(C)}(\gamma)$ for all $\gamma(x) \neq 0$ as

$$W_m^{(C)} = \min_{\gamma \neq 0} W_m^{(C)}(\gamma).$$

Then all the $\ell^{(Z)}$ -type logical operators can be supported inside $\bar{E}$ if $|E| < W_m^{(C)}$. One can repeat a similar analysis for $r^{(Z)}$ -type logical operators, and ends up with the following equation

$$\gamma(x) = \sum_{ij} x^i \bar{g}(x)^j \bar{f}(x)^\ell D_{ij}^{(\ell)}, \quad \text{for all } \ell. \quad (\text{A19})$$

By considering all the possible solutions to Eq. (A19) for all $\gamma \neq 0$ and denoting their minimal weight as $W_m^{(D)}$, one notices that all the $r^{(Z)}$ -type logical operators can be supported inside $\bar{E}$ if $|E| < W_m^{(D)}$.

We denote the minimum of $W_m^{(C)}$ and $W_m^{(D)}$ as W_m . Then, all the logical operators can be supported inside $\bar{E}$ if $|E| < W_m$. Then there is no logical operator whose weight is smaller than W_m [45], and one has

$$d_m \geq W_m \quad (\text{A20})$$

where d_m represents the code distance for $L = p^m$. This leads to the proof of Eq. (A12).

-
- [1] A. Y. Kitaev, Ann. Phys. **303**, 2 (2003).
 - [2] X. G. Wen and Q. Niu, Phys. Rev. B **41**, 9377 (1990).
 - [3] D. C. Tsui, H. L. Stormer, and A. C. Gossard, Phys. Rev. Lett. **48**, 1559 (1982).
 - [4] R. B. Laughlin, Phys. Rev. Lett. **50**, 1395 (1983).
 - [5] X.-G. Wen, *Quantum Field Theory of Many-Body Systems* (Oxford University Press, Oxford, 2004).
 - [6] X. G. Wen, F. Wilczek, and A. Zee, Phys. Rev. B **39**, 11413 (1989), URL <http://link.aps.org/doi/10.1103/PhysRevB.39.11413>.
 - [7] N. Read and S. Sachdev, Phys. Rev. Lett. **66**, 1773 (1991).
 - [8] X. G. Wen, Phys. Rev. B **44**, 2664 (1991).
 - [9] P. W. Anderson, Science **235**, 1196 (1987).

- [10] I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki, Phys. Rev. Lett. **59**, 799 (1987).
- [11] T. Senthil and M. P. A. Fisher, Phys. Rev. B **62**, 7850 (2000).
- [12] R. Moessner and S. L. Sondhi, Phys. Rev. Lett. **86**, 1881 (2001).
- [13] A. Kitaev, Ann. Phys. **321**, 2 (2006).
- [14] E. Witten, Commun. Math. Phys. **117**, 353 (1988).
- [15] M. Atiyah, Publ. Math. I.H.E.S. **68**, 175 (1988).
- [16] M. A. Levin and X.-G. Wen, Phys. Rev. B **71**, 045110 (2005).
- [17] X. Chen, Z.-C. Gu, and X.-G. Wen, Phys. Rev. B **82**, 155138 (2010).
- [18] J. Haah, Phys. Rev. A **83**, 042330 (2011).
- [19] S. Bravyi and J. Haah, arXiv:1112.3252.
- [20] I. H. Kim, arXiv:1202.0052.
- [21] C. Castelnovo and C. Chamon, Philosophical Magazine **92**, 304 (2012).
- [22] M. E. J. Newman and C. Moore, Phys. Rev. E **60**, 5068 (1999).
- [23] B. Yoshida, arXiv:1111.3275.
- [24] C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, Rev. Mod. Phys. **80**, 1083 (2008).
- [25] J. Haah, arXiv:1204.1063.
- [26] V. Efimov, Physics Letters B **33**, 563 (1970).
- [27] K. G. Wilson, Phys. Rev. D **3**, 1818 (1971).
- [28] P. F. Bedaque, H. W. Hammer, and U. van Kolck, Phys. Rev. Lett. **82**, 463 (1999).
- [29] A. B. Zmolodchikov, JETP Lett. **43**, 730 (1986).
- [30] J. Polchinski, Nucl. Phys. B **303**, 226 (1988).
- [31] J.-F. Fortin, B. Grinstein, and A. Stergiou, Physics Letters B **704**, 74 (2011).
- [32] X.-G. Wen, Phys. Rev. Lett. **90**, 016803 (2003).
- [33] S. Bravyi, M. Hastings, and S. Michalakis, J. Math. Phys. **51**, 093512 (2010).
- [34] B. Yoshida, Ann. Phys. **326**, 2566 (2011).
- [35] S. Wolfram, *A new kind of science* (Wolfram Media Inc. Champaign., 2002).
- [36] A. P. Ramirez, Annu. Rev. Mater. Sci. **24**, 453 (1994).
- [37] M. J. Harris, S. T. Bramwell, D. F. McMorrow, T. Zeiske, and K. W. Godfrey, Phys. Rev. Lett. **79**, 2554 (1997).
- [38] R. Moessner and J. T. Chalker, Phys. Rev. B **58**, 12049 (1998).
- [39] C. L. Henley, Phys. Rev. B **71**, 014424 (2005).
- [40] C. Castelnovo, R. Moessner, and S. L. Sondhi, Nature **451**, 42 (2008).
- [41] S. Bravyi, M. B. Hastings, and F. Verstraete, Phys. Rev. Lett. **97**, 050401 (2006).
- [42] M. B. Hastings and X.-G. Wen, Phys. Rev. B **72**, 045141 (2005).
- [43] B. Yoshida, Ann. Phys. **326**, 15 (2011).
- [44] S. J. Willson, Physica D **24**, 190 (1987).
- [45] B. Yoshida and I. L. Chuang, Phys. Rev. A **81**, 052302 (2010).
- [46] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).
- [47] S. Bravyi and J. Haah, Phys. Rev. Lett. **107**, 150504 (2011).
- [48] A. Hama, C. Castelnovo, and C. Chamon, Phys. Rev. B **79**, 245122 (2009).
- [49] K. Michnicki, arXiv:1208.3496.
- [50] M. Levin and X.-G. Wen, Phys. Rev. Lett. **96**, 110405 (2006).
- [51] A. Kitaev and J. Preskill, Phys. Rev. Lett. **96**, 110404 (2006).
- [52] G. Vidal, Phys. Rev. Lett. **99**, 220405 (2007).
- [53] F. Verstraete and J. I. Cirac, cond-mat/0407066 (2004).
- [54] Z.-C. Gu, M. Levin, and X.-G. Wen, Phys. Rev. B **78**, 205116 (2008).
- [55] G. Evenbly and G. Vidal, Phys. Rev. Lett. **102**, 180406 (2009).
- [56] S. A. Ocko and B. Yoshida, Phys. Rev. Lett. **107**, 250502 (2011).
- [57] S. Bravyi, D. Poulin, and B. Terhal, Phys. Rev. Lett. **104**, 050503 (2010).
- [58] S. B. Bravyi and A. Y. Kitaev, quant-ph/9811052 (1998).