

Existence of strong solutions to the steady Navier-Stokes equations for a compressible heat-conductive fluid with large forces

Changsheng Dou^{1,*}, Fei Jiang^{2,3}, Song Jiang², Yongfu Yang⁴

¹ School of Statistics, Capital University of Economics and Business,
Beijing 100070, P.R. China

² Institute of Applied Physics and Computational Mathematics,
P.O. Box 8009, Beijing 100088, P.R. China

³ College of Mathematics and Computer Science, Fuzhou University,
Fuzhou 350108, China

⁴Department of Mathematics, College of Sciences, Hohai University,
Nanjing 210098, Jiangsu Province, P.R. China

Abstract

We prove that there exists a strong solution to the Dirichlet boundary value problem for the steady Navier-Stokes equations of a compressible heat-conductive fluid with large external forces in a bounded domain $\Omega \subset \mathbb{R}^d$ ($d = 2, 3$), provided that the Mach number is appropriately small. At the same time, the low Mach number limit is rigorously verified. The basic idea in the proof is to split the equations into two parts, one of which is similar to the steady incompressible Navier-Stokes equations with large forces, while another part corresponds to the steady compressible heat-conductive Navier-Stokes equations with small forces. The existence is then established by dealing with these two parts separately, establishing uniform in the Mach number a priori estimates and exploiting the known results on the steady incompressible Navier-Stokes equations.

MSC: 76N99; 35M33; 35Q30

Keywords. Steady compressible heat-conductive Navier-Stokes equations, large external forces, existence of strong solutions, low Mach number limit, Dirichlet boundary condition.

1 Introduction

This paper is mainly concerned with the existence of strong solutions to the steady Navier-Stokes equations of a compressible heat-conductive fluid in a bounded domain $\Omega \subset \mathbb{R}^d$ ($d = 2, 3$) with large external forces:

$$\begin{cases} \operatorname{div}(\varrho \mathbf{u}) = 0, \\ \varrho \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p = \operatorname{div} \mathbb{S}(\nabla \mathbf{u}) + \varrho \mathbf{f} + \mathbf{g}, \\ c_v \varrho \mathbf{u} \cdot \nabla \Theta + p \operatorname{div} \mathbf{u} = \kappa \Delta \Theta + \Psi. \end{cases} \quad (1.1)$$

Here ϱ denotes the density, $\mathbf{u} \in \mathbb{R}^d$ the velocity, Θ the temperature, $p = R\varrho\Theta$ the pressure with $R > 0$ being the gas constant, $c_v > 0$ is the heat capacity at constant volume; \mathbf{f} is the density

*Corresponding author. E-mails: douchangsheng@163.com douchangsheng@cueb.edu.cn (C. Dou), jiangfei0591@163.com (F. Jiang), jiang@iapcm.ac.cn (S. Jiang), fudanyoung@gmail.com (Y. Yang)

of external body force and \mathbf{g} is a given external force. The stress tensor \mathbb{S} and the dissipation function Ψ are defined by

$$\mathbb{S} = 2\mu D(\mathbf{u}) + \lambda \operatorname{div} \mathbf{u} \mathbb{I}, \quad \Psi = 2\mu D(\mathbf{u}) : D(\mathbf{u}) + \lambda (\operatorname{div} \mathbf{u})^2 \geq 0,$$

$D(\mathbf{u}) = (\nabla \mathbf{u} + \nabla \mathbf{u}^t)/2$ is the deformation tensor. The viscosity coefficients μ, λ satisfy $2\mu + d\lambda \geq 0$ and $\mu > 0, \kappa > 0$ is the heat conductivity coefficient. Moreover, the total mass is prescribed:

$$\int_{\Omega} \varrho dx = M > 0.$$

We impose that the velocity \mathbf{u} satisfies no-slip boundary condition and the temperature Θ keeps constant on the boundary of Ω , i.e.,

$$\mathbf{u} = 0, \quad \Theta = \vartheta_0 \quad \text{on } \partial\Omega. \quad (1.2)$$

In the last decades, the steady compressible heat-conductive Navier-Stokes equations have been studied by many mathematicians and there are a lot of results on the existence in the literature, here we recall some of them for both small and large external forces which are related to our study in this paper, and we refer to the monograph [27] for more details. When external forces are sufficiently small, Matsumura and Nishida in 1982/83 proved the existence of a solution with potential forces near a rest state [20, 21], while Valli and Zajackowski [34, 36] used the existence of global non-stationary solutions to get the existence of stationary solutions. Later, Valli [35] showed the existence of stationary solutions in the case of general forces by using an idea of Padula [28] to decompose the equations into two parts that are governed by the Stokes equations and a transport equation, respectively. Beirão da Veiga [2] obtained more general existence results in the L^p -setting by decomposing the equations into three parts that are governed by the Stokes equations, a transport equation and the Laplace equation, respectively. Another decomposition was studied in the paper [26]. In 1989, Farwig [7] showed the existence of solutions to the steady compressible heat-conductive Navier-Stokes equations for small forces with slip boundary condition.

When external forces are of arbitrary size, the existence of strong solutions was proved in [25, 22] for the case of potential forces. When the equations of state and the viscosity coefficients satisfy certain (growth) conditions, Novotný and Pokorný [23, 24] showed that weak or strong solutions to the steady compressible heat-conductive Navier-Stokes equations exist. Unfortunately, their results exclude the case of ideal polytropic gases, for which the existence of strong solutions, to our best knowledge, still remains open.

The aim of the present paper is to establish the existence of strong solutions to the steady compressible heat-conductive Navier-Stokes system (1.1) without any smallness assumption on the external forces \mathbf{f} and \mathbf{g} , when the Mach number is small.

We mention that in the isentropic flow case, the existence of weak solutions or strong solutions for large external forces has been extensively investigated. Lions [19] first proved the existence of weak solutions under the assumption that the specific heat ratio $\gamma > 1$ in two dimensions and $\gamma > 5/3$ in three dimensions. The restriction on γ actually comes from the integrability of the density ϱ in L^p , and in fact, the higher integrability of ϱ has, the smaller γ can be allowed. In [27] Novotný and Straškraba showed the existence of weak solutions for any $\gamma > 3/2$ if \mathbf{f} is potential and $\mathbf{g} = 0$. By deriving a new weighted estimate of the pressure, Frehse, Goj and Steinhauer [11], Plotnikov and Sokolowski [30] established an improved integrability for the density under the assumption of the L^1 -boundedness of $\varrho \mathbf{u}^2$ which was not shown to hold unfortunately. Plotnikov and Sokolowski [32] proved the existence of renormalized solutions to the Dirichlet boundary value problem for the compressible Navier-Stokes equations for all $\gamma > 4/3$. In 2008, Březina and Novotný [4] was able to prove the existence of weak solution to the spatially periodic problem for any $\gamma > (3 + \sqrt{41})/8$ when \mathbf{f} is potential and $\mathbf{g} = 0$, or for any $\gamma > (1 + \sqrt{13})/3 \approx 1.53$

when $\mathbf{f}, \mathbf{g} \in L^\infty$, without assuming the L^1 -boundedness of $\varrho \mathbf{u}^2$, by combining the L^∞ -estimate of $\Delta^{-1}P$ with the (usual) energy and density bounds. Then, in the framework of [31], Frehse, Steinhauer and Weigant [12, 13] established the existence of weak solutions to the Dirichlet boundary value problem for any $\gamma > 4/3$ in three dimensions and to the spatially periodic or mixed boundary value problem for $\gamma = 1$ (isothermal flow) in two dimensions. Recently, Jiang and Zhou [17] proved the existence of weak solutions to the spatially periodic problem in \mathbb{R}^3 for any $\gamma > 1$ by establishing a new weighted estimate. More recently, the existence for the slip and Dirichlet boundary value problems for $\gamma > 1$ was shown by Jesslé and Novotný [16], and Plotnikov and Weigant [33], respectively. Furthermore, we emphasize that the existence of strong solutions was shown by Choe and Jin [5] when the Mach number is small, by exploiting the known results for the incompressible steady Navier-Stokes equations.

Now, we rewrite (1.1) in the form of the Mach number. After scaling and a straightforward calculation we obtain the following dimensionless form of the steady full compressible Navier-Stokes equations:

$$\begin{cases} \operatorname{div}(\varrho \mathbf{u}) = 0, \\ \varrho \mathbf{u} \cdot \nabla \mathbf{u} + \frac{\nabla p}{\epsilon^2} = \operatorname{div} \mathbb{S}(\nabla \mathbf{u}) + \varrho \mathbf{f} + \mathbf{g}, \\ \varrho \mathbf{u} \cdot \nabla \Theta + p \operatorname{div} \mathbf{u} = \kappa \Delta \Theta + \epsilon^2 \Psi, \end{cases} \quad (1.3)$$

where ϵ is the Mach number.

Since the total mass of the fluid is given, we impose the condition

$$\bar{\varrho} := \frac{1}{|\Omega|} \int_{\Omega} \varrho(x) dx > 0,$$

which can be renormalized to $\bar{\varrho} = 1$ without loss of generality. Similarly, we also assume that $\bar{\Theta} = 1$, $R = c_V = 1$, $\vartheta_0 = 1$.

To show the existence, we take the transformation

$$\varrho = 1 + \epsilon \rho, \quad \Theta = 1 + \epsilon \theta \quad (1.4)$$

to rewrite the system (1.3) in the form:

$$\begin{cases} \operatorname{div} \mathbf{u} + \epsilon \operatorname{div}(\rho \mathbf{u}) = 0, \\ (1 + \epsilon \rho)(\mathbf{u} \cdot \nabla \mathbf{u}) + \frac{(1 + \epsilon \theta) \nabla \rho}{\epsilon} + \frac{(1 + \epsilon \rho) \nabla \theta}{\epsilon} = \operatorname{div} \mathbb{S}(\nabla \mathbf{u}) + (1 + \epsilon \rho) \mathbf{f} + \mathbf{g}, \\ \epsilon(1 + \epsilon \rho) \mathbf{u} \cdot \nabla \theta + \operatorname{div} \mathbf{u} + (\epsilon \rho + \epsilon \theta + \epsilon^2 \rho \theta) \operatorname{div} \mathbf{u} = \epsilon \kappa \Delta \theta + \epsilon^2 \Psi, \end{cases} \quad (1.5)$$

with boundary conditions

$$\mathbf{u} = 0, \quad \theta = 0 \quad \text{on } \partial\Omega. \quad (1.6)$$

The low Mach number limit for the corresponding evolutionary equations was investigated extensively. Here, we only refer to the non-isentropic Navier-Stokes equations, whose analysis is more difficult due to the complexity of the system structure. Hagstrom and Lorenz [15] used a similar transformation to (1.4), the standard energy arguments and decay estimates for heat kernels, and constructed a special symmetrizer for linear hyperbolic-parabolic systems with large hyperbolic part satisfying the interaction condition to get the low Mach number limit of compressible Navier-Stokes equations in \mathbb{R}^n . Bresch et al [3] analyzed the acoustic waves by a method of characteristic expansions and gave a formal asymptotics as $\epsilon \rightarrow 0$ in a periodic domain under the assumption that the viscous heating and thermal diffusion are negligible. Concerning the full compressible Navier-Stokes equations, Alazard [1] studied this singular limit for local H^s solutions ($s > 2 + \frac{n}{2}$) in \mathbb{R}^n for "ill-prepared" initial data by employing the technique of pseudo-differential operators which does not apply to the cases with boundary due to the restriction of the Fourier transform. As an improvement of [3], Feireisl and Novotný [9] considered the

low Mach number limit for the periodic "variational solutions" to the full Navier-Stokes-Fourier equations of certain radiative gases for "ill-prepared" initial data. Related progresses for bounded domains with various boundary conditions can be found in [8, 10, 6]. On the other hand, when the thermal conductivity vanishes, the incompressible limit of the non-isentropic Navier-Stokes equations in bounded domains has been studied recently in [18] where the strong solutions are shown to be bounded uniformly in a local time interval for "well-prepared" initial data, thus implying the limit. However, to our best knowledge, the low Mach number limit for the non-isentropic Navier-Stokes equations governing polytropic gases in bounded domains is not yet proved so far, and the aim of the current paper is thus to show this limit.

Now, we state the main result of this paper.

Theorem 1.1. *Let $\mathbf{f}, \mathbf{g} \in H^2(\Omega)$. Then there is an ϵ_0 depending on $\|(\mathbf{f}, \mathbf{g})\|_{H^2}$ and Ω , such that for any $\epsilon \in (0, \epsilon_0)$, there exists a solution $(\rho^\epsilon, \mathbf{u}^\epsilon, \theta^\epsilon) \in \bar{H}^2 \times (H^3 \cap H_0^1) \times (H^3 \cap H_0^1)$ to the boundary value problem (1.5), (1.6), satisfying*

$$\lim_{\epsilon \rightarrow 0} \inf_{U, P \in \mathbf{L}} \|\mathbf{u}^\epsilon - U\|_3 + \|\rho^\epsilon\|_2 + \|\theta^\epsilon\|_3 + \left\| \frac{\rho^\epsilon + \theta^\epsilon}{\epsilon} - P \right\|_2 = 0,$$

where $(U, P) \in \mathbf{L} := \{(U, P) \in (H^4 \cap H_0^1) \times \bar{H}^3 \mid (U, P) \text{ is a solution of the incompressible steady Navier-Stokes equations (1.7) with external force } \mathbf{f} + \mathbf{g}\}$, i.e.,

$$\begin{cases} U \cdot \nabla U - \mu \Delta U + \nabla P = \mathbf{f} + \mathbf{g}, \\ \operatorname{div} U = 0, \\ U = 0 \text{ on } \partial\Omega, \quad \int_{\Omega} P dx = 0. \end{cases} \quad (1.7)$$

Remark 1.1. *If considering the existence of spatially periodic solutions to (1.5) in a periodic domain, we can also obtain a existence result similar to Theorem 1.1.*

Remark 1.2. *In general, we could not get the uniqueness of strong solutions to the boundary value problem (1.1) due to lack of the uniqueness of strong solutions to the corresponding incompressible Navier-Stokes equations (1.7). As for the results of the incompressible Navier-Stokes equations, one can refer to Galdi's book [14, Chapter IX].*

The system (1.5) is complicated mixed-type nonlinear equations containing such structures as elliptic and hyperbolic systems, for which the usual approach of the fixed point arguments used to prove the existence of classical solutions requires the smallness of data. To show Theorem 1.1 we split the system (1.5) into two parts, one of which is similar to the steady incompressible Navier-Stokes equations with large force $\mathbf{f} + \mathbf{g}$, while another part corresponds to the steady compressible heat-conductive Navier-Stokes equations with small force $\epsilon \mathbf{f}$, provided the Mach number ϵ is small. Then, as noted in [5], we modify and elaborately combine the arguments in [14] where the existence of strong solutions to the incompressible Navier-Stokes equations for large forces was presented, and in [7] where strong solutions of the compressible viscous heat-conductive equations with small forces were dealt with, to establish Theorem 1.1.

Compared with the isentropic case studied in [5], due to presence of the energy equation, the main difficulties here lie in obtaining the existence of weak solutions to the linearized system (2.10), dealing with the coupling terms between the velocity, density and temperature, and deriving the uniform-in- ϵ estimates in a bounded domain, for example, how to control the energy norm $\|\mathbf{u} - U\|_3 + \|\eta\|_2 + \|\theta\|_3$ uniformly in ϵ under the no-slip boundary condition. To circumvent such difficulties, we take the transform of $\varrho = 1 + \epsilon\rho$, $\Theta = 1 + \epsilon\theta$ for the system (1.3) (see also [15]), instead of the transform $(\varrho = 1 + \epsilon^2\rho, \Theta = 1 + \epsilon^2\theta)$ used in [5], carefully construct an approximate linear problem by using a cut-off function and employ a mollifier technique to get the existence and uniqueness of solutions to (2.10), and utilize the lower order terms to control

the higher order terms. We should remark that in [5] the linearized problems are decoupled, while, to obtain the uniform a priori estimates in this paper, the linearized problems are strongly coupled that gives rise to more difficulties than the isentropic case in [5].

Let us briefly explain the main steps of our proof. First, we construct the approximate linear problem, derive the uniform estimates and employ a compactness argument to get the existence of a weak solution to the system (2.10) in Section 2.2. Second, we exploit the property of the momentum equations and the regularity of the Stokes problem to establish the estimates of $|\eta + \theta|$, which, combined with an estimate for θ , implies the boundedness for η . Due to presence of boundary here, some difficulties involved with controlling the boundary terms arise. To overcome such difficulties, the crucial step is to get a H^2 -bound of $\operatorname{div} \mathbf{u}$ near the boundary, for which we shall adopt the local isothermal coordinates used in [34, 36]. This strategy has also been used in [18] to study the low Mach number limit of the compressible Navier-Stokes equations with non-slip boundary condition. Then, summing up all the estimates for (\mathbf{v}, θ) and η , we can establish the desired a priori uniform in ϵ estimates in view of the smallness of ϵ (see Section 3). Finally, we apply the Tikhonov fixed point theorem to obtain the existence of a strong solution. Moreover, with the help of the uniform a priori estimates, one can take the limit to show the incompressible limit. We point out here that due to the splitting, we have to impose that the energy equation should not possess an external heat source.

The rest of this paper is organized as follows. In the next section, we prove the existence of weak solutions and regularity to the linearized incompressible and compressible problems. Section 3 is devoted to establishing the existence for the nonlinear problem. Finally, the incompressible limit of solutions to the steady compressible heat-conductive Navier-Stokes equations is presented in Section 4.

Notations: We denote by L^2 the Lebesgue space $L^2(\Omega)$ with norm $\|\cdot\|_0$, by H^m the Sobolev spaces $H^m(\Omega)$ with norm $\|\cdot\|_m$. Define the spaces

$$\bar{H}^m = \left\{ \rho \in H^m \mid \int_{\Omega} \rho(x) dx = 0 \right\}, \quad H_{0,\sigma}^1 = \left\{ \mathbf{u} \in H_0^1 \mid \operatorname{div} \mathbf{u} = 0 \right\}, \quad H_{0,\sigma}^m := H^m \cap H_{0,\sigma}^1.$$

We denote by H^{-1} the dual space of H_0^1 with the dual product $\langle \cdot, \cdot \rangle$ and the norm $\|\cdot\|_{-1} = \sup_{\|h\|_1=1} |\langle \cdot, h \rangle|$. We shall use the abbreviation:

$$\int \cdot dx := \int_{\Omega} \cdot dx.$$

2 Existence of solutions to the linearized problem

We first split the system (1.5) into two parts, so that one part looks like the incompressible Navier-Stokes equations, while the other part behaves like the compressible Navier-Stokes equations. More precisely, let (U, P) and (\mathbf{v}, η) be the solutions to the following systems, respectively:

$$\begin{cases} U \cdot \nabla U + \mathbf{v} \cdot \nabla U - \mu \Delta U + \nabla P = \mathbf{f} + \mathbf{g}, \\ \operatorname{div} U = 0, \\ U = 0 \quad \text{on } \partial\Omega \quad \text{and} \quad \int_{\Omega} P dx = 0; \end{cases} \quad (2.1)$$

and

$$\begin{cases} U \cdot \nabla \eta + \frac{\operatorname{div} \mathbf{v}}{\epsilon} = -\mathbf{v} \cdot \nabla \eta - \eta \operatorname{div} \mathbf{v} - \epsilon \operatorname{div}(P(U + \mathbf{v})), \\ U \cdot \nabla \mathbf{v} - \mu \Delta \mathbf{v} - (\mu + \lambda) \nabla \operatorname{div} \mathbf{v} + \frac{\nabla \eta + \nabla \theta}{\epsilon} = \epsilon F - \mathbf{v} \cdot \nabla \mathbf{v} - \theta \nabla \eta - \eta \nabla \theta, \\ U \cdot \nabla \theta - \kappa \Delta \theta + \frac{\operatorname{div} \mathbf{v}}{\epsilon} = \epsilon G - \mathbf{v} \cdot \nabla \theta - \eta \operatorname{div} \mathbf{v} - \theta \operatorname{div} \mathbf{v}, \\ \mathbf{v} = 0, \quad \theta = 0 \quad \text{on } \partial\Omega \quad \text{and} \quad \int_{\Omega} \eta dx = 0, \end{cases} \quad (2.2)$$

where the new force F and heat source G are defined by

$$\begin{aligned} F &= (\epsilon P + \eta)\mathbf{f} - (\epsilon P + \eta)(U + \mathbf{v}) \cdot \nabla(U + \mathbf{v}) - \theta \nabla P - P \nabla \theta, \\ G &= \Psi - (\epsilon P + \eta)(U + \mathbf{v}) \cdot \nabla \theta - (\epsilon P + \eta)\theta \operatorname{div} \mathbf{v} - P \operatorname{div} \mathbf{v}. \end{aligned}$$

It is clear to observe that $\mathbf{u} := U + \mathbf{v}$, $\rho := \epsilon P + \eta$ and θ are a solution to (1.5). Thus, we can obtain a solution of the system (1.5) if we can solve the systems (2.1) and (2.2). First, we will give the existence of weak solutions to the linearized incompressible problem (2.1) and derive a priori estimates of higher order derivatives of the unknowns (U, P) . Then, we shall show the existence of weak solutions to the linearized compressible problem (2.2) and establish uniform estimates of higher order derivatives of the unknowns $(\eta, \mathbf{v}, \theta)$.

In what follows, we assume that $\operatorname{meas}(\Omega) = 1$ without loss of generality.

2.1 Linearized incompressible equations

Let \tilde{U} and $\tilde{\mathbf{v}}$ be given functions satisfying $\tilde{U} \in H^4 \cap H_{0,\sigma}^1$ and $\tilde{\mathbf{v}} \in H^3 \cap H_0^1$. At first, we consider the linearized equations to (2.1) for given \tilde{U} and $\tilde{\mathbf{v}}$ as follows.

$$\begin{cases} (\tilde{U} + \tilde{\mathbf{v}}) \cdot \nabla U - \mu \Delta U + \nabla P = \mathbf{h}, \\ \operatorname{div} U = 0, \\ U = 0 \text{ on } \partial\Omega \quad \text{and} \quad \int_{\Omega} P dx = 0. \end{cases} \quad (2.3)$$

where $\mathbf{h} = \mathbf{f} + \mathbf{g}$.

The problem (2.3) is a Stokes problem which is solvable for arbitrarily large forces. In fact, (2.3) can be solved by using the Lax-Milgram theorem for small $\tilde{\mathbf{v}}$, and we can obtain the following existence result, the proof of which can be found, for example, in [5], and is therefore omitted here.

Lemma 2.1. *Let $\mathbf{h} \in H^{-1}$, $\tilde{U} \in H_{0,\sigma}^1$ and $\tilde{\mathbf{v}} \in H^3 \cap H_0^1$. There exists a constant a_0 depending only on μ and Ω , such that if $\|\tilde{\mathbf{v}}\|_3 < a_0$, then there exists a weak solution $(U, P) \in H_0^1 \times \bar{H}^0$ of (2.3), satisfying*

$$\|U\|_1 \leq C_0 \|\mathbf{h}\|_{-1}, \quad (2.4)$$

$$\|P\|_0 \leq C_1 \|\mathbf{h}\|_{-1} (1 + \|\mathbf{h}\|_{-1}), \quad (2.5)$$

where C_0 and C_1 are positive constants which depend only on Ω , μ and a_0 .

As for the regularity of solutions, we consider the Stokes equations:

$$\begin{aligned} -\mu \Delta U + \nabla P &= \mathbf{h} - (\tilde{U} + \tilde{\mathbf{v}}) \cdot \nabla U, \\ \operatorname{div} U &= 0. \end{aligned}$$

Then we can derive the following estimates by employing bootstrap arguments similar to those in [5].

Lemma 2.2. *Let $\mathbf{h} \in H^m$, $\tilde{U} \in H^{m+1} \cap H_{0,\sigma}^1$, $m = 0, 1, 2$, and $\tilde{\mathbf{v}}$ be the same as in Lemma 2.1. There are positive constants C_2 , C_3 and C_4 , depending only on Ω , μ and a_0 , such that if $\tilde{U} \in H_0^1$ satisfies the inequality (2.4), then*

$$\|U\|_2 + \|\nabla P\|_0 \leq C_2 \|\mathbf{h}\|_0 (\|\mathbf{h}\|_0 + 1)^4. \quad (2.6)$$

If $\tilde{U} \in H^2 \cap H_0^1$ satisfies (2.6), then

$$\|U\|_3 + \|\nabla P\|_1 \leq C_3 \|\mathbf{h}\|_1 (\|\mathbf{h}\|_1 + 1)^8, \quad (2.7)$$

and if $\tilde{U} \in H^3 \cap H_0^1$ satisfies (2.7), then

$$\|U\|_4 + \|\nabla P\|_2 \leq C_4 \|\mathbf{h}\|_2 (\|\mathbf{h}\|_2 + 1)^{12}. \quad (2.8)$$

Let $\mathbf{f}, \mathbf{g} \in H^2(\Omega)$, then it is obvious that $\mathbf{h} \in H^2(\Omega)$. We define a space K_0 by

$$K_0 := \left\{ U \in H_{0,\sigma}^4(\Omega) : \|U\|_1 \leq C_1 \|\mathbf{h}\|_1, \|U\|_2 \leq C_2 \|\mathbf{h}\|_0 (\|\mathbf{h}\|_0 + 1)^4, \right. \\ \left. \|U\|_3 \leq C_3 \|\mathbf{h}\|_1 (\|\mathbf{h}\|_1 + 1)^8, \|U\|_4 \leq C_4 \|\mathbf{h}\|_2 (\|\mathbf{h}\|_2 + 1)^{12} \right\}. \quad (2.9)$$

Thus, by Lemma 2.2 we see that the solution U of the system (2.3) also lies in K_0 for any given $\tilde{U} \in K_0$, since the constants C_1, \dots, C_4 do not depend on \tilde{U} .

2.2 Linearized compressible equations

Let $(\tilde{U}, \tilde{\mathbf{v}}, \tilde{\theta}) \in (H^4 \cap H_{0,\sigma}^1) \times (H^3 \cap H_0^1) \times (H^3 \cap H_0^1)$ be given functions. Next, we give the existence of weak solutions and derive some a priori estimates for solutions to the linearized equations of the system (2.2). For simplicity, we only consider the three-dimensional case. As aforementioned, we shall apply the Tikhonov fixed point theorem to show the existence of strong solutions to (1.5), (1.6). To this end, for given $(\tilde{\mathbf{v}}, \tilde{\theta}) \in (H^3 \cap H_0^1) \times (H^3 \cap H_0^1)$, let $(\eta, \mathbf{v}, \theta)$ be the unique solution of the following linearized system of (2.2) the existence of which will be shown below:

$$\begin{cases} U \cdot \nabla \eta + \frac{\operatorname{div} \mathbf{v}}{\epsilon} + \tilde{\mathbf{v}} \cdot \nabla \eta + \eta \operatorname{div} \tilde{\mathbf{v}} = -\epsilon \operatorname{div}(P(U + \tilde{\mathbf{v}})), \\ U \cdot \nabla \mathbf{v} - \mu \Delta \mathbf{v} - \zeta \nabla \operatorname{div} \mathbf{v} + \frac{\nabla \eta + \nabla \theta}{\epsilon} + \tilde{\theta} \nabla \eta + \eta \nabla \tilde{\theta} = \epsilon \tilde{F} - \tilde{\mathbf{v}} \cdot \nabla \tilde{\mathbf{v}}, \\ U \cdot \nabla \theta - \kappa \Delta \theta + \frac{\operatorname{div} \mathbf{v}}{\epsilon} + \eta \operatorname{div} \tilde{\mathbf{v}} = \epsilon \tilde{G} - \tilde{\mathbf{v}} \cdot \nabla \tilde{\theta} - \tilde{\theta} \operatorname{div} \tilde{\mathbf{v}}, \\ \mathbf{v} = 0, \quad \theta = 0 \quad \text{on } \partial\Omega \quad \text{and} \quad \int_{\Omega} \eta = 0, \end{cases} \quad (2.10)$$

where (U, P) be the solution of (2.3) established in Section 2.1 and the new force \tilde{F} and heat source \tilde{G} are defined by

$$\begin{aligned} \tilde{F} &= (\epsilon P + \eta) \mathbf{f} - (\epsilon P + \eta)(U + \tilde{\mathbf{v}}) \cdot \nabla(U + \tilde{\mathbf{v}}) - \tilde{\theta} \nabla P - P \nabla \tilde{\theta}, \\ \tilde{G} &= \tilde{\Psi} - (\epsilon P + \eta)(U + \tilde{\mathbf{v}}) \cdot \nabla \tilde{\theta} - (\epsilon P + \eta) \tilde{\theta} \operatorname{div} \tilde{\mathbf{v}} - P \operatorname{div} \tilde{\mathbf{v}}, \end{aligned}$$

with $\tilde{\Psi} = 2\mu D(U + \tilde{\mathbf{v}}) : D(U + \tilde{\mathbf{v}}) + \lambda(\operatorname{div}(U + \tilde{\mathbf{v}}))^2$ and $\zeta = \mu + \lambda$.

Thus, for given $\tilde{U}, \tilde{\mathbf{v}}$ and $\tilde{\theta}$, we can construct a map N :

$$N(\tilde{U}, \tilde{\mathbf{v}}, \tilde{\theta}) := (U, \mathbf{v}, \theta).$$

And, we have to show that N maps some space into itself and is weak continuous to get a fixed point of the mapping N .

In order to obtain the existence of weak solutions to (2.10), we set

$$\begin{aligned} \tilde{F}' &= \epsilon P \mathbf{f} - \epsilon P(U + \tilde{\mathbf{v}}) \cdot \nabla(U + \tilde{\mathbf{v}}) - \tilde{\theta} \nabla P - P \nabla \tilde{\theta}, \\ \tilde{G}' &= \tilde{\Psi} - \epsilon P(U + \tilde{\mathbf{v}}) \cdot \nabla \tilde{\theta} - \epsilon P \tilde{\theta} \operatorname{div} \tilde{\mathbf{v}} - P \operatorname{div} \tilde{\mathbf{v}}. \end{aligned}$$

So, $\tilde{F} = \tilde{F}' + \eta \mathbf{f} + \eta(U + \tilde{\mathbf{v}}) \cdot \nabla(U + \tilde{\mathbf{v}})$, $\tilde{G} = \tilde{G}' - \eta(U + \tilde{\mathbf{v}}) \cdot \nabla \tilde{\theta} - \eta \tilde{\theta} \operatorname{div} \tilde{\mathbf{v}}$.

2.2.1 Existence of weak solutions

Lemma 2.3. *Let $\tilde{F}', \tilde{G}' \in H^{-1}$, $\mathbf{f} \in H^2$, and (U, P) be a solution of (2.3) established in Lemma 2.1. If $\|\tilde{\mathbf{v}}\|_3 + \|\tilde{\theta}\|_3$ is sufficiently small, then there exists a unique weak solution $(\eta, \mathbf{v}, \theta) \in \tilde{H}^0 \times H_0^1 \times H_0^1$ to the problem (2.10).*

Remark 2.1. *We remark here that a weak solution of (2.10) is defined through testing the system (2.10)₁–(2.10)₃ by $\phi \in H^1$, $\mathbf{V} \in H_0^1$ and $\Theta \in H_0^1$, respectively.*

Remark 2.2. In [29] the authors used the Bergman projection to reduce the linearized problem to a boundary value problem for the transport operator equation and then proved the existence of strong solutions and the incompressible limit as $\omega = \text{Re}/(\gamma \text{Ma}^2)$ goes to ∞ . Here we shall employ the techniques of approximate systems, the Leray-Schauder fixed theorem and compactness arguments and mollifiers to show the existence and uniqueness of weak solutions directly.

Proof. The proof is broken up into three steps.

Step 1: Construction of strong solutions to the approximate problems.

We first suppose that $\tilde{F}', \tilde{G}' \in L^2$ and the case $\tilde{F}', \tilde{G}' \in H^{-1}$ will then be dealt with by a density argument later. Define the cut-off function

$$1_\alpha(\eta) = \begin{cases} 1/\alpha, & \text{if } \eta > 1/\alpha; \\ \eta, & \text{if } \eta \leq 1/\alpha. \end{cases}$$

Now, we consider the following approximate system to (2.10) for arbitrary but fixed positive numbers β, α :

$$\begin{cases} -\beta\Delta\eta = -\text{div}(U\eta) - \frac{\text{div}\mathbf{v}}{\epsilon} - \text{div}[1_\alpha(\eta)\tilde{\mathbf{v}}] - \epsilon\text{div}(P(U + \tilde{\mathbf{v}})) =: \text{div}R_1(\eta, \mathbf{v}), \\ -\mu\Delta\mathbf{v} - \zeta\nabla\text{div}\mathbf{v} = -\frac{\nabla(\eta + \theta)}{\epsilon} - U \cdot \nabla\mathbf{v} - \nabla[\tilde{\theta}1_\alpha(\eta)] \\ \quad + \epsilon[\tilde{F}' + 1_\alpha(\eta)\mathbf{f} + 1_\alpha(\eta)(U + \tilde{\mathbf{v}}) \cdot \nabla(U + \tilde{\mathbf{v}})] - \tilde{\mathbf{v}} \cdot \nabla\tilde{\mathbf{v}} =: R_2(\eta, \mathbf{v}, \theta), \\ -\kappa\Delta\theta = -U \cdot \nabla\theta - \frac{\text{div}\mathbf{v}}{\epsilon} - 1_\alpha(\eta)\text{div}\tilde{\mathbf{v}} \\ \quad + \epsilon\{\tilde{G}' - 1_\alpha(\eta)[(U + \tilde{\mathbf{v}}) \cdot \nabla\tilde{\theta} + \tilde{\theta}\text{div}\tilde{\mathbf{v}}]\} - \tilde{\mathbf{v}} \cdot \nabla\tilde{\theta} - \tilde{\theta}\text{div}\tilde{\mathbf{v}} := R_3(\eta, \mathbf{v}, \theta), \\ \mathbf{v} = 0, \quad \theta = 0, \quad \frac{\partial\eta}{\partial\mathbf{n}} = 0 \quad \text{on } \partial\Omega, \quad \text{and } \int_\Omega \eta dx = 0, \end{cases} \quad (2.11)$$

where \mathbf{n} denotes the outer normal vector of the boundary $\partial\Omega$. We point out that the introduction of the cut-off function makes it possible to establish an important estimate (2.18) below.

Next, we show the strong solvability of (2.11) by employing the Leray-Schauder fixed point theorem. To this end, we first study the following boundary problem:

$$\begin{cases} -\beta\Delta\eta = \text{div}R_1(t\chi, t\mathbf{u}), \\ -\mu\Delta\mathbf{v} - \zeta\nabla\text{div}\mathbf{v} = R_2(t\chi, t\mathbf{u}, t\vartheta), \\ -\kappa\Delta\theta = R_3(t\chi, t\mathbf{u}, t\vartheta), \\ \mathbf{v} = 0, \quad \theta = 0, \quad \frac{\partial\eta}{\partial\mathbf{n}} = 0 \quad \text{on } \partial\Omega, \quad \text{and } \int_\Omega \eta = 0 \end{cases} \quad (2.12)$$

for given $t \in [0, 1]$ and $(\chi, \mathbf{u}, \vartheta) \in \bar{H}^1 \times H_0^1 \times H_0^1$.

Recalling that $(\tilde{U}, \tilde{\mathbf{v}}, \tilde{\theta}) \in (H^4 \cap H_{0,\sigma}^1) \times (H^3 \cap H_0^1) \times (H^3 \cap H_0^1)$ and $1_\alpha(\eta) \in H^1$, by the elliptic theory (see [27, Lemma 4.32 and Lemma 4.27]), we know that there exists a unique strong solution $(\eta, \mathbf{v}, \theta) \in H^2$ satisfying (2.12) and

$$\|(\eta, \theta, \mathbf{v})\|_2 \leq C(\|\text{div}R_1(t\chi, t\mathbf{u})\|_0 + \|(R_1(t\chi, t\mathbf{u}), R_2(t\chi, t\mathbf{u}, t\vartheta), R_3(t\chi, t\mathbf{u}, t\vartheta))\|_0). \quad (2.13)$$

Then, we derive estimates for strong solutions $(\eta, \mathbf{v}, \theta)$ of the problem:

$$\begin{cases} -\beta\Delta\eta = \text{div}R_1(t\eta, t\mathbf{v}), \\ -\mu\Delta\mathbf{v} - \zeta\nabla\text{div}\mathbf{v} = R_2(t\eta, t\mathbf{v}, t\theta), \\ -\kappa\Delta\theta = R_3(t\eta, t\mathbf{v}, t\theta), \\ \mathbf{v} = 0, \quad \theta = 0, \quad \frac{\partial\eta}{\partial\mathbf{n}} = 0 \quad \text{on } \partial\Omega, \quad \text{and } \int_\Omega \eta = 0, \quad t \in [0, 1]. \end{cases} \quad (2.14)$$

Using L^2 energy estimates, we can easily obtain the following identities on $(\eta, \mathbf{v}, \theta)$:

$$\begin{aligned} \int \mu |\nabla \mathbf{v}|^2 + \zeta (\operatorname{div} \mathbf{v})^2 dx &= \frac{1}{\epsilon} \int t(\eta + \theta) \operatorname{div} \mathbf{v} dx + \int 1_\alpha(t\eta) \tilde{\theta} \operatorname{div} \mathbf{v} dx \\ &+ \int \{ \epsilon [\tilde{F}' + 1_\alpha(t\eta) \mathbf{f} + 1_\alpha(t\eta)(U + \tilde{\mathbf{v}}) \cdot \nabla(U + \tilde{\mathbf{v}})] - \tilde{\mathbf{v}} \cdot \nabla \tilde{\mathbf{v}} \} \mathbf{v} dx, \end{aligned} \quad (2.15)$$

$$\beta \int |\nabla \eta|^2 dx + \frac{1}{\epsilon} \int \eta \operatorname{div} \mathbf{v} dx = -\epsilon \int \operatorname{div}(P(U + \tilde{\mathbf{v}})) \eta dx - \frac{1}{2} \int \eta 1_\alpha(t\eta) \operatorname{div} \tilde{\mathbf{v}} dx, \quad (2.16)$$

$$\begin{aligned} \int \kappa |\nabla \theta|^2 dx + \frac{1}{\epsilon} \int \theta \operatorname{div} \mathbf{v} dx &= \epsilon \int \tilde{G}' \theta dx - \epsilon \int (1_\alpha(t\eta)(U + \tilde{\mathbf{v}}) \cdot \nabla \tilde{\theta} + 1_\alpha(t\eta) \tilde{\theta} \operatorname{div} \tilde{\mathbf{v}}) \theta dx \\ &- \int (\tilde{\mathbf{v}} \cdot \nabla \tilde{\theta} + \tilde{\theta} \operatorname{div} \tilde{\mathbf{v}}) \theta dx - \int 1_\alpha(t\eta) \theta \operatorname{div} \tilde{\mathbf{v}} dx. \end{aligned} \quad (2.17)$$

Keeping in mind that $t \leq 1$, $\|1_\alpha(\eta)\|_0^2 \leq |\Omega|/\alpha$ and $\|\eta\|_0 \leq C\|\nabla \eta\|_0$, summing up the identities (2.15)–(2.17), we deduce the following estimate:

$$\|(\eta, \mathbf{v}, \theta)\|_1 < K \equiv K(1 + \|(\tilde{F}', \tilde{G}')\|_0 + \|\mathbf{f}\|_1 + \|(U, \tilde{\mathbf{v}}, \tilde{\theta}, P)\|_2^2), \quad (2.18)$$

where the constant K may depend on other known quantities, but is independent of t .

Now for given $(\chi, \mathbf{u}, \vartheta) \in \bar{H}^1 \times H_0^1 \times H_0^1$, we define a mapping T_t by

$$T_t(\chi, \mathbf{u}, \vartheta) := (\eta_t, \mathbf{v}_t, \theta_t) \quad \text{that is the strong solution of (2.12).}$$

Then the solution operator T_t enjoys the following properties (in what follows, C denotes a general constant independent of t):

- By virtue of (2.18),

$$0 \notin (I - T_t)(\partial B_K), \quad (2.19)$$

where

$$B_K = \left\{ (\eta, \mathbf{v}, \theta) \in H^1 \times H_0^1 \times H_0^1 \mid \|(\eta, \mathbf{v}, \theta)\|_1 < K, \quad \int_\Omega \eta = 0 \right\}.$$

- Let B be a ball of center 0 in $\bar{H}^1 \times H_0^1 \times H_0^1$, then for any $t \in [0, 1]$, $T_t B$ is a precompact set of $\bar{H}^1 \times H_0^1 \times H_0^1$, since one has by (2.13) that

$$\|T_t(\chi, \mathbf{u}, \vartheta)\|_2 \leq C(\|(\chi, \mathbf{u}, \vartheta)\|_1 + 1)(1 + \|(\tilde{F}', \tilde{G}')\|_0 + \|\mathbf{f}\|_1 + \|(U, \tilde{\mathbf{v}}, \tilde{\theta}, P)\|_2^2).$$

- For any $t, s \in [0, 1]$, one has by the elliptic regularity theory (see [27, Lemma 4.32 and Lemma 4.27]) that

$$\begin{aligned} \|T_t(\chi, \mathbf{u}, \vartheta) - T_s(\chi, \mathbf{u}, \vartheta)\|_1 &\leq C(\|R_1(t\chi, t\mathbf{u}) - R_1(s\chi, s\mathbf{u})\|_0 \\ &+ \|(R_2(t\chi, t\mathbf{u}, t\vartheta), R_3(t\chi, t\mathbf{u}, t\vartheta)) - (R_2(s\chi, s\mathbf{u}, s\vartheta), R_3(s\chi, s\mathbf{u}, s\vartheta))\|_{-1}) \\ &\leq C|t - s| \|(\chi, \mathbf{u}, \vartheta)\|_1 (1 + \|(U, \tilde{\mathbf{v}}, \tilde{\theta}, P)\|_2^2). \end{aligned} \quad (2.20)$$

Consequently, we can applying the Leray-Schauder fixed point theorem [27, Section 1.4.11.7] to (2.12) to get a strong solution $(\eta, \mathbf{v}, \theta)$ of the approximate system (2.11). In what follow we write such strong solution by $(\eta_\beta^\alpha, \mathbf{v}_\beta^\alpha, \theta_\beta^\alpha)$ to indicate the dependence upon α and $\beta > 0$.

Step 2: Uniform-in- (α, β) estimates and the limit process

Recalling that the strong solutions $(\eta_\beta^\alpha, \mathbf{v}_\beta^\alpha, \theta_\beta^\alpha)$ constructed above satisfy (2.15)–(2.17) for $t = 1$, and the fact $\left|1_\alpha(\eta_\beta^\alpha)\right| \leq |\eta_\beta^\alpha|$, we find that

$$\begin{aligned} \|\mathbf{v}_\beta^\alpha\|_1^2 + \|\theta_\beta^\alpha\|_1^2 + \beta\|\nabla\eta_\beta^\alpha\|_0^2 &\leq C\|\eta_\beta^\alpha\|_0^2\left[\|\tilde{\theta}\|_2 + \epsilon(\|\mathbf{f}\|_1 + \|U + \tilde{\mathbf{v}}\|_2^4)\right] \\ &+ \epsilon\|\tilde{\mathbf{v}}\|_3 + \epsilon\|\tilde{\theta}\|_2^2(\|U + \tilde{\mathbf{v}}\|_{H^1}^2 + \|\tilde{\mathbf{v}}\|_2^2) + \|\tilde{\mathbf{v}}\|_2^2] + C\epsilon\|\eta_\beta^\alpha\|_0\|P\|_2\|U + \tilde{\mathbf{v}}\|_2 \\ &+ C\epsilon(\|\tilde{F}'\|_{-1}^2 + \|\tilde{G}'\|_{-1}^2) + C\|\tilde{\mathbf{v}}\|_1^4 + C\|\tilde{\mathbf{v}}\|_1^2\|\tilde{\theta}\|_1^2, \end{aligned} \quad (2.21)$$

provided $\|\tilde{\mathbf{v}}\|_3 + \|\tilde{\theta}\|_3$ and ϵ are sufficiently small. We remark that throughout the proof of Lemma 2.3, the smallness of $\|\tilde{\mathbf{v}}\|_3 + \|\tilde{\theta}\|_3$ and ϵ are independent of α and β .

We proceed to bound $\|\eta_\beta^\alpha\|_0$. According to the content on page 123 in [29], for a function $\xi \in L^2(\Omega)$ with $\int \xi dx = 0$, one can choose a vector field $q \in H_0^1(\Omega)$, such that

$$\operatorname{div} q = \xi, \quad \|q\|_1 \leq C(\Omega)\|\xi\|_0.$$

Multiplying (2.10)₂ by this q and integrating the resulting equation over Ω , we obtain

$$\begin{aligned} \frac{1}{\epsilon} \int (\eta_\beta^\alpha + \theta_\beta^\alpha) \operatorname{div} q dx &= - \int (\eta_\beta^\alpha \tilde{\theta}) \operatorname{div} q dx - \epsilon \int [\tilde{F}' + \eta_\beta^\alpha f + \eta_\beta^\alpha (U + \tilde{\mathbf{v}}) \cdot \nabla (U + \tilde{\mathbf{v}})] \cdot q dx \\ &+ \int \tilde{\mathbf{v}} \cdot \nabla \tilde{\mathbf{v}} \cdot q dx + \int (U \cdot \nabla \mathbf{v}) \cdot q dx + \int \mu \nabla \mathbf{v} \cdot \nabla q + \lambda \operatorname{div} \mathbf{v} \cdot \operatorname{div} q dx, \end{aligned}$$

which, by taking $\xi = \eta_\beta^\alpha + \theta_\beta^\alpha - \frac{1}{|\Omega|} \int \theta_\beta^\alpha dx$ and applying Young's inequality, gives

$$\begin{aligned} \left\| \eta_\beta^\alpha + \theta_\beta^\alpha - \frac{1}{|\Omega|} \int \theta_\beta^\alpha dx \right\|_0^2 &\leq C\epsilon(\|\mathbf{v}_\beta^\alpha\|_{H_0^1}^2 \|U\|_1^2 + \|\eta_\beta^\alpha\|_0^2 \|\tilde{\theta}\|_2^2 + \|\mathbf{v}_\beta^\alpha\|_2^2 + \|\tilde{\mathbf{v}}\|_2^4) \\ &+ C\epsilon^2[\|\tilde{F}'\|_{-1}^2 + \|\eta_\beta^\alpha\|_0^2(\|\mathbf{f}\|_1^2 + \|U + \tilde{\mathbf{v}}\|_2^4)]. \end{aligned} \quad (2.22)$$

It is easy to see that

$$\left\| \frac{1}{|\Omega|} \int \theta_\beta^\alpha dx \right\|_0^2 = \frac{1}{|\Omega|} \left(\int \theta_\beta^\alpha dx \right)^2 \leq \|\theta_\beta^\alpha\|_0^2 \quad (2.23)$$

Hence, from (2.21)–(2.23) and Poincaré's inequality we get

$$\begin{aligned} \|\eta_\beta^\alpha\|_0^2 &\leq 2\|\eta_\beta^\alpha + \theta_\beta^\alpha\|_0^2 + 2\|\theta_\beta^\alpha\|_0^2 \\ &\leq \left\| \eta_\beta^\alpha + \theta_\beta^\alpha - \frac{1}{|\Omega|} \int \theta_\beta^\alpha dx \right\|_0^2 + \left\| \frac{1}{|\Omega|} \int \theta_\beta^\alpha dx \right\|_0^2 + 2\|\theta_\beta^\alpha\|_0^2 \\ &\leq C\epsilon(\|\mathbf{v}_\beta^\alpha\|_0^2 \|U\|_1^2 + \|\eta_\beta^\alpha\|_0^2 \|\tilde{\theta}\|_2^2 + \|\mathbf{v}_\beta^\alpha\|_1^2 + \|\tilde{\mathbf{v}}\|_1^4) \\ &+ C\epsilon^2[\|\tilde{F}'\|_{-1}^2 + \|\eta_\beta^\alpha\|_0^2(\|\mathbf{f}\|_1^2 + \|U + \tilde{\mathbf{v}}\|_2^4)] \\ &+ \|\eta_\beta^\alpha\|_0^2 C[\|\tilde{\theta}\|_2 + \epsilon(\|\mathbf{f}\|_1 + \|U + \tilde{\mathbf{v}}\|_2^4) + \epsilon\|\tilde{\mathbf{v}}\|_3] \\ &+ \epsilon\|\tilde{\theta}\|_2^2(\|U + \tilde{\mathbf{v}}\|_1^2 + \|\tilde{\mathbf{v}}\|_2^2) + \|\tilde{\mathbf{v}}\|_2^2] + C\epsilon\|\eta_\beta^\alpha\|_0\|P\|_2\|U + \tilde{\mathbf{v}}\|_2 \\ &+ C\epsilon(\|\tilde{F}'\|_{-1}^2 + \|\tilde{G}'\|_{-1}^2) + C\|\tilde{\mathbf{v}}\|_1^4 + C\|\tilde{\mathbf{v}}\|_1^2\|\tilde{\theta}\|_1^2. \end{aligned} \quad (2.24)$$

If we combine (2.21) with (2.24), we see that there is a small constant $\epsilon_1 > 0$, such that for any $\epsilon \leq \epsilon_1$ and sufficiently small $\|(\tilde{\mathbf{v}}, \tilde{\theta})\|_3$,

$$\|\eta_\beta^\alpha\|_{H^0}^2 + \|\mathbf{v}_\beta^\alpha\|_0^2 + \|\theta_\beta^\alpha\|_1^2 + \beta\|\nabla\eta_\beta^\alpha\|_0 \leq C_1, \quad (2.25)$$

where C_1 is a positive constant independent of ϵ , α and β . Applying the elliptic regularity theory to (2.11)₁, we have

$$\|\eta_\beta^\alpha\|_2 \leq C(\beta) \|(\operatorname{div} R_1(\eta_\beta^\alpha, \mathbf{v}_\beta^\alpha), R_1(\eta_\beta^\alpha, \mathbf{v}_\beta^\alpha))\|_0 \leq C \equiv C(\beta, C_1),$$

where the constant C is independent of α . Therefore, by the embedding theorem $H^2 \hookrightarrow L^\infty$, we see that

$$1_\alpha(\eta_\beta^\alpha) = \eta_\beta^\alpha,$$

if α is sufficiently small while β is fixed.

With the help of this fact and the uniform estimate (2.25), we can take limit as $\alpha \rightarrow 0$ (while keeping β fixed) in (2.11) and use the standard weak convergence arguments to obtain a weak solution $(\eta_\beta, \mathbf{v}_\beta, \theta_\beta)$ of the problem (2.11) with 1 in place of 1_α , and moreover by (2.25), we have

$$\|\eta_\beta\|_0^2 + \|\mathbf{v}_\beta\|_1^2 + \|\theta_\beta\|_1^2 + \beta \|\nabla \eta_\beta\|_0 \leq C_1.$$

Recalling that the above estimate is uniform in β , we can thus take limit once more as $\beta \rightarrow 0$ in (2.11) with 1 in place of 1_α to get a weak solution $(\eta, \mathbf{v}, \theta)$ of the original problem (2.10), which is the weak limit of $(\eta_\beta, \mathbf{v}_\beta, \theta_\beta)$ and enjoys the estimate (i.e. (2.25)):

$$\|\eta\|_0^2 + \|\mathbf{v}\|_1^2 + \|\theta\|_1^2 \leq C_1. \quad (2.26)$$

Keeping in mind that the constant C_1 in (2.26) depends only on $\|(\tilde{F}', \tilde{G}')\|_{-1}$ (cf. (2.25)) and L^2 is dense in H^{-1} , we can thus obtain a weak solution $(\eta, \mathbf{v}, \theta)$ of (2.10) for $\tilde{F}', \tilde{G}' \in H^{-1}$ by a density argument. We remark that the obtained solution $(\eta, \mathbf{v}, \theta)$ satisfies the weak form of (2.10)₁, i.e.,

$$\int_\Omega \left[-\operatorname{div}(\varphi U) \eta + \left(\frac{\operatorname{div} \mathbf{v}}{\epsilon} + \tilde{\mathbf{v}} \cdot \nabla \eta + \eta \operatorname{div} \tilde{\mathbf{v}} \right) \varphi \right] dx = - \int_\Omega \epsilon \operatorname{div}(P(U + \tilde{\mathbf{v}})) \varphi dx, \quad \forall \varphi \in H^1.$$

Moreover, extending $(\eta, P, U, \mathbf{v}, \tilde{\mathbf{v}})$ outside of Ω by zero and still denoting the extended functions by $(\eta, P, U, \mathbf{v}, \tilde{\mathbf{v}})$, we find that the above identity still holds in \mathbb{R}^3 , i.e.,

$$\int_{\mathbb{R}^3} \left[-\operatorname{div}(\varphi U) \eta + \left(\frac{\operatorname{div} \mathbf{v}}{\epsilon} + \tilde{\mathbf{v}} \cdot \nabla \eta + \eta \operatorname{div} \tilde{\mathbf{v}} \right) \varphi \right] dx = - \int_{\mathbb{R}^3} \epsilon \operatorname{div}(P(U + \tilde{\mathbf{v}})) \varphi dx \quad (2.27)$$

for any $\varphi \in H^1(\mathbb{R}^3)$. This property is important in the proof of uniqueness in the next step.

Step 3: Uniqueness of weak solutions to (2.10).

Letting $(\eta_1, \mathbf{v}_1, \theta_1), (\eta_2, \mathbf{v}_2, \theta_2) \in \bar{H}^0 \times H_0^1 \times H_0^1$ be two weak solutions of the problem (2.10) and denoting $\bar{\eta} = \eta_1 - \eta_2, \bar{\mathbf{v}} = \mathbf{v}_1 - \mathbf{v}_2, \bar{\theta} = \theta_1 - \theta_2$, we find that $(\bar{\eta}, \bar{\mathbf{v}}, \bar{\theta})$ is a weak solution of the following boundary value problem:

$$\begin{cases} U \cdot \nabla \bar{\eta} + \frac{\operatorname{div} \bar{\mathbf{v}}}{\epsilon} + \tilde{\mathbf{v}} \cdot \nabla \bar{\eta} + \bar{\eta} \operatorname{div} \tilde{\mathbf{v}} = 0, \\ U \cdot \nabla \bar{\mathbf{v}} - \mu \Delta \bar{\mathbf{v}} - \zeta \nabla \operatorname{div} \bar{\mathbf{v}} + \frac{\nabla \bar{\eta} + \nabla \bar{\theta}}{\epsilon} + \tilde{\theta} \nabla \bar{\eta} + \bar{\eta} \nabla \tilde{\theta} = \epsilon (\bar{\eta} \mathbf{f} + \bar{\eta} (U + \tilde{\mathbf{v}}) \cdot \nabla (U + \tilde{\mathbf{v}})), \\ U \cdot \nabla \bar{\theta} - \kappa \Delta \bar{\theta} + \frac{\operatorname{div} \bar{\mathbf{v}}}{\epsilon} + \bar{\eta} \operatorname{div} \tilde{\mathbf{v}} = \epsilon (-\bar{\eta} (U + \tilde{\mathbf{v}}) \cdot \nabla \tilde{\theta} - \bar{\eta} \tilde{\theta} \operatorname{div} \tilde{\mathbf{v}}), \\ \bar{\mathbf{v}} = 0, \quad \bar{\theta} = 0 \quad \text{on } \partial\Omega \quad \text{and} \quad \int_\Omega \bar{\eta} = 0. \end{cases} \quad (2.28)$$

We want to test (2.28) with $\bar{\eta}$ which is unfortunately not in H^1 . To circumvent this difficulty, we use the technique of mollifiers. For a function w of x ($x \in \Omega$) we denote

$$(w)_\delta(x) = \int_\Omega \eta_\delta(x-y) w(y) dy, \quad x \in \Omega,$$

where η_δ is the Friedrichs mollifier.

Now, we take the test function φ in (2.27) to be the mollifier to find that the equation (2.28)₁ can be regularized as

$$\begin{aligned} U \cdot \nabla(\bar{\eta})_\delta + \frac{(\operatorname{div}\bar{\mathbf{v}})_\delta}{\epsilon} + \tilde{\mathbf{v}} \cdot \nabla(\bar{\eta})_\delta + (\bar{\eta}\operatorname{div}\tilde{\mathbf{v}})_\delta \\ = (U \cdot \nabla(\bar{\eta})_\delta - (U \cdot \nabla\bar{\eta})_\delta) + (\tilde{\mathbf{v}} \cdot \nabla(\bar{\eta})_\delta - (\tilde{\mathbf{v}} \cdot \nabla\bar{\eta})_\delta), \quad \text{a.e. in } \Omega, \end{aligned} \quad (2.29)$$

where $U \cdot \nabla\bar{\eta} = \operatorname{div}(U\bar{\eta}) - \bar{\eta}\operatorname{div}U$ and $\tilde{\mathbf{v}} \cdot \nabla\bar{\eta} = \operatorname{div}(\tilde{\mathbf{v}}\bar{\eta}) - \bar{\eta}\operatorname{div}\tilde{\mathbf{v}}$.

Since $\eta \in L^2$, and $U, \tilde{\mathbf{v}} \in H_0^1 \cap W^{1,\infty}$ by Sobolev's imbedding theorem, we have by Friedrichs' lemma on commutators [27, Lemma 3.1] that

$$U \cdot \nabla(\bar{\eta})_\delta - (U \cdot \nabla\bar{\eta})_\delta, \quad \tilde{\mathbf{v}} \cdot \nabla(\bar{\eta})_\delta - (\tilde{\mathbf{v}} \cdot \nabla\bar{\eta})_\delta \rightarrow 0 \quad \text{strongly in } L^2(\Omega). \quad (2.30)$$

We should point here that (2.30) is shown in the whole space \mathbb{R}^3 . However, if we extend $U, \tilde{\mathbf{v}}$ and η outside of Ω by zero, we still have $\eta \in L^2(\mathbb{R}^3)$ and $U, \tilde{\mathbf{v}} \in H_0^1(\mathbb{R}^3) \cap W^{1,\infty}(\mathbb{R}^3)$, and hence we can apply Lemma 3.1 of [27] to get (2.30).

Now, we test the equations (2.29), (2.28)₂ and (2.28)₃ by $(\bar{\eta})_\delta, \bar{\mathbf{v}}$ and $\bar{\theta}$, respectively, to deduce that

$$\begin{aligned} & \int (\mu|\nabla\bar{\mathbf{v}}|^2 + \zeta|\nabla\bar{\mathbf{v}}|^2 + \kappa|\nabla\bar{\theta}|^2) dx \\ & = \epsilon \int \bar{\eta}\bar{\mathbf{v}}[\mathbf{f} + (U + \tilde{\mathbf{v}}) \cdot \nabla(U + \tilde{\mathbf{v}})] - \bar{\eta}\bar{\theta}[(U + \tilde{\mathbf{v}}) \cdot \nabla\bar{\theta} + \bar{\theta}\operatorname{div}\tilde{\mathbf{v}}] dx \\ & + \int \left[\frac{1}{2}\operatorname{div}\tilde{\mathbf{v}}|(\bar{\eta})_\delta|^2 + \bar{\eta}\bar{\theta}\operatorname{div}\mathbf{v} - (\bar{\eta}\operatorname{div}\tilde{\mathbf{v}})_\delta(\eta)_\delta - \bar{\eta}\bar{\theta}\operatorname{div}\tilde{\mathbf{v}} \right] dx \\ & + \int \left\{ \left(\frac{\operatorname{div}\bar{\mathbf{v}}}{\epsilon}\eta - \frac{(\operatorname{div}\bar{\mathbf{v}})_\delta}{\epsilon}(\eta)_\delta \right) + [(U \cdot \nabla(\bar{\eta})_\delta - (U \cdot \nabla\bar{\eta})_\delta) + (\tilde{\mathbf{v}} \cdot \nabla(\bar{\eta})_\delta - (\tilde{\mathbf{v}} \cdot \nabla\bar{\eta})_\delta)](\eta)_\delta \right\} dx. \end{aligned}$$

Letting $\delta \rightarrow 0$ in the above identity, using (2.30) and noting that (cf. (2.30))

$$\int \left(\frac{1}{2}\operatorname{div}\tilde{\mathbf{v}}|(\bar{\eta})_\delta|^2 - (\bar{\eta}\operatorname{div}\tilde{\mathbf{v}})_\delta(\eta)_\delta + \frac{\operatorname{div}\bar{\mathbf{v}}}{\epsilon}\eta - \frac{(\operatorname{div}\bar{\mathbf{v}})_\delta}{\epsilon}(\eta)_\delta \right) dx \rightarrow 0,$$

we conclude that

$$\begin{aligned} & \int (\mu|\nabla\bar{\mathbf{v}}|^2 + \zeta|\nabla\bar{\mathbf{v}}|^2 + \kappa|\nabla\bar{\theta}|^2) dx \leq C(\|\bar{\eta}\|_0^2 + \|\bar{\theta}\|_0^2)[(\|U\|_2 + \|\tilde{\mathbf{v}}\|_2)\|\bar{\theta}\|_3 + \|\tilde{\mathbf{v}}\|_3(1 + \|\bar{\theta}\|_2)] \\ & + \epsilon C(\|\bar{\eta}\|_0^2 + \|\bar{\mathbf{v}}\|_0^2)(\|\mathbf{f}\|_2 + \|U\|_2 + \|\tilde{\mathbf{v}}\|_2). \end{aligned} \quad (2.31)$$

On the other hand, for the inhomogeneous Stokes problem:

$$\begin{cases} -\mu\Delta\bar{\mathbf{v}} + \frac{\nabla\bar{\eta} + \nabla\bar{\theta}}{\epsilon} = -U \cdot \nabla\bar{\mathbf{v}} + \zeta\nabla\operatorname{div}\bar{\mathbf{v}} + \epsilon[\bar{\eta}\mathbf{f} + \bar{\eta}(U + \tilde{\mathbf{v}}) \cdot \nabla(U + \tilde{\mathbf{v}})] - (\bar{\theta}\nabla\bar{\eta} + \bar{\eta}\nabla\bar{\theta}), \\ \operatorname{div}\bar{\mathbf{v}} = \operatorname{div}\bar{\mathbf{v}}, \\ \bar{\mathbf{v}} = 0, \end{cases}$$

we have the estimate

$$\begin{aligned} & \epsilon\|\bar{\mathbf{v}}\|_1 + \|\bar{\eta} + \bar{\theta}\|_0 \\ & \leq \epsilon C \left\{ (\|U\|_2 + 1)\|\bar{\mathbf{v}}\|_1 + \|\bar{\mathbf{v}}\|_1 + \epsilon\|\bar{\eta}\|_0 \left[\|\mathbf{f}\|_2 + (\|U\|_2 + \|\tilde{\mathbf{v}}\|_2)^2 \right] + \|\bar{\eta}\|_0\|\bar{\theta}\|_2 \right\}. \end{aligned} \quad (2.32)$$

Combining (2.31) with (2.32), making use of Poincaré's inequality, and recalling the smallness of $\epsilon, \|\tilde{\mathbf{v}}\|_3$ and $\|\bar{\theta}\|_3$, we find that

$$\|\bar{\eta}\|_0 + \|\bar{\mathbf{v}}\|_1 + \|\bar{\theta}\|_1 \leq 0, \quad (2.33)$$

which implies $\eta_1 = \eta_2, \mathbf{v}_1 = \mathbf{v}_2, \theta_1 = \theta_2$. This completes the proof of Lemma 2.3. \square

In order to get higher order uniform-in- ϵ estimates of the $(\eta, \mathbf{v}, \theta)$, we have to bound $\|\mathbf{v}\|_1$ and $\|\theta\|_1$ as shown in the following lemma.

Lemma 2.4. *Let (U, P) be the solution of (2.3) given in Lemma 2.1. Let $\tilde{F}, \tilde{G} \in H^{-1}$. Then we have the following uniform-in- ϵ estimate:*

$$\begin{aligned} \|\mathbf{v}\|_1^2 + \|\theta\|_1^2 \leq & C_5 \left[\|\tilde{\mathbf{v}}\|_3 \|\eta\|_0^2 + \epsilon^2 (\|\tilde{F}\|_{-1}^2 + \|\tilde{G}\|_{-1}^2) + \epsilon \|P\|_2 (\|U\|_2 + \|\tilde{\mathbf{v}}\|_2) \|\eta\|_0 \right. \\ & \left. + \|\tilde{\mathbf{v}}\|_1^4 + \|\eta\|_1^2 \|\tilde{\theta}\|_1^2 + \|\tilde{\mathbf{v}}\|_1^2 (\|\tilde{\theta}\|_1^2 + \|\eta\|_1^2) \right], \end{aligned} \quad (2.34)$$

where the constant $C_5 > 0$ is independent of ϵ .

Proof. Multiplying (2.10)₁, (2.10)₂ and (2.10)₃ by η , v and θ in L^2 respectively, and summing up the resulting equations, we find that

$$\begin{aligned} & C' (\|\mathbf{v}\|_1^2 + \|\theta\|_1^2) \\ &= -\frac{1}{\epsilon} \int (\operatorname{div} \mathbf{v} (\eta + \theta) + v \cdot (\nabla \eta + \nabla \theta)) dx - \int [(U + \tilde{\mathbf{v}}) \cdot \nabla \eta \cdot \eta + \eta^2 \operatorname{div} \tilde{\mathbf{v}} + \epsilon \operatorname{div} (P(U + \tilde{\mathbf{v}})) \eta] dx \\ & \quad + \int (\epsilon \tilde{F} - \tilde{\mathbf{v}} \cdot \nabla \tilde{\mathbf{v}} - \tilde{\theta} \nabla \eta - \eta \nabla \tilde{\theta}) \cdot \mathbf{v} dx + \int (\epsilon \tilde{G} - \tilde{\mathbf{v}} \cdot \nabla \tilde{\theta} - (\eta + \tilde{\theta}) \operatorname{div} \tilde{\mathbf{v}}) \theta dx \\ &\leq \frac{1}{2} \|\tilde{\mathbf{v}}\|_3 \|\eta\|_0^2 + \delta (\|\mathbf{v}\|_1^2 + \|\theta\|_1^2) + C_\delta [\epsilon^2 (\|\tilde{F}\|_{-1}^2 + \|\tilde{G}\|_{-1}^2) + \epsilon \|P\|_2 (\|U\|_2 + \|\tilde{\mathbf{v}}\|_2) \|\eta\|_0 \\ & \quad + \|\tilde{\mathbf{v}}\|_1^4 + \|\eta\|_1^2 \|\tilde{\theta}\|_1^2 + \|\tilde{\mathbf{v}}\|_1^2 (\|\tilde{\theta}\|_1^2 + \|\eta\|_1^2)], \end{aligned} \quad (2.35)$$

where we have used integration by parts, Sobolev's inequality and the fact that

$$-\frac{1}{\epsilon} \int ((\eta + \theta) \operatorname{div} \mathbf{v} + \mathbf{v} \cdot (\nabla \eta + \nabla \theta)) dx = \frac{1}{\epsilon} \int [(\eta + \theta) \operatorname{div} \mathbf{v} - (\eta + \theta) \operatorname{div} \mathbf{v}] dx = 0.$$

Finally, if we take δ in (2.35) suitably small and apply Poincaré's inequality, we obtain the estimate (2.34). \square

2.2.2 Stokes problem

We rewrite the momentum equations (2.10)₂ as an inhomogeneous Stokes problem to derive the desired bounds for $\|\mathbf{v}\|_3$ and $\left\| \frac{\nabla(\eta + \theta)}{\epsilon} \right\|_1$:

$$\begin{cases} -\mu \Delta \mathbf{v} + \frac{\nabla \eta + \nabla \theta}{\epsilon} = \epsilon \tilde{F} - \tilde{\mathbf{v}} \cdot \nabla \tilde{\mathbf{v}} - \tilde{\theta} \nabla \eta - \eta \nabla \tilde{\theta} - U \cdot \nabla \mathbf{v} + \zeta \nabla \operatorname{div} \mathbf{v}, \\ \operatorname{div} \mathbf{v} = \operatorname{div} \mathbf{v}, \\ \mathbf{v} = 0, \quad \text{on } \Omega. \end{cases} \quad (2.36)$$

By the usual estimates for the steady Stokes problem (cf. Galdi's book [14, Chapter IV]), Sobolev's embedding $H^2 \hookrightarrow L^\infty$ and the inequality

$$\|\operatorname{div} \mathbf{v}\|_1^2 \leq \delta \|\mathbf{v}\|_3^2 + C_\delta \|\mathbf{v}\|_1^2, \quad (2.37)$$

we have

$$\begin{aligned} \|\mathbf{v}\|_2 + \left\| \frac{\nabla \eta + \nabla \theta}{\epsilon} \right\|_0 \leq & C (\|\epsilon \tilde{F}\|_0 + \|\tilde{\mathbf{v}} \cdot \nabla \tilde{\mathbf{v}}\|_0 + \|\tilde{\theta} \nabla \eta\|_0 + \|\eta \nabla \tilde{\theta}\|_0 \\ & + \|U \cdot \nabla \mathbf{v}\|_0 + \|\operatorname{div} \mathbf{v}\|_1). \end{aligned} \quad (2.38)$$

and

$$\begin{aligned} \|\mathbf{v}\|_3 + \left\| \frac{\nabla\eta + \nabla\theta}{\epsilon} \right\|_1 \leq & C(\|\epsilon\tilde{F}\|_1 + \|\tilde{\mathbf{v}} \cdot \nabla\tilde{\mathbf{v}}\|_1 + \|\tilde{\theta}\nabla\eta\|_1 + \|\eta\nabla\tilde{\theta}\|_1 \\ & + \|U \cdot \nabla\mathbf{v}\|_1 + \|\operatorname{div}\mathbf{v}\|_2), \end{aligned}$$

which together with (2.34), (2.37) and (2.38) yields

$$\begin{aligned} \|\mathbf{v}\|_3 + \left\| \frac{\nabla\eta + \nabla\theta}{\epsilon} \right\|_1 \leq & C_6(1 + \|U\|_2^2) \left\{ \epsilon(\|\tilde{F}\|_1 + \|\tilde{G}\|_{-1}) + \|\tilde{\mathbf{v}}\|_3^2 + \|\tilde{\theta}\|_3\|\eta\|_2 \right. \\ & \left. + \|\tilde{\mathbf{v}}\|_3^{1/2}\|\eta\|_0 + \left[\epsilon\|P\|_2(\|U\|_2 + \|\tilde{\mathbf{v}}\|_2)\|\eta\|_0 \right]^{1/2} + \|\tilde{\mathbf{v}}\|_1(\|\tilde{\theta}\|_1 + \|\eta\|_1) \right\} + \|\nabla^2\operatorname{div}\mathbf{v}\|_0, \end{aligned} \quad (2.39)$$

where C_6 is a positive constant independent of ϵ .

2.2.3 Estimate of $\|\nabla^2\operatorname{div}\mathbf{v}\|_0$

As in [34, 36, 18], in order to control the term $\|\nabla^2\operatorname{div}\mathbf{v}\|_0$ we divide it into the interior part and the part near the boundary. We remark that here we have to carefully deal with the terms which involve with the large parameter $1/\epsilon$ in (2.10).

I. Interior estimate

First, we derive the interior estimate of $\nabla^2\operatorname{div}\mathbf{v}$ by using the estimate (2.38). Let χ_0 be a C_0^∞ -function, then we have

Lemma 2.5. *There is a positive constant C_7 independent of ϵ , such that*

$$\begin{aligned} & \mu\|\chi_0\nabla^2\mathbf{v}\|_0^2 + \zeta\|\chi_0\nabla\operatorname{div}\mathbf{v}\|_0^2 + \kappa\|\chi_0\nabla^2\theta\|_0^2 \\ & \leq C_7 \left[(\|U\|_3 + \|\tilde{\mathbf{v}}\|_3)\|\eta\|_1^2 + \epsilon^2(\|\tilde{F}\|_0^2 + \|\tilde{G}\|_0^2) + \epsilon\|P\|_2(\|U\|_2 + \|\tilde{\mathbf{v}}\|_2)\|\eta\|_1 + \|\tilde{\mathbf{v}}\|_2^4 \right. \\ & \quad \left. + \|\eta\|_2^2\|\tilde{\theta}\|_2^2 + \|U\|_3(\|\mathbf{v}\|_1^2 + \|\theta\|_1^2) + \|\tilde{\mathbf{v}}\|_2^2(\|\tilde{\theta}\|_1^2 + \|\eta\|_1^2) + \|\mathbf{v}\|_1^2 \right] + \delta \left\| \frac{\nabla\eta + \nabla\theta}{\epsilon} \right\|_0^2. \end{aligned} \quad (2.40)$$

Proof. We differentiate (2.10) with respect to x to get that

$$\begin{cases} U^j\partial_{ij}^2\eta + \frac{\partial_i\operatorname{div}\mathbf{v}}{\epsilon} = -\tilde{v}^j\partial_{ij}^2\eta - \partial_i(U^j + \tilde{v}^j)\partial_j\eta - \tilde{v}^j\partial_{ij}^2\eta - \partial_i(\eta\operatorname{div}\tilde{\mathbf{v}}) - \epsilon\partial_i\operatorname{div}(P(U + \tilde{\mathbf{v}})), \\ U^j\partial_{ij}^2v^k + \partial_iU^j\partial_jv^k - \mu\partial_{ijj}^3v^k - \zeta\partial_{ik}^2\operatorname{div}\mathbf{v} + \frac{\partial_{ik}^2\eta + \partial_{ik}^2\theta}{\epsilon} = \epsilon\partial_i\tilde{F}^k - \partial_i(\tilde{v}^j\partial_j\tilde{v}^k) - \partial_{ik}^2(\tilde{\theta}\eta), \\ U^j\partial_{ij}^2\theta + \partial_iU^j\partial_j\theta - \kappa\partial_{ijj}^3\theta + \frac{\partial_i\operatorname{div}\mathbf{v}}{\epsilon} = \epsilon\partial_i\tilde{G} - \partial_i(\tilde{v}^j\partial_j\tilde{\theta}) - \partial_i((\eta + \tilde{\theta})\operatorname{div}\tilde{\mathbf{v}}). \end{cases} \quad (2.41)$$

Multiplying (2.41)₁, (2.41)₂ and (2.41)₃ by $\chi_0^2\partial_i\eta$, $\chi_0^2\partial_iv^k$ and $\chi_0^2\partial_i\theta$ in L^2 respectively, and summing up the resulting equations, we find that

$$\begin{aligned} & \mu\|\chi_0\partial_{ij}^2v^k\|_0^2 + \zeta\|\chi_0\partial_i\operatorname{div}\mathbf{v}\|_0^2 + \kappa\|\chi_0\partial_{ij}^2\theta\|_0^2 \\ & = -\frac{1}{\epsilon} \int \chi_0^2\partial_i\operatorname{div}\mathbf{v}(\partial_i\eta + \partial_i\theta) + \chi_0^2\partial_iv^k\partial_k(\partial_i\eta + \partial_i\theta)dx \\ & \quad - \int (2\mu\chi_0\partial_j\chi_0\partial_{ij}^2v^k\partial_iv^k + 2\zeta\chi_0\partial_k\chi_0\partial_i\operatorname{div}\mathbf{v}\partial_iv^k + 2\kappa\chi_0\partial_j\chi_0\partial_{ij}^2\theta\partial_i\theta)dx \\ & \quad - \int \chi_0^2 \left[\partial_i(U^j + \tilde{v}^j)\partial_j\eta + (U^j + \tilde{v}^j)\partial_{ji}^2\eta + \partial_i\eta\operatorname{div}\tilde{\mathbf{v}} + \eta\partial_i\operatorname{div}\tilde{\mathbf{v}} + \epsilon\partial_i\operatorname{div}(P(U + \tilde{\mathbf{v}})) \right] \partial_i\eta dx \\ & \quad + \int \chi_0^2 (\epsilon\partial_i\tilde{F}^k - \partial_i(\tilde{\mathbf{v}} \cdot \nabla\tilde{v}^k) - \partial_{ik}^2(\tilde{\theta}\eta)) \partial_iv^k dx \\ & \quad + \int \chi_0^2 (\epsilon\partial_i\tilde{G} - \partial_i(\tilde{v}^j\partial_j\tilde{\theta}) - \partial_i((\eta + \tilde{\theta})\operatorname{div}\tilde{\mathbf{v}})) \partial_i\theta dx. \end{aligned}$$

If we apply partial integrations to the above identity, employ Sobolev's and Young's inequalities and the fact that

$$\begin{aligned}
& -\frac{1}{\epsilon} \int \chi_0^2 \partial_i \operatorname{div} \mathbf{v} (\partial_i \eta + \partial_i \theta) + \chi_0^2 \partial_i v^k \partial_k (\partial_i \eta + \partial_i \theta) dx \\
&= \frac{1}{\epsilon} \int 2\chi_0 \partial_k \chi_0 \partial_i v^k (\partial_i \eta + \partial_i \theta) dx + \frac{1}{\epsilon} \int \chi_0^2 \partial_i v^k \partial_k (\partial_i \eta + \partial_i \theta) - \chi_0^2 \partial_i v^k \partial_k (\partial_i \eta + \partial_i \theta) dx \\
&= \frac{1}{\epsilon} \int 2\chi_0 \partial_k \chi_0 \partial_i v^k (\partial_i \eta + \partial_i \theta) dx \leq \delta \left\| \frac{\nabla \eta + \nabla \theta}{\epsilon} \right\|_0^2 + C_\delta \|\mathbf{v}\|_1^2,
\end{aligned}$$

we infer by summing up i, j, k that

$$\begin{aligned}
& \mu \|\chi_0 \nabla^2 \mathbf{v}\|_0^2 + \zeta \|\chi_0 \nabla \operatorname{div} \mathbf{v}\|_0^2 + \kappa \|\chi_0 \nabla^2 \theta\|_0^2 \\
& \leq (\|U\|_3 + \|\tilde{\mathbf{v}}\|_3) \|\eta\|_1^2 + \delta (\|\mathbf{v}\|_2^2 + \|\theta\|_2^2 + \left\| \frac{\nabla \eta + \nabla \theta}{\epsilon} \right\|_0^2) \\
& \quad + C_\delta \left[\epsilon^2 (\|\tilde{F}\|_0^2 + \|\tilde{G}\|_0^2) + \epsilon \|P\|_2 (\|U\|_2 + \|\tilde{\mathbf{v}}\|_2) \|\eta\|_1 + \|\tilde{\mathbf{v}}\|_2^4 + \|\eta\|_2^2 \|\tilde{\theta}\|_2^2 \right. \\
& \quad \left. + \|U\|_3 (\|\mathbf{v}\|_1^2 + \|\theta\|_1^2) + \|\tilde{\mathbf{v}}\|_2^2 (\|\tilde{\theta}\|_1^2 + \|\eta\|_1^2) + \|\mathbf{v}\|_1^2 \right],
\end{aligned}$$

which, by using Poincaré's inequality and choosing δ appropriately small, implies the lemma. \square

Lemma 2.6. *There is a positive constant C_8 independent of ϵ , such that*

$$\begin{aligned}
& \mu \|\chi_0 \nabla^3 \mathbf{v}\|_0^2 + \zeta \|\chi_0 \nabla^2 \operatorname{div} \mathbf{v}\|_0^2 + \kappa \|\chi_0 \nabla^3 \theta\|_0^2 \\
& \leq C_8 \left[(\|U\|_3 + \|\tilde{\mathbf{v}}\|_3) \|\eta\|_2^2 + \epsilon^2 (\|\tilde{F}\|_1^2 + \|\tilde{G}\|_1^2) + \epsilon \|P\|_3 (\|U\|_3 + \|\tilde{\mathbf{v}}\|_3) \|\eta\|_2 + \|\tilde{\mathbf{v}}\|_3^4 \right. \\
& \quad \left. + \|\eta\|_2^2 \|\tilde{\theta}\|_2^2 + \|U\|_3 (\|\mathbf{v}\|_2^2 + \|\theta\|_2^2) + \|\tilde{\mathbf{v}}\|_2^2 (\|\tilde{\theta}\|_2^2 + \|\eta\|_2^2) + \|\mathbf{v}\|_2^2 \right] + \delta \left\| \frac{\nabla \eta + \nabla \theta}{\epsilon} \right\|_1^2.
\end{aligned} \tag{2.42}$$

Proof. We differentiate (2.10) twice with respect to x to get that

$$\left\{ \begin{aligned}
& U^j \partial_{ij}^3 \eta + \frac{\partial_i \operatorname{div} \mathbf{v}}{\epsilon} = -\tilde{v}^j \partial_{ij}^3 \eta - \partial_{il}^2 (U^j + \tilde{v}^j) \partial_j \eta - \partial_i (U^j + \tilde{v}^j) \partial_{jl}^2 \eta \\
& \quad - \partial_l (U^j + \tilde{v}^j) \partial_{ij}^2 \eta - \partial_{il} (\eta \operatorname{div} \tilde{\mathbf{v}}) - \epsilon \partial_{il}^2 \operatorname{div} (P(U + \tilde{\mathbf{v}})), \\
& U^j \partial_{ijl}^3 v^k + \partial_l U^j \partial_{ij}^2 v^k + \partial_{il}^2 U^j \partial_j v^k + \partial_i U^j \partial_{lj}^2 v^k - \mu \partial_{iljj}^4 v^k - \zeta \partial_{ilk}^3 \operatorname{div} \mathbf{v} + \frac{\partial_{ilk}^3 \eta + \partial_{ilk}^3 \theta}{\epsilon} \\
& \quad = \epsilon \partial_{il}^2 \tilde{F}^k - \partial_{il}^2 (\tilde{v}^j \partial_j \tilde{v}^k) - \partial_{ilk}^3 (\tilde{\theta} \eta), \\
& U^j \partial_{ijl}^3 \theta + \partial_l U^j \partial_{ij}^2 \theta + \partial_{il}^2 U^j \partial_j \theta + \partial_i U^j \partial_{lj}^2 \theta - \kappa \partial_{iljj}^4 \theta + \frac{\partial_{il}^2 \operatorname{div} \mathbf{v}}{\epsilon} \\
& \quad = \epsilon \partial_{il}^2 \tilde{G} - \partial_{il}^2 (\tilde{v}^j \partial_j \tilde{\theta}) - \partial_{il}^2 ((\eta + \tilde{\theta}) \operatorname{div} \tilde{\mathbf{v}}).
\end{aligned} \right. \tag{2.43}$$

Multiplying (2.43)₁, (2.43)₂ and (2.43)₃ again by $\chi_0^2 \partial_{il}^2 \eta$, $\chi_0^2 \partial_{il}^2 v^k$ and $\chi_0^2 \partial_{il}^2 \theta$ respectively, and summing up the resulting equations, we deduce that

$$\begin{aligned}
& \mu \|\chi_0 \partial_{ij}^3 v^k\|_0^2 + \zeta \|\chi_0 \partial_{il}^2 \operatorname{div} \mathbf{v}\|_0^2 + \kappa \|\chi_0 \partial_{ilj}^3 \theta\|_0^2 \\
&= -\frac{1}{\epsilon} \int \chi_0^2 \partial_{il} \operatorname{div} \mathbf{v} (\partial_{il} \eta + \partial_{il}^2 \theta) + \chi_0^2 \partial_{il}^2 v^k \partial_k (\partial_{il}^2 \eta + \partial_{il}^2 \theta) dx \\
& \quad - \int 2\chi_0 \partial_j \chi_0 (\mu \partial_{ilj}^3 v^k \partial_{il}^2 v^k + \kappa \partial_{ilj}^3 \theta \partial_{il}^2 \theta) + 2\zeta \chi_0 \partial_k \chi_0 \partial_{il} \operatorname{div} v \partial_{il} v^k dx \\
& \quad - \int \left[U^j \partial_{ij}^3 \eta + \tilde{v}^j \partial_{ij}^3 \eta + \partial_{il}^2 (U^j + \tilde{v}^j) \partial_j \eta + \partial_i (U^j + \tilde{v}^j) \partial_{jl}^2 \eta \right. \\
& \quad \left. + \partial_l (U^j + \tilde{v}^j) \partial_{ij}^2 \eta + \partial_{il}^2 (\eta \operatorname{div} \tilde{\mathbf{v}}) + \epsilon \partial_{il}^2 \operatorname{div} (P(U + \tilde{\mathbf{v}})) \right] \chi_0^2 \partial_{il}^2 \eta dx
\end{aligned}$$

$$\begin{aligned}
& + \int \left[\epsilon \partial_{\tilde{u}}^2 \tilde{F}^k - \partial_{\tilde{u}}^2 (\tilde{v}^j \partial_j \tilde{v}^k) - \partial_{\tilde{u}k}^3 (\tilde{\theta} \eta) - (U^j \partial_{\tilde{u}jl}^3 v^k + \partial_l U^j \partial_{\tilde{u}j}^2 v^k \right. \\
& \quad \left. + \partial_{\tilde{u}}^2 U^j \partial_j v^k + \partial_i U^j \partial_{\tilde{u}j}^2 v^k) \right] \chi_0^2 \partial_{\tilde{u}}^2 v^k dx \\
& + \int \left[\epsilon \partial_{\tilde{u}}^2 \tilde{G} - \partial_{\tilde{u}}^2 (\tilde{v}^j \partial_j \tilde{\theta}) - \partial_{\tilde{u}}^2 ((\eta + \tilde{\theta}) \operatorname{div} \tilde{\mathbf{v}}) - (U^j \partial_{\tilde{u}jl}^3 \theta + \partial_l U^j \partial_{\tilde{u}j}^2 \theta \right. \\
& \quad \left. + \partial_{\tilde{u}}^2 U^j \partial_j \theta + \partial_i U^j \partial_{\tilde{u}j}^2 \theta) \right] \chi_0^2 \partial_{\tilde{u}} \theta dx.
\end{aligned}$$

We integrate by parts the above identity, utilize Sobolev's inequality and the fact that

$$\begin{aligned}
& -\frac{1}{\epsilon} \int \chi_0^2 \partial_{\tilde{u}}^2 \operatorname{div} \mathbf{v} (\partial_{\tilde{u}}^2 \eta + \partial_{\tilde{u}}^2 \theta) + \chi_0^2 \partial_{\tilde{u}}^2 v^k \partial_k (\partial_{\tilde{u}}^2 \eta + \partial_{\tilde{u}}^2 \theta) dx \\
& = \frac{1}{\epsilon} \int \chi_0^2 \partial_{\tilde{u}}^2 v^k \partial_k (\partial_{\tilde{u}}^2 \eta + \partial_{\tilde{u}}^2 \theta) - \chi_0^2 \partial_{\tilde{u}}^2 v^k \partial_k (\partial_{\tilde{u}}^2 \eta + \partial_{\tilde{u}}^2 \theta) dx + \frac{1}{\epsilon} \int 2\chi_0 \partial_k \chi_0 \partial_{\tilde{u}}^2 v^k (\partial_{\tilde{u}}^2 \eta + \partial_{\tilde{u}}^2 \theta) dx \\
& = \frac{1}{\epsilon} \int 2\chi_0 \partial_k \chi_0 \partial_{\tilde{u}}^2 v^k (\partial_{\tilde{u}}^2 \eta + \partial_{\tilde{u}}^2 \theta) dx \leq \delta \left\| \frac{\nabla \eta + \nabla \theta}{\epsilon} \right\|_1^2 + C_\delta \|\mathbf{v}\|_2^2,
\end{aligned}$$

and sum up i, j, k to infer that

$$\begin{aligned}
& \mu \|\chi_0 \nabla^3 \mathbf{v}\|_0^2 + \zeta \|\chi_0 \nabla^2 \operatorname{div} \mathbf{v}\|_0^2 + \kappa \|\chi_0 \nabla^3 \theta\|_0^2 \\
& \leq (\|U\|_3 + \|\tilde{\mathbf{v}}\|_3) \|\eta\|_2^2 + \delta (\|\mathbf{v}\|_3^2 + \|\theta\|_3^2 + \left\| \frac{\nabla \eta + \nabla \theta}{\epsilon} \right\|_1^2) \\
& \quad + C_\delta [\epsilon^2 (\|\tilde{F}\|_1^2 + \|\tilde{G}\|_1^2) + \epsilon \|P\|_3 (\|U\|_3 + \|\tilde{\mathbf{v}}\|_3) \|\eta\|_2 + \|\tilde{\mathbf{v}}\|_3^4 + \|\eta\|_2^2 \|\tilde{\theta}\|_2^2 \\
& \quad + \|U\|_3 (\|\mathbf{v}\|_2^2 + \|\theta\|_2^2) + \|\tilde{\mathbf{v}}\|_2^2 (\|\tilde{\theta}\|_2^2 + \|\eta\|_2^2) + \|\mathbf{v}\|_2^2],
\end{aligned}$$

which, by employing Poincaré's inequality and choosing δ suitably small, gives the lemma. \square

II. Boundary estimate

Next, we shall use the method of local coordinates to bound $\nabla^2 \operatorname{div} \mathbf{v}$ in the vicinity of the boundary (also see [34, 36, 18]). For completeness, we briefly describe the local coordinates as follows. First, one constructs the local coordinates by the isothermal coordinates $\lambda(\varphi, \phi)$ to derive an estimate near the boundary (see also [34, 36]), where

$$\lambda_\varphi \cdot \lambda_\varphi > 0, \quad \lambda_\phi \cdot \lambda_\phi > 0, \quad \lambda_\varphi \cdot \lambda_\phi = 0.$$

The boundary $\partial\Omega$ can be covered by a finite number of bounded open sets $W^k \subset \mathbb{R}^3$, $k = 1, 2, \dots, L$, such that for any $x \in W^k \cap \Omega$,

$$x = \Lambda^k(\varphi, \phi, r) \equiv \lambda^k(\varphi, \phi) + r \mathbf{n}(\lambda^k(\varphi, \phi)), \quad (2.44)$$

where $\lambda^k(\varphi, \phi)$ is the isothermal coordinates and \mathbf{n} is the unit outer normal to $\partial\Omega$.

Without confusion, we will omit the superscript k in each W^k in the following. We construct the orthonormal system corresponding to the local coordinates by

$$e_1 := \frac{\lambda_\varphi}{|\lambda_\varphi|}, \quad e_2 := \frac{\lambda_\phi}{|\lambda_\phi|}, \quad e_3 := e_1 \times e_2 \equiv \mathbf{n}(\lambda). \quad (2.45)$$

By a straightforward calculation, we see that for sufficiently small r ,

$$\begin{aligned}
C^2 \ni J & := \det \operatorname{Jac} \Lambda = \det \frac{\partial x}{\partial(\varphi, \phi, r)} = \Lambda_\varphi \times \Lambda_\phi \cdot e_3 \\
& = |\lambda_\varphi| |\lambda_\phi| + r (|\lambda_\varphi| \mathbf{n}_\phi \cdot e_2 + |\lambda_\phi| \mathbf{n}_\varphi \cdot e_1) + r^2 [(\mathbf{n}_\varphi \cdot e_1)(\mathbf{n}_\phi \cdot e_2) - (\mathbf{n}_\varphi \cdot e_2)(\mathbf{n}_\phi \cdot e_1)] > 0.
\end{aligned}$$

And, we can easily derive the following relations as $(\text{Jac}\Lambda^{-1}) \circ \Lambda = (\text{Jac}\Lambda)^{-1}$ (also see [34]):

$$[\nabla(\Lambda^{-1})^1] \circ \Lambda = J^{-1}(\Lambda_\phi \times e_3), \quad (2.46)$$

$$[\nabla(\Lambda^{-1})^2] \circ \Lambda = J^{-1}(e_3 \times \Lambda_\varphi), \quad (2.47)$$

$$[\nabla(\Lambda^{-1})^3] \circ \Lambda = J^{-1}(\Lambda_\varphi \times \Lambda_\phi) = e_3, \quad (2.48)$$

where the symbol \circ stands for the composite of operators. Set $y := (\varphi, \phi, r)$, and denote by D_i the partial derivative with respect to y_i in local coordinates. We set the unknowns in local coordinates

$$\hat{\eta}(t, y) := \eta(t, \Lambda(y)), \quad \hat{\mathbf{v}}(t, y) := \mathbf{v}(t, \Lambda(y)), \quad \hat{\theta}(t, y) := \theta(t, \Lambda(y)),$$

and the knowns

$$\hat{U}(t, y) := U(t, \Lambda(y)), \quad \hat{\mathbf{v}}(t, y) := \hat{\mathbf{v}}(t, \Lambda(y)), \quad \hat{\theta}(t, y) := \hat{\theta}(t, \Lambda(y)).$$

Then, we rewrite the system (2.10) in $[0, T] \times \tilde{\Omega}$, where $\tilde{\Omega} := \Lambda^{-1}(W \cap \Omega)$, as follows.

$$\left\{ \begin{array}{l} \hat{U}^j a_{kj} D_k \hat{\eta} + \frac{a_{kj} D_k \hat{v}^j}{\epsilon} = -\hat{v}^j a_{kj} D_k \hat{\eta} - \hat{\eta} a_{kj} D_k \hat{v}^j - \epsilon a_{kj} D_k (\hat{P}(\hat{U}^j + \hat{v}^j)), \\ \hat{U}^j a_{kj} D_k \hat{v}^i - \mu a_{kj} D_k (a_{lj} D_l \hat{v}^i) - \zeta a_{ki} D_k (a_{lj} D_l \hat{v}^j) + \frac{a_{ki} D_k (\hat{\eta} + \hat{\theta})}{\epsilon} \\ \quad = \epsilon \hat{F}^i - \hat{v}^j a_{kj} D_k \hat{v}^i - \hat{\theta} a_{ki} D_k \hat{\eta} - \hat{\eta} a_{ki} D_k \hat{\theta}, \\ \hat{U}^j a_{kj} D_k \hat{\theta} - \kappa a_{kj} D_k (a_{lj} D_l \hat{\theta}) + \frac{a_{kj} D_k \hat{v}^j}{\epsilon} = \epsilon \hat{G} - \hat{v}^j a_{kj} D_k \hat{\theta} - \hat{\eta} a_{kj} D_k \hat{v}^j - \hat{\theta} a_{kj} D_k \hat{v}^j, \end{array} \right. \quad (2.49)$$

with boundary conditions

$$\hat{\mathbf{v}}(t, y) = 0, \quad \hat{\theta}(t, y) = 0 \quad \text{on } \partial\tilde{\Omega}, \quad (2.50)$$

where a_{ij} is the (i, j) -th entry of the matrix $\text{Jac}(\Lambda^{-1}) = \frac{\partial y}{\partial x}$. Clearly, a_{ij} is a C^2 -function, and it follows from (2.46)–(2.48) that

$$\sum_{j=1}^3 a_{3j} a_{3j} = |\mathbf{n}|^2 = 1, \quad \sum_{j=1}^3 a_{1j} a_{3j} = \sum_{j=1}^3 a_{2j} a_{3j} = 0. \quad (2.51)$$

Moreover, this localized system has the following properties (see also [34]):

Proposition 2.1. $D_i(Ja_{ij}) = 0$, for $j = 1, 2, 3$; $\varsigma D_\tau \hat{\mathbf{v}} = 0$, $\varsigma D_\tau D_\xi \hat{\mathbf{v}} = 0$ on $\partial\tilde{\Omega}$ in the tangential directions $\tau, \xi = 1, 2$, where $\varsigma \in C_0^\infty(\Lambda^{-1}(W))$. Similarly, $\varsigma D_\tau \hat{\theta} = 0$, $\varsigma D_\tau D_\xi \hat{\theta} = 0$ on $\partial\tilde{\Omega}$.

Recalling $D_j = \sum_{i=1}^3 a_{ji} \partial_i$, we will frequently make use of the following relations without pointing out explicitly in subsequent calculations:

$$\|D_y \hat{\mathbf{v}}\|_{L^p(\Omega)} \leq C \|\nabla_x \mathbf{v}\|_{L^p(\Omega)}, \quad \|D_y^2 \hat{\mathbf{v}}\|_{L^p(\Omega)} \leq C \|\nabla_x \mathbf{v}\|_{W^{1,p}(\Omega)}, \quad 1 \leq p \leq \infty. \quad (2.52)$$

The above inequalities apply to η , θ and U , $\hat{\mathbf{v}}$, $\hat{\theta}$, too.

By virtue of the interpolation $\|\cdot\|_{H^2}^2 \leq \delta \|\cdot\|_{H^3}^2 + C_\delta \|\cdot\|_{H^1}^2$, the boundary estimate of $\|\nabla^2 \text{div} \mathbf{u}\|_{L_t^2(L^2)}$ can be reduced to the boundedness of

$$\int_0^t \int_{\tilde{\Omega}} J \chi^2 |D_y^2 (a_{ji} D_j U^i)| dy ds,$$

where χ is a $C_0^\infty(\Lambda^{-1}(W))$ -function. So, we can split the estimate of derivatives on the boundary into two parts: the estimate of derivatives in the tangential directions and in the normal

direction.

Part 1. Estimate of derivatives in the tangential directions

First, we apply $D_{\tau\xi}^2$ to (2.49) with τ, ξ being the tangential directions to $\partial\tilde{\Omega}$ to get

$$\left\{ \begin{aligned} & \hat{U}^j a_{kj} D_{k\tau\xi}^3 \hat{\eta} + \frac{1}{\epsilon} D_{\tau\xi}^2 [a_{kj} D_k \hat{v}^j] \\ & = D_\xi (\hat{U}^j a_{kj}) D_{k\tau}^2 \hat{\eta} + D_{\tau\xi}^2 (\hat{U}^j a_{kj}) D_k \hat{\eta} + D_\tau (\hat{U}^j a_{kj}) D_{k\xi}^2 \hat{\eta} \\ & \quad - D_{\tau\xi}^2 [\hat{v}^j a_{kj} D_k \hat{\eta} + \hat{\eta} a_{kj} D_k \hat{v}^j + \epsilon a_{kj} D_k (\hat{P}(\hat{U}^j + \hat{v}^j))], \\ & \hat{U}^j a_{kj} D_{k\tau\xi}^3 \hat{v}^i - D_{\tau\xi}^2 [\mu a_{kj} D_k (a_{lj} D_l \hat{v}^i) + \zeta a_{ki} D_k (a_{lj} D_l \hat{v}^j)] + \frac{1}{\epsilon} D_{\tau\xi}^2 [a_{ki} D_k (\hat{\eta} + \hat{\theta})] \\ & = D_\xi (\hat{U}^j a_{kj}) D_{k\tau}^2 \hat{v}^i + D_{\tau\xi}^2 (\hat{U}^j a_{kj}) D_k \hat{v}^i + D_\tau (\hat{U}^j a_{kj}) D_{k\xi}^2 \hat{v}^i \\ & \quad + D_{\tau\xi}^2 [\epsilon \hat{F}^i - \hat{v}^j a_{kj} D_k \hat{v}^i - \hat{\theta} a_{ki} D_k \hat{\eta} - \hat{\eta} a_{ki} D_k \hat{\theta}], \\ & \hat{U}^j a_{kj} D_{k\tau\xi}^3 \hat{\theta} - \kappa D_{\tau\xi}^2 [a_{kj} D_k (a_{lj} D_l \hat{\theta})] + \frac{1}{\epsilon} D_{\tau\xi}^2 [a_{kj} D_k \hat{v}^j] \\ & = D_\xi (\hat{U}^j a_{kj}) D_{k\tau}^2 \hat{\theta} + D_{\tau\xi}^2 (\hat{U}^j a_{kj}) D_k \hat{\theta} + D_\tau (\hat{U}^j a_{kj}) D_{k\xi}^2 \hat{\theta} \\ & \quad + D_{\tau\xi}^2 [\epsilon \hat{G} - \hat{v}^j a_{kj} D_k \hat{\theta} - \hat{\eta} a_{kj} D_k \hat{v}^j - \hat{\theta} a_{kj} D_k \hat{v}^j]. \end{aligned} \right. \quad (2.53)$$

We multiply (2.53)₁, (2.53)₂ and (2.53)₃ by $J\chi^2 D_{\tau\xi} \hat{\eta}$, $J\chi^2 D_{\tau\xi} \hat{v}^i$ and $J\chi^2 D_{\tau\xi} \hat{\theta}$ respectively, and integrate the resulting identities to deduce that

$$\begin{aligned} & - \int_{\tilde{\Omega}} D_{\tau\xi}^2 [\mu a_{kj} D_k (a_{lj} D_l \hat{v}^i) + \zeta a_{ki} D_k (a_{lj} D_l \hat{v}^j)] \cdot J\chi^2 D_{\tau\xi}^2 \hat{v}^i dy \\ & - \int_{\tilde{\Omega}} \kappa D_{\tau\xi}^2 [a_{kj} D_k (a_{lj} D_l \hat{\theta})] \cdot J\chi^2 D_{\tau\xi}^2 \hat{\theta} dy \\ & + \int_{\tilde{\Omega}} \hat{U}^j a_{kj} (D_{k\tau\xi}^3 \hat{\eta} \cdot J\chi^2 D_{\tau\xi}^2 \hat{\eta} + D_{k\tau\xi}^3 \hat{v}^i \cdot J\chi^2 D_{\tau\xi}^2 \hat{v}^i + D_{k\tau\xi}^3 \hat{\theta} \cdot J\chi^2 D_{\tau\xi}^2 \hat{\theta}) dy \\ & + \frac{1}{\epsilon} \int_{\tilde{\Omega}} \{ D_{\tau\xi}^2 [a_{kj} D_k \hat{v}^j] \cdot J\chi^2 D_{\tau\xi}^2 \hat{\eta} + D_{\tau\xi}^2 [a_{ki} D_k (\hat{\eta} + \hat{\theta})] \cdot J\chi^2 D_{\tau\xi}^2 \hat{v}^i + D_{\tau\xi}^2 [a_{kj} D_k \hat{v}^j] \cdot J\chi^2 D_{\tau\xi}^2 \hat{\theta} \} dy \\ = & \int_{\tilde{\Omega}} \{ D_\xi (\hat{U}^j a_{kj}) D_{k\tau}^2 \hat{\eta} + D_{\tau\xi}^2 (\hat{U}^j a_{kj}) D_k \hat{\eta} + D_\tau (\hat{U}^j a_{kj}) D_{k\xi}^2 \hat{\eta} \\ & - D_{\tau\xi}^2 [\hat{v}^j a_{kj} D_k \hat{\eta} + \hat{\eta} a_{kj} D_k \hat{v}^j + \epsilon a_{kj} D_k (\hat{P}(\hat{U}^j + \hat{v}^j))] \} \cdot J\chi^2 D_{\tau\xi}^2 \hat{\eta} dy \\ & + \int_{\tilde{\Omega}} \{ D_\xi (\hat{U}^j a_{kj}) D_{k\tau}^2 \hat{v}^i + D_{\tau\xi}^2 (\hat{U}^j a_{kj}) D_k \hat{v}^i + D_\tau (\hat{U}^j a_{kj}) D_{k\xi}^2 \hat{v}^i \\ & + D_{\tau\xi}^2 [\epsilon \hat{F}^i - \hat{v}^j a_{kj} D_k \hat{v}^i - \hat{\theta} a_{ki} D_k \hat{\eta} - \hat{\eta} a_{ki} D_k \hat{\theta}] \} \cdot J\chi^2 D_{\tau\xi}^2 \hat{v}^i dy \\ & + \int_{\tilde{\Omega}} \{ D_\xi (\hat{U}^j a_{kj}) D_{k\tau}^2 \hat{\theta} + D_{\tau\xi}^2 (\hat{U}^j a_{kj}) D_k \hat{\theta} + D_\tau (\hat{U}^j a_{kj}) D_{k\xi}^2 \hat{\theta} \\ & + D_{\tau\xi}^2 [\epsilon \hat{G} - \hat{v}^j a_{kj} D_k \hat{\theta} - \hat{\eta} a_{kj} D_k \hat{v}^j - \hat{\theta} a_{kj} D_k \hat{v}^j] \} \cdot J\chi^2 D_{\tau\xi}^2 \hat{\theta} dy. \end{aligned} \quad (2.54)$$

Now, we denote LHS of (2.54) $:= L'_1 + L'_2 + L'_3 + L'_4$ and have to deal with each term due to integration by part and the boundary conditions.

$$\begin{aligned} L'_1 = & - \int_{\tilde{\Omega}} \left\{ D_{\tau\xi}^2 (\mu a_{kj}) D_k (a_{lj} D_l \hat{v}^i) + D_\tau (\mu a_{kj}) D_{k\xi}^2 (a_{lj} D_l \hat{v}^i) \right. \\ & + D_\xi (\mu a_{kj}) D_k [D_\tau (a_{lj}) D_l \hat{v}^i + a_{lj} D_{\tau l}^2 \hat{v}^i] + \mu a_{kj} D_{k\xi} [D_\tau (a_{lj}) D_l \hat{v}^i + a_{lj} D_{\tau l}^2 \hat{v}^i] \\ & \left. + D_{\tau\xi}^2 (\zeta a_{ki}) D_k (a_{lj} D_l \hat{v}^j) + D_\tau (\zeta a_{ki}) D_{k\xi}^2 (a_{lj} D_l \hat{v}^j) \right\} \end{aligned}$$

$$\begin{aligned}
& + D_\xi(\zeta a_{ki}) D_k [D_\tau(a_{lj}) D_l \hat{v}^j + a_{lj} D_{l\tau}^2 \hat{v}^j] \\
& + \zeta a_{ki} D_{k\xi}^2 [D_\tau(a_{lj}) D_l \hat{v}^j + a_{lj} D_{l\tau} \hat{v}^j] \} \cdot J \chi^2 D_{\tau\xi}^2 \hat{v}^i dy
\end{aligned}$$

where

$$\begin{aligned}
& - \int_{\tilde{\Omega}} [\mu a_{kj} D_{k\xi}^2 (a_{lj} D_{l\tau}^2 \hat{v}^i) + \zeta a_{ki} D_{k\xi}^2 (a_{lj} D_{l\tau}^2 \hat{v}^j)] \cdot J \chi^2 D_{\tau\xi}^2 \hat{v}^i dy \\
= & \int_{\tilde{\Omega}} \mu J \chi^2 a_{kj} D_{k\tau\xi}^3 \hat{v}^i a_{lj} D_{l\tau\xi}^3 \hat{v}^i + \zeta J \chi^2 a_{ki} D_{k\tau\xi}^3 \hat{v}^i a_{lj} D_{l\tau\xi}^3 \hat{v}^j dy \\
& + \int_{\tilde{\Omega}} [D_k(\mu J \chi^2 a_{kj}) D_\xi(a_{lj} D_{l\tau} \hat{v}^i) \cdot D_{\tau\xi}^2 \hat{v}^i + \mu J \chi^2 a_{kj} D_{k\tau\xi}^3 \hat{v}^i D_\xi(a_{lj}) D_{l\tau}^2 \hat{v}^i \\
& + D_k(\zeta J \chi^2 a_{ki}) D_\xi(a_{lj} D_{l\tau}^2 \hat{v}^i) \cdot D_{\tau\xi}^2 \hat{v}^i + \mu J \chi^2 a_{ki} D_{k\tau\xi}^3 \hat{v}^i D_\xi(a_{lj}) D_{l\tau}^2 \hat{v}^j] dy,
\end{aligned}$$

and

$$\begin{aligned}
L'_2 = & \int_{\tilde{\Omega}} \kappa J \chi^2 a_{kj} D_{k\tau\xi}^3 \hat{\theta} a_{lj} D_{l\tau\xi}^3 \hat{\theta} dy \\
& + \int_{\tilde{\Omega}} \left[D_k(\kappa J \chi^2 a_{kj}) D_\xi(a_{lj} D_{l\tau}^2 \hat{\theta}) D_{\tau\xi}^2 \hat{\theta} + \kappa J \chi^2 a_{kj} D_{k\tau\xi}^3 \hat{\theta} D_\xi(a_{lj}) D_{l\tau}^2 \hat{\theta} \right] dy \\
& - \int_{\tilde{\Omega}} \left\{ D_{\tau\xi}^2(\kappa a_{kj}) D_k(a_{lj} D_l \hat{\theta}) + D_\tau(\kappa a_{kj}) D_{k\xi}^2(a_{lj} D_l \hat{\theta}) \right. \\
& \left. + D_\xi(\kappa a_{kj}) D_k \left[D_\tau(a_{lj}) D_l \hat{\theta} + a_{lj} D_{l\tau}^2 \hat{\theta} \right] + \kappa a_{kj} D_{k\xi}^2 D_\tau(a_{lj}) D_l \hat{\theta} \right\} \cdot J \chi^2 D_{\tau\xi}^2 \hat{\theta} dy.
\end{aligned}$$

On the other hand, recalling that $a_{kj} D_k \hat{U}^j = 0$, we have

$$\begin{aligned}
L'_3 = & - \frac{1}{2} \int_{\tilde{\Omega}} J \chi^2 a_{kj} D_k \hat{U}^j (|D_{\tau\xi}^2 \hat{\eta}|^2 + |D_{\tau\xi}^2 \hat{v}^i|^2 + |D_{\tau\xi}^2 \hat{\theta}|^2) dy \\
& - \frac{1}{2} \int_{\tilde{\Omega}} D_k(J \chi^2 a_{kj}) \hat{U}^j (|D_{\tau\xi}^2 \hat{\eta}|^2 + |D_{\tau\xi}^2 \hat{v}^i|^2 + |D_{\tau\xi}^2 \hat{\theta}|^2) dy \\
= & - \frac{1}{2} \int_{\tilde{\Omega}} D_k(J \chi^2 a_{kj}) \hat{U}^j (|D_{\tau\xi}^2 \hat{\eta}|^2 + |D_{\tau\xi}^2 \hat{v}^i|^2 + |D_{\tau\xi}^2 \hat{\theta}|^2) dy.
\end{aligned}$$

As for L'_4 , in view of the following identity

$$\begin{aligned}
& \frac{1}{\epsilon} \int_{\tilde{\Omega}} D_{\tau\xi}^2 [a_{ki} D_k(\hat{\eta} + \hat{\theta})] \cdot J \chi^2 D_{\tau\xi}^2 \hat{v}^i dy = - \frac{1}{\epsilon} \int_{\tilde{\Omega}} J \chi^2 D_{\tau\xi}^2 (a_{ki} D_k \hat{v}^i) D_{\tau\xi}^2 (\hat{\eta} + \hat{\theta}) dy \\
& + \frac{1}{\epsilon} \int_{\tilde{\Omega}} \left[D_{\tau\xi}^2(a_{ki}) D_k(\hat{\eta} + \hat{\theta}) + D_\tau(a_{ki}) D_{k\xi}^2(\hat{\eta} + \hat{\theta}) + D_\xi(a_{ki}) D_{k\tau}^2(\hat{\eta} + \hat{\theta}) \right] \cdot J \chi^2 D_{\tau\xi}^2 \hat{v}^i dy \\
& + \frac{1}{\epsilon} \int_{\tilde{\Omega}} D_{\xi\tau}^2(\hat{\eta} + \hat{\theta}) [D_{\xi\tau}^2(a_{ki}) D_k \hat{v}^i + D_\tau(a_{ki}) D_{k\xi} \hat{v}^i + D_\xi(a_{ki}) D_{k\tau} \hat{v}^i \\
& - D_k(J \chi^2 a_{ki}) D_{\tau\xi}^2 \hat{v}^i - D_k(a_{ki}) J \chi^2 D_{\tau\xi}^2 \hat{v}^i] dy,
\end{aligned}$$

we deduce that

$$\begin{aligned}
L'_4 = & \frac{1}{\epsilon} \int_{\tilde{\Omega}} \left[D_{\tau\xi}^2(a_{ki}) D_k(\hat{\eta} + \hat{\theta}) + D_\tau(a_{ki}) D_{k\xi}^2(\hat{\eta} + \hat{\theta}) + D_\xi(a_{ki}) D_{k\tau}^2(\hat{\eta} + \hat{\theta}) \right] \cdot J \chi^2 D_{\tau\xi}^2 \hat{v}^i dy \\
& + \frac{1}{\epsilon} \int_{\tilde{\Omega}} D_{\xi\tau}^2(\hat{\eta} + \hat{\theta}) [D_{\xi\tau}^2(a_{ki}) D_k \hat{v}^i + D_\tau(a_{ki}) D_{k\xi}^2 \hat{v}^i + D_\xi(a_{ki}) D_{k\tau}^2 \hat{v}^i \\
& - D_k(J \chi^2 a_{ki}) D_{\tau\xi}^2 \hat{v}^i - D_k(a_{ki}) J \chi^2 D_{\tau\xi}^2 \hat{v}^i] dy.
\end{aligned}$$

Substituting the above estimates into (2.54), using Sobolev's and Young's inequalities and taking into account the property (2.52), we deduce that

$$\begin{aligned}
& \int_{\tilde{\Omega}} \mu J \chi^2 a_{kj} D_{k\tau\xi}^3 \hat{v}^i a_{lj} D_{l\tau\xi}^3 \hat{v}^i dy + \int_{\tilde{\Omega}} \zeta J \chi^2 a_{ki} D_{k\tau\xi}^3 \hat{v}^i a_{lj} D_{l\tau\xi}^3 \hat{v}^j dy + \int_{\tilde{\Omega}} \kappa J \chi^2 a_{kj} D_{k\tau\xi}^3 \hat{\theta} a_{lj} D_{l\tau\xi}^3 \hat{\theta} dy \\
& \leq C_9 \left[\|U\|_3 \|\eta\|_2^2 + \|\tilde{\mathbf{v}}\|_3 \|\eta\|_2^2 + \epsilon \|P\|_3 (\|U\|_3 + \|\tilde{\mathbf{v}}\|_3) \|\eta\|_2 + \|U\|_3^2 (\|\mathbf{v}\|_1 + \|\theta\|_1) \right. \\
& \quad + \epsilon^2 (\|\tilde{F}\|_1^2 + \|\tilde{G}\|_1^2) + \|\tilde{\mathbf{v}}\|_2^4 + \|\eta\|_2^2 (\|\tilde{\mathbf{v}}\|_2^2 + \|\tilde{\theta}\|_2^2) + \|\tilde{\mathbf{v}}\|_2^2 \|\tilde{\theta}\|_2^2 \\
& \quad \left. + (\|\mathbf{v}\|_1^2 + \|\theta\|_1^2) \right] + \delta \left(\|\mathbf{v}\|_3^2 + \|\theta\|_3^2 + \left\| \frac{\nabla \eta + \nabla \theta}{\epsilon} \right\|_1^2 \right), \tag{2.55}
\end{aligned}$$

where C_9 is a constant.

Part 2. Estimate of derivatives in the normal direction

We multiply (2.49)₂ by a_{3i} to obtain that

$$\begin{aligned}
& -(\mu + \zeta) D_3(a_{lj} D_l \hat{v}^j) + \frac{1}{\epsilon} D_3(\hat{\eta} + \hat{\theta}) \\
& = -a_{3i} \hat{U}^j a_{kj} D_k \hat{v}^i + \epsilon a_{3i} \hat{F}^i - a_{3i} \hat{v}^j a_{kj} D_k \hat{v}^i - \hat{\theta} D_3 \hat{\eta} - \hat{\eta} D_3 \hat{\theta} \\
& \quad + \mu \left[a_{3i} a_{kj} D_k (a_{lj} D_l \hat{v}^i) - D_3(a_{lj} D_l \hat{v}^j) \right], \tag{2.56}
\end{aligned}$$

where the last term in RHS of (2.56) can be written as follows.

$$\begin{aligned}
& \mu \left[a_{3i} a_{kj} D_k (a_{lj} D_l \hat{v}^i) - D_3(a_{lj} D_l \hat{v}^j) \right] \\
& = \mu \left[D_3(a_{3j}) D_3 \hat{v}^j + D_3(a_{\tau j}) D_\tau \hat{v}^j + a_{\tau j} D_{3\tau}^2 \hat{v}^j - a_{3j} D_3(a_{3j}) a_{3i} D_3 \hat{v}^i \right. \\
& \quad \left. - a_{\tau j} a_{3i} D_\tau a_{lj} D_l \hat{v}^i - a_{\tau j} a_{\xi j} a_{3i} D_{\tau\xi}^2 \hat{v}^i - a_{3j} a_{3i} D_3(a_{\tau j}) D_\tau \hat{v}^i \right], \quad \tau, \xi = 1, 2, \tag{2.57}
\end{aligned}$$

which does not include the term $D_{33} \hat{\mathbf{v}}$.

Step 1. To continue our estimate, we show the following lemma.

Lemma 2.7. *There are a constant C_{10} and a small $\delta > 0$, such that*

$$\begin{aligned}
& \frac{\mu + \zeta}{2} \int_{\tilde{\Omega}} J \chi^2 |D_{\tau 3}^2(a_{lj} D_l \hat{v}^j)|^2 dy + \frac{\kappa}{2} \int_{\tilde{\Omega}} J \chi^2 |D_{\tau 3}^2(a_{kj} D_k \hat{\theta})|^2 dy \\
& \leq C_{10} \left\{ \|U\|_4^2 \|\mathbf{v}\|_1^2 + \epsilon^2 \|\tilde{F}\|_1^2 + \|\tilde{\mathbf{v}}\|_2^4 + \|\tilde{\theta}\|_2^2 \|\eta\|_2^2 + \int_{\tilde{\Omega}} J \chi^2 |D_{\tau\xi y}^3 \mathbf{v}|^2 dy \right. \\
& \quad + \left[\|U\|_3 \|\eta\|_2^2 + \|\tilde{\mathbf{v}}\|_3 \|\eta\|_2^2 + (1 + \|U\|_3) \|\theta\|_2^2 + \epsilon \|P\|_3 (\|U\|_3 + \|\tilde{\mathbf{v}}\|_3) \|\eta\|_2 \right. \\
& \quad \left. \left. + \epsilon^2 \|\tilde{G}\|_1^2 + \|\tilde{\mathbf{v}}\|_2^2 \|\tilde{\theta}\|_2^2 + \|\tilde{\mathbf{v}}\|_2^2 \|\eta\|_2^2 \right] \right\} + \delta (\|\mathbf{v}\|_3^2 + \|\theta\|_3^2). \tag{2.58}
\end{aligned}$$

Proof. We differentiate (2.56) with respect to y_τ ($\tau = 1, 2$), then multiply $-J \chi^2 D_{\tau 3} (a_{lj} D_l \hat{v}^j)$ in $L^2(\tilde{\Omega})$ to get

$$\begin{aligned}
& \frac{\mu + \zeta}{2} \int_{\tilde{\Omega}} J \chi^2 |D_{\tau 3}^2(a_{lj} D_l \hat{v}^j)|^2 dy - \frac{1}{\epsilon} \int_{\tilde{\Omega}} J \chi^2 D_{\tau 3}^2(\hat{\eta} + \hat{\theta}) D_{\tau 3}^2(a_{lj} D_l \hat{v}^j) dy \\
& \leq C (\|a_{3i} \hat{U}^j a_{kj} D_k \hat{v}^i\|_1^2 + \epsilon^2 \|\hat{F}^i\|_1^2 + \|a_{3i} \hat{v}^j a_{kj} D_k \hat{v}^i\|_1^2 \\
& \quad + \|\hat{\theta} D_3 \hat{\eta} + \hat{\eta} D_3 \hat{\theta}\|_1^2) + C \int_{\tilde{\Omega}} J \chi^2 |D_{\tau\xi y}^3 v|^2 dy \\
& \leq C (\|U\|_2^2 \|v\|_2^2 + \epsilon^2 \|\tilde{F}\|_1^2 + \|\tilde{v}\|_2^4 + \|\tilde{\theta}\|_2^2 \|\hat{\eta}\|_2^2) + C \int_{\tilde{\Omega}} J \chi^2 |D_{\tau\xi y}^3 v|^2 dy. \tag{2.59}
\end{aligned}$$

In the mean while, we apply D_{τ_3} to (2.49)₃ and (2.49)₁, take the product of the resulting equations with $J\chi^2 D_{\tau_3}^2 \hat{\theta}$ and $J\chi^2 D_{\tau_3}^2 \hat{\eta}$ in $L^2(\tilde{\Omega})$, and sum then two identities to get

$$\begin{aligned}
& - \int_{\tilde{\Omega}} \kappa D_{\tau_3} \left[a_{kj} D_k (a_{lj} D_l \hat{\theta}) \right] \cdot J\chi^2 D_{\tau_3}^2 \hat{\theta} dy + \frac{1}{\epsilon} \int_{\tilde{\Omega}} D_{\tau_3}^2 (a_{kj} D_k \hat{v}^j) \cdot J\chi^2 (D_{\tau_3}^2 \hat{\theta} + D_{\tau_3}^2 \hat{\eta}) dy \\
& + \int_{\tilde{\Omega}} \left[D_{\tau_3}^2 (\hat{U}^j a_{kj} D_k \hat{\eta}) \cdot J\chi^2 D_{\tau_3}^2 \hat{\eta} + D_{\tau_3}^2 (\hat{U}^j a_{kj} D_k \hat{\theta}) \cdot J\chi^2 D_{\tau_3}^2 \hat{\theta} \right] dy \\
& = - \int_{\tilde{\Omega}} D_{\tau_3}^2 \left\{ \hat{v}^j a_{kj} D_k \hat{\eta} + \hat{\eta} a_{kj} D_k \hat{v}^j + \epsilon a_{kj} D_k \left[\hat{P} (\hat{U}^j + \hat{v}^j) \right] \right\} \cdot J\chi^2 D_{\tau_3}^2 \hat{\eta} \\
& + \int_{\tilde{\Omega}} D_{\tau_3}^2 \left(\epsilon \hat{G} - \hat{v}^j a_{kj} D_k \hat{\theta} - \hat{\eta} a_{kj} D_k \hat{v}^j - \hat{\theta} a_{kj} D_k \hat{v}^j \right) \cdot J\chi^2 D_{\tau_3}^2 \hat{\theta} dy.
\end{aligned} \tag{2.60}$$

We denote LHS of (2.60) := $L_1'' + L_2'' + L_3''$. To control L_k'' , we integrate by part to deduce that

$$\begin{aligned}
L_1'' & = \kappa \int_{\tilde{\Omega}} J\chi^2 D_{\tau_3}^2 (a_{kj} D_k \hat{\theta}) D_{\tau_3}^2 (a_{lj} D_l \hat{\theta}) dy \\
& - \int_{\tilde{\Omega}} \kappa J\chi^2 D_{\tau_3}^2 \hat{\theta} \left(D_{\tau_3}^2 (a_{kj}) D_k (a_{lj} D_l \hat{\theta}) + D_{\tau} (a_{kj}) D_{3k}^2 (a_{lj} D_l \hat{\theta}) + D_3 (a_{kj}) D_{k\tau}^2 (a_{lj} D_l \hat{\theta}) \right) dy \\
& + \kappa \int_{\tilde{\Omega}} D_k (J\chi^2 a_{kj}) D_{\tau_3}^2 (\hat{\theta}) D_{\tau_3}^2 (a_{lj} D_l \hat{\theta}) dy \\
& - \kappa \int_{\tilde{\Omega}} J\chi^2 \left(D_{\tau_3}^2 (a_{kj}) D_k \hat{\theta} + D_{\tau} (a_{kj}) D_{3k}^2 \hat{\theta} + D_3 (a_{kj}) D_{k\tau}^2 \hat{\theta} \right) \cdot D_{\tau_3}^2 (a_{lj} D_l \hat{\theta}) dy
\end{aligned}$$

and

$$\begin{aligned}
L_3'' & = \int_{\tilde{\Omega}} J\chi^2 D_{\tau_3}^2 \hat{\eta} \cdot \left(D_{\tau_3}^2 (\hat{U}^j a_{kj}) D_k \hat{\eta} + D_{\tau} (\hat{U}^j a_{kj}) D_{k3}^2 \hat{\eta} + D_3 (\hat{U}^j a_{kj}) D_{k\tau}^2 \hat{\eta} \right) dy \\
& - \frac{1}{2} \int_{\tilde{\Omega}} D_k (J\chi^2 a_{kj}) \hat{U}^j |D_{\tau_3}^2 \hat{\eta}|^2 + J\chi^2 a_{kj} D_k \hat{U}^j |D_{\tau_3}^2 \hat{\eta}|^2 dy \\
& + \int_{\tilde{\Omega}} \left(D_{\tau_3}^2 (\hat{U}^j a_{kj}) D_k \hat{\theta} + D_{\tau} (\hat{U}^j a_{kj}) D_{k3}^2 \hat{\theta} + D_3 (\hat{U}^j a_{kj}) D_{k\tau}^2 \hat{\theta} \right) \cdot J\chi^2 D_{\tau_3}^2 \hat{\theta} dy \\
& - \frac{1}{2} \int_{\tilde{\Omega}} D_k (J\chi^2 a_{kj}) \hat{U}^j |D_{\tau_3}^2 \hat{\theta}|^2 + J\chi^2 a_{kj} D_k \hat{U}^j |D_{\tau_3}^2 \hat{\theta}|^2 dy.
\end{aligned}$$

Inserting the estimates for L_1'' and L_3'' into (2.60), we obtain

$$\begin{aligned}
& \frac{\kappa}{2} \int_{\tilde{\Omega}} J\chi^2 |D_{\tau_3}^2 (a_{kj} D_k \hat{\theta})|^2 dy + \frac{1}{\epsilon} \int_{\tilde{\Omega}} J\chi^2 D_{\tau_3}^2 (\hat{\eta} + \hat{\theta}) D_{\tau_3}^2 (a_{lj} D_l \hat{v}^j) dy \\
& \leq \|\theta\|_2^2 + \delta \left(\|D_{3k}^2 (a_{lj} D_l \hat{\theta})\|_0^2 + \|D_{k\tau}^2 (a_{lj} D_l \hat{\theta})\|_0^2 + \|\theta\|_3^2 \right) \\
& + C \left[\|U\|_3 \|\eta\|_2^2 + \|U\|_3 \|\theta\|_2^2 + \|\tilde{v}\|_3 \|\eta\|_2^2 + \epsilon \|P\|_3 (\|U\|_3 + \|\tilde{v}\|_3) \|\eta\|_2 \right. \\
& \left. + \epsilon^2 \|\tilde{G}\|_1^2 + \|\tilde{v}\|_2^2 \|\tilde{\theta}\|_2^2 + \|\tilde{v}\|_2^2 \|\eta\|_2^2 \right].
\end{aligned} \tag{2.61}$$

Thanks to Sobolev's and Young's inequalities, we take the sum of (2.59) and (2.61) to deduce the estimate (2.58). \square

Step 2. Now, it suffices to bound $\|D_{33}^2 (a_{ij} D_i \hat{v}^j)\|_0$ in order to close the estimate for div_v . We apply D_3 to (2.56) to find that

$$-(\mu + \zeta) D_{33}^2 (a_{lj} D_l \hat{v}^j) + \frac{1}{\epsilon} D_{33}^2 (\hat{\eta} + \hat{\theta})$$

$$\begin{aligned}
&= D_3(a_{3i})(-\hat{U}^j a_{kj} D_k \hat{v}^i + \epsilon \hat{F}^i - \hat{v}^j a_{kj} D_k \hat{v}^i) - a_{3i} D_3(\hat{U}^j a_{kj} D_k \hat{v}^i - \epsilon \hat{F}^i + \hat{v}^j a_{kj} D_k \hat{v}^i) \\
&\quad - D_3(\hat{\theta} D_3 \hat{\eta} + \hat{\eta} D_3 \hat{\theta}) + O(1)(D_{33\tau}^3 \hat{v}^j + D_{3l}^2 \hat{v}^j + D_l \hat{v}^j). \tag{2.62}
\end{aligned}$$

Now, multiplying the above equality (2.62) by $-J\chi^2 D_{33}^2(a_{lj} D_l \hat{v}^j)$ in $L^2(\tilde{\Omega})$, one infers that

$$\begin{aligned}
&\frac{\mu + \zeta}{2} \int_{\tilde{\Omega}} J\chi^2 |D_{33}^2(a_{lj} D_l \hat{v}^j)|^2 dy - \frac{1}{\epsilon} \int_{\tilde{\Omega}} J\chi^2 D_{33}^2(a_{lj} D_l \hat{v}^j) \cdot D_{33}^2(\hat{\eta} + \hat{\theta}) dy \\
&\leq C(\|U\|_2^2 \|v\|_2^2 + \epsilon^2 \|\tilde{F}\|_1^2 + \|\tilde{v}\|_2^4 + \|\tilde{\theta}\|_2^2 \|\tilde{\eta}\|_2^2) + \|v\|_2^2 + \|v\|_1^2 + \int_{\tilde{\Omega}} J\chi^2 |D_{33\tau}^3 \hat{v}|^2 dy. \tag{2.63}
\end{aligned}$$

Correspondingly, applying D_{33}^2 to (2.49)₁ and (2.49)₃ and multiplying the resulting equations by $J\chi^2 D_{33}^2 \hat{\eta}$ and $J\chi^2 D_{33}^2 \hat{\theta}$ respectively, we get

$$\begin{aligned}
&\frac{\kappa}{2} \int_{\tilde{\Omega}} J\chi^2 |D_{33}^2(a_{kj} D_k \hat{\theta})|^2 dy + \frac{1}{\epsilon} \int_{\tilde{\Omega}} J\chi^2 D_{33}^2(a_{lj} D_l \hat{v}^j) \cdot D_{33}^2(\hat{\eta} + \hat{\theta}) dy \\
&\leq C\|\theta\|_2^2(1 + \|U\|_3) + \delta\|\theta\|_3^2 + \|U\|_3 \|\eta\|_2^2 + \|\tilde{v}\|_3 \|\eta\|_2^2 + \epsilon\|P\|_3(\|U\|_3 + \|\tilde{v}\|_3) \|\eta\|_2 \\
&\quad + \epsilon^2 \|\tilde{G}\|_1^2 + \|\tilde{v}\|_2^2 \|\tilde{\theta}\|_2^2 + \|\tilde{v}\|_2^2 \|\eta\|_2^2. \tag{2.64}
\end{aligned}$$

Combining (2.63) with (2.64), we see that there are a constant C_{11} and a small δ , such that

$$\begin{aligned}
&\frac{\mu + \zeta}{2} \int_{\tilde{\Omega}} J\chi^2 |D_{33}^2(a_{lj} D_l \hat{v}^j)|^2 dy + \frac{\kappa}{2} \int_{\tilde{\Omega}} J\chi^2 |D_{33}^2(a_{kj} D_k \hat{\theta})|^2 dy \\
&\leq \delta(\|\theta\|_3^2 + \|\mathbf{v}\|_3^2) + C_{11} \left\{ (\|U\|_4^2 \|\mathbf{v}\|_1^2 + \epsilon^2 \|\tilde{F}\|_1^2 + \|\tilde{v}\|_2^4 + \|\tilde{\theta}\|_2^2 \|\tilde{\eta}\|_2^2) \right. \\
&\quad + \left[\|\mathbf{v}\|_1^2 + \int_{\tilde{\Omega}} J\chi^2 |D_{33\tau}^3 \hat{v}|^2 dy + \|\theta\|_1^2(1 + \|U\|_3) + \|U\|_3 \|\eta\|_2^2 + \|\tilde{v}\|_3 \|\eta\|_2^2 \right. \\
&\quad \left. \left. + \epsilon\|P\|_3(\|U\|_3 + \|\tilde{v}\|_3) \|\eta\|_2 + \epsilon^2 \|\tilde{G}\|_1^2 + \|\tilde{v}\|_2^2 \|\tilde{\theta}\|_2^2 + \|\tilde{v}\|_2^2 \|\eta\|_2^2 \right] \right\}. \tag{2.65}
\end{aligned}$$

Step 3. To control the term $D_{33\tau}^3 \hat{v}$ on RHS of (2.65), we introduce an auxiliary Stokes problem in the original coordinates in the region near the boundary:

$$\begin{cases} -\mu \Delta_x [(\chi D_\tau \hat{v}) \circ \Lambda^{-1}] + \frac{1}{\epsilon} \nabla_x [(\chi D_\tau (\hat{\eta} + \hat{\theta})) \circ \Lambda^{-1}] = G_1 & \text{in } W \cap \Omega, \\ \operatorname{div}_x [(\chi D_\tau \hat{v}) \circ \Lambda^{-1}] = G_2 & \text{in } W \cap \Omega, \\ (\chi D_\tau \hat{v}) \circ \Lambda^{-1} = 0 & \text{in } W \cap \Omega, \end{cases}$$

where

$$\begin{aligned}
G_1^i &= \chi D_\tau \left[\zeta a_{ki} D_k (a_{lj} D_l \hat{v}^j) + \epsilon \hat{F}^i - \hat{v}^j a_{kj} D_k \hat{v}^i - \hat{\theta} a_{ki} D_k \hat{\eta} - \hat{\eta} a_{ki} D_k \hat{\theta} \right. \\
&\quad \left. - \hat{U}^j a_{kj} D_k \hat{v}^i \right] + o(1) \left[D_l \hat{v}^i + D_{kl}^2 \hat{v}^i + \frac{1}{\epsilon} D_k (\hat{\eta} + \hat{\theta}) \right]
\end{aligned}$$

and

$$G_2^i = o(1)(D_\tau \hat{v}^j + D_k \hat{v}^j + D_{\tau k}^2 \hat{v}^j),$$

which can be bounded as follows.

$$\begin{aligned}
\|G_1\|_{L^2}^2 &\leq C \left(\left\| \frac{\nabla(\eta + \theta)}{\epsilon} \right\|_{L^2}^2 + \int_{\tilde{\Omega}} J\chi^2 |D_{\tau k}^2(a_{lj} D_l \hat{v}^j)|^2 dy \right) + \delta \|\mathbf{v}\|_3^2 \\
&\quad + C_\delta \|\mathbf{v}\|_1^2 + C(\epsilon^2 \|\tilde{F}\|_1^2 + \|\tilde{v}\|_2^4 + \|\tilde{\theta}\|_2^2 \|\tilde{\eta}\|_2^2 + \|U\|_3^4 \|\mathbf{v}\|_1^2) \tag{2.66}
\end{aligned}$$

and

$$\|G_2\|_{H^1}^2 \leq \delta \|\mathbf{v}\|_3^2 + C_\delta \|\mathbf{v}\|_1^2 + C \int_{\tilde{\Omega}} J\chi^2 |D_{\tau k}(a_{lj} D_l \hat{v}^j)|^2 dy. \tag{2.67}$$

Due to the regularity theory of the Stokes problem (see [14]), one has

$$\int_{W \cap \Omega} |\Delta_x(\chi D_\tau \hat{\mathbf{v}}) \circ \Lambda^{-1}(x)|^2 dx \leq C(\|G_1\|_{L^2(W \cap \Omega)}^2 + \|G_2\|_{H^1(W \cap \Omega)}^2), \quad (2.68)$$

where the left-hand side of (2.68) is equal to

$$\begin{aligned} \int_{W \cap \Omega} |\Delta_x(\chi D_\tau \hat{\mathbf{v}}) \circ \Lambda^{-1}(x)|^2 dx &= \int_{\tilde{\Omega}} J \left| \sum_{j=1}^3 \sum_{k=1}^3 a_{kj} D_k \left(\sum_{l=1}^3 a_{lj} D_l(\chi D_\tau \hat{\mathbf{v}}) \right) \right|^2 dy \\ &= \int_{\tilde{\Omega}} J \chi^2 \left| \sum_{j,k,l=1}^3 a_{kj} a_{lj} D_{kl\tau}^3 \hat{\mathbf{v}} \right|^2 dy + o(1) \int_{\tilde{\Omega}} (|D_\tau \hat{\mathbf{v}}|^2 + |D_{y\tau}^2 \hat{\mathbf{v}}|^2) dy. \end{aligned}$$

And we use (2.51) to get

$$D_{33\tau}^3 \hat{\mathbf{v}} = \sum_{j,k,l=1}^3 a_{kj} a_{lj} D_{kl\tau}^3 \hat{\mathbf{v}} - \sum_{1 \leq k, l \leq 2} \sum_{j=1}^3 a_{kj} a_{lj} D_{kl\tau}^3 \hat{\mathbf{v}},$$

from which, (2.66) and (2.67) it follows that the inequality (2.68) gives

$$\begin{aligned} (C_{11} + 1) \int_{\tilde{\Omega}} J \chi^2 |D_{33\tau}^3 \hat{\mathbf{v}}|^2 dy &\leq C(\|G_1\|_{L^2(W \cap \Omega)}^2 + \|G_2\|_{H^1(W \cap \Omega)}^2) + C \int_{\tilde{\Omega}} J \chi^2 |D_{\tau\xi\zeta}^3 \hat{\mathbf{v}}|^2 dy + C_\delta \|\nabla \mathbf{v}\|_{L^2}^2 + \delta \|\mathbf{v}\|_3^2 \\ &\leq \delta \|\mathbf{v}\|_3^2 + C_{12} \left[\|\mathbf{v}\|_1^2 + \left\| \frac{\nabla(\eta + \theta)}{\epsilon} \right\|_0^2 + \int_{\tilde{\Omega}} J \chi^2 (|D_{\xi\tau y}^3 \hat{\mathbf{v}}|^2 + |D_{3\tau}^2 (a_{lj} D_l \hat{\mathbf{v}}^j)|^2) dy \right. \\ &\quad \left. + (\epsilon^2 \|\tilde{F}\|_1^2 + \|\tilde{\mathbf{v}}\|_2^4 + \|\tilde{\theta}\|_2^2 \|\eta\|_2^2 + \|U\|_3^4 \|\mathbf{v}\|_1^2) \right]. \end{aligned} \quad (2.69)$$

Now, letting

$$\Phi_\chi := \int_{\tilde{\Omega}} J \chi^2 (|D_{\tau\xi y}^3 \hat{\mathbf{v}}|^2 + |D_{\tau 3}^2 (a_{lj} D_l \hat{\mathbf{v}}^j)|^2 + |D_{33}^2 (a_{lj} D_l \hat{\mathbf{v}}^j)|^2 + |D_{33\tau}^3 \hat{\mathbf{v}}|^2) dy$$

and

$$\Psi_\chi := \int_{\tilde{\Omega}} J \chi^2 (a_{kj} |D_{k\tau\xi}^3 \hat{\theta}|^2 + |D_{\tau 3}^2 (a_{kj} D_k \hat{\theta})|^2 + |D_{33}^2 (a_{kj} D_k \hat{\theta})|^2) dy,$$

we can apply Cauchy-Schwarz's and Young's inequalities as well as the estimate (2.38) to deduce from (2.55), (2.58), (2.65) and (2.69) that

$$\begin{aligned} \Phi_\chi + \Psi_\chi &\leq \|U\|_3 \|\eta\|_2^2 + (\|\tilde{\mathbf{v}}\|_3 + \|\tilde{\mathbf{v}}\|_2^2) \|\eta\|_2^2 + \epsilon \|P\|_3 (\|U\|_3 + \|\tilde{\mathbf{v}}\|_3) \|\eta\|_2 \\ &\quad + C(1 + \|U\|_3^4) (\|\mathbf{v}\|_1^2 + \|\theta\|_1^2) + \delta \left(\|\mathbf{v}\|_3^2 + \|\theta\|_3^2 + \left\| \frac{\nabla(\eta + \theta)}{\epsilon} \right\|_1^2 \right) \\ &\quad + C_\delta \left[\epsilon^2 (\|\tilde{F}\|_1^2 + \|\tilde{G}\|_1^2) + \|\tilde{\mathbf{v}}\|_2^4 + (\|\tilde{\theta}\|_2^2 + \|\tilde{\mathbf{v}}\|_2^2) \|\eta\|_2^2 + \|\tilde{\mathbf{v}}\|_2^2 \|\tilde{\theta}\|_2^2 \right], \end{aligned} \quad (2.70)$$

which, together with (2.39)–(2.42), results in

$$\begin{aligned} &\|\mathbf{v}\|_3^2 + \|\theta\|_3^2 + \left\| \frac{\nabla \eta + \nabla \theta}{\epsilon} \right\|_1^2 \\ &\leq C_{12} \left\{ \|U\|_3 \|\eta\|_2^2 + (1 + \|U\|_3^4) \left[\epsilon \|P\|_3 (\|U\|_3 + \|\tilde{\mathbf{v}}\|_3) + (\|\tilde{\theta}\|_2^2 + \|\tilde{\mathbf{v}}\|_2^2 + \|\tilde{\mathbf{v}}\|_3) \right] \|\eta\|_2^2 \right. \\ &\quad \left. + (1 + \|U\|_3^4) (\|\mathbf{v}\|_1^2 + \|\theta\|_1^2) + (1 + \|U\|_3^4) \left[\epsilon^2 (\|\tilde{F}\|_1^2 + \|\tilde{G}\|_1^2) + \|\tilde{\mathbf{v}}\|_2^4 + \|\tilde{\mathbf{v}}\|_2^2 \|\tilde{\theta}\|_2^2 \right] \right\}. \end{aligned} \quad (2.71)$$

2.2.4 Boundedness of η

In the next lemma, we derive upper bounds of $\|\eta\|_1$ and $\|\eta\|_2$.

Lemma 2.8. *There are a small $\delta > 0$, and two positive constants C_{13} and C_{14} independent of ϵ , such that*

$$\begin{aligned} \|\eta\|_1 \leq & C_{13} \left[\epsilon^2 \|\tilde{F}\|_0 + \epsilon (\|\tilde{\mathbf{v}}\|_2 \|\tilde{\mathbf{v}}\|_1 + \|\tilde{\theta}\|_2 \|\eta\|_2 + \|U\|_2 \|\mathbf{v}\|_2 + \|\operatorname{div} \mathbf{v}\|_1 + (\|\tilde{F}\|_{-1} + \|\tilde{G}\|_{-1})) \right. \\ & \left. + \epsilon \|P\|_2 (\|U\|_2 + \|\tilde{\mathbf{v}}\|_2) + \|\tilde{\mathbf{v}}\|_3^{\frac{1}{2}} \|\eta\|_0 + \|\tilde{\mathbf{v}}\|_2^2 + \|\eta\|_1 \|\tilde{\theta}\|_1 + \|\tilde{\mathbf{v}}\|_1 (\|\tilde{\theta}\|_1 + \|\eta\|_1) \right] \end{aligned} \quad (2.72)$$

and

$$\begin{aligned} \|\eta\|_2 \leq & C_{14} (1 + \epsilon) (1 + \|U\|_2^2) \left\{ \epsilon (\|\tilde{F}\|_1 + \|\tilde{G}\|_{-1}) + \|\tilde{\mathbf{v}}\|_3^2 + \|\tilde{\theta}\|_3 \|\eta\|_2 + \|\tilde{\mathbf{v}}\|_3^{\frac{1}{2}} \|\eta\|_0 \right. \\ & \left. + \left[\epsilon \|P\|_2 (\|U\|_2 + \|\tilde{\mathbf{v}}\|_2) \|\eta\|_0 \right]^{\frac{1}{2}} + \|\tilde{\mathbf{v}}\|_1 (\|\tilde{\theta}\|_1 + \|\eta\|_1) \right\} + \epsilon \|\mathbf{v}\|_3 + \delta \|\theta\|_3. \end{aligned} \quad (2.73)$$

Proof. From (2.38) we get

$$\epsilon \|\mathbf{v}\|_2 + \|\nabla \eta + \nabla \theta\|_0 \leq C \left[\epsilon^2 \|\tilde{F}\|_0 + \epsilon (\|\tilde{\mathbf{v}}\|_2 \|\tilde{\mathbf{v}}\|_1 + \|\tilde{\theta}\|_2 \|\eta\|_2 + \|U\|_2 \|\mathbf{v}\|_2 + \|\operatorname{div} \mathbf{v}\|_1) \right],$$

which together with Lemma 2.4 gives

$$\begin{aligned} \|\nabla \eta\|_1 & \leq \|\nabla \eta + \nabla \theta\|_0 + \|\nabla \theta\|_0 \\ & \leq C \left[\epsilon^2 \|\tilde{F}\|_0 + \epsilon (\|\tilde{\mathbf{v}}\|_2 \|\tilde{\mathbf{v}}\|_1 + \|\tilde{\theta}\|_2 \|\eta\|_2 + \|U\|_2 \|\mathbf{v}\|_2 + \|\operatorname{div} \mathbf{v}\|_1) \right] \\ & \quad + C \left[\|\tilde{\mathbf{v}}\|_3 \|\eta\|_0^2 + \epsilon^2 (\|\tilde{F}\|_{-1}^2 + \|\tilde{G}\|_{-1}^2) + \epsilon \|P\|_2 (\|U\|_2 + \|\tilde{\mathbf{v}}\|_2) \|\eta\|_0 \right. \\ & \quad \left. + \|\tilde{\mathbf{v}}\|_1^4 + \|\eta\|_1^2 \|\tilde{\theta}\|_1^2 + \|\tilde{\mathbf{v}}\|_1^2 (\|\tilde{\theta}\|_1^2 + \|\eta\|_1^2) \right]^{1/2}. \end{aligned}$$

If we apply Poincaré's and Young's inequalities to the above inequality, and use the fact that

$$(A_1 + A_2 + \dots + A_n)^{1/2} \leq A_1^{1/2} + A_2^{1/2} + \dots + A_n^{1/2} \quad \text{for } A_i \geq 0 \quad (i = 1, \dots, n), \quad (2.74)$$

we obtain the estimate (2.72) immediately.

On the other hand, from the estimate (2.39) we conclude that

$$\begin{aligned} \|\nabla \eta\|_1 & \leq \|\nabla \eta + \nabla \theta\|_1 + \|\nabla \theta\|_1 \\ & \leq C \epsilon (1 + \|U\|_2^2) \left\{ \epsilon (\|\tilde{F}\|_1 + \|\tilde{G}\|_{-1}) + \|\tilde{\mathbf{v}}\|_3^2 + \|\tilde{\theta}\|_3 \|\eta\|_2 + \|\tilde{\mathbf{v}}\|_3^{\frac{1}{2}} \|\eta\|_0 \right. \\ & \quad \left. + \left[\epsilon \|P\|_2 (\|U\|_2 + \|\tilde{\mathbf{v}}\|_2) \|\eta\|_0 \right]^{\frac{1}{2}} + \|\tilde{\mathbf{v}}\|_1 (\|\tilde{\theta}\|_1 + \|\eta\|_1) \right\} + \epsilon \|\mathbf{v}\|_3 + \delta \|\theta\|_3 + C_\delta \|\theta\|_1, \end{aligned}$$

which, together with Poincaré's inequality, (2.72) and (2.74), implies (2.73). \square

3 Existence of the nonlinear problem

In this section, we give the proof of the existence for the nonlinear problem (1.5) by using the Tikhonov theorem which can be found in [27]. For completeness, we state the theorem in the following.

Theorem 3.1. *(Tikhonov Theorem, [27, P72, 1.2.11.6]) Let M be a nonempty bounded closed convex subset of a separable reflexive Banach space X and let $F : M \rightarrow M$ be a weakly continuous mapping (i.e., if $x_n \in M$, $x_n \rightharpoonup x$ weakly in X , then $F(x_n) \rightharpoonup F(x)$ weakly in X as well). Then F has at least one fixed point in M .*

Define a Banach space X by

$$X = \bar{H}^1 \times H_0^1 \times H_0^1,$$

which can be easily verified to be separable and reflexive.

A convex subset $K_1(E)$ of X is defined by

$$K_1(E) = \{(\mathbf{v}, \theta) \in (H^3 \cap H_0^1) \times (H^3 \cap H_0^1) \mid \|\mathbf{v}\|_3 + \|\theta\|_3 \leq E\},$$

where $E < 1$ is a small positive constant. By the lower semi-continuity of norms, we easily see that the subset $K_1(E)$ is also closed in X .

We define a space K by

$$K = K_0 \times K_1(E),$$

where K_0 is defined by (2.9). Note that K is a nonempty bounded closed convex subset of X .

Now, we define a nonlinear operator N from K to X by

$$N(\tilde{U}, \tilde{\mathbf{v}}, \tilde{\theta}) := (U, \mathbf{v}, \theta),$$

where U and (\mathbf{v}, θ) are the solutions of (2.3) and (2.10) for given $(\tilde{U}, \tilde{\mathbf{v}}, \tilde{\theta})$, respectively.

Next, we want to find a fixed point (U, \mathbf{v}, θ) of N in K , such that $(U, \mathbf{v}, \theta) = N(U, \mathbf{v}, \theta)$, which, together with the existence of weak solutions in Lemmas 2.1 and 2.4, gives that (U, P) and $(\eta, \mathbf{v}, \theta)$ are solutions of the boundary value problems (2.1) and (2.2), respectively. So $(U + \mathbf{v}, \epsilon P + \eta, \theta)$ will be a solution to (1.5). For this purpose, we have to show that N maps K into itself and $N : K \rightarrow K$ is a weakly continuous mapping.

Lemma 3.1. *There is a small constant $\epsilon_0 > 0$, depending only on $\Omega, \mu, \lambda, \mathbf{f}$ and \mathbf{g} , such that for any $\epsilon \in (0, \epsilon_0)$, K is a nonempty bounded closed convex subset of X and $N(K) \subset K$.*

Proof. By virtue of the definition, it is obvious that $K \subset X$ is a nonempty, bounded, closed convex set. Now, we will show that the operator N maps K into itself, i.e., $N(K) \subset K$. To this end, let $(\tilde{U}, \tilde{\mathbf{v}}, \tilde{\theta}) \in K$ and $(U, \mathbf{v}, \theta) = N(\tilde{U}, \tilde{\mathbf{v}}, \tilde{\theta})$. By Lemmas 2.1 and 2.2, we see that $U \in K_0$ for all $\tilde{U} \in K_0$. Thus, it suffices to check that $(\mathbf{v}, \theta) \in K_1(E)$ for $(\tilde{\mathbf{v}}, \tilde{\theta}) \in K_1(E)$. By (2.7) and (2.8), we have

$$\|U\|_3 + \|\nabla P\|_1 \leq M_1 \quad \text{and} \quad \|U\|_4 + \|\nabla P\|_2 \leq M_2, \quad (3.1)$$

where $M_1 = C_3 \|\mathbf{h}\|_1 (\|\mathbf{h}\|_1 + 1)^8$ and $M_2 = C_4 \|\mathbf{h}\|_2 (\|\mathbf{h}\|_2 + 1)^{12}$.

On the other hand, recalling the definition of \tilde{F} and \tilde{G} , we get from (3.1) that

$$\begin{aligned} \|\tilde{F}\|_1 &= \|(\epsilon P + \eta)\mathbf{f} - (\epsilon P + \eta)(U + \tilde{\mathbf{v}}) \cdot \nabla(U + \tilde{\mathbf{v}}) - \tilde{\theta} \nabla P - P \nabla \tilde{\theta}\|_1 \\ &\leq C [\|\eta\|_2 (\|\mathbf{f}\|_1 + \|U\|_2^2 + \|\tilde{\mathbf{v}}\|_2^2) + \epsilon \|P\|_2 (\|\mathbf{f}\|_1 + \|U\|_2^2 + \|\tilde{\mathbf{v}}\|_2^2 + \|\tilde{\theta}\|_2)] \\ &\leq C \|\eta\|_2 (\|\mathbf{f}\|_1 + (M_1 + 1)^2) + \epsilon C M_1 (\|\mathbf{f}\|_1 + (M_1 + 1)^2), \end{aligned} \quad (3.2)$$

$$\begin{aligned} \|\tilde{G}\|_1 &= \|\tilde{\Psi} - (\epsilon P + \eta)(U + \tilde{\mathbf{v}}) \cdot \nabla \tilde{\theta} + (\epsilon P + \eta) \tilde{\theta} \operatorname{div} \tilde{\mathbf{v}} + P \operatorname{div} \tilde{\mathbf{v}}\|_1 \\ &\leq C [\epsilon (\|U\|_2^2 + \|\tilde{\mathbf{v}}\|_2^2) + (\|\eta\|_2 + \epsilon \|P\|_2) (\|U\|_2 + \|\tilde{\mathbf{v}}\|_2) \|\tilde{\theta}\|_2 + \|P\|_2 \|\tilde{\mathbf{v}}\|_2] \\ &\leq C \|\eta\|_2 (M_1 + 1) + C (\epsilon (M_1 + 1)^2 + (M_1 + 1)). \end{aligned} \quad (3.3)$$

As a result of Poincaré's inequality and Lemma 2.4, we have

$$\begin{aligned} \|\mathbf{v}\|_1 + \|\theta\|_1 &\leq C \left\{ (E^{\frac{1}{2}} + E) \|\eta\|_2 + \epsilon \left[\|\eta\|_2 (M_1 + 1) + C (\|\eta\|_2 (\|\mathbf{f}\|_1 + (M_1 + 1)^2) \right. \right. \\ &\quad \left. \left. + \epsilon C M_1 (\|\mathbf{f}\|_1 + (M_1 + 1)^2) + \epsilon (M_1 + 1)^2 + (M_1 + 1) \right] \right. \\ &\quad \left. + \epsilon C_\delta (M_1 + 1)^2 + E^2 \right\} + \delta \|\eta\|_2 \end{aligned}$$

$$\begin{aligned} &\leq C[E^{\frac{1}{2}} + \epsilon(\|\mathbf{f}\|_1 + (M_1 + 1)^2)]\|\eta\|_2 + \epsilon^2 CM_1(\|\mathbf{f}\|_1 + (M_1 + 1)^2) + CE^2 \\ &\quad + \delta\|\eta\|_2 + \epsilon C_\delta(M_1 + 1)^2, \end{aligned} \quad (3.4)$$

where we have used the estimates (3.1), (3.2) and (3.3).

On the other hand, in view of Poincaré's and Young's inequalities, (3.4) and (2.73) in Lemma 2.8, we find that

$$\begin{aligned} \|\eta\|_2 &\leq C(1 + \epsilon)(1 + M_1^2) \left\{ \epsilon \left[\|\eta\|_2(\|\mathbf{f}\|_1 + (M_1 + 1)^2) + M_1(\|\mathbf{f}\|_1 + (M_1 + 1)^2) \right. \right. \\ &\quad \left. \left. + \|\eta\|_2(M_1 + 1) + (\epsilon(M_1 + 1)^2 + (M_1 + 1)) \right] + E^2 + (E + E^{\frac{1}{2}})\|\eta\|_2 \right. \\ &\quad \left. + \epsilon^{\frac{1}{2}} M_1^{\frac{1}{2}} (M_1 + E)^{\frac{1}{2}} \|\eta\|_2^{\frac{1}{2}} + E(E + \|\eta\|_2) \right\} + \epsilon\|\mathbf{v}\|_3 + \delta\|\theta\|_3 \\ &\leq C_{15}(1 + \epsilon)(1 + M_1^2) \left\{ \epsilon \left[\|\mathbf{f}\|_1 + (M_1 + 1)^2 + (M_1 + 1) \right] + (E + E^{\frac{1}{2}}) + \delta \right\} \|\eta\|_2 \\ &\quad + \epsilon\|\mathbf{v}\|_3 + \delta\|\theta\|_3 + C_{15}(1 + \epsilon)(1 + M_1^2) \left\{ \epsilon(M_1 + 1)^2 \right. \\ &\quad \left. + \epsilon \left[M_1(\|\mathbf{f}\|_1 + (M_1 + 1)^2) + (\epsilon(M_1 + 1)^2 + (M_1 + 1)) \right] + E^2 \right\}, \end{aligned} \quad (3.5)$$

where C_{15} is a positive constant.

Combining (2.71) with (3.2)–(3.5), we conclude that there is a constant C_{16} , such that

$$\begin{aligned} \|\mathbf{v}\|_3 + \|\theta\|_3 + \|\eta\|_2 &\leq C_{16}(1 + M_1)^5 \left\{ \epsilon \left[\|\mathbf{f}\|_1 + (M_1 + 1)^2 \right] + (E + E^{\frac{1}{2}}) + \delta \right\} \|\eta\|_2 \\ &\quad + \epsilon M_1^{\frac{1}{2}} \|\mathbf{v}\|_3 + \delta M_1^{\frac{1}{2}} \|\theta\|_3 + C_{16}(1 + M_1)^5 \left\{ \epsilon(M_1 + 1)^2 \right. \\ &\quad \left. + \epsilon(1 + M_1) \left[\|\mathbf{f}\|_1 + (M_1 + 1)^2 \right] + E^2 \right\}. \end{aligned} \quad (3.6)$$

Thus, first taking δ small enough and then choosing ϵ_0 and E suitably small, such that

$$\begin{aligned} C_{16}(1 + M_1)^5 \left\{ \epsilon_0 \left[\|\mathbf{f}\|_1 + (M_1 + 1)^2 \right] + (E + E^{\frac{1}{2}}) \right\} &< 1, \quad \epsilon_0 M_1^{\frac{1}{2}} < 1, \\ C_{16}(1 + M_1)^5 \left\{ \epsilon_0(M_1 + 1)^2 + \epsilon_0(1 + M_1) \left[\|\mathbf{f}\|_1 + (M_1 + 1)^2 \right] + E^2 \right\} &< E, \end{aligned}$$

we deduce from (3.6) that for all $\epsilon \in (0, \epsilon_0)$,

$$\|\mathbf{v}\|_3 + \|\theta\|_3 + \|\eta\|_2 \leq E,$$

which gives $\|\mathbf{v}\|_3 + \|\theta\|_3 \leq E$ immediately. This completes the proof. \square

Lemma 3.2. *Let N, X, K_0 and $K_1(E)$ be the same as in Lemma 3.1. Then $N : K \rightarrow K$ is a weakly continuous mapping.*

Proof. By the definition of weakly continuous mapping (see, for example, [27, P72,1.4.11.6]), it suffices to prove that N is continuous on K in the norm of X .

Let $(U_i, \mathbf{v}_i, \theta_i) = N(\tilde{U}_i, \tilde{\mathbf{v}}_i, \tilde{\theta}_i)$, $i = 1, 2$. In particular, let $(U_i, P_i) \in (H^4 \cap H_{0,\sigma}^1) \times \bar{H}^3$ and $(\eta_i, \mathbf{v}_i, \theta_i) \in \bar{H}^2 \times (H^3 \cap H_0^1) \times (H^3 \cap H_0^1)$ be the solutions of (2.3) and (2.10) for given $(\tilde{U}_i, \tilde{\mathbf{v}}_i, \tilde{\theta}_i)$ respectively, i.e.,

$$\begin{cases} (\tilde{U}_i + \tilde{\mathbf{v}}_i) \cdot \nabla U_i - \mu \Delta U_i + \nabla P_i = \mathbf{h}, & \int P_i dx = 0, \\ \operatorname{div} U_i = 0; \end{cases} \quad (3.7)$$

and

$$\begin{cases} U_i \cdot \nabla \eta_i + \frac{\operatorname{div} \mathbf{v}_i}{\epsilon} = -\tilde{\mathbf{v}}_i \cdot \nabla \eta_i - \eta_i \operatorname{div} \tilde{\mathbf{v}}_i - \epsilon \operatorname{div}(P_i(U_i + \tilde{\mathbf{v}}_i)), \\ U_i \cdot \nabla \mathbf{v}_i - \mu \Delta \mathbf{v}_i - \zeta \nabla \operatorname{div} \mathbf{v}_i + \frac{\nabla \eta_i + \nabla \theta_i}{\epsilon} = \epsilon \tilde{F}_i - \tilde{\mathbf{v}}_i \cdot \nabla \tilde{\mathbf{v}}_i - \tilde{\theta}_i \nabla \eta_i - \eta_i \nabla \tilde{\theta}_i, \\ U_i \cdot \nabla \theta_i - \kappa \Delta \theta_i + \frac{\operatorname{div} \mathbf{v}_i}{\epsilon} = \epsilon \tilde{G}_i - \tilde{\mathbf{v}}_i \cdot \nabla \tilde{\theta}_i - \eta_i \operatorname{div} \tilde{\mathbf{v}}_i - \tilde{\theta}_i \operatorname{div} \tilde{\mathbf{v}}_i, \end{cases} \quad (3.8)$$

where the force \tilde{F}_i and heat source \tilde{G}_i are given by

$$\begin{aligned} \tilde{F}_i &= (\epsilon P_i + \eta_i) \mathbf{f} - (\epsilon P_i + \eta_i)(U_i + \tilde{\mathbf{v}}_i) \cdot \nabla(U_i + \tilde{\mathbf{v}}_i) - \tilde{\theta}_i \nabla P_i - P_i \nabla \tilde{\theta}_i, \\ \tilde{G}_i &= \tilde{\Psi}_i - (\epsilon P_i + \eta_i)(U_i + \tilde{\mathbf{v}}_i) \cdot \nabla \tilde{\theta}_i + (\epsilon P_i + \eta_i) \tilde{\theta}_i \operatorname{div} \tilde{\mathbf{v}}_i + P_i \operatorname{div} \tilde{\mathbf{v}}_i. \end{aligned}$$

Now, if we set

$$\begin{aligned} W &= U_2 - U_1, \quad \tilde{W} = \tilde{U}_2 - \tilde{U}_1, \quad Q = P_2 - P_1, \quad \xi = \eta_2 - \eta_1, \\ \mathbf{w} &= \mathbf{v}_2 - \mathbf{v}_1, \quad \tilde{\mathbf{w}} = \tilde{\mathbf{v}}_2 - \tilde{\mathbf{v}}_1, \quad \beta = \theta_2 - \theta_1, \quad \tilde{\beta} = \tilde{\theta}_2 - \tilde{\theta}_1, \\ J &= \tilde{F}_2 - \tilde{F}_1, \quad I = \tilde{G}_2 - \tilde{G}_1, \end{aligned}$$

then, we can have the following systems:

$$\begin{cases} (\tilde{U}_1 + \tilde{v}_1) \cdot \nabla W - \mu \Delta W + \nabla Q = -(\tilde{W} + \tilde{\mathbf{w}}) \cdot \nabla U_2, & \int Q dx = 0, \\ \operatorname{div} W = 0, \end{cases} \quad (3.9)$$

and

$$\begin{cases} U_1 \cdot \nabla \xi + \frac{\operatorname{div} \mathbf{w}}{\epsilon} = -\operatorname{div}(\tilde{\mathbf{v}}_1 \xi + \tilde{\mathbf{w}} \eta_2) - \epsilon \operatorname{div}(P_1(W + \tilde{\mathbf{w}}) + Q(U_2 + \tilde{\mathbf{v}}_2)) - W \cdot \nabla \eta_2, \\ U_1 \cdot \nabla \mathbf{w} - \mu \Delta \mathbf{w} - \zeta \nabla \operatorname{div} \mathbf{w} + \frac{\nabla \xi + \nabla \beta}{\epsilon} \\ \quad = \epsilon J - \tilde{\mathbf{w}} \cdot \nabla \tilde{\mathbf{v}}_2 - \tilde{\mathbf{v}}_1 \cdot \nabla \tilde{\mathbf{w}} - W \cdot \nabla \mathbf{v}_2 - \nabla(\tilde{\theta}_1 \xi + \tilde{\beta} \eta_2), \\ U_1 \cdot \nabla \beta - \kappa \Delta \beta + \frac{\operatorname{div} \mathbf{w}}{\epsilon} \\ \quad = \epsilon I - \tilde{\mathbf{w}} \cdot \nabla \tilde{\theta}_2 - \tilde{\mathbf{v}}_1 \cdot \nabla \tilde{\beta} - W \cdot \nabla \theta_2 - \xi \operatorname{div} \tilde{\mathbf{v}}_1 - \eta_2 \operatorname{div} \tilde{\mathbf{w}} - \tilde{\beta} \operatorname{div} \tilde{\mathbf{v}}_1 - \tilde{\theta}_2 \operatorname{div} \tilde{\mathbf{w}}, \end{cases} \quad (3.10)$$

where J and I read as

$$\begin{aligned} J &= (\epsilon Q + \xi) \mathbf{f} - (\epsilon Q + \xi)(U_2 + \tilde{\mathbf{v}}_2) \cdot \nabla(U_2 + \tilde{\mathbf{v}}_2) \\ &\quad - (\epsilon P_1 + \eta_1)((W + \tilde{\mathbf{w}}) \cdot \nabla(U_2 + \tilde{\mathbf{v}}_2) + (U_1 + \tilde{\mathbf{v}}_1) \cdot \nabla(W + \tilde{\mathbf{w}})) - \nabla(\tilde{\beta} P_1 + Q \tilde{\theta}_2), \\ I &= 2\mu D(W + \tilde{\mathbf{w}}) : D(U_2 + \tilde{\mathbf{v}}_2) + 2\mu D(U_1 + \tilde{\mathbf{v}}_1) : D(W + \tilde{\mathbf{w}}) \\ &\quad + \lambda \operatorname{div}(W + \tilde{\mathbf{w}}) \cdot \operatorname{div}(U_2 + \tilde{\mathbf{v}}_2) + \lambda \operatorname{div}(U_1 + \tilde{\mathbf{v}}_1) \cdot \operatorname{div}(W + \tilde{\mathbf{w}}) \\ &\quad - (\epsilon Q + \xi)(U_2 + \tilde{\mathbf{v}}_2) \cdot \nabla \tilde{\theta}_2 - (\epsilon P_1 + \eta_1)((W + \tilde{\mathbf{w}}) \cdot \nabla \tilde{\theta}_2 + (U_1 + \tilde{\mathbf{v}}_1) \cdot \nabla \tilde{\beta}) \\ &\quad + (\epsilon Q + \xi) \tilde{\theta}_2 \operatorname{div} \tilde{\mathbf{v}}_2 - (\epsilon P_1 + \eta_1)(\tilde{\beta} \operatorname{div} \tilde{\mathbf{v}}_2 + \tilde{\theta}_1 \operatorname{div} \tilde{\mathbf{w}}) + Q \operatorname{div} \tilde{\mathbf{v}}_2 + P_1 \operatorname{div} \tilde{\mathbf{w}}. \end{aligned}$$

Note that J and I can be bounded as follows.

$$\begin{aligned} \|J\|_0 &\leq (\epsilon \|Q\|_1 + \|\xi\|_1)(\|\mathbf{f}\|_2 + \|U_2\|_2^2 + \|\tilde{\mathbf{v}}_2\|_2^2) + (\epsilon \|P_1\|_2 + \|\eta_1\|_2)(\|W\|_1 + \|\tilde{\mathbf{w}}\|_1) \\ &\quad \cdot (\|U_2\|_2 + \|\tilde{\mathbf{v}}_2\|_2 + \|U_1\|_2 + \|\tilde{\mathbf{v}}_1\|_2) + \|\tilde{\theta}\|_2 \|Q\|_1 + \|\tilde{\beta}\|_1 \|P_1\|_2 \end{aligned} \quad (3.11)$$

and

$$\begin{aligned}
\|I\|_0 &\leq C\{(\|W\|_1 + \|\tilde{\mathbf{w}}\|_1)(\|U_1\|_2 + \|U_2\|_2 + \|\tilde{\mathbf{v}}_1\|_2 + \|\tilde{\mathbf{v}}_2\|_2) + (\epsilon\|Q\|_1 + \|\xi\|_1)(\|U_2\|_2 + \|\tilde{\mathbf{v}}_2\|_2) \\
&\quad \cdot \|\tilde{\theta}_2\|_2 + (\epsilon\|P_1\|_2 + \|\eta_1\|_2)[(\|W\|_1 + \|\tilde{\mathbf{w}}\|_1)\|\tilde{\theta}_2\|_2 + \|U_1\|_2 + \|\tilde{\mathbf{v}}_1\|_2\|\beta\|_1] \\
&\quad + (\epsilon\|P_1\|_2 + \|\eta_1\|_2)(\|\tilde{\beta}\|_1\|\tilde{\mathbf{v}}_2\|_2 + \|\tilde{\theta}_1\|_2\|\tilde{\mathbf{w}}\|_1) + \|Q\|_1\|\tilde{\mathbf{v}}_2\|_2 + \|P_1\|_2\|\tilde{\mathbf{w}}\|_1\}
\end{aligned} \tag{3.12}$$

On the one hand, we multiply (3.9)₁ by W and make use of Poincaré's inequality to deduce that

$$\left(\mu - \frac{C\|\tilde{\mathbf{v}}\|_3}{2}\right)\|\nabla W\|_0^2 \leq C(\|\tilde{W}\|_1 + \|\tilde{\mathbf{w}}\|_1)\|U_2\|_3\|W\|_1,$$

where C is a positive constant depending only on Ω and μ . Consequently,

$$\|W\|_1 \leq C(\|\tilde{W}\|_1 + \|\tilde{\mathbf{w}}\|_1) \tag{3.13}$$

for some positive constant C depending only on $\Omega, \mu, \lambda, \mathbf{f}, E$ and ϵ_0 .

By the classical estimates for the Stokes equations

$$\begin{cases} -\mu\Delta W + \nabla Q = -(\tilde{W} + \tilde{\mathbf{w}}) \cdot \nabla U_2 - (\tilde{U}_1 + \tilde{\mathbf{v}}_1) \cdot \nabla W, \\ \operatorname{div} W = 0, \end{cases}$$

we obtain that

$$\begin{aligned}
\|W\|_2 + \|\nabla Q\|_0 &\leq C(\|\tilde{W} + \tilde{\mathbf{w}}\|_1\|U_2\|_2 + \|\tilde{U}_1 + \tilde{\mathbf{v}}_1\|_2\|W\|_1) \\
&\leq C(\|\tilde{W}\|_1 + \|\tilde{\mathbf{w}}\|_1),
\end{aligned} \tag{3.14}$$

where the estimate (3.13) has been used.

On the other hand, if we multiply (3.10)₁, (3.10)₂ and (3.10)₃ by ξ, w and β in L^2 respectively, we find that

$$\begin{aligned}
&\mu\|\nabla \mathbf{w}\|_0^2 + \zeta\|\operatorname{div} \mathbf{w}\|_0^2 + \kappa\|\nabla \beta\|_0^2 \\
&= - \int \left[\xi \operatorname{div} \tilde{\mathbf{v}}_1 + \eta_2 \operatorname{div} \tilde{\mathbf{w}} + \tilde{\mathbf{v}}_1 \cdot \nabla \xi + \tilde{\mathbf{w}} \cdot \nabla \eta_2 + \epsilon \operatorname{div}(P_1(W + \tilde{\mathbf{w}}) + Q(U_2 + \tilde{\mathbf{v}}_2)) + W \cdot \nabla \eta_2 \right] \xi dx \\
&\quad + \int \left[\epsilon J - \tilde{\mathbf{w}} \cdot \nabla \tilde{\mathbf{v}}_2 - \tilde{\mathbf{v}}_1 \cdot \nabla \tilde{\mathbf{w}} - W \cdot \nabla \mathbf{v}_2 - \nabla(\tilde{\theta}_1 \xi + \tilde{\beta} \eta_2) \right] \cdot w dx \\
&\quad + \int \left[\epsilon D - \tilde{\mathbf{w}} \cdot \nabla \tilde{\theta}_2 - \tilde{\mathbf{v}}_1 \cdot \nabla \tilde{\beta} - W \cdot \nabla \theta_2 - \xi \operatorname{div} \tilde{\mathbf{v}}_1 - \eta_2 \operatorname{div} \tilde{\mathbf{w}} - \tilde{\beta} \operatorname{div} \tilde{\mathbf{v}}_1 - \tilde{\theta}_1 \operatorname{div} \tilde{\mathbf{w}} \right] \beta dx \\
&\leq C \left\{ \|\xi\|_0^2 \|\tilde{\mathbf{v}}_1\|_3 + \|\xi\|_1 \|\tilde{\mathbf{w}}\|_1 \|\eta_2\|_1 + \epsilon \|\xi\|_0 [\|P_1\|_2(\|W\|_1 + \|\tilde{\mathbf{w}}\|_1) + \|Q\|_1(\|U_2\|_2 + \|\tilde{\mathbf{v}}_2\|_2)] \right. \\
&\quad + \|W\|_1 \|\eta_2\|_3 \|\xi\|_1 \left. \right\} + C \left\{ \epsilon \|J\|_0^2 + \|\tilde{\mathbf{w}}\|_1^2 (\|\tilde{\mathbf{v}}_2\|_2^2 + \|\tilde{\mathbf{v}}_1\|_2^2) + \|W\|_1^2 \|\mathbf{v}_2\|_2^2 + \|\tilde{\theta}_2\|_1^2 \|\xi\|_1^2 \right. \\
&\quad + \|\tilde{\beta}\|_1^2 \|\eta\|_2^2 \left. \right\} + C \left\{ \epsilon \|I\|_0^2 + \|\tilde{\mathbf{w}}\|_1^2 \|\tilde{\theta}_2\|_2^2 + \|\tilde{\mathbf{v}}_1\|_2^2 \|\tilde{\beta}\|_1^2 + \|W\|_1^2 \|\theta_2\|_2^2 + \|\xi\|_1^2 \|\tilde{\mathbf{v}}_1\|_2^2 \right. \\
&\quad \left. + \|\eta_2\|_2^2 \|\tilde{\mathbf{w}}\|_1^2 + \|\tilde{\beta}\|_1^2 \|\tilde{\mathbf{v}}_1\|_2^2 + \|\tilde{\theta}_1\|_2^2 \|\tilde{\mathbf{w}}\|_1^2 \right\} + \delta(\|\mathbf{w}\|_0^2 + \|\beta\|_0^2).
\end{aligned} \tag{3.15}$$

Also, from the Stokes equations

$$\begin{cases} -\mu\epsilon\Delta \mathbf{w} + \nabla \xi = \epsilon(\epsilon J - \tilde{\mathbf{w}} \cdot \nabla \tilde{\mathbf{v}}_2 - \tilde{\mathbf{v}}_1 \cdot \nabla \tilde{\mathbf{w}} - W \cdot \nabla \mathbf{v}_2 - \nabla(\tilde{\theta}_1 \xi + \tilde{\beta} \eta_2) + \zeta \nabla \operatorname{div} \mathbf{w}) - \nabla \beta, \\ \operatorname{div} \mathbf{w} = \operatorname{div} \mathbf{w}, \end{cases}$$

we get the following estimate

$$\begin{aligned}
\epsilon\|\mathbf{w}\|_2 + \|\nabla \xi\|_0 &\leq \epsilon C \left(\|\operatorname{div} \mathbf{w}\|_2 + \epsilon \|J\|_0 + \|\tilde{\mathbf{w}}\|_1 (\|\tilde{\mathbf{v}}_2\|_2 + \|\tilde{\mathbf{v}}_1\|_2) + \|W\|_1 \|\mathbf{v}_2\|_2 \right. \\
&\quad \left. + \|\tilde{\theta}_1\|_2 \|\xi\|_1 + \|\tilde{\beta}\|_1 \|\eta_2\|_2 \right) + C\|\beta\|_1.
\end{aligned} \tag{3.16}$$

Applying Poincaré's inequality and substituting (3.16) into (3.15), employing the estimates (3.11)–(3.14) and recalling the smallness of ϵ_0 and E , we conclude that

$$\|W\|_1 + \|\mathbf{w}\|_1 + \|\beta\|_1 \leq C(\|\tilde{W}\|_1 + \|\tilde{\mathbf{w}}\|_1 + \|\tilde{\beta}\|_1),$$

where C is a positive constant depending only on $\Omega, \mu, \lambda, \mathbf{f}, E$ and ϵ_0 . This completes the proof. \square

Finally, having had Lemmas 3.1 and 3.2, we can apply the Tikhonov fixed point theorem to find a fixed point $(U, \mathbf{v}, \theta) = N(U, \mathbf{v}, \theta)$ in the set K . Moreover, the pressure $P \in \bar{H}^2$ satisfies

$$\nabla P = \mathbf{f} + \mathbf{g} + \mu \Delta U - (U + \mathbf{v}) \cdot \nabla U,$$

and $(U + \mathbf{v}, \epsilon P + \eta, \theta)$ is a solution to (1.5). Thus, we have shown the following proposition.

Proposition 3.1. *Let $\mathbf{f}, \mathbf{g} \in H^2(\Omega)$. Then, there exists an ϵ_0 depending only on $\Omega, \mu, \lambda, \mathbf{f}$ and \mathbf{g} , such that for all $\epsilon \in (0, \epsilon_0)$, there is a solution $(U, P, \mathbf{v}, \eta, \theta) \in (H^4 \cap H_{0,\sigma}^1) \times \bar{H}^3 \times (H^3 \cap H_0^1) \times \bar{H}^2 \times (H^3 \cap H_0^1)$ of (2.1) and (2.2), satisfying*

$$\|\mathbf{v}\|_3 + \|\eta\|_2 + \|\theta\|_3 \leq E,$$

where E is a small positive constant depending only on $\Omega, \mu, \lambda, \mathbf{f}$ and \mathbf{g} . Moreover, $(U + \mathbf{v}, \epsilon P + \eta, \theta)$ is a solution of the system (1.5) for any $\epsilon \in (0, \epsilon_0)$.

4 Incompressible limit

Let $\epsilon < \epsilon_0$ and $(U^\epsilon, \mathbf{v}^\epsilon, \theta^\epsilon) \in K$ be the solution established in Proposition 3.1. We take $\mathbf{v} = \tilde{\mathbf{v}} = \mathbf{v}^\epsilon$, $\theta = \tilde{\theta} = \theta^\epsilon$ and $\eta = \eta^\epsilon$ in (3.6) to get that

$$\begin{aligned} \|\mathbf{v}^\epsilon\|_3 + \|\theta^\epsilon\|_3 + \|\eta^\epsilon\|_2 &\leq C_{16}(1 + M_1)^5 \left\{ \epsilon \left[\|\mathbf{f}\|_1 + (M_1 + 1)^2 \right] + (E + E^{\frac{1}{2}}) + \delta \right\} \|\eta\|_2 \\ &\quad + (\epsilon M_1^{\frac{1}{2}} + E) \|\mathbf{v}\|_3 + (\delta M_1^{\frac{1}{2}} + E) \|\theta\|_3 \\ &\quad + C_{16}(1 + M_1)^5 \left\{ \epsilon (M_1 + 1)^2 + \epsilon (1 + M_1) \left[\|\mathbf{f}\|_1 + (M_1 + 1)^2 \right] \right\}. \end{aligned}$$

Thus, by taking ϵ_0 and E so small that

$$C_{16}(1 + M_1)^5 \left\{ \epsilon \left[\|\mathbf{f}\|_1 + (M_1 + 1)^2 \right] + (E + E^{\frac{1}{2}}) \right\} < 1, \quad \epsilon M_1^{\frac{1}{2}} + E < 1,$$

we obtain

$$\|\mathbf{v}^\epsilon\|_3 + \|\theta^\epsilon\|_3 + \|\eta^\epsilon\|_2 \leq C_{16}(1 + M_1)^5 \left\{ \epsilon (M_1 + 1)^2 + \epsilon (1 + M_1) \left[\|\mathbf{f}\|_1 + (M_1 + 1)^2 \right] \right\},$$

whence,

$$\|\mathbf{v}^\epsilon\|_3 + \|\theta^\epsilon\|_3 + \|\eta^\epsilon\|_2 \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0. \quad (4.1)$$

Furthermore, from (2.2)₁, i.e.,

$$U^\epsilon \cdot \nabla \eta^\epsilon + \frac{\operatorname{div} \mathbf{v}^\epsilon}{\epsilon} = -\mathbf{v}^\epsilon \cdot \nabla \eta^\epsilon - \eta^\epsilon \operatorname{div} \mathbf{v}^\epsilon - \epsilon \operatorname{div} (P^\epsilon (U^\epsilon + \mathbf{v}^\epsilon))$$

and (4.1) we get that as $\epsilon \rightarrow 0$,

$$\left\| \frac{\operatorname{div} \mathbf{v}^\epsilon}{\epsilon} \right\|_1 \leq \|\mathbf{v}^\epsilon \cdot \nabla \eta^\epsilon\|_1 + \|\eta^\epsilon \operatorname{div} \mathbf{v}^\epsilon\|_1 + \|\epsilon \operatorname{div} (P^\epsilon (U^\epsilon + \mathbf{v}^\epsilon))\|_1 + \|U^\epsilon \cdot \nabla \eta^\epsilon\|_1 \rightarrow 0. \quad (4.2)$$

Due to (4.1) and

$$\frac{\nabla\eta^\epsilon + \nabla\theta^\epsilon}{\epsilon} = \epsilon F^\epsilon - \mathbf{v}^\epsilon \cdot \nabla \mathbf{v}^\epsilon - \theta^\epsilon \nabla \eta^\epsilon - \eta^\epsilon \nabla \theta^\epsilon - U^\epsilon \cdot \nabla \mathbf{v}^\epsilon + \mu \Delta \mathbf{v}^\epsilon + (\mu + \lambda) \nabla \operatorname{div} \mathbf{v}^\epsilon$$

with $F^\epsilon = (\epsilon P^\epsilon + \eta^\epsilon) \mathbf{f} - (\epsilon P^\epsilon + \eta^\epsilon)(U^\epsilon + \mathbf{v}^\epsilon) \cdot \nabla(U^\epsilon + \mathbf{v}^\epsilon) - \theta^\epsilon \nabla P^\epsilon - P^\epsilon \nabla \theta^\epsilon$, which comes from the transform of (2.2)₂, one deduces, recalling Poincaré's inequality, that

$$\left\| \frac{\eta^\epsilon + \theta^\epsilon}{\epsilon} \right\|_2 \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0. \quad (4.3)$$

On the other hand, in view of Lemma 2.2, we observe that (U^ϵ, P^ϵ) is a uniform-in- ϵ bounded sequence in $(H^4 \cap H_0^1) \times \bar{H}^3$. Hence, there are a subsequence of $(U^{\epsilon_k}, P^{\epsilon_k})$, still denoted by $(U^{\epsilon_k}, P^{\epsilon_k})$ for simplicity, and $(\bar{U}, \bar{P}) \in (H^4 \cap H_0^1) \times \bar{H}^3$, such that as $\epsilon \rightarrow 0$,

$$(U^\epsilon, P^\epsilon) \rightharpoonup (\bar{U}, \bar{P}), \quad \text{weakly in } (H^4 \cap H_0^1) \times \bar{H}^3,$$

and

$$(U^\epsilon, P^\epsilon) \rightarrow (\bar{U}, \bar{P}), \quad \text{strongly in } (H^3 \cap H_0^1) \times \bar{H}^2.$$

Thus, if we take to the limit as $\epsilon \rightarrow 0$ in (2.1) and (2.2), we conclude that (\bar{U}, \bar{P}) is a solution of the steady incompressible Navier-Stokes equations (1.7).

In conclusion, we have that

$$\liminf_{\epsilon \rightarrow 0} \inf_{U, P \in \mathbf{L}} \|U^\epsilon + \mathbf{v}^\epsilon - U\|_3 + \left\| P^\epsilon + \frac{\eta^\epsilon + \theta^\epsilon}{\epsilon} - P \right\|_2 + \|\theta^\epsilon\|_3 = 0,$$

where \mathbf{L} is the same as in Theorem 1.1. Thus, the proof of the low Mach number limit is completed.

Acknowledgements. The authors would like to thank Prof. Qiangchang Ju and Prof. Yue-Jun Peng for helpful discussions. This work was supported by the National Basic Research Program under the Grant 2011CB309705, NSFC (Grant Nos. 11229101, 11201115 and 11301083) and China Postdoctoral Science Foundation (Nos. 2012M510365, 2012M520205).

References

- [1] T. Alazard, Low Mach number limit of the full Navier-Stokes equations, *Arch. Ration. Mech. Anal.*, 180 (2006) 1-73.
- [2] H. Beirão da Veiga, An L^p -theory for the n -dimensional, stationary, compressible Navier-Stokes equations, and the incompressible limit for compressible fluids. The equilibrium solutions, *Comm. Math. Phys.*, 109 (1987) 229-248.
- [3] D. Bresch, B. Desjardins, E. Grenier, C.-K. Lin, Low Mach number limit of viscous polytropic flows: formal asymptotics in the periodic case, *Stud. Appl. Math.*, 109 (2002) 125-149.
- [4] J. Březina, A. Novotný, On weak solutions of steady Navier-Stokes equations for monoatomic gas, *Comment. Math. Univ. Carolinae*, 49 (2008) 611-632.
- [5] H. Choe and B. Jin, Existence of solutions of stationary compressible Navier-Stokes Equations with large force, *J. Func. Anal.*, 177 (2000) 54-88.
- [6] B. Dorin, E. Feireisl, The incompressible limit of the full Navier-Stokes-Fourier system on domains with rough boundaries, *Nonlinear Anal. Real World Appl.*, 10 (2009) 3203-3229.

- [7] R. Farwig, Stationary silutions of compressible Navier-Stokes equations with slip boundary condition, *Comm. Partial Differential Equations*, 14 (1989) 1579-1606.
- [8] E. Feireisl, J. Malek, A. Novotny, Navier's slip and incompressible limits in domains with variable bottoms, *Discrete Contin. Dyn. Syst. Ser. S*, 1 (2008) 427-460.
- [9] E. Feireisl, A. Novotny, The low Mach number limit for the full Navier-Stokes-fourier system, *Arch. Ration. Mech. Anal.*, 186 (2007) 77-107.
- [10] E. Feireisl, A. Novotny, H. Petzeltova, On the incompressible limit for the Navier-Stokes-Fourier system in domains with wavy bottoms, *Math. Models Methods Appl. Sci.*, 18 (2008) 291-324.
- [11] J. Frehse, S. Goj, M. Steinhauer, L^p -estimates for the Navier-Stokes equations for steady compressible flow, *Manuscripta Math.*, 116 (2005) 265-275.
- [12] J. Frehse, M. Steinhauer, W. Weigant, On stationary solutions for 2-D viscous compressible isothermal Navier-Stokes equations, *J. Math. Fluid Mech.*, 13 (2010) 55-63.
- [13] J. Frehse, M. Steinhauer, W. Weigant, The Dirichlet problem for steady viscous compressible flow in three dimensions, *J. Math. Pures Appl.*, Vol. 97, Issue 2, (2012) 85-97.
- [14] G.P. Galdi, An Introduction to the Mathematical Theory of the Navier-Stokes Equations. Vol. I. Linearized Steady Problems, Springer-Verlag, New York, 1994.
- [15] T. Hagstrom and J. Lorenz, On the stability of approximate solutions of hyperbolic-parabolic systems and the all-time existence of smooth, slightly compressible flows, *Indiana Univ. Math. J.*, 51 No. 6 (2002) 1339-1387.
- [16] Didier Jesslé and Antonín Novotný, Existence of renormalized weak solutions to the steady equations describing compressible fluids in barotropic regime *J. Math. Pures Appl.*, Vol. 99, Issue 3, (2013) 280-296.
- [17] S. Jiang and C.H. Zhou, Existence of weak solutions to the three-dimensional steady compressible Navier-Stokes equations, *Ann. Inst. H. Poincaré-AN*, 28 (2011) 485-498.
- [18] S. Jiang and Y.B. Ou, Incompressible limit of the non-isentropic Navier-Stokes equations with well-prepared initial data in three-dimensional bounded domains, *J. Math. Pures Appl.*, 96 (2011) 1-28.
- [19] P.-L. Lions, Mathematical topics in fluid dynamics, Vol 2, Compressible models, Oxford Science Publication, Oxford, 1998.
- [20] A. Matsumura and T. Nishida, Initial boundary value problems for the equations of motion of general fluids, *Computing Methods in Applied Sciences and Engineering V* (eds. R. Glowinski, J.L. Lions), North Holland, Amsterdam 1982, 389-406.
- [21] A. Matsumura and T. Nishida, Initial boundary value problems for the equations of motions of compressible viscous and heat-conductive fluids, *Comm. Math. Phys.*, 89 (1983) 445-464.
- [22] A. Matsumura and M. Padula, Stability of stationary flow of compressible fluids subject to large external potential forces, *Stab. Anal. Cont. Media*, (1992) 2183-2202.
- [23] A. Novotný and M. Pokorný, Weak and variational solutions to steady equations for compressible heat conducting fluids, *SIAM J. Math. Anal.*, 43 (2011) 1158-1188.
- [24] A. Novotný and M. Pokorný, Steady compressible Navier-Stokes-Fourier system for monoatomic gas and its generalizations, *J. Diff. Eqns.*, 251 (2011) 270-315.

- [25] A. Novotný and M. Padula, Existence and uniqueness of stationary solutions for viscous compressible heat conductive fluid with large potential and small nonpotential external forces, *Siberian Math. J.*, 34 (1991) 120-146.
- [26] A. Novotný and M. Padula, L^p -approach to steady flows of viscous compressible fluids in exterior domains, *Arch. Rational Meth. Anal.*, 126 (1994) 243-297.
- [27] A. Novotný and I. Straškraba, Introduction to the theory of compressible flow, *Oxford University Press*, Oxford, 2004.
- [28] M. Padula, Existence and uniqueness for viscous steady compressible motions, *Arch. Rational Mech. Anal.*, 97 (1987) 89-102.
- [29] P.I. Plotnikov, E.V. Rubana, J. Sokolowski, Inhomogeneous boundary value problems for compressible Navier-Stokes and transport equations, *J. Math. Pures Appl.*, 92 (2009) 113-162.
- [30] P.I. Plotnikov, J. Sokolowski, Concentrations of solutions to time-discretized compressible Navier-Stokes equations, *Comm. Math. Phys.*, 258 (2005) 567-608.
- [31] P.I. Plotnikov, J. Sokolowski, On compactness, domain dependence and existence of steady state solutions to compressible isothermal Navier-Stokes equations, *J. Math. Fluid Meth.*, 7 (2005) 529-573.
- [32] P.I. Plotnikov, J. Sokolowski, Stationary solutions of Navier-Stokes equations for diatomic gases, (Russian) *Uspekhi Mat. Nauk* 62 (2007) no.3(375) 117-148; English translation in *Russian Math. Surveys*, 62 No.3 (2007) 561-593.
- [33] P.I. Plotnikov, W. Weigant, Steady 3D viscous compressible flows with adiabatic exponent $\gamma \in (1, \infty)$, arXiv:1312.5829, Dec. 2013.
- [34] A. Valli, Periodic and stationary solutions for compressible Navier-Stokes equations by stability method, *Ann. Scuola Norm. Sup. Pisa*, (1983) 607-647.
- [35] A. Valli, On the existence of stationary solutions for compressible Navier-Stokes equation, *Ann. Inst. H. Poincaré-AN*, 4 (1987) 99-113.
- [36] A. Valli and W.M. Zajaczkowski, Navier-Stokes equations for compressible fluids: Global existence and qualitative properties of the solutions in the general case, *Comm. Math. Phys.*, 103 (1986) 259-296.