

The shape of spins

Christoph Englert,^{1,*} Dorival Gonçalves,^{2,3,†} Graeme Nail,^{1,‡} and Michael Spannowsky^{1,§}

¹*Institute for Particle Physics Phenomenology, Department of Physics,
Durham University, DH1 3LE, United Kingdom*

²*Institut für Theoretische Physik, Universität Heidelberg, 69120 Heidelberg, Germany*

³*Max-Planck-Institut für Physik, Föhringer Ring 6, 80805 München, Germany*

After the discovery of a Higgs-like particle at the LHC, the determination of its spin quantum numbers across different channels will be the next step in arriving at a more precise understanding of the new state and its role in electroweak symmetry breaking. Event shape observables have been shown to provide extremely sensitive observables for the discrimination of the scalar Higgs boson's \mathcal{CP} quantum numbers as a consequence of the different radiation patterns of Higgs production via gluon fusion vs. weak boson fusion in the $pp \rightarrow X + 2j$ selection. We show that a similar strategy serves to constrain the spin quantum numbers of the discovered particle as a function of the involved couplings. We also discuss the prospects of applying a similar strategy to future discoveries of Higgs-like particles.

I. INTRODUCTION

After the discovery of a Standard Model Higgs boson-like particle [1] at the LHC [2, 3], the measurement its spin is the next step in arriving at a more complete picture of this discovery. There is a theoretical prejudice from Lorentz invariance against spin $J = 1$ [4] as the particle is observed in the decay to photons, which leaves scalar $J = 0$ as the well-defined option in terms of our current understanding of perturbative Quantum Field Theory.

There is a known caveat in analyzing spin hypotheses $J \geq 2$ that arises when we investigate tensor particles and beyond. As a matter of fact, there is no well-behaved QFT which predicts the interactions of such a state with SM matter from first principles. In particular, there are certain indirect constraints on the spin $J = 2$ options if we take into account the non-observation of large excesses in $VV + 2j$ final states ($V = W^\pm, Z$) at the LHC so far, while there is consistency in $X \rightarrow VV$ with the SM within errors. The latter implies that the observed particle is involved in the unitarization of $V_L V_L$ scattering and probably provides the dominant share to the saturation of the unitarity sum rules. In simple realizations, this cannot be achieved with a spin 2 particle [5] and the worsened unitarity problem in longitudinal gauge boson scattering would manifest in a large cross section in the $VV + 2j$ final state at large invariant masses.

On the other hand, we can perform spin analyses beyond indirect constraints in model-independent ways in the fully reconstructible final states $X \rightarrow ZZ, \gamma\gamma$ [6–9]. Many of the direct measurement analysis strategies originate from similar questions addressed in hadron physics [10]. Doing so, one typically treats the X decay

independent from X production.¹ Indeed, recent LHC measurements along these lines seem to favor $J^{\mathcal{CP}} = 0^+$ searches [12, 13].

However, treating the resonance's decay independent from its production does not allow one to draw a more complete picture of Higgs couplings because momentum dependencies are typically encoded in off-shell effects that cannot be studied in this way. It is precisely the momentum dependence of higher dimensional operators that leaves footprints in the $X + 2j$ channel [14], *i.e.*, the t channel gauge bosons in the weak boson fusion (WBF) topologies are always virtual. In this sense, adapted search strategies for the $X + 2j$ selection do not only provide additional sensitivity, which can be used in a global spin hypothesis test across various channels, but also include orthogonal information that cannot be accessed via more traditional spin measurements.

In this letter we show that the global energy flow structure that follows from typical representatives of alternative spin structures provides a highly sensitive observable to study these properties. We select combinations of couplings, right from the beginning, that lead to a SM Higgs-like phenomenology. As Refs. [15–17] explain, the “tagging” jet kinematics in $X + 2j$ final states can be a strong discriminant for the spin of the produced particle X . It should be noted that this typically results from the involved (higher-dimensional) operator structures, which are determined by the spin hypotheses. With this in mind, we specifically analyze spin 2 models that have p_T distributions similar to the SM Higgs [16]. In doing so, we complement the analyzes of [15–17] by answering how much sensitivity hides beyond the tagging jet level and how it carries over to experimental reality.

We will also investigate the strategy's prospects for heavier “Higgs” masses. This latter point is motivated

*Electronic address: christoph.englert@durham.ac.uk

†Electronic address: dorival@mpp.mpg.de

‡Electronic address: graeme.nail@durham.ac.uk

§Electronic address: michael.spannowsky@durham.ac.uk

¹The simulation of such final states, however, needs to include the full matrix element because, *e.g.*, for a graviton-like object the only source of deviation is the propagator, see also [11].




FIG. 1: Sample Feynman diagrams that contribute to $X + 2j$ production via gluon fusion (a) and weak boson fusion (b). We do not show the X decay.

by the fact that similar questions, as to those we currently face for the 125 GeV particle, will arise if additional Higgs-like states are discovered in the future. Such states are predicted by many extensions of the SM Higgs sector.

Event shapes as electroweak-sensitive observables

The azimuthal angle between the two tagging jets in the $pp \rightarrow X + 2j$ selection $\Delta\Phi_{jj}$ [18–22] defined according to rapidity y

$$\Delta\Phi_{jj} = \phi(p_{>}) - \phi(p_{<}), \quad (1)$$

where $p_{\leq}^{\mu} = \sum_{j \in \{\text{jets: } y_j \leq y_X\}} p_j^{\mu}$, is known to be a highly sensitive observable to the \mathcal{CP} quantum numbers of the produced X state. This finding is not limited to the WBF channels [23], but is known to also work in the gluon fusion channel [22, 24]. The latter production mechanism can give rise to \mathcal{CP} odd Higgs production via tree-level \mathcal{CP} odd couplings to the heavy fermion sector, Fig. 1. Such a state is typically present in any non-singlet Higgs sector extension that feature fields transforming in non-trivial representations under $SU(2)_L$. In the light of recent measurements, the fields of these extensions need to be heavier, with suppressed cross sections.

Another way to understand the sensitivity encoded in $\Delta\Phi_{jj}$ is that the amplitude as a whole is sensitive to the \mathcal{CP} quantum numbers. Hence, any additional \mathcal{CP} -preserving QCD leg that is attached to diagrams in Fig. 1 will still give rise to an amplitude which encodes the \mathcal{CP} -specific properties reflected in $\Delta\Phi_{jj}$ for two-jet configurations. As a result, the entire QCD activity that results from the hard interactions in Fig. 1 can be considered a probe of the produced state X .² Finding the “proper” jets of Fig. 1 that reflect the nature of the produced state in a multi-jet environment amounts to a combinatorial

and quantum-interference-governed problem; this results in reduced sensitivity in the $\geq 3j$ selection [18].

A way to circumvent this was outlined in Ref. [16]: Since QCD radiation implies energy-momentum flow, the entire energy distribution in the detector (upon reconstructing and removing X from the list of calorimeter hits) can be expected to provide a superior discriminant compared to $\Delta\Phi_{jj}$ in an inclusive selection. The energy momentum flow of an LHC event is commonly quantified by means of hadronic event shape observables [25].³ Indeed, Ref. [27] found an increase in sensitivity that follows from investigating event shapes for discrete \mathcal{CP} measurements.⁴ The interplay between event shapes and Higgs physics was further studied in Ref. [28].

Recently in Refs. [15–17] a substantial discriminative power was revealed in the $pp \rightarrow X + 2j$ final state for different spin hypotheses $J(X)$. This sensitivity is driven by the energy-dependence of operators which mimic the Higgs boson’s interactions. The differences in the observed phenomenology can be manifold and depends on the specific higher spin scenario that one investigates. However, a rather generic finding is that spin 1 and 2 operators tend to populate the central region of the detector, thus leading to a departure from a WBF-like signature; consequentially central jet vetos [29, 30] need to be relaxed to be sensitive to such an event topology. This means that backgrounds need to be suppressed by a combination of stiff b vetos [31] and state-of-the-art signal vs. background (S/B) discriminators, such as the matrix element method [32], depending on the final state.⁵

In the following we will consider $pp \rightarrow X + 2j$ with X decay to fully leptonic taus for a toy-level signal vs. background study to compare the performance of various event shape-based observables. The details of the Higgs reconstruction are inconsequential in this comparison, as all observables are affected in the same way, and the Higgs candidate does not enter our analysis apart from reconstructing the signal within a window cut around the candidate mass of $m_X \simeq 125$ GeV. Hence, we do not include any tau reconstruction efficiencies that can also vary across the different exclusive tau decay modes [33, 34]. We also note that our analysis strategy is insensitive to the specifics of the “Higgs” decay channel, and our methods straightforwardly generalize to other decay channels such as, *e.g.*, the $\gamma\gamma + 2j$ selection.

³See *e.g.* Ref. [26] for publicly available implementations within the RIVET analysis package.

⁴Since the sensitivity does not follow from a specific angular distribution $\Delta\Phi_{jj}$ still remains the observable of choice for mixed \mathcal{CP} states, which can be straightforwardly extracted by fitting trigonometrical functions for an essentially background-free selection [23, 24]. This procedure only becomes available at high integrated luminosities.

⁵Another finding of [16, 17] is that the sensitivity observed, in the combination of transverse momentum and rapidity difference, points to the invariant dijet mass as single discriminant.

²In principle this argument extends also to the soft coherent radiation down to the hadronization scale. These effects are however subleading.

II. ANALYSIS SETUP

For the purpose of comparability, we closely follow Ref. [27]. We model our signal hypotheses with a combination of MADGRAPH [35] and HERWIG++ [36]. For the simulation of the backgrounds we generate matched events with SHERPA [37] and in the following limit ourselves to the $t\bar{t}$ +jets and Z +jets backgrounds [23]; normalizing these event samples to the NNLO [38, 39] and NLO cross sections [40–42], respectively.

We apply a typical WBF selection [23, 27] and cluster jets with the anti-kT algorithm [43] as implemented in FASTJET [44] with $D = 0.4$ and define jets with the thresholds

$$p_{T,j} \geq 40 \text{ GeV}, \text{ and } |y_j| \leq 4.5. \quad (2)$$

We impose an invariant mass cut on the two hardest tagging jets in the event of

$$m_{jj} = \sqrt{(p_{j,1} + p_{j,2})^2} \geq 600 \text{ GeV}, \quad (3)$$

and reconstruct the Higgs from taus with

$$p_\tau \geq 20 \text{ GeV and } |\eta_\tau| \leq 2.5 \quad (4)$$

within a 50 GeV window around 125 GeV. The Higgs candidate has to fall between the tagging jets

$$\min(y_1, y_2) < y_X < \max(y_1, y_2). \quad (5)$$

We further suppress the $t\bar{t}$ +jets background by imposing a central b veto with an efficiency of 80% [31]. The additional signal reduction due to mistagging is negligible within the approximations we make. When normalizing *all signal samples to the SM Higgs cross section* after cuts (*i.e.* we treat the $J = 2$ hypotheses as Higgs-lookalikes) we have signal cross sections $\sigma(X + 2j) = 3.82 \text{ fb}$. The combined background is $\sigma(\text{bkg}) = 6.54 \text{ fb}$.⁶

We proceed further by setting up two different track-selections that eventually enter the evaluation of the considered event shape observables. One of which is more robust against pile-up that can cause issues when we want to study the global event properties in the context of this paper.

- (i) For the events that pass the above selection criteria we feed all calorimeter hits with $p_T \geq 1 \text{ GeV}$ and $|\eta| \leq 4.5$ into the definition of the event shapes. This amounts to the most inclusive definition of the event shapes that is possible in the light of the above cuts. Selecting events according to the requirements Eq. (2)-(5) is already at odds with continuous globalness [25], which guarantees good re-summation properties [25]. However, the used selection is the most inclusive possible in the light of

unavoidable signal vs. background discrimination. To this end, we note that the analysis of Ref. [25] also shows that matched shower MCs reproduce the analytically resummed results well, so that we can expect our simulation to be under sufficient control. Quite obviously, this selection will be affected by pile-up activity.

- (ii) The pile-up conditions for $\sqrt{s} = 14 \text{ TeV}$ will need to be assessed when the LHC turns on again, but it can be expected that pile-up suppression in the central part of the detector is going to allow to lower jet thresholds in the rapidity region of the tracker $|\eta| \leq 2.5$ [45]. Currently, there is no tracking available for the more forward rapidity regions, so we will need to rely on hard jets to reduce in- and out-of time pile-up and underlying event.

To reflect the effect of pile-up suppression to achieve a more robust definition of our observables we modify our event selection. We cluster jets as before, with the anti-kT algorithm and $D = 0.4$, but this time we use the constituents of the jets obeying

$$p_{T,j} \geq \begin{cases} 40 \text{ GeV}, & 2.5 < |\eta_j| \leq 4.5 \\ 10 \text{ GeV}, & |\eta_j| \leq 2.5 \end{cases} \quad (6)$$

as input for the event shapes instead of all particle tracks as considered in (i). This also allows one to enhance pile-up suppression by *e.g.* using the method of Ref. [46]. Furthermore, we explicitly require additional jet activity (specifically $n_j \geq 3$) which probes the spin structure induced radiation pattern. Since we are requiring at least three jets according to these modified criteria, the sensitivity we will find can be straightforwardly enhanced by including sensitivity from $\Delta\Phi_{jj}$, $\Delta\eta_{jj}$, p_{T_j} (or equivalently m_{jj}) for the *exclusive* two jet category [16] in a hybrid observable approach. We find cross sections for these cuts of $\sigma(\text{signal}) = 1.89 \text{ fb}$, while the background remains unchanged.⁷

For the spin 2 hypothesis

$$\mathcal{L}_2 = -g_1 G_{\mu\nu} T_V^{\mu\nu} - g_2 G_{\mu\nu} T_G^{\mu\nu} - g_3 G_{\mu\nu} T_f^{\mu\nu}, \quad (7)$$

where $G_{\mu\nu}$ is the spin 2 resonance and $T_{V,G,f}^{\mu\nu}$ is the energy-momentum tensor for the EW gauge bosons, gluons and fermions, we consider two representative scenarios [16].

- 2⁺: The ordinary graviton-like tensor particle paradigm (*i.e.* $g_1 = g_2 = g_3 = 1/\Lambda$), as considered in many other publications (see *e.g.* Ref. [15, 47]), has jet

⁶The details of the cutflow are identical to Ref. [27] and can be found in this earlier publication.

⁷Note that this motivated central jet vetos [29, 30] in the first place. The sensitivity we are going to find is lost in employing CJV-bases analysis strategies.




FIG. 2: Event shape distribution for the different event shapes calculated from all particle tracks in $|\eta| < 4.5$ with $p_T \geq 1$ GeV for the selection (i). We also show $\Delta\Phi_{jj}$.




FIG. 3: Event shape distribution calculated from jet constituents of selection (ii). We also show $\Delta\Phi_{jj}$.




FIG. 4: Result of the binned log-likelihood hypotheses test based on the input of selection (i).




FIG. 5: Result of the binned log-likelihood hypotheses test based on the input of selection (ii).

kinematics in the $X + 2j$ final state that are close to the SM Higgs, once the additional selection cuts are imposed [16]. The tagging jets are well-separated in η and their p_T distribution is not too different from the SM Higgs boson.

2_{ew+q}^+ : We also study a model which has considerably harder jets while the WBF rapidity gap (and hence WBF-likeness) is still preserved. This specific model constrains the tensor couplings to weak bosons and fermions (*i.e.* $g_1 = g_3 = 1/\Lambda$ and $g_2 = 0$). This specific operator selection is therefore a less “reasonable” representative of a spin 2 Higgs-lookalike.

Our two choices will be clear from the discussion below, and are also motivated by our findings for heavier Higgs-like particles in Sec. III.

The results of a number of event shape observables (for their definition we refer the reader to the appendix and Ref. [25]) are depicted in Fig. 2 for selection (i). This figure should be compared to Fig. 3, which displays

the same distributions subject to the modified requirements (ii).

To quantify the statistical discriminative power of the event shape observables we perform a binned log-likelihood hypothesis test [48] in Figs. 4 and 5; this provides a statistically well-defined estimate of the luminosity (upon dividing out all reconstruction efficiencies) that is required to reject the spin 2 hypotheses at the 5 sigma level using the CL_S method [49].

The results of this analysis are shown in Fig. 4 for the 2_{ew+q}^+ and in Fig. 5 for the 2^+ cases. As already expected from Figs. 2 and 3, the broadening observables perform best. Depending on the specific scenario, these observables are robust against pile-up as discussed in (ii). Fig. 5, however, also shows that, when the jet kinematics become SM-like, this will be reflected in a lower sensitivity of the event shapes to the involved spin hypothesis. This especially holds when discriminative power at smaller broadening is lost due to soft radiation not taken into account for selection (ii) vs. (i). This also explains




FIG. 6: Event shape and Φ_{jj} distributions for selection (i) and $m_X = 300$ GeV.

our initial choice of the spin 2 hypotheses: 2^+ is characterized by soft radiation and therefore suitable to be studied using event shape observables. We find broadening observables to provide the strongest statistical sensitivity. However, while this model can be formidably constrained using event shapes if pile-up is under sufficient control, *i.e.*, when the actual selection can be chosen closer to (i), the discriminative power of the broadening observables is severely reduced for selection (ii). On the other hand, 2^+_{ew+q} which has a slightly harder spectrum is robust in our comparison (i) vs (ii) and the event shape observables provide a statistically appealing single-valued discriminant.

III. SPIN DISCRIMINATION OF FUTURE HIGGS-LIKE RESONANCES

Let us finally comment on the prospect of using the methods of the previous section also in the context of spin analyses of Higgs-like states that might be discovered in the future with a heavier mass. This is not immediately clear since the higher mass scale implies a different (soft) radiation pattern. As a representative example we discuss $m_X \simeq 300$ GeV.

In general we can expect relatively small couplings of this additional state to the electroweak gauge bosons Z and W , as current measurements seem to suggest that unitarity cancellations, which characteristically determine the couplings of additional massive scalars with corresponding couplings, are saturated by the 125 GeV state. The standard technique in $X \rightarrow ZZ$ [9, 52] might hence not be applicable and an investigation of the $X + 2j$ final state could well be the only phenomenologically available channel to constrain the spin and CP structure

of such a discovery.

We consider these reasons as enough motivation to limit ourselves for scalar boson candidates to the gluon fusion channel Fig. 1 (a). For the spin 2 candidates we will again adopt the scenarios of the previous section, which will have quite different phenomenology as compared to the $m_X = 125$ GeV case.

For spin 1 candidates our above arguments constrain the interactions of copies of the SM gauge bosons. The phenomenology of a Kaluza-Klein excitation spectrum as encountered in *e.g.* warped extra dimensions (and their dual interpretation as vectorial and axial vector resonances of a strongly-interacting sector) is therefore heavily suppressed in the SM vector boson final states. There is an exception to the unitarity argument which are $Z'ZZ$ interactions as determined in the generalized Landau Yang theorem [50]. The structure of the interaction vertices does not introduce an energy-dependent unitarity violation and hence, is not constrained by current measurements. We include this interaction to model a WBF (Fig. 1 (b)) spin 1 candidate $J(X) = 1^-_Z$.

Gluon-fusion contributions for spin 1 degrees of freedom analogous to Fig. 1 (a) are more difficult to model. Furry's theorem [51] guarantees the exact cancellation of vector current from $J^{CP}(X) = 1^-$ hypothesis in $gg \rightarrow X$. Axial vector currents still have to obey the Landau Yang theorem [4]. This renders an observation of prompt gluon fusion impossible; on-shell production exactly vanishes and gluon fusion becomes a function of the $J^{CP}(X) = 1^+$ particle's width and the virtuality of the gluon. These small effects are at odds with conventional bump searches and leave gluon fusion, as depicted in Fig. 1 (a), as the only production mechanism when such a state has suppressed couplings to the SM Z 's (these couplings are again determined by the generalized Landau Yang theo-




FIG. 7: Event shape comparison for the SM Higgs and 1_q^+ for the qq -induced channels including the full mass dependence $m_X = 300$ GeV.

rem). While the particle X , in Fig. 1 (a), can be considered on-shell for resonance-driven searches, the t channel gluons are always off-shell: this enables $J^{\mathcal{CP}}(X) = 1^+$ production via gluon fusion (see also Ref. [53]). For the moment we are not interested in a survey of the effects of $d > 6$ operators that are involved in these interactions [54] on the events' energy momentum flow. We however note that different effective operators will contribute to the gluon-gluon, gluon-quark, and quark-quark channels.

Instead, we will model axial vector particles in gluon fusion plus two jets by introducing a doublet of heavy fermions, which couple to the axial vector boson with couplings chosen such that anomaly cancellation is manifest. We keep the full mass dependence by simulating $qq' \rightarrow 1_q^+ + qq'$ events with a modified version of MADGRAPH v4.4 [55] that links a customized one-loop capable HELAS [56] library. To gain a qualitative picture we compare the energy momentum flow of this model against the corresponding full one-loop SM Higgs events $qq' \rightarrow 0_{\text{SM}}^+ + qq'$. The gluon-induced channels will populate more the central region, but do not change the overall picture.⁸

The (normalized) results are presented in Fig. 6 for the identical jet cut setup of Sec. II. We do not include the backgrounds as these depend on the specific decay channel in which such a future resonance will be discovered. Typical QCD background suppression will however always be centered around the cuts of the previous section, independent of the specific exclusive decay channel of X . From the shown distributions it is clear that there is substantial discriminative power in separating the scalar options from $1_{\bar{Z}}$ and 2^+ in the event shape observables. A combination with ordinary jet-based observables such as $\Delta\Phi_{jj}$ will serve to discriminate these options further for tighter selections if feasible.

In Fig. 7 we finally show the comparison of the quark channels for the 1_q^+ vs. 1_{SM}^+ , which also provides insights how different partonic channels (and hence effective op-

erators) will influence our findings. Indeed the shapes are rather identical to Fig. 6 for the scalar boson; we can therefore expect that the event shapes also serve to discriminate between 0^+ and 1^+ for various spin template combinations, beyond the approximations we have made. Note also that our spin 2 hypotheses behave completely opposite compared to the $m_h = 125$ GeV case due to the changed momentum dependence of the cross section on the tagging jets. In this sense, $2_{\text{ew}+q}^+$ provides a better alternative hypothesis than 2^+ when such a measurement is performed in the future.

IV. SUMMARY AND CONCLUSIONS

The recent discovery of a Higgs-like particle at the LHC and further measurements of it seem to suggest that we have indeed discovered a particle which is consistent with the $J^{\mathcal{CP}}(X) = 0^+$ SM Higgs boson prediction. Analyses with increased statistics across many different channels will allow to answer the $J^{\mathcal{CP}}$ question more reliably. The $pp \rightarrow X + 2j$ mode, when analyzed in inclusive selections, provides a valuable channel to discriminate between different spin (and \mathcal{CP}) hypotheses when the events' global QCD energy-momentum flow pattern is analyzed. The latter is most efficiently captured in event shape distributions. While thrust provides a straightforward handle to discriminate discrete \mathcal{CP} values [27], the broadening observables reflect the spin-induced radiation patterns. Issues that may arise from challenging pile-up conditions can be counteracted with adopted definitions of the event shape observables and hybrid exclusive/inclusive definitions of the employed single valued discriminants. Depending on the spin 2 scenario (no spin 2 scenario is theoretically motivated but merely invoked as an alternative hypothesis to be excluded) we find large discriminative power in the accompanied energy momentum flow. This generalizes the results of Refs.[15–17, 27]. Pile-up, as for many analyses, can become a challenge of the discussed analysis strategy to the point where discriminative power in all collider observables is lost in the $X + 2j$ final state. This again highly depends on the chosen hypothesis.

⁸The quark-gluon and gluon-gluon- induced channels do not introduce a different $\Delta\Phi_{jj}$ radiation pattern for instance [21, 24], Fig. 7.

Given the consistency of the observed cross sections in $pp \rightarrow X \rightarrow ZZ, W^+W^-$ with the SM Higgs boson, it is likely that spin analyses of an additional resonance as predicted by many BSM scenarios cannot be straightforwardly performed in the $X \rightarrow \gamma\gamma, ZZ$ channels. In this case an event shape based analysis of the QCD energy momentum flow might be crucial since it does not rely on a particular exclusive final state decay of X . Indeed, we find significant discriminative power of the event shape observables for heavier ‘‘Higgs’’ masses, which allows to discriminate various J^{CP} hypotheses in combination with exclusive 2-jet measurements in the same channel [16]. As shown in this work, the advantages of event shape-based analyses are not limited to the study of pure QCD events but clearly generalize to the interplay of QCD with the (BSM) electroweak sector.

Acknowledgments — We thank Andrea Banfi, Gavin Salam, and Giulia Zanderighi for providing the CAESAR event shape library. C.E. acknowledges funding by the Durham International Junior Research Fellowship scheme.

Appendix A: Definitions of the studied event shapes

Event shapes are widely used observables to investigate geometrical properties of particle collisions at lepton and hadron colliders [57–61], which can be described to very high theoretical accuracy, see *e.g.* [25, 61]. At hadron colliders one typically defines the observables in the beam transverse plane. *Transverse thrust* is therefore defined as the maximization procedure in the transverse plane

$$T_{\perp,g} = \max_{\mathbf{n}_T} \frac{\sum_i |\mathbf{p}_{\perp,i} \cdot \mathbf{n}_T|}{\sum_i |\mathbf{p}_{\perp,i}|}, \quad |\mathbf{n}_T| = 1, \quad (\text{A1})$$

where $p_{T,i}$ denotes the transverse momentum of the track i . The transverse thrust value of circularly symmetric event is $T_{\perp,g} = 2/\pi \simeq 0.64$, while an ideal alignment is characterized by $T_{\perp,g} = 1$.

As a result of the maximization procedure we obtain the transverse thrust axis \mathbf{n}_T which enters the definition of transverse *thrust minor*

$$T_{m,g} = \frac{\sum_i |\mathbf{p}_{\perp,i} \times \mathbf{n}_T|}{\sum_i |\mathbf{p}_{\perp,i}|}, \quad (\text{A2})$$

which measures the energy-momentum flow perpendicular to the transverse thrust axis.

Observables that are particularly helpful in the context of spin analyses are broadening observables [59]. For their definitions we first specify a central region, C , in terms of pseudorapidity; here C corresponds to $|\eta| \leq 4.5$. Then we split this region according to transverse thrust axis

$$\text{region } \begin{matrix} C_U \\ C_D \end{matrix} \quad \mathbf{p}_{\perp,i} \cdot \mathbf{n}_T \gtrless 0 \quad (\text{A3a})$$

and subsequently compute the weighted pseudorapidity and azimuthal angle

$$\eta_\sigma = \frac{\sum_i |\mathbf{p}_{\perp,i}| \eta_i}{\sum_i |\mathbf{p}_{\perp,i}|}, \quad \phi_\sigma = \frac{\sum_i |\mathbf{p}_{\perp,i}| \phi_i}{\sum_i |\mathbf{p}_{\perp,i}|}, \quad \sigma = C_U, C_D. \quad (\text{A3b})$$

The broadening of the above regions is then defined as

$$B_\sigma = \frac{1}{2Q_T} \sum_{i \in \sigma} |\mathbf{p}_{\perp,i}| \sqrt{(\eta_i - \eta_\sigma)^2 + (\phi_i - \phi_\sigma)^2}, \quad \sigma = C_U, C_D \quad (\text{A3c})$$

with $Q_T = \sum_i |\mathbf{p}_{\perp,i}|$. The *central total broadening* and *central wide broadening* observables are

$$\begin{aligned} \text{central total broadening: } B_T &= B_{C_U} + B_{C_D}, \\ \text{central wide broadening: } B_W &= \max\{B_{C_U}, B_{C_D}\}. \end{aligned} \quad (\text{A3d})$$

-
- [1] P. W. Higgs, Phys. Lett. **12** (1964) 132 and Phys. Rev. Lett. **13** (1964) 508. F. Englert and R. Brout, Phys. Rev. Lett. **13** (1964) 321. G. S. Guralnik, C. R. Hagen and T. W. B. Kibble, Phys. Rev. Lett. **13** (1964) 585.
 - [2] The ATLAS collaboration, Phys. Lett. B **716** (2012) 1 and ATLAS-CONF-2012-170.
 - [3] The CMS collaboration, Phys. Lett. B **716** (2012) 30 and CMS-PAS-HIG-12-045.
 - [4] L.F. Landau, Dok. Akad. Nauk USSR **60** (1948) 207. C.N. Yang, Phys. Rev. **77** (1950) 242.
 - [5] A. Alboteanu, W. Kilian and J. Reuter, JHEP **0811** (2008) 010.
 - [6] Y. Gao, A. V. Gritsan, Z. Guo, K. Melnikov, M. Schulze and N. V. Tran, Phys. Rev. D **81** (2010) 075022. A. De Rujula, J. Lykken, M. Pierini, C. Rogan and M. Spiropulu, Phys. Rev. D **82** (2010) 013003.
 - [7] S. Y. Choi, M. M. Muhlleitner and P. M. Zerwas, Phys. Lett. B **718** (2013) 1031.
 - [8] J. Ellis and D. S. Hwang, arXiv:1202.6660 [hep-ph].
 - [9] D. J. Miller, 2, S. Y. Choi, B. Eberle, M. M. Muhlleitner and P. M. Zerwas, Phys. Lett. B **505** (2001) 149. S. Y. Choi, D. J. Miller, M. M. Muhlleitner and P. M. Zerwas, Phys. Lett. B **553**, 61 (2003). C. P. Buszello, I. Fleck, P. Marquard and J. J. van der Bij, Eur. Phys. J. C **32** (2004) 209.
 - [10] N. Cabibbo and A. Maksymowicz, Phys. Rev. **137**, B438 (1965) [Erratum-ibid. **168**, 1926 (1968)]. T. L. Trueman, Phys. Rev. D **18** (1978) 3423. J. R. Dell’Aquila and C. A. Nelson, Phys. Rev. D **33** (1986) 80.
 - [11] S. Banerjee, J. Kalinowski, W. Kotlarski, T. Przedzinski and Z. Was, arXiv:1212.2873 [hep-ph].
 - [12] The Atlas collaboration, ATLAS-CONF-2013-029,

- ATLAS-CONF-2013-031. The CMS collaboration, CMS-PAS-HIG-13-002.
- [13] A. Djouadi, G. Moreau, arXiv:1303.6591 [hep-ph].
- [14] V. Hankele, G. Klamke, D. Zeppenfeld and T. Figy, Phys. Rev. D **74** (2006) 095001.
- [15] J. Frank, M. Rauch and D. Zeppenfeld, arXiv:1211.3658 [hep-ph].
- [16] C. Englert, D. Goncalves-Netto, K. Mawatari and T. Plehn, JHEP **1301** (2013) 148
- [17] A. Djouadi, R. M. Godbole, B. Mellado and K. Mohan, arXiv:1301.4965 [hep-ph].
- [18] J. R. Andersen, K. Arnold and D. Zeppenfeld, JHEP **1006**, 091 (2010).
- [19] K. Hagiwara, Q. Li and K. Mawatari, JHEP **0907** (2009) 101.
- [20] V. Del Duca, W. Kilgore, C. Oleari, C. R. Schmidt and D. Zeppenfeld, Phys. Rev. D **67**, 073003 (2003).
- [21] F. Campanario, M. Kubocz and D. Zeppenfeld, Phys. Rev. D **84** (2011) 095025.
- [22] T. Figy, C. Oleari and D. Zeppenfeld, Phys. Rev. D **68**, 073005 (2003). T. Figy and D. Zeppenfeld, Phys. Lett. B **591**, 297 (2004). V. Del Duca, G. Klamke, D. Zeppenfeld, M. L. Mangano, M. Moretti, F. Piccinini, R. Pittau and A. D. Polosa, JHEP **0610** (2006) 016. J. M. Campbell, R. K. Ellis and G. Zanderighi, JHEP **0610** (2006) 028.
- [23] T. Plehn, D. L. Rainwater and D. Zeppenfeld, Phys. Rev. Lett. **88** (2002) 051801.
- [24] G. Klamke and D. Zeppenfeld, JHEP **0704** (2007) 052.
- [25] A. Banfi, G. P. Salam and G. Zanderighi, JHEP **1006** (2010) 038. A. Banfi, G. P. Salam and G. Zanderighi, JHEP **0201** (2002) 018. A. Banfi, G. P. Salam and G. Zanderighi, JHEP **0408** (2004) 062.
- [26] A. Buckley, J. Butterworth, L. Lonnblad, H. Hoeth, J. Monk, H. Schulz, J. E. von Seggern and F. Siegert *et al.*, arXiv:1003.0694 [hep-ph].
- [27] C. Englert, M. Spannowsky and M. Takeuchi, JHEP **1206** (2012) 108.
- [28] C. Bernaciak, M. S. A. Buschmann, A. Butter and T. Plehn, arXiv:1212.4436 [hep-ph].
- [29] Y. L. Dokshitzer, S. I. Troian and V. A. Khoze, Sov. J. Nucl. Phys. **46**, 712 (1987) [Yad. Fiz. **46**, 1220 (1987)]. Y. L. Dokshitzer, V. A. Khoze and T. Sjostrand, Phys. Lett. B **274** (1992) 116. J. D. Bjorken, Phys. Rev. D **47** (1993) 101.
- [30] U. Baur and E. W. N. Glover, Phys. Lett. B **252** (1990) 683. V. D. Barger, K. -m. Cheung, T. Han and D. Zeppenfeld, Phys. Rev. D **44** (1991) 2701 [Erratum-ibid. D **48** (1993) 5444]. D. L. Rainwater, R. Szalapski and D. Zeppenfeld, Phys. Rev. D **54** (1996) 6680. B. E. Cox, J. R. Forshaw and A. D. Pilkington, Phys. Lett. B **696** (2011) 87. S. Ask, J. H. Collins, J. R. Forshaw, K. Joshi and A. D. Pilkington, JHEP **1201** (2012) 018.
- [31] see *e.g.* The ATLAS collaboration ATLAS-CONF-2012-043.
- [32] K. Kondo, J. Phys. Soc. Jap. **57**, 4126 (1988). D. E. Soper and M. Spannowsky, Phys. Rev. D **84**, 074002 (2011). J. M. Campbell, W. T. Giele and C. Williams, JHEP **1211** (2012) 043. J. R. Andersen, C. Englert and M. Spannowsky, Phys. Rev. D **87** (2013) 015019. A. Freitas and J. S. Gainer, arXiv:1212.3598 [hep-ph]. J. M. Campbell, R. K. Ellis, W. T. Giele and C. Williams, arXiv:1301.7086 [hep-ph].
- [33] M. Heldmann, D. Cavalli, ATL-PHYS-PUB-2006-008, ATL-COM-PHYS-2006-010.
- [34] The CMS Collaboration, CMS-PAS-TAU-11-001, 2011, and Phys. Lett. B **713** (2012) 68.
- [35] J. Alwall, M. Herquet, F. Maltoni, O. Mattelaer and T. Stelzer, JHEP **1106** (2011) 128.
- [36] M. Bahr, S. Gieseke, M. A. Gigg, D. Grellscheid, K. Hamilton, O. Latunde-Dada, S. Platzer and P. Richardson *et al.*, Eur. Phys. J. C **58** (2008) 639.
- [37] T. Gleisberg, S. Hoeche, F. Krauss, M. Schonherr, S. Schumann, F. Siegert and J. Winter, JHEP **0902** (2009) 007. S. Schumann, F. Krauss, JHEP **0803** (2008) 038. T. Gleisberg and S. Hoeche, JHEP **0812** (2008) 039. S. Hoeche, F. Krauss, S. Schumann, F. Siegert, JHEP **0905** (2009) 053.
- [38] P. Nason, S. Dawson and R. K. Ellis, Nucl. Phys. B **327**, 49 (1989) [Erratum -ibid. B **335**, 260 (1990)]. W. Beenakker, W.L. van Neerven, R. Menge, G.A. Schuler and J. Smith, Nucl. Phys. B **351**, 507 (1991). M. L. Mangano, P. Nason and G. Ridolfi, Nucl. Phys. B **373**, 295 (1992). S. Frixione, M. L. Mangano, P. Nason and G. Ridolfi, Phys. Lett. B **351**, 555 (1995).
- [39] S. Moch and P. Uwer, Nucl. Phys. Proc. Suppl. **183** (2008) 75. P. Baernreuther, M. Czakon and A. Mitov, Phys. Rev. Lett. **109** (2012) 132001. M. Czakon and A. Mitov, JHEP **1212** (2012) 054. M. Czakon, P. Fiedler, A. Mitov arXiv:1303.6254 [hep-ph].
- [40] J. M. Campbell and R. K. Ellis, Nucl. Phys. Proc. Suppl. **205-206** (2010) 10. R. K. Ellis, Nucl. Phys. Proc. Suppl. **160**, 170 (2006).
- [41] K. Arnold, M. Bahr, G. Bozzi, F. Campanario, C. Englert, T. Figy, N. Greiner and C. Hackstein *et al.*, Comput. Phys. Commun. **180** (2009) 1661.
- [42] J. M. Campbell, R. K. Ellis, Phys. Rev. D **65** (2002) 113007. C. Oleari and D. Zeppenfeld, Phys. Rev. D **69** (2004) 093004.
- [43] M. Cacciari, G. P. Salam and G. Soyez, JHEP **0804** (2008) 063.
- [44] M. Cacciari, G. P. Salam and G. Soyez, arXiv:1111.6097 [hep-ph].
- [45] The CMS collaboration, CMS PAS PFT-09/001.
- [46] G. Soyez, G. P. Salam, J. Kim, S. Dutta and M. Cacciari, arXiv:1211.2811 [hep-ph].
- [47] J. Ellis, D. S. Hwang, V. Sanz and T. You, JHEP **1211** (2012) 134. C. -Q. Geng, D. Huang, Y. Tang and Y. -L. Wu, Phys. Lett. B **719** (2013) 164. J. Ellis, R. Fok, D. S. Hwang, V. Sanz and T. You, arXiv:1210.5229 [hep-ph]. J. Ellis, V. Sanz and T. You, arXiv:1211.3068 [hep-ph]. T. Modak, D. Sahoo, R. Sinha and H. -Y. Cheng, arXiv:1301.5404 [hep-ph].
- [48] T. Junk, Nucl. Instrum. Meth. A **434** (1999) 435. T. Junk, CDF Note 8128 [cdf/doc/statistics/public/8128]. T. Junk, CDF Note 7904 [cdf/doc/statistics/public/7904]. H. Hu and J. Nielsen, in 1st Workshop on Confidence Limits, CERN 2000-005 (2000).
- [49] A. L. Read, CERN-OPEN-2000-205. A. L. Read, J. Phys. G **G28** (2002) 2693-2704.
- [50] W. -Y. Keung, I. Low and J. Shu, Phys. Rev. Lett. **101** (2008) 091802.
- [51] W. H. Furry, Phys. Rev., **51**, 125 (1937).
- [52] C. Hackstein and M. Spannowsky, Phys. Rev. D **82**, 113012 (2010). C. Englert, C. Hackstein and M. Spannowsky, Phys. Rev. D **82**, 114024 (2010).
- [53] J. J. van der Bij, and E. W. N. Glover, Nucl. Phys. B **313** (1989) 237.

- [54] W. Buchmuller, D. Wyler and , Nucl. Phys. B **268** (1986) 621.
- [55] J. Alwall, P. Demin, S. de Visscher, R. Frederix, M. Herquet, F. Maltoni, T. Plehn and D. L. Rainwater *et al.*, JHEP **0709** (2007) 028.
- [56] H. Murayama, I. Watanabe and K. Hagiwara, KEK-Report 91-11, 1992.
- [57] S. Brandt, C. Peyrou, R. Sosnowski and A. Wroblewski, Phys. Lett. **12**, 57 (1964). E. Fahri, Phys. Rev. Lett. **39**, (1977) 1587.
- [58] V. Khachatryan *et al.* [CMS Collaboration], Phys. Lett. B **699** (2011) 48.
- [59] S. Catani, G. Turnock and B. R. Webber, Phys. Lett. B **295** (1992) 269. Y. L. Dokshitzer, A. Lucenti, G. Marchesini and G. P. Salam, JHEP **9801** (1998) 011.
- [60] A. Heister *et al.* [ALEPH Collaboration], Eur. Phys. J. C **35**, 457 (2004). J. Abdallah *et al.* [DELPHI Collaboration], Eur. Phys. J. C **37** (2004) 1. P. Achard *et al.* [L3 Collaboration], Phys. Rept. **399** (2004) 71.
- [61] A. Gehrmann-De Ridder, T. Gehrmann, E. W. N. Glover and G. Heinrich, Phys. Rev. Lett. **99** (2007) 132002. T. Becher and M. D. Schwartz, JHEP **0807** (2008) 034.