The characterization of Hermitian surfaces by the number of points

Masaaki Homma *
Department of Mathematics and Physics
Kanagawa University
Hiratsuka 259-1293, Japan
homma@kanagawa-u.ac.jp

Seon Jeong Kim †
Department of Mathematics and RINS
Gyeongsang National University
Jinju 660-701, Korea
skim@gnu.kr

Abstract

The nonsingular Hermitian surface of degree $\sqrt{q} + 1$ is characterized by its number of \mathbb{F}_q -points among the surfaces over \mathbb{F}_q of degree $\sqrt{q} + 1$ in the projective 3-space without \mathbb{F}_q -plane components. Key Words: Finite field, Hermitian surface, Weil-Deligne bound MSC: 11G25, 14G15, 14J70, 14N05

1 Introduction

Hermitian varieties are known as ones having particular properties over finite fields. Throughout this paper, except Section 3, q is an even power of a prime number p. A Hermitian variety over \mathbb{F}_q is a hypersurface in \mathbb{P}^n defined by

$$(X_0^{\sqrt{q}}, \dots, X_n^{\sqrt{q}}) A^t(X_0, \dots, X_n) = 0,$$
 (1)

where A is a square matrix of order n+1 whose entries are in \mathbb{F}_q with the property ${}^tA = A^{(\sqrt{q})}$; here $A^{(\sqrt{q})}$ means taking entry-wise the \sqrt{q} -th power, and tA is the transposed matrix of A. We refer this kind of polynomial as a Hermitian polynomial over \mathbb{F}_q . For a homogeneous polynomial F of degree $\sqrt{q}+1$ whose coefficients are

^{*}Partially supported by Grant-in-Aid for Scientific Research (24540056), JSPS.

 $^{^\}dagger Partially$ supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education, Science and Technology (2012R1A1A2042228).

in \mathbb{F}_q , the hypersurface given by F=0 is Hermitian if and only if there is an element $\rho \in \mathbb{F}_q^*$ such that ρF is a Hermitian polynomial. The family of Hermitian polynomials over \mathbb{F}_q forms an $\mathbb{F}_{\sqrt{q}}$ vector space. It is obvious that the Hermitian polynomial (1) defines a nonsingular Hermitian variety if and only if $\det A \neq 0$. By the standard argument, a nonsingular Hermitian variety is projectively equivalent to the variety

$$H_{n-1}: X_0^{\sqrt{q}+1} + X_1^{\sqrt{q}+1} + \dots + X_n^{\sqrt{q}+1} = 0$$
 (2)

over \mathbb{F}_q .

The number $N_q(H_{n-1})$ of \mathbb{F}_q -points of H_{n-1} is

$$N_q(H_{n-1}) = (\sqrt{q}^{n+1} + (-1)^n)(\sqrt{q}^n + (-1)^{n-1})/(q-1), \tag{3}$$

which is due to [1].

This number is remarkable in the following sense. The Weil conjecture [15] established by Deligne [2] implies that the number N of \mathbb{F}_q -points of a nonsingular hypersurface of degree d in \mathbb{P}^n defined over \mathbb{F}_q is bounded by

$$N \le \frac{q^n - 1}{q - 1} + \frac{d - 1}{d} \left((d - 1)^n - (-1)^n \right) \sqrt{q}^{n - 1}. \tag{4}$$

Furthermore, if equality holds in (4) for a certain nonsingular hypersurface of degree d, then $d \leq \sqrt{q} + 1$.

As for this additional claim, see [12, Corollary 2.2] or [8, Corollary 4.3]. The number $N_q(H_{n-1})$ achieves the equality in (4) for $d = \sqrt{q} + 1$.

There is a characterization of the Hermitian curve among the family of plane curves over \mathbb{F}_q of degree $\sqrt{q}+1$ without \mathbb{F}_q -line components [3]. Note that $N_q(H_1) = \sqrt{q^3} + 1$.

Theorem 1.1 (Hirschfeld-Storme-Thas-Voloch) Suppose that $q \neq 4$. Let C be a plane curve over \mathbb{F}_q of degree $\sqrt{q}+1$ without \mathbb{F}_q -line components. If C has \sqrt{q}^3+1 points over \mathbb{F}_q , then C is a Hermitian curve.

The purpose of this paper is to show a similar fact for surfaces in \mathbb{P}^3 .

Theorem 1.2 Let S be a surface in \mathbb{P}^3 defined over \mathbb{F}_q without \mathbb{F}_q -plane components. If the degree of S is $\sqrt{q} + 1$ and $N_q(S) = (\sqrt{q}^3 + 1)(q + 1)$, then S is a nonsingular Hermitian surface over \mathbb{F}_q .

After Hirschfeld-Storme-Thas-Voloch's characterization of the Hermitian curve established, Rück-Stichtenoth gave another characterization of the Hermitian curve among the family of nonsingular curves defined over \mathbb{F}_q of genus $\frac{\sqrt{q}(\sqrt{q}-1)}{2}$ [10]. A connection of those two characterization will be mentioned in Appendix.

Notation • When X is an algebraic set in \mathbb{P}^n defined by equations over \mathbb{F}_q , the set of \mathbb{F}_q -points in X is denoted by $X(\mathbb{F}_q)$, and the cardinality of $X(\mathbb{F}_q)$ by $N_q(X)$.

- When coordinates X_0, \ldots, X_n of \mathbb{P}^n are given, for a homogeneous polynomial h in those variables, $\{h=0\}$ means the hypersurface defined by h=0.
- Let D and E are curves in \mathbb{P}^2 without common components. For a point $Q \in D \cap E$, i(D.E;Q) denotes the local intersection multiplicity of D and E at Q, and $(D.E) = \sum_{Q \in D \cap E} i(D.E;Q)$.
- For a plane curve $C \subset \mathbb{P}^2$ and a nonsingular point $P \in C$, $T_P(C)$ denotes the embedded tangent line to C at P.
- The multiplicative set $\mathbb{F}_q \setminus \{0\}$ is denoted by \mathbb{F}_q^* .
- For a finite set Y, #Y denotes the number of elements of Y.

2 Plane curves

For a plane curve C over \mathbb{F}_q , we proved a simple bound for $N_q(C)$ in a series of papers [5, 6, 7], which had been originally conjectured by Sziklai [13].

Theorem 2.1 (Sziklai bound) Let d be an integer with $2 \le d \le q + 2$, and C a curve of degree d in \mathbb{P}^2 defined over \mathbb{F}_q without \mathbb{F}_q -linear components. Then

$$N_q(C) \le (d-1)q + 1 \tag{5}$$

except for curves over \mathbb{F}_4 which are projectively equivalent to the curve defined by

$$K: (X_0 + X_1 + X_2)^4 + (X_0X_1 + X_1X_2 + X_2X_0)^2 + X_0X_1X_2(X_0 + X_1 + X_2) = 0.$$

For the exceptional curve K above, $N_4(K) = 14$.

The number $N_q(H_1)$ attains the equality of the Sziklai bound too.

This bound implies a sufficient condition on a plane curve over \mathbb{F}_q to be absolutely irreducible.

Corollary 2.2 Let C be a curve over \mathbb{F}_q of degree d in \mathbb{P}^2 without \mathbb{F}_q -linear components. If $N_q(C) \geq (d-2)q+3$, then C is absolutely irreducible.

Proof. First note that the range of d is $2 \le d \le q + 2$ by assumptions.

(Step 1) Suppose C is decomposed into two curves C_1 and C_2 over \mathbb{F}_q . For each i=1,2, let d_i be the degree of C_i . If either " $q \neq 4$ " or "q=4 and none of the C_i 's is projectively equivalent to K," then

$$N_a(C) \le N_a(C_1) + N_a(C_2) \le ((d_1 - 1)q + 1) + ((d_2 - 1)q + 1) \le (d - 2)q + 2.$$

We consider the exceptional case, namely "q=4 and one of the C_i 's is projectively equivalent to K oner \mathbb{F}_4 ." Applying an \mathbb{F}_4 -linear transformation if necessary, we may assume that $C_1=K$. Then d=6 and $d_2=2$, because $2 \leq d \leq 6$ and C

has no \mathbb{F}_4 -linear components. Since C_2 is \mathbb{F}_4 -irreducible of degree 2, $N_4(C_2)=5$ or 1. When $N_4(C_2)=1$, $N_4(C)\leq 14+1<(6-2)4+3$. If C_2 is an absolutely irreducible conic, then $\#(C_2\cap\mathbb{P}^2(\mathbb{F}_2))\leq 4$, because the maximum length of \mathbb{F}_2 -arc is 4. Since $K(\mathbb{F}_4)=\mathbb{P}^2(\mathbb{F}_4)\setminus\mathbb{P}^2(\mathbb{F}_2)$ (see [5, §3]), $\#(K(\mathbb{F}_4)\cap C_2(\mathbb{F}_4))\geq 1$. Hence $N_q(C)\leq 14+5-1=(6-2)4+2$.

Therefore C is irreducible over \mathbb{F}_q .

(Step 2) Suppose that the \mathbb{F}_q -irreducible curve C is not absolutely irreducible. Let D be an irreducible component of C. Then D is defined over an extension \mathbb{F}_{q^t} of \mathbb{F}_q with $t \geq 2$, and $C = D \cup D^{(q)} \cup \cdots \cup D^{(q^{t-1})}$, where $D^{(q)}, \ldots, D^{(q^{t-1})}$ are conjugates of D over \mathbb{F}_q . Hence $C(\mathbb{F}_q) \subset D \cap D^{(q)}$ and $\deg D = \deg D^{(q)} \leq \frac{d}{2}$. Hence $N_q(C) \leq \left(\frac{d}{2}\right)^2$. Since $2 \leq d \leq q+2$, we have $\left(\frac{d}{2}\right)^2 < (d-2)q+3$. Indeed, consider the quadratic $P(d) = \left(\frac{d}{2}\right)^2 - (d-2)q-3$. Then P(2) < 0 and P(q+2) < 0, which implies P(d) < 0 for any d with $2 \leq d \leq q+2$.

Therefore the curve C is absolutely irreducible too.

An important step of the proof of Theorem 1.2 is to study plane sections of S and to apply the Hirschfeld-Storme-Thas-Voloch theorem to some of them, but the exception in the assertion of this theorem harms this step for q=4. So we take a detour in this case.

Lemma 2.3 Let C be a plane curve of degree 3 over \mathbb{F}_4 without \mathbb{F}_4 -linear components. Suppose $N_4(C) = 9$. Then C is absolutely irreducible and all \mathbb{F}_4 -points are nonsingular. In addition, suppose that all \mathbb{F}_4 -points are flexes, i.e., $i(T_p(C).C; P) = 3$ for any $P \in C(\mathbb{F}_4)$. Then C is a Hermitian curve.

Proof. From (2.2), C is absolutely irreducible. Let $P \in C(\mathbb{F}_4)$. Assume that P is a singular point. Then for any \mathbb{F}_4 -line passing through P, $\#(l(\mathbb{F}_4) \cap C \setminus \{P\}) \leq 1$ because (l.C) = 3 and $i(l.C; P) \geq 2$. Hence $N_4(C) \leq 6$ because $C(\mathbb{F}_4) = \bigcup_{l \in \check{P}} (l(\mathbb{F}_4) \cap C)$, which contradicts to the assumption $N_4(C) = 9$. Hence all \mathbb{F}_4 -points are nonsingular.

Next we show the additional statement. Since $\deg C = 3$ and $i(T_p(C), C; P) = 3$ for any $P \in C(\mathbb{F}_4)$, a line joining two \mathbb{F}_4 -points of C meets with C at the third \mathbb{F}_4 -point. Hence the 9 \mathbb{F}_4 -points $C(\mathbb{F}_4)$ can be divided three triples

$${P_{01}, P_{02}, P_{03}} {P_{11}, P_{12}, P_{13}} {P_{21}, P_{22}, P_{23}}$$

such that three points of each triple are collinear. Let l_i be the line on which P_{i1}, P_{i2} and P_{i3} lie, and $l_i(\mathbb{F}_4) = \{P_{i1}, P_{i2}, P_{i3}, Q_i, R_i\}$. Assume that l_0, l_1 and l_2 are concurrent. We may assume that $Q_0 = Q_1 = Q_2$. Hence $T_{P_{0\mu}} \cap l_1$ must be R_1 for any $\mu = 1, 2, 3$, and also $T_{P_{2\nu}} \cap l_1 = R_1$ for any $\nu = 1, 2, 3$ because $P_{i\alpha}$ $(i, \alpha = 1, 2, 3)$ are flexes. But, since the number of \mathbb{F}_4 -lines passing through R_1 is 5 and $T_{P_{i\mu}} \neq T_{P_{j\nu}}$ for any pair $(P_{i\mu}, P_{j\nu})$, it is impossible.

Therefore $\{l_0, l_1, l_2\}$ forms a frame of \mathbb{P}^2 , that is, we may assume that $l_i = \{X_i = 0\}$, where X_0, X_1, X_2 are coordinates of \mathbb{P}^2 . Let $C = \{F(X_0, X_1, X_2) = 0\}$. Since

$$F(X_0, X_1, 0) = c_2(X_0^3 + X_1^3) \quad (c_2 \in \mathbb{F}_4^*)$$

$$F(X_0, 0, X_2) = c_1(X_0^3 + X_2^3) \quad (c_1 \in \mathbb{F}_4^*)$$

$$F(0, X_1, X_2) = c_0(X_1^3 + X_2^3) \quad (c_0 \in \mathbb{F}_4^*),$$

we have $F(X_0, X_1, X_2) = c(X_0^3 + X_1^3 + X_2^3) + \gamma X_0 X_1 X_2$ with $c \in \mathbb{F}_4^*$ and $\gamma \in \mathbb{F}_4$. If $\gamma \neq 0$, then $(1, 1, \frac{c}{\gamma})$ is also an \mathbb{F}_4 -point of C, which is not any of the $P_{i,\alpha}$'s. This is a contradiction. Hence $\gamma = 0$, that is, C is Hermitian.

We close this section with a remark on plane curves that are used in the proof of the main theorem.

Lemma 2.4 (Segre) Let d be an integer with $1 \le d \le q+1$, and C be a curve of degree d in \mathbb{P}^2 defined over \mathbb{F}_q , which may have \mathbb{F}_q -linear components. Then $N_q(C) \le dq+1$, and equality holds if and only if C is a pencil of d \mathbb{F}_q -lines.

Proof. See Segre [11, II, $\S 6$ Observation IV] or Homma-Kim [5, Remark 1.2]. \square

3 An elementary bound

In this section, q is simply a power of p, that is, it need not be an even power of p. In [8], we established an upper bound for $N_q(X)$ of a hypersurface X over \mathbb{F}_q without \mathbb{F}_q -linear components, particularly a surface in \mathbb{P}^3 without \mathbb{F}_q -plane components

Theorem 3.1 Let S be a surface of degree d in \mathbb{P}^3 defined over \mathbb{F}_q without \mathbb{F}_q -plane components. Then

$$N_q(S) \le (d-1)q^2 + dq + 1.$$
 (6)

We refer the bound (6) as the elementary bound. When $d = \sqrt{q} + 1$, the Weil-Deligne bound (4) for n = 3 agrees with the elementary bound. In this section, we investigate the geometry of a surface in \mathbb{P}^3 whose number of \mathbb{F}_q -points achieves the bound (6). There are at least two examples other than the Hermitian surface each of which attains the equality in (6) 1 .

From now on, we keep the following setup until the end of this section.

Setup 3.2 Let d be an integer with $2 \le d \le q + 1$. Let S be a surface of degree d in \mathbb{P}^3 defined over \mathbb{F}_q without \mathbb{F}_q -plane components. Furthermore we suppose that $N_q(S)$ achieves the equality in (6).

¹ After wrote up the first draft of this paper, we proved that these two examples together with the Hermitian surface are only examples having this property [9], in which we referred this paper. Recently, Tironi [14] has settled the classification of hypersurfaces in \mathbb{P}^n that achieve the elementary bound stated in [8].

Lemma 3.3 The surface S contains an \mathbb{F}_q -line.

Proof. Suppose S does not contain any \mathbb{F}_q -lines. Let H be any \mathbb{F}_q -plane in \mathbb{P}^3 . Then $S \cap H$ is a plane curve of degree d over \mathbb{F}_q in $H = \mathbb{P}^2$, and has no \mathbb{F}_q -line as a component. Hence

$$N_q(S \cap H) \le \begin{cases} (d-1)q + 1 & \text{if } (d,q) \ne (4,4) \\ 14 & \text{if } (d,q) = (4,4) \end{cases}$$
 (7)

by Lemma 2.1. In a term of [4], defining the s-degree δ of $S(\mathbb{F}_q)$ by

$$\delta = \max\{N_q(S \cap H) \mid H \text{ is an } \mathbb{F}_q\text{-plane}\},$$

we have

$$N_q(S) \le (\delta - 1)q + 1 + \left| \frac{\delta - 1}{q + 1} \right|$$

by [4, Proposition 2.2]. Hence if $(d, q) \neq (4, 4)$, then

$$N_{q}(S) \leq (d-1)q^{2} + 1 + \left\lfloor \frac{(d-1)q}{q+1} \right\rfloor$$

$$= (d-1)q^{2} + 1 + \left\lfloor \frac{(d-2)(q+1) + q + 2 - d}{q+1} \right\rfloor$$

$$= (d-1)(q^{2} + 1);$$

and if (d,q)=(4,4), then $N_q(S)\leq 55$. In either case, $N_q(S)$ can't be $(d-1)q^2+dq+1$. \square

Definition 3.4 Let l_1, \ldots, l_d be \mathbb{F}_q -lines in \mathbb{P}^3 with $d \geq 2$. The union of those d-lines $Z = \bigcup_{i=1}^d l_i$ is called a planar \mathbb{F}_q -pencil of degree d if those d-lines lie on a plane simultaneously and $l_1 \cap \cdots \cap l_d \neq \emptyset$. The \mathbb{F}_q -point $\{v_Z\} = l_1 \cap \cdots \cap l_d$ is called the vertex of Z.

Notation 3.5 For an \mathbb{F}_q -line l of \mathbb{P}^3 , the set of \mathbb{F}_q -planes containing the line l is denoted by $\check{l}(\mathbb{F}_q)$.

Lemma 3.6 Let l be an \mathbb{F}_q -line on the surface S.

- (i) If an \mathbb{F}_q -plane H contains l, then $S \cap H$ is a planar \mathbb{F}_q -pencil of degree d.
- (ii) The map $\check{l}(\mathbb{F}_q) \ni H \mapsto v_{S \cap H} \in l(\mathbb{F}_q)$ is bijective.

Proof. (i) Since S has no \mathbb{F}_q -plane components, $S \cap H$ is a plane curve of degree d in H, and $N_q(S \cap H) \leq dq + 1$ by Lemma 2.4. Counting the cardinality of $S(\mathbb{F}_q)$ by the decomposition

$$S(\mathbb{F}_q) = \bigcup_{H \in \check{l}(\mathbb{F}_q)} \left((S \cap H)(\mathbb{F}_q) \setminus l(\mathbb{F}_q) \right) \bigcup l(\mathbb{F}_q),$$

we have

$$(d-1)q^2 + dq + 1 = N_q(S) \le (q+1)(dq+1-(q+1)) + (q+1)$$

= $(d-1)q^2 + dq + 1.$

Hence $N_q(S \cap H) = dq + 1$. So $S \cap H$ is a planar \mathbb{F}_q -pencil of degree d by the latter part of Lemma 2.4.

(ii) Let $\check{l}(\mathbb{F}_q) = \{H_1, H_2, \dots, H_{q+1}\}$. Since $S \cap H_i$ is a planar \mathbb{F}_q -pencil and has the line l as a component, we may set notations as

$$S \cap H_i = l \cup l_{i,1} \cup \cdots \cup l_{i,d-1}$$

and

$$v_i = v_{S \cap H_i}$$
.

It is obvious that $v_i \in l(\mathbb{F}_q)$. Since $\check{l}(\mathbb{F}_q)$ and $l(\mathbb{F}_q)$ have the same cardinality, it is enough to show that the map $\check{l}(\mathbb{F}_q) \ni H_i \mapsto v_i \in l(\mathbb{F}_q)$ is surjective.

Contrary, suppose this map is not surjective. Pick a point $Q \in l(\mathbb{F}_q) \setminus \{v_1, \dots, v_{q+1}\}$, and choose an \mathbb{F}_q -plane K such that $K \ni Q$ and $K \not\supset l$. Then we face two consequences:

- (α) $S \cap K$ does not contain any \mathbb{F}_q -line;
- (β) $N_q(S \cap K) \ge (d-1)q + d$,

as we verify them below. Since $S \cap K$ is a plane curve of degree d in K, those two conditions are incompatible each other by Lemma 2.1.

The verification of (α) . If $S \cap K$ contains an \mathbb{F}_q -line, then it must be a planar \mathbb{F}_q -pencil by (i). Since Q is a point of S, there is an \mathbb{F}_q -line m passing through Q among d lines of $S \cap K$. Hence l and m spans an \mathbb{F}_q -plane which is one of the H_i 's, say H_i . Then $v_i \in l \cap m = \{Q\}$, which contradicts to the choice of Q.

The verification of (β) . Let $Q_{i,j}$ be the intersection point of $l_{i,j}$ with K. Then the plane containing l and $Q_{i,j}$ is H_i and the line containing v_i and $Q_{i,j}$ is $l_{i,j}$. Hence $Q_{i,j} = Q_{i',j'}$ implies (i,j) = (i',j'). Since

$$(S \cap K)(\mathbb{F}_q) \supset \{Q_{i,j} \mid 1 \le i \le q+1, \ 1 \le j \le d-1\} \cup \{Q\},\$$

we have $N_q(S \cap K) \ge (d-1)(q+1) + 1$.

Lemma 3.7 Let H be an \mathbb{F}_q -plane such that $S \cap H$ is a planar \mathbb{F}_q -pencil of degree d. If an \mathbb{F}_q -line $l \subset S$ passes through $v_{S \cap H}$, then l is a component of $S \cap H$.

Proof. Since $d \geq 2$, there are two distinct components l_1 and l_2 of $S \cap H$. Suppose that l is not contained in H. Then l and l_i span an \mathbb{F}_q -plane, say H_i . Hence $S \cap H_i$ is also a planar \mathbb{F}_q -pencil of degree d by Lemma 3.6 (i). By the construction of H_1 and H_2 ,

$$v_{S \cap H_1} = l \cap l_1 = v_{S \cap H} = l \cap l_2 = v_{S \cap H_2},$$

which contradicts (ii) of Lemma 3.6. Hence l is contained in H, and hence it is a component of $S \cap H$.

Corollary 3.8 For any \mathbb{F}_q -point P of S, there is a unique \mathbb{F}_q -plane H such that $S \cap H$ is a planar \mathbb{F}_q -pencil of degree d with $v_{S \cap H} = P$.

Proof. From Lemma 3.3 and Lemma 3.6 (i), there is a line l on S containing P. Indeed the existence of an \mathbb{F}_q -line l' on S is guaranteed by Lemma 3.3; if the point P in question lies on l', there is nothing to do; when $P \notin l'$, consider the \mathbb{F}_q -plane H' containing l' and P, then $S \cap H'$ is a planar \mathbb{F}_q -pencil by Lemma 3.6 (i), which contains P; hence there is a line l on S containing P. Hence, by (ii) of Lemma 3.6 there is a desired \mathbb{F}_q -plane, and such a plane is uniquely determined by P.

4 A characterization of Hermitian surfaces

In this section, q is assumed to be an even power of a prime number p again, and S is a surface in \mathbb{P}^3 of degree $\sqrt{q} + 1$ defined over \mathbb{F}_q without \mathbb{F}_q -plane components such that $N_q(S) = (\sqrt{q}^3 + 1)(q+1)$. Note that this number $N_q(S)$ achieves both the Weil-Deligne bound (4) for n = 3 and the elementary bound (6).

Proposition 4.1 With the above situation, let H be an \mathbb{F}_q -plane of \mathbb{P}^3 . Then either

- (1) $S \cap H$ is a planar \mathbb{F}_q -pencil of degree $\sqrt{q} + 1$, or
- (2) $S \cap H$ is a nonsingular Hermitian curve of degree $\sqrt{q} + 1$ over \mathbb{F}_q .

Furthermore, let ν_1 denote the number of \mathbb{F}_q -planes having the property (1) above, and ν_2 the property (2). Then

$$\begin{cases} \nu_1 &= N_q(S) = (\sqrt{q^3} + 1)(q+1) \\ \nu_2 &= \sqrt{q^3}(q+1)(\sqrt{q} - 1). \end{cases}$$

Proof. Let $\check{\mathbb{P}}^3(\mathbb{F}_q)$ denote the set of \mathbb{F}_q -planes of \mathbb{P}^3 , and \mathcal{P}_1 the set of \mathbb{F}_q -planes having the property (1). The map $\mathcal{P}_1 \ni H \mapsto v_{S \cap H} \in S(\mathbb{F}_q)$ is bijective by Corollary 3.8. Hence $\nu_1 = N_q(S)$. Let $\mathcal{P}_2 = \check{\mathbb{P}}^3(\mathbb{F}_q) \setminus \mathcal{P}_1$. Note that if $H \in \mathcal{P}_2$, then $S \cap H$ has no \mathbb{F}_q -liner components by Lemma 3.6 (i). Consider the correspondence

$$\mathcal{A} = \{(P, H) \mid P \in H\} \subset S(\mathbb{F}_q) \times \check{\mathbb{P}}^3(\mathbb{F}_q)$$

with two projections $\pi_1: \mathcal{A} \to S(\mathbb{F}_q)$ and $\pi_2: \mathcal{A} \to \check{\mathbb{P}}^3(\mathbb{F}_q)$. Counting the cardinality of \mathcal{A} by π_1 , we have

$$^{\#}\mathcal{A} = N_q(S) \cdot (q^2 + q + 1).$$

If $H \in \mathcal{P}_1$, then $\#\pi_2^{-1}(H) = N_q(S \cap H) = \sqrt{q^3} + q + 1$, and if $H \in \mathcal{P}_2$, then $\#\pi_2^{-1}(H) = N_q(S \cap H) \leq \sqrt{q^3} + 1$ by (2.1). Hence, counting the cardinality of \mathcal{A} by π_2 , we have

$${}^{\#}\mathcal{A} \leq N_q(S) \cdot (\sqrt{q}^3 + q + 1) + \left(\frac{q^4 - 1}{q - 1} - N_q(S)\right) (\sqrt{q}^3 + 1)$$

= $N_q(S) \cdot (q^2 + q + 1)$.

Therefore, if $H \in \mathcal{P}_2$, then $N_q(S \cap H) = \sqrt{q^3} + 1$. Furthermore, when $H \in \mathcal{P}_2$, since $S \cap H$ has no \mathbb{F}_q -line as a component, $S \cap H$ is a nonsingular Hermitian curve by (1.1) if $q \neq 4$.

We need a little more argument if q=4. We want to apply Lemma 2.3. If an \mathbb{F}_4 -point is not a flex of $S\cap H$, i.e., $i(T_P(S\cap H).S\cap H;P)=2$, then $T_P(S\cap H)$ meets with $S\cap H$ at exactly two \mathbb{F}_4 -points, because $\deg S\cap H=3$. Consider all \mathbb{F}_4 -planes $\{H_\lambda\}$ in \mathbb{P}^3 such that $H_\lambda\supset T_P(S\cap H_\lambda)$. Let $a=\#\{\lambda\mid H_\lambda\in \mathcal{P}_1\}$. Then $\#\{\lambda\mid H_\lambda\in \mathcal{P}_2\}=5-a$. Since $N_4(S\cap H_\lambda)$ is either 13 or 9, we have 45=(13-2)a+(9-2)(5-a)+2. Hence there is an \mathbb{F}_4 -plane H_λ such that $S\cap H_\lambda$ is a planar \mathbb{F}_4 -pencil of degree 3, and the line $T_P(S\cap H)$ lies on H_λ . But it is impossible, because any \mathbb{F}_4 -line on H_λ meets with the planar pencil at either 1, or 3, or 5 points.

Last Step of the Proof of Theorem 1.2. From Proposition 4.1, there is an \mathbb{F}_q -plane $H_{\infty} \subset \mathbb{P}^3$ such that $S \cap H_{\infty}$ is a nonsingular Hermitian curve. Choose a system of homogeneous coordinates X_0, X_1, X_2, X_3 of \mathbb{P}^3 such that

- (i) H_{∞} is given by $X_0 = 0$; and
- (ii) the plane curve $S \cap H_{\infty}$ in $H_{\infty} = \mathbb{P}^2$ is given by

$$\bar{X}_1 \bar{X}_2^{\sqrt{q}} + \bar{X}_1^{\sqrt{q}} \bar{X}_2 + \bar{X}_3^{\sqrt{q}+1} = 0,$$

where $\bar{X}_1, \bar{X}_2, \bar{X}_3$ are coordinates on H_∞ induced by X_1, X_2, X_3 respectively.

Let $P_1=(0,0,1,0)$ and $P_2=(0,1,0,0)$, both of which are points on $S\cap H_{\infty}$. For $\alpha=1,2$, the tangent line L_{α} at P_{α} to $S\cap H_{\infty}$ in $H_{\infty}=\mathbb{P}^2$ is given by $\bar{X}_{\alpha}=0$. It is easy to see that $S\cap L_{\alpha}=\{P_{\alpha}\}$. Hence for an \mathbb{F}_q -plane $H\supset L_{\alpha}$,

$${}^{\#}(S(\mathbb{F}_q)\cap (H\setminus L_{\alpha})) = \left\{ \begin{array}{ll} \sqrt{q^3} + 1 - 1 & \text{if } S\cap H \text{ is Hermitian} \\ \sqrt{q^3} + q + 1 - 1 & \text{if } S\cap H \text{ is an } \mathbb{F}_q\text{-pencil.} \end{array} \right.$$

Counting the number $N_q(S)-1$ by using all \mathbb{F}_q -planes containing L_{α} , we know that there is a unique plane $H_{\alpha,0}\supset L_{\alpha}$ such that $S\cap H_{\alpha,0}$ is a planar \mathbb{F}_q -pencil of degree $\sqrt{q}+1$; and $S\cap H_{\alpha,\lambda}$ is a nonsingular Hermitian curve for other plane $H_{\alpha,\lambda}\supset L_{\alpha}$. By changing coordinates of type

$$\begin{cases} X_1 & \mapsto & X_1 + aX_0 \\ X_2 & \mapsto & X_2 + bX_0 \end{cases}$$

if necessary, we may suppose that $H_{\alpha,0}$ is defined by $X_{\alpha}=0$ for $\alpha=1$ and 2 respectively. But the situation on H_{∞} never change.

Summing up, S is defined by

$$F(X_0, \dots, X_3) = X_0 f(X_0, \dots, X_3) + h(X_1, X_2, X_3) = 0$$

where

$$h(X_1, X_2, X_3) = X_1 X_2^{\sqrt{q}} + X_1^{\sqrt{q}} X_2 + X_3^{\sqrt{q}+1}$$

and $\deg f = \sqrt{q}$.

Since $S \cap H_{1,0} = S \cap \{X_1 = 0\}$ is a planar \mathbb{F}_q -pencil with the vertex $P_1 = (0,0,1,0)$,

$$F(X_0, 0, X_2, X_3) = X_0 f(X_0, 0, X_2, X_3) + X_3^{\sqrt{q}+1}$$

does not contain X_2 , because this polynomial must have the form $c \prod_j (X_0 + \gamma_j X_3)$, with $c \in \mathbb{F}_q^*$. Hence, in $F(X_0, X_1, X_2, X_3)$, any monomial containing X_2 also contains X_1 . By the same argument on $S \cap H_{2,0} = S \cap \{X_2 = 0\}$, any monomial containing X_1 also contains X_2 . Therefore $f(X_0, X_1, X_2, X_3)$ is written as

$$f(X_0, X_1, X_2, X_3) = g_1(X_0, X_3) + g_2(X_0, \dots, X_3)X_1X_2,$$

where $\deg g_1 = \sqrt{q}$ and $\deg g_2 = \sqrt{q} - 2$.

For the plane $H_{1,\lambda} = \{X_0 = \lambda X_1\}$ $(\lambda \in \mathbb{F}_q^*)$, since $S \cap \{X_0 = \lambda X_1\}$ is a Hermitian curve,

$$\rho F(\lambda X_1, X_1, X_2, X_3) = \rho(\lambda X_1(g_1(\lambda X_1, X_3) + g_2(\lambda X_1, X_1, X_2, X_3)X_1X_2) + h(X_1, X_2, X_3))$$
(8)

is a Hermitian polynomial for some $\rho \in \mathbb{F}_q^*$. Since $X_3^{\sqrt{q}+1}$ appears only in $h(X_1, X_2, X_3)$, the constant ρ must be an element of $\mathbb{F}_{\sqrt{q}}^*$. Hence (8) is a Hermitian polynomial even if $\rho = 1$. Since h itself Hermitian,

$$\lambda g_1(\lambda X_1, X_3)X_1 + \lambda g_2(\lambda X_1, X_1, X_2, X_3)X_1^2X_2$$

must be Hermitian.

Any monomial in $g_2(\lambda X_1, X_1, X_2, X_3)X_1^2X_2$ never appear in $g_1(\lambda X_1, X_3)X_1$. So $g_2(\lambda X_1, X_1, X_2, X_3))X_1^2X_2$ contains the monomials only of types $X_i^{\sqrt{q}+1}$ or $X_i^{\sqrt{q}}X_j$. Hence

$$g_2(\lambda X_1, X_1, X_2, X_3)X_1^2 X_2 = \mu X_1^{\sqrt{q}} X_2 \ (\mu \in \mathbb{F}_q).$$

But the monomial $X_1X_2^{\sqrt{q}+1}$ can't appear in

$$g_1(\lambda X_1, X_3)X_1 + g_2(\lambda X_1, X_1, X_2, X_3)X_1^2X_2.$$

Hence $g_2(\lambda X_1, X_1, X_2, X_3) = 0$, that is, $X_0 - \lambda X_1$ is a factor of $g_2(X_0, X_1, X_2, X_3)$ for any $\lambda \in \mathbb{F}_q^*$. But $\deg g_2 = \sqrt{q} - 2$ (q - 1), it is impossible. Therefore $g_2(X_0, X_1, X_2, X_3) = 0$ and $\lambda g_1(\lambda X_1, X_3) X_1$ is Hermitian.

$$g_1(X_0, X_3) = \sum_{i=0}^{\sqrt{q}} a_i X_0^i X_3^{\sqrt{q}-i}.$$

Then

$$\lambda g_1(\lambda X_1, X_3) X_1 = \sum_{i=0}^{\sqrt{q}} a_i \lambda^{i+1} X_1^{i+1} X_3^{\sqrt{q}-i},$$

which is Hermitian. Hence $a_i = 0$ for $i \neq 0$, $\sqrt{q} - 1$, \sqrt{q} ; and also $a_0^{\sqrt{q}} = a_{\sqrt{q}-1}$ and $a_{\sqrt{q}} \in \mathbb{F}_{\sqrt{q}}$, that is,

$$g_1(X_0, X_3) = a_0 X_3^{\sqrt{q}} + a_0^{\sqrt{q}} X_0^{\sqrt{q}-1} X_3 + a_{\sqrt{q}} X_0^{\sqrt{q}} \text{ with } a_{\sqrt{q}} \in \mathbb{F}_{\sqrt{q}}.$$

Hence

$$F(X_0, \dots, X_3) = a_0 X_0 X_3^{\sqrt{q}} + a_0^{\sqrt{q}} X_0^{\sqrt{q}} X_3 + a_{\sqrt{q}} X_0^{\sqrt{q}+1} + h(X_1, X_2, X_3),$$

which is Hermitian. If $a_{\sqrt{q}} = 0$, then the surface is a cone of a nonsingular Hermitian curve with vertex (1,0,0,0), and then $N_q(S) = (\sqrt{q}^3 + 1)q + 1$, which is not the given number. So $a_{\sqrt{q}} \in \mathbb{F}_{\sqrt{q}}^*$, and hence S is a nonsingular Hermitian surface. \square

Appendix

In this appendix, we consider a possibility of generalizing Rück-Stichtenoth's characterization of the Hermitian curve to the Hermitian surface.

Theorem A.1 (Rück-Stichtenoth) Let C be a nonsingular curve of genus $g = \frac{1}{2}\sqrt{q}(\sqrt{q}-1)$ defined over \mathbb{F}_q . If $N_q(C) = \sqrt{q}^3 + 1$, then the function field $\mathbb{F}_q(C)$ of C coincides with that of the Hermitian curve over \mathbb{F}_q .

Remark A.2 If one considers only nonsingular plane curves, Rück-Stichtenoth's characterization implies Hirschfeld et al.'s, because the linear system $g_{\sqrt{q}+1}^2$ is unique if $g = \frac{1}{2}\sqrt{q}(\sqrt{q}-1) > 1$. When q = 4, the genus in question is 1. As was mentioned in [3], the plane curve $\{X_0 + \omega X_1 + \omega^2 X_2 = 0\}$ over \mathbb{F}_4 is not Hermitian, but has 9 \mathbb{F}_4 -points, where $\mathbb{F}_4 = \{0, 1, \omega, \omega^2\}$. However the function field of this curve over \mathbb{F}_4 coincides with that of the Hermitian curve over \mathbb{F}_4 .

As has been remarked above, the Rück-Stichtenoth theorem gives a characterization of the Hermitian curve in wider family than the plane curves if one considers only nonsingular curves, namely among the family of nonsingular curves with the same "topological" invariant as the Hermitian curve. Since the number \sqrt{q}^3+1 achieves the Weil bound for $g=\frac{1}{2}\sqrt{q}(\sqrt{q}-1)$, the zeta function of the nonsingular curve C in (A.1) is $\frac{(1+\sqrt{q}t)^{2g}}{(1-t)(1-qt)}$. So Theorem A.1 can be interpreted as:

Theorem A.3 (Rück-Stichtenoth) Let C be a nonsingular curve of genus $g = \frac{1}{2}\sqrt{q}(\sqrt{q}-1) > 1$ defined over \mathbb{F}_q whose zeta function is $\frac{(1+\sqrt{q}t)^{2g}}{(1-t)(1-qt)}$. Then C is the Hermitian curve.

On the other hand, the number of \mathbb{F}_q -points of the Hermitian surface attains the Weil-Deligne bound, and its zeta function is:

$$\frac{1}{(1-t)(1-qt)^{b_2}(1-q^2t)} \qquad (b_2 = q\sqrt{q} - q + \sqrt{q} + 1). \tag{9}$$

It is natural to ask whether a nonsingular surface over \mathbb{F}_q with the zeta function (9) is Hermitian or not. The example below shows the answer of this question is negative.

From now on, when X is a nonsingular surface over \mathbb{F}_q , $Z_X(t)$ denotes the zeta function of X.

Lemma A.4 Let \tilde{X} be a blowing-up of a nonsingular surface X over \mathbb{F}_q with center an \mathbb{F}_q -point of X. Then $Z_{\tilde{X}}(t) = Z_X(t) \frac{1}{(1-qt)}$.

Proof. Let $N_m(X)$ and $N_m(\tilde{X})$ denote the number of \mathbb{F}_{q^m} -points of X and \tilde{X} respectively. Then $N_m(\tilde{X}) = N_m(X) + q^m$. Hence

$$Z_{\tilde{X}}(t) = \exp\left(\sum_{m=1}^{\infty} (N_m(X) + q^m) \frac{t^m}{m}\right)$$
$$= \exp\left(\sum_{m=1}^{\infty} N_m(X) \frac{t^m}{m}\right) \exp\left(\sum_{m=1}^{\infty} q^m \frac{t^m}{m}\right)$$
$$= Z_X(t) \frac{1}{(1 - qt)}.$$

Example A.5 Let $X_1 = \mathbb{P}^2$ over \mathbb{F}_q . It is obvious that $Z_{X_1}(t) = \frac{1}{(1-t)(1-qt)(1-q^2t)}$. Define a nonsingular surface X_n over \mathbb{F}_q by a blowing-up of X_{n-1} with center an \mathbb{F}_q -point, successively. Then the zeta function of X_{b_2} is the same as Hermitian surface's. However, if $\sqrt{q}+1 \geq 4$, the Hermitian surface of degree $\sqrt{q}+1$ is not rational. So, if $q>2^2$, then the function field of X_{b_2} does not coincides with that of the Hermitian surface over \mathbb{F}_q .

References

- [1] R. C. Bose and I. M. Chakravarti, Hermitian varieties in a finite projective space $PG(N, q^2)$, Canad. J. Math. 18 (1966) 1161–1182.
- [2] P. Deligne, La conjecture de Weil, I, Inst. Hautes Études Sci. Publ. Math. 43 (1974), 273–307.
- [3] J. W. P. Hirschfeld, L. Storme, J. A. Thas and J. F. A. Voloch, A characterization of Hermitian curves, J. Geom. 41 (1991) 72–78.
- [4] M. Homma, A bound on the number of points of a curve in a projective space over a finite field, in: M. Lavrauw, G.L. Mullen, S. Nikova, D. Panario, and L. Storme (Eds.), Theory and Applications of Finite Fields, 103–110 Contemp. Math., vol. 579, AMS, Providence, 2012.
- [5] M. Homma and S. J. Kim, Around Sziklai's conjecture on the number of points of a plane curve over a finite field, Finite Fields Appl. 15 (2009), 468-474.

- [6] M. Homma and S. J. Kim, Sziklai's conjecture on the number of points of a plane curve over a finite field II, in: G. McGuire, G.L. Mullen, D. Panario, I.E. Shparlinski (Eds.), Finite Fields: Theory and Applications, 225–234, Contemp. Math., vol. 518, AMS, Providence, 2010. (An update is available at arXiv:0907.1325v2.)
- [7] M. Homma and S. J. Kim, Sziklai's conjecture on the number of points of a plane curve over a finite field III, Finite Fields Appl. 16 (2010) 315–319.
- [8] M. Homma and S. J. Kim, An elementary bound for the number of points of a hypersurface over a finite field, Finite Fields Appl. 20 (2013) 76–83.
- [9] M. Homma and S. J. Kim, Numbers of points of surfaces in the projective 3-space over finite fields, Finite Fields Appl. 35 (2015) 52–60 (A previous version is available at arXiv:1409.5842v1).
- [10] H.-G. Rück and H. Stichtenoth, A characterization of Hermitian function fields over finite fields, J. Reine Angew. Math. 457 (1994) 185–188.
- [11] B. Segre, Le geometrie di Galois, Ann. Mat. Pura Appl. (4) 48 (1959) 1–96.
- [12] A. B. Sørensen, On the number of rational points on codimension-1 algebraic sets in $\mathbf{P}^n(\mathbf{F}_q)$, Discrete Math. 135 (1994), 321–334.
- [13] P. Sziklai, A bound on the number of points of a plane curve, Finite Fields Appl. 14 (2008) 41–43.
- [14] A. L. Tironi, Hypersurfaces achieving the Homma-Kim bound, preprint, 2014, available at arXiv: 1410.7320v2
- [15] A. Weil, Numbers of solutions of equations in finite fields, Bull. Amer. Math. Soc. 55 (1949) 497–508.