arXiv:1304.0516v1 [physics.optics] 2 Apr 2013

Unusual Brownian motion of photons in open absorbing media

Li-Yi Zhao¹, Chu-Shun Tian¹, Zhao-Qing Zhang² and Xiang-Dong Zhang³

¹Institute for Advanced Study, Tsinghua University, Beijing 100084, China

²Department of Physics and William Mong Institute of Nano Science and Technology,

Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong

³School of Physics, Beijing Institute of Technology, Beijing 100081, China

Very recent experiments have discovered that localized light in strongly absorbing media displays intriguing diffusive phenomena. Here we develop a first-principles theory of light propagation in open media with arbitrary absorption strength and sample length. We show analytically that photons in localized open absorbing media exhibit unusual Brownian motion. Specifically, wave transport follows the diffusion equation with the diffusion coefficient exhibiting spatial resolution. Most strikingly, despite that the system is controlled by two parameters – the ratio of the localization (absorption) length to the sample length – the spatially resolved diffusion coefficient displays novel single parameter scaling: it depends on the space via the returning probability. Our analytic predictions for this diffusion coefficient are confirmed by numerical simulations. In the strong absorption limit they agree well with the experimental results.

PACS numbers: 42.25.Dd,71.23.An

Many complex systems exhibit fundamentally different physics in various scales. A canonical example, thoroughly studied by Einstein in one of his revolutionary papers in 1905 [1], is the motion of pollens suspended in water (the Brownian motion). In short time scales, pollens undergo bombardment by water molecules and exhibits random motion. The resulting complexity of microscopic dynamics notwithstanding, in large time scales emerges substantially simpler physics: the evolution of the pollen concentration profile fully characterizes the dynamics and is governed by a diffusion equation. Such emergent diffusive phenomenon is universal, and microscopic dynamics only affect the diffusivity namely the value of the diffusion coefficient.

The classical electromagnetic wave in infinite random media is the very system where universal macrosocpic diffusive phenomena emerge from complicated microscopic dynamics. In short (time) scales, the system's physics is dominated by light scattering off random dielectric field described by the Maxwell equation. In scales larger than the elastic mean free time, as noticed by astrophysicists long time ago [2], light propagation may be viewed as random motion of Brownian particles, the photons. Yet, light displays the particle-wave duality, and the wave nature has far-reaching consequences. Specifically, wave interference suppresses the diffusivity [3]. In the presence of strong dielectric fluctuations, the (bare) diffusion coefficient is strongly renormalized and vanishes in a much larger time scale. This is a hallmark of Anderson localization of light (see Ref. [4] for a review), and the corresponding time scale is called the localization time.

In realistic experiments or photonic devices the medium is open: photons escape the medium from the interfaces. Therefore, diffusivity of waves cannot be fully suppressed even disorders are strong. A fundamental question thereby is: do localized waves in open media exhibit certain macroscopic diffusive phenomena? This is a long-standing issue (see Ref. [5] for a recent review). Conceptually, a prominent difficulty arises from the following. On the one hand long-time transport of localized waves in open media is dominated by rare long-lived, high transmission (resonant) states [6, 7], while on the other hand macroscopic diffusion of waves - if exists - characterizes statistical behavior of wave propagation: these are two seemingly contradictory facets. In fact, experiments and numerical simulations [8-10] have shown that due to the failure of capturing resonant states the prevailing macroscopic diffusion model developed in Ref. [11] cannot describe transport of strongly localized waves in open media. This had inspired the opinion that the resonant state and macroscopic diffusion of photons are two incompatible concepts.

Contrary to this common belief, the results achieved recently [5, 9] show that the rare resonant states do not wash out macroscopic diffusion of photons. Rather, they are reconciled in the way that the spatial resolution of the diffusion coefficient exhibits a novel scaling. Specifically, the diffusion coefficient depends on the space via the returning probability. (One may show – both analytically and numerically [9] – that this novel scaling is completely missed in the prevailing model [11] due to the unjustified self-consistent treatments employed there.) This novel scaling was analytically found by using a first-principles theory [9, 12] and has been fully confirmed by numerical simulations [9].

So far we have focused on passive media. Yet, absorption inevitably exists in realistic environments [13]. In open media the strong interplay between absorption and localization has been experimentally shown to cause surprising phenomena [14], and may even find practical applications in optical devices [15]. Very recently, both realistic and numerical experiments have observed that localized waves exhibit unusual macroscopic diffusion in open absorbing media [16, 17]. However, these experiments (both realistic and numerical) focus on short samples and strong absorption. For such systems resonant states are unimportant. In the opposite limit, where samples are sufficiently long and absorption is weak, resonant states dominate transport of localized waves. As these states are very sensitive to absorption due to long lifetimes [18], a very fundamental issue arises: how do absorption interplay with macroscopic diffusion of localized waves in open media? The purpose of this Letter is to present the first systematic first-principles study of this issue for arbitrary sample length and absorption strength.

Main results and experimental relevance. — Specifically, we consider propagation of light in quasi-onedimensional open (uniformly) absorbing media of length L. This system is controlled by two dimensionless parameters, ξ/L and ξ_a/L , where ξ (ξ_a) is the localization (absorption) length. We find that – even in localized samples ($\xi/L \ll 1$) – photons exhibit macroscopic diffusion irrespective of the absorption rate γ . More precisely, in the presence of a steady monochromatic light source located at x' the wave intensity (namely energy density) profile, $\mathcal{Y}(x, x')$, obeys

$$\{\gamma - \partial_x D(x)\partial_x\} \mathcal{Y}(x, x') = \delta(x - x').$$
(1)

It resembles the normal diffusion equation, but, the diffusion coefficient exhibits a number of anomalies. First of all, it has a spatial resolution, D(x), decreasing monotonically from the air-medium interface to the sample midpoint and varying over several orders. Most strikingly, despite that the system is controlled by two parameters, D(x) exhibits a novel single parameter scaling,

$$D(x)/D(0) = D_{\infty}(\lambda(x)) \tag{2}$$

similar to the passive medium case. Here, the scaling factor $\lambda(x)$ is proportional to the (static) returning probability which depends on x/L as well as ξ/L and ξ_a/L , and the scaling function $D_{\infty}(\lambda)$ is the same as that of passive media. (The explicit form of both $\lambda(x)$ and $D_{\infty}(\lambda)$ will be given below.) Bearing these anomalies and the analogy to the normal diffusion equation, Eq. (1) describes 'unusual Brownian motion' of photons. Furthermore, from Eq. (2) we are able to predict analytically the spatial resolution of the diffusion coefficient which is confirmed by numerical experiments. We stress that our analytic results are very general: they are valid for arbitrary sample length and absorption strength and, additionally, for both time-reversal (orthogonal symmetry) and broken time-reversal (unitary symmetry) systems.

In the case of strong absorption, our analytic results are simplified and fully agree with the recent experimental findings shown in Refs. [16, 17]. Specifically, if absorption is strong such that $\xi_a/L \ll 1$, we find that D(x) exhibits a plateau in the regime $\min(x, L-x) \gtrsim \xi_a$. Moreover, the plateau value is determined by single parameter, $\xi_a/\xi,$

$$D(L/2)/D(0) = D_{\infty}(\xi_a/2\xi)$$
 (3)

(cf. the first line of Table I and II). We stress that the plateau as well as Eq. (3) exists only if $\xi_a/L \ll 1$.

Origin of unusual Brownian motion. — We begin with a physical explanation of the main results. Since closer to the interface more easily escape photons from the medium (cf. dashed lines of Fig. 1), the returning probability, $\mathcal{Y}_0(x, x)$, is inhomogeneous in space (i.e., depending on x as well as the parameters ξ/L and ξ_a/L). As such, wave interference effects are inhomogeneous also. Near the interface (deep inside the sample) they are weak (strong). For diffusive samples ($\xi/L \gg 1$ where $\xi \sim \pi \nu D_0$ [19] with ν the photon density of states), $\mathcal{Y}_0(x, x)$ gives rise to an inhomogeneous (one-loop) weak localization correction $\sim \mathcal{O}(\mathcal{Y}_0(x, x)/(\pi\nu)) \sim \mathcal{O}(\lambda(x))$.



FIG. 1: Examples of wave interference between two optical paths (red and blue) leading to the novel scaling behavior. The paths trace the three loops with different orders $(B \rightarrow A \rightarrow C \text{ for the red and } C \rightarrow B \rightarrow A \text{ for the blue})$. Notice that these loops may not be complete (e.g., dashed parts of A and C) due to photon leakage.

For localized samples $(\xi/L \ll 1)$, as waves penetrate deeply into the sample photons easily return to its departure point. This causes more complicated wave interference essential to strong localization physics. For example, two optical paths may take the same n(=3 inFig. 1) loops as their routes and trace them with different orders. Therefore, they constructively interfere with each other. This leads to a weak localization correction $\sim \mathcal{O}(\lambda^n(x))$. Therefore, the spatial resolution D(x) depends on x/L (as well as ξ/L and ξ_a/L) via the factor $\lambda(x)$, justifying Eq. (2).

Importantly, the optical paths in Fig. 1 may propagate in the same direction – clockwise or counterclockwise – during tracing each loop, suggesting that interference picture beyond what leads to usual one-loop weak localization does not necessarily require time-reversal symmetry. This is crucial to localization physics of systems with unitary symmetry (which is completely beyond the reach of the self-consistent theory [11].) Therefore, the novel single parameter scaling is an intrinsic wave phenomenon unrelated to time-reversal symmetry.

First-principles theory. — The derivations are largely parallel to those of passive media [9, 12]. Therefore, we shall outline the key steps below with an emphasis on the key differences, and refer the readers to Ref. [5]

for technical details. We first introduce the microscopic formalism valid for arbitrary dimensions. Consider the point-like source located at \mathbf{r}' inside the medium with the spectral decomposition $J_{\omega}(\mathbf{r}')$ (ω being the wave angular frequency). The time-integrated wave intensity – upon disorder averaging – is given by [20] $I(\mathbf{r}) = \int \frac{d\omega}{2\pi} \mathcal{Y}(\mathbf{r},\mathbf{r}') |J_{\omega}(\mathbf{r}')|^2$. Here the (static) spatial correlation function $\mathcal{Y}(\mathbf{r},\mathbf{r}') = \langle G^A_{\omega^2}(\mathbf{r},\mathbf{r}') G^R_{\omega^2}(\mathbf{r}',\mathbf{r}) \rangle$, with $\langle \cdots \rangle$ standing for the disorder average and the advanced (retarded) Green function $G^A(G^R)$ defined as

$$[\nabla^2 + \omega^2 (1 + \epsilon(\mathbf{r}) \pm i\epsilon'')] G^{R,A}_{\omega^2}(\mathbf{r},\mathbf{r}') = \delta(\mathbf{r} - \mathbf{r}'), \quad (4)$$

where the dielectric fluctuation, $\epsilon(\mathbf{r})$, is gaussian, $\epsilon'' > 0$ causes uniform absorption, and the light velocity in the air is set to unity.

Then, it is a standard procedure to cast the above spatial correlation function in terms of the functional integral over a supermatrix field $Q(\mathbf{r})$,

$$\mathcal{Y}(\mathbf{r},\mathbf{r}') = \left(\frac{\pi\nu}{8\omega}\right)^2 \int D[Q] \mathrm{str}(A_+Q(\mathbf{r})A_-Q(\mathbf{r}'))e^{-F[Q]}.(5)$$

Here A_{\pm} are some constant supermatrices and 'str' the supertrace [5, 19]. The action,

$$F[Q] = \frac{\pi\nu}{8} \int d\mathbf{r} \operatorname{str} \left\{ D_0 (\nabla Q)^2 - 2\gamma \Lambda Q \right\}, \qquad (6)$$

differs from the passive one [5] in the second term accounting for the absorption. Notice that the (bare) diffusion coefficient $D_0 \equiv D(0)$ and for $\epsilon'' \ll 1$ the absorption rate $\gamma = \omega \epsilon''$ [21]. The supermatrix $Q = T^{-1}\Lambda T$ where Λ is the so-called metallic saddle point and T takes the value from the coset space of $UOSP(2,2|4)/UOSP(2|2) \otimes UOSP(2|2)$ for orthogonal symmetry and of $U(1,1|2)/U(1|1) \otimes U(1|1)$ for unitary symmetry. It is very important that if the air-medium interface is transparent, the supermatrix field is fixed to be Λ at the interface. This reflects that on the interface there is no photon accumulation [5, 9, 12].

In quasi one dimension the microscopic formalism is simplified. Specifically, the supermatrix Q field is homogeneous in the transverse plane. F[Q] reduces to $\frac{\pi\nu}{8} \int_0^L dx \operatorname{str} \{D_0(\partial_x Q)^2 - 2\gamma \Lambda Q\}, (\nu \text{ includes a factor of}$ the cross section area.) and the boundary condition to

$$Q(x=0) = Q(x=L) = \Lambda.$$
(7)

As a result, \mathcal{Y} depend only on the longitudinal coordinate, x. Then, we follow the procedures of Refs. [5, 12] to calculate $\mathcal{Y}(x, x')$ explicitly. We find that $\mathcal{Y}(x, x')$ satisfies Eq. (1) with the boundary condition: $\mathcal{Y}(x = 0, x') =$ $\mathcal{Y}(x = L, x') = 0$. Furthermore, we find that the diffusion coefficient has a spatial resolution D(x), and the latter is a functional of the factor $\lambda(x) = \mathcal{Y}_0(x, x' = x)/(\pi\nu)$. This justifies Eq. (2). The correlator $\mathcal{Y}_0(x, x')$ solves the normal diffusion equation: $(\gamma - D_0\partial_x^2)\mathcal{Y}_0(x, x') =$ $\delta(x-x')$ implemented by the boundary condition $\mathcal{Y}_0(x = 0, x') = \mathcal{Y}_0(x = L, x') = 0$. Solving this equation we obtain

$$\lambda(x) = \frac{\xi_a}{2\xi} \frac{\cosh(L/\xi_a) - \cosh((L-2x)/\xi_a)}{\sinh(L/\xi_a)}, \qquad (8)$$

where the diffusive absorption length $\xi_a \equiv \sqrt{D_0/\gamma}$.

For $\lambda \ll 1$ we find that the perturbative expansion of the scaling function $D_{\infty}(\lambda)$ is identical to that at $\gamma = 0$. This implies that $D_{\infty}(\lambda)$ is the same as the one at $\gamma = 0$. On the other hand, at $\gamma = 0$ the scaling function has been found analytically and fully confirmed by numerical experiments [9], which has the following asymptotic form:

$$D_{\infty}(\lambda) \sim \begin{cases} 1 + c_1 \lambda + c_2 \lambda^2 + \cdots, & \lambda \ll 1, \\ e^{-\lambda}, & \lambda \gtrsim 1. \end{cases}$$
(9)

Notice that the coefficients c_i depend on the system's symmetry. In particular, for orthogonal symmetry $c_1 < 0$ while for unitary symmetry $c_1 = 0$, $c_2 < 0$.

Eqs. (1), (2), (8) and (9) constitute a complete description of unusual Brownian motion of photons in open absorbing media. In essence, it is a macroscopic phenomenon emerging from the microscopic Helmholtz equation (4) in large time scales. The present macroscopic theory differs from the previous one for passive media [5, 9] in the absorption term and that the spatial resolution D(x) depends also on a new parameter ξ_a/L . Below we provide numerical evidence of this intriguing phenomenon.



FIG. 2: In localized open absorbing media the spatial resolution of the diffusion coefficient exhibits novel single parameter scaling. Analytic predictions for D(x) (solid lines) are in good agreement with the simulation results (circles). From bottom to top the absorption strength increases (see the text for the value of ϵ'' as well as other parameters).

Evidence from numerical experiments. — We perform numerical simulations of the (microscopic) Helmholtz equation. To this end we prepare a randomly layered medium where the analytic results remain valid. The relative permittivity in each layer is a random number with a real part uniformly distributed in the interval [0.3, 1.7]. The imaginary part, ϵ'' , is assumed to be a constant. The random medium is embedded in the air background so that there exist no internal reflections when ϵ'' vanishes. We launch a plane wave of frequency $\omega = 1.65a^{-1}$ into the system, where the sample length L = 200a with athe layer thickness, and use the standard transfer matrix method to calculate the wave intensity profile, $I_{\beta}(x)$, for each dielectric disorder configuration β . For a given value of ϵ'' , we calculate the ensemble-averaged wave intensity profile, $I(x) \equiv \langle I_{\beta}(x) \rangle$, of 5 000 000 disorder configurations. Then, by presuming Eq. (1), which gives

$$D(x) = \frac{-\gamma \int_x^L I(x')dx' + D(0)\partial_x|_{x=L}I(x)}{\partial_x I(x)}, \qquad (10)$$

we obtain numerical results of D(x) from the measurements of I(x).

The results of simulations are shown in Fig. 2 for $\epsilon'' = 0, 1 \times 10^{-4}, 5 \times 10^{-4}, 1 \times 10^{-3}$, and 2×10^{-3} (from bottom to top). First of all, in the passive limit ($\epsilon'' = 0$, the bottom curve) we recover the result of Ref. [9] for deeply localized (Indeed, in this limit $L/\xi = 200a/22a \approx 9$ is large.) samples. The effects of absorption become significant when $\xi_a/L \sim \mathcal{O}(1)$, i.e., $\epsilon'' \approx \xi/(L^2\omega) \approx 3 \times 10^{-4}$. Having this estimation in mind, we may adjust the value of ϵ'' to systematically explore the effects of absorption. The results are compared with the analytic predictions obtained from the novel single parameter scaling theory namely Eqs. (2), (8) and (9). As shown in Fig. 2, they are in good agreement. Now we turn to analytic discussions of the behavior of D(x) based on the developed first-principles theory.

Behavior of D(x) in semi-infinite samples. — In this case $L \rightarrow \infty$, the scaling factor (8) is simplified to $\lambda(x) = \frac{\xi_a}{2\xi} (1 - e^{-2x/\xi_a})$. If the absorption is weak such that $\xi/\xi_a \ll 1$, then $\lambda \ll 1$ for $x \ll \xi$ and $\lambda \gtrsim 1$ for $x \gtrsim \xi$. Applying the first and second line of Eq. (9), respectively, we find distinct behavior of $D(x)/D_0$ in various regimes, which is summarized in Table I. The first line indicates that D(x) reaches an exponentially small residual value at $x \gtrsim \xi_a/2$. This reflects that deep inside the medium the ordinary Brownian motion of photons is restored, albeit with dramatically smaller diffusion coefficient. Qualitatively, the plateau arises from that due to absorption photons could not propagate for a distance larger than ξ_a and as such, photons deep inside the medium could not 'see' the air-medium interface. Indeed, similar results were achieved by using the scaling theory of light localization in infinite absorbing media long time ago [3]. The second and third lines indicate an intermediate scale, $\sqrt{\xi_a\xi}$, between ξ and ξ . For x smaller than this scale the local diffusion coefficient behaves essentially the same as that of (semi-infinite) passive media (see the third and

TABLE I: Behavior of D(x) at $\xi \ll \xi_a \ll L$

regime	D(x)/D(0)
$x \gtrsim \xi_a/2$	$e^{-\xi_a/(2\xi)}$
$\sqrt{\xi_a \xi} \ll x \ll \xi_a/2$	$e^{x^2/(\xi_a\xi)}e^{-x/\xi}$
$\xi \lesssim x \ll \sqrt{\xi_a \xi}$	$e^{-x/\xi}$
$x \ll \xi$	$1 + c_1(x/\xi) + c_2(x/\xi)^2 + \cdots$

fourth lines), while the deviation starts at $x \sim \sqrt{\xi_a \xi}$. A physical explanation for this deviation may be as follows. Consider light incident from the interface. It effectively penetrates into the medium with a depth ξ_a . As such, the Lyapunov exponent (inverse localization length), $\bar{\gamma}$, fluctuates, following a distribution $\sim e^{-\frac{\xi_a \xi}{4}(\bar{\gamma}-\xi^{-1})^2}$ [22]. Averaging $e^{-\bar{\gamma}x}$ with respect to this distribution we recover the result given in the second line.

If absorption is strong such that $\xi/\xi_a \gg 1$, then $\lambda \ll 1$ irrespective of x. By using the first line of Eq. (9) we find the results summarized in Table II.

Behavior of D(x) in finite localized samples. — These samples $(\xi/L \ll 1)$ have an essential difference from semiinfinite samples in the existence of resonant states most of which reside near the sample center [6, 7]. As shown in Ref. [9], they play decisive roles in establishing the highly unconventional macroscopic diffusion of localized waves. Due to long lifetime the resonant state is very sensitive to absorption. Below we shall discuss separately the cases of weak $(\xi_a/L \gg 1)$ and strong $(\xi_a/L \ll 1)$ absorption. In the former case, the scaling factor (8) is simplified to $\lambda(x) \approx \frac{x(L-x)}{L\xi} [1 - (\frac{x(L-x)}{3\xi_a^2})]$. For $\lambda(x) \gg 1$ by using the second line of Eq. (9), we find the local diffusion coefficient $D(x) \sim e^{-\lambda(x)}$. It enhances from its value in the passive case, which is $\sim e^{-\frac{x(L-x)}{L\xi}}$ [9], by a factor of $\sim \exp[\frac{1}{3}(\frac{x(L-x)}{\xi_a\sqrt{L\xi}})^2]$. The enhancement factor increases monotonically from the interface to the sample mid-point. Such inhomogeneous enhancement reflects that upon turning on absorption resonant states are 'killed', and the portion is determined by the lifetime of resonant states. Indeed, near the sample mid-point these states have the longest lifetime and are most easily to be 'killed', and this accounts for the strongest enhancement at the mid-point.

The interplay of localization and absorption is even stronger in the latter case $(\xi_a/L \ll 1)$. For x sufficiently away from the sample center, $x \ll \xi_a/2$, the scaling factor (8) reduces to that of semi-infinite samples, $\lambda(x) = \frac{\xi_a}{2\xi}(1 - e^{-2x/\xi_a})$, and D(x) is the same as that of semi-infinite samples. (We consider only $x \leq L/2$ since D(x) is symmetric with respect to the sample midpoint.) Its behavior is summarized in Table I and II. Near the sample center, $x \gtrsim \xi_a/2$, the scaling factor (8) is simplified to $\lambda(x) \approx \lambda(L/2)$. Thus, $D(x) \approx D(L/2)$ exhibiting a plateau. Strikingly, despite that the system is controlled by two parameters, ξ/L and ξ_a/L , the

TABLE II: Behavior of D(x) at $\xi_a \ll \xi \ll L$

regime	D(x)/D(0)
$x \gtrsim \xi_a/2$	$1 + c_1(\xi_a/2\xi) + c_2(\xi_a/2\xi)^2 + \cdots$
$x \ll \xi_a/2$	$1 + c_1(x/\xi) + c_2(x/\xi)^2 + \cdots$

plateau value depends on single parameter, ξ_a/ξ . More precisely, Eqs. (2) and (8) give Eq. (3). Moreover, from Eq. (9) we obtain the expression of D(L/2) for $\xi_a/2\xi \gg 1$ $(\xi_a/2\xi \ll 1)$ which is given by the first line of Table I (II). Note that in the strong absorption limit, $\xi/\xi_a \to \infty$, the homogeneity of the diffusion coefficient is restored, $D(x) = D_0$, reflecting the absence of localization effects.

That the spatial resolution of D(x) for $x \leq L/2$ is identical to that of semi-infinite samples when $\xi_a/L \ll 1$ reflects an important fact: for strong absorption waves could not propagate from the left-half to the right-half part of the sample and *vice versa*. As such, resonance states play no roles. Therefore, one might expect that the phenomenological model of Ref. [11] is valid, as observed in numerical experiments [17].

In summary, we present the first microscopic theory showing that in localized open absorbing media photons display unusual Brownian motion. First of all, the diffusion coefficient is inhomogeneous in space; most strikingly, despite that the system is controlled by two parameters $(\xi/L \text{ and } \xi_a/L)$, it exhibits novel single parameter scaling namely Eq. (2). The analytic predictions for the spatial resolution of the diffusion coefficient are confirmed by numerical simulations. We stress that our theory is very general and, particularly, valid for arbitrary absorption strength. In the limiting case of strong absorption realized experimentally [16], our results agree well with experimental measurements. It is very interesting to generalize the present theory to the gained system in the future, and this may have direct applications in random lasers [23].

We thank A. Z. Genack and A. G. Yamilov for useful discussions. This work is supported by the NSFC (No. 11174174) and by the Tsinghua University Initiative Scientific Research Program (No. 2011Z02151).

- [1] A. Einstein, Annalen der Physik **17**, 549 (1905).
- [2] S. Chandrasekhar, *Radiative Transfer* (Dover, New York, 1960).
- [3] S. John, Phys. Rev. Lett. 53, 2169 (1984).
- [4] A. Lagendijk, B. van Tiggelen, and D. S. Wiersma, Physics Today 62, 24 (2009).
- [5] C. Tian, Physica E **49**, 124 (2013).
- [6] M. Ya. Azbel, Phys. Rev. B 28, 4106 (1983).
- [7] K. Yu. Bliokh, Yu. P. Bliokh, and V. D. Freilikher, J. Opt. Soc. Am. B 21, 113 (2004).
- [8] Z. Q. Zhang, A. A. Chabanov, S. K. Cheung, C. H. Wong, and A. Z. Genack, Phys. Rev. B 79, 144203 (2009).
- [9] C. S. Tian, S. K. Cheung, and Z. Q. Zhang, Phys. Rev. Lett. 105, 263905 (2010).
- [10] B. Payne, A. Yamilov, and S. E. Skipetrov, Phys. Rev. B 82, 024205 (2010).
- [11] B. A. van Tiggelen, A. Lagendijk, and D. S. Wiersma, Phys. Rev. Lett. 84, 4333 (2000).
- [12] C. Tian, Phys. Rev. B 77, 064205 (2008).
- [13] J. Wang and A. Z. Genack, Nature 471, 345 (2011).
- [14] K. Yu. Bliokh, Yu. P. Bliokh, V. Freilikher, A. Z. Genack, B. Hu., and P. Sebbah, Phys. Rev. Lett. 97, 243904 (2006).
- [15] K. Yu. Bliokh, Yu. P. Bliokh, V. Freilikher, S. Savel'ev, and F. Nori, Rev. Mod. Phys. 80, 1201 (2008).
- [16] A. G. Yamilov, R. Sarma, B. Redding, B. Payne, H. Noh and H. Cao, arXiv: 1303.3244.
- [17] A. G. Yamilov and B. Payne, arXiv:1303.2717.
- [18] V. D. Freilikher, M. Pustilnik, and I. Yurkevich, Phys. Rev. B 50, 56 (1997).
- [19] K. B. Efetov, Supersymmetry in Disorder and Chaos (Cambridge University Press, Cambridge, England, 1997).
- [20] V. Ya. Chernyak, K. I. Grigoshin, E. I. Ogievetsky, and V. M. Agranovich, Solid State Commun. 84, 209 (1992).
- [21] For $\epsilon'' \sim \mathcal{O}(1)$ or in strictly one dimensions both D_0 and γ strongly depends on ϵ'' . However, the detailed forms are unimportant for the present work, and therefore we shall not discuss this issue further.
- [22] P. W. Anderson, D. J. Thouless, E. Abrahams, and D. S. Fisher, Phys. Rev. B 22, 3519 (1980).
- [23] H. Cao, Y. G. Zhao, S. T. Ho, E. W. Seelig, Q. H. Wang, and R. P. H. Chang, Phys. Rev. Lett. 82, 2278 (1999).