

Uniform approximation of paraxial flat-topped beams

Riccardo Borghi

Dipartimento di Elettronica, Università “Roma Tre”

Via Volterra 62, I-00146 Rome, Italy

A uniform asymptotic theory of the free-space paraxial propagation of coherent flattened Gaussian beams is proposed in the limit of nonsmall Fresnel numbers. The pivotal role played by the error function in the mathematical description of the related wavefield is stressed.

© 2019 Optical Society of America

OCIS codes: 000.3860, 260.1960, 350.5500

1. Introduction

Several analytical models aimed at describing flat-topped profiles were proposed in the past. Although several of them, like for instance the supergaussian and the Fermi-Dirac profiles, present a very simple mathematical structure, their free-space propagation cannot be described without resorting to numerical integration, even within paraxial approximation. To overcome such difficulties, in 1994 Gori introduced the first model, called flattened Gaussian (FG henceforth) [1], able to describe the paraxial propagation of coherent flat-topped beams in exact terms [2]. Differently from the supergaussian or the Fermi-Dirac, FG profiles are expressed through a finite sum in the following way:

$$\text{FG}_N(\xi) = \exp(-\xi^2) \sum_{m=0}^N \frac{1}{m!} \xi^{2m}, \quad (1)$$

where N denotes the so-called FG profile order [3]. On further rescaling the argument ξ by a factor $\sqrt{N+1}$, the FG profile assumes a flat-topped shape which for $N = 0$ reduces to a Gaussian distribution whereas for $N \rightarrow \infty$ tends to coincide with the characteristic function of the unitary disk. Later, Li [4] found that for a typical profile to display a flat-topped shape it is necessary for its first $2N$ ξ -derivatives to be null. On using such condition, which is satisfied by the FG profile in Eq. (1), Li introduced a new flat-topped profile based on the superposition of N fundamental Gaussian functions having different widths suitably chosen. Both Gori's and Li's approaches allow the paraxial propagation problem of flat-topped beams to be solved without approximations, as the resulting fields turns out to be described in terms of finite sums. However, although no computational difficulties are encountered, the behavior of those sums upon free propagation is really hard to grasp.

The aim of the present Letter is to replace the N -term sum by a uniform [5], with respect to the transverse radial position, analytical approximation whenever the Fresnel number [6] is not too small.

2. Theoretical Analysis

We start our analysis from the field distribution of an axisymmetric FG beam of order N at the plane $z = 0$ of a cylindrical reference frame (\mathbf{r}, z) , given by [1]

$$U_N(r; 0) = \text{FG}_N \left(\frac{r}{w_0} \sqrt{N+1} \right), \quad (2)$$

where an unessential amplitude constant has been set to one for simplicity and the symbol w_0 denotes the spot-size of the FG beam. The field $U_N(\mathbf{r}; z)$ propagated in free space at the

transverse plane $z > 0$ can be expressed as follows [7]:

$$\begin{aligned}
U_N(r; z) &= \frac{\exp(ikz)}{1 + i\frac{N+1}{\pi N_F}} \exp \left[-\frac{N+1}{1 + i\frac{N+1}{\pi N_F}} \left(\frac{r}{w_0} \right)^2 \right] \\
&\times \mathcal{G}_N \left[\frac{1}{1 + i\frac{N+1}{\pi N_F}}, \frac{N+1}{1 + i\frac{N+1}{\pi N_F}} \left(\frac{r}{w_0} \right)^2 \right],
\end{aligned} \tag{3}$$

where $N_F = w_0^2/\lambda z$ denotes the Fresnel number [6] and the function $\mathcal{G}_N(\cdot, \cdot)$ is defined as

$$\mathcal{G}_N(t, s) = \sum_{n=0}^N (-1)^n \binom{N+1}{n+1} t^n L_n(s), \tag{4}$$

with $L_n(\cdot)$ being the n th-order Laguerre polynomial [8]. To derive a uniform asymptotic approximation of $U_N(r; z)$ with respect to r we start from the following integral representation of Laguerre polynomials [9, Eq. (5.4.1)]:

$$L_n(s) = \frac{\exp(s)}{n!} \int_0^\infty d\xi \exp(-\xi) \xi^n J_0 \left(2\sqrt{s\xi} \right), \tag{5}$$

where $J_n(\cdot)$ denotes the n th-order Bessel function of the first kind [8]. On substituting from Eq. (5) into Eq. (4) and on taking into account that

$$\sum_{n=0}^N \frac{(-1)^n}{n!} \binom{N+1}{n+1} x^n = L_N^{(1)}(x), \tag{6}$$

with $L_n^{(\alpha)}(\cdot)$ denoting the generalized Laguerre polynomial of order n and degree α [8], we obtain

$$\begin{aligned}
\exp(-s) \mathcal{G}_N(t, s) &= \\
&= \int_0^\infty d\xi \exp(-\xi) J_0 \left(2\sqrt{s\xi} \right) L_N^{(1)}(\xi t).
\end{aligned} \tag{7}$$

For nonsmall values of N , the Laguerre polynomial $L_N^{(1)}$ can be estimated by using Eq. (8.22.4) of [9] which gives

$$\frac{\exp\left(-\frac{\xi t}{2}\right)}{N+1} L_N^{(1)}(\xi t) \simeq \frac{2 J_1 \left[2\sqrt{(N+1)\xi t} \right]}{2\sqrt{(N+1)\xi t}}. \tag{8}$$

that, once substituted into Eq. (7), on taking into account that, from Eq. (3), $s = \frac{r^2}{w_0^2} (N+1) t$, and after letting $\eta = (N + 1) \xi t$, leads to

$$\begin{aligned} \exp(-s) \mathcal{G}_N(t, s) &\simeq \\ &\simeq \frac{2}{\sqrt{t}} \int_0^\infty d\eta \exp(-p\eta^2) J_0\left(2\frac{r}{w_0}\eta\sqrt{t}\right) J_1\left(2\eta\sqrt{t}\right), \end{aligned} \quad (9)$$

where $p = (1 - t/2)/(N + 1)$. Finally, on using formula 2.12.39.1 of [10] to estimate the integral in Eq. (9), after long but straightforward algebra it is not difficult to recast Eq. (3) as follows:

$$U_N(\mathbf{r}; z) \simeq \exp(ikz) \mathcal{J}\left(\frac{2(N+1)}{1 + 2i\frac{N+1}{\pi N_F}}, \frac{r}{w_0}\right), \quad (10)$$

where the function $\mathcal{J}(u, \rho)$ is defined by

$$\mathcal{J}(u, \rho) = 1 - \exp(-u\rho^2) \int_u^\infty d\xi \exp(-\xi) I_0(2\rho\sqrt{\xi u}), \quad (11)$$

with $I_0(\cdot)$ denoting the zeroth-order modified Bessel function of the first kind [8]. To derive a uniform, with respect to ρ , approximation of $\mathcal{J}(u, \rho)$, we use an approach originally employed by Schwarzschild in studying the Fresnel diffraction from a circular hole [11, Sec. 4.3]. First of all we replace the modified Bessel function by its asymptotic expansion $I_0(x) \sim \exp(x)/\sqrt{2\pi x}$ to obtain, after the integration variable change $\xi \rightarrow u\xi^2$,

$$\mathcal{J}(u, \rho) \sim 1 - \sqrt{\frac{u}{\pi\rho}} \int_1^\infty \sqrt{\xi} d\xi \exp[-u(\xi - \rho)^2]. \quad (12)$$

Then we replace the factor $\sqrt{\xi}$ by $(\sqrt{\xi} - \sqrt{\rho}) + \sqrt{\rho}$, thus obtaining

$$\begin{aligned} \mathcal{J}(u, \rho) &\sim 1 - \sqrt{\frac{u}{\pi}} \int_1^\infty d\xi \exp[-u(\xi - \rho)^2] \\ &\quad - \sqrt{\frac{u}{\pi\rho}} \int_1^\infty d\xi (\sqrt{\xi} - \sqrt{\rho}) \exp[-u(\xi - \rho)^2]. \end{aligned} \quad (13)$$

On using formula 2.3.15.4 of [10] the first integral becomes

$$\sqrt{\frac{u}{\pi}} \int_1^\infty d\xi \exp[-u(\xi - \rho)^2] = \frac{1}{2} \operatorname{erfc}[\sqrt{u}(1 - \rho)], \quad (14)$$

where $\text{erfc}(\cdot)$ denotes the complementary error function [8, Eq. (7.2.2)]. As far as the second integral in Eq. (13) is concerned, a first partial integration gives at once

$$\begin{aligned}
& -\sqrt{\frac{u}{\pi\rho}} \int_1^\infty d\xi (\sqrt{\xi} - \sqrt{\rho}) \exp[-u(\xi - \rho)^2] = \\
& -\frac{1}{2\sqrt{u\pi\rho}} \frac{\exp[-u(1 - \rho)^2]}{1 + \sqrt{\rho}} \\
& + \frac{1}{4\sqrt{u\pi\rho}} \int_1^\infty \frac{d\xi}{\sqrt{\xi}} \frac{\exp[-u(\xi - \rho)^2]}{(\sqrt{\xi} + \sqrt{\rho})^2},
\end{aligned} \tag{15}$$

while further partial integrations would give rise to an asymptotic expansion in negative powers of u whose single terms can easily find by using standard techniques [12]. However, in the following we shall keep only the leading term in Eq. (15) and write \mathcal{J} as follows:

$$\begin{aligned}
\mathcal{J}(u, \rho) & \simeq 1 - \frac{1}{2} \text{erfc}[\sqrt{u}(1 - \rho)] \\
& - \frac{1}{2\sqrt{u\pi\rho}} \frac{\exp[-u(1 - \rho)^2]}{1 + \sqrt{\rho}}, \quad |u| \gg 1.
\end{aligned} \tag{16}$$

Equations (10) and (16) constitutes the main results of the present Letter and suggest an intriguing connection between the error function and FG profiles.

3. Discussion

In fact, we note that for $N_F \rightarrow \infty$ the propagated field $U_N(\mathbf{r}; z)$ must coincide with the initial field $U_N(\mathbf{r}; 0)$, so that from Eq. (10) we have $U_N(\mathbf{r}; 0) \simeq \mathcal{J}[2(N + 1), r/w_0]$. Accordingly, the following asymptotic relation holds:

$$\text{FG}_N(\xi) \simeq 1 - \frac{1}{2} \text{erfc} \left[\sqrt{2}(\sqrt{N + 1} - \xi) \right], \tag{17}$$

for $N \gg 1$. Figure 1 shows a visual comparison of FG profiles obtained through the definition in Eq. (1) (solid curve) and through the asymptotic estimate in Eq. (17) (dashed curves). Except for the case $N = 1$, the agreement is remarkably good and the two curves are practically undistinguishable for $N > 20$. A possible intuitive explanation is to prove that the initial erfc-based profile in the r.h.s. of Eq. (17), evaluated at the scaled variable $\xi/\sqrt{N + 1}$, satisfies the above quoted Li's flatness condition given by Eq. (4) of [4]. On using formula 1.5.1.1 of [13] it can be shown that the ξ -derivative of order $2n$ of the scaled




Fig. 1. FG profiles $f_N(\xi)$ evaluated, for different values of N , through Eq. (1) (solid curves) and through the asymptotic estimate in Eq. (17) (dashed curves).

erfc-based profile, evaluated at $\xi = 0$, is given by

$$-\frac{2}{\sqrt{\pi}} \left(\frac{2}{N+1} \right)^n \exp[2(N+1)] H_{2n-1}(\sqrt{2}\sqrt{N+1}), \quad (18)$$

where $H_n(\cdot)$ denotes the n th-order Hermite polynomial [8]. Then, on using formula 8.22.7 of [9] and the Stirling's approximation of factorials, after some algebra the following estimate of the modulus of the above derivative holds:

$$\frac{(n-1)!}{\pi} \left(\frac{8}{N+1} \right)^n \exp[-(N+1)], \quad (19)$$

which, for sufficiently high values of N , turns out to be negligible for $n < N$, as required. To complete the scenario, in Fig. 2 the amplitudes of a FG beam of order $N = 30$ propagated at the Fresnel number $N_F = 2$ (a) and $N_F = 5$ (b) are shown as functions of the normalized transverse radial distance r/w_0 (the dotted curve is the initial FG profile). In each figure the open circles represent the exact values of the field amplitude provided by Eq. (3), the dashed curve is the uniform approximation obtained by keeping only the erfc-based term in Eq. (16), and the solid curve is obtained by adding the first-order correction in the second row of Eq. (16). On further increasing N_F the relative weight of such a correction becomes negligible and the error function alone well describes the propagated field. In such limit there is a sort of “invariance” in the mathematical description of the FG beam propagation. The functional form remains the same describing the field at the initial plane and the effect of the propagation is represented by a simple (complex) scaling of its argument.




Fig. 2. Behavior of the amplitude of a FG beam of order $N = 30$ propagated at the Fresnel number $N_F = 2$ (a) and $N_F = 5$ (b). Open circles: exact values provided by Eq. (3). Dashed curve: erfc-based asymptotic estimate through the sole first row of Eq. (1). Solid curve: asymptotic estimate obtained by adding the correction given in the second row of Eq. (1). The dotted curve represents the initial FG profile.

4. Conclusions

While the use of the error function is well known in the study of diffraction by large hard-edge apertures [5], our results seem to suggest its involvement also in the mathematical description of the propagation of smooth initial field distributions, which is worth being further explored.

References

1. F. Gori, “Flattened gaussian beams,” *Opt. Commun.* **107**, 335-341 (1994).

2. V. Bagini, R. Borghi, F. Gori, A. M. Pacileo, M. Santarsiero, D. Ambrosini, and G. Schirripa Spagnolo, "Propagation of axially symmetric flattened Gaussian beams," *J. Opt. Soc. Am. A* **13**, 1385-1394 (1996).
3. Equation (1) was originally derived starting from the identity $1 = \exp(-\xi^2) \exp(\xi^2)$ and on truncating the Taylor expansion of the second exponential up to N .
4. Y. Li, "Light beams with flat-topped profiles," *Opt. Lett.* **27**, 1007-1009 (2002).
5. J. Stamnes, "Uniform asymptotic theory of diffraction by apertures," *J. Opt. Soc. Am.* **73**, 96-109 (1983).
6. M. Born and E. Wolf, *Principles of Optics* (Cambridge University Press, Cambridge, 1999).
7. R. Borghi, "Elegant Laguerre-Gauss beams as a new tool for describing axisymmetric flattened Gaussian beams," *J. Opt. Soc. Am. A* **18**, 1627-1633 (2001).
8. Digital Library of Mathematical Functions, National Institute of Standards and Technology <http://dlmf.nist.gov/>.
9. G. Szegő, *Orthogonal Polynomials* (AMS, Providence, 1939).
10. A. P. Prudnikov, Y. A. Brychkov, and O. I. Marichev, *Integrals and Series* (Gordon Breach, 1986), Vol. I.
11. K. D. Mielenz, "Algorithms for Fresnel diffraction at rectangular and circular apertures," *J. Res. Natl. Inst. Stand. Technol.* **103**, 497 (1998).
12. N. M. Temme, "Uniform asymptotic expansions of integrals: a selection of problems," *J. Comput. Appl. Math.* **65**, 395-417 (1995).
13. Y. A. Brichkov, *Handbook of Special Functions* (CRC Press, London, 2008).