

AVERAGING ON MANIFOLDS BY EMBEDDING ALGORITHM

Petre Birtea*, Dan Comănescu* and Călin-Adrian Popa**

*Department of Mathematics, West University of Timișoara

Bd. V. Pârvan, No 4, 300223 Timișoara, România

birtea@math.uvt.ro, comanescu@math.uvt.ro

**Department of Computer Science, "Politehnica" University of Timișoara

Bd. V. Pârvan, No 2, 300223 Timișoara, România

calin.popa@cs.upt.ro

Abstract

We will propose a new algorithm for finding critical points of cost functions defined on a differential manifold. We will lift the initial cost function to a manifold that can be embedded in a Riemannian manifold (Euclidean space) and will construct a vector field defined on the ambient space whose restriction to the embedded manifold is the gradient vector field of the lifted cost function. The advantage of this method is that it allows us to do computations in Cartesian coordinates instead of using local coordinates and covariant derivatives on the initial manifold. We will exemplify the algorithm in the case of $SO(3)$ averaging problems and will rediscover a few well known results that appear in literature.

MSC: 43A07, 53B21, 58A05.

Keywords: Averaging, Distance functions, Rotations, Quaternions, Metriplectic dissipation, Gradient vector field.

1 Introduction

A problem which one frequently encounters in applications is to find an average for a finite set of sample points $\{y_1, y_2, \dots, y_r\}$ belonging to a manifold N . A method to solve this problem is to construct a cost function $G_N : N \rightarrow \mathbb{R}$ associated with this set of sample points and then the average is defined by

$$\arg \min_{y \in N} G_N(y).$$

Of special interest are the cost functions of least square type $G_N(y) := \sum_{i=1}^r d^2(y, y_i)$, where d is a distance function on N . The average of the set of sample points $\{y_1, y_2, \dots, y_r\}$ on the manifold N is the set defined by

$$\arg \min_{y \in N} \sum_{i=1}^r d^2(y, y_i).$$

When the function G_N is differentiable, this is equivalent with solving the equation $dG_N(y) = 0$ and with testing for which solutions the minimum value is attained. In order to write this equation we need the knowledge of a local system of coordinates on the manifold N , a requirement that might be difficult to fulfill in many practical cases. Another problem that we may encounter when this approach is that the set of sample points $\{y_1, \dots, y_r\}$ might not be entirely included in the domain of a single local system of coordinates. One way to overcome these difficulties is to lift the problem on a simpler manifold S .

Let $\mathbf{P} : S \rightarrow N$ be a surjective submersion. The lifted cost function is $G_S : S \rightarrow \mathbb{R}$ defined by $G_S = G_N \circ \mathbf{P}$. The set equality $\mathbf{P}(\{x \in S \mid dG_S(x) = 0\}) = \{y \in N \mid dG_N(y) = 0\}$ shows that it is sufficient to solve the equation $dG_S(x) = 0$ on the simpler manifold S and project these solutions on the manifold N .

A way to solve this new problem is to endow S with a Riemannian metric τ and compute the critical points of the gradient vector field $\nabla_{\tau}G_S$ and verify which one of them are local/global minima. We note that if $x_0 \in S$ is a critical point of the vector field $\nabla_{\tau}G_S$, then it is a critical point of the vector field $\nabla_{\tau'}G_S$, where τ' is any other Riemannian metric on S . Although the manifold S might have a simpler geometrical structure than the initial manifold N , we still need a local system of coordinates or the knowledge of covariant derivatives on the manifold S in order to be able to compute the critical points of the function G_S , see [2], [3], [9], [11], [19], [24]. To further avoid the use of Riemannian geometry of the manifold S we will embed S in a larger space M .

In the current paper we propose a solution for finding the critical points of the lifted cost function G_S by constructing a vector field $\mathbf{v}_0 \in \mathcal{X}(M)$ on the ambient space M (usually an Euclidean space), that is tangent to the submanifold S and by also constructing a Riemannian metric τ on S such that $\mathbf{v}_0|_S = \nabla_{\tau}G_S$. Consequently, the critical points of the vector field \mathbf{v}_0 (usually written in Cartesian coordinates) that belong to S are critical points of the lifted cost function G_S and their projection through the surjective submersion \mathbf{P} gives the critical points of the initial cost function G_N . This construction will be illustrated in details for the averaging problem on the Lie group $SO(3)$ associated with four different cost functions.

In the literature the problem of averaging on Lie groups is often solved by using the exponential map as a tool to introduce local coordinates and so lifting the problem on the tangent space, see [3], [4], [11], [12], [13], [21]. This method can be generalized to the context of Riemannian manifolds as there also exists an exponential map. The exponential map can be further replaced by the more general notion of retraction developed in [3], [10]. Averaging problems coming from real world applications have been studied in [2], [3], [16], [24], [25], [26] by using the notions of covariant derivatives and the geometry of geodesics on various Riemannian manifolds. In this paper we will propose a different algorithm in order to deal with averaging problems on general differentiable manifolds.

Other interesting cost functions are the ones that come from the Fermat-Torricelli problem. This averaging problem has been studied on various spaces, see [7], [8], [15], [20], and it will be of interest to apply the techniques developed in this paper to such problems.

2 Averaging on a differentiable manifold

We will solve the averaging problem presented in the Introduction by embedding the manifold S as a submanifold of a Riemannian manifold (M, g) . Further more, we will assume in this paper that S is the preimage of a regular value for a smooth function $\mathbf{F} := (F_1, \dots, F_k) : M \rightarrow \mathbb{R}^k$, i.e. $S = \mathbf{F}^{-1}(c_0)$, for c_0 a regular value of \mathbf{F} .

As we have already stated, our approach is to find a Riemannian metric τ on the submanifold S and a vector field $\mathbf{v}_0 \in \mathcal{X}(M)$ such that

$$\mathbf{v}_0|_S = \nabla_{\tau}G_S. \quad (2.1)$$

Consequently, our initial problem is equivalent with finding the critical points of the vector field \mathbf{v}_0 , which is defined on the ambient space M , and with choosing the ones that belong to S .

The vector field that we are looking for is the standard control vector field \mathbf{v}_0 introduced in [6], which in turn is a continuation of the study we have begun in [5]. Let (M, g) be a m dimensional Riemannian manifold and $F_1, \dots, F_k, G : M \rightarrow \mathbb{R}$ be $k + 1$ smooth functions. The standard control vector field is defined by

$$\mathbf{v}_0 = \sum_{i=1}^k (-1)^{i+k+1} \det \Sigma_{(F_1, \dots, \widehat{F}_i, \dots, F_k, G)}^{(F_1, \dots, F_k)} \nabla F_i + \det \Sigma_{(F_1, \dots, F_k)}^{(F_1, \dots, F_k)} \nabla G, \quad (2.2)$$

where $\det \Sigma_{(F_1, \dots, F_k)}^{(F_1, \dots, F_k)}$ is the determinant of the Gramian matrix formed with the vector fields $\nabla F_1, \dots, \nabla F_k$. The vector field \mathbf{v}_0 conserves the regular leaves of the map $\mathbf{F} := (F_1, \dots, F_k) : M \rightarrow \mathbb{R}^k$ and dissipates the function G , more precisely $L_{\mathbf{v}_0}G = \det \Sigma_{(F_1, \dots, F_k, G)}^{(F_1, \dots, F_k, G)} \geq 0$.

In what follows, we will describe the geometry of the standard control vector field. Let $\mathbf{T} : \Omega^1(M) \times$

$\Omega^1(M) \rightarrow \mathbb{R}$ be the degenerate symmetric contravariant 2-tensor given by

$$\mathbf{T} := \sum_{i,j=1}^k (-1)^{i+j+1} \det \Sigma_{(F_1, \dots, \hat{F}_j, \dots, F_k)}^{(F_1, \dots, \hat{F}_j, \dots, F_k)} \nabla F_i \otimes \nabla F_j + \det \Sigma_{(F_1, \dots, F_k)}^{(F_1, \dots, F_k)} g^{-1}. \quad (2.3)$$

The following result shows that \mathbf{v}_0 looks like the gradient vector field of the function G , with respect to the degenerate symmetric contravariant 2-tensor \mathbf{T} .

Theorem 2.1. [6] *On the manifold (M, g) , the standard control vector field \mathbf{v}_0 is given by the following formula*

$$\mathbf{v}_0 = \mathbf{i}_{dG}(\mathbf{T}).$$

From the above theorem, computing the critical points of \mathbf{v}_0 (which is written in local coordinates on M) that belong also to S is equivalent with solving the following system of $m+k$ equations on M :

$$\begin{cases} \begin{bmatrix} \mathbf{T}^{11}(x) & \dots & \mathbf{T}^{1m}(x) \\ \vdots & \dots & \vdots \\ \mathbf{T}^{m1}(x) & \dots & \mathbf{T}^{mm}(x) \end{bmatrix} \begin{bmatrix} \frac{\partial G}{\partial x^1}(x) \\ \vdots \\ \frac{\partial G}{\partial x^m}(x) \end{bmatrix} = \mathbf{0} \\ \mathbf{F}(x) = c \end{cases}, \quad (2.4)$$

where $\mathbf{T} = \mathbf{T}^{pq} \frac{\partial}{\partial x^p} \otimes \frac{\partial}{\partial x^q}$ and $dG = \frac{\partial G}{\partial x^i} dx^i$. In general, a system of $m+k$ equations with m unknowns might not have any solutions. But the above system that has m unknowns, namely the coordinates of x in M , has exactly m functional independent equations due to the fact that the rank of the tensor \mathbf{T} is $m-k$ for every regular point in the open set in M of regular points of \mathbf{F} .

The standard control vector field \mathbf{v}_0 is tangent to every regular leaf and next we will recall its geometry when restricted to a regular leaf. Let $i_c : L_c \rightarrow M$ be the canonical inclusion of the regular leaf L_c into the manifold M . The contravariant symmetric 2-tensor \mathbf{T} is non-degenerate when restricted to a regular leaf L_c and consequently, the restriction can be inverted and it thus generates the Riemannian metric, see [6], $\tau_c : \mathcal{X}(L_c) \times \mathcal{X}(L_c) \rightarrow \mathcal{C}^\infty(L_c)$ defined by

$$\tau_c(X^c, Y^c) := \mathbf{T}^{-1}((i_c)_* X^c, (i_c)_* Y^c).$$

Theorem 2.2. [6] *On a regular leaf L_c we have the following characterizations.*

- (i) $\tau_c = \frac{1}{\det \Sigma_{(F_1, \dots, F_k)}^{(F_1, \dots, F_k)} \circ i_c} (i_c)^* g$
- (ii) $\mathbf{v}_0|_{L_c} = (i_c)_* \nabla_{\tau_c} (G \circ i_c)$.

The Riemannian metric on S is $\tau = \tau_{c_0}$ and the above theorem also shows that the equality (2.1) holds, where $G_S = G|_S$.

In conclusion, in order to solve our initial averaging problem on the manifold N associated to the cost function G_N , by using the standard control vector field, we apply the following **embedding algorithm**:

- (i) Choose the geometrical setting of a manifold S , a surjective submersion $\mathbf{P} : S \rightarrow N$ and construct the lifted cost function $G_S = G_N \circ \mathbf{P}$.
- (ii) Find a Riemannian ambient space (M, g) and a differentiable function $\mathbf{F} : M \rightarrow \mathbb{R}^k$ such that $S = \mathbf{F}^{-1}(c_0)$, where c_0 is a regular value of \mathbf{F} . Construct the contravariant symmetric 2-tensor \mathbf{T} .
- (iii) Find a prolongation function $G : M \rightarrow \mathbb{R}$ such that $G|_S = G_S$ and construct the standard control vector field \mathbf{v}_0 with the initial data \mathbf{F}, G .
- (iv) Solve the system $\mathbf{v}_0|_S(x) = \mathbf{0}$ (i.e. the system (2.4)) which is equivalent with finding the critical points of the lifted cost function G_S . Project through \mathbf{P} these solutions and thus obtain the critical points of the initial cost function G_N .

(v) Find critical points that are local/global minima for the cost function G_N .

Examples that fit the geometry underlying this embedding algorithm are given by the rich cases of quotient manifolds, $N = S/K$ with K a group that acts on S . In the case when N can be written as a preimage of a regular value of a smooth map \mathbf{F} , then (i) is not a necessary step of the algorithm ($S = N$ and \mathbf{P} is the identity).

In the case when $\mathbf{P}_*\mathbf{v}_{\mathbf{0}|S}$ is a vector field on the manifold N , then we have the set inclusion $\mathbf{P}(\{x \in S \mid \mathbf{v}_{\mathbf{0}|S}(x) = \mathbf{0}\}) \subset \{y \in N \mid \mathbf{P}_*\mathbf{v}_{\mathbf{0}|S}(y) = \mathbf{0}\}$. If \mathbf{P} is also a local diffeomorphism we will obtain equality between the above two sets and consequently, the critical points of the cost function G_N are characterized by the equation $\mathbf{P}_*\mathbf{v}_{\mathbf{0}|S}(y) = \mathbf{0}$. In this particular setting the step (iv) can be replaced by the following:

(iv') Suppose that $\mathbf{P}_*\mathbf{v}_{\mathbf{0}|S} \in \mathcal{X}(N)$ and \mathbf{P} is a local diffeomorphism, then solve the equation $\mathbf{P}_*\mathbf{v}_{\mathbf{0}|S}(y) = \mathbf{0}$ (the solutions are critical points for the cost function G_N).

For solving point (v) one can study the second order derivatives of the cost function G_N (see [9], [16], [19]) or study the convexity properties of the cost functions as in [15].

3 Averaging on $SO(3)$

A problem frequently encountered in practice is finding a rotation that is in some sense an average of a finite set of rotations. The group of special rotations is not an Euclidean space, but a differential manifold. Consequently, an averaging problem on such a space has to be considered in the realm of differential geometry.

The special orthogonal group $SO(3)$ can be identified by a double covering map with the sphere S^3 in \mathbb{R}^4 , see [1], [23]. Our manifold S will be S^3 and the ambient space (M, g) will be the Euclidean space \mathbb{R}^4 . We will consider different distance functions on $SO(3)$ that generate different cost functions and will solve the associated averaging problem, as we have described in the previous section, by using the embedding algorithm.

From a historical perspective, we will use the quaternion notation on \mathbb{R}^4 , although the quaternion structure will not be specifically used in our computations. The unit quaternions $\mathbf{q} = (q^0, q^1, q^2, q^3) \in S^3 \subset \mathbb{R}^4$ and $-\mathbf{q} \in S^3 \subset \mathbb{R}^4$ correspond to the following rotation in $SO(3)$:

$$\mathbf{R}^{\mathbf{q}} = \begin{pmatrix} (q^0)^2 + (q^1)^2 - (q^2)^2 - (q^3)^2 & 2(q^1q^2 - q^0q^3) & 2(q^1q^3 + q^0q^2) \\ 2(q^1q^2 + q^0q^3) & (q^0)^2 - (q^1)^2 + (q^2)^2 - (q^3)^2 & 2(q^2q^3 - q^0q^1) \\ 2(q^1q^3 - q^0q^2) & 2(q^2q^3 + q^0q^1) & (q^0)^2 - (q^1)^2 - (q^2)^2 + (q^3)^2 \end{pmatrix}$$

This gives rise to the smooth double covering map $\mathbf{P} : S^3 \rightarrow SO(3)$, $\mathbf{P}(\mathbf{q}) = \mathbf{R}^{\mathbf{q}}$. The covering map is a local diffeomorphism and consequently, instead of working with the distance function (cost function) on $SO(3)$, we will work with cost functions on the unit sphere $S^3 \subset \mathbb{R}^4$ as it has been explained in the previous section. In this case, for the step (ii) of the embedding algorithm, we have $S^3 = F^{-1}(1)$, with $F : \mathbb{R}^4 \rightarrow \mathbb{R}$, $F(\mathbf{q}) = (q^0)^2 + (q^1)^2 + (q^2)^2 + (q^3)^2$. The tensor \mathbf{T} is given by the formula, see [6] eq. (4.2), $\mathbf{T} = -\nabla F \otimes \nabla F + \|\nabla F\|^2 g^{-1}$, where g is the Euclidean metric on \mathbb{R}^4 . Writing this explicitly, the matrix associated to the symmetric contravariant 2-tensor \mathbf{T} is given by:

$$[\mathbf{T}(\mathbf{q})] = 4 \begin{pmatrix} (q^1)^2 + (q^2)^2 + (q^3)^2 & -q^0q^1 & -q^0q^2 & -q^0q^3 \\ -q^1q^0 & (q^0)^2 + (q^2)^2 + (q^3)^2 & -q^1q^2 & -q^1q^3 \\ -q^2q^0 & -q^2q^1 & (q^0)^2 + (q^1)^2 + (q^3)^2 & -q^2q^3 \\ -q^3q^0 & -q^3q^1 & -q^3q^2 & (q^0)^2 + (q^1)^2 + (q^2)^2 \end{pmatrix}.$$

In order to compute the standard control vector field $\mathbf{v}_{\mathbf{0}}$ according to the formula given in Theorem 2.1, we need the following elementary computation: if $\omega(\mathbf{q}) = \omega_0(\mathbf{q})dq^0 + \omega_1(\mathbf{q})dq^1 + \omega_2(\mathbf{q})dq^2 + \omega_3(\mathbf{q})dq^3$ is a one form on \mathbb{R}^4 , then we have the vector field

$$\mathbf{i}_{\omega} \mathbf{T}(\mathbf{q}) = 4(\langle \mathbf{q}, \mathbf{q} \rangle \bar{\omega}(\mathbf{q}) - \langle \mathbf{q}, \bar{\omega}(\mathbf{q}) \rangle \mathbf{q}), \quad (3.1)$$

where we have made the notation $\bar{\omega}(\mathbf{q}) := (\omega_0(\mathbf{q}), \omega_1(\mathbf{q}), \omega_2(\mathbf{q}), \omega_3(\mathbf{q}))$.

We will exemplify our construction by using various distance functions used in literature for measuring distances between Euclidean transformations. An extensive list is presented in [17], where their equivalence and functional dependence is also studied.

I. On the Lie group $SO(3)$ we will consider the distance function $d_1 : SO(3) \times SO(3) \rightarrow \mathbb{R}_+$, $d_1(\mathbf{R}_1, \mathbf{R}_2) = \|\mathbf{R}_1 - \mathbf{R}_2\|_F$, where $\|\cdot\|_F$ is the Frobenius norm. The associated cost function is $G_{1_{SO(3)}} : SO(3) \rightarrow \mathbb{R}$,

$$G_{1_{SO(3)}}(\mathbf{R}) = \sum_{i=1}^r \|\mathbf{R} - \mathbf{R}_i\|_F^2,$$

where $\mathbf{R}_1, \dots, \mathbf{R}_r$ are the sample rotation matrices, see [19], [22].

According to the step (i) of the embedding algorithm, we will lift the problem of finding the critical points of the cost function $G_{1_{SO(3)}}$ to the problem of finding the critical points of the lifted cost function $G_{1_{S^3}} : S^3 \rightarrow \mathbb{R}$, $G_{1_{S^3}} := G_{1_{SO(3)}} \circ \mathbf{P}$.

In order to compute the lifted cost function $G_{1_{S^3}}$, by using the surjectivity of the covering map \mathbf{P} , for each sample matrix \mathbf{R}_i we will choose a sample quaternion $\mathbf{q}_i \in S^3$. We have the following computation:

$$\begin{aligned} G_{1_{S^3}}(\mathbf{q}) &= \sum_{i=1}^r \|\mathbf{R}^{\mathbf{q}} - \mathbf{R}^{\mathbf{q}_i}\|_F^2 = \sum_{i=1}^r \text{tr}((\mathbf{R}^{\mathbf{q}} - \mathbf{R}^{\mathbf{q}_i})(\mathbf{R}^{\mathbf{q}} - \mathbf{R}^{\mathbf{q}_i})^T) \\ &= 2 \sum_{i=1}^r (3 - \text{tr}(\mathbf{R}^{\mathbf{q}T} \mathbf{R}^{\mathbf{q}_i})) \\ &= 8 \sum_{i=1}^r (1 - \langle \mathbf{q}, \mathbf{q}_i \rangle^2), \end{aligned}$$

where we have used the equality

$$\langle \mathbf{q}, \mathbf{q}_i \rangle^2 = \frac{1}{4}(\text{tr}(\mathbf{R}^{\mathbf{q}T} \mathbf{R}^{\mathbf{q}_i}) + 1). \quad (3.2)$$

For implementing the step (iii) we need to construct the prolongation function $G_1 : \mathbb{R}^4 \rightarrow \mathbb{R}$, $G_1(\mathbf{q}) = \sum_{i=1}^r 8(1 - \langle \mathbf{q}, \mathbf{q}_i \rangle^2)$. Note that $G_{1_{S^3}}$ and G_1 are even functions so they do not depend on the choice of the sample quaternions \mathbf{q}_i or $-\mathbf{q}_i$ which represent the same sample rotation \mathbf{R}_i . This subtler problem has also been addressed in a different way in [18]. Computing the differential 1-form dG_1 we obtain:

$$\begin{aligned} \overline{dG_1}(\mathbf{q}) &= \left(\frac{\partial G_1}{\partial q^0}(\mathbf{q}), \dots, \frac{\partial G_1}{\partial q^3}(\mathbf{q}) \right) = \left(-16 \sum_{i=1}^r q_i^0 \langle \mathbf{q}, \mathbf{q}_i \rangle, \dots, -16 \sum_{i=1}^r q_i^3 \langle \mathbf{q}, \mathbf{q}_i \rangle \right) \\ &= -16 \sum_{i=1}^r \langle \mathbf{q}, \mathbf{q}_i \rangle \mathbf{q}_i. \end{aligned}$$

Using (3.1), where the 1-form ω is dG_1 , the system of equations (2.4) becomes:

$$\begin{cases} \sum_{i=1}^r \langle \mathbf{q}, \mathbf{q}_i \rangle (\langle \mathbf{q}, \mathbf{q} \rangle \mathbf{q}_i - \langle \mathbf{q}, \mathbf{q}_i \rangle \mathbf{q}) = \mathbf{0} \\ \langle \mathbf{q}, \mathbf{q} \rangle = 1 \end{cases}, \quad (3.3)$$

and this represents the equation $\mathbf{v}_0|_{S^3}(\mathbf{q}) = \mathbf{0}$. The above system has four equations of third degree in the four unknowns $\mathbf{q} = (q^0, q^1, q^2, q^3)$ corresponding to the standard control vector field $\mathbf{v}_0 = \mathbf{0}$ plus the constraint equation that describes S^3 .

The four equations corresponding to $\mathbf{v}_0 = \mathbf{0}$ are not functionally independent because, as we have stated in Section 2, the tensor \mathbf{T} is degenerate. Nevertheless, since $\mathbf{v}_0(\mathbf{q}) \in T_{\mathbf{q}}S^3$, for any $\mathbf{q} \in S^3$, the system (3.3) can be described only by three equations plus the constraint equation.

In order to write $\mathbf{v}_{\mathbf{0}|S^3}(\mathbf{q}) = \mathbf{0}$ as a system of three independent equations, we will push forward the vector field $\mathbf{v}_{\mathbf{0}|S^3}$ through the covering map \mathbf{P} , which will also place us in the hypotheses of the step (iv') of the embedding algorithm. By definition of the push-forward operator, we have that

$$\mathbf{P}_* \mathbf{v}_{\mathbf{0}|S^3}(\mathbf{R}^{\mathbf{q}}) = T_{\mathbf{q}}\mathbf{P}(\mathbf{v}_{\mathbf{0}|S^3}(\mathbf{q})) \in T_{\mathbf{R}^{\mathbf{q}}}SO(3).$$

As \mathbf{P} is not an injective map, in order to obtain a vector field on the target space $SO(3)$, the equality $T_{\mathbf{q}}\mathbf{P}(\mathbf{v}_{\mathbf{0}|S^3}(\mathbf{q})) = T_{-\mathbf{q}}\mathbf{P}(\mathbf{v}_{\mathbf{0}|S^3}(-\mathbf{q}))$ must be satisfied. We will show in what follows that this is the case.

We need to compute the tangent map of the covering map $\mathbf{P} : S^3 \rightarrow SO(3)$. This map can be written as a restriction of the map $\tilde{\mathbf{P}} : \mathbb{R}^4 \rightarrow \mathbb{R}^9$ defined by $\tilde{\mathbf{P}}(q^0, q^1, q^2, q^3) = ((q^0)^2 + (q^1)^2 - (q^2)^2 - (q^3)^2, 2(q^1q^2 - q^0q^3), 2(q^1q^3 + q^0q^2), 2(q^1q^2 + q^0q^3), (q^0)^2 - (q^1)^2 + (q^2)^2 - (q^3)^2, 2(q^2q^3 - q^0q^1), 2(q^1q^3 - q^0q^2), 2(q^2q^3 + q^0q^1), (q^0)^2 - (q^1)^2 - (q^2)^2 + (q^3)^2)$, where a rotation matrix $\mathbf{R}^{\mathbf{q}}$ has been identified with a point in \mathbb{R}^9 (the first line is identified with the first three components and so on).

The matrix corresponding to the linear map $T_{\mathbf{q}}\tilde{\mathbf{P}} : \mathbb{R}^4 \rightarrow \mathbb{R}^9$ is given by

$$\text{Jac}_{\mathbf{q}}\tilde{\mathbf{P}} = 2 \begin{bmatrix} q^0 & q^1 & -q^2 & -q^3 \\ -q^3 & q^2 & q^1 & -q^0 \\ q^2 & q^3 & q^0 & q^1 \\ q^3 & q^2 & q^1 & q^0 \\ q^0 & -q^1 & q^2 & -q^3 \\ -q^1 & -q^0 & q^3 & q^2 \\ -q^2 & q^3 & -q^0 & q^1 \\ q^1 & q^0 & q^3 & q^2 \\ q^0 & -q^1 & -q^2 & q^3 \end{bmatrix}.$$

By direct computation, where we identify a vector in \mathbb{R}^9 with a tangent vector $\mathbf{R}^{\mathbf{q}}\Delta \in T_{\mathbf{R}^{\mathbf{q}}}SO(3) = \{\mathbf{R}^{\mathbf{q}}\Delta \mid \Delta \text{ is a } 3 \times 3 \text{ skew symmetric matrix}\}$, for $\mathbf{q} \in S^3$ we have that

$$\begin{aligned} T_{\mathbf{q}}\mathbf{P} \cdot \mathbf{v}_{\mathbf{0}|S^3}(\mathbf{q}) &= T_{\mathbf{q}}\tilde{\mathbf{P}} \cdot \mathbf{v}_{\mathbf{0}|S^3}(\mathbf{q}) \\ &= \sum_{i=1}^r \langle \mathbf{q}, \mathbf{q}_i \rangle T_{\mathbf{q}}\tilde{\mathbf{P}} \cdot (\langle \mathbf{q}, \mathbf{q} \rangle \mathbf{q}_i - \langle \mathbf{q}, \mathbf{q}_i \rangle \mathbf{q}) \\ &= \sum_{i=1}^r \langle \mathbf{q}, \mathbf{q}_i \rangle \text{Jac}_{\mathbf{q}}\tilde{\mathbf{P}} \cdot (\langle \mathbf{q}, \mathbf{q} \rangle \mathbf{q}_i - \langle \mathbf{q}, \mathbf{q}_i \rangle \mathbf{q}) \\ &\cong \sum_{i=1}^r \langle \mathbf{q}, \mathbf{q}_i \rangle \mathbf{R}^{\mathbf{q}} \Delta_i(\mathbf{q}) \\ &= \mathbf{R}^{\mathbf{q}} \sum_{i=1}^r \langle \mathbf{q}, \mathbf{q}_i \rangle \Delta_i(\mathbf{q}), \end{aligned}$$

where

$$\Delta_i(\mathbf{q}) = \begin{pmatrix} 0 & -q^0q_i^3 + q^1q_i^2 - q^2q_i^1 + q^3q_i^0 & q^0q_i^2 + q^1q_i^3 - q^2q_i^0 - q^3q_i^1 \\ q^0q_i^3 - q^1q_i^2 + q^2q_i^1 - q^3q_i^0 & 0 & -q^0q_i^1 + q^1q_i^0 + q^2q_i^3 - q^3q_i^2 \\ -q^0q_i^2 - q^1q_i^3 + q^2q_i^0 + q^3q_i^1 & q^0q_i^1 - q^1q_i^0 - q^2q_i^3 + q^3q_i^2 & 0 \end{pmatrix}.$$

By identifying the vector $\text{Jac}_{\mathbf{q}}\tilde{\mathbf{P}} \cdot (\langle \mathbf{q}, \mathbf{q} \rangle \mathbf{q}_i - \langle \mathbf{q}, \mathbf{q}_i \rangle \mathbf{q}) \in \mathbb{R}^9$ with a 3×3 matrix, the skew-symmetric matrix is given by the formula

$$\Delta_i(\mathbf{q}) = \mathbf{R}^{\mathbf{q}T} \text{Jac}_{\mathbf{q}}\tilde{\mathbf{P}} \cdot (\langle \mathbf{q}, \mathbf{q} \rangle \mathbf{q}_i - \langle \mathbf{q}, \mathbf{q}_i \rangle \mathbf{q}).$$

By noticing that $\Delta_i(-\mathbf{q}) = -\Delta_i(\mathbf{q})$, we obtain the equality $T_{\mathbf{q}}\mathbf{P}(\mathbf{v}_{\mathbf{0}|S^3}(\mathbf{q})) = T_{-\mathbf{q}}\mathbf{P}(\mathbf{v}_{\mathbf{0}|S^3}(-\mathbf{q}))$ and consequently, $\mathbf{P}_*\mathbf{v}_{\mathbf{0}|S^3}$ is a vector field on the manifold $SO(3)$.

As a result we obtain that the critical points of the cost function $G_{1_{S^3}}$ are the solutions of the following system (which is equivalent with (3.3)):

$$\begin{cases} \sum_{i=1}^r \langle \mathbf{q}, \mathbf{q}_i \rangle \Delta_i(\mathbf{q}) = \mathbf{0} \\ \langle \mathbf{q}, \mathbf{q} \rangle = 1 \end{cases} . \quad (3.4)$$

The advantage of the above system, compared with the system (3.3), is that we have only three equations of second degree instead of four equations of third degree plus the constraint equation.

If we transform the above system into rotations, by direct computation, we obtain:

$$\langle \mathbf{q}, \mathbf{q}_i \rangle \Delta_i(\mathbf{q}) = \frac{1}{4} (\mathbf{R}^{\mathbf{q}_i T} \mathbf{R}^{\mathbf{q}} - \mathbf{R}^{\mathbf{q} T} \mathbf{R}^{\mathbf{q}_i}) \quad (3.5)$$

and the above system becomes:

$$\sum_{i=1}^r (\mathbf{R}^{\mathbf{q}_i T} \mathbf{R}^{\mathbf{q}} - \mathbf{R}^{\mathbf{q} T} \mathbf{R}^{\mathbf{q}_i}) = \mathbf{0},$$

which is equivalent with

$$\overline{\mathbf{R}}^T \mathbf{R}^{\mathbf{q}} - \mathbf{R}^{\mathbf{q} T} \overline{\mathbf{R}} = \mathbf{0}, \quad (3.6)$$

where $\overline{\mathbf{R}} = \frac{1}{r} \sum_{i=1}^r \mathbf{R}^{\mathbf{q}_i} = \frac{1}{r} \sum_{i=1}^r \mathbf{R}_i$. Solving this equation corresponds to step (iv') of the embedding algorithm and the solutions are the critical points of the cost function $G_{1_{SO(3)}}$. Also, this equation is the same as the characterization for the Euclidean mean obtained in [19]. Thus, we have obtained that searching for critical points of the cost function $G_{1_{SO(3)}}$ can be performed in two ways. More precisely, we can solve (3.6) or we can transform the initial problem into quaternions and solve the system (3.4) and then transform the solutions into rotations.

II. Next, we will consider the geodesic distance on the Lie group $SO(3)$ which is defined by the angle of two rotations, see [1], [4], [14], [19], [21], [22]. For two rotations $\mathbf{R}_1, \mathbf{R}_2 \in SO(3)$, $d_2(\mathbf{R}_1, \mathbf{R}_2) = \|\text{Log}(\mathbf{R}_1^T \mathbf{R}_2)\|_F = \sqrt{2}|\theta|$, where $\theta \in (-\pi, \pi)$ is the angle between the rotations \mathbf{R}_1 and \mathbf{R}_2 . The associated cost function is $G_{2_{SO(3)}} : SO(3) \rightarrow \mathbb{R}$,

$$G_{2_{SO(3)}}(\mathbf{R}) = \sum_{i=1}^r \|\text{Log}(\mathbf{R}_i^T \mathbf{R})\|_F^2,$$

where $\mathbf{R}_1, \dots, \mathbf{R}_r$ are the sample rotation matrices.

As before, we will compute the lifted cost function $G_{2_{S^3}}$ as stated in step (i) of the embedding algorithm. More precisely, we have¹

$$\begin{aligned} G_{2_{S^3}}(\mathbf{q}) &= \sum_{i=1}^r \|\text{Log}(\mathbf{R}^{\mathbf{q}_i T} \mathbf{R}^{\mathbf{q}})\|_F^2 = 2 \sum_{i=1}^r \theta_i^2 \\ &= 2 \sum_{i=1}^r \arccos^2 \left(\frac{\text{tr}(\mathbf{R}^{\mathbf{q}_i T} \mathbf{R}^{\mathbf{q}}) - 1}{2} \right) \\ &\stackrel{(3.2)}{=} 2 \sum_{i=1}^r \arccos^2(2\langle \mathbf{q}_i, \mathbf{q} \rangle^2 - 1) \\ &= 2 \sum_{i=1}^r \arccos^2(|\langle \mathbf{q}_i, \mathbf{q} \rangle|). \end{aligned}$$

The lifted cost function $G_{2_{S^3}}$ does not depend on the choice of sample quaternions \mathbf{q}_i or $-\mathbf{q}_i$ that represent the same sample rotation \mathbf{R}_i .

¹ $|\theta_i| = \arccos \left(\frac{\text{tr}(\mathbf{R}^{\mathbf{q}_i T} \mathbf{R}^{\mathbf{q}}) - 1}{2} \right)$, if $x \in [-1, 1]$ then $\arccos(2x^2 - 1) = 2 \arccos(|x|)$.

For step (iii) of the embedding algorithm, the prolongation function is given by $G_2 : \mathbb{R}^4 \setminus \{\mathbf{0}\} \rightarrow \mathbb{R}$, $G_2(\mathbf{q}) = 2 \sum_{i=1}^r \arccos^2 \left(\frac{|\langle \mathbf{q}, \mathbf{q}_i \rangle|}{\|\mathbf{q}\| \cdot \|\mathbf{q}_i\|} \right)$. The coefficients of the differential 1-form dG_2 are given by:

$$\begin{aligned} \overline{dG_2}(\mathbf{q}) &= \left(\frac{\partial G_2}{\partial q^0}(\mathbf{q}), \dots, \frac{\partial G_2}{\partial q^3}(\mathbf{q}) \right) \\ &= \left(\dots, -4 \sum_{i=1}^r \frac{\operatorname{sgn}(\langle \mathbf{q}, \mathbf{q}_i \rangle) \|\mathbf{q}\|^2 q_i^j - |\langle \mathbf{q}, \mathbf{q}_i \rangle| q^j}{\|\mathbf{q}\|^2 \sqrt{\|\mathbf{q}\|^2 \|\mathbf{q}_i\|^2 - \langle \mathbf{q}, \mathbf{q}_i \rangle^2}} \arccos \left(\frac{|\langle \mathbf{q}, \mathbf{q}_i \rangle|}{\|\mathbf{q}\| \cdot \|\mathbf{q}_i\|} \right), \dots \right) \\ &= -4 \sum_{i=1}^r \arccos \left(\frac{|\langle \mathbf{q}, \mathbf{q}_i \rangle|}{\|\mathbf{q}\| \cdot \|\mathbf{q}_i\|} \right) \frac{\operatorname{sgn}(\langle \mathbf{q}, \mathbf{q}_i \rangle)}{\|\mathbf{q}\|^2 \sqrt{\|\mathbf{q}\|^2 \|\mathbf{q}_i\|^2 - \langle \mathbf{q}, \mathbf{q}_i \rangle^2}} (\langle \mathbf{q}, \mathbf{q} \rangle \mathbf{q}_i - \langle \mathbf{q}, \mathbf{q}_i \rangle \mathbf{q}). \end{aligned}$$

In order to solve the differentiability problem we will eliminate from the domain of definition of the function G_2 the hyperplanes $\Pi_i = \{\mathbf{q} \in \mathbb{R}^4 \mid \langle \mathbf{q}, \mathbf{q}_i \rangle = 0\}$ and the lines d_i that go through points $\mathbf{0}$ and \mathbf{q}_i .

In this case, the equation $\mathbf{v}_{\mathbf{0}|S^3}(\mathbf{q}) = \mathbf{0}$ is equivalent with

$$\begin{cases} \sum_{i=1}^r \frac{\operatorname{sgn}(\langle \mathbf{q}, \mathbf{q}_i \rangle) \arccos(|\langle \mathbf{q}, \mathbf{q}_i \rangle|)}{\sqrt{1 - \langle \mathbf{q}, \mathbf{q}_i \rangle^2}} (\langle \mathbf{q}, \mathbf{q} \rangle \mathbf{q}_i - \langle \mathbf{q}, \mathbf{q}_i \rangle \mathbf{q}) = \mathbf{0} \\ \mathbf{q} \in S^3 \setminus \bigcup_{i=1}^r \Pi_i \\ \mathbf{q} \neq \pm \mathbf{q}_i \end{cases}$$

By using the same arguments as before, the equation $\mathbf{v}_{\mathbf{0}|S^3}(\mathbf{q}) = \mathbf{0}$ is equivalent with $\mathbf{P}_* \mathbf{v}_{\mathbf{0}|S^3} = \mathbf{0}$ which has the following form

$$\begin{cases} \sum_{i=1}^r \frac{\operatorname{sgn}(\langle \mathbf{q}, \mathbf{q}_i \rangle) \arccos(|\langle \mathbf{q}, \mathbf{q}_i \rangle|)}{\sqrt{1 - \langle \mathbf{q}, \mathbf{q}_i \rangle^2}} \Delta_i(\mathbf{q}) = \mathbf{0} \\ \mathbf{q} \in S^3 \setminus \bigcup_{i=1}^r \Pi_i \\ \mathbf{q} \neq \pm \mathbf{q}_i \end{cases} \quad (3.7)$$

The above expression also shows that $\mathbf{P}_* \mathbf{v}_{\mathbf{0}|S^3}$ is a vector field on the manifold $SO(3)$.

In order to transform (3.7) into rotations we need the following computation:

$$\begin{aligned} \frac{\operatorname{sgn}(\langle \mathbf{q}, \mathbf{q}_i \rangle) \arccos(|\langle \mathbf{q}, \mathbf{q}_i \rangle|)}{\sqrt{1 - \langle \mathbf{q}, \mathbf{q}_i \rangle^2}} &= \frac{\arccos(|\langle \mathbf{q}, \mathbf{q}_i \rangle|) \langle \mathbf{q}, \mathbf{q}_i \rangle}{|\langle \mathbf{q}, \mathbf{q}_i \rangle| \sqrt{1 - \langle \mathbf{q}, \mathbf{q}_i \rangle^2}} = \frac{\frac{|\theta_i|}{2}}{|\cos(\frac{\theta_i}{2})| \sqrt{1 - \cos^2(\frac{\theta_i}{2})}} \langle \mathbf{q}, \mathbf{q}_i \rangle \\ &= \frac{|\theta_i|}{|\sin(\theta_i)|} \langle \mathbf{q}, \mathbf{q}_i \rangle = \frac{\theta_i}{\sin(\theta_i)} \langle \mathbf{q}, \mathbf{q}_i \rangle, \end{aligned}$$

where we have used the property $\mathbf{q} \in S^3 \setminus \bigcup_{i=1}^r \Pi_i$ which implies that $\langle \mathbf{q}, \mathbf{q}_i \rangle \neq 0$. Using (3.5) we obtain the equivalent system in rotations, and this corresponds to the step (iv') of the embedding algorithm,

$$\sum_{i=1}^r (\mathbf{R}^{\mathbf{q}_i T} \mathbf{R}^{\mathbf{q}} - \mathbf{R}^{\mathbf{q} T} \mathbf{R}^{\mathbf{q}_i}) \frac{\theta_i}{\sin \theta_i} = \mathbf{0} \Leftrightarrow \sum_{i=1}^r \operatorname{Log}(\mathbf{R}^{\mathbf{q}_i T} \mathbf{R}^{\mathbf{q}}) = \mathbf{0}. \quad (3.8)$$

The above equation has been also obtained in [19] as a characterization of the Riemannian mean. This was to be expected, as we are averaging the same cost function by two different methods that naturally lead to the same result. Also note that the cost function $G_{2SO(3)}$ is not well defined for angles $\theta_i = \pm\pi$ or equivalently $\langle \mathbf{q}, \mathbf{q}_i \rangle = 0$, a situation in which the prolongation function G_2 is not differentiable.

III. Another distance used on the Lie group $SO(3)$ is given by $d_3(\mathbf{R}_1, \mathbf{R}_2) = 1 - \frac{1}{2}\sqrt{\text{tr}(\mathbf{R}_1^T \mathbf{R}_2) + 1}$. This distance appears in [17], where the functional dependence between d_3 and d_2 is also given. We notice that this distance function on $SO(3)$ can be obtained from the pseudodistance $d_3 : S^3 \times S^3 \rightarrow \mathbb{R}$, $d_3(\mathbf{q}_1, \mathbf{q}_2) = 1 - |\langle \mathbf{q}_1, \mathbf{q}_2 \rangle|$. The cost function $G_{3_{SO(3)}}(\mathbf{R}) = \sum_{i=1}^r d_3^2(\mathbf{R}, \mathbf{R}_i)$ is lifted to the cost function

$$G_{3_{S^3}}(\mathbf{q}) = \sum_{i=1}^r (1 - |\langle \mathbf{q}, \mathbf{q}_i \rangle|)^2.$$

The prolongation function is $G_3 : \mathbb{R}^4 \rightarrow \mathbb{R}$, $G_3(\mathbf{q}) = \sum_{i=1}^r (1 - |\langle \mathbf{q}, \mathbf{q}_i \rangle|)^2$. The coefficients of the differential 1-form dG_3 are given by:

$$\overline{dG_3}(\mathbf{q}) = -2 \sum_{i=1}^r (1 - |\langle \mathbf{q}, \mathbf{q}_i \rangle|) \text{sgn}(\langle \mathbf{q}, \mathbf{q}_i \rangle) \mathbf{q}_i,$$

where, in order to have differentiability, we restrict our selves to the open set $\mathbf{q} \in S^3 \setminus \bigcup_{i=1}^r \Pi_i$ (the set Π_i is the hyperplane determined by \mathbf{q}_i as before).

The system of equations (2.4) becomes

$$\begin{cases} \sum_{i=1}^r (1 - |\langle \mathbf{q}, \mathbf{q}_i \rangle|) \text{sgn}(\langle \mathbf{q}, \mathbf{q}_i \rangle) (\langle \mathbf{q}, \mathbf{q} \rangle \mathbf{q}_i - \langle \mathbf{q}, \mathbf{q}_i \rangle \mathbf{q}) = \mathbf{0} \\ \mathbf{q} \in S^3 \setminus \bigcup_{i=1}^r \Pi_i \end{cases} \quad (3.9)$$

and this represents the equation $\mathbf{v}_0 = \mathbf{0}$ restricted to the open set $S^3 \setminus \bigcup_{i=1}^r \Pi_i$ of the sphere S^3 .

As in the previous cases, after applying the projection operator \mathbf{P}_* , we generate the vector field $\mathbf{P}_* \mathbf{v}_0|_{S^3}$ and consequently, we obtain the equivalent system of equations:

$$\begin{cases} \sum_{i=1}^r (1 - |\langle \mathbf{q}, \mathbf{q}_i \rangle|) \text{sgn}(\langle \mathbf{q}, \mathbf{q}_i \rangle) \Delta_i(\mathbf{q}) = \mathbf{0} \\ \mathbf{q} \in S^3 \setminus \bigcup_{i=1}^r \Pi_i \end{cases} \quad (3.10)$$

Transforming the above system into rotations, after some algebraic manipulations using (3.5) and (3.2), we obtain the equation from the step (\mathbf{iv}') of the embedding algorithm

$$\sum_{i=1}^r \left(\frac{2}{\sqrt{\text{tr}(\mathbf{R}^{\mathbf{q}_i T} \mathbf{R}^{\mathbf{q}_i})} + 1} - 1 \right) (\mathbf{R}^{\mathbf{q}_i T} \mathbf{R}^{\mathbf{q}} - \mathbf{R}^{\mathbf{q} T} \mathbf{R}^{\mathbf{q}_i}) = \mathbf{0}.$$

IV. Next we will apply the techniques developed so far for a cost function that is of L^p -type but not of least square type, see [15]. We start with the distance $d_1 : SO(3) \times SO(3) \rightarrow \mathbb{R}_+$, $d_1(\mathbf{R}_1, \mathbf{R}_2) = \|\mathbf{R}_1 - \mathbf{R}_2\|_F$. The associated L^4 -mean cost function is $G_{4_{SO(3)}} : SO(3) \rightarrow \mathbb{R}$,

$$G_{4_{SO(3)}}(\mathbf{R}) = \sum_{i=1}^r \|\mathbf{R} - \mathbf{R}_i\|_F^4.$$

The lifted cost function is given by $G_{4_{S^3}} : S^3 \rightarrow \mathbb{R}$,

$$G_{4_{S^3}}(\mathbf{q}) = 64 \sum_{i=1}^r (1 - \langle \mathbf{q}, \mathbf{q}_i \rangle)^2.$$

The prolongation function is $G_4 : \mathbb{R}^4 \rightarrow \mathbb{R}$, $G_4(\mathbf{q}) = 64 \sum_{i=1}^r (1 - \langle \mathbf{q}, \mathbf{q}_i \rangle)^2$.

As in the case of the function G_1 , the system (2.4) corresponding to the function G_4 becomes:

$$\begin{cases} \sum_{i=1}^r (\langle \mathbf{q}, \mathbf{q}_i \rangle - \langle \mathbf{q}, \mathbf{q}_i \rangle^3) (\langle \mathbf{q}, \mathbf{q} \rangle \mathbf{q}_i - \langle \mathbf{q}, \mathbf{q}_i \rangle \mathbf{q}) = \mathbf{0} \\ \langle \mathbf{q}, \mathbf{q} \rangle = 1 \end{cases} \quad (3.11)$$

By using the same arguments as before, in the case of function G_4 , the equation $\mathbf{v}_{0|S^3}(\mathbf{q}) = \mathbf{0}$ is equivalent with $\mathbf{P}_* \mathbf{v}_{0|S^3} = \mathbf{0}$, which has the following form

$$\begin{cases} \sum_{i=1}^r (1 - \langle \mathbf{q}, \mathbf{q}_i \rangle^2) \langle \mathbf{q}, \mathbf{q}_i \rangle \Delta_i(\mathbf{q}) = \mathbf{0} \\ \langle \mathbf{q}, \mathbf{q} \rangle = 1 \end{cases} \quad (3.12)$$

Using again (3.2) and (3.5) we obtain, according to step (iv') of the embedding algorithm, the equation in rotations that gives the critical points of the cost function $G_{4_{SO(3)}}$,

$$\sum_{i=1}^r (3 - \text{tr}(\mathbf{R}^{\mathbf{q}T} \mathbf{R}^{\mathbf{q}_i})) (\mathbf{R}^{\mathbf{q}_i T} \mathbf{R}^{\mathbf{q}} - \mathbf{R}^{\mathbf{q}T} \mathbf{R}^{\mathbf{q}_i}) = \mathbf{0}.$$

Acknowledgements. Petre Birtea and Dan Comănescu have been supported by the grant of the Romanian National Authority for Scientific Research, CNCS UEFISCDI, project number PN-II-RU-TE-2011-3-0006.

References

- [1] **S.L. Altman**, *Rotations, quaternions, and double groups*, Oxford University Press, 1986.
- [2] **P.A. Absil, R. Mahony, R. Sepulchre**, *Riemannian geometry of Grassmann manifolds with a view on algorithmic computation*, Acta Applicandae Mathematicae, Vol. 80, Issue 2 (2004), pp. 199-220.
- [3] **P.A. Absil, R. Mahony, R. Sepulchre**, *Optimization Algorithms on Matrix Manifolds*, Princeton University Press, 2008.
- [4] **V. Arsigny, X. Pennec, and N. Ayache**, *Bi-invariant means in Lie groups. Application to left-invariant polyaffine transformations*, INRIA research report, 5885 (2006).
- [5] **P. Birtea, D. Comănescu**, *Asymptotic Stability of Dissipated Hamilton-Poisson Systems*, SIAM J. Appl. Dyn. Syst., Vol. 8, Issue 3 (2009), pp. 967-976.
- [6] **P. Birtea, D. Comănescu**, *Geometric Dissipation for dynamical systems*, Comm. Math. Phys., Vol. 316, Issue 2 (2012), pp. 375-394.
- [7] **D. Comănescu, S.S. Dragomir**, *On the Torricellian Point in Inner Product Spaces*, Demonstratio Mathematica, Vol. XLI, Issue 3 (2008), pp. 639-650.
- [8] **D. Comănescu, S.S. Dragomir, E. Kikianty**, *On the Torricellian Points in Normed Linear Spaces and Applications*, (2012), <http://ajmaa.org/RGMIA/papers/v15/v15a3.pdf>
- [9] **A. Edelman, T.A. Arias, S.T. Smith**, *The geometry of algorithms with orthogonality constraints*, SIAM J. Matrix Anal. Appl., Vol. 20, Issue 2 (1998), pp. 303-353.
- [10] **S. Fiori**, *Lie-group-type neural system learning by manifold retractions*, Neural Networks, Vol. 21 (2008), pp. 1524-1529.

- [11] **S. Fiori**, *Solving Minimal-Distance Problems over the Manifold of Real-Symplectic Matrices*, SIAM J. Matrix Anal. & Appl., Vol. 32, Issue 3 (2011), pp. 938-968.
- [12] **S. Fiori, T. Tanaka**, *An algorithm to compute averages on matrix Lie groups*, IEEE Trans. Signal Process., Vol. 57, no. 12 (2009), pp. 4734-4743.
- [13] **V. M. Govindu**, *Lie-Algebraic Averaging For Globally Consistent Motion Estimation*, Proc. IEEE Conf. on Computer Vision and Pattern Recognition, Washington, DC, Vol. 1 (2004), pp. 684-691.
- [14] **C. Gramkow**, *On Averaging Rotations*, J. Math. Imaging Vis., Vol. 15 (2001), pp. 7-16.
- [15] **R. Hartley, J. Trumpf, Y. Dai**, *Rotation averaging and weak convexity*, International Symposium on Mathematical Theory of Networks and Systems, Piscataway, USA (2010), pp. 2435-2442.
- [16] **U. Helmke, M.A. Shayman**, *Critical Points of Matrix Least Squares Distance Functions*, Linear Algebra and its Applications, Vol. 215 (1995), pp. 1-19.
- [17] **D.Q. Huynh**, *Metrics for 3D Rotations: Comparison and Analysis*, J. Math. Imaging Vis., Vol. 35 (2009), pp. 155-164.
- [18] **F.L. Markley, Y. Cheng, J.L. Crassidis, Y. Oshman**, *Averaging Quaternions*, Journal of Guidance, Control, and Dynamics, Vol. 30, Issue 4 (2007), pp. 1193-1197.
- [19] **M. Moakher**, *Means and averaging in the group of rotations*, SIAM J. Matrix Anal. Appl., Vol. 24, Issue 1 (2002), pp. 1-16.
- [20] **P.L. Papini, J. Puerto**, *Averaging the k largest distances among n : k -centra in Banach spaces*, J. Math. Anal. Appl., Vol. 291, Issue 2 (2004), pp. 477-487.
- [21] **X. Pennec**, *Computing the mean of geometric features - Application to the mean rotation*, Institut National de Recherche en Informatique et en Automatique, Rapport des Recherche 3371, 1998.
- [22] **I. Sharf, A. Wolf, M.B. Rubin**, *Arithmetic and geometric solutions for average rigid-body rotation*, Mechanism and Machine Theory, Vol. 45 (2010), pp. 1239-1251.
- [23] **M.D. Shuster**, *A Survey of Attitude Representations*, Journal of the Astronautical Sciences, Vol. 41, Issue 4 (1993), pp. 439-517.
- [24] **C. Samir, P.A. Absil, A. Srivastava, E. Klassen**, *A Gradient-Descent Method for Curve Fitting on Riemannian Manifolds*, Foundations of Computational Mathematics, Vol. 12, Issue 1 (2012), pp. 49-73.
- [25] **R. Subbarao, P. Meer**, *Nonlinear Mean Shift over Riemannian Manifolds*, Int. J. Comput. Vis., Vol. 84 (2009), pp. 1-20.
- [26] **B. Vandereycken, P.A. Absil, S. Vandewalle**, *A Riemannian geometry with complete geodesics for the set of positive semidefinite matrices of fixed rank*, IMA Journal of Numerical Analysis, (2012), doi:10.1093/imanum/drnxxx