CLASSIFYING τ -TILTING MODULES OVER PREPROJECTIVE ALGEBRAS OF DYNKIN TYPE

YUYA MIZUNO

ABSTRACT. We study support τ -tilting modules over preprojective algebras of Dynkin type. We classify basic support τ -tilting modules by giving a bijection with elements in the corresponding Weyl groups. Moreover we show that they are in bijection with the set of torsion classes, the set of torsion-free classes and other important objects in representation theory. We also study g-matrices of support τ -tilting modules, which show terms of minimal projective presentations of indecomposable direct summands. We give an explicit description of g-matrices and prove that cones given by g-matrices coincide with chambers of the associated root systems.

CONTENTS

Introduction	1
1. Preliminaries	4
1.1. Preprojective algebras	4
1.2. Support τ -tilting modules	5
1.3. Torsion classes	6
2. Support τ -tilting ideals and the Weyl group	7
2.1. Support τ -tilting ideals	7
2.2. Bijection between support τ -tilting modules and the Weyl group	12
2.3. Partial orders of support τ -tilting modules and the Weyl group	16
3. <i>g</i> -matrices and cones	18
4. Further connections	22
References	24

INTRODUCTION

Preprojective algebras first appeared in the work of Gelfand-Ponomarev [GP]. Since then, they have been one of the important objects in the representation theory of algebras, for example [DR1, BGL, DR2]. Preprojective algebras appear also in many branches of mathematics. For instance, they play central roles in Lusztig's Lagrangian construction of semicanonical basis [L1, L2].

In [BIRS] (see also [IR]), the authors studied preprojective algebras of non-Dynkin quivers via tilting theory. As an application, they succeeded to construct a large class of 2-Calabi-Yau categories which have close connections with cluster algebras. On the other hand, in Dynkin cases (i.e. the underlying graph of a quiver is A_n $(n \ge 1)$, D_n $(n \ge 4)$ and E_n (n = 6, 7, 8)), the preprojective algebras are selfinjective, so that all tilting modules are trivial (i.e. projective) and they are not effective tools. In this paper, instead of tilting

The author is supported by Grant-in-Aid for JSPS Fellowships No.23.5593.

modules, we will study support τ -tilting modules (Definition 1.3) over the preprojective algebras of Dynkin type.

The notion of support τ -tilting modules was introduced in [AIR], as a generalization of tilting modules. Support τ -tilting modules have several nice properties. For example, it is shown that there are deep connections between τ -tilting theory, torsion theory, silting theory and cluster tilting theory. Support τ -tilting modules also have nicer mutation properties than tilting modules, that is, mutation of support τ -tilting modules is always possible (Definition-Theorem 1.5). Moreover, support τ -tilting modules over selfinjective algebras are useful to provide tilting complexes [M]. It is therefore fruitful to investigate these remarkable modules for preprojective algebras. To explain our results, we give the following set-up.

Let Q be a finite connected acyclic quiver with the set $Q_0 = \{1, \ldots, n\}$ of vertices, Λ the preprojective algebra of Q and I_i the two-sided ideal of Λ generated by $1 - e_i$, where e_i is an idempotent of Λ corresponding to $i \in Q_0$. These ideals are quite useful to study structure of categories [IR, BIRS, AIRT, ORT]. They also play important roles in Geiss-Leclerc-Schröer's construction of cluster monomials of certain types of cluster algebras [GLS1, GLS3] and Baumann-Kamnitzer-Tingley's theory of affine MV polytopes [BKT].

If Q is non-Dynkin, then I_i is a tilting module. On the other hand, if Q is Dynkin, then I_i is never a tilting module. Instead, we show that I_i is a support τ -tilting module (Lemma 2.3) and this observation is crucial in this paper. We denote by $\langle I_1, \ldots, I_n \rangle$ the set of ideals of Λ of the form $I_{i_1}I_{i_2}\cdots I_{i_k}$ for some $k \geq 0$ and $i_1, \ldots, i_k \in Q_0$.

Our main goal is to give the following bijections which provide a complete description of support τ -tilting modules over preprojective algebras of Dynkin type.

Theorem 0.1 (Theorem 2.21). Let Q be a Dynkin quiver with the set $Q_0 = \{1, \ldots, n\}$ of vertices and Λ the preprojective algebra of Q. There are bijections between the following objects.

- (a) The elements of the Weyl group W_Q associated with Q.
- (b) The set $\langle I_1, \ldots, I_n \rangle$.
- (c) The set $s\tau$ -tilt Λ of isomorphism classes of basic support τ -tilting Λ -modules.
- (d) The set $s\tau$ -tilt Λ^{op} of isomorphism classes of basic support τ -tilting Λ^{op} -modules.

These bijections give a strong relationship between the representation theory of algebras and root systems as below. It is known that support τ -tilting modules have a natural structure of partially ordered set (Definition 1.4). We relate this partial order with that of the Weyl group.

Theorem 0.2 (Theorem 2.30). The bijection $W_Q \to s\tau$ -tilt Λ in Theorem 0.1 gives an isomorphism of partially ordered sets (W_Q, \leq_L) and $(s\tau$ -tilt $\Lambda, \leq)^{\text{op}}$, where \leq_L is the left weak order on W_Q .

We also study the *g*-matrix (Definition 3.1) of support τ -tilting module T, which shows terms of minimal projective presentations of each indecomposable direct summand of T. By *g*-matrices, we define *cones*, which provide a geometric realization of the simplicial complex of support τ -tilting modules. We give a complete description of *g*-matrices in terms of reflection transformations and show that their cones coincide with chambers of the associated root system. The results are summarized as follows, where $\mathbf{e}_1, \ldots, \mathbf{e}_n$ is the canonical basis of \mathbb{Z}^n , $C_0 := \{a_1\mathbf{e}_1 + \cdots + a_n\mathbf{e}_n \mid a_i \in \mathbb{R}_{>0}\}$ and $\sigma^* : W_Q \to GL_n(\mathbb{Z})$ is the contragradient of the geometric representation (Definition 3.5). **Theorem 0.3** (Theorem 3.8). Let Q be a Dynkin quiver with the set $Q_0 = \{1, ..., n\}$ of vertices and Λ the preprojective algebra of Q.

- (a) The set of g-matrices of support τ -tilting Λ -modules coincides with $\sigma^*(W_O)$.
- (b) For any $w \in W_Q$, we have

$$C(w) = \sigma^*(w)(C_0),$$

where C(w) denotes the cone of the support τ -tilting module I_w (Definition 3.1). In particular, cones of basic support τ -tilting Λ -modules give chambers of the associated root system.

The above results give several important applications. Among others we conclude that there are only finitely many basic support τ -tilting Λ -modules. This fact yields bijections between basic support τ -tilting Λ -modules, torsion classes in mod Λ and torsion-free classes in mod Λ by a result in [IJ]. Then we can give complete descriptions of these two classes in terms of W_Q (Proposition 4.2). This seems to be interesting by itself since almost all preprojective algebras are of wild representation type.

As another application, by combining with results of [AIR], [KY, BY] and [ORT], we extend bijections of Theorem 0.1 as follows.

Corollary 0.4 (Corollary 4.1). *Objects of Theorem 0.1 are in bijection with the following objects.*

- (e) The set of torsion classes in $mod\Lambda$.
- (f) The set of torsion-free classes in $mod\Lambda$.
- (g) The set of isomorphism classes of basic two-term silting complexes in $\mathsf{K}^{\mathsf{b}}(\mathsf{proj}\Lambda)$.
- (h) The set of intermediate bounded co-t-structures in $K^{b}(\text{proj}\Lambda)$ with respect to the standard co-t-structure.
- (i) The set of intermediate bounded t-structures in D^b(modΛ) with length heart with respect to the standard t-structure.
- (j) The set of isomorphism classes of two-term simple-minded collections in $D^{b}(mod\Lambda)$.
- (k) The set of quotient closed subcategories in modKQ.
- (1) The set of subclosed subcategories in modKQ.

We now describe the organization of this paper.

In section 1, we recall definitions of preprojective algebras, support τ -tilting modules and torsion classes.

In section 2.1, we introduce support τ -tilting ideals (Definition 2.1), which play key roles in this paper. Moreover, we prove that any element of $\langle I_1, \ldots, I_n \rangle$ is a basic support τ -tilting ideal (Theorem 2.2). In section 2.2, we prove that elements of W_Q correspond bijectively with support τ -tilting modules (Theorem 2.21). Using this, we give a description of mutations of support τ -tilting modules in terms of the Weyl group W_Q . We also determine the annihilators of all support τ -tilting modules. In section 2.3, we give an (anti)isomorphism between a partial order of W_Q and that of $s\tau$ -tilt Λ (Theorem 2.30).

In section 3, we deal with g-matrices of support τ -tilting modules. We give an explicit description of them in terms of reflection transformations. Furthermore, we study cones of support τ -tilting modules and show that they coincide with chambers of the associated root system.

In section 4, we show Corollary 0.4. In particular, we give complete descriptions of torsion classes and torsion-free classes (Proposition 4.2). Furthermore, we explain a relationship between our results and other works.

Notations. Let K be an algebraically closed field and we denote by $D := \operatorname{Hom}_K(-, K)$. By a finite dimensional algebra Λ , we mean a basic finite dimensional algebra over K. By a module, we mean a right module unless stated otherwise. We denote by $\operatorname{mod}\Lambda$ the category of finitely generated Λ -modules and by $\operatorname{proj}\Lambda$ the category of finitely generated projective Λ -modules. The composition gf means first f, then g. We denote by Q_0 the set of vertices and by Q_1 the set of arrows of a quiver Q. For an arrow $a \in Q_1$, we denote by s(a) and e(a) the start and end vertices of a respectively. For $X \in \operatorname{mod}\Lambda$, we denote by $\operatorname{Sub}X$ (respectively, $\operatorname{Fac}X$) the subcategory of $\operatorname{mod}\Lambda$ whose objects are submodules (respectively, factor modules) of finite direct sums of copies of X. We denote by $\operatorname{add}M$ the subcategory of $\operatorname{mod}\Lambda$ consisting of direct summands of finite direct sums of copies of $M \in \operatorname{mod}\Lambda$.

Acknowledgement. First and foremost, the author would like to thank Osamu Iyama for his support and patient guidance. He would like to express his gratitude to Steffen Oppermann, who kindly explain results of his paper. He is grateful to Joseph Grant, Laurent Demonet and Dong Yang for answering questions and helpful comments. He thanks Kota Yamaura, Takahide Adachi and Gustavo Jasso for their help and stimulating discussions. The author is very grateful to the anonymous referee for valuable comments, especially for suggesting the better terms and sentences.

1. Preliminaries

1.1. **Preprojective algebras.** In this subsection, we recall definitions and some properties of preprojective algebras. We refer to [BBK, Ri] for basic properties and background information.

Definition 1.1. Let Q be a finite connected acyclic quiver with vertices $Q_0 = \{1, \ldots, n\}$. The preprojective algebra associated to Q is the algebra

$$\Lambda = K \overline{Q} / \langle \sum_{a \in Q_1} (aa^* - a^*a) \rangle$$

where \overline{Q} is the double quiver of Q, which is obtained from Q by adding for each arrow $a: i \to j$ in Q_1 an arrow $a^*: i \leftarrow j$ pointing in the opposite direction.

If Q is a non-Dynkin quiver, the bounded derived category $\mathsf{D}^{\mathsf{b}}(\mathsf{fd}\,\Lambda)$ of the category $\mathsf{fd}\,\Lambda$ of finite dimensional modules are 2-Calabi-Yau (2-CY for short), that is, there is a functorial isomorphism

$$D\operatorname{Hom}_{\mathsf{D}^{\mathrm{b}}(\mathsf{fd}\,\Lambda)}(X,Y)\cong\operatorname{Hom}_{\mathsf{D}^{\mathrm{b}}(\mathsf{fd}\,\Lambda)}(Y,X[2]).$$

We refer to [AR2, Boc, CB] and see [GLS2] for the detailed proofs. On the other hand, the algebra Λ is finite-dimensional selfinjective if Q is a Dynkin quiver.

In this paper, we use the two-sided ideal I_i of Λ generated by $1 - e_i$, where e_i is a primitive idempotent of Λ for $i \in Q_0$. It is easy to see that the ideal has the following property, where we denote by $S_i := \Lambda/I_i$.

Lemma 1.2. Let Q be a finite connected acyclic quiver, Λ the preprojective algebra of Q and $X \in \text{mod}\Lambda$. Then XI_i is maximal amongst submodules Y of X such that any composition factor of X/Y is isomorphic to a simple module S_i .

1.2. Support τ -tilting modules. We recall the definition of support τ -tilting modules. We refer to [AIR] for the details about support τ -tilting modules.

Let Λ be a finite dimensional algebra and τ denote the AR translation [ARS].

Definition 1.3. (a) We call X in mod $\Lambda \tau$ -rigid if Hom_{Λ} $(X, \tau X) = 0$.

- (b) We call X in mod $\Lambda \tau$ -tilting (respectively, almost complete τ -tilting) if X is τ -rigid and $|X| = |\Lambda|$ (respectively, $|X| = |\Lambda| 1$), where |X| denotes the number of non-isomorphic indecomposable direct summands of X.
- (c) We call X in mod Λ support τ -tilting if there exists an idempotent e of Λ such that X is a τ -tilting $(\Lambda/\langle e \rangle)$ -module.
- (d) We call a pair (X, P) of $X \in \mathsf{mod}\Lambda$ and $P \in \mathsf{proj}\Lambda \tau$ -rigid if X is τ -rigid and $\operatorname{Hom}_{\Lambda}(P, X) = 0$.
- (e) We call a τ -rigid pair (X, P) a support τ -tilting (respectively, almost complete support τ -tilting) pair if $|X| + |P| = |\Lambda|$ (respectively, $|X| + |P| = |\Lambda| 1$).

We call (X, P) basic if X and P are basic, and we say that (X, P) is a direct summand of (X', P') if X is a direct summand of X' and P is a direct summand of P'. Note that (X, P) is a τ -rigid (respectively, support τ -tilting) pair for Λ if and only if X is a τ -rigid (respectively, τ -tilting) $(\Lambda/\langle e \rangle)$ -module, where e is an idempotent of Λ such that addP = adde Λ [AIR, Proposition 2.3]. Moreover, if (X, P) and (X, P') are support τ tilting pairs for Λ , then we get addP = addP'. Thus, a basic support τ -tilting module X determines a basic support τ -tilting pair (X, P) uniquely and we can identify basic support τ -tilting modules with basic support τ -tilting pairs.

We denote by $s\tau$ -tilt Λ the set of isomorphism classes of basic support τ -tilting Λ -modules.

Next we recall some properties of support τ -tilting modules. The set of support τ -tilting modules has a natural partial order as follows.

Definition 1.4. Let Λ be a finite dimensional algebra. For $T, T' \in s\tau$ -tilt Λ , we write

 $T' \ge T$

if $\operatorname{Fac} T' \supset \operatorname{Fac} T$. Then \geq gives a partial order on $s\tau$ -tilt Λ [AIR, Theorem 2.18].

Then we give the following results, which play significant roles in this paper.

Definition-Theorem 1.5. [AIR, Theorem 2.18, 2.28 and 2.30] Let Λ be a finite dimensional algebra. Then

(i) any basic almost support τ -tilting pair (U, Q) is a direct summand of exactly two basic support τ -tilting pairs (T, P) and (T', P'). Moreover, we have T > T' or T < T'.

Under the above setting, let X be an indecomposable Λ -module satisfying either $T = U \oplus X$ or $P = Q \oplus X$. We write $(T', P') = \mu_{(X,0)}(T, P)$ if X is a direct summand of T and $(T', P') = \mu_{(0,X)}(T, P)$ if X is a direct summand of P, and we say that (T', P') is a *mutation* of (T, P). In particular, we say that (T', P') is a *left mutation* (respectively, right mutation) of (T, P) if T > T' (respectively, if T < T') and write μ^- (respectively, μ^+). By (i), exactly one of the left mutation or right mutation occurs.

Now, assume that X is a direct summand of T and $T = U \oplus X$. In this case, for simplicity, we write a left mutation $T' = \mu_X^-(T)$ and a right mutation $T' = \mu_X^+(T)$.

Then, the following conditions are equivalent.

(ii) (a) T > T' (i.e. $T' = \mu_X^-(T)$).

(b) $X \notin \mathsf{Fac}U$.

Furthermore, if T is a τ -tilting Λ -module and (ii) holds, then we obtain the following result.

- (iii) Let $X \xrightarrow{f} U' \to Y \to 0$ be an exact sequence, where f is a minimal left $(\mathsf{add}U)$ -approximation. Then one of the following holds.
 - (a) Y = 0 and $\mu_X^-(T) \cong U$ is a basic support τ -tilting Λ -module.
 - (b) $Y \neq 0$ and Y is a direct sum of copies of an indecomposable Λ -module Y_1 , which is not in addT, and $\mu_X^-(T) \cong Y_1 \oplus U$ is a basic τ -tilting Λ -module.

Finally, we define the following useful quiver.

Definition 1.6. Let Λ be a finite dimensional algebra. We define the support τ -tilting quiver $\mathcal{H}(s\tau$ -tilt Λ) as follows.

- The set of vertices is $s\tau$ -tilt Λ .
- Draw an arrow from T to T' if T' is a left mutation of T (i.e. $T' = \mu_X^-(T)$).

The following theorem relates $\mathcal{H}(s\tau-\text{tilt}\Lambda)$ with partially orders of $s\tau-\text{tilt}\Lambda$.

Theorem 1.7. [AIR, Corollary 2.34] The support τ -tilting quiver $\mathcal{H}(s\tau$ -tilt Λ) is the Hasse quiver of the partially ordered set $s\tau$ -tilt Λ .

1.3. Torsion classes. The notion of torsion classes has been extensively studied in the representation theory. As we will see below, support τ -tilting modules have a close relationship with torsion classes.

Definition 1.8. Let Λ be a finite dimensional algebra and \mathcal{T} a full subcategory in mod Λ .

- (a) We call \mathcal{T} torsion class (respectively, torsion-free class) if it is closed under factor modules (respectively, submodules) and extensions.
- (b) We call \mathcal{T} a contravariantly finite subcategory in mod Λ if for each $X \in \text{mod}\Lambda$, there is a map $f: T \to X$ with $T \in \mathcal{T}$ such that $\text{Hom}_{\Lambda}(T', T) \xrightarrow{\cdot f} \text{Hom}_{\Lambda}(T', X)$ is surjective for all $T' \in \mathcal{T}$. Dually, a covariantly finite subcategory is defined. We call \mathcal{T} a functorially finite subcategory if it is contravariantly and covariantly finite.

We denote by f-tors Λ the set of functorially finite torsion classes in mod Λ .

Then, we have the following result, where we denote by $P(\mathcal{T})$ the direct sum of one copy of each of the indecomposable *Ext-projective* objects in \mathcal{T} (i.e. $\{X \in \mathcal{T} \mid \text{Ext}^{1}_{\Lambda}(X, \mathcal{T}) = 0\}$) up to isomorphism.

Theorem 1.9. [AIR] For a finite dimensional algebra Λ , there are mutually inverse bijections

 $\mathrm{s}\tau\text{-tilt}\Lambda\longleftrightarrow\mathrm{f}\text{-tors}\Lambda$

given by $s\tau$ -tilt $\Lambda \ni T \mapsto \mathsf{Fac}T \in \mathrm{f-tors}\Lambda$ and $\mathrm{f-tors}\Lambda \ni T \mapsto P(T) \in s\tau$ -tilt Λ .

Thus, support τ -tilting modules are quite useful to investigate functorially finite torsion classes. Moreover, we have the following much stronger result under the assumption that $|s\tau$ -tilt $\Lambda| < \infty$.

Theorem 1.10. [IJ] For a finite dimensional algebra Λ , the following conditions are equivalent.

- (a) $|s\tau \text{tilt}\Lambda| < \infty$.
- (b) Any torsion class in $mod\Lambda$ is functorially finite.
- (c) Any torsion-free class in $mod\Lambda$ is functorially finite.

Therefore, if there are only finitely many basic support τ -tilting modules, then Theorem 1.9 give a bijection between $s\tau$ -tilt Λ and torsion classes of mod Λ .

2. Support τ -tilting ideals and the Weyl group

The aim of this section is to give a complete description of all support τ -tilting modules over preprojective algebras of Dynkin type. We also study their several properties.

Throughout this section, unless otherwise specified, let Q be a Dynkin quiver with $Q_0 = \{1, \ldots, n\}$, Λ the preprojective algebra of Q and $I_i := \Lambda(1 - e_i)\Lambda$ for $i \in Q_0$. We denote by $\langle I_1, \ldots, I_n \rangle$ the set of ideals of Λ which can be written as

$$I_{i_1}I_{i_2}\cdots I_{i_k}$$

for some $k \geq 0$ and $i_1, \ldots, i_k \in Q_0$.

2.1. Support τ -tilting ideals. In this subsection, we introduce support τ -tilting ideals. We will show that any element of $\langle I_1, \ldots, I_n \rangle$ is a basic support τ -tilting ideal. This fact plays a key role in this paper.

We start with the following definition.

Definition 2.1. We call a two-sided ideal I of Λ support τ -tilting if I is a left support τ -tilting Λ -module and a right support τ -tilting Λ -module.

The aim of this subsection is to prove the following result.

Theorem 2.2. Any $T \in \langle I_1, \ldots, I_n \rangle$ is a basic support τ -tilting ideal of Λ .

First, we recall some properties for later use (see [BBK]). For any $i \in Q_0$, we can take a minimal projective presentation of $S_i := \Lambda/I_i$ as follows

$$e_i\Lambda \xrightarrow{p_2} \bigoplus_{a \in \overline{Q}_1, e(a)=i} e_{s(a)}\Lambda \xrightarrow{p_1} e_i\Lambda \xrightarrow{p_0} S_i \longrightarrow 0.$$

Since $\text{Im} p_1 = e_i I_i$, we have the following exact sequences

(1)
$$0 \longrightarrow e_i I_i \xrightarrow{\iota} e_i \Lambda \xrightarrow{p_0} S_i \longrightarrow 0.$$

(2)
$$e_i \Lambda \xrightarrow{p_2} \bigoplus_{a \in \overline{Q}_1, e(a)=i} e_{s(a)} \Lambda \xrightarrow{\pi} e_i I_i \longrightarrow 0.$$

As a direct sum of (1) and $0 \to (1 - e_i)\Lambda \xrightarrow{id} (1 - e_i)\Lambda \to 0$, we have an exact sequence

(3)
$$0 \longrightarrow I_i \stackrel{\begin{pmatrix} \iota & 0\\ 0 & \mathrm{id} \end{pmatrix}}{\longrightarrow} \Lambda \stackrel{(p_0 & 0)}{\longrightarrow} S_i \longrightarrow 0.$$

Moreover, we have Ker $p_2 = S_{\sigma(i)}$, where $\sigma : Q_0 \to Q_0$ is a Nakayama permutation of Λ (i.e. $D(\Lambda e_{\sigma(i)}) \cong e_i \Lambda$) [BBK, Proposition 4.2]. Therefore, we have the following exact sequences

(4)
$$0 \longrightarrow S_{\sigma(i)} \longrightarrow e_i \Lambda \xrightarrow{p_2} \bigoplus_{a \in \overline{Q}_1, e(a) = i} e_{s(a)} \Lambda.$$

Then we give the following lemma.

Lemma 2.3. I_i is a basic support τ -tilting ideal of Λ for any $i \in Q_0$.

Proof. It is clear that I_i is basic. We will show that I_i is a right support τ -tilting Λ -module; the proof for a left Λ -module is similar. If $e_i\Lambda$ is a simple module, then Λ is a simple algebra since Λ is a preprojective algebra. Thus, in this case, we have $I_i = 0$ and it is a support τ -tilting module.

Assume that $e_i\Lambda$ is not a simple module and hence $e_iI_i \neq 0$. Since Λ is selfinjective, $e_j\Lambda$ has a simple socle for any $j \in Q_0$ and these simple modules are mutually non-isomorphic. Therefore, e_iI_i is not isomorphic to $e_j\Lambda$ for any $j \in Q_0$. Hence, we obtain $|I_i| = |\Lambda|$. Thus, in this case, it is enough to show that $\operatorname{Hom}_{\Lambda}(I_i, \tau(I_i)) = 0$.

Applying the functor $\nu := D \operatorname{Hom}_{\Lambda}(-, \Lambda)$ to (2), we have the following exact sequence

(5)
$$0 \longrightarrow \tau(e_i I_i) \longrightarrow \nu(e_i \Lambda) \xrightarrow{\nu p_2} \bigoplus_{a \in \overline{Q}_1, e(a) = i} \nu(e_{s(a)} \Lambda).$$

On the other hand, applying the functor ν to (4), we have the following exact sequence

(6)
$$0 \longrightarrow \nu(S_{\sigma(i)}) \longrightarrow \nu(e_i\Lambda) \xrightarrow{\nu p_2} \bigoplus_{a \in \overline{Q}_1, e(a) = i} \nu(e_{s(a)}\Lambda).$$

Since Λ is selfinjective, we have $\nu(S_{\sigma(i)}) \cong \nu(\operatorname{Soc}(e_i\Lambda)) \cong \operatorname{Top}(e_i\Lambda) \cong S_i$. Therefore, comparing (5) with (6), we get $\tau(I_i) \cong \tau(e_iI_i) \cong S_i$. Since $S_i \notin \operatorname{add}(\operatorname{Top}(I_i))$, we have $\operatorname{Hom}_{\Lambda}(I_i, \tau(I_i)) = 0$.

The next two statements are analogous to results of [BIRS, Lemma II.1.1, Proposition II.1.5], where tilting modules over preprojective algebras of non-Dynkin type are treated. Though the proofs given here are essentially the same, we include them for convenience.

Lemma 2.4. Let T be a support τ -tilting Λ -module. For a simple Λ^{op} -module S, at least one of the statements $T \otimes_{\Lambda} S = 0$ and $\text{Tor}_{1}^{\Lambda}(T, S) = 0$ holds.

Proof. By [AIR, Proposition 2.5], we can take a minimal projective presentation $P_1 \rightarrow P_0 \rightarrow T \rightarrow 0$ such that P_0 and P_1 do not have a common summand. Thus, at least one of the statement $P_0 \otimes_{\Lambda} S = 0$ and $P_1 \otimes_{\Lambda} S = 0$ holds. Since we have $\operatorname{Tor}_1^{\Lambda}(T, S) \cong P_1 \otimes_{\Lambda} S$ and $T \otimes_{\Lambda} S \cong P_0 \otimes_{\Lambda} S$, the assertion follows.

Corollary 2.5. Let T be a support τ -tilting Λ -module. If $TI_i \neq T$, then we have $T \otimes_{\Lambda} I_i \cong TI_i$.

Proof. By the assumption $TI_i \neq T$ and Lemma 1.2, we have $T \otimes_{\Lambda} S_i \neq 0$. Then, by Lemma 2.4, we obtain $\operatorname{Tor}_1^{\Lambda}(T, S_i) = 0$. By applying the functor $T \otimes_{\Lambda} -$ to the exact sequence (3), we get the following exact sequence

$$0 = \operatorname{Tor}_{1}^{\Lambda}(T, S_{i}) \to T \otimes_{\Lambda} I_{i} \to T \otimes_{\Lambda} \Lambda = T.$$

Hence we have $T \otimes_{\Lambda} I_i \cong \operatorname{Im}(T \otimes_{\Lambda} I_i \to T) = TI_i$.

Next we give the following easy observation.

Lemma 2.6. Let I be a two-sided ideal of Λ . For a primitive idempotent e_i with $i \in Q_0$, e_iI is either indecomposable or zero.

Proof. Since Λ is selfinjective, $e_i\Lambda$ has a simple socle. Thus the submodule e_iI of $e_i\Lambda$ has a simple socle or zero.

The next two lemmas are crucial.

Lemma 2.7. Let I be a two-sided ideal of Λ . Then there exists a surjective map

$$\Lambda \to \operatorname{End}_{\Lambda}(I), \quad \lambda \mapsto (I \ni x \mapsto \lambda x \in I).$$

In particular, for an idempotent e_i , $i \in Q_0$, we have a surjective map

$$e_i \Lambda(1-e_i) \to \operatorname{Hom}_{\Lambda}((1-e_i)I, e_iI).$$

Proof. We have to show that, for any $f \in \operatorname{End}_{\Lambda}(I)$, there exists $\lambda \in \Lambda$ such that $f = (\lambda \cdot) : I \to I$. Since Λ is selfinjective, there exists a Λ -module map $h : \Lambda \to \Lambda$ making the diagram



commutative, where ι is the canonical inclusion and $h : \Lambda \to \Lambda$ is the Λ -module map. Put $h(1) = \lambda \in \Lambda$. Then h is given by left multiplication with λ . Thus, we obtain $f(x) = h(x) = \lambda x$ for any $x \in I$ and we have proved our claim. The second statement easily follows from the first one.

Lemma 2.8. Let T be a support τ -tilting ideal of Λ . If $I_iT \neq T$, then we have $e_iT \notin Fac((1-e_i)T)$. In particular, T has a left mutation $\mu_{e_iT}^-(T)$.

Proof. By Lemma 2.6 and the assumption, $e_i T \neq 0$ is indecomposable. We assume that $e_i T \in \mathsf{Fac}((1-e_i)T)$. Then, there exist an integer d > 0 and a surjective map

$$\{(1-e_i)T\}^d \to e_iT.$$

By Lemma 2.7, there exist $\lambda_m \in \Lambda$ $(1 \leq m \leq d)$ such that

$$\sum_{m=1}^{d} e_i \lambda_m (1-e_i) \cdot (1-e_i)T = e_i T.$$

On the other hand, since $I_i = \Lambda(1 - e_i)\Lambda$, we get

$$e_i\Lambda(1-e_i)\cdot(1-e_i)T = e_iI_iT.$$

Thus we have

$$e_i T = \sum_{m=1}^d e_i \lambda_m (1 - e_i) T \subset e_i I_i T,$$

which is a contradiction. The second statement given in this lemma follows from Definition-Theorem 1.5 (ii). $\hfill \Box$

We also need to prove the following statement.

Lemma 2.9. Let T be a support τ -tilting ideal of Λ . For the map p_2 in the sequence (2), the map

$$e_i \Lambda \otimes_{\Lambda} T \xrightarrow{p_2 \otimes_{\Lambda} T} \bigoplus_{a \in \overline{Q}_1, e(a) = i} e_{s(a)} \Lambda \otimes_{\Lambda} T$$

is a left $(add((1 - e_i)T))$ -approximation.

Proof. We will show that the map

$$(7) \operatorname{Hom}_{\Lambda}(\bigoplus_{a \in \overline{Q}_{1}, e(a)=i} e_{s(a)} \Lambda \otimes_{\Lambda} T, (1-e_{i})T) \xrightarrow{(p_{2} \otimes_{\Lambda} T, (1-e_{i})T)} \operatorname{Hom}_{\Lambda}(e_{i} \Lambda \otimes_{\Lambda} T, (1-e_{i})T)$$

is surjective.

Let $E := \operatorname{End}_{\Lambda}(T)$. Then, we have

$$\operatorname{Hom}_{\Lambda}(e_{i}\Lambda \otimes_{\Lambda} T, (1-e_{i})T) \cong \operatorname{Hom}_{\Lambda}(e_{i}\Lambda, \operatorname{Hom}_{\Lambda}(T, (1-e_{i})T)) \\ \cong \operatorname{Hom}_{\Lambda}(e_{i}\Lambda, (1-e_{i})E),$$

and similarly

$$\operatorname{Hom}_{\Lambda}(\bigoplus_{a\in\overline{Q}_{1},e(a)=i}e_{s(a)}\Lambda\otimes_{\Lambda}T,(1-e_{i})T)\cong\operatorname{Hom}_{\Lambda}(\bigoplus_{a\in\overline{Q}_{1},e(a)=i}e_{s(a)}\Lambda,(1-e_{i})E).$$

Then, from the functoriality, we write the map (7) as follows.

$$\operatorname{Hom}_{\Lambda}(\bigoplus_{a\in\overline{Q}_{1},e(a)=i}e_{s(a)}\Lambda,(1-e_{i})E) \xrightarrow{(p_{2},(1-e_{i})E)} \operatorname{Hom}_{\Lambda}(e_{i}\Lambda,(1-e_{i})E).$$

On the other hand, by Lemma 2.7, there exists a surjective Λ -module map

$$g:\Lambda\longrightarrow E.$$

Then, we have the following commutative diagram

It is clear that $\operatorname{Hom}_{\Lambda}(\bigoplus_{a \in \overline{Q}_1, e(a)=i} e_{s(a)}\Lambda, g)$ and $\operatorname{Hom}_{\Lambda}(e_i\Lambda, g)$ are surjective since $e_{s(a)}\Lambda$ and $e_i\Lambda$ are projective. Moreover, by the sequence (4), $\operatorname{Hom}_{\Lambda}(p_2, (1-e_i)\Lambda)$ is also surjective. Consequently, $\operatorname{Hom}_{\Lambda}(p_2, (1-e_i)E)$ is also surjective. Then the statement follows from the additivity of the functors. \Box

Now we apply the above results to the following key proposition.

Proposition 2.10. Let $T \in \langle I_1, \ldots, I_n \rangle$ and assume that T is a basic support τ -tilting ideal of Λ . Then I_iT is a basic support τ -tilting Λ -module.

Proof. There is nothing to show if $I_iT = T$. Assume that $I_iT \neq T$. Then, by Lemma 2.8, there exists a left mutation $\mu_{e,T}^-(T)$.

Now let e be an idempotent of Λ such that T is a τ -tilting $(\Lambda/\langle e \rangle)$ -module. By applying the functor $-\otimes_{\Lambda} T$ to (2), we get the following exact sequence of $(\Lambda/\langle e \rangle)$ -module

$$e_i\Lambda\otimes_{\Lambda}T \xrightarrow{p_2\otimes_{\Lambda}T} \bigoplus_{a\in\overline{Q}_1,e(a)=i} e_{s(a)}\Lambda\otimes_{\Lambda}T \xrightarrow{\pi\otimes_{\Lambda}T} e_iI_i\otimes_{\Lambda}T \longrightarrow 0.$$

By applying Corollary 2.5 to the support τ -tilting left Λ -module T, we have $e_i I_i \otimes_{\Lambda} T \cong e_i I_i T$. Moreover, Lemma 2.6 implies that $e_i I_i T$ is indecomposable. On the other hand, from Lemma 2.9, $p_2 \otimes_{\Lambda} T$ is a left $(\mathsf{add}((1 - e_i)T))$ -approximation.

If $e_i I_i T = 0$, then $p_2 \otimes_{\Lambda} T$ is clearly left minimal. Assume $e_i I_i T \neq 0$. Since Λ is selfinjective, $e_j T$ has a simple socle for any $j \in Q_0$ and these simple modules are mutually non-isomorphic. Hence $e_{s(a)} \Lambda \otimes_{\Lambda} T$ and $e_i I_i T$ are not isomorphic for any $a \in \overline{Q}_1$ such that e(a) = i. Thus, $\pi \otimes_{\Lambda} T$ is a radical map and hence $p_2 \otimes_{\Lambda} T$ is left minimal. Then, applying Definition-Theorem 1.5 (iii), we conclude that $\mu_{e_iT}^-(T) \cong e_i I_i T \oplus (1-e_i)T = I_i T$ is a basic support τ -tilting $(\Lambda/\langle e \rangle)$ -module.

Now we are ready to prove Theorem 2.2.

Proof of Theorem 2.2. We use induction on the number of products of the ideal I_i , $i \in Q_0$. By Lemma 2.3, I_i is a basic support τ -tilting ideal of Λ for any $i \in Q_0$. Assume that $I_{i_1} \cdots I_{i_k}$ is a basic support τ -tilting ideal for $i_1, \ldots, i_k \in Q_0$ and k > 1. By Proposition 2.10, $I_{i_0}(I_{i_1} \cdots I_{i_k})$ is a basic support τ -tilting Λ -module for any vertex $i_0 \in Q_0$. Similarly we can show that it is a basic support τ -tilting left Λ -module.

At the end of this subsection, we give the following lemma for later use.

Lemma 2.11. Let $T \in \langle I_1, \ldots, I_n \rangle$. If $I_i T \neq T$, then there is a left mutation of T:

$$\mu_{e_iT}^{-}(T) \cong I_iT.$$

Proof. This follows from Lemma 2.8 and Proposition 2.10.

Now we show examples of support τ -tilting modules and their mutations.

Example 2.12. (a) Let Λ be the preprojective algebra of type A_2 . In this case, $\mathcal{H}(s\tau\text{-tilt}\Lambda)$ is given as follows.



Here we represent modules by their radical filtrations and we write a direct sum $X \oplus Y$ by X Y. Note that I_i denotes a left multiplication.

(b) Let Λ be the preprojective algebra of type A_3 . In this case, $\mathcal{H}(s\tau\text{-tilt}\Lambda)$ is given as follows



In these two examples, $\mathcal{H}(s\tau\text{-tilt}\Lambda)$ consists of a finite connected component. We will show that this is the case for preprojective algebras of Dynkin type in the sequel. Thus, all support τ -tilting modules can be obtained by mutations from Λ .

2.2. Bijection between support τ -tilting modules and the Weyl group. In the previous subsection, we have shown that elements of $\langle I_1, \ldots, I_n \rangle$ are basic support τ -tilting ideals. In this subsection, we will show that any basic support τ -tilting Λ -module is given as an element of $\langle I_1, \ldots, I_n \rangle$. Moreover, we show that there exists a bijection between the Weyl group and basic support τ -tilting Λ -modules, which implies that there are only finitely many basic support τ -tilting Λ -modules.

First we recall the following definition.

Definition 2.13. Let Q be a finite connected acyclic quiver with vertices $Q_0 = \{1, \ldots, n\}$. The *Coxeter group* W_Q associated to Q is defined by the generators s_1, \ldots, s_n and relations

- $s_i^2 = 1$,
- $s_i s_j = s_j s_i$ if there is no arrow between *i* and *j* in *Q*,
- $s_i s_j s_i = s_j s_i s_j$ if there is precisely one arrow between *i* and *j* in *Q*.

Each element $w \in W_Q$ can be written in the form $w = s_{i_1} \cdots s_{i_k}$. If k is minimal among all such expressions for w, then k is called the *length* of w and we denote by l(w) = k. In this case, we call $s_{i_1} \cdots s_{i_k}$ a *reduced expression* of w. In particular W_Q is the Weyl group if Q is Dynkin. Then, we give the following important result.

Theorem 2.14. Let Q be a finite connected acyclic quiver with vertices $Q_0 = \{1, \ldots, n\}$. There exists a bijection $W_Q \to \langle I_1, \ldots, I_n \rangle$. It is given by $w \mapsto I_w = I_{i_1}I_{i_2}\cdots I_{i_k}$ for any reduced expression $w = s_{i_1}\cdots s_{i_k}$.

The statement is given in [BIRS, III.1.9] for non-Dynkin quivers. The same result holds for Dynkin quivers. For the convenience of the reader, we give a complete proof of Theorem 2.14 for Dynkin quivers.

For a proof, we provide the following set-up.

Notation 2.15. Let \widetilde{Q} be an extended Dynkin quiver obtained from Q by adding a vertex 0 (i.e. $\widetilde{Q}_0 = \{0\} \cup Q_0$) and the associated arrows. We denote by $\widetilde{\Lambda}$ the completion of the associated preprojective algebra and $\widetilde{I}_i := \widetilde{\Lambda}(1 - e_i)\widetilde{\Lambda}$ for $i \in \widetilde{Q}$. For each $w \in W_{\widetilde{Q}}$, let $\widetilde{I}_w := \widetilde{I}_{i_1} \cdots \widetilde{I}_{i_k}$, where $w = s_{i_1} \cdots s_{i_k}$ is a reduced expression (this is well-defined [BIRS, Theorem III.1.9]). Note that we have

$$\Lambda = \widetilde{\Lambda} / \langle e_0 \rangle, \quad I_i = \widetilde{I}_i / \langle e_0 \rangle.$$

Proof of Theorem 2.14. One can check that the map

$$W_Q \to \langle I_1, \dots, I_n \rangle, \ w = s_{i_1} \cdots s_{i_k} \mapsto I_w := I_{i_1} I_{i_2} \cdots I_{i_k}$$

is well-defined and surjective by [BIRS, Theorem III.1.9], where the assumption that Q is non-Dynkin is not used. We only have to show the injectivity.

Since $i_1, \dots, i_k \in Q_0$, we have $s_{i_j} \neq s_0$ for any $1 \leq j \leq k$. Thus we have

$$I_w = I_{i_1} \cdots I_{i_k} = (\widetilde{I}_{i_1} / \langle e_0 \rangle) \cdots (\widetilde{I}_{i_k} / \langle e_0 \rangle) = (\widetilde{I}_{i_1} \cdots \widetilde{I}_{i_k}) / \langle e_0 \rangle = \widetilde{I}_w / \langle e_0 \rangle.$$

Therefore, if $I_w = I_{w'}$ for $w, w' \in W_Q$, then we have $\widetilde{I}_w/\langle e_0 \rangle = \widetilde{I}_{w'}/\langle e_0 \rangle$. Since s_0 does not appear in reduced expressions of w and w', we have $e_0 \in \widetilde{I}_w$ and $e_0 \in \widetilde{I}_{w'}$. Hence, we obtain $\widetilde{I}_w = \widetilde{I}_{w'}$ and conclude w = w' by [BIRS, Theorem III.1.9].

Thus, for any $w \in W_Q$, we have a support τ -tilting module I_w . Now we give an explicit description of mutations using this correspondence.

Let (I_w, P_w) be a basic support τ -tilting pair, where P_w is a basic projective Λ -module which is determined from I_w (subsection 1.2). Because $e_i\Lambda$ has simple socle $S_{\sigma(i)}$, where $\sigma: Q_0 \to Q_0$ is a Nakayama permutation of Λ , if $e_iI_w = 0$, then $e_{\sigma(i)}\Lambda \in \operatorname{add} P_w$ and hence P_w is given by $\bigoplus_{i \in Q_0, e_iI_w = 0} e_{\sigma(i)}\Lambda$. For any $i \in Q_0$, we define a mutation

$$\mu_i(I_w, P_w) := \begin{cases} \mu_{(e_i I_w, 0)}(I_w, P_w) & \text{if } e_i I_w \neq 0\\ \mu_{(0, e_{\sigma(i)} \Lambda)}(I_w, P_w) & \text{if } e_i I_w = 0. \end{cases}$$

Then we have the following result.

Theorem 2.16. For any $w \in W_Q$ and $i \in Q_0$, a mutation of basic support τ -tilting Λ -module I_w is given as follows :

$$\mu_i(I_w, P_w) \cong (I_{s_iw}, P_{s_iw})$$

In particular, W_Q acts transitively and freely on $s\tau$ -tilt Λ by $s_i(I_w, P_w) := \mu_i(I_w, P_w)$.

Proof. By Theorem 2.2 and 2.14, I_w and I_{s_iw} are basic support τ -tilting modules, which are not isomorphic.

Let $w = s_{i_1} \cdots s_{i_k}$ be a reduced expression. If $l(s_i w) > l(w)$, then $s_i w = s_i s_{i_1} \cdots s_{i_k}$ is a reduced expression. Hence, we get $(1 - e_i)I_{s_i w} = (1 - e_i)I_i I_w = (1 - e_i)I_w$. Therefore we have

$$(I_{s_iw}, P_{s_iw}) \cong \begin{cases} (e_i I_i I_w \oplus (1 - e_i) I_w, P_w) & \text{if } e_i I_i I_w \neq 0\\ ((1 - e_i) I_w, P_w \oplus e_{\sigma(i)} \Lambda) & \text{if } e_i I_i I_w = 0. \end{cases}$$

Thus (I_w, P_w) and (I_{s_iw}, P_{s_iw}) have a common almost complete support τ -tilting pair $((1 - e_i)I_w, P_w)$ as a direct summand.

On the other hand, if $l(s_iw) < l(w)$, then $u := s_iw$ satisfies $l(u) < l(s_iu)$. Then, similarly, we can show that $(I_u, P_u) = (I_{s_iw}, P_{s_iw})$ and $(I_{s_iu}, P_{s_iu}) = (I_w, P_w)$ have a common almost complete support τ -tilting pair as a direct summand.

Therefore, by Definition-Theorem 1.5 (i), the first statement follows. The second statement given in this theorem follows from Theorem 2.14. \Box

Using the above result, we give the following crucial lemma.

Lemma 2.17. The support τ -tilting quiver $\mathcal{H}(s\tau\text{-tilt}\Lambda)$ has a finite connected component.

Proof. Let C be the connected component of $\mathcal{H}(s\tau\text{-tilt}\Lambda)$ containing Λ . By Theorem 2.2, I_w is a basic support τ -tilting Λ -module for any $w \in W_Q$. We will show that $\{I_w \mid w \in W_Q\}$ gives the set of vertices of C.

First, we show that every I_w belongs to C by induction on the length of $w \in W_Q$. Assume that $\{I_w \mid w \in W_Q \text{ with } l(w) \leq k\}$ belong to C. Take $w \in W_Q$ with l(w) = k + 1and decompose $w = s_i w'$, where l(w') = k. Then, we have $I_w = I_i I_{w'} \neq I_{w'}$ by Theorem 2.14 and we get $\mu_{e_i I_{w'}}^-(I_{w'}) \cong I_i I_{w'}$ by Lemma 2.11. Thus, I_w belongs to C and our claim follows inductively.

Next, we show that any vertex in C has a form I_w for some $w \in W_Q$. This follows the fact that the neighbours of I_w are given as I_{s_iw} for $i \in Q$ by Theorem 2.16. Therefore we obtain the assertion.

Now we recall the following useful result.

Proposition 2.18. [AIR, Corollary 2.38] If $\mathcal{H}(s\tau\text{-tilt}\Lambda)$ has a finite connected component C, then $C = \mathcal{H}(s\tau\text{-tilt}\Lambda)$.

As a conclusion, we can obtain the following statement.

Theorem 2.19. Any basic support τ -tilting Λ -module is isomorphic to an element of $\langle I_1, \ldots, I_n \rangle$. In particular, it is isomorphic to a support τ -tilting ideal of Λ .

Proof. By Lemma 2.17, $\mathcal{H}(s\tau\text{-tilt}\Lambda)$ has a finite connected component, and Proposition 2.18 implies that it coincides with $\mathcal{H}(s\tau\text{-tilt}\Lambda)$. Thus, any support $\tau\text{-tilting }\Lambda\text{-module}$ is given by I_w for some $w \in W_Q$. In particular, it is a support $\tau\text{-tilting}$ ideal from Theorem 2.2.

Furthermore, we give the following lemma.

Lemma 2.20. If right ideals T and U are isomorphic as Λ -modules, then T = U.

Proof. Assume that $f: T \to U$ is an isomorphism of Λ -modules. Since Λ is selfinjective, there exists a Λ -module map $h: \Lambda \to \Lambda$ making the diagram



commutative, where ι is the canonical inclusion. Put $h(1) = \lambda \in \Lambda$. By the commutative diagram, we have $\iota f(T) = h\iota(T)$ and hence we get $U = h(T) = \lambda T \subset T$. By the similar argument for f^{-1} , we can show that $T \subset U$. Thus we conclude that T = U.

Now we are ready to prove the main result of this paper.

Theorem 2.21. There exist bijections between the isomorphism classes of basic support τ -tilting Λ -modules, basic support τ -tilting Λ^{op} -modules and the elements of W_Q .

Proof. By Theorem 2.19, it is enough to show a bijection between $s\tau$ -tilt Λ and W_Q .

By Theorem 2.2 and 2.14, we have a map $W_Q \ni w \mapsto I_w \in s\tau$ -tilt Λ . This map is surjective since any support τ -tilting Λ -module is isomorphic to I_w for some $w \in W_Q$ by Theorem 2.19. Moreover, it is injective by Theorem 2.14 and Lemma 2.20. Thus we get the conclusion.

Remark 2.22. It is known that the order of W_Q is finite and the explicit number is determined by the type of the underlying graph of Q as follows [H, Section 2.11].

\overline{Q}	A_n	D_n	E_6	E_7	E_8
	(n+1)!	$2^{n-1}n!$	51840	2903040	696729600

At the end of this subsection, we give an application related to annihilators of $s\tau$ -tilt Λ . Note that a tilting module is a *faithful* τ -tilting modules (i.e. its (right) annihilator vanishes) [AIR, Proposition 2.2]. We can completely determine annihilators of support τ -tilting Λ -modules. Here, we denote by ann $X := \{\lambda \in \Lambda \mid X\lambda = 0\}$ the annihilator of $X \in \mathsf{mod}\Lambda$.

Corollary 2.23. We have ann $I_w = I_{w^{-1}w_0}$, where w_0 is the longest element in W_Q . Thus, we have a bijection

$$s\tau$$
-tilt $\Lambda \to s\tau$ -tilt $\Lambda, T \mapsto annT$.

For a proof, we recall the following useful result.

Proposition 2.24. [ORT, Proposition 6.4] Let w_0 be the longest element in W_Q . Then we have the following isomorphisms

- (a) $DI_w \cong \Lambda/I_{w^{-1}w_0}$ in mod Λ^{op} .
- (b) $DI_w \cong \Lambda/I_{w_0w^{-1}}$ in mod Λ .

Then we give a proof of Corollary 2.23.

Proof of Corollary 2.23. It is clear that the right annihilator of I_w coincides with the left annihilator of DI_w . On the other hand, we have isomorphism $DI_w \cong \Lambda/I_{w^{-1}w_0}$ of Λ^{op} -modules by Proposition 2.24. Thus, we get ann $I_w = I_{w^{-1}w_0}$. The second statement given in this corollary follows from Theorem 2.14.

2.3. Partial orders of support τ -tilting modules and the Weyl group. In this subsection, we study a relationship between a partial order of support τ -tilting modules and that of W_Q . In particular, we show that the bijection $W_Q \to s\tau$ -tilt Λ given in Theorem 2.21 induces an (anti)isomorphism as partially ordered sets.

Definition 2.25. Let $u, w \in W_Q$. We write $u \leq_L w$ if there exist s_{i_1}, \ldots, s_{i_k} such that $w = s_{i_k} \ldots s_{i_1} u$ and $l(s_{i_j} \ldots s_{i_1} u) = l(u) + j$ for $0 \leq j \leq k$. Clearly \leq_L gives a partial order on W_Q , and we call \leq_L the (*left*) weak order. We denote by $\mathcal{H}(W_Q, \leq_L)$ the Hasse quiver induced by weak order on W_Q .

The following assertion is immediate from the definition of the weak order.

Lemma 2.26. An arrow in $\mathcal{H}(W_Q, \leq_L)$ is given by

 $w \to s_i w \quad (if \ l(w) > l(s_i w)), \\ s_i w \to w \quad (if \ l(w) < l(s_i w))$

for any $w \in W_Q$ and $i \in Q_0$.

Example 2.27. (a) Let Q be a quiver of type A_2 . Then $\mathcal{H}(W_Q, \leq_L)$ is given as follows.



(b) Let Q be a quiver of type A_3 . Then $\mathcal{H}(W_Q, \leq_L)$ is given as follows.



Next we give the following observation.

Lemma 2.28. Let $w \in W_Q$ and $i \in Q_0$.

- (i) If $l(w) < l(s_iw)$, then $I_i I_w = I_{s_iw} \subsetneq I_w$ and we have a left mutation $\mu_i^-(I_w, P_w)$.
- (ii) If $l(w) > l(s_i w)$, then $I_i I_w = I_w \subsetneq I_{s_i w}$ and we have a right mutation $\mu_i^+(I_w, P_w)$.

Proof. Let $w = s_{i_1} \cdots s_{i_k}$ be a reduced expression. If $l(s_i w) > l(w)$, then $s_i w = s_i s_{i_1} \cdots s_{i_k}$ is a reduced expression. Thus, we have $I_i I_w = I_{s_i w} \subsetneq I_w$ by Theorem 2.14, and we get a left mutation $\mu_i^-(I_w, P_w)$ by Lemma 2.11.

On the other hand, if $l(s_iw) < l(w)$, then $u := s_iw$ satisfies $l(u) < l(s_iu)$. Then, similarly, we get a left mutation $\mu_i^-(I_u, P_u) = (I_{s_iu}, P_{s_iu}) = (I_w, P_w)$, or equivalently $\mu_i^+(I_w, P_w) = (I_{s_iw}, P_{s_iw})$.

Summarizing previous results, we give the following equivalent conditions that a left (or right) mutation occurs.

Corollary 2.29. Let $w \in W_Q$ and $i \in Q_0$.

- (a) The following conditions are equivalent:
 - (i) $l(w) < l(s_i w)$.
 - (ii) $I_i I_w \neq I_w$.
 - (iii) We have a left mutation $\mu_i^-(I_w, P_w)$.
- (b) The following conditions are equivalent :
 - (i) $l(w) > l(s_i w)$.
 - (ii) $I_i I_w = I_w$.
 - (iii) We have a right mutation $\mu_i^+(I_w, P_w)$.

Proof. It is enough to show (a).

By Lemma 2.11 and 2.28, (i) implies (ii) and (ii) implies (iii). Assume that we have a left mutation $\mu_i^-(I_w, P_w)$. If $l(w) > l(s_i w)$, then we get a right mutation $\mu_i^+(I_w, P_w)$ by Lemma 2.28, which contradicts Definition-Theorem 1.5. Thus (iii) implies (i).

Finally we give the following results.

Theorem 2.30. The bijection in $W_Q \to s\tau$ -tilt Λ in Theorem 2.21 gives an isomorphism of partially ordered sets (W_Q, \leq_L) and $(s\tau$ -tilt $\Lambda, \leq)^{\text{op}}$.

Proof. It is enough to show that $\mathcal{H}(W_Q, \leq_L)$ coincides with $\mathcal{H}(s\tau\text{-tilt}\Lambda)^{\text{op}}$ since the Hasse quivers determine the partial orders.

By Theorem 1.7 and Corollary 2.29, arrows starting from I_w are given by $I_w \to I_i I_w$ in $\mathcal{H}(s\tau\text{-tilt}\Lambda)$, where $i \in Q_0$ satisfies $l(w) < l(s_i w)$.

On the other hand, Lemma 2.26 implies that arrows to w are given by $s_i w \to w$ in $\mathcal{H}(W_Q, \leq_L)$, where s_i satisfies $l(w) < l(s_i w)$. Therefore $\mathcal{H}(s\tau\text{-tilt}\Lambda)^{\text{op}}$ coincides with $\mathcal{H}(W_Q, \leq_L)$ and the statement follows.

We remark that the *Bruhat order* on W_Q coincides with the reverse inclusion relation on $\langle I_1, \dots, I_n \rangle$ [ORT, Lemma 6.5].

3. g-matrices and cones

In this section, we study *g*-vectors [DK], which is also called *index* [P] (see also [AR1]), and *g*-matrices of support τ -tilting modules. We give a complete description of *g*-matrices in terms of reflection transformations. Moreover, we study cones defined by *g*-matrices, which provide a geometric realization of support τ -tilting modules, and we show that they coincide with chambers of the associated root system.

Throughout this section, unless otherwise specified, let Q be a Dynkin quiver with vertices $Q_0 = \{1, \ldots, n\}$, Λ the preprojective algebra of Q and $I_i = \Lambda(1 - e_i)\Lambda$ for $i \in Q_0$.

First, we define g-vectors in our setting. We refer to [AIR, section 5] for a background of this notion.

Definition 3.1. By Krull-Schmidt theorem, we identify the set of isomorphism classes of projective Λ -modules with submonoid $\mathbb{Z}_{>0}^n$ of the free abelian group \mathbb{Z}^n .

For a Λ -module X, take a minimal projective presentation

 $P_1(X) \longrightarrow P_0(X) \longrightarrow X \longrightarrow 0$

and let $g(X) = (g_1(X), \dots, g_n(X))^t := P_0(X) - P_1(X) \in \mathbb{Z}^n$, where t denotes the transpose matrix. Then, for any $w \in W_Q$ and $i \in Q_0$, we define a g-vector by

$$\mathbb{Z}^{n} \ni g^{i}(w) = \begin{cases} g(e_{i}I_{w}) & \text{if } e_{i}I_{w} \neq 0\\ \sigma^{(i)} \\ (0, \dots, 0, -1, 0, \dots, 0)^{t} & \text{if } e_{i}I_{w} = 0, \end{cases}$$

where $\sigma: Q_0 \to Q_0$ is a Nakayama permutation of Λ (i.e. $D(\Lambda e_{\sigma(i)}) \cong e_i \Lambda$)).

We define a *g*-matrix of support τ -tilting Λ -module I_w by

$$g(w) := (g^1(w), \cdots, g^n(w)) \in GL_n(\mathbb{Z})$$

and its cone by

$$C(w) := \{a_1g^1(w) + \dots + a_ng^n(w) \mid a_i \in \mathbb{R}_{>0}\}.$$

Next, we define g-vectors for extended Dynkin cases. We follow Notation 2.15. By Theorem 2.14, there exists a bijection

$$W_{\widetilde{Q}} \to \langle \widetilde{I}_0, \widetilde{I}_1, \dots, \widetilde{I}_n \rangle, w \mapsto \widetilde{I}_w = \widetilde{I}_{i_1} \widetilde{I}_{i_2} \cdots \widetilde{I}_{i_k},$$

where $w = s_{i_1} \cdots s_{i_k}$ is a reduced expression of w. Note that \widetilde{I}_w is a tilting $\widetilde{\Lambda}$ -modules of projective dimension at most one [IR] (hence $e_i \widetilde{I}_w \neq 0$ for any $i \in \widetilde{Q}_0$). For any $w \in W_{\widetilde{Q}}$ and $i \in \widetilde{Q}_0$, take a minimal projective presentation of $\widetilde{\Lambda}$ -module $e_i \widetilde{I}_w$

$$P_1(e_i \widetilde{I}_w) \longrightarrow P_0(e_i \widetilde{I}_w) \longrightarrow e_i \widetilde{I}_w \longrightarrow 0.$$

Then we define a g-vector by

$$\tilde{g}^i(w) = (g_0(e_i \widetilde{I}_w), \cdots, g_n(e_i \widetilde{I}_w))^t := P_0(e_i \widetilde{I}_w) - P_1(e_i \widetilde{I}_w) \in \mathbb{Z}^{n+1},$$

and we define a g-matrix by

$$\tilde{g}(w) := (\tilde{g}^0(w), \cdots, \tilde{g}^n(w)) \in GL_{n+1}(\mathbb{Z}).$$

We use these notations in this section. We also denote by $\overline{(-)} := - \otimes_{\widetilde{\Lambda}} \Lambda$ for simplicity. Then, we give the following results.

Proposition 3.2. Let $w \in W_Q$ and $i \in Q_0$. Take a minimal projective presentation of $\widetilde{\Lambda}$ -module $e_i \widetilde{I}_w$

(8)
$$0 \longrightarrow P_1 \xrightarrow{f_1} P_0 \xrightarrow{f_0} e_i \widetilde{I}_w \longrightarrow 0.$$

Then, applying the functor $-\otimes_{\widetilde{\Lambda}} \Lambda$ to (8), we have the following exact sequence

$$0 \longrightarrow \nu^{-1}(e_i \Lambda/e_i I_w) \longrightarrow \overline{P_1} \xrightarrow{\overline{f_1}} \overline{P_0} \xrightarrow{\overline{f_0}} e_i I_w \longrightarrow 0 ,$$

where $\nu^{-1} := \operatorname{Hom}_{\Lambda}(D\Lambda, -)$. Moreover, if $e_i I_w \neq 0$, then the sequence

$$\overline{P_1} \xrightarrow{\overline{f_1}} \overline{P_0} \xrightarrow{\overline{f_0}} e_i I_w \longrightarrow 0$$

is a minimal projective presentation of Λ -module $e_i I_w$.

Proof. Since $e_i \neq e_0$ and $\Lambda = \widetilde{\Lambda}/\langle e_0 \rangle$, we have $e_i \widetilde{I}_w \otimes_{\widetilde{\Lambda}} \Lambda \cong e_i \widetilde{I}_w/e_i \widetilde{I}_w \langle e_0 \rangle \cong e_i \widetilde{I}_w/e_i \langle e_0 \rangle \cong e_i I_w$. Then, by applying the functor $- \otimes_{\widetilde{\Lambda}} \Lambda$ to (8), we have the following exact sequence

$$0 \longrightarrow \operatorname{Tor}_{1}^{\widetilde{\Lambda}}(e_{i}\widetilde{I}_{w},\Lambda) \xrightarrow{g} \overline{P_{1}} \xrightarrow{\overline{f_{1}}} \overline{P_{0}} \xrightarrow{\overline{f_{0}}} e_{i}I_{w} \longrightarrow 0$$

On the other hand, we have the following exact sequence

$$0 \longrightarrow e_i \widetilde{I}_w \longrightarrow e_i \widetilde{\Lambda} \longrightarrow e_i \widetilde{\Lambda}/e_i \widetilde{I}_w \longrightarrow 0.$$

Then, we obtain Λ -module isomorphisms

$$\operatorname{Tor}_{1}^{\Lambda}(e_{i}\widetilde{I}_{w},\Lambda) \cong D\operatorname{Ext}_{\widetilde{\Lambda}}^{1}(e_{i}\widetilde{I}_{w},D\Lambda) \quad ([\operatorname{CE}])$$

$$\cong D\operatorname{Ext}_{\widetilde{\Lambda}}^{2}(e_{i}\widetilde{\Lambda}/e_{i}\widetilde{I}_{w},D\Lambda)$$

$$\cong \operatorname{Hom}_{\widetilde{\Lambda}}(D\Lambda,e_{i}\widetilde{\Lambda}/e_{i}\widetilde{I}_{w}) \quad (2\text{-CY property})$$

$$\cong \operatorname{Hom}_{\Lambda}(D\Lambda,e_{i}\Lambda/e_{i}I_{w}) \quad (e_{i}\widetilde{\Lambda}/e_{i}\widetilde{I}_{w}\cong e_{i}\Lambda/e_{i}I_{w})$$

$$\cong \nu^{-1}(e_{i}\Lambda/e_{i}I_{w}).$$

Now assume that $e_i I_w \neq 0$. Then, clearly $\nu^{-1}(e_i \Lambda/e_i I_w)$ is not projective and hence g is a radical map. Moreover, it is clear that $\overline{f_0}$ is a projective cover by the assumption of f_0 . Thus we obtain the conclusion.

Remark 3.3. It is known that, for every non-projective indecomposable module X over a preprojective algebra of Dynkin type, its third syzygy $\Omega^3 X$ is isomorphic to $\nu^{-1}(X)$ [AR2, BBK]. This property appears in the above proposition.

Using the above proposition, we give the following key result.

Proposition 3.4. For any $w \in W_Q$ and $i \in Q_0$, the g-vector $g^i(w)$ equals the image of $\tilde{g}^i(w)$ under the projection $\mathbb{Z}^{n+1} \to \mathbb{Z}^n$ which forgets the 0-th coordinate.

Proof. Take a minimal projective presentation of Λ -module $e_i I_w$

(9)
$$0 \longrightarrow P_1 \xrightarrow{f_1} P_0 \xrightarrow{f_0} e_i \widetilde{I}_w \longrightarrow 0.$$

We decompose $P_0 = P'_0 \oplus (e_0 \widetilde{\Lambda})^\ell$ and $P_1 = P'_1 \oplus (e_0 \widetilde{\Lambda})^m$, where $(e_0 \widetilde{\Lambda})^\ell$ (respectively, $(e_0 \widetilde{\Lambda})^m$) is a maximal direct summand of P_0 (respectively, P_1) which belongs to $\mathsf{add}(e_0 \widetilde{\Lambda})$. For $\tilde{g}^i(w) = (g_0(e_i \widetilde{I}_w), g_1(e_i \widetilde{I}_w), \cdots, g_n(e_i \widetilde{I}_w))^t$, it is clear that $(g_1(e_i \widetilde{I}_w), \ldots, g_n(e_i \widetilde{I}_w))^t$ is determined by P'_0 and P'_1 . We will show that it coincides with $g^i(w)$.

By Proposition 3.2, applying $-\otimes_{\widetilde{\Lambda}} \Lambda$ to (9), we have the following exact sequence

(10)
$$0 \longrightarrow \nu^{-1}(e_i \Lambda/e_i I_w) \longrightarrow \overline{P'_1} \xrightarrow{\overline{f_1}} \overline{P'_0} \xrightarrow{\overline{f_0}} e_i I_w \longrightarrow 0.$$

(i) Assume that $P'_0 \neq 0$. It implies that $\overline{P'_0} \neq 0$ and hence $e_i I_w \neq 0$. Therefore, by Proposition 3.2, the sequence (10) gives a minimal projective presentation of $e_i I_w$. Thus $g^i(w)$ coincides with $(g_1(e_i \tilde{I}_w), \ldots, g_n(e_i \tilde{I}_w))^t$.

(ii) Assume that $P'_0 = 0$. It implies that $\overline{P'_0} = 0$ and hence $e_i I_w = 0$. Thus, $g^i(w) = (0, \ldots, 0, -1, 0, \cdots, 0)^t$ by definition. On the other hand, by the sequence (10), we have

 $\overline{P'_1} \cong \nu^{-1}(e_i\Lambda) \cong e_{\sigma(i)}\Lambda$ and hence $P'_1 \cong e_{\sigma(i)}\widetilde{\Lambda}$. Thus $\tilde{g}^i(w) = (g_0(e_i\widetilde{I}_w), 0, \dots, 0, -1, 0, \dots, 0)^t$ and we obtain the conclusion.

Next we introduce the following notations [Bou, BB].

Definition 3.5. Let Q be a finite connected acyclic quiver with vertices $Q_0 = \{1, \ldots, n\}$. For $i, j \in Q_0$, let $m_{ij} := \sharp\{a \in Q_1 | s(a) = i, e(a) = j\} + \sharp\{a \in Q_1 | s(a) = j, e(a) = i\}$. Let $\mathbf{e}_1, \ldots, \mathbf{e}_n$ be the canonical basis of \mathbb{Z}^n .

The geometric representation $\sigma: W_Q \to GL_n(\mathbb{Z})$ of W_Q is defined by

$$\sigma(s_i)(\mathbf{e}_j) = \sigma_i(\mathbf{e}_j) := \mathbf{e}_j + (m_{ij} - 2\delta_{ij})\mathbf{e}_i,$$

where δ_{ij} denotes the Kronecker delta. On the other hand, the contragradient $\sigma^* : W_Q \to GL_n(\mathbb{Z})$ of the geometric representation is defined by

$$\sigma^*(s_i)(\mathbf{e}_j) = \sigma^*_i(\mathbf{e}_j) = \begin{cases} -\mathbf{e}_j + \sum_{a \in \overline{Q}_1, s(a)=i}^{\mathbf{e}_j} \mathbf{e}_{e(a)} & i=j \end{cases}$$

For an arbitrary expression $w = s_{i_1} \dots s_{i_k}$ of w, we define $\sigma(w) := \sigma_{i_k} \dots \sigma_{i_1}$ and $\sigma^*(w) = \sigma^*_{i_k} \dots \sigma^*_{i_1}$.

Then, we give the following key result.

CLASSIFYING τ -TILTING MODULES OVER PREPROJECTIVE ALGEBRAS OF DYNKIN TYPE 21

Proposition 3.6. For any $w \in W_Q$, we have $g(w) = \sigma^*(w)$.

Proof. By Proposition 3.4, the *g*-matrix $g(w) = (g^1(w), \dots, g^n(w))$ is given by 1-st to *n*-th rows and columns of $\tilde{g}(w) = (\tilde{g}^0(w), \tilde{g}^1(w), \dots, \tilde{g}^n(w))$.

On the other hand, by [IR, Theorem 6.6], we have $\tilde{g}(w) = \tilde{\sigma}^*(w)$, where $\tilde{\sigma}^*$ is the contragradient $W_{\tilde{Q}} \to GL_{n+1}(\mathbb{Z})$ of the geometric representation.

For $w = s_{i_1} \dots s_{i_k}$, the 0-th column of $\tilde{\sigma}_{i_j}^*$ is $(1, 0, \dots, 0)^t$ by the assumption of $s_{i_j} \neq s_0$ for any $1 \leq j \leq k$. Because 1-st to *n*-th rows and columns of $\tilde{\sigma}_{i_j}^*$ is equal to $\sigma_{i_j}^*$ by definition, we conclude that $g(w) = \sigma_{i_k}^* \dots \sigma_{i_1}^* = \sigma^*(w)$.

Finally we give the following settings.

Definition 3.7. An element of $\Phi := \{\sigma(w)(\mathbf{e}_i)\}_{i \in Q_0, w \in W_Q}$ is called a *root*. For $a, b \in \mathbb{R}^n$, we write a natural pairing $\langle a, b \rangle := a \cdot b^t$. To each root $x \in \Phi$, we define a hyperplane

$$H_x := \{ y \in \mathbb{R}^n \mid \langle y, x \rangle = 0 \}$$

A chamber of Φ is a connected component of $\mathbb{R}^n \setminus \bigcup_{x \in \Phi} H_x$. We denote by C_0 the chamber $\{a_1 \mathbf{e}_1 + \cdots + a_n \mathbf{e}_n \mid a_i \in \mathbb{R}_{>0}\}$. Note that chambers are given by $\coprod_{w \in W_Q} \sigma^*(w)(C_0)$ and there is a bijection between W_Q and chambers via $w \mapsto \sigma^*(w)(C_0)$.

Then we obtain the following conclusion.

Theorem 3.8. (a) The set of g-matrices of support τ -tilting Λ -modules coincides with the subgroup $\langle \sigma_1^*, \ldots, \sigma_n^* \rangle$ of $GL(\mathbb{Z}^n)$ generated by σ_i^* for all $i \in Q_0$. (b) For $w \in W_Q$, we have

$$C(w) = \sigma^*(w)(C_0),$$

where C(w) denotes the cone of I_w . In particular, cones of basic support τ -tilting Λ -modules coincide with chambers of the associated root system Φ .

Proof. From Proposition 3.6, (a) follows. Moreover, we have

$$C(w) = \{a_1g^1(w) + \dots + a_ng^n(w) \mid a_i \in \mathbb{R}_{>0}\} = \{a_1\sigma^*(w)(\mathbf{e}_1) + \dots + a_n\sigma^*(w)(\mathbf{e}_n) \mid a_i \in \mathbb{R}_{>0}\} = \sigma^*(w)(C_0).$$

Thus, (b) follows.

Example 3.9. (a) Let Λ be the preprojective algebra of type A_2 . In this case, the *g*-matrices of Example 2.12 (a) are given as follows, where $\sigma_1^* = \begin{pmatrix} -1 & 0 \\ 1 & 1 \end{pmatrix}$ and $\sigma_2^* = \begin{pmatrix} 1 & 1 \\ 0 & -1 \end{pmatrix}$.



Moreover, their chambers are given as follows.



(b) Let Λ be the preprojective algebra of type A_3 . In this case, the *g*-matrices of Example 2.12 (b) are given as follows.



4. Further connections

In this section, we extend our bijections by combining with other works. We also explain some relationships with other results.

We have the following bijections (We refer to [BY] for the definitions (g), (h), (i) and (j)).

Theorem 4.1. Let Q be a Dynkin quiver with vertices $Q_0 = \{1, \ldots, n\}$ and Λ the preprojective algebra of Q. There are bijections between the following objects.

- (a) The elements of the Weyl group W_Q .
- (b) The set $\langle I_1, \ldots, I_n \rangle$.
- (c) The set of isomorphism classes of basic support τ -tilting Λ -modules.
- (d) The set of isomorphism classes of basic support τ -tilting Λ^{op} -modules.
- (e) The set of torsion classes in $mod\Lambda$.
- (f) The set of torsion-free classes in $mod\Lambda$.
- (g) The set of isomorphism classes of basic two-term silting complexes in $\mathsf{K}^{\mathsf{b}}(\mathsf{proj}\Lambda)$.
- (h) The set of intermediate bounded co-t-structures in $K^{b}(\text{proj}\Lambda)$ with respect to the standard co-t-structure.
- (i) The set of intermediate bounded t-structures in D^b(modΛ) with length heart with respect to the standard t-structure.
- (j) The set of isomorphism classes of two-term simple-minded collections in $D^{b}(mod\Lambda)$.
- (k) The set of quotient closed subcategories in modKQ.
- (1) The set of subclosed subcategories in modKQ.

We have given bijections between (a), (b), (c) and (d). Bijections between (g), (h), (i) and (j) are the restriction of [KY] and it is given in [BY, Corollary 4.3] (it is stated for Jacobian algebras, but it holds for any finite dimensional algebra as they point out). Note that compatibilities of mutations and partial orders of these objects are discussed in [KY].

We will give bijections between (a), (e) and (f) by showing the following statement, which provides complete descriptions of torsion classes and torsion-free classes in $mod\Lambda$.

Proposition 4.2. Any torsion class in $\text{mod}\Lambda$ is given as $\text{Fac}I_w$ and any torsion-free class in $\text{mod}\Lambda$ is given as $\text{Sub}\Lambda/I_w$ for some $w \in W_Q$. Moreover, there exist bijections between the elements of W_Q , torsion classes and torsion-free classes.

Proof. Since there exists only finitely many support τ -tilting modules by Theorem 2.21, Theorem 1.10 implies that any torsion class (torsion-free class) is functorially finite. Thus, Theorem 1.9 gives a bijection between basic support τ -tilting Λ -modules and torsion classes in mod Λ , which is given by the map $I_w \mapsto \mathsf{Fac} I_w$.

Similarly, there exists a bijection between basic support τ -tilting Λ^{op} -modules and torsion classes in $\mathsf{mod}\Lambda^{\text{op}}$. By the duality $D : \mathsf{mod}\Lambda^{\text{op}} \to \mathsf{mod}\Lambda$, any torsion-free class of $\mathsf{mod}\Lambda$ is given as $D(\mathsf{Fac}I_w)$, where $\mathsf{Fac}I_w$ is a torsion class in $\mathsf{mod}\Lambda^{\text{op}}$. Then, by Proposition 2.24, we have $D(\mathsf{Fac}I_w) \cong \mathsf{Sub}(DI_w) \cong \mathsf{Sub}\Lambda/I_{w_0w^{-1}}$, where w_0 is the longest element in W_Q .

Remark 4.3. It is shown that objects $\operatorname{Fac} I_w$ and $\operatorname{Sub} \Lambda/I_w$ have several nice properties. For example, $\operatorname{Fac} I_w$ and $\operatorname{Sub} \Lambda/I_w$ are *Frobenius* and, moreover, *stable 2-CY* categories which have cluster-tilting objects [BIRS, GLS3].

It is known that weak order of W_Q is a *lattice* (i.e. any two elements $x, y \in W_Q$ have a greatest lower bound, called *meet* $x \wedge y$ and a least upper bound, called *join* $x \vee y$). From the viewpoint of Theorem 4.2, it is natural to consider what is a join and meet in torsion classes. We give an answer to the question as follows.

Proposition 4.4. Let \mathcal{T} and \mathcal{T}' be torsion classes of mod Λ . Then we have $\mathcal{T} \wedge \mathcal{T}' = \mathcal{T} \cap \mathcal{T}'$ and $\mathcal{T} \vee \mathcal{T}' = \text{Tors}\{\mathcal{T}, \mathcal{T}'\}$, where $\text{Tors}\{\mathcal{T}, \mathcal{T}'\}$ is the smallest torsion class in mod Λ containing \mathcal{T} and \mathcal{T}' .

Proof. It is clear that $\mathcal{T} \cap \mathcal{T}'$ and $\mathsf{Tors}\{\mathcal{T}, \mathcal{T}'\}$ are torsion classes. Since any torsion class in mod Λ is functorially finite by Theorem 1.10 and 2.21, the statement follows.

Next we will see bijections between (a), (k) and (l), which are given by [ORT]. We briefly recall their results and explain a relationship with our results.

Let X be a Λ -module. We denote by X_{KQ} the KQ-module by the restriction, that is, we forget the action of the arrows $a^* \in \overline{Q}$. Then we define $(-)_{KQ} : \operatorname{mod} \Lambda \to \operatorname{mod} KQ, X \mapsto (X)_{KQ} := \operatorname{add} X_{KQ}$. Take I_w for $w \in W_Q$. In [ORT], the authors show that $(I_w)_{KQ}$ (respectively, $(\Lambda/I_w)_{KQ}$) is a quotient closed subcategory (respectively, subclosed subcategory) of $\operatorname{mod} KQ$, and any quotient closed subcategory (respectively, subclosed subcategory) is given in this form.

Now, we can conclude that the functor $(-)_{KQ}$ gives a bijection between (e) and (k) (similarly (f) and (l)). Indeed, any torsion class is given by $\mathsf{Fac}I_w$ for some $w \in W_Q$ from Proposition 4.2 and we have $(\mathsf{Fac}I_w)_{KQ} = (I_w)_{KQ}$ by a result of [ORT].

Moreover, it is natural to consider when a quotient closed subcategory $(I_w)_{KQ}$ is a torsion class. The answer is given by [ORT, Proposition 10.4] and they show that elements of W_Q satisfying a certain condition called being *c-sortable* [Re] are in bijection with torsion classes of modKQ by this correspondence. We emphasize that there exist bijections between torsion classes of modKQ and several important objects such as clusters in the cluster algebra given by Q (we refer to [IT, Theorem 1.1] for more details). Thus, there exists an inclusion from the objects of [IT, Theorem 1.1] to the objects of Theorem 4.1.

Finally, we provide a bijection between (c) and (g), which is given by [AIR, Theorem 3.2], and explain a definition of g-vectors. Let $\mathsf{K}^{\mathsf{b}}(\mathsf{proj}\Lambda)$ be the homotopy category of bounded complexes of $\mathsf{proj}\Lambda$. For a Λ -module X, take a minimal projective presentation of X

$$P_1(X) \longrightarrow P_0(X) \longrightarrow X \longrightarrow 0.$$

We denote by $P_X := (P_1(X) \to P_0(X)) \in \mathsf{K}^{\mathsf{b}}(\mathsf{proj}\Lambda)$. Then, for a support τ -tilting pair (X, P) for Λ , the map $(X, P) \mapsto P_X \oplus P[1] \in \mathsf{K}^{\mathsf{b}}(\mathsf{proj}\Lambda)$ gives a two-term silting complex in $\mathsf{K}^{\mathsf{b}}(\mathsf{proj}\Lambda)$ and it provides a bijection between (c) and (g). Note that the *g*-vector of (X, P) is given as a class of the corresponding silting complex $P_X \oplus P[1]$ in the Grothendieck group $K_0(\mathsf{K}^{\mathsf{b}}(\mathsf{proj}\Lambda))$ [AIR, section 5], and Definition 3.1 is given by this correspondence. We also remark that, since Λ is selfnjective, it is shown that a ν -stable support τ -tilting Λ -module X (i.e $X \cong \nu(X)$ for $\nu := D \operatorname{Hom}_{\Lambda}(-, \Lambda)$) gives a two-term tilting complex by this correspondence [M]. From Theorem 2.21, there exist only finitely many basic two-term tilting complexes in $\mathsf{K}^{\mathsf{b}}(\mathsf{proj}\Lambda)$.

References

[AIR] T. Adachi, O. Iyama, I. Reiten, τ -tilting theory, arXiv: 1210.1036, to appear in Compos. Math.

- [AIRT] C. Amiot, O. Iyama, I. Reiten, G. Todorov, Preprojective algebras and c-sortable words, Proc. Lond. Math. Soc. (3) 104 (2012), no. 3, 513–539.
- [AR1] M. Auslander, I. Reiten, Modules determined by their composition factors, Ill. J. of Math 29 (1985), 280–301.
- [AR2] M. Auslander, I. Reiten, DTr-periodic modules and functors, Representation theory of algebras (Cocoyoc, 1994), 39–50, CMS Conf. Proc., 18, Amer. Math. Soc., Providence, RI, 1996.
- [ARS] M. Auslander, I. Reiten, S. O. Smalø, *Representation Theory of Artin Algebras*, Cambridge Studies in Advanced Mathematics, 36. Cambridge University Press, Cambridge, 1995.
- [BGL] D. Baer, W. Geigle, H. Lenzing, *The preprojective algebra of a tame hereditary Artin algebra*, Comm. Algebra 15 (1987), no. 1-2, 425–457.

- [BKT] P. Baumann, J. Kamnitzer, P. Tingley, *Affine Mirković-Vilonen polytopes*, arXiv:1110.3661, to appear in Publ. IHES.
- [BB] A. Björner, F. Brenti, *Combinatorics of Coxeter groups*, Graduate Texts in Mathematics, vol. 231, Springer, New York, 2005.
- [Boc] R. Bocklandt, Graded Calabi Yau algebras of dimension 3, J. Pure Appl. Algebra 212 (2008), no. 1, 14–32.
- [Bou] N. Bourbaki, *Lie groups and Lie algebras. Chapters* 4–6, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 2002, Translated from the 1968 French original by Andrew Pressley.
- [BBK] S. Brenner, M. C. R. Butler, A. D. King, *Periodic Algebras which are Almost Koszul*, Algebr. Represent. Theory 5 (2002), no. 4, 331–367.
- [BY] T. Brüstle, D. Yang, *Ordered exchange graphs*, arXiv:1302.6045, to appear in Advances in Representation Theory of Algebras (ICRA Bielefeld 2012).
- [BIRS] A. B. Buan, O. Iyama, I. Reiten, J. Scott, *Cluster structures for 2-Calabi-Yau categories and unipotent groups*, Compos. Math. 145 (2009), 1035–1079.
- [CE] H. Cartan, S. Eilenberg, Homological algebra, Princeton University Press, Princeton, NJ, 1956.
- [CB] W. Crawley-Boevey, On the exceptional fibres of Kleinian singularities, Amer. J. Math. 122 (2000), no. 5, 1027–1037.
- [DK] R. Dehy, B. Keller, On the combinatorics of rigid objects in 2-Calabi-Yau categories, Int. Math. Res. Not. IMRN 2008, no. 11, Art. ID rnn029.
- [DR1] V. Dlab, C. M. Ringel, The preprojective algebra of a modulated graph, Representation theory, II (Proc. Second Internat. Conf., Carleton Univ., Ottawa, Ont., 1979), pp. 216–231, Lecture Notes in Math., 832, Springer, Berlin-New York, 1980.
- [DR2] V. Dlab, C. M. Ringel, The module theoretical approach to quasi-hereditary algebras, Representations of algebras and related topics (Kyoto, 1990), 200–224, London Math. Soc. Lecture Note Ser., 168, Cambridge Univ. Press, Cambridge, 1992.
- [GLS1] C. Geiss, B. Leclerc, J. Schröer, *Rigid modules over preprojective algebras*, Invent. Math. 165 (2006), no. 3, 589–632.
- [GLS2] C. Geiss, B. Leclerc, J. Schröer, Semicanonical bases and preprojective algebras II: A multiplication formula, Compos. Math. 143 (2007), no. 5, 1313–1334.
- [GLS3] C. Geiss, B. Leclerc, J. Schröer, *Kac-Moody groups and cluster algebras*, Adv. Math. 228 (2011), no. 1, 329–433.
- [GP] I. M. Gelfand, V. A. Ponomarev, *Model algebras and representations of graphs*, Funktsional. Anal. i Prilozhen. 13 (1979), no. 3, 1–12.
- [H] J. E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics, 29. Cambridge University Press, Cambridge, 1990.
- [IT] C. Ingalls, H. Thomas, Noncrossing partitions and representations of quivers, Compos. Math. 145 (2009), no. 6, 1533–1562.
- [IJ] O. Iyama, G. Jasso, in preparation.
- [IR] O. Iyama, I. Reiten, Fomin-Zelevinsky mutation and tilting modules over Calabi-Yau algebras, Amer. J. Math. 130 (2008), no. 4, 1087–1149.
- [KY] S. Koenig, D. Yang, Silting objects, simple-minded collections, t-structures and co-t-structures for finite-dimensional algebras, arXiv:1203.5657.
- [L1] G. Lusztig, Quivers, perverse sheaves, and quantized enveloping algebras, J. Amer. Math. Soc. 4 (1991), no. 2, 365–421.
- [L2] G. Lusztig, Semicanonical bases arising from enveloping algebras, Adv. Math. 151 (2000), no. 2, 129–139.
- [M] Y, Mizuno, ν -stable τ -tilting modules, arXiv:1210.8322, to appear in Comm. Algebra.
- [ORT] S. Oppermann, I. Reiten, H. Thomas, *Quotient closed subcategories of quiver representations*, arXiv:1205.3268.
- [P] Y. Palu, Cluster characters for 2-Calabi-Yau triangulated categories, Ann. Inst. Fourier (Grenoble) 58 (2008), no. 6, 2221–2248.
- [Re] N. Reading, Clusters, Coxeter-sortable elements and noncrossing partitions, Trans. Amer. Math. Soc., 359 (2007), no. 12, 5931–5958.
- [Ri] C. M. Ringel, The preprojective algebra of a quiver, Algebras and modules, II (Geiranger, 1996), 467–480, CMS Conf. Proc., 24, Amer. Math. Soc., Providence, RI, 1998.

Graduate School of Mathematics, Nagoya University, Frocho, Chikusaku, Nagoya, 464-8602, Japan

E-mail address: yuya.mizuno@math.nagoya-u.ac.jp