

SYMPLECTIC AND POISSON GEOMETRY OF THE MODULI SPACES OF FLAT CONNECTIONS OVER QUILTED SURFACES.

DAVID LI-BLAND AND PAVOL ŠEVERA

ABSTRACT. In this paper we study the symplectic and Poisson geometry of moduli spaces of flat connections over *quilted surfaces*. These are surfaces where the structure group varies from region to region in the surface, and where a reduction (or relation) of structure occurs along the boundaries of the regions. Our main theoretical tool is a new form moment-map reduction in the context of Dirac geometry. This reduction framework allows us to extend the results of [26, 36] to allow more general relations of structure groups, and to investigate both the symplectic and Poisson geometry of the resulting moduli spaces from a unified perspective.

The moduli spaces we construct in this way include a number of important examples, including Poisson Lie groups and their Homogeneous spaces, moduli spaces for meromorphic connections over Riemann surfaces (following the work of Philip Boalch), and various symplectic groupoids. Realizing these examples as moduli spaces for *quilted surfaces* provides new insights into their geometry.

CONTENTS

1. Introduction and Summary of Results	2
1.1. Notation and terminology	3
1.2. The construction	3
1.3. Colouring Edges	9
1.4. Domain walls and branched surfaces	12
1.5. Poisson structures	19
2. Background	26
2.1. Courant algebroids	26
2.2. Courant relations and morphisms of Manin pairs	29
2.3. Quasi-Hamiltonian manifolds	32
3. Quasi-Hamiltonian reduction	33
3.1. Reduction theorems	33
3.2. Bivector fields and quasi-Poisson structures	35
3.3. Exact morphisms of Manin pairs and 2-forms	36
3.4. Commutativity of reductions	41
4. Quasi-Hamiltonian structures on moduli spaces of flat connections	42

Date: May 5, 2022.

D.L-B. was supported by the National Science Foundation under Award No. DMS-1204779.

P.Š. was partially supported by the Swiss National Science Foundation (grants 140985 and 141329).

4.1. Sewing construction	42
4.2. The quasi-Hamiltonian structure on $\mathcal{M}_{\Sigma,V}(G)$	45
5. Poisson structures on the moduli space of flat connections	46
5.1. Fusion	47
5.2. The quasi-Hamiltonian structure for quasi-triangular structure Lie algebras	49
Appendix A. Proofs of reduction theorems	51
References	58

1. INTRODUCTION AND SUMMARY OF RESULTS

Suppose that G is a Lie group whose Lie algebra, \mathfrak{g} , is endowed with a G -invariant inner product, $\langle \cdot, \cdot \rangle$. Suppose that Σ is a closed oriented surface, and $P \rightarrow \Sigma$ is a principal G -bundle. Let $\mathcal{A}_{flat}(P \rightarrow \Sigma)$ denote the space of flat connections on P . Atiyah and Bott [6] showed that the moduli space

$$\mathcal{M}(P \rightarrow \Sigma) := \mathcal{A}_{flat}(P \rightarrow \Sigma) / \text{Aut}(P)$$

of flat connections on P carries a symplectic structure. Their construction involves infinite dimensional symplectic reduction. In this paper, we will describe a finite dimensional construction of this moduli space inspired by the quasi-Hamiltonian approach of Alekseev, Malkin and Meinrenken [4]. Our approach generalizes easily to construct symplectic structures on related moduli spaces:

- surfaces with boundary, where segments of the boundary are labelled by coisotropic subalgebras of \mathfrak{g} (generalizing some results found in [36]),
- surfaces with domains labelled by distinct structure groups and domain walls labelled by coisotropic relations between the structure groups (also called *quilted surfaces*),
- branched surfaces, where the branch locus is labelled by a coisotropic interactions between the branches.

Even more generally, our construction will produce Poisson structures, and a natural generalization of quasi-Hamiltonian and quasi-Poisson structures. As a result, we are able to provide a moduli-theoretic interpretation of a number of well known spaces including: Lu's symplectic double groupoid integrating a Poisson Lie group [28], Boalch's Fission spaces [8, 9], Poisson Lie groups [14, 32], and Poisson homogeneous spaces [30], among others.

Our construction requires two new techniques in the theory of moment maps:

- A slight generalization of group-valued moment maps, so that the moduli space on a surface with several marked points on every boundary component is equipped with such a moment map.
- A new type of symplectic reduction, generalizing the ordinary Marsden-Weinstein reduction and the quasi-Hamiltonian reduction. In particular, the moduli space

of a triangulated surface is obtained via reduction from the moduli spaces of the triangles.

Some of these results appeared in [26], where the (quasi-)Poisson structures on moduli spaces are constructed in terms of an intersection pairing. Here we present the reduction theorems in full generality (unifying both the twists and reductions found in (quasi-)Poisson geometry) and with an emphasis on symplectic structures. We also formulate the results in more natural way, as morphisms of Manin pairs. Among the morphisms of Manin pairs, we introduce the class of *exact morphisms*, corresponding to (quasi-)symplectic structures.

1.1. Notation and terminology. At this point, we would like to introduce some notation. Suppose V_i is a family of vector spaces (or manifolds) indexed by a set I and $f : J \rightarrow I$ is a map. We use the notation

$$\begin{aligned} f^! : \prod_{i \in I} V_i &\rightarrow \prod_{j \in J} V_{f(j)} \\ \{v_i\}_{i \in I} &\mapsto \{v_{f(j)}\}_{j \in J} \end{aligned}$$

for the induced pull-back map.

For any oriented graph Γ , we let E_Γ denote the set of edges, V_Γ the set of vertices and $\text{in}, \text{out} : E_\Gamma \rightarrow V_\Gamma$ the incidence maps. Γ is called a *permutation graph*¹ if both in and out are bijections.

A *quadratic Lie algebra* is a Lie algebra endowed with an invariant non-degenerate symmetric pairing.

To simplify our presentation, we will assume that the Lie group G is connected throughout this paper. The generalization to disconnected Lie groups is straightforward.

1.2. The construction.

1.2.1. Motivating example: The symplectic form from a triangulation. Let Σ be a closed oriented surface and let

$$\mathcal{M}_\Sigma(G) = \text{Hom}(\pi_1(\Sigma), G) / \mathbf{Ad}(G)$$

be the moduli space of flat connections. Let us recall how to compute the Atiyah-Bott symplectic form ω on $\mathcal{M}_\Sigma(G)$ in terms of a triangulation of Σ .

Let \mathcal{T} be a triangulation of Σ . Let \mathcal{T}_0 denote the set of its vertices, \mathcal{T}_1 the set of (unoriented) edges and \mathcal{T}_2 the set of triangles. We let $\tilde{\mathcal{T}}_1$ denote the set of oriented edges (we thus have a 2-to-1 map $\tilde{\mathcal{T}}_1 \rightarrow \mathcal{T}_1$) we let

$$(e \rightarrow \bar{e}) : \tilde{\mathcal{T}}_1 \rightarrow \tilde{\mathcal{T}}_1$$

denote the map which reverses the orientation of the edges.

Let $\mathcal{A}_{flat}(\mathcal{T})$ be the space of “combinatorial flat connections” on Σ :

$$\mathcal{A}_{flat}(\mathcal{T}) = \{g \in G^{\tilde{\mathcal{T}}_1} \mid g_{\bar{e}} = g_e^{-1} \text{ for all } e \in \tilde{\mathcal{T}}_1, \text{ and } \prod_{e \in \partial t} g_e = 1 \text{ for all } t \in \mathcal{T}_2\},$$

¹Such graphs are also called *directed cycle graphs* in the literature.




FIGURE 1. In the figure, $t \in \mathcal{T}_2$ is a triangle, $e_1, e_2, e_3 \in \tilde{\mathcal{T}}_1$ are oriented edges, and $v_1, v_2, v_3 \in \mathcal{T}_0$ are vertices. We have $\partial t = \{e_1, e_2, e_3\}$, and $v_2 = \text{in}(e_1)$ and $v_1 = \text{out}(e_1)$.

here $\partial t \subset \tilde{\mathcal{T}}_1$ denotes the oriented boundary and the product is taken in the natural (cyclic) order (cf. Fig. 1). We have an action of $G^{\mathcal{T}^0}$ on $\mathcal{A}_{flat}(\mathcal{T})$ by “gauge transformations”

$$(1.1) \quad (g' \cdot g)_e = g'_{\text{in}(e)} g_e (g'_{\text{out}(e)})^{-1}, \quad g' \in G^{\mathcal{T}^0}, \quad g \in G^{\tilde{\mathcal{T}}_1}$$

and

$$\mathcal{M}_\Sigma(G) = \mathcal{A}_{flat}(\mathcal{T}) / G^{\mathcal{T}^0}.$$

If t is an oriented triangle with edges e_1, e_2, e_3 (in their cyclic order), let

$$(1.2) \quad \mathcal{M}_t(G) = \{(g_{e_1}, g_{e_2}, g_{e_3}) \in G \times G \times G \mid g_{e_1} g_{e_2} g_{e_3} = 1\}.$$

We have an inclusion

$$i : \mathcal{A}_{flat}(\mathcal{T}) \subset \prod_{t \in \mathcal{T}_2} \mathcal{M}_t(G),$$

where the subset $\mathcal{A}_{flat}(\mathcal{T})$ is given by the condition $g_e = g_{\bar{e}}^{-1}$.

Let

$$\omega_t = \frac{1}{2} \langle g_{e_2}^{-1} dg_{e_2}, dg_{e_1} g_{e_1}^{-1} \rangle \in \Omega^2(\mathcal{M}_t(G)).$$

The 2-form ω_t is invariant under cyclic permutations of the edges.

The symplectic form ω on $\mathcal{M}_\Sigma(G)$ is given by

$$(1.3) \quad p^* \omega = i^* \sum_{t \in \mathcal{T}_2} \omega_t,$$

where $p : \mathcal{A}_{flat}(\mathcal{T}) \rightarrow \mathcal{M}_\Sigma(G)$ is the projection [41].

We shall interpret Equation (1.3) in the following way: $\mathcal{M}_\Sigma(G)$ is obtained from $\prod_{t \in \mathcal{T}_2} \mathcal{M}_t(G)$ by a variant of Hamiltonian reduction. The subset $\mathcal{A}_{flat}(\mathcal{T}) \subset \prod_{t \in \mathcal{T}_2} \mathcal{M}_t(G)$ is given by a moment map condition, and then we need to take the quotient by the residual group $G^{\mathcal{T}^0}$ to get a symplectic manifold. To do it, we need to explain this (quasi-)Hamiltonian reduction and the (quasi-)Hamiltonian structure on $\mathcal{M}_t(G)$.

1.2.2. *Quasi-Hamiltonian reduction.* Let \mathfrak{d} be a quadratic Lie algebra and $\mathfrak{h} \subset \mathfrak{d}$ a Lagrangian subalgebra (i.e. $\mathfrak{h}^\perp = \mathfrak{h}$). In other words, $(\mathfrak{d}, \mathfrak{h})$ is a Manin pair.

Suppose that \mathfrak{d} acts on a manifold N so that all the stabilizers are coisotropic Lie subalgebras of \mathfrak{d} . We shall recall below the following notions (introduced by Alekseev, Malkin and Meinrenken in [4] and by Alekseev, Kosmann-Schwarzbach and Meinrenken in [3], slightly generalized in this paper):

- A *quasi-Hamiltonian* $(\mathfrak{d}, \mathfrak{h}) \times N$ -manifold (or quasi-Hamiltonian \mathfrak{h} -manifold, if \mathfrak{d} and N are clear from the context) is a manifold M with an action of \mathfrak{h} , an \mathfrak{h} -equivariant map $\mu : M \rightarrow N$ (*moment map*), and a bivector field π on M , satisfying certain conditions.²
- Among the moment maps there are *exact moment maps*. In this case the bivector field π can be replaced by a 2-form (M is “quasi-symplectic”).

One of our main results is the following reduction theorem:

Theorem 1.1. *Let M be a quasi-Hamiltonian $(\mathfrak{d}, \mathfrak{h}) \times N$ -manifold, $\mathfrak{l} \subset \mathfrak{d}$ a Lagrangian Lie subalgebra, and $S \subset N$ an \mathfrak{l} -invariant submanifold.*

- (1) *There is a natural Poisson bracket on the algebra $C^\infty(M)^{\mathfrak{l} \cap \mathfrak{h}} \subset C^\infty(M)$ of $\mathfrak{l} \cap \mathfrak{h}$ -invariant functions. In particular, if $M/(\mathfrak{l} \cap \mathfrak{h})$ is a manifold, it is a Poisson manifold.*
- (2) *The ideal $I \subset C^\infty(M)^{\mathfrak{l} \cap \mathfrak{h}}$ of functions vanishing on $\mu^{-1}(S)$ is a Poisson ideal. In particular, $\mu^{-1}(S)/(\mathfrak{l} \cap \mathfrak{h})$ is a Poisson manifold, provided it is a manifold.*
- (3) *If the moment map μ is exact and S is an \mathfrak{l} -orbit then the Poisson manifold $\mu^{-1}(S)/(\mathfrak{l} \cap \mathfrak{h})$ is symplectic.*

More generally, if in place of the Lagrangian subalgebra \mathfrak{l} we use a coisotropic subalgebra, we have a similar result, where the reduced manifold is still quasi-Hamiltonian. This result is contained in Theorems 3.1 and 3.2, expressed in the more appropriate language of morphisms of Manin pairs.

1.2.3. *The quasi-Hamiltonian structure on moduli spaces.* Let e be an (abstract) oriented edge, let $N_e = G$ and $\mathfrak{d}_e = \bar{\mathfrak{g}} \oplus \mathfrak{g}$, where $\bar{\mathfrak{g}}$ is \mathfrak{g} with the inner product negated. The corresponding group $D_e = G \times G$ acts on $N_e = G$ via

$$(1.4) \quad (g_1, g_2) \cdot g = g_1 g g_2^{-1}.$$

$N_e = G$ should be imagined as the space of possible holonomies along e , and the action of $D_e = G \times G$ as gauge transformations at the endpoints of e .

Let Σ be a compact oriented surface and $V \subset \partial\Sigma$ a finite subset such that every component of both Σ and $\partial\Sigma$ intersects V non-trivially. We shall call (Σ, V) a *marked surface*. The boundary circles of Σ are cut into a sequence of oriented edges with endpoints in V . Together these edges and vertices form a permutation graph Γ , the *boundary graph* of

²Strictly speaking the bivector field π depends in an inessential way on a choice of a vector space complement $\mathfrak{k} \subset \mathfrak{d}$ to \mathfrak{h} , as in [2]. Similarly, in the exact case, the 2-form depends in an inessential way on some other choice. These choices can be made canonically in our cases of interest, and so we will ignore this subtlety until Section 2.3.




FIGURE 2. The marked surface (Σ, V) , with $V = \{v_1, v_2, v_3\}$. The boundary graph Γ has edges $E_\Gamma = \{e_1, e_2, e_3\}$ and vertices $V_\Gamma = V$.

(Σ, V) (cf. Fig. 2). Let $\Pi_1(\Sigma, V)$ denote the fundamental groupoid of Σ with the base set V . Let

$$\mathcal{M}_{\Sigma, V}(G) = \text{Hom}(\Pi_1(\Sigma, V), G)$$

be the moduli space of flat connections on G -bundles over Σ trivialized at V . This moduli space is quasi-Hamiltonian in the following way:

We have an action of the group $H = G^V$ on $\mathcal{M}_{\Sigma, V}(G)$ by (residual) gauge transformations,

$$(h \cdot f)(e) = h_{\text{in}(e)} f(e) h_{\text{out}(e)}^{-1},$$

for $h \in G^V$, $f \in \mathcal{M}_{\Sigma, V}(G)$, and $e \in \Pi_1(\Sigma, V)$. We also have a map

$$\mu : \mathcal{M}_{\Sigma, V}(G) \rightarrow N := \prod_{e \in E_\Gamma} N_e.$$

where the components of μ are given by

$$\mu(f)_e = f(e)$$

(in other words, μ is the list of holonomies along the boundary arcs). Notice that the map μ is H -equivariant, where $H = G^V$ embeds as a subgroup

$$G^V \subseteq D := \prod_{e \in E_\Gamma} D_e = \prod_{e \in E_\Gamma} (G \times G).$$

Here $g \in G^V$ is included as the element $\prod_{e \in E_\Gamma} (g_{\text{in}(e)}, g_{\text{out}(e)})$. Letting \mathfrak{d} and \mathfrak{h} denote the Lie algebras of D and H , we have:

Theorem 1.2. *There is a natural $(\mathfrak{d}, \mathfrak{h}) \times N$ -quasi-Hamiltonian structure on $\mathcal{M}_{\Sigma, V}(G)$ with the moment map μ . The moment map is exact and the quasi-symplectic form ω on $\mathcal{M}_{\Sigma, V}(G)$ is given by the formula (1.3), where \mathcal{T} is any triangulation of Σ such that $\mathcal{T}_0 \cap \partial\Sigma = V$.*

We prove this theorem in Section 4.

Remark 1.1. In the case where every boundary component of Σ contains exactly one element of V , the theorem (except for the triangulation part) was proved by Alekseev, Malkin and Meinrenken in [4], and became the motivation for quasi-Hamiltonian structures.

1.2.4. *Reduction applied to moduli spaces.* We can combine Theorems 1.1 and 1.2 to produce Poisson and symplectic manifolds: we choose a collection (Σ_i, V_i) of marked surfaces with boundary graphs Γ_i , and a collection G_i of Lie groups with quadratic Lie algebras. The manifold

$$\mathcal{M} := \prod_i \mathcal{M}_{\Sigma_i, V_i}(G_i)$$

is quasi-Hamiltonian, with the moment map $\mu : \mathcal{M} \rightarrow N = \prod_i N_i$. We choose a Lagrangian Lie subalgebra $\mathfrak{l} \subset \mathfrak{d}$ and a \mathfrak{l} -invariant submanifold $S \subset N$. Then by Theorem 1.1, if the transversality conditions are satisfied, the manifold

$$\mathcal{M}_{red} = \mu^{-1}(S)/(\mathfrak{l} \cap \mathfrak{h})$$

is symplectic or Poisson.


The reduced manifold \mathcal{M}_{red} can be again seen as a moduli space of flat connections, with certain boundary (or sewing) conditions. Below we shall give various examples for simple choices of \mathfrak{l} and S .

Example 1.1. As the first example, let Σ be a closed surface with a triangulation \mathcal{T} . Let (Σ', V') be the disjoint union of the triangles, with V' consisting of the vertices, and let


$$\mathcal{M} = \mathcal{M}_{\Sigma', V'}(G) = \prod_{t \in \mathcal{T}_2} M_t(G).$$

in the notation of Eq. (1.2).

Let us now identify our data on a picture (showing just two triangles, with the parallel edges identified in Σ):



The Lie algebra \mathfrak{d} is the direct sum of all the \mathfrak{g} 's and $\bar{\mathfrak{g}}$'s, situated at the half-edges of the triangles. N is the product of all G 's. The Lie algebra $\mathfrak{h} \subset \mathfrak{d}$ is the direct sum of all the diagonal Lie subalgebras, $\mathfrak{g}_\Delta \subset \mathfrak{g} \oplus \bar{\mathfrak{g}}$, situated at the vertices of the triangles. Let the Lie algebra $\mathfrak{l} \subset \mathfrak{d}$ be the direct sum of all the diagonals $\mathfrak{g}_\Delta \subset \mathfrak{g} \oplus \bar{\mathfrak{g}}$ situated at the pairs of half-edges that are identified in Σ . Notice that $\mathfrak{h} \cap \mathfrak{l} = \mathfrak{g}^{\mathcal{T}_0}$.



(A) Elements of \mathfrak{h} lie in the direct sum of all the diagonals $\mathfrak{g}_\Delta \subset \mathfrak{g} \oplus \bar{\mathfrak{g}}$ at the vertices of the triangles.

(B) Elements of \mathfrak{l} lie in the direct sum of all the diagonals $\mathfrak{g}_\Delta \subset \mathfrak{g} \oplus \bar{\mathfrak{g}}$ at the pairs of half-edges that are identified in Σ . The \mathfrak{l} -orbit, S , consists of those elements $g \in \prod_{e \in \tilde{\mathcal{T}}_1} \mathfrak{g}_e$ such that $g_e = g_{\bar{e}}^{-1}$.

For the \mathfrak{l} -orbit $S \subset N$ we take the subset given by the conditions $g_{\bar{e}} = g_e^{-1}$ for any pair of edges e, \bar{e} that are identified in Σ . We have

$$\mathcal{M}_\Sigma(G) = \mathcal{M}_{red} := \mu^{-1}(S)/(\mathfrak{h} \cap \mathfrak{l}).$$

Thus we are able to obtain $\mathcal{M}_\Sigma(G)$ by quasi-Hamiltonian reduction from triangles.

So far we have not explicitly described the symplectic or Poisson structure on \mathcal{M}_{red} . In a special case it is very simple. Let

$$\mu_i : \mathcal{M}_{\Sigma_i, V_i}(G_i) \rightarrow N_i$$

denote the (exact) moment map, and let ω_i be the quasi-symplectic 2-form on $\mathcal{M}_{\Sigma_i, V_i}(G_i)$ (given explicitly in Theorem 1.2). For every boundary arc e of Σ_i we have the involution of $\mathfrak{d}_e = \bar{\mathfrak{g}}_i \oplus \mathfrak{g}_i$ given by

$$(\xi, \eta) \mapsto (\eta, \xi).$$

If we apply the involution simultaneously at all the boundary arcs, we get an involution of

$$\mathfrak{d} = \bigoplus_i \bigoplus_{e \in \partial \Sigma_i} \bar{\mathfrak{g}}_i \oplus \mathfrak{g}_i.$$

We shall say that a subalgebra $\mathfrak{l} \subset \mathfrak{d}$ is *symmetric* if it is invariant with respect to this involution.

Theorem 1.3. *If $\mathfrak{l} \subset \mathfrak{d}$ is a symmetric Lagrangian subalgebra and $S \subset N$ is the \mathfrak{l} -orbit through the identity element*

$$1 \in N = \prod_i G_i^{E_{\Gamma_i}},$$

then the symplectic form ω_{red} on

$$\mathcal{M}_{red} = \mu^{-1}(S)/\mathfrak{l} \cap \mathfrak{h}$$

is given by

$$p^* \omega_{red} = \sum_i \omega_i \Big|_{\mu^{-1}(S)}$$

where $p : \mu^{-1}(S) \rightarrow \mathcal{M}_{red}$ is the projection.

As explained in Remark 3.3, Theorem 1.3 will follow as a corollary to Proposition 3.1.

1.3. Colouring Edges. Suppose that $\mathfrak{c} \subseteq \mathfrak{g}$ is a coisotropic subalgebra (i.e. $\mathfrak{c}^\perp \subseteq \mathfrak{c}$). Then the subalgebra

$$\mathfrak{l}_{\mathfrak{c}} := \{(\xi, \eta) \in (\bar{\mathfrak{g}} \oplus \mathfrak{g}) \mid \xi, \eta \in \mathfrak{c} \text{ and } \xi - \eta \in \mathfrak{c}^\perp\}$$

is both Lagrangian and symmetric. The orbit of $\mathfrak{l}_{\mathfrak{c}}$ through the identity of G , with respect to the action Eq. (1.4), can be identified with the simply connected Lie group C^\perp integrating the Lie algebra \mathfrak{c}^\perp .

Let (Σ, V) be a marked surface. For every boundary arc e (i.e. for every edge of the permutation graph $\Gamma_{\Sigma, V}$ with the vertex set V), let $\mathfrak{c}_e \in \mathfrak{g}$ be a coisotropic subalgebra, and consider the Lie subalgebra

$$\mathfrak{l} := \bigoplus_e \mathfrak{l}_{\mathfrak{c}_e} \subset \bigoplus_e \bar{\mathfrak{g}} \oplus \mathfrak{g} = \mathfrak{d}.$$

It is clear that \mathfrak{l} is both Lagrangian and symmetric. Let $S \subset N = \prod_e G$ be the \mathfrak{l} -orbit passing through $1 \in \prod_e G$. Theorem 1.3 implies that if the quotient space

$$\mathcal{M}_{red} = \mu^{-1}(S)/\mathfrak{l} \cap \mathfrak{g}^V$$

is a manifold, it is symplectic.

Concretely,

$$(1.5) \quad \mathcal{M}_{red} = \{f : \Pi_1(\Sigma, V) \rightarrow G \mid f(e) \in C_e^\perp \text{ for every } e\}/\mathfrak{l} \cap \mathfrak{g}^V,$$

and $\mathfrak{l} \cap \mathfrak{g}^V \subset \mathfrak{g}^V$ is given by the conditions

$$(1.6a) \quad \xi_v \in \mathfrak{c}_{e_1} \cap \mathfrak{c}_{e_2} \text{ where } v = \text{in}(e_1) = \text{out}(e_2)$$

$$(1.6b) \quad \xi_{\text{in}(e)} - \xi_{\text{out}(e)} \in \mathfrak{c}_e^\perp.$$

Notice that if \mathfrak{c}_e 's are Lagrangian then the first condition implies the second one. If, moreover, $\mathfrak{c}_{e_1} \cap \mathfrak{c}_{e_2} = 0$ for any pair of consecutive boundary arcs then $\mathfrak{l} \cap \mathfrak{g}^V = 0$. Under these conditions the moduli space \mathcal{M}_{red} was considered in [36].




FIGURE 3. The symplectic double groupoid integrating the Lie-Poisson structures on E and F .




FIGURE 4. The symplectic double groupoid integrating the Lie-Poisson structure on G .

Example 1.2 ([35, 36]). Suppose that $\mathfrak{e}, \mathfrak{f} \subseteq \mathfrak{g}$ are transverse Lagrangian subalgebras, and let $E, F \subset G$ denote the corresponding connected Lie groups. We may colour alternate edges of a rectangle with \mathfrak{e} and \mathfrak{f} , as in Fig. 3. From Eq. (1.5) we see that

$$\mathcal{M}_{red} = \{(e_1, e_2, f_1, f_2) \in E^2 \times F^2 \mid e_1 f_1 e_2 f_2 = 1\}.$$

By Theorem 1.3 the moduli space \mathcal{M}_{red} carries the symplectic form

$$\omega = \frac{1}{2} \langle e_1^{-1} de_1, df_1 f_1^{-1} \rangle + \frac{1}{2} \langle e_2^{-1} de_2, df_2 f_2^{-1} \rangle.$$

Here, the upper-left triangle in Fig. 3 contributed the term $\frac{1}{2} \langle e_1^{-1} de_1, df_1 f_1^{-1} \rangle$ to this expression while the bottom-right triangle in Fig. 3 contributed the term $\frac{1}{2} \langle e_2^{-1} de_2, df_2 f_2^{-1} \rangle$.

As explained in [35, 36], the symplectic manifold (\mathcal{M}, ω) is the Lu-Weinstein symplectic double groupoid integrating the Lie-Poisson structures on E and F [29].

Example 1.3 ([34, 36]). Let $\mathfrak{e}, \mathfrak{f} \subseteq \mathfrak{g}$ be as above. Divide each boundary component of the annulus into two segments and colour alternate edges with \mathfrak{e} and \mathfrak{f} , as in Fig. 4. From Eq. (1.5) we see that

$$\mathcal{M}_{red} = \{(e_1, e_2, f_1, f_2, g) \in E^2 \times F^2 \times G \mid g e_1 f_1 g^{-1} f_2 e_2 = 1\}.$$




FIGURE 5. The symplectic groupoid integrating the Lu-Yakimov Poisson structure on G/C .

The moduli space \mathcal{M}_{red} carries the symplectic form

$$\begin{aligned} \omega = \frac{1}{2} \langle e_2^{-1} de_2, dg g^{-1} \rangle + \frac{1}{2} \langle (e_2 g)^{-1} d(e_2 g), de_1 e_1^{-1} \rangle \\ + \frac{1}{2} \langle g f_1^{-1} d(f_1 g^{-1}), g f_2 f_2^{-1} \rangle - \frac{1}{2} \langle dg g^{-1}, df_1 f_1^{-1} \rangle, \end{aligned}$$

which can be computed from the triangulation pictured in Fig. 4.

As explained in [34,36], the symplectic manifold (\mathcal{M}, ω) is the symplectic double groupoid integrating the Lie-Poisson structure on G .

Example 1.4. Suppose that $\mathfrak{e}, \mathfrak{f} \subseteq \mathfrak{g}$ are transverse Lagrangian subalgebras and $\mathfrak{c} \subseteq \mathfrak{g}$ is a coisotropic subalgebra. Let $E, F, C, C^\perp \subset G$ denote the corresponding connected Lie subgroups, and suppose that $C \subset G$ is closed. Consider the annulus whose outer boundary is divided into two segments. Colour the outer boundary by the two Lie subalgebras \mathfrak{e} and \mathfrak{f} and the inner boundary by the inner boundary by the Lie subalgebra \mathfrak{c} , as in Fig. 5. We have

$$\mu^{-1}(S) = \{(e, f, g, c) \in E \times F \times G \times C^\perp \mid e f g c g^{-1} = 1\}.$$

Meanwhile, Eq. (1.6) yields

$$\mathfrak{l} \cap \mathfrak{g}^V = \{\xi \in \mathfrak{g}^V \mid \xi_{v_0} \in \mathfrak{c} \text{ and } \xi_v = 0 \text{ for all } v \neq v_0\},$$

where v_0 is the vertex labelled in Fig. 5. Thus the Lie group of residual gauge transformations is C , acting as

$$c' \cdot (e, f, g, c) = (e, f, g c'^{-1}, c' c c'^{-1}), \quad c' \in C, \quad (e, f, g, c) \in E \times F \times G \times C^\perp.$$

Since, by assumption this acts freely and properly on $\mu^{-1}(N)$, Theorem 1.3 implies that the moduli space

$$\mathcal{M}_{red} = \{(e, f, g, c) \in E \times F \times G \times C^\perp \mid e f g c g^{-1} = 1\} / C$$

carries the symplectic form

$$\omega = -\frac{1}{2} \langle dg g^{-1}, de e^{-1} \rangle + \frac{1}{2} \langle c^{-1} dc, d(g^{-1} e) e^{-1} g \rangle + \frac{1}{2} \langle f^{-1} df, dg g^{-1} \rangle.$$

The symplectic manifold $(\mathcal{M}_{red}, \omega)$ is the symplectic groupoid integrating the Lu-Yakimov Poisson structure on the homogeneous space G/C [30]. The source and target maps are

$$s(e, f, g, c) = g, \quad t(e, f, g, c) = fg,$$

and the multiplication is

$$(e', f', g', c') \cdot (e, f, g, c) = (ee', f'f, g, c'), \quad g' = fg.$$

1.4. Domain walls and branched surfaces. Let (Σ_i, V_i) be a finite collection of marked surfaces with boundary graphs Γ_i , and G_i a collection of Lie groups with quadratic Lie algebras \mathfrak{g}_i . As we observed above, the space

$$\mathcal{M} = \prod_i \mathcal{M}_{\Sigma_i, V_i}(G_i)$$

is a $(\mathfrak{d}, \mathfrak{h}) \times N$ -quasi-Hamiltonian for appropriate $(\mathfrak{d}, \mathfrak{h}, N)$, and if we choose a Lagrangian Lie subalgebra $\mathfrak{l} \subset \mathfrak{d}$ and a \mathfrak{l} -orbit $S \subset N$, then

$$\mathcal{M}_{red} = \mu^{-1}(S)/\mathfrak{l} \cap \mathfrak{h}$$

is symplectic. If the subalgebra $\mathfrak{l} \subset \mathfrak{d}$ is symmetric then Theorem 1.3 gives us a simple formula for the symplectic form on \mathcal{M}_{red} .

Let us now choose a symmetric $\mathfrak{l} \subset \mathfrak{d}$ in the following way. We first glue the boundary arcs of (Σ_i, V_i) in an arbitrary way. More precisely, let W be a finite collection of (disjoint) unit intervals called *domain walls*, let

$$\kappa : \sqcup_i E_{\Gamma_i} \rightarrow W$$

be a surjective map assigning to every edge of every boundary graph Γ_i a domain wall, and let

$$\phi_e : e \rightarrow \kappa(e)$$

be a homeomorphism for every boundary edge e (not required to preserve the orientation). Let Σ be the topological space obtained from Σ_i 's and the domain walls after we identify every boundary arc e with $\kappa(e)$ via the map ϕ_e .

For every boundary arc $e \in E_{\Gamma_i}$ let $i(e) = i$, and

$$\text{sign}(e) = \begin{cases} +1 & \text{if } \phi_e \text{ is orientation-preserving} \\ -1 & \text{otherwise.} \end{cases}$$

For every domain wall $w \in W$, let

$$\mathfrak{g}_w = \bigoplus_{\substack{e \in \kappa^{-1}(w) \\ \text{sign}(e)=+1}} \mathfrak{g}_{i(e)} \oplus \bigoplus_{\substack{e \in \kappa^{-1}(w) \\ \text{sign}(e)=-1}} \bar{\mathfrak{g}}_{i(e)}$$

and

$$\mathfrak{d}_w = \bar{\mathfrak{g}}_w \oplus \mathfrak{g}_w.$$

Notice that

$$\mathfrak{d} := \bigoplus_{w \in W} \bar{\mathfrak{g}}_w \oplus \mathfrak{g}_w = \bigoplus_i (\bar{\mathfrak{g}}_i \oplus \mathfrak{g}_i)^{E_{\Gamma_i}}.$$




FIGURE 6. Our surface is divided into domains with distinct structure groups, and the domain walls are coloured by coisotropic relations between the structure groups. As before, coisotropic boundary conditions are also chosen.

For every domain wall $w \in W$ we now choose a coisotropic Lie subalgebra

$$\mathfrak{c}_w \subset \mathfrak{g}_w.$$

Using \mathfrak{c}_w we construct the symmetric Lagrangian Lie subalgebra $\mathfrak{l}_w \subset \mathfrak{d}_w$,

$$\mathfrak{l}_w := \{(\xi, \eta) \in \bar{\mathfrak{g}}_w \oplus \mathfrak{g}_w \mid \xi, \eta \in \mathfrak{c}_w, \xi - \eta \in \mathfrak{c}_w^\perp\}.$$

Finally we set

$$\mathfrak{l} = \bigoplus_{w \in W} \mathfrak{l}_w.$$

For every domain wall $w \in W$, let $C_w^\perp \subset G_w$ denote the connected Lie subgroup with Lie algebra \mathfrak{c}_w^\perp , and

$$C^\perp = \prod_{w \in W} C_w^\perp \subseteq \prod_i G_i^{E_{\Gamma_i}}.$$

Then $S := C^\perp \subset N$ is the \mathfrak{l} -orbit passing through $1 \in \prod_i G_i^{E_{\Gamma_i}} = N$. As before, we have

$$(1.7) \quad \mu^{-1}(S) = \left\{ \{f_i : \Pi_1(\Sigma_i, V_i) \rightarrow G_i\}_i \mid \prod_i \{f_i(e)^{\text{sign}(e)}\}_{e \in E_{\Gamma_i}} \in C^\perp \right\}$$

and

$$\mathcal{M}_{red} = \mu^{-1}(S) / \mathfrak{l} \cap \bigoplus_i \mathfrak{g}_i^{V_i}.$$

Example 1.5 (Oriented surfaces with coloured boundaries). If we have just one domain and the gluing map κ is injective, then we are in the case described in Section 1.3.

1.4.1. Domain walls. Suppose that the glued topological space Σ is still a (not necessarily oriented) surface. Equivalently, every domain wall $w \in W$ borders either one or two domains (i.e. the preimage $\kappa^{-1}(w)$ has cardinality one or two). The resulting surface Σ was called a *quilted surface* in [26] (following [40]).

Remark 1.2. Quantizations of these moduli spaces have been studied in the physics community [17, 20, 21] for abelian structure groups and Lagrangian relations on the domain walls.




FIGURE 7. On the surface pictured above, the structure group in the yellow domain is G while the structure group in the blue domain is H . Along the boundary of the two domains, blue edges are coloured with \mathfrak{c}_+ while the red edges are coloured with \mathfrak{c}_- . Cutting along the dotted line in the first picture yields the second picture. Acting by H at the vertices v_1, \dots, v_{2r} allows us to set the holonomies h_0, \dots, h_{2r-1} to the identity.

Example 1.6 ([9]). Suppose that $\mathfrak{g} = \mathfrak{u}_+ \oplus \mathfrak{h} \oplus \mathfrak{u}_-$ as a vector space (but not as a Lie algebra), where $\mathfrak{p}_\pm := \mathfrak{h} \oplus \mathfrak{u}_\pm \subseteq \mathfrak{g}$ are coisotropic subalgebras satisfying $\mathfrak{p}_\pm^\perp = \mathfrak{u}_\pm$. Suppose further that the Lie subalgebras $\mathfrak{u}_\pm, \mathfrak{p}_\pm, \mathfrak{h}$ all integrate to closed subgroups $U_\pm, P_\pm, H \subseteq G$ such that $H = P_+ \cap P_-$. The metric on \mathfrak{g} descends to a non-degenerate invariant metric on $\mathfrak{h} \subseteq \mathfrak{g}$, and

$$(1.8) \quad \mathfrak{c}_\pm := \{(\xi; \xi + \mu) \in \mathfrak{h} \oplus \mathfrak{g} \mid \xi \in \mathfrak{h} \text{ and } \mu \in \mathfrak{u}_\pm\}$$

is a coisotropic subalgebra (in fact, it is Lagrangian).

As in Fig. 7, let Σ denote the annulus, and let $\gamma \subset (\Sigma \setminus \partial\Sigma)$ be a simple closed curve representing the generator of the fundamental group. Cutting Σ along γ yields two annuli, $\Sigma_G, \Sigma_H \subset \Sigma$, which we label with the structure groups G and H , respectively. We divide γ into $2r$ segments with endpoints labelled v_1, \dots, v_{2r} , and colour alternating segments with the coisotropic Lie subalgebras \mathfrak{c}_+ and \mathfrak{c}_- . Finally, we mark the respective components of $\partial\Sigma$ with points x_G and x_H , and label the ends of the $2r$.

The points $x_G, x_H, v_1, \dots, v_{2r}$ form the vertices of a triangulation of Σ , as pictured in Fig. 7. Now the orbit of \mathfrak{l}_\pm through the identity is P_\pm . Thus, from Eq. (1.7), we see that

$$\begin{aligned} \mu^{-1}S = \{ & (h, h_0, \dots, h_{2r-1}; C_0, C_1, \dots, C_{2r}) \in H^{2r+1} \times G^{2r+1} \\ & \mid h_{2i+1}^{-1} C_{2i+1} C_{2i}^{-1} h_{2i} \in U_+ \text{ and } h_{2i}^{-1} C_{2i} C_{2i-1}^{-1} h_{2i-1} \in U_-, \} \end{aligned}$$

where the elements $h, h_0, \dots, h_{2r-1} \in H$ and $C_0, C_1, \dots, C_{2r} \in G$ denote the appropriately labelled holonomies in Fig. 7.

On the other hand,

$$\mathfrak{l} \cap \bigoplus_i \mathfrak{g}_i^{V_i} \cong \prod_{v_1, \dots, v_{2r}} \mathfrak{h},$$

acting at the appropriate vertices. Thus, up to a gauge transformation, we may assume that $h_0 = h_1 = \dots = h_{2r-1} = 1$. Setting $S_i = C_i C_{i-1}^{-1}$, we see that the quotient space, $\text{hol}^{-1}(\mathfrak{l} \cdot 1) / (\mathfrak{l} \cap \prod_{t \in \mathcal{T}_2} (\mathfrak{g}_t)_{\Gamma_{P_3}})$, can be identified with

$${}_G \mathcal{A}_H^r := \{(h; S_{2r}, \dots, S_1; C_0) \in H \times (U_- \times U_+)^r \times G\}.$$

We compute the two form to be

$$\begin{aligned} \omega = -\frac{1}{2} & (\langle d(hC_{2r}) (hC_{2r})^{-1}, dC_0 C_0^{-1} \rangle + \langle (hC_{2r})^{-1} d(hC_{2r}), C_{2r-1}^{-1} dC_{2r-1} \rangle \\ & + \sum_{i=1}^{2r-1} \langle C_i^{-1} dC_i, C_{i-1}^{-1} dC_{2i-1} \rangle). \end{aligned}$$

Substituting $bC_0 = hC_{2r}$ in the first term and simplifying yields

$$\begin{aligned} \omega = \frac{1}{2} & (\langle dC_0 C_0^{-1}, \mathbf{A} d_b dC_0 C_0^{-1} \rangle + \langle dC_0 C_0^{-1}, db b^{-1} \rangle + \langle dC_{2r} C_{2r}^{-1}, h^{-1} dh \rangle \\ & + \sum_{i=1}^{2r} \langle C_i^{-1} dC_i, C_{i-1}^{-1} dC_{2i-1} \rangle), \end{aligned}$$

(here we have used the fact that $\langle dS_{2r} S_{2r}^{-1}, h^{-1} dh \rangle = 0$). Now, we haven't coloured the boundary of $\partial\Sigma$, so Theorem 1.3 does not imply that ω is symplectic. Nevertheless, as we shall see later, Theorem 3.2 implies that ω defines a quasi-Hamiltonian $G \times H$ structure on ${}_G \mathcal{A}_H^r$, where the moment map ${}_G \mathcal{A}_H^r \rightarrow G \times H$ is given by the holonomy along the (oriented) boundary components:

$$(h; S_{2r}, \dots, S_1; C_0) \rightarrow (C_0^{-1} h S_{2r} \dots S_1 C_0, h^{-1}),$$

and the G and H actions on ${}_G \mathcal{A}_H^r$ are precisely the residual gauge transformations. These act by G at x_G and by H at x_H :

$$(g, k) \cdot (h; S_{2r}, \dots, S_1; C_0) = (khk^{-1}, kS_{2r}k^{-1}, \dots, kS_1k^{-1}, kC_0g^{-1}), \quad g \in G, \quad k \in H.$$

Remark 1.3. This quasi-Hamiltonian $G \times H$ -space was first discovered by Boalch [7–9], who used it to study meromorphic connections on Riemann surfaces. Realizing it as a moduli space is one of the motivations for considering quilted surfaces.

1.4.2. Branched Surfaces. We can now consider examples where Σ is not a topological surface, i.e. where the domain walls may border more than two domains.

Remark 1.4. Since our gauge fields (connections on Σ) are constrained to lie in $\mathfrak{c}_w \subseteq \bigoplus_{e \in \kappa^{-1}(w)} \mathfrak{g}_e$ along the domain wall $w \in W$, one may interpret \mathfrak{c}_w as a ‘‘conservation law’’




FIGURE 8. Pictured above are three domains with structure Lie algebras \mathfrak{g}_1 , \mathfrak{g}_2 and \mathfrak{g}_3 . The three domains intersect at a branch locus, which we must colour by a coisotropic subalgebra $\mathfrak{c} \subseteq \oplus_{i=1}^3 \mathfrak{g}_i$.




FIGURE 9

for an interaction between the structure groups of the various domains glued to the domain wall w .

Example 1.7 ([7,9]). Let $V = \oplus_{i=1}^n V_i$ be a direct sum decomposition of a finite dimensional vector space, $G = \text{Gl}(V)$, and $P_+ \subseteq G$ the stabilizer for the flag

$$F_1 \subset F_2 \subset \cdots \subset F_n = V,$$

where $F_k = \oplus_{i=1}^k V_i$. Similarly, let $P_- \subseteq G$ be the stabilizer for the flag

$$\tilde{F}_n \subset \cdots \subset \tilde{F}_2 \subset \tilde{F}_1 = V,$$

where $\tilde{F}_k = \oplus_{i=k}^n V_i$. Finally, let $H_i = \text{Gl}(V_i)$ so that $\prod_{i=1}^n H_i = P_+ \cap P_-$. Let $H = \prod_{i=1}^n H_i$, let U_{\pm} denote the unipotent radicals of P_{\pm} , and let $\mathfrak{g}, \mathfrak{p}_{\pm}, \mathfrak{u}_{\pm}, \mathfrak{h}, \mathfrak{h}_i$ denote the Lie algebras corresponding to the various Lie groups.




FIGURE 10. Each of the edges above are coloured by \mathbf{c}_n , where n is the number of domains branching off the given edge. As we cross a branch locus, the orientation of the domains reverses. As depicted, we may move branch loci past each other.

Now consider the moduli space

$${}_G\mathcal{A}_H^r := \left(\prod_{i=1}^n H_i \right) \times (U_- \times U_+)^r \times G$$

described in Example 1.6. The coisotropic Lie algebra

$$\mathbf{c}_\pm := \left\{ \left(\sum_{i=1}^n \xi_i; \sum_{i=1}^n \xi_i + \mu \right) \mid \xi_i \in \mathfrak{h}_i \text{ and } \mu \in \mathfrak{u}_\pm \right\} \subset (\oplus_{i=1}^n \mathfrak{h}_i) \oplus \mathfrak{g}$$

defined in Eq. (1.8) can be used to colour the branch locus of $n + 1$ domains with the structure groups H_1, \dots, H_n and G . Thus we may interpret ${}_G\mathcal{A}_H^r$ as the moduli space of flat connections for the branched surface pictured in Fig. 9.

Remark 1.5. The quasi-Hamiltonian space ${}_G\mathcal{A}_H^r$ first appeared in the work of Philip Boalch [7, 9]. Realizing it as a moduli space is one of the motivations for considering branched surfaces.

Example 1.8 (quasi-triangular structures). Let \mathfrak{g} be a quasi-triangular Lie quasi-bialgebra, i.e. \mathfrak{g} is a Lie algebra with a chosen element $s \in (S^2\mathfrak{g})^\mathfrak{g}$. Let \mathfrak{d} be the Drinfel'd double of \mathfrak{g} . This means that \mathfrak{d} is a quadratic Lie algebra, $\mathfrak{g} \subset \mathfrak{d}$ is a Lagrangian subalgebra, and $\mathfrak{p} \subset \mathfrak{d}$ is an ideal such that $\mathfrak{d} = \mathfrak{g} \oplus \mathfrak{p}$ as a vector space. Additionally, the restriction of the quadratic form on \mathfrak{d} to $\mathfrak{p} \cong \mathfrak{g}^*$ is s .³

³These properties uniquely define \mathfrak{d} . In particular, the Lie bracket is given by

$$[\xi + \alpha, \eta + \beta] = [\xi, \eta] + \mathbf{ad}_\xi^* \alpha - \mathbf{ad}_\eta^* \beta, \quad \xi, \eta \in \mathfrak{g}, \quad \alpha \in \mathfrak{p}, \beta \in \mathfrak{p}^\perp$$

where \mathbf{ad}^* denotes the contragredient representation of \mathfrak{g} .

There is a natural groupoid structure on \mathfrak{d} , where \mathfrak{g} is the space of objects and composition is defined by

$$(\xi + \alpha)(\xi + \beta) = \xi + \alpha + \beta \quad \forall \xi \in \mathfrak{g}, \alpha \in \mathfrak{p}, \beta \in \mathfrak{p}^\perp,$$

and the source and target maps are

$$s(\xi + \alpha) = \xi, \quad t(\xi + \alpha) = \xi + s(\alpha, \cdot), \quad \xi \in \mathfrak{g}, \alpha \in \mathfrak{p}$$

The graph of multiplication,

$$(1.9) \quad \text{gr}(\text{Mult}) = \{(\xi\eta, \xi, \eta) \mid \xi, \eta \in \mathfrak{d} \text{ are composable}\} \subseteq \mathfrak{d} \oplus \bar{\mathfrak{d}} \oplus \bar{\mathfrak{d}},$$

is a Lagrangian Lie subalgebra. See [15] and [24] for more details.

Similarly, the graph of iterated multiplication

$$\mathfrak{c}_n := \{(\xi_1, \dots, \xi_n) \mid \xi_1 \xi_2 \cdots \xi_n \in \mathfrak{g}\} \subseteq \mathfrak{d}^n$$

is also a Lagrangian Lie subalgebra. As such it can be used to colour the branch locus of n domains each with structure group D (a connected Lie group with the Lie algebra \mathfrak{d}). Note that crossing such a branch locus reverses the orientation of the domain.

The associativity of multiplication on \mathfrak{d} plays out as follows: paying attention to the orientations, if (as in Fig. 10) we

- move two branch loci past each other, or
- break a \mathfrak{c}_{m+n-2} -coloured branch locus into two separate \mathfrak{c}_m and \mathfrak{c}_n coloured branch loci (or vice-versa),

the resulting moduli spaces are canonically symplectomorphic.

In fact, there is a clear interpretation of the the moduli spaces constructed by sewing domains together using the Lagrangian relations \mathfrak{c}_n . One may identify them with certain traditional moduli spaces in the following way: Suppose that $\cup \Sigma_d \rightarrow \Sigma$ is our \mathfrak{c}_n -coloured surface with domains Σ_d . First we form a two sheeted (branched cover) $\tilde{\Sigma}$ of Σ as follows: double each domain Σ_d to two sheets $\Sigma_d^+ \cup \Sigma_d^-$, where the sheet Σ_d^+ is canonically identified with Σ_d , while the sheet Σ_d^- is also canonically identified with Σ_d but with the opposite orientation. At each \mathfrak{c}_n -coloured domain wall with incident domains $\Sigma_{d_1}, \dots, \Sigma_{d_n}$, cyclically glue the sheets $\Sigma_{d_1}^\pm, \dots, \Sigma_{d_n}^\pm$ together along their corresponding boundary segment, respecting the orientations, as in Fig. 11. In this way, one constructs the oriented surface $\tilde{\Sigma}$.

The groupoid inversion $\text{Inv} : \mathfrak{d} \dashrightarrow \bar{\mathfrak{d}}$, being a morphism of Lie algebras, integrates to an involution of the Lie group D . This involution in turn lifts to an involution

$$\mathcal{A}_{\tilde{\Sigma}}(D) \rightarrow \mathcal{A}_{\tilde{\Sigma}}(D)$$

of the connections on $\tilde{\Sigma}$, mapping the fibre of the principal bundle over the $+$ -sheet to the fibre over the $-$ -sheet via $\text{Inv} : D \rightarrow D$. The involution is a symplectomorphism which is compatible with the gauge transformations, and thus descends to a symplectomorphic involution on the moduli space, $\mathcal{M}_{\tilde{\Sigma}}(D)$. The fixed points of this involution are naturally identified with the moduli space $\mathcal{M}_{\Sigma}(D)$ of flat connections on the original \mathfrak{c}_n -coloured surface.




FIGURE 11. We replace each domain in our branched surface by two copies, one with the same orientation (labelled $+$), and one with the opposite orientation (labelled $-$). Each \mathfrak{c}_n -coloured domain wall is replaced by cyclically gluing the incident sheets together, respecting the orientations. In this way we obtain an oriented surface from our \mathfrak{c}_n -coloured (branched) surface.

1.5. Poisson structures. In this section, we will describe some Poisson structures which may be constructed using our approach. Later, in Section 5 we will generalize these results to the case where \mathfrak{g} is a quasi-triangular Lie quasi-bialgebra rather than a quadratic Lie algebra.

Let (Σ, V) be a marked surface with boundary graph Γ . First, recall from Theorem 1.2 that the moduli space $\mathcal{M}_{\Sigma, V}(G)$ for a marked surface (Σ, V) carries a $(\mathfrak{d}^{V_\Gamma}, \mathfrak{g}^{V_\Gamma}, G^{E_\Gamma})$ -quasi-Hamiltonian structure, where $\mathfrak{d} = \mathfrak{g} \oplus \bar{\mathfrak{g}}$, the Lagrangian Lie subalgebra $\mathfrak{g}^{V_\Gamma} = \mathfrak{g}_\Delta^{V_\Gamma} \subset \mathfrak{d}^{V_\Gamma}$ is embedded as the diagonal, and \mathfrak{d}^{V_Γ} acts on the $e \in E_\Gamma$ -th factor of G^{E_Γ} via the vector field

$$(1.10) \quad \xi_{\text{out}(e)}^L - \eta_{\text{in}(e)}^R, \quad (\xi, \eta) \in \mathfrak{g}^{V_\Gamma} \oplus \bar{\mathfrak{g}}^{V_\Gamma} = \mathfrak{d}^{V_\Gamma}.$$

Here the superscripts $^L, ^R$ denote left, right invariant vector fields. The bivector field on $\mathcal{M}_{\Sigma, V}(G)$ is computed in Section 5.2 and leads to the result of [26, Theorem 3], which we summarize briefly.⁴

⁴In fact, the computation in Section 5.2 results in minus the bivector field described in [26], due to us orienting $\partial\Sigma$ in the opposite way.

Strictly speaking, the bivector field on $\mathcal{M}_{\Sigma, V}(G)$ depends on the choice of a complement $\mathfrak{k} \subset \mathfrak{d}^{V_\Gamma}$ to $\mathfrak{h} = \mathfrak{g}^{V_\Gamma}$. In this case, \mathfrak{k} can be chosen canonically as $\mathfrak{k} := \mathfrak{g}_\Delta^{V_\Gamma}$, where

$$\mathfrak{g}_\Delta := \{(\xi, -\xi) \in (\mathfrak{g} \oplus \bar{\mathfrak{g}})\}.$$

If $a, b \in \Pi_1(\Sigma, V)$, let us represent them by transverse smooth paths α, β . For any point A in their intersection, let

$$\lambda(A) = \begin{cases} 1 & \text{if } A \in \partial\Sigma \\ 2 & \text{otherwise} \end{cases}$$

$$\text{sign}(A) := \text{sign}(\alpha, \beta; A) = \begin{cases} 1 & \text{if } (\beta'|_A, \alpha'|_A) \text{ is positively oriented} \\ -1 & \text{otherwise.} \end{cases}$$

as in Fig. 12.




FIGURE 12. $\text{sign}(A) = \pm 1$ is determined by comparing the orientation of α and β with that of Σ . The path $[\alpha_A^{-1} \beta_A]$ is shown to the right.

Let α_A denote the portion of α parametrized from the beginning up to the point A . Finally, let

$$(a, b) := \sum_A \lambda(A) \text{sign}(A) [\alpha_A^{-1} \beta_A] \in \mathbb{Z} \Pi_1(\Sigma, V).$$

(When V contains only one point, this is a skew symmetrized version of the intersection form described in [38]).

Then for any $a, b \in \Pi_1(\Sigma, V)$,

$$(1.11) \quad \pi(\text{ev}_a^*(g^{-1}dg), \text{ev}_b^*(g^{-1}dg)) = \frac{1}{2}(\mathbf{Ad}_{\text{ev}_{(a,b)}} \otimes 1) s,$$

where $\text{ev}_a, \text{ev}_b : \mathcal{M}_{\Sigma, V}(G) = \text{Hom}(\Pi_1(\Sigma, V), G) \rightarrow G$ denotes evaluation, $s \in \mathfrak{g} \otimes \mathfrak{g}$ is the inverse of the quadratic form, and $g^{-1}dg$ denotes the left invariant Maurer-Cartan form on G .

Example 1.9 (The two sided polygon, P_2). Suppose $(\Sigma, V) = P_2$ is the disk with two marked points and $E_\Gamma = \{e_1, e_2\}$, as in Fig. 13. Then we may identify $\mathcal{M}_{P_2}(G)$ with G via g_{e_1} (since $g_{e_2} = g_{e_1}^{-1}$). Under this identification, the bivector field is trivial, $\pi_{P_2} = 0$ (cf. [26]).




FIGURE 13. The cyclic product of holonomies, $g_{e_2}g_{e_1}$, is trivial, so $g_{e_2} = g_{e_1}^{-1}$.

1.5.1. *Colouring edges.* Suppose now that we colour each marked point $v \in V_\Gamma$ with a Lagrangian subalgebra $\mathfrak{l}_v \subseteq \mathfrak{d}$, and each edge $e \in E_\Gamma$ with a submanifold $S_e \subseteq G$ in a compatible way: Specifically, we require that

$$S := \prod_{e \in E_\Gamma} S_e \subseteq G^{E_\Gamma}$$

be \mathfrak{l} -invariant, where

$$\mathfrak{l} := \bigoplus_{v \in V_\Gamma} \mathfrak{l}_v \subseteq \mathfrak{d}^{V_\Gamma},$$

and the action is given in Eq. (1.10). Then Theorem 1.1 implies that

$$\mathcal{M}_{red} := \mu^{-1}(S)/(\mathfrak{l} \cap \mathfrak{g}^{V_\Gamma})$$

is a Poisson manifold (provided it is a manifold).

It is not difficult to describe the bivector field on $\mu^{-1}(S)/(\mathfrak{l} \cap \mathfrak{g}^{V_\Gamma})$. First, let $\mathfrak{g}_\Delta \subseteq \mathfrak{d}$ denote the diagonal and $\mathfrak{g}_{\bar{\Delta}} \subseteq \mathfrak{d}$ the off-diagonal:

$$\mathfrak{g}_{\bar{\Delta}} := \{(\xi, -\xi) \in (\mathfrak{g} \oplus \bar{\mathfrak{g}})\}.$$

Each Lagrangian Lie subalgebra $\mathfrak{l}_v \subseteq \mathfrak{d}^{V_\Gamma}$ defines an element $\tau_v \in \wedge^2(\mathfrak{h}/(\mathfrak{l} \cap \mathfrak{h}))$ by the equation

$$\mathfrak{l}_v = \{(\alpha + \tau_v^\sharp \alpha) \mid \alpha \in \mathfrak{g}_{\bar{\Delta}}, \langle \alpha, \mathfrak{l}_v \cap \mathfrak{g}_\Delta \rangle = 0\} + \mathfrak{l}_v \cap \mathfrak{g}_\Delta.$$

Here $\tau_v^\sharp \alpha \in \mathfrak{g}_\Delta/(\mathfrak{l}_v \cap \mathfrak{g}_\Delta)$ is defined by

$$\tau_v^\sharp \alpha := \frac{1}{2} \sum_{i,j} \tau_v^{ij} \langle \alpha, \xi_i \rangle \xi_j - \frac{1}{2} \sum_{i,j} \tau_v^{ij} \langle \alpha, \xi_j \rangle \xi_i$$

when we represent τ_v as $\tau_v = \frac{1}{2} \sum_{i,j} \tau_v^{ij} \xi_i \wedge \xi_j$.

We have the following theorem:

Theorem 1.4. *If the intersection, $\mu^{-1}(S)$, of $S \times \mathcal{M}_{\Sigma,V}(G)$ with the graph of μ is clean, and the $\mathfrak{l} \cap \mathfrak{g}^{V_\Gamma}$ -orbits of $\mu^{-1}(S)$ form a regular foliation, then the bivector field*

$$\pi + \sum_{v \in V_\Gamma} \rho_v(\tau_v) \in \Gamma(\wedge^2 T(\mu^{-1}(S))/\rho(\mathfrak{l} \cap \mathfrak{g}^{V_\Gamma}))$$

is $\mathfrak{l} \cap \mathfrak{g}^{V_\Gamma}$ invariant and descends to define the Poisson structure on $\mu^{-1}(S)/(\mathfrak{l} \cap \mathfrak{g}^{V_\Gamma})$. Here $\rho : \mathfrak{g}^{V_\Gamma} \rightarrow \mathfrak{X}(\mathcal{M}_{\Sigma,V}(G))$ denotes the action by infinitesimal gauge transformations at the marked points, and ρ_v is the restriction of ρ to the $v \in V_\Gamma$ -th factor.

Moreover, for any \mathfrak{l} -orbit $O \subseteq S$, the image of $\mu^{-1}(O)$ in $\mu^{-1}(S)/(\mathfrak{l} \cap \mathfrak{g}^{V_\Gamma})$ will be a symplectic leaf.

Proof. This will follow from Theorem 3.3, while the statement for the symplectic leaves will follow from Theorem 3.2. \square

Remark 1.6. As in Section 1.4, one may also sew domains together to obtain Poisson structures on the moduli spaces of branched surfaces. The general reduction statement is Theorem 3.3.

Example 1.10 (Double Poisson Lie group [26]). Suppose that $\mathfrak{g} = \mathfrak{e} \oplus \mathfrak{f}$ as a vector space, where $\mathfrak{e}, \mathfrak{f} \subseteq \mathfrak{g}$ are Lagrangian Lie subalgebras, and that $\mathfrak{e}, \mathfrak{f} \subseteq \mathfrak{g}$ integrate to Lie subgroups $E, F \subseteq G$ such that $E \cap F = 1$. Let Σ be a disk with two marked points labelled as in Fig. 14. We colour the edges with the full group G ,

$$S_{e_1} = G, \quad S_{e_2} = G,$$

and the vertices as

$$\mathfrak{l}_{v_1} = \mathfrak{e} \oplus \mathfrak{f}, \quad \mathfrak{l}_{v_2} = \mathfrak{f} \oplus \mathfrak{e}.$$

Therefore,

$$\mu^{-1}(S) = \mathcal{M}_{P_2} = \{(g_1, g_2) \in G \times G \mid g_1 g_2 = 1\}.$$

Meanwhile the residual gauge transformations,

$$\bigoplus_{v \in V_\Gamma} \mathfrak{l}_v \cap \mathfrak{g}_\Delta = 0$$

(since $\mathfrak{e} \cap \mathfrak{f} = 0$). Thus we may identify the moduli space $\mathcal{M}_{red} \cong G$, via the map $(g_1, g_2) \rightarrow g_1$.




FIGURE 14. The double Poisson Lie group. The holonomies $g_i \in G$ satisfy $g_1 g_2 = 1$.




FIGURE 15. The Poisson Lie group. The holonomies $g \in G$ and $e \in E$ satisfy $eg = 1$.

Next, we compute the bivector field, $\pi = \pi_{P_2} + \rho_{v_1}(\tau_{v_1}) + \rho_{v_2}(\tau_{v_2})$. Now, as explained in Example 1.9, $\pi_{P_2} = 0$ so only the term $\sum_i \rho_{v_i}(\tau_{v_i})$ contributes. Now,

$$\tau_{v_1} = \frac{1}{2} \sum_i (\zeta^i, \zeta^i) \wedge (\eta_i, \eta_i), \quad \tau_{v_2} = \frac{1}{2} \sum_i (\eta_i, \eta_i) \wedge (\zeta_i, \zeta_i),$$

where $\{\eta_i\} \subset \mathfrak{e}$ and $\{\zeta^i\} \subset \mathfrak{f}$ are basis in duality. Therefore,

$$\pi = \frac{1}{2} \sum_i (\zeta^i)^L \wedge (\eta_i)^L + (\eta_i)^R \wedge (\zeta^i)^R.$$

In fact, π defines the Poisson Lie group structure on G corresponding the double Lie bialgebra structure on \mathfrak{g} resulting from the Manin triple $(\mathfrak{g}, \mathfrak{e}, \mathfrak{f})$ [26, 28]. The symplectic leaves are computed as the restriction of the \mathfrak{l} -orbits, which in this case can be seen to correspond to the orbits of the dressing action on G .⁵

Example 1.11 (Poisson Lie group [26]). Suppose the Lie groups G, E, F and their Lie algebras are as in Example 1.10, and let Σ be a disk with two marked points, as in Fig. 15. We colour the vertices as in Example 1.10, but we colour the edges as

$$S_{e_1} = E, \quad S_{e_2} = G.$$

Therefore,

$$\mu^{-1}(S) = \{(e, g) \in E \times G \mid eg = 1\},$$

while the residual gauge transformations are trivial, as before. Thus we may identify the moduli space $\mathcal{M}_{red} \cong E$, via the map $(e, g) \rightarrow e$.

The bivector field, π , on E is computed to be the restriction of the bivector field

$$\frac{1}{2} \sum_i (\zeta^i)^L \wedge (\eta_i)^L + (\eta_i)^R \wedge (\zeta^i)^R.$$

on G to $E \subseteq G$. Thus, π defines the Poisson Lie group structure on E corresponding the Manin triple $(\mathfrak{g}, \mathfrak{e}, \mathfrak{f})$ [26, 28]. As before, the symplectic leaves are computed as the restriction of the \mathfrak{l} -orbits. Once again, they are precisely the orbits of the dressing action on E .

⁵In fact computing the symplectic leaves via Theorem 1.4 is precisely the computation found in [23].




FIGURE 16. The Poisson homogeneous space corresponding to the Lagrangian Lie subalgebra $\mathfrak{h} \subseteq \mathfrak{g}$. The holonomies $g \in G$ and $e \in E$ satisfy $eg = 1$.

Example 1.12 (Poisson homogenous spaces). Suppose the Lie groups G, E, F and their Lie algebras are as in Example 1.10, and let Σ be a disk with two marked points, as in Fig. 16. Suppose further that $\mathfrak{h} \subseteq \mathfrak{g}$ is a Lagrangian subalgebra such that $\mathfrak{k} := \mathfrak{h} \cap \mathfrak{e}$ integrates to a closed Lie subgroup $K \subseteq E$. We colour the edges as in Example 1.11, but we colour the vertices as

$$\mathfrak{l}_{v_1} = \mathfrak{e} \oplus \mathfrak{h}, \quad \mathfrak{l}_{v_2} = \mathfrak{f} \oplus \mathfrak{e}.$$

Therefore,

$$\mu^{-1}(S) = \{(e, g) \in E \times G \mid eg = 1\},$$

while the residual gauge transformations are $G \times K$ acting as

$$(g, k) : (e, g) \rightarrow (ek^{-1}, kg).$$

We may identify the moduli space $\mathcal{M}_{red} \cong E/K$, via the map $(e, g) \rightarrow [e]$. The Poisson structure on \mathcal{M}_{red} is the Poisson homogenous structure corresponding to the Lagrangian Lie subalgebra $\mathfrak{h} \subseteq \mathfrak{g}$ in Drinfel'd's classification [16]. We leave it to the reader to compute the bivector field and symplectic leaves on E/K via Theorem 1.4.

Example 1.13 (Affine Poisson structure on G [28]). Generalizing the setup found in Example 1.10, we suppose that $\mathfrak{h} \subseteq \mathfrak{g}$ is a Lagrangian subalgebra which is also complementary (as a vector space) to $\mathfrak{e} \subseteq \mathfrak{g}$. As in Example 1.10, we colour the edges with the full group G , but the vertices as

$$\mathfrak{l}_{v_1} = \mathfrak{e} \oplus \mathfrak{h}, \quad \mathfrak{l}_{v_2} = \mathfrak{f} \oplus \mathfrak{e},$$

(cf. Fig. 17). As before, we have $\mu^{-1}(S) = \mathcal{M}_{P_2}$, and the residual gauge transformations are trivial. Thus, we may identify the moduli space $\mathcal{M}_{red} \cong G$, via the map $(g_1, g_2) \rightarrow g_1$.

Meanwhile the bivector field, π , is

$$\pi = \frac{1}{2} \sum_i (\zeta_{\mathfrak{f}}^i)^L \wedge (\eta_i)^L + (\eta_i)^R \wedge (\zeta_{\mathfrak{h}}^i)^R,$$

where the bases $\{\zeta_{\mathfrak{f}}^i\} \subseteq \mathfrak{f}$ and $\{\zeta_{\mathfrak{h}}^i\} \subseteq \mathfrak{h}$ are both dual to $\{\eta_i\} \subseteq \mathfrak{g}$. In fact, π defines the affine Poisson structure on G corresponding to the Manin triples $(\mathfrak{g}, \mathfrak{e}, \mathfrak{f})$ and $(\mathfrak{g}, \mathfrak{e}, \mathfrak{h})$ (as described by Lu [28]). The symplectic leaves are computed as the restriction of the \mathfrak{l} -orbits.




FIGURE 17. The affine Poisson structure on G . The holonomies $g_i \in G$ satisfy $g_1 g_2 = 1$.




FIGURE 18. Lu-Yakimov Poisson homogenous spaces. The holonomies $g_i \in G$ satisfy $g_1 g_2 = 1$.

Example 1.14 (Lu Yakimov Poisson homogenous spaces [26]). Suppose the Lie groups G, E, F and their Lie algebras are as in Example 1.10, and that $C \subseteq G$ is a closed Lie subgroup whose Lie algebra $\mathfrak{c} \subseteq \mathfrak{g}$ is coisotropic. Let Σ be a disk with two marked points and edges and vertices labelled as in Fig. 18. We colour the edges with the full group G ,

$$S_{e_1} = G, \quad S_{e_2} = G,$$

and the vertices as

$$\mathfrak{l}_{v_1} = \mathfrak{l}_{\mathfrak{c}} = \{(\xi, \xi') \in \mathfrak{c} \oplus \mathfrak{c} \mid \xi - \xi' \in \mathfrak{c}^\perp\}, \quad \mathfrak{l}_{v_2} = \mathfrak{f} \oplus \mathfrak{e}$$

(cf. Fig. 18). Therefore,

$$\mu^{-1}(S) = \mathcal{M}_{P_2} = \{(g_1, g_2) \in G \times G \mid g_1 g_2 = 1\}.$$

Meanwhile the residual gauge transformations,

$$\bigoplus_{v \in V_\Gamma} \mathfrak{l}_v \cap \mathfrak{g}_\Delta = \mathfrak{c}_{v_0},$$

where $\mathfrak{c}_{v_0} = \{(\xi, \xi) \in \mathfrak{c} \oplus \mathfrak{c}\} \subseteq \mathfrak{l}_{v_0}$. Thus, up to a gauge transformation, $(g_1, g_2) \sim (g_1 c^{-1}, c g_2)$ (for any $c \in C$), and we may identify the moduli space $\mathcal{M}_{red} \cong G/C$, via the map $(g_1, g_2) \rightarrow [g_1]$.

The bivector field, π , on G/C can be computed to be the projection of the bivector field

$$\frac{1}{2} \sum_i (\eta_i)^R \wedge (\zeta^i)^R.$$

on G to G/C . Thus, π defines the Lu-Yakimov Poisson structure on G/C corresponding the Manin triple $(\mathfrak{g}, \mathfrak{e}, \mathfrak{f})$ [26, 30].

2. BACKGROUND

2.1. Courant algebroids. Courant algebroids and Dirac structures are the basic tools in the theory of generalized moment maps.

Definition 2.1 ([27]). A *Courant algebroid* is a vector bundle $\mathbb{E} \rightarrow M$ endowed with a non-degenerate symmetric bilinear form $\langle \cdot, \cdot \rangle$ on the fibres, a bundle map $\mathbf{a} : \mathbb{E} \rightarrow TM$ called the *anchor* and a bracket $[\![\cdot, \cdot]\!] : \Gamma(\mathbb{E}) \times \Gamma(\mathbb{E}) \rightarrow \Gamma(\mathbb{E})$ called the *Courant bracket* satisfying the following axioms for sections $e_1, e_2, e_3 \in \Gamma(\mathbb{E})$ and functions $f \in C^\infty(M)$:

- c1) $[\![e_1, [e_2, e_3]]\!] = [[e_1, e_2], e_3] + [e_2, [e_1, e_3]],$
- c2) $\mathbf{a}(e_1)\langle e_2, e_3 \rangle = \langle [e_1, e_2], e_3 \rangle + \langle e_2, [e_1, e_3] \rangle,$
- c3) $[\![e_1, e_2]\!] + [e_2, e_1] = \mathbf{a}^* d\langle e_1, e_2 \rangle.$

Here $\mathbf{a}^* : T^*M \rightarrow \mathbb{E}^* \cong \mathbb{E}$ is the map dual to the anchor.

A subbundle $E \subseteq \mathbb{E}|_S$ along a submanifold $S \subseteq M$ is called a *Dirac structure with support on S* if

$$e_1|_S, e_2|_S \in \Gamma(E) \Rightarrow [\![e_1, e_2]\!]|_S \in \Gamma(E),$$

(it is *involutive*) and $E^\perp = E$ (it is *Lagrangian*). If $S = M$, then E is simply called a *Dirac structure*.

Remark 2.1. As shown in [31, 39], one may also derive the following useful identities from the Courant axioms:

- c4) $[\![e_1, f e_2]\!] = f [e_1, e_2] + (\mathbf{a}(e_1)f) e_2$
- c5) $[\![f e_1, e_2]\!] = f [e_1, e_2] - (\mathbf{a}(e_2)f) e_1 + \langle e_1, e_2 \rangle \mathbf{a}^* df$
- c6) $\mathbf{a}[\![e_1, e_2]\!] = [\mathbf{a}(e_1), \mathbf{a}(e_2)]$

For any Courant algebroid \mathbb{E} , we denote by $\overline{\mathbb{E}}$ the Courant algebroid with the same bracket and anchor, but with the metric negated.⁶

Example 2.1. A Courant algebroid over a point is a quadratic Lie algebra. Dirac structures are Lagrangian Lie subalgebras.

Example 2.2 (Standard Courant algebroid [12, 13]). The vector bundle $\mathbb{T}M := TM \oplus T^*M$ is a Courant algebroid with metric

$$\langle v_1 + \mu_1, v_2 + \mu_2 \rangle = \mu_1(v_2) + \mu_2(v_1), \quad v_1, v_2 \in TM, \quad \mu_1, \mu_2 \in T^*M$$

and bracket

$$[X + \alpha, Y + \beta] = [X, Y] + \mathcal{L}_X \beta - \iota_Y d\alpha, \quad X, Y \in \mathfrak{X}(M), \alpha, \beta \in \Omega^1(M).$$

⁶Note that this also negates the map $\mathbf{a}^* : T^*M \rightarrow \mathbb{E}^* \cong \mathbb{E}$, so axiom c3) still holds.

The standard Courant algebroid is an example of an important class of Courant algebroids called *exact Courant algebroids*.

Definition 2.2 (Exact Courant algebroids [33, 37]). A Courant algebroid $\mathbb{E} \rightarrow M$ is called *exact* if the sequence

$$(2.1) \quad 0 \rightarrow T^*M \xrightarrow{\mathbf{a}^*} \mathbb{E} \xrightarrow{\mathbf{a}} TM \rightarrow 0$$

is exact.

If a Dirac structure $E \subseteq \mathbb{E}$ is supported on $S \subseteq M$, then $\mathbf{a}(E) \subseteq TS$, and $\mathbf{a}^*(\text{ann}(TS)) \subseteq E$.

Definition 2.3 (Exact Dirac structures). Suppose $\mathbb{E} \rightarrow M$ is a Courant algebroid (not necessarily exact), and $E \subseteq \mathbb{E}$ is a Dirac structure with support on $S \subseteq M$. We say that E is an *exact Dirac structure* if the sequence

$$(2.2) \quad 0 \rightarrow \text{ann}(TS) \xrightarrow{\mathbf{a}^*} E \xrightarrow{\mathbf{a}} TS \rightarrow 0$$

is exact.

Lemma 2.1. Suppose $\mathbb{E} \rightarrow M$ is a Courant algebroid, and $E \subseteq \mathbb{E}$ is a Dirac structure with support on $S \subseteq M$. Then the following two statements are equivalent:

- (1) E is an exact Dirac structure.
- (2) The Courant algebroid \mathbb{E} is exact along S (that is, the sequence Eq. (2.1) is exact at every $x \in S$), and $\mathbf{a} : E \rightarrow TS$ is surjective.

Proof. Let $x \in S$, and consider the commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \text{ann}(T_x S) & \xrightarrow{\mathbf{a}^*} & E_x & \xrightarrow{\mathbf{a}} & T_x S & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & T_x^* M & \xrightarrow{\mathbf{a}^*} & \mathbb{E}_x & \xrightarrow{\mathbf{a}} & T_x M & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & T_x^* M / \text{ann}(T_x S) & \xrightarrow{\mathbf{a}^*} & E_x^* & \xrightarrow{\mathbf{a}} & T_x M / T_x S & \longrightarrow & 0 \end{array}$$

Note that the vertical sequences are exact.

Suppose that E is an exact Dirac structure. Then the top horizontal sequence is exact, by assumption. The lower horizontal sequence is dual to the top sequence, and hence also exact. The five lemma then implies that all terms in the long exact sequence vanish. In particular, the central horizontal sequence is exact.

Conversely, suppose that the Courant algebroid \mathbb{E} is exact along S and $\mathbf{a} : E_x \rightarrow T_x S$ is surjective (and hence $T_x^* M / \text{ann}(T_x S) \rightarrow E_x^*$ is injective). Once again, the five lemma implies that all terms in the long exact sequence vanish. We conclude that E is an exact Dirac structure. \square

Example 2.3 (Action Courant algebroids [23]). Suppose \mathfrak{d} is a Lie algebra equipped with an invariant metric. Given a Lie algebra action $\rho: \mathfrak{d} \rightarrow \mathfrak{X}(M)$ on a manifold M , let $\mathbb{E} = \mathfrak{d} \times M$ with anchor map $\mathbf{a}(\xi, m) = \rho(\xi)_m$, and with the bundle metric coming from the metric on \mathfrak{d} . As shown in [23], the Lie bracket on constant sections $\mathfrak{d} \subseteq C^\infty(M, \mathfrak{d}) = \Gamma(\mathbb{E})$ extends to a Courant bracket *if and only if* the stabilizers $\mathfrak{d}_m \subseteq \mathfrak{d}$ are coisotropic, i.e. $\mathfrak{d}_m \supseteq \mathfrak{d}_m^\perp$. Explicitly, for $\xi_1, \xi_2 \in \Gamma(\mathbb{E}) = C^\infty(M, \mathfrak{d})$ the Courant bracket reads (see [23, § 4])

$$(2.3) \quad \llbracket \xi_1, \xi_2 \rrbracket = [\xi_1, \xi_2] + \mathcal{L}_{\rho(\xi_1)}\xi_2 - \mathcal{L}_{\rho(\xi_2)}\xi_1 + \rho^* \langle d\xi_1, \xi_2 \rangle.$$

Here $\rho^*: T^*M \rightarrow \mathfrak{d} \times M$ is the dual map to the action map $\rho: \mathfrak{d} \times M \rightarrow TM$, using the metric to identify $\mathfrak{d}^* \cong \mathfrak{d}$. We refer to $\mathfrak{d} \times M$ with bracket (2.3) as an *action Courant algebroid*.

Example 2.4 (Cartan Courant algebroid [37]). Suppose \mathfrak{g} is a Lie group endowed with an invariant metric, $\langle \cdot, \cdot \rangle$. We let $\bar{\mathfrak{g}}$ denote the Lie algebra \mathfrak{g} with the metric negated, $-\langle \cdot, \cdot \rangle$. Suppose G is a Lie group with Lie algebra \mathfrak{g} and which preserves the metric. The Lie algebra $\bar{\mathfrak{g}} \oplus \mathfrak{g}$ acts on G by $\rho: (\bar{\mathfrak{g}} \oplus \mathfrak{g}) \times G \rightarrow TG$,

$$\rho(\xi, \eta) = -\xi^R + \eta^L, \quad \xi, \eta \in \mathfrak{g},$$

where $\xi^L, \xi^R \in \mathfrak{X}(G)$ denotes the left/right-invariant vector field on G which is equal to $\xi \in \mathfrak{g}$ at the identity element.

The stabilizer at the identity element is the diagonal subalgebra, $\mathfrak{g}_\Delta \subseteq \bar{\mathfrak{g}} \oplus \mathfrak{g}$ which is Lagrangian. Now ρ is equivariant with respect to the G -action on $(\bar{\mathfrak{g}} \oplus \mathfrak{g}) \times G$ given by

$$g' : (\xi, \eta; g) \rightarrow (\mathbf{Ad}_{g'}\xi, \eta, g' \cdot g), \quad g' \in G, (\xi, \eta; g) \in (\bar{\mathfrak{g}} \oplus \mathfrak{g}) \times G,$$

and the left action of G on TG . Since this action is transitive on the base of the vector bundles and G preserves the metric on \mathfrak{g} , it follows that all stabilizers are Lagrangian. Thus $(\bar{\mathfrak{g}} \oplus \mathfrak{g}) \times G$ is an action Courant algebroid, called the *Cartan Courant algebroid*.

The diagonal subalgebra $\mathfrak{g}_\Delta \subseteq (\bar{\mathfrak{g}} \oplus \mathfrak{g})$ defines a Dirac structure

$$\mathfrak{g}_\Delta \times G \subseteq (\bar{\mathfrak{g}} \oplus \mathfrak{g}) \times G$$

called the *Cartan Dirac structure*.

Remark 2.2. The Cartan Courant algebroid was first introduced in [37], and later simplified to the above description in [1]. The Cartan Dirac structure was discovered independently by Alekseev, Ševera and Strobl [3, 22, 37]. The description given above was found in [1]


The Dirac structure of central focus in this paper is the following generalization of the Cartan Dirac structure.

Example 2.5 (Γ -twisted Cartan-Dirac structure). Suppose Γ is a permutation graph, with edge set E_Γ , vertex set V_Γ , and (bijective) incidence maps


$$\text{in}, \text{out} : E_\Gamma \rightarrow V_\Gamma.$$

The diagonal subalgebra $\mathfrak{g}_\Delta^{V_\Gamma} \subseteq (\bar{\mathfrak{g}} \oplus \mathfrak{g})^{V_\Gamma}$ is Lagrangian, and hence so is its image

$$\mathfrak{g}_\Gamma := (\text{in} \oplus \text{out})^! (\mathfrak{g}_\Delta^{V_\Gamma}) \subseteq (\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_\Gamma}$$



(A) The Γ -twisted Cartan Courant algebroid. Here $\text{in}(e) = v = \text{out}(e')$, where $v \in V_\Gamma$ is the vertex and $e, e' \in E_\Gamma$ are edges.



(B) The element $\xi^{(\cdot)} \in \mathfrak{g}_\Delta^{V_\Gamma} \cong \mathfrak{g}_\Gamma$ of the Γ -twisted Cartan Dirac structure acts diagonally by ξ^v at the vertex v .

under the isomorphism $(\text{in} \oplus \text{out})^! : (\bar{\mathfrak{g}} \oplus \mathfrak{g})^{V_\Gamma} \rightarrow (\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_\Gamma}$.

Thus

$$\mathfrak{g}_\Gamma \times G^{E_\Gamma} \subseteq ((\bar{\mathfrak{g}} \oplus \mathfrak{g}) \times G)^{E_\Gamma}$$

is a Dirac structure, called the Γ -twisted Cartan Dirac structure.

The following picture can be helpful. We associate a copy of the Courant algebroid $(\bar{\mathfrak{g}} \oplus \mathfrak{g}) \times G$ to each edge, as in Fig. 19a. The Dirac structure \mathfrak{g}_Γ acts diagonally at each vertex, as pictured in Fig. 19b.

Remark 2.3. Γ -twisted Cartan-Dirac structures were discovered independently by Alejandro Cabrera, who provides their construction in terms of Dirac reduction of the Lie-Poisson structure on the dual of the loop Lie algebra, $\Omega^1(S^1, \mathfrak{g})$.

Remark 2.4. The following was explained to the authors by Eckhard Meinrenken. Suppose $\sigma_\Gamma : V_\Gamma \rightarrow V_\Gamma$ is the permutation induced by Γ (by the discrete flow along the edges of Γ), and consider the group $G_{\text{big}} := G^{V_\Gamma} \rtimes \mathbb{Z}$,

$$(g, i) \cdot (g', i') = (g(\sigma_\Gamma^i)^!(g'), i + i'),$$

where $((\sigma_\Gamma^i)^!(g'))_v = (g')_{\sigma_\Gamma^i(v)}$ for any $v \in V_\Gamma$. Consider the embedding of manifolds

$$((\text{in}^{-1})^!, -1) : G^{E_\Gamma} \rightarrow G^{V_\Gamma} \rtimes \mathbb{Z} = G_{\text{big}}.$$

The Lie algebra of G_{big} is \mathfrak{g}^{V_Γ} , and for every $\xi \in \mathfrak{g}^{V_\Gamma}$, the left and right invariant vector fields $\xi^L, \xi^R \in \mathfrak{X}(G_{\text{big}})$ restrict to the $e \in E_\Gamma$ 'th factor of $((\text{in}^{-1})^!, -1)(G^{E_\Gamma})$ as $\xi_{\text{in}(e)}^R$ and $\xi_{\text{out}(e)}^L$, (respectively). Thus, the Γ -twisted Cartan-Dirac structure on G^{E_Γ} may be canonically identified with the restriction of the Cartan-Dirac structure on G_{big} to $((\text{in}^{-1})^!, -1)(G^{E_\Gamma})$.

2.2. Courant relations and morphisms of Manin pairs.

2.2.1. Relations. A smooth relation $S : M_1 \dashrightarrow M_2$ between manifolds is an immersed submanifold $S \subseteq M_2 \times M_1$. We will write $m_1 \sim_S m_2$ if $(m_2, m_1) \in S$. Given smooth relations $S : M_1 \dashrightarrow M_2$ and $S' : M_2 \dashrightarrow M_3$, the set-theoretic composition $S' \circ S$ is the image of

$$(2.4) \quad S' \diamond S = (S' \times S) \cap (M_3 \times (M_2)_\Delta \times M_1)$$

under projection to $M_3 \times M_1$, where $(M_2)_\Delta \subseteq M_2 \times M_2$ denotes the diagonal.

We say that the two relations *compose cleanly* if (2.4) is a clean intersection in the sense of Bott (i.e. it is smooth, and the intersection of the tangent bundles is the tangent bundle of the intersection), and the map from $S' \diamond S$ to $M_2 \times M_1$ has constant rank. In this case, the composition $S' \circ S : M_1 \dashrightarrow M_3$ is a well-defined smooth relation. See [24, Appendix A] for more information on the composition of smooth relations. For background on clean intersections of manifolds, see e.g. [18, page 490].

For any relation $S : M_1 \dashrightarrow M_2$, we let $S^\top : M_2 \dashrightarrow M_1$ denote the *transpose* relation,

$$S^\top = \{(m_1, m_2) \in M_1 \times M_2 \mid (m_2, m_1) \in S\}.$$

2.2.2. Courant relations. As popularized by the second author [33, 35], Dirac structures can be interpreted as the ‘canonical relations’ between Courant algebroids:

Definition 2.4 (Courant relations and morphisms [5, 11, 35]). Suppose $\mathbb{E}_1 \rightarrow M_1$ and $\mathbb{E}_2 \rightarrow M_2$ are two Courant algebroids. A relation

$$R : \mathbb{E}_1 \dashrightarrow \mathbb{E}_2$$

is called a *Courant relation* if $R \subseteq \mathbb{E}_2 \times \overline{\mathbb{E}_1}$ is a Dirac structure supported on a submanifold $S \subseteq M_2 \times M_1$. A Courant relation is called *exact* if the underlying Dirac structure is exact.

When $S = \text{gr}(\mu)$ is the graph of a smooth map $\mu : M_1 \rightarrow M_2$, R is called a *Courant morphism*.

We define the range $\text{ran}(R) \subseteq \mathbb{E}_2|_S$ and the kernel $\ker(R) \subseteq \mathbb{E}_2|_S$ of R by

$$\begin{aligned} \text{ran}(R) &:= \{e \in \mathbb{E}_2|_S \mid e' \sim_R e \text{ for some } e' \in \mathbb{E}_1\} \\ \ker(R) &:= \{e \in \mathbb{E}_1|_S \mid e \sim_R 0\}. \end{aligned}$$

As an example, any Dirac structure $E \subseteq \mathbb{E}$ defines a Courant morphism

$$E : \mathbb{E} \dashrightarrow *$$

to the trivial Courant algebroid (or a Courant relation from the trivial Courant algebroid). Similarly, the diagonal $\mathbb{E}_\Delta \subseteq \mathbb{E} \times \overline{\mathbb{E}}$ defines the Courant morphism

$$\mathbb{E}_\Delta : \mathbb{E} \dashrightarrow \mathbb{E}$$

corresponding to the identity map.

The key property of Courant relations is the ability to compose them:

Proposition 2.1 ([24, Proposition 1.4]). *Suppose $R : \mathbb{E}_1 \dashrightarrow \mathbb{E}_2$ and $R' : \mathbb{E}_2 \dashrightarrow \mathbb{E}_3$ are two Courant relations which compose cleanly, then their composition,*

$$R' \circ R : \mathbb{E}_1 \dashrightarrow \mathbb{E}_3,$$

is a Courant relation.

Example 2.6 (Standard lift). Suppose $S : M_1 \rightarrow M_2$ is a relation, then the *standard lift* of S ,

$$R_S := TS \oplus \text{ann}(TS) \subseteq \mathbb{T}M_2 \times \overline{\mathbb{T}M_1},$$

defines a Courant relation

$$R_S : \mathbb{T}M_1 \dashrightarrow \mathbb{T}M_2.$$

Example 2.7 (Coisotropic subalgebras). Suppose \mathfrak{d} is a Lie algebra equipped with an invariant metric. A subalgebra $\mathfrak{c} \subseteq \mathfrak{d}$ is said to be *coisotropic* if $\mathfrak{c}^\perp \subseteq \mathfrak{c}$. In this case, $\mathfrak{c}^\perp \subseteq \mathfrak{c}$ is an ideal, and the metric on \mathfrak{d} descends to define a metric on

$$\mathfrak{d}_{\mathfrak{c}} := \mathfrak{c}/\mathfrak{c}^\perp.$$

The natural relation

$$R_{\mathfrak{c}} : \mathfrak{d} \dashrightarrow \mathfrak{d}_{\mathfrak{c}}, \quad \xi \sim_{R_{\mathfrak{c}}} \xi + \mathfrak{c}^\perp \quad \text{for } \xi \in \mathfrak{c},$$

is a Courant relation, where $(\xi + \mathfrak{c}^\perp) \in \mathfrak{c}/\mathfrak{c}^\perp$ denotes the equivalence class of $\xi \in \mathfrak{c}$.

For any Lagrangian subalgebra $\mathfrak{h} \subseteq \mathfrak{d}$, Proposition 2.1 implies that

$$\mathfrak{h}_{\mathfrak{c}} := R_{\mathfrak{c}} \circ \mathfrak{h} = (\mathfrak{h} \cap \mathfrak{c})/(\mathfrak{h} \cap \mathfrak{c}^\perp)$$

is a Lagrangian subalgebra of $\mathfrak{d}_{\mathfrak{c}}$.

2.2.3. Morphisms of Manin pairs. A pair (\mathbb{E}, E) consisting of a Courant algebroid, \mathbb{E} , together with a Dirac structure $E \subseteq \mathbb{E}$ is known as a *Manin pair* [2, 11].

Definition 2.5 ([11]). Suppose $\mathbb{E}_1 \rightarrow M_1$ and $\mathbb{E}_2 \rightarrow M_2$ are two Courant algebroids. A Courant morphism

$$R : \mathbb{E}_1 \dashrightarrow \mathbb{E}_2,$$

supported on the graph of a map $\mu : M_1 \rightarrow M_2$, defines a *morphism of Manin pairs*,

$$(2.5) \quad R : (\mathbb{E}_1, E_1) \dashrightarrow (\mathbb{E}_2, E_2)$$

if

- m1) $R \circ E_1 \subseteq E_2$, and
- m2) $\ker(R) \cap E_1 = 0$

Here $\ker(R) := (0 \times \overline{\mathbb{E}_1}) \cap R$.

The morphism of Manin pairs, (2.5), is said to be *exact* if the underlying Dirac structure is exact.

Suppose

$$(2.6) \quad R' : (\mathbb{E}_2, E_2) \dashrightarrow (\mathbb{E}_3, E_3)$$

is a second morphism of Manin pairs. Conditions (m1) and (m2) imply that the composition of relations $R' \circ R$ is clean. Moreover, the composition defines a morphism of Manin pairs

$$(2.7) \quad R' \circ R : (\mathbb{E}_1, E_1) \dashrightarrow (\mathbb{E}_3, E_3),$$

(cf. [11]).

Remark 2.5. In [25], a morphism of Manin pairs, (2.5) was said to be *full* if the map

$$\mathbf{a}|_R : R \rightarrow T\text{gr}(\mu)$$

was a surjection. The concept of exact morphisms of Manin pairs is a stronger, but more natural, condition.

If (2.5) is a morphism of Manin pairs, then there exists map $\rho_R : \mu^* E_2 \rightarrow E_1$ uniquely determined by the condition

$$(2.8) \quad \rho_R(e) \sim_R e, \quad e \in E_2.$$

The induced map of section $\rho_R : \mathbf{\Gamma}(E_2) \rightarrow \mathbf{\Gamma}(E_1)$ is a morphism of Lie algebras. Thus (2.5) defines an action of E_2 on M_1 which factors through the action of E_1 .

2.3. Quasi-Hamiltonian manifolds. If (\mathbb{E}, E) is a Manin pair, a *quasi-Hamiltonian* (\mathbb{E}, E) -manifold in the sense of [11] is a manifold M together with a morphism of Manin pairs

$$(\mathbb{T}M, TM) \dashrightarrow (\mathbb{E}, E).$$

We shall say that the quasi-Hamiltonian space is *exact* (or *quasi-symplectic*) if the morphism of Manin pairs is exact. To simplify notation (when M is a complicated expression), we will often denote the Manin pair on the left as $(\mathbb{T}M, TM) = (\mathbb{T}, T)M$.

Example 2.8 (Poisson and symplectic structures). Let 0 denote the trivial Courant algebroid over a point. Consider a morphism of Manin pairs

$$(2.9) \quad R : (\mathbb{T}, T)M \dashrightarrow (0, 0).$$

In this case, $R \subseteq \mathbb{T}M$ is just a Dirac structure with support on all of M . Condition (m1) is vacuous, while condition (m2) is equivalent to $R \cap TM = 0$. As explained in [12, 13], it follows that

$$R = \text{gr}(\pi^\sharp) := \{(\pi(\alpha, \cdot) + \alpha) \mid \alpha \in \Omega^1(M)\} \subseteq \mathbb{T}M$$

is the graph of a Poisson bivector field $\pi \in \mathfrak{X}^2(M)$. In this way, there is a one-to-one correspondence between morphisms of Manin pairs of the form (2.9) and Poisson structures on M [11].

Equation (2.9) is an exact morphism of Manin pairs if $\mathbf{a}|_R : \text{gr}(\pi^\sharp) \rightarrow TM$ is a surjection. Equivalently,

$$\pi^\sharp : T^*M \rightarrow TM$$

is an isomorphism, or the Poisson structure on M is symplectic. In this way, there is a one-to-one correspondence between exact morphisms of Manin pairs of the form (2.9) and symplectic structures on M [25].

The Poisson and symplectic structures that appear in this paper will all arise in this way.

Example 2.9 (E -invariant submanifolds). Let (\mathbb{E}, E) be a Manin pair over a manifold N , and suppose $M \subseteq N$ is an E -invariant submanifold, i.e. $\mathbf{a}(E|_M) \subseteq TM$. Then

$$\mathbf{a}^*|_M : T^*N|_M \rightarrow \mathbb{E}$$

descends to a map

$$\mathbf{a}^*|_M : T^*M \cong T^*N/\text{ann}(TM) \rightarrow \mathbb{E}.$$

As in [24, Example 1.6], define the Courant relation $R_{E,M} : \mathbb{T}M \dashrightarrow \mathbb{E}$ by

$$\mathbf{a}(e) + i^* \alpha \sim_{R_{E,M}} e + \mathbf{a}^* \alpha, \quad e \in E, \quad \alpha \in T^*N,$$

where $i : M \rightarrow N$ denotes the inclusion. Then

$$(2.10) \quad R_{E,M} : (\mathbb{T}, T)M \dashrightarrow (\mathbb{E}, E)$$

is a morphism of Manin pairs.

Moreover, (2.10) is an *exact* morphism of Manin pairs if and only if $\mathbf{a}(E|_M) = TM$ and the Courant algebroid $\mathbb{E} \rightarrow N$ is exact along M .

Remark 2.6. In fact, Eq. (2.10) is the unique morphism of Manin pairs supported on $\mathbf{gr}(i)$. To see why, suppose

$$R : (\mathbb{T}, T)M \dashrightarrow (\mathbb{E}, E)$$

is such a morphism of Manin pairs. Since R is supported on $\mathbf{gr}(i)$, we must have

$$i^* \alpha \sim_R \mathbf{a}^* \alpha,$$

for any $\alpha \in T^*N$. On the other hand, as explained in [11, Proposition 3.3], for any $e \in E|_M$ there exists a unique $X \in TM$ such that

$$X \sim_R e.$$

Since R is supported on $\mathbf{gr}(i)$, we have $i_* X = \mathbf{a}(e)$, or $X = \mathbf{a}(e)$. Thus $R = R_{E,M}$.

3. QUASI-HAMILTONIAN REDUCTION

3.1. Reduction theorems. Let \mathfrak{d} be a quadratic Lie algebra acting on a manifold N so that all the stabilizers are coisotropic, and let $\mathfrak{h} \subset \mathfrak{d}$ be a Lagrangian Lie subalgebra. We shall consider the following special case of general quasi-Hamiltonian (\mathbb{E}, E) -manifolds.

Definition 3.1. A *quasi-Hamiltonian* $(\mathfrak{d}, \mathfrak{h}) \times N$ -manifold is a manifold M together with a morphism of Manin pairs

$$(\mathbb{T}, T)M \dashrightarrow (\mathfrak{d}, \mathfrak{h}) \times N.$$

It is *exact* (or *quasi-symplectic*) if the morphism is exact.

Example 3.1. If $\mathfrak{d} = 0$ and N is a point then (as we saw in Example 2.8) a quasi-Hamiltonian structure is the same as a Poisson or (in the exact case) symplectic structure. More generally, a quasi-Hamiltonian $(0, 0) \times N$ -structure on M is equivalent to a Poisson structure on M and to a map $\mu : M \rightarrow N$ such that $\mu^*(C^\infty(N)) \subset C^\infty(M)$ is in the Poisson centre.

If \mathfrak{g} is a quadratic Lie algebra then an *exact* quasi-Hamiltonian $(\mathfrak{g} \oplus \bar{\mathfrak{g}}, \mathfrak{g}_\Delta) \times G$ -structure on M is equivalent to a quasi-Hamiltonian G -structure in the sense of Alekseev, Malkin and Meinrenken. If \mathfrak{h} is a Lie algebra then a quasi-Hamiltonian $(\mathfrak{h} \ltimes \mathfrak{h}^*) \times \mathfrak{h}^*$ -structure on M is equivalent to a Poisson structure on M together with a moment map $M \rightarrow \mathfrak{h}^*$ generating an action of \mathfrak{h} . In the exact case the Poisson structure is symplectic.

In this subsection we present a reduction procedure, which will be the main tool used in our study of the moduli spaces of flat connections.

Definition 3.2. Let M be a quasi-Hamiltonian $(\mathfrak{d}, \mathfrak{h}) \times N$ -manifold. *Reductive data* (\mathfrak{c}, S) consists of a coisotropic Lie subalgebra $\mathfrak{c} \subseteq \mathfrak{d}$ together with a \mathfrak{c} -invariant submanifold $S \subseteq N$ such that

- r1) the \mathfrak{c}^\perp -orbits in S form a regular foliation⁷ with quotient $q_N : S \rightarrow N_{\mathfrak{c},S}$,
- r2) the graph $\text{gr}(\mu)$, where μ is the underlying map $M \rightarrow N$, intersects $S \times M$ cleanly, and the $\mathfrak{h} \cap \mathfrak{c}^\perp$ -orbits in $\mu^{-1}(S)$ form a regular foliation with quotient $q_M : \mu^{-1}(S) \rightarrow N_{\mathfrak{c},S}$.

Theorem 3.1 (Quasi-Hamiltonian reduction). *Suppose (\mathfrak{c}, S) is reductive data for a morphism of Manin pairs*

$$(3.1) \quad R : (\mathbb{T}, T)M \dashrightarrow (\mathfrak{d}, \mathfrak{h}) \times N.$$

Then

$$(3.2) \quad R_{\mathfrak{c},S} : (\mathbb{T}, T)M_{\mathfrak{c},S} \dashrightarrow (\mathfrak{d}_{\mathfrak{c}}, \mathfrak{h}_{\mathfrak{c}}) \times N_{\mathfrak{c},S}$$

is a morphism of Manin pairs, where

$$M_{\mathfrak{c},S} = \mu^{-1}(S) / \mathfrak{h} \cap \mathfrak{c}^\perp$$

$$R_{\mathfrak{c},S} := R_2 \circ R \circ R_1^\top,$$

$$R_2 := R_{\mathfrak{c}} \times (\text{gr}(q_N) \circ \text{gr}(i_N)^\top),$$

$$R_1 := R_{q_M} \circ R_{i_M}^\top,$$

$i_N : S \rightarrow N$ and $i_M : \mu^{-1}(S) \rightarrow M$ are the canonical inclusions, and $R_{\mathfrak{c}} : \mathfrak{d} \dashrightarrow \mathfrak{d}_{\mathfrak{c}}$ and $\mathfrak{h}_{\mathfrak{c}}$ are as in Example 2.7.

The first two statements in Theorem 1.1 follow as consequences of Theorem 3.1.

Theorem 3.2 (Quasi-Hamiltonian reduction in the exact case). *If, in the setup of Theorem 3.1, the following additional assumptions hold:*

- \mathfrak{c} acts transitively on S ,
- (3.1) is an exact morphism of Manin pairs,

then

- (3.2) is an exact morphism of Manin pairs.

The third statement in Theorem 1.1 follows as a consequence of Theorem 3.2.

We delay the proof of both these theorems to Appendix A.

Notice that if $\mathfrak{c} \subset \mathfrak{d}$ is Lagrangian then the reduced manifold is Poisson or (in the exact case) symplectic, as $\mathfrak{d}_{\mathfrak{c}} = 0$.

⁷By a *regular foliation*, we mean that the leaf space carries the structure of a smooth manifold for which the quotient map is a surjective submersion.

3.2. Bivector fields and quasi-Poisson structures. In this section we shall explain Theorem 3.1 in more traditional terms, using bivector fields.

Suppose that

$$R : (\mathbb{T}, T)M \dashrightarrow (\mathfrak{d}, \mathfrak{h}) \times N$$

is a morphism of Manin pairs over $\mu : M \rightarrow N$, and the subspace $\mathfrak{k} \subseteq \mathfrak{d}$ is a Lagrangian complement to $\mathfrak{h} \subseteq \mathfrak{d}$: that is, $\mathfrak{d} = \mathfrak{h} \oplus \mathfrak{k}$. Then axiom (m1) of Definition 2.5 implies that R composes transversely with \mathfrak{k} , while property (m2) implies that $\mathfrak{k} \circ R \subseteq \mathbb{T}M$ is a Lagrangian complement to TM . Thus there exists a unique bivector field $\pi_{\mathfrak{k}} \in \mathfrak{X}^2(M)$ such that

$$\mathfrak{k} \circ R = \text{gr}(\pi^{\sharp}) := \{(\pi_{\mathfrak{k}}(\alpha, \cdot) + \alpha) \mid \alpha \in T^*M\}.$$

The triple $(\mathfrak{d}, \mathfrak{h}; \mathfrak{k})$ is called a quasi-Manin triple, and $(M, \pi_{\mathfrak{k}}, \rho_R)$ is called a Hamiltonian quasi-Poisson $(\mathfrak{d}, \mathfrak{h}; \mathfrak{k})$ -space with moment map $\mu : M \rightarrow N$ (cf. [11, 19]). The bivector field $\pi_{\mathfrak{k}}$ is called a quasi-Poisson structure.

Suppose that (\mathfrak{c}, S) is reductive data for the morphism of Manin pairs R . We want to reinterpret Theorem 3.1 using the language of quasi-Poisson geometry. Thus we will be interested in Lagrangian complements to the reduced Lie algebra $\mathfrak{h}_{\mathfrak{c}}$.

Lemma 3.1. *Each Lagrangian complement $\mathfrak{k}'_{\mathfrak{c}} \subseteq \mathfrak{d}_{\mathfrak{c}}$ to $\mathfrak{h}_{\mathfrak{c}}$ defines an element*

$$\tau \in \wedge^2(\mathfrak{h}/(\mathfrak{h} \cap \mathfrak{c}^{\perp})).$$

Proof. Suppose $\mathfrak{k}'_{\mathfrak{c}} \subseteq \mathfrak{d}_{\mathfrak{c}} := \mathfrak{c}/\mathfrak{c}^{\perp}$ is a Lagrangian complement to $\mathfrak{h}_{\mathfrak{c}}$. Let $R_{\mathfrak{c}} : \mathfrak{d} \dashrightarrow \mathfrak{d}_{\mathfrak{c}}$ be the relation described in Example 2.7, and define $\mathfrak{k}' := \mathfrak{k}'_{\mathfrak{c}} \circ R_{\mathfrak{c}} \subseteq \mathfrak{c}$. Now \mathfrak{k}' can be seen as the graph

$$\mathfrak{k}' = \{(\xi + \tau^{\sharp}(\xi) + \eta) \mid \xi \in \mathfrak{k}, \langle \xi, \cdot \rangle|_{\mathfrak{h} \cap \mathfrak{k}'} = 0, \text{ and } \eta \in \mathfrak{h} \cap \mathfrak{k}'\}$$

of a map

$$\tau^{\sharp} : \text{ann}(\mathfrak{h} \cap \mathfrak{k}') \cong (\mathfrak{h}/(\mathfrak{h} \cap \mathfrak{k}'))^* \rightarrow \mathfrak{h}/(\mathfrak{h} \cap \mathfrak{k}'),$$

where $\text{ann}(\mathfrak{h} \cap \mathfrak{k}') := (\mathfrak{h} \cap \mathfrak{k}')^{\perp} \cap \mathfrak{k}$. Let $\tau \in \mathfrak{h}/(\mathfrak{h} \cap \mathfrak{k}') \otimes \mathfrak{h}/(\mathfrak{h} \cap \mathfrak{k}')$ be the element defined by $\tau^{\sharp}(\xi) = \tau(\xi, \cdot)$, then the fact that \mathfrak{k}' is Lagrangian forces τ to be skew-symmetric. Finally, since $\mathfrak{k}'_{\mathfrak{c}} + \mathfrak{h}_{\mathfrak{c}} = \mathfrak{c}/\mathfrak{c}^{\perp}$ we have $\mathfrak{h} + \mathfrak{k}' = \mathfrak{h} + \mathfrak{c}$. Hence $\mathfrak{h} \cap \mathfrak{k}' = \mathfrak{h} \cap \mathfrak{c}^{\perp}$. Thus $\tau \in \wedge^2(\mathfrak{h}/(\mathfrak{h} \cap \mathfrak{c}^{\perp}))$. \square

Theorem 3.3. *Suppose that (\mathfrak{c}, S) is reductive data for the morphism of Manin pairs*

$$(3.3) \quad R : (\mathbb{T}, T)M \dashrightarrow (\mathfrak{d}, \mathfrak{h}) \times N,$$

and $(M, \pi_{\mathfrak{k}}, \rho_R)$ is the Hamiltonian quasi-Poisson $(\mathfrak{d}, \mathfrak{h}; \mathfrak{k})$ -space corresponding to the Lagrangian complement $\mathfrak{k} \subseteq \mathfrak{d}$ to \mathfrak{h} . Let

$$R_{\mathfrak{c}, S} : (\mathbb{T}, T)M_{\mathfrak{c}, S} \dashrightarrow (\mathfrak{d}_{\mathfrak{c}}, \mathfrak{h}_{\mathfrak{c}}) \times N_{\mathfrak{c}, S}$$

denote the reduced morphism of Manin pairs described in Theorem 3.1, and let $(M_{\mathfrak{c}, S}, \pi_{\mathfrak{k}'_{\mathfrak{c}}}, \rho_{R_{\mathfrak{c}, S}})$ be the Hamiltonian $(\mathfrak{d}_{\mathfrak{c}}, \mathfrak{h}_{\mathfrak{c}}; \mathfrak{k}'_{\mathfrak{c}})$ -quasi Poisson manifold corresponding to a chosen Lagrangian complement $\mathfrak{k}'_{\mathfrak{c}}$ to $\mathfrak{h}_{\mathfrak{c}}$.

As in Lemma 3.1, let $\tau \in \wedge^2(\mathfrak{h}/(\mathfrak{h} \cap \mathfrak{c}^{\perp}))$ be the element corresponding to $\mathfrak{k}'_{\mathfrak{c}}$. Then

$$(\pi_{\mathfrak{k}} + \rho_R(\tau))|_{\mu^{-1}(S)}$$

is an $\mathfrak{h} \cap \mathfrak{c}^\perp$ -invariant section of $\wedge^2(T(\mu^{-1}(S))/\rho_R(\mathfrak{h} \cap \mathfrak{c}^\perp))$ which is mapped to $\pi_{\mathfrak{k}'_\mathfrak{c}}$ under the surjective submersion $\mu^{-1}(S) \rightarrow M_{\mathfrak{c},S}$.

Proof. Let $R_\mathfrak{c} : \mathfrak{d} \dashrightarrow \mathfrak{d}_\mathfrak{c}$ be the relation described in Example 2.7. Let $\mu : M \rightarrow N$ denote the map supporting R , and let

$$R_{i_M} : \mathbb{T}\mu^{-1}(S) \dashrightarrow \mathbb{T}M, \quad R_{q_M} : \mathbb{T}\mu^{-1}(S) \dashrightarrow \mathbb{T}M_{\mathfrak{c},S}$$

denote the standard lifts of the inclusion and projection, respectively. Then

$$\mathrm{gr}(\pi_{\mathfrak{k}'_\mathfrak{c}}^\sharp) = \mathfrak{k}'_\mathfrak{c} \circ R_\mathfrak{c} \circ R \circ R_{i_M} \circ R_{q_M}^\top.$$

Now, as explained in Lemma 3.1,

$$(3.4a) \quad \mathfrak{k}'_\mathfrak{c} \circ R_\mathfrak{c} = \mathfrak{k}' := \{(\xi + \tau^\sharp(\xi) + \eta) \mid \xi \in \mathfrak{k}, \langle \xi, \cdot \rangle|_{\mathfrak{h} \cap \mathfrak{c}^\perp} = 0, \text{ and } \eta \in \mathfrak{h} \cap \mathfrak{c}^\perp\}.$$

Meanwhile, since $\mathrm{gr}(\pi_{\mathfrak{k}}^\sharp) = \mathfrak{k} \circ R$, it follows that

$$(3.4b) \quad X + \alpha \sim_R \xi + \eta, \quad \xi \in \mathfrak{k}, \eta \in \mathfrak{h} \Leftrightarrow X = \rho_R(\eta) + \pi_{\mathfrak{k}}^\sharp \alpha, \text{ and } \xi = j \circ \rho_R^* \alpha,$$

where $j : \mathfrak{h}^* \rightarrow \mathfrak{k}$ inverts the isomorphism $\mathfrak{k} \rightarrow \mathfrak{d}/\mathfrak{h} \cong \mathfrak{h}^*$.

Using Eqs. 3.4 we compute

$$\mathfrak{k}'_\mathfrak{c} \circ R_\mathfrak{c} \circ R = \{(\rho_R(\eta) + (\pi_{\mathfrak{k}} + \rho(\tau))^\sharp(\alpha) + \alpha) \mid \eta \in \mathfrak{h} \cap \mathfrak{c}^\perp \text{ and } \alpha \in \mathrm{ann}(\rho_R(\mathfrak{h} \cap \mathfrak{c}^\perp))\}.$$

This shows that $\pi_{\mathfrak{k}} + \rho(\tau)$ is $\mathfrak{h} \cap \mathfrak{c}^\perp$ -invariant. Moreover, $\mathfrak{k}'_\mathfrak{c} \circ R_\mathfrak{c} \subseteq \mathfrak{c}$ and thus elements of $\mathfrak{k}'_\mathfrak{c} \circ R_\mathfrak{c}$ act to preserve S . In turn, this implies that

$$(\pi_{\mathfrak{k}} + \rho(\tau))|_{\mu^{-1}(S)} \in \Gamma\left(\wedge^2(T\mu^{-1}(S)/\rho_R(\mathfrak{h} \cap \mathfrak{c}^\perp))\right)$$

is in fact a section of $\wedge^2(T\mu^{-1}(S)/\rho_R(\mathfrak{h} \cap \mathfrak{c}^\perp))$.

Now, $\mathrm{gr}(\pi_{\mathfrak{k}'_\mathfrak{c}}^\sharp) = \mathfrak{k}'_\mathfrak{c} \circ R_\mathfrak{c} \circ R \circ R_{i_M} \circ R_{q_M}^\top$, i.e.

$$(q_M)_*(\pi_{\mathfrak{k}} + \rho(\tau))|_{\mu^{-1}(S)} = \pi_{\mathfrak{k}'_\mathfrak{c}}.$$

□

3.3. Exact morphisms of Manin pairs and 2-forms. In this section, we will examine exact morphisms of Manin pairs in more detail. We recall from [11, 25] that once isotropic splittings are chosen, these are uniquely determined by a map between the underlying spaces, and a 2-form on the domain. This description in terms of 2-forms can be useful for simplifying calculations.

Suppose \mathbb{E} is an exact Courant algebroid over N . That is, the sequence

$$(3.5) \quad 0 \rightarrow T^*N \xrightarrow{\mathbf{a}^*} \mathbb{E} \xrightarrow{\mathbf{a}} TN \rightarrow 0$$

is exact. Let $s : TN \rightarrow \mathbb{E}$ be a splitting of Eq. (3.5) such that $s(TN) \subseteq \mathbb{E}$ is isotropic (such splittings are called *isotropic splittings*). Then, as explained in [37], the formula

$$\iota_X \iota_Y \iota_Z \gamma := \langle \llbracket s(X), s(Y) \rrbracket, s(Z) \rangle, \quad X, Y, Z \in \mathfrak{X}(N)$$

defines a closed 3-form, $\gamma \in \Omega^3(N)$, called the *curvature 3-form of the splitting* s . The isomorphism $s \oplus \mathbf{a}^* : TN \oplus T^*N \xrightarrow{\cong} \mathbb{E}$ identifies the metric on \mathbb{E} with

$$\langle X + \alpha, Y + \beta \rangle = \alpha(Y) + \beta(X), \quad X, Y \in TN, \quad \alpha, \beta \in T^*N,$$

and the bracket with

$$(3.6) \quad \llbracket X + \alpha, Y + \beta \rrbracket = [X, Y] + \mathcal{L}_X \beta - \iota_Y d\alpha + \iota_X \iota_Y \gamma, \quad X, Y \in \mathfrak{X}(N), \quad \alpha, \beta.$$

The Courant algebroid with underlying bundle $TN \oplus T^*N$, and bracket given by Eq. (3.6) is called the γ -*twisted exact Courant algebroid* over N , and denoted by $\mathbb{T}_\gamma N$.

Example 3.2 (The Cartan Courant algebroid [37]). The Cartan Courant algebroid

$$(\bar{\mathfrak{g}} \oplus \mathfrak{g}) \times G$$

reviewed in Example 2.4 is exact. In [1] it is shown that the map $s : TG \rightarrow (\bar{\mathfrak{g}} \oplus \mathfrak{g}) \times G$ defined as

$$s : X \rightarrow \frac{1}{2}(-\iota_X(dg g^{-1}), \iota_X(g^{-1}dg)), \quad X \in TG$$

is a $(\bar{\mathfrak{g}} \oplus \mathfrak{g})$ -invariant isotropic splitting of the Cartan Courant algebroid. The corresponding curvature 3-form is computed as $\gamma = \frac{1}{24}\langle [g^{-1}dg, g^{-1}dg], g^{-1}dg \rangle$ (note that the normalization differs from that in [1]).

Suppose $L \subseteq \mathbb{E}$ is a Lagrangian subbundle with support on $S \subseteq N$, such that $\mathbf{a}(L) = TS$, then there is a 2-form $\omega \in \Omega^2(S)$ uniquely determined by the formula

$$\iota_X \iota_Y \omega = \langle s(X), e \rangle, \quad X, Y \in TS, \quad e \in L \text{ and } \mathbf{a}(e) = Y.$$

Thus, we may identify L with

$$L = \mathbf{gr}(\omega^\flat) := \{(s(X) + \mathbf{a}^*(\iota_X \omega + \alpha)) \mid X \in TS, \alpha \in \mathbf{ann}(TS)\} \subseteq \mathbb{E}.$$

A quick calculation using Eq. (3.6) shows that L is a Dirac structure with support on S if and only if $d\omega = i^* \gamma$, where $i : S \rightarrow N$ is the inclusion (cf. [37] and [11, Proposition 2.8]).

Suppose

$$R : \mathbb{T}M \dashrightarrow \mathbb{E}$$

is a Courant morphism supported on the graph of a map $\mu : M \rightarrow N$. Since \mathbb{E} is an exact Courant algebroid, R is *exact* if $\mathbf{a}(R) = T\mathbf{gr}(\mu)$ (cf. Lemma 2.1), in which case the considerations above show that

$$R = \mathbf{gr}(\omega^\flat) \subseteq \mathbb{E} \times \overline{\mathbb{T}M},$$

where $\omega \in \Omega^2(M) \cong \Omega^2(\mathbf{gr}(\mu))$ satisfies $d\omega = \mu^* \gamma$ [11, 25]. That is, $R = R_{\mu, \omega}$, where $R_{\mu, \omega}$ is defined by

$$(3.7) \quad X - \iota_X \omega + \mu^* \alpha \sim_{R_{\mu, \omega}} s(\mu_* X) + \mathbf{a}^* \alpha, \quad X \in TM, \quad \alpha \in T^*N.$$

In particular, if $E \subseteq \mathbb{E}$ is a Dirac structure, once an isotropic splitting $s : TN \rightarrow \mathbb{E}$ is chosen, a morphism of Manin pairs

$$R : (\mathbb{T}, T)M \dashrightarrow (\mathbb{E}, E)$$

is entirely determined by the underlying map $\mu : M \rightarrow N$ between the spaces, and a 2-form $\omega \in \Omega^2(M)$ (satisfying certain conditions).

Remark 3.1 (Twisted quasi-Hamiltonian structures). Recall the Γ -twisted Cartan Dirac structure described in Example 2.5,

$$((\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_\Gamma}, \mathfrak{g}_\Gamma) \times G^{E_\Gamma}.$$

Suppose we use the splitting $TG^{E_\Gamma} \rightarrow ((\bar{\mathfrak{g}} \oplus \mathfrak{g}) \times G)^{E_\Gamma}$ described in Example 3.2 to identify

$$((\bar{\mathfrak{g}} \oplus \mathfrak{g}) \times G)^{E_\Gamma} \cong \mathbb{T}_\gamma G^{E_\Gamma},$$

where $\gamma = \frac{1}{24} \langle [g^{-1}dg, g^{-1}dg], g^{-1}dg \rangle$. Then exact morphisms of Manin pairs

$$R : (\mathbb{T}, T)M \dashrightarrow ((\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_\Gamma}, \mathfrak{g}_\Gamma) \times G^{E_\Gamma}$$

are in one-to-one correspondence with quadruples (M, μ, ρ, ω) , where

- $\mu : M \rightarrow G^{E_\Gamma}$ is a smooth map,
- $\rho : \mathfrak{g}^{V_\Gamma} \rightarrow \mathfrak{X}(M)$ is a Lie algebra action, and
- $\omega \in \Omega^2(M)$ is a 2-form,

such that

- (1) $d\omega = \mu^* \gamma$,
- (2) $\mu : M \rightarrow G^{E_\Gamma}$ is \mathfrak{g}^{V_Γ} -equivariant with respect to the \mathfrak{g}^{V_Γ} action on G^{E_Γ} given on the $e \in E_\Gamma$ -th factor by

$$\xi \rightarrow -\xi_{\text{in}(e)}^R + \xi_{\text{out}(e)}^L, \quad \xi \in \mathfrak{g}^{V_\Gamma},$$

- (3) $\ker(d\mu)_x \cap \ker(\omega^\flat)_x = 0$, for every $x \in M$, and
- (4)

$$\iota_{\rho(\xi)} \omega = \frac{1}{2} \mu^* \sum_{e \in E_\Gamma} \langle g_e^{-1} dg_e, \xi_{\text{out}(e)} \rangle + \langle dg_e g_e^{-1}, \xi_{\text{in}(e)} \rangle.$$

(Note that conditions (2),(3) and (4) determine ρ uniquely in terms of μ and ω). The quadruple (M, μ, ρ, ω) corresponds to the morphism of Manin pairs Eq. (3.7), supported on the graph of μ .

The quadruples (M, μ, ρ, ω) generalize quasi-Hamiltonian G^{E_Γ} -structures in the sense that they incorporate an automorphism of G^{E_Γ} into their definition. Here the automorphism is simply the permutation of factors described by the permutation graph, Γ . Allowing arbitrary automorphisms leads to a definition of twisted quasi-Hamiltonian spaces along the lines of the one given in [26] for twisted quasi-Poisson structures (cf. [26, Definition 3]).

Let $G_{\text{big}} := G^{V_\Gamma} \rtimes \mathbb{Z}$, where \mathbb{Z} acts by permuting the factors according to the graph Γ (cf. Remark 2.4). As explained to the authors by Eckhard Meinrenken, quasi-Hamiltonian $((\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_\Gamma}, \mathfrak{g}_\Gamma) \times G^{E_\Gamma}$ -structures are quasi-Hamiltonian G_{big} -structures in the (original) sense of Alekseev-Malkin-Meinrenken [4] for which the moment map takes values in $G^{V_\Gamma} \times \{-1\} \subset G_{\text{big}}$ (cf. Remark 2.4).

We now examine the behaviour of twisted exact Courant algebroids under the partial reduction procedure described in Theorem 3.1. Suppose that $\mathfrak{d} \times N$ is an exact Courant algebroid, and a \mathfrak{d} -invariant isotropic splitting $s : TN \rightarrow \mathfrak{d} \times N$ is chosen, defining an isomorphism

$$\mathfrak{d} \times N \cong \mathbb{T}_\gamma N,$$

We let $E \subseteq \mathbb{T}_\gamma N$ denote the Dirac structure corresponding to $\mathfrak{h} \times N \subseteq \mathfrak{d} \times N$ under this isomorphism. Suppose

$$R_{\mu,\omega} : (\mathbb{T}, T)M \dashrightarrow (\mathbb{T}_\gamma N, E) \cong (\mathfrak{d}, \mathfrak{h}) \times N$$

is an exact morphism of Manin pairs, and $S \subseteq N$ is an orbit of the coisotropic subalgebra $\mathfrak{c} \subseteq \mathfrak{d}$, and the assumptions of both Theorems 3.1 and 3.2 hold for the reductive data (\mathfrak{c}, S) . Let $i_N : S \rightarrow N$ and $i_M : \mu^{-1}(S) \rightarrow M$ denote the inclusions, and $q_N : S \rightarrow N_{\mathfrak{c},S}$ and $q_M : \mu^{-1}(S) \rightarrow M_{\mathfrak{c},S}$ the quotients (by \mathfrak{c}^\perp and $\mathfrak{h} \cap \mathfrak{c}^\perp$, respectively).

We would like to define a splitting of the reduced Courant algebroid $\mathfrak{d}_{\mathfrak{c}^\perp} \times N_{\mathfrak{c},S}$. As explained in [10, Proposition 3.6], unless $s(TS) \subseteq \mathfrak{c}$, this will depend on a choice of a \mathfrak{c}^\perp -invariant connection 1-form $\theta \in \Omega^1(S, \mathfrak{c}^\perp)$ for the bundle $q_N : S \rightarrow N_{\mathfrak{c},S}$. For any $X \in \mathfrak{X}(N_{\mathfrak{c},S})$, let $X^h \in \mathfrak{X}(S)$ denote its horizontal lift with respect to the chosen connection. The map $X \rightarrow s(X^h) \in \Gamma(\mathfrak{d} \times S)$ may not take values in $\Gamma(\mathfrak{c} \times S)$, but

$$s_\theta : X \rightarrow s(X^h) + \mathbf{a}^*(\iota_{X^h} \langle \theta, \vartheta_s \rangle)$$

does, where $\vartheta_s \in \Omega^1(S, (\mathfrak{c}^\perp)^*)$ is defined by $\langle \xi, \vartheta_s \rangle := s^* \xi$, for $\xi \in \mathfrak{c}^\perp$. Note also that $s_\theta(X)$ is \mathfrak{c}^\perp -invariant, and hence descends to a unique section of $\Gamma(\mathfrak{c} \times N_{\mathfrak{c},S})$. Thus, the composition

$$\mathfrak{X}(N_{\mathfrak{c},S}) \xrightarrow{s_\theta} \Gamma(\mathfrak{c} \times S) \rightarrow \Gamma(\mathfrak{c}/\mathfrak{c}^\perp \times N_{\mathfrak{c},S}) = \Gamma(\mathfrak{d}_{\mathfrak{c}} \times N_{\mathfrak{c},S})$$

defines an isotropic splitting $\tilde{s}_\theta : TN_{\mathfrak{c},S} \rightarrow \mathfrak{d}_{\mathfrak{c}} \times N_{\mathfrak{c},S}$. We define $\tilde{\gamma}_\theta \in \Omega^3(N_{\mathfrak{c},S})$ to be the associated curvature 3-form. Let $E_{\mathfrak{c},S} \subseteq \mathbb{T}_{\tilde{\gamma}_\theta} N_{\mathfrak{c},S}$ denote the Dirac structure corresponding to $\mathfrak{h}_{\mathfrak{c},S}$ under the isomorphism defined by \tilde{s}_θ .

Proposition 3.1 (Partial reduction for split exact Courant algebroids). *Suppose that*

$$(3.8) \quad R_{\mu,\omega} : (\mathbb{T}, T)M \dashrightarrow (\mathbb{T}_\gamma N, E) \cong (\mathfrak{d}, \mathfrak{h}) \times N$$

is an exact morphism of Manin pairs, that the assumptions of both Theorems 3.1 and 3.2 hold for the reductive data (\mathfrak{c}, S) . Let $\theta \in \Omega^1(S, \mathfrak{c}^\perp)$ be a \mathfrak{c}^\perp -invariant connection 1-form for the bundle $q_N : S \rightarrow N_{\mathfrak{c},S}$.

Then, under the isomorphism $\mathfrak{d}_{\mathfrak{c}} \times N_{\mathfrak{c},S} \cong \mathbb{T}_{\tilde{\gamma}_\theta} N_{\mathfrak{c},S}$ defined by the isotropic splitting \tilde{s}_θ , the reduced morphism of Manin pairs (3.2), described in Theorem 3.1, is identified with

$$(3.9) \quad R_{\tilde{\mu}, \tilde{\omega}_\theta} : (\mathbb{T}, T)M_{\mathfrak{c},S} \dashrightarrow (\mathbb{T}_{\tilde{\gamma}_\theta} N_{\mathfrak{c},S}, E_{\mathfrak{c},S}),$$

where $\tilde{\omega}_\theta \in \Omega^2(M_{\mathfrak{c},S})$ is defined by the equation

$$(3.10) \quad q_M^* \tilde{\omega}_\theta = i_M^* \omega - \mu^* \langle \theta, \vartheta_s - \frac{1}{2} s \circ \mathbf{a}(\theta) \rangle,$$

and $\tilde{\mu} : M_{\mathfrak{c},S} \rightarrow N_{\mathfrak{c},S}$ is the unique map such that

$$\begin{array}{ccc} \mu^{-1}(S) & \xrightarrow{\mu} & S \\ q_M \downarrow & & \downarrow q_N \\ M_{\mathfrak{c},S} & \xrightarrow{\tilde{\mu}} & N_{\mathfrak{c},S} \end{array}$$

commutes.

Moreover, the 2-form $\tilde{\omega}_\theta$ is independent of θ if $s(TS) \subseteq \mathfrak{c}$, more precisely, the term $\mu^* \langle \theta, \vartheta_s - \frac{1}{2} s \circ \mathbf{a}(\theta) \rangle$ in Eq. (3.10) vanishes.

We defer the proof of Proposition 3.1 to Appendix A

Remark 3.2. In the special case where $\mathfrak{c} = \mathfrak{c}^\perp$ is Lagrangian, then \mathfrak{c}^\perp acts transitively on S , so Eq. (3.10) simplifies to

$$q_M^* \tilde{\omega}_\theta = i_M^* \omega - \frac{1}{2} \mu^* \langle \theta, s \circ \mathbf{a}(\theta) \rangle.$$

Remark 3.3. Consider the Γ -twisted Cartan Dirac structure $((\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_\Gamma}, \mathfrak{g}_\Gamma) \times G^{E_\Gamma}$. The conditions that $\mathfrak{c} \subseteq (\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_\Gamma}$ be symmetric and that $S \subseteq G^{E_\Gamma}$ be the \mathfrak{c} -orbit through the identity imply that

$$(3.11) \quad s(TS) \subseteq \mathfrak{c}$$

Indeed, Eq. (3.11) is easily checked at the identity of G^{E_Γ} , and the invariance of s implies that it holds at every other point in the \mathfrak{c} -orbit, S .

Thus, Eq. (3.10) simplifies to

$$q_M^* \tilde{\omega}_\theta = i_M^* \omega,$$

and Theorem 1.3 follows as a corollary to Proposition 3.1.

Example 3.3 (Conjugacy classes and quasi-Hamiltonian geometry). Recall the splitting $s : TG \rightarrow (\bar{\mathfrak{g}} \oplus \mathfrak{g}) \times G$ of the Cartan Courant algebroid described in Example 3.2. Suppose $\mathfrak{c} = \mathfrak{g}_\Delta \subseteq (\bar{\mathfrak{g}} \oplus \mathfrak{g})$ is the diagonal subalgebra, and $S \subseteq G$ is a conjugacy class. Then,

$$s \circ \mathbf{a}(\xi, \xi) = (\xi - \mathbf{Ad}_g \xi, -\mathbf{Ad}_{g^{-1}} \xi + \xi), \quad \xi \in \mathfrak{g}, \quad g \in G.$$

So

$$\iota_{\mathbf{a}(\eta, \eta)} \iota_{\mathbf{a}(\xi, \xi)} \frac{1}{2} \mu^* \langle \theta, s \circ \mathbf{a}(\theta) \rangle = \langle \eta, \mathbf{Ad}_g \xi \rangle - \langle \mathbf{Ad}_g \eta, \xi \rangle.$$

Thus $\frac{1}{2} \langle \theta, s \circ \mathbf{a}(\theta) \rangle$ is proportional to the quasi-Hamiltonian 2-form on the conjugacy class, S , described in [4, Proposition 3.1].

Suppose now that M is a quasi-Hamiltonian G space with moment map $\mu : M \rightarrow G$, and 2-form ω . It follows that partial reduction of the morphism of Manin pairs

$$R_{\mu, \omega} : (\mathbb{T}, T)M \dashrightarrow (\bar{\mathfrak{g}} \oplus \mathfrak{g}, \mathfrak{g}_\Delta) \times G$$

by (\mathfrak{c}, S) , yields the same symplectic structure as the quasi-Hamiltonian reduction

$$M \otimes S //_1 G,$$

where $M \otimes S$ denotes the fusion of M with S , as described in [4].

3.4. Commutativity of reductions. Suppose that $S_1, S_2 \subseteq N$ intersect cleanly, and that (\mathfrak{c}_1, S_1) and (\mathfrak{c}_2, S_2) are both reductive data for a morphism of Manin pairs

$$(3.12) \quad R : (\mathbb{T}, T)M \dashrightarrow (\mathfrak{d}, \mathfrak{h}) \times N.$$

Notice that the Lie algebra $\mathfrak{c}_{2,1} := R_{\mathfrak{c}_1} \circ \mathfrak{c}_2 \subseteq \mathfrak{d}_{\mathfrak{c}_1}$ is coisotropic. Let $S_{2,1}$ denote the image of $S_2 \cap S_1$ under the quotient map $S_1 \rightarrow N_{\mathfrak{c}_1, S_1}$. In practise, $(\mathfrak{c}_{2,1}, S_{2,1})$ will often form reductive data for

$$R_{\mathfrak{c}_{2,1}, S_{2,1}} : (\mathbb{T}, T)M_{\mathfrak{c}_{2,1}, S_{2,1}} \dashrightarrow (\mathfrak{d}_{\mathfrak{c}_1}, \mathfrak{h}_{\mathfrak{c}_1}) \times N_{\mathfrak{c}_1, S_1}.$$

Similarly, with $\mathfrak{c}_{1,2} := R_{\mathfrak{c}_2} \circ \mathfrak{c}_1 \subseteq \mathfrak{d}_{\mathfrak{c}_2}$ and $S_{1,2}$ the image of $S_1 \cap S_2$ under the quotient map $S_2 \rightarrow N_{\mathfrak{c}_2, S_2}$, the pair $(\mathfrak{c}_{1,2}, S_{1,2})$ will often form reductive data for

$$R_{\mathfrak{c}_{1,2}, S_{1,2}} : (\mathbb{T}, T)M_{\mathfrak{c}_{1,2}, S_{1,2}} \dashrightarrow (\mathfrak{d}_{\mathfrak{c}_2}, \mathfrak{h}_{\mathfrak{c}_2}) \times N_{\mathfrak{c}_2, S_2}.$$

Proposition 3.2 (Commutativity of reductions). *Suppose that $S_1, S_2 \subseteq N$ intersect cleanly, that (\mathfrak{c}_1, S_1) , (\mathfrak{c}_2, S_2) and $(\mathfrak{c}_1 \cap \mathfrak{c}_2, S_1 \cap S_2)$ all form reductive data for the morphism of Manin pairs*

$$R : (\mathbb{T}, T)M \dashrightarrow (\mathfrak{d}, \mathfrak{h}) \times N,$$

while $(\mathfrak{c}_{2,1}, S_{2,1})$ and $(\mathfrak{c}_{1,2}, S_{1,2})$ form reductive data for

$$R_{\mathfrak{c}_{2,1}, S_{2,1}} : (\mathbb{T}, T)M_{\mathfrak{c}_{2,1}, S_{2,1}} \dashrightarrow (\mathfrak{d}_{\mathfrak{c}_1}, \mathfrak{h}_{\mathfrak{c}_1}) \times N_{\mathfrak{c}_1, S_1}$$

and

$$R_{\mathfrak{c}_{1,2}, S_{1,2}} : (\mathbb{T}, T)M_{\mathfrak{c}_{1,2}, S_{1,2}} \dashrightarrow (\mathfrak{d}_{\mathfrak{c}_2}, \mathfrak{h}_{\mathfrak{c}_2}) \times N_{\mathfrak{c}_2, S_2},$$

respectively.

Then

$$(3.13) \quad (R_{\mathfrak{c}_1, S_1})_{\mathfrak{c}_{2,1}, S_{2,1}} = R_{\mathfrak{c}_1 \cap \mathfrak{c}_2, S_1 \cap S_2} = (R_{\mathfrak{c}_2, S_2})_{\mathfrak{c}_{1,2}, S_{1,2}}.$$

The proof of this proposition is deferred to Appendix A.0.2.

In this paper, Proposition 3.2 will always be applied as the following corollary:

Corollary 3.1 (Reductions of distinct factors commute). *Suppose $\mathfrak{d} \times N, \mathfrak{d}' \times N'$, and $\mathfrak{d}'' \times N''$ are all action Courant algebroids, $S \subseteq N$ and $S' \subseteq N'$ are submanifolds and $(\mathfrak{c} \oplus \mathfrak{d}' \oplus \mathfrak{d}'', S \times N' \times N'')$, $(\mathfrak{d} \oplus \mathfrak{c}' \oplus \mathfrak{d}'', N \times S' \times N'')$ and $(\mathfrak{c} \oplus \mathfrak{c}' \oplus \mathfrak{d}'', S \times S' \times N'')$ each form reductive data for a morphism of Manin pairs*

$$R : (\mathbb{T}, T)M \dashrightarrow (\mathfrak{d} \oplus \mathfrak{d}' \oplus \mathfrak{d}'', \mathfrak{h}) \times (N \times N' \times N'').$$

Then

$$\begin{aligned} (R_{\mathfrak{c} \oplus \mathfrak{d}' \oplus \mathfrak{d}'', S \times N' \times N''})_{(\mathfrak{d} \oplus \mathfrak{c}' \oplus \mathfrak{d}'', N_{\mathfrak{c}, S \times S' \times N''})} &= R_{\mathfrak{c} \oplus \mathfrak{c}' \oplus \mathfrak{d}'', S \times S' \times N''} \\ &= (R_{\mathfrak{d} \oplus \mathfrak{c}' \oplus \mathfrak{d}'', N \times S' \times N''})_{(\mathfrak{c} \oplus \mathfrak{d}'_{\mathfrak{c}'} \oplus \mathfrak{d}'', S \times N'_{\mathfrak{c}', S'} \times N'')}. \end{aligned}$$




FIGURE 19. Sewing two edges from Σ . The two edges must have opposite orientation to ensure that the resulting surface is orientable.

4. QUASI-HAMILTONIAN STRUCTURES ON MODULI SPACES OF FLAT CONNECTIONS

Suppose that (Σ, V) is a marked surface, i.e. Σ is a compact oriented surface and $V \subset \partial\Sigma$ a finite set which intersects each component of Σ and $\partial\Sigma$ non-trivially. Let Γ be the boundary graph of Σ with the vertex set V (see Section 1.2.3 for details). In this section we shall prove Theorem 1.2 using quasi-Hamiltonian reduction. In other words, we want to construct an exact morphism of Manin pairs

$$(4.1) \quad R_{\Sigma, V} : (\mathbb{T}, T) \mathcal{M}_{\Sigma, V} \dashrightarrow ((\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_{\Gamma}}, \mathfrak{g}_{\Gamma}) \times G^{E_{\Gamma}}$$

over the map

$$\mu : \mathcal{M}_{\Sigma, V} \rightarrow G^{E_{\Gamma}}$$

given by the boundary holonomies.

Let us start with a simple case:


Proposition 4.1 (union of polygons). *Let Σ be a disjoint union of discs and $V \subset \partial\Sigma$ a finite subset meeting every boundary circle. Then there is a unique exact morphism*

$$(\mathbb{T}, T) \mathcal{M}_{\Sigma, V} \dashrightarrow ((\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_{\Gamma}}, \mathfrak{g}_{\Gamma}) \times G^{E_{\Gamma}}$$

over μ .

Proof. The map μ is in this case an embedding. As explained in Example 2.9, this implies that there exists a unique exact morphism of Manin pairs \square

4.1. Sewing construction. Suppose Σ is a (possibly disconnected) marked surface, and $e_1, e_2 \in \Gamma$ are two distinct edges from the boundary graph, we may ‘sew’ the surface



(A) The graph Γ' , with chosen edges $e_1, e_2 \in E_{\Gamma'}$. (B) The graph with edges e_1 and e_2 identified with opposite orientation. (C) Γ is the graph Γ' with edges e_1 and e_2 identified, and the identified edges removed along with any isolated vertices.

together along e_1 and e_2 to form a new surface

$$\Sigma' := \frac{\Sigma}{e_1 \sim e_2},$$

as pictured in Fig. 19. In this section, we describe the analogous procedure for the corresponding morphisms of Manin pairs, Eq. (4.1).

First, note that

$$\mathfrak{l}_{sew} := \{((\xi, \eta); (\eta, \xi)) \mid \xi, \eta \in \mathfrak{g}\} \subseteq (\bar{\mathfrak{g}} \oplus \mathfrak{g}) \bigoplus (\bar{\mathfrak{g}} \oplus \mathfrak{g})$$

is a Lagrangian subalgebra. Since G is connected, the \mathfrak{l}_{sew} -orbit through the identity of $G \times G$ is

$$G_{\Delta}^{\natural} := \{(g, g^{-1}) \mid g \in G\}.$$

Definition 4.1. Suppose that

$$((\bar{\mathfrak{g}} \oplus \mathfrak{g}) \bigoplus (\bar{\mathfrak{g}} \oplus \mathfrak{g}) \bigoplus \mathfrak{d}') \times (G \times G \times N)$$

is the product of two Cartan Courant algebroids with an action Courant algebroid, $\mathfrak{d}' \times N$, and that $\mathfrak{h} \subseteq (\bar{\mathfrak{g}} \oplus \mathfrak{g}) \bigoplus (\bar{\mathfrak{g}} \oplus \mathfrak{g}) \bigoplus \mathfrak{d}'$ is a Lagrangian subalgebra. Further suppose that

$$(4.2) \quad R : (\mathbb{T}, T)M \dashrightarrow ((\bar{\mathfrak{g}} \oplus \mathfrak{g}) \bigoplus (\bar{\mathfrak{g}} \oplus \mathfrak{g}) \bigoplus \mathfrak{d}', \mathfrak{h}) \times (G \times G \times N)$$

is a morphism of Manin pairs.

Let $\mathfrak{c}_{sew} = \mathfrak{l}_{sew} \oplus \mathfrak{d}'$ and let

$$S_{sew} = G_{\Delta}^{\natural} \times N \subseteq G \times G \times N.$$

If $(\mathfrak{c}_{sew}, S_{sew})$ is reduction data for (4.2), then the reduction,

$$R_{\mathfrak{c}_{sew}, S_{sew}} : (\mathbb{T}, T)M_{\mathfrak{c}_{sew}, S_{sew}} \dashrightarrow (\mathfrak{d}', \mathfrak{h}_{\mathfrak{c}_{sew}, S_{sew}}) \times N$$

is called the *sewing* of (4.2).

In a typical example of sewing, Γ' will be a permutation graph, $e_1, e_2 \subseteq E_{\Gamma'}$ will be two (distinct) edges, and

$$R : (\mathbb{T}, T)M \dashrightarrow ((\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_{\Gamma'}}, \mathfrak{g}_{\Gamma'}) \times G^{E_{\Gamma'}}$$

will be a morphism of Manin pairs.

We let Γ denote the graph with edge set $E_\Gamma := E_{\Gamma'} \setminus \{e_1, e_2\}$ and vertex set

$$V_\Gamma := \frac{\text{out}(E_\Gamma)}{\text{in}(e_1) \sim \text{out}(e_2) \text{ and } \text{in}(e_2) \sim \text{out}(e_1)},$$

as pictured in Fig. 20c.

Let

$$\begin{aligned} \mathfrak{c}_{sew}^{e_1, e_2} &:= \{(\xi, \eta) \in (\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_{\Gamma'}} \mid (\xi, \eta)_{\{e_1, e_2\}} \in \mathfrak{l}_{sew}\} \\ S_{sew}^{e_1, e_2} &:= \{g \in G^{E_{\Gamma'}} \mid g_{\{e_1, e_2\}} \in G_\Delta^{\natural}\} \end{aligned}$$

as in Definition 4.1. Here we are using the notation $(\xi, \eta)_{\{e_1, e_2\}} := ((\xi_{e_1}, \eta_{e_1}); (\xi_{e_2}, \eta_{e_2}))$, as described in Section 1.1. The morphism of Manin pairs

$$R_{\mathfrak{c}_{sew}^{e_1, e_2}, S_{sew}^{e_1, e_2}} : (\mathbb{T}, T)M_{\mathfrak{c}_{sew}^{e_1, e_2}, S_{sew}^{e_1, e_2}} \dashrightarrow ((\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_\Gamma}, \mathfrak{g}_\Gamma) \times G^{E_\Gamma}$$

is called the result of *sewing edges e_1 and e_2 together*. Here we have used the following lemma to simplify the right hand side:

Lemma 4.1. *The Lie subalgebras $\mathfrak{g}_\Gamma \subseteq (\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_\Gamma}$ and $(\mathfrak{g}_{\Gamma'})_{\mathfrak{c}_{sew}^{e_1, e_2}} \subseteq (\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_{\Gamma'}}$ are equal.*

Proof. Let us recall that

$$\mathfrak{g}_{\Gamma'} = (\text{in} \oplus \text{out})^! \mathfrak{g}_\Delta^{V_{\Gamma'}} \cong \mathfrak{g}^{V_{\Gamma'}}$$

and

$$(\mathfrak{g}_{\Gamma'})_{\mathfrak{c}_{sew}^{e_1, e_2}} = \mathfrak{g}_{\Gamma'} \cap \mathfrak{c}_{sew}^{e_1, e_2} / \mathfrak{g}_{\Gamma'} \cap (\mathfrak{c}_{sew}^{e_1, e_2})^\perp.$$

By definition of $\mathfrak{c}_{sew}^{e_1, e_2}$, $\mathfrak{g}_{\Gamma'} \cap \mathfrak{c}_{sew}^{e_1, e_2}$ is the subalgebra of $\mathfrak{g}^{V_{\Gamma'}}$ where the components corresponding to identified vertices are equal. After we divide by $\mathfrak{g}_{\Gamma'} \cap (\mathfrak{c}_{sew}^{e_1, e_2})^\perp$ we obtain the Lie algebra $\mathfrak{g}_\Gamma = \mathfrak{g}^{V_\Gamma}$. \square

The morphism of Manin pairs (4.1) which we assign to the surface Σ will satisfy the following sewing property:

Sewing property. Let (Σ, V) be obtained out of (Σ', V') by sewing edges e_1 and e_2 . There is a canonical isomorphism

$$(4.3) \quad \mathcal{M}_{\Sigma, V} \xrightarrow{\cong} (\mathcal{M}_{\Sigma', V'})_{\mathfrak{c}_{sew}^{e_1, e_2}, S_{sew}^{e_1, e_2}},$$

which identifies the following two morphisms of Manin pairs:

$$R_{\Sigma, V} : (\mathbb{T}, T)\mathcal{M}_{\Sigma, V} \dashrightarrow ((\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_\Gamma}, \mathfrak{g}_\Gamma) \times G^{E_\Gamma},$$

and

$$(R_{\Sigma', V'})_{\mathfrak{c}_{sew}^{e_1, e_2}, S_{sew}^{e_1, e_2}} : (\mathbb{T}, T)\mathcal{M}_{\Sigma, V} \dashrightarrow ((\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_\Gamma}, \mathfrak{g}_\Gamma) \times G^{E_\Gamma}.$$

Notice that

$$\mu^{-1}(S_{sew}^{e_1, e_2}) = \{f \in \mathcal{M}_{\Sigma', V'}(G) \mid f(e_1)f(e_2) = 1\}$$

and that

$$\mu^{-1}(S_{sew}^{e_1, e_2}) / \mathfrak{g}^{V'} \cap (\mathfrak{c}_{sew}^{e_1, e_2})^\perp = \mathcal{M}_{\Sigma, V}(G).$$

Thus, the isomorphism 4.3 is clear. The non-trivial part of the statement is the behaviour of the $R_{\Sigma, V}$'s

4.1.1. *Commutativity of sewing.* Suppose $\{e_1, e_2\}, \{e'_1, e'_2\} \subseteq E_\Gamma$ are two distinct pairs of distinct edges (i.e. $\{e_1, e_2\} \cap \{e'_1, e'_2\} = \emptyset$), and let $i : \{e_1, e_2\} \rightarrow E_\Gamma$ and $i' : \{e'_1, e'_2\} \rightarrow E_\Gamma$ denote the inclusions. We may form the graph Γ'' , with edge set $E_{\Gamma''} := E_\Gamma \setminus \{e_1, e_2, e'_1, e'_2\}$ and vertex set

$$V_{\Gamma''} := \frac{\text{out}(E_{\Gamma''})}{\text{in}(e_1) \sim \text{out}(e_2), \text{in}(e_2) \sim \text{out}(e_1), \text{ and } \text{in}(e'_1) \sim \text{out}(e'_2), \text{in}(e'_2) \sim \text{out}(e'_1)}.$$

It is the graph obtained from Γ by identifying the oppositely directed edges $e_1 \sim e_2$ and $e'_1 \sim e'_2$, and then deleting these newly identified edges and any isolated vertices.

Let

$$\begin{aligned} \mathfrak{c}_1 &:= \mathfrak{c}_{sew}^{e_1, e_2} = i(\mathfrak{l}_{sew}) \oplus (\bar{\mathfrak{g}} \oplus \mathfrak{g})^{\{e'_1, e'_2\}} \oplus (\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_{\Gamma''}}, \\ \mathfrak{c}_2 &:= \mathfrak{c}_{sew}^{e'_1, e'_2} = (\bar{\mathfrak{g}} \oplus \mathfrak{g})^{\{e_1, e_2\}} \oplus i'(\mathfrak{l}_{sew}) \oplus (\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_{\Gamma''}}, \\ \mathfrak{c}_{1,2} &:= \mathfrak{c}_1 \cap \mathfrak{c}_2 = i(\mathfrak{l}_{sew}) \oplus i'(\mathfrak{l}_{sew}) \oplus (\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_{\Gamma''}}, \end{aligned}$$

and

$$\begin{aligned} S_1 &:= S_{sew}^{e_1, e_2} = i(G_\Delta^\natural) \times G^{\{e'_1, e'_2\}} \times G^{E_{\Gamma''}}, \\ S_2 &:= S_{sew}^{e'_1, e'_2} = G^{\{e_1, e_2\}} \times i'(G_\Delta^\natural) \times G^{E_{\Gamma''}}, \\ S_{1,2} &:= S_1 \cap S_2 = i(G_\Delta^\natural) \times i'(G_\Delta^\natural) \times G^{E_{\Gamma''}}, \end{aligned}$$

If the pairs $(\mathfrak{c}_{sew}^{e_1, e_2}, S_{sew}^{e_1, e_2})$, $(\mathfrak{c}_{sew}^{e'_1, e'_2}, S_{sew}^{e'_1, e'_2})$ and $(\mathfrak{c}_1 \cap \mathfrak{c}_2, S_1 \cap S_2)$ all form reductive data for R , then the assumptions of Corollary 3.1 are satisfied. Therefore,

$$(R_{\mathfrak{c}_{sew}^{e_1, e_2}, S_{sew}^{e_1, e_2}})_{\mathfrak{c}_{sew}^{e'_1, e'_2}, S_{sew}^{e'_1, e'_2}} = R_{\mathfrak{c}_{1,2}, S_{1,2}} = (R_{\mathfrak{c}_{sew}^{e'_1, e'_2}, S_{sew}^{e'_1, e'_2}})_{\mathfrak{c}_{sew}^{e_1, e_2}, S_{sew}^{e_1, e_2}}$$

as morphisms of Manin pairs

$$(\mathbb{T}, T)M'' \dashrightarrow ((\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_{\Gamma''}}, \mathfrak{g}_{\Gamma''}) \times G^{E_{\Gamma''}},$$

where $M'' = M_{\mathfrak{c}_{1,2}, S_{1,2}}$. That is to say, it makes no difference in which order we sew pairs of edges: the results are all naturally isomorphic.

4.2. The quasi-Hamiltonian structure on $\mathcal{M}_{\Sigma, V}(G)$.

Theorem 4.1. *There is a unique way to assign to every marked surface (Σ, V) an exact morphism of Manin pairs*


$$R_\Sigma : (\mathbb{T}, T)\mathcal{M}_{\Sigma, V} \dashrightarrow ((\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_\Gamma}, \mathfrak{g}_\Gamma) \times G^{E_\Gamma}$$

supported on the graph of $\mu : \mathcal{M}_{\Sigma, V} \rightarrow G^{E_\Gamma}$ such that the assignment satisfies the sewing property.


Proof. If Σ' is a disjoint union of disks then $R_{\Sigma'}$ exists and is unique by Proposition 4.1.

Suppose that Σ is triangulated and $V = \mathcal{T}^0 \cap \partial\Sigma$. Let Σ' be the disjoint union of the triangles and V' the set of its vertices. Then, by sewing the respective edges of $\mathcal{M}_{\Sigma', V'}(G)$ according to the triangulation, we get an exact morphism of Manin pairs

$$R_{\mathcal{T}} : (\mathbb{T}, T)\mathcal{M}_{\Sigma, V} \dashrightarrow ((\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_\Gamma}, \mathfrak{g}_\Gamma) \times G^{E_\Gamma}.$$



(D) A triangulation of the 3-gon.



(E) A Triangulation of the 4-gon.

FIGURE 20. Triangulations of 3 and 4-gons. The dashed lines indicate internal edges of the triangulation along which we sew.

By commutativity of sewing these morphisms satisfy the sewing property. It remains to show that they are independent of the triangulation \mathcal{T} .

Thus, we need to prove that $R_{\mathcal{T}}$ is invariant under Pachner moves applied to \mathcal{T} . Pachner moves are, however, simply changes of triangulations of a polygon (either a triangle or a square, see Theorem 4.1). The independence thus follows from Proposition 4.1. \square

5. POISSON STRUCTURES ON THE MODULI SPACE OF FLAT CONNECTIONS


Suppose G is a (connected) Lie group with Lie algebra \mathfrak{g} and $s \in S^2(\mathfrak{g})^G$ is a G -invariant symmetric 2-tensor. We let \mathfrak{d} denote the Drinfel'd double of \mathfrak{g} , as in Example 1.8. Suppose now that Γ is a permutation graph. Then $\mathfrak{d}^{V_{\Gamma}}$ acts on $G^{E_{\Gamma}}$ with coisotropic stabilizers, as follows: $\mathfrak{d}^{V_{\Gamma}}$ acts on the $e \in E_{\Gamma}$ -th factor $G^{E_{\Gamma}}$ via the vector field

$$\rho(\xi)_e = -s(\xi_{\text{in}(e)})^R + \mathfrak{t}(\xi_{\text{out}(e)})^L, \quad \xi \in \mathfrak{d}^{V_{\Gamma}},$$


(cf. Fig. 21b). Thus

$$\mathfrak{g}^{V_{\Gamma}} \times G^{E_{\Gamma}} \subseteq \mathfrak{d}^{V_{\Gamma}} \times G^{E_{\Gamma}}$$

is a Dirac structure.



(A) Here $\text{in}(e) = v = \text{out}(e')$, where $v \in V_{\Gamma}$ is the vertex and $e, e' \in E_{\Gamma}$ are edges.



(B) The element $\xi_{(\cdot)} \in \mathfrak{d}^{V_{\Gamma}}$ acts diagonally by ξ_v at the vertex v .

Suppose (Σ, V) is a marked surface (where we now allow components of $\partial\Sigma$ to intersect V trivially), with boundary graph Γ . In this section, we prove there exists a natural quasi-Hamiltonian $(\mathfrak{d}, \mathfrak{g})^{V_\Gamma} \times G^{E_\Gamma}$ structure on $\mathcal{M}_{\Sigma, V}(G)$, i.e a morphism of Manin pairs

$$R_{\Sigma, V} : (\mathbb{T}, T)\mathcal{M}_{\Sigma, V}(G) \dashrightarrow (\mathfrak{d}, \mathfrak{g})^{V_\Gamma} \times G^{E_\Gamma}$$

over the map

$$\mu : \mathcal{M}_{\Sigma, V}(G) \rightarrow G^{E_\Gamma}$$

given by the boundary holonomies.

As before, for disjoint unions of polygons the quasi-Hamiltonian structure is uniquely defined.

Proposition 5.1 (union of polygons). *Let Σ be a disjoint union of discs and $V \subset \partial\Sigma$ a finite subset. Then there is a unique exact morphism of Manin pairs*

$$(\mathbb{T}, T)\mathcal{M}_{\Sigma, V} \dashrightarrow ((\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_\Gamma}, \mathfrak{g}_\Gamma) \times G^{E_\Gamma}$$

over μ .

Proof. As in Proposition 4.1, μ is an embedding, so this follows from Example 2.9. \square

5.1. Fusion. Suppose that $P, Q \in V_\Gamma$ are two distinct vertices. The operation of *fusion* at the ordered pair of vertices (P, Q) described in [26] (following [3, 4]), can be understood in terms of a morphism of Manin pairs

$$R_{(P, Q)} : (\mathfrak{d}, \mathfrak{g})^{V_\Gamma} \times G^{E_\Gamma} \dashrightarrow (\mathfrak{d}, \mathfrak{g})^{V_{\Gamma^*}} \times G^{E_{\Gamma^*}}$$

where the graph Γ^* is a permutation graph constructed from Γ , as we shall now explain:

Let $\overleftarrow{e}_P = \text{in}^{-1}(P)$ and $e_{\overleftarrow{P}} := \text{out}^{-1}(P)$ denote the edges entering and exiting P . Similarly, let $\overleftarrow{e}_Q = \text{in}^{-1}(Q)$ and $e_{\overleftarrow{Q}} := \text{out}^{-1}(Q)$ denote the edges entering and exiting Q .

Case 1: $\overleftarrow{e}_Q = e_{\overleftarrow{P}}$: In this case, Γ^* is obtained by discarding the edge $\overleftarrow{e}_Q = e_{\overleftarrow{P}}$ and identifying the vertices P and Q (cf. Fig. 21).

Meanwhile for $(\xi, g) \in \mathfrak{d}^{V_\Gamma} \times G^{E_\Gamma}$ and $(\xi^*, g^*) \in \mathfrak{d}^{V_{\Gamma^*}} \times G^{E_{\Gamma^*}}$

$$(\xi, g) \sim_{R_{(P, Q)}} (\xi^*, g^*)$$

if and only if $g_e^* = g_e$ for every $e \in E_{\Gamma^*}$ and

$$(5.1) \quad \xi_v^* = \begin{cases} \xi_Q \circ \xi_P & \text{if } v \text{ is the vertex obtained by identifying } P \text{ and } Q \\ \xi_v & \text{otherwise,} \end{cases}$$

(in particular, we assume that ξ_Q and ξ_P are composable elements of the Lie groupoid \mathfrak{d}).

Case 2: $\overleftarrow{e}_Q \neq e_{\overleftarrow{P}}$: In this case, Γ^* is obtained by identifying the vertices P and Q and composing the edges $e_{\overleftarrow{P}}$ and \overleftarrow{e}_Q to form a new edge $e_{\overleftarrow{PQ}}$ (cf. Fig. 22).

Meanwhile for $(\xi, g) \in \mathfrak{d}^{V_\Gamma} \times G^{E_\Gamma}$ and $(\xi^*, g^*) \in \mathfrak{d}^{V_{\Gamma^*}} \times G^{E_{\Gamma^*}}$

$$(\xi, g) \sim_{R_{(P, Q)}} (\xi^*, g^*)$$

if and only if

$$g_e^* = \begin{cases} g_{e_{\overline{P}}} g_{\overleftarrow{Q}e} & \text{if } e = e_{\overline{P}\overline{Q}} \\ g_e & \text{otherwise,} \end{cases}$$

while ξ^* and ξ satisfy Eq. (5.1), as before.




FIGURE 21. The graph Γ^* is the permutation graph obtained from Γ by deleting the edge, $\overleftarrow{Q}e_{\overline{P}}$, passing from P to Q , and identifying the vertices P and Q . (The graphs are isomorphic outside the pictured regions.) Heuristically, we have obtained the graph Γ^* by gluing short sections from both ends of $\overleftarrow{Q}e_{\overline{P}}$ together (effectively divorcing it from the graph).

Meanwhile, the Courant morphism, $R_{(P,Q)}$, is defined by composing the corresponding elements labelling the vertices and by forgetting the element which labelled the deleted edge.

Now suppose that M is a quasi-Hamiltonian $(\mathfrak{d}, \mathfrak{g})^{V_\Gamma} \times G^{E_\Gamma}$ -space defined by the morphism of Manin pairs

$$R : (\mathbb{T}, T)M \dashrightarrow (\mathfrak{d}, \mathfrak{g})^{V_\Gamma} \times G^{E_\Gamma}.$$

Then the morphism of Manin pairs

$$R^* := R_{(P,Q)} \circ R : (\mathbb{T}, T)M \dashrightarrow (\mathfrak{d}, \mathfrak{g})^{V_{\Gamma^*}} \times G^{E_{\Gamma^*}}$$

defines a quasi-Hamiltonian $(\mathfrak{d}, \mathfrak{g})^{V_{\Gamma^*}} \times G^{E_{\Gamma^*}}$ -structure on M which we call the *fusion* of R at the ordered pair (P, Q) of vertices.

Remark 5.1 (Associativity of Fusion). Since the Courant morphism $R_{(P,Q)}$ is defined in terms of the groupoid structure on \mathfrak{d} and the group structure on G , it follows that fusion is an associative operation.




FIGURE 22. The graph Γ^* is the permutation graph obtained from Γ by identifying the vertices P and Q and composing the edge entering Q with the edge leaving P . (The graphs are isomorphic outside the pictured regions.) Heuristically, we have obtained the graph Γ^* by gluing a short section of the edges e_P and e_Q together.

Meanwhile, the Courant morphism, $R_{(P,Q)}$, is defined by composing the corresponding elements labelling the edges and vertices.

5.2. The quasi-Hamiltonian structure for quasi-triangular structure Lie algebras. Let (Σ, V) be a marked surface. If we choose an ordered pair (P, Q) of marked points ($P \neq Q \in V$) then the corresponding *fused surface* Σ^* is obtained by gluing a short piece of the arc starting at P with a short piece of the arc ending at Q (so that P and Q get identified). The subset $V^* \subset \partial\Sigma^*$ is obtained from V by identifying P and Q . The map

$$M_{\Sigma^*, V^*}(G) \rightarrow M_{\Sigma, V}(G),$$

coming from the map $(\Sigma, V) \rightarrow (\Sigma^*, V^*)$, is a diffeomorphism.

Theorem 5.1. *There is unique way to assign to every marked surface (Σ, V) a morphism of Manin pairs*

$$R_{\Sigma, V} : (\mathbb{T}, T)\mathcal{M}_{\Sigma, V}(G) \dashrightarrow (\mathfrak{d}, \mathfrak{g})^{V_\Gamma} \times G^{E_\Gamma}$$

supported on the graph of $\mu : \mathcal{M}_{\Sigma, V}(G) \rightarrow G^{E_\Gamma}$ such that if (Σ^, V^*) is obtained from (Σ, V) by fusion, then R_{Σ^*, V^*} is obtained from $R_{\Sigma, V}$ by the corresponding fusion.*

We defer the proof until the next section.

Remark 5.2. When $s \in S^2(\mathfrak{g})^G$ is non-degenerate (i.e. \mathfrak{g} is quadratic), then $\mathfrak{d} = \mathfrak{g} \oplus \bar{\mathfrak{g}}$ is the pair groupoid and

$$(\text{in} \oplus \text{out})^!(s \oplus t) : \mathfrak{d}^{V_\Gamma} \rightarrow (\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_\Gamma}$$

is an isomorphism. In this case, it is not difficult to convince oneself that the morphisms of Manin pairs

$$R_{\Sigma, V} : (\mathbb{T}, T)\mathcal{M}_{\Sigma, V}(G) \dashrightarrow ((\bar{\mathfrak{g}} \oplus \mathfrak{g})^{E_\Gamma}, \mathfrak{g}_\Delta^{V_\Gamma}) \times G^{E_\Gamma}$$

described in Theorem 5.1 satisfy the sewing property. That is they are precisely the ones described in Theorem 4.1.

5.2.1. *quasi-Hamiltonian $(\mathfrak{d}, \mathfrak{g})^{V_\Gamma} \times G^{E_\Gamma}$ -manifolds and quasi-Poisson geometry.* Using the material in Section 3.2, we intend to relate quasi-Hamiltonian $(\mathfrak{d}, \mathfrak{g})^{V_\Gamma} \times G^{E_\Gamma}$ -spaces to the quasi-Poisson spaces studied in [3, 26]. A canonical choice of complement to $\mathfrak{g} \subseteq \mathfrak{d}$ is $\mathfrak{g}_\Delta^* := (\mathfrak{s} + \mathfrak{t})^*(\mathfrak{g}^*) \subseteq \mathfrak{d}$, explicitly

$$\mathfrak{g}_\Delta^* = \left\{ \left(\alpha - \frac{1}{2}s(\alpha, \cdot) \right) \mid \alpha \in \mathfrak{p} \subseteq \mathfrak{d} \right\}.$$

Similarly, $\mathfrak{k} = (\mathfrak{g}_\Delta^*)^{V_\Gamma}$ is a canonical choice of complement to $\mathfrak{g}^{V_\Gamma} \subseteq \mathfrak{d}^{V_\Gamma}$.

Let $\bar{\sigma}_\Gamma : V_\Gamma \rightarrow V_\Gamma$ be the permutation given by walking along the graph Γ *against* the direction of each edge. Suppose that (M, ρ, π) is a quasi-Poisson G^{V_Γ} -manifold, in the sense of [3, 26] and $\mu : M \rightarrow G^{V_\Gamma}$ is a $\bar{\sigma}_\Gamma^{-1}$ -twisted moment map in the sense of [26]. Let $\tilde{\mu} : M \rightarrow G^{E_\Gamma}$ be defined by

$$\tilde{\mu}(m)_e = (\mu(m)_{\text{in}(e)})^{-1}, \quad m \in M, e \in E_\Gamma.$$

Proposition 5.2. *There exists a unique morphism of Manin pairs*

$$R : (\mathbb{T}, T)M \dashrightarrow (\mathfrak{d}^{V_\Gamma}, \mathfrak{g}^{V_\Gamma}) \times G^{E_\Gamma}$$

over the map $\tilde{\mu}$ which is compatible with the \mathfrak{g}^{V_Γ} action on M and such that $\mathfrak{k} \circ R = \text{gr}(\pi^\#)$.

Moreover, the converse holds whenever the action of \mathfrak{g}^{V_Γ} on M integrates to an action of G^{V_Γ} .

Proof. This follows from a direct application of [11, Proposition 3.5]. (cf. Section 3.2). \square

In this sense there is a one-to-one correspondence between quasi-Poisson G^{V_Γ} -manifolds with $\bar{\sigma}_\Gamma^{-1}$ -twisted moment maps and quasi-Hamiltonian $(\mathfrak{d}^{V_\Gamma}, \mathfrak{g}^{V_\Gamma}) \times G^{E_\Gamma}$ -manifolds.

Proposition 5.3. *Suppose (M, ρ, π) is a quasi-Poisson G^{V_Γ} -manifold with $\bar{\sigma}_\Gamma^{-1}$ -twisted moment map, and let*

$$R : (\mathbb{T}, T)M \dashrightarrow (\mathfrak{d}^{V_\Gamma}, \mathfrak{g}^{V_\Gamma}) \times G^{E_\Gamma}$$

be the corresponding quasi-Hamiltonian $(\mathfrak{d}^{V_\Gamma}, \mathfrak{g}^{V_\Gamma}) \times G^{E_\Gamma}$ -structure on M . Let R^* denote the fusion of R at the ordered pair of vertices $(P, Q) \subset V_\Gamma$. Then the bivector field for the quasi-Poisson $G^{V_\Gamma^*}$ -structure corresponding to R^* is

$$\pi^* := \pi + \rho(\tau),$$

where $\tau \in \wedge^2(\mathfrak{g}^{V_\Gamma})$ is the insertion of $\psi \in \wedge^2(\mathfrak{g}^P \oplus \mathfrak{g}^Q)$,

$$\psi = \frac{1}{2} \sum_{i,j} s^{ij} (\xi_i, 0) \wedge (0, \xi_j)$$

at the P, Q -th factors. Here $s = \sum_{i,j} s^{ij} \xi_i \otimes \xi_j$ in some basis ξ_i of \mathfrak{g} .

Thus, (up to a sign difference) fusion in the sense of Section 5.1 is precisely the same as fusion in the sense of [3, 26].

Proof. Let $\mathfrak{k}_* = (\mathfrak{g}_\Delta^*)^{V_{\Gamma^*}}$. A straightforward computation shows that

$$\mathfrak{k}_* \circ R_{(P,Q)} = \{(\xi + \tau^\sharp(\xi)) \mid \xi \in \mathfrak{k}\}.$$

By definition, $\text{gr}((\pi^*)^\sharp) = \mathfrak{k}_* \circ R^*$. Thus, we see from

$$\mathfrak{k}_* \circ R_{(P,Q)} \circ R = \text{gr}(\pi) + \text{gr}(\rho(\tau))$$

that $\pi^* = \pi + \rho(\tau)$. \square

Remark 5.3 (Sign differences). In [26], the bivector field resulting from fusing the ordered pair (P, Q) is defined to be $\pi^* = \pi - \rho(\tau)$. This difference is essentially due to the fact that we orient our boundary graph to agree with the orientation of $\partial\Sigma$, whereas the opposite convention is used in [26].

Proof of Theorem 5.1. Propositions 5.2 and 5.3 show that it suffices to prove the equivalent statement for quasi-Poisson G^{V_Γ} -structures. However, [26, Theorem 2] and [26, Theorem 4] shows there exists a unique quasi-Poisson G^{V_Γ} -structure on $\mathcal{M}_{\Sigma,V}(G)$ with $\overline{\sigma}_\Gamma^!$ -twisted moment map which is compatible with fusion (notice that Proposition 5.1 implies the first two properties of [26, Theorem 2] are automatically satisfied). \square

Corollary 5.1. *The proof of Theorem 5.1 also shows that the bivector field on $\mathcal{M}_{\Sigma,V}$ is given by Eq. (1.11).*

APPENDIX A. PROOFS OF REDUCTION THEOREMS

Before proving Theorems 3.1 and 3.2 and Proposition 3.1, we first establish some lemmas.

Lemma A.1. *Suppose that $(\eta; Z) \in R_{\mathfrak{c},S}$, where $\eta \in \mathfrak{d}_{\mathfrak{c}}$, $Z \in TM_{\mathfrak{c},S}$ and $R_{\mathfrak{c},S}$ is as in Theorem 3.1. Then*

- (1) $\eta \in \mathfrak{h}_{\mathfrak{c}}$, and
- (2) $\eta = 0$ only if $Z = 0$.

Proof. Let $\xi \in \mathfrak{d}$, $X \in TM$ and $\alpha \in T^*M$ be chosen so that $(\xi; X + \alpha) \in R$ and

$$(\xi; X + \alpha) \sim_{(R_2 \times R_1)} (\eta; Z).$$

Since $R_1 = R_{q_M} \circ R_{i_M}^\top$, it follows that $\alpha \in \text{ann}(T\mu^{-1}S)$. Consequently, there exists $\tilde{\alpha} \in \text{ann}(TS)$ such that $\alpha = \mu^*\tilde{\alpha}$. Since S is \mathfrak{c} invariant,

$$\zeta_\alpha := \mathbf{a}^*\tilde{\alpha} \in \mathfrak{c}^\perp.$$

Moreover, since R is supported on the graph of μ , $(\zeta_\alpha, \alpha) = \mathbf{a}^*(-\tilde{\alpha}, \mu^*\tilde{\alpha}) \in R$. Thus

$$(A.1) \quad (\xi - \zeta_\alpha; X) \in R$$

and

$$(A.2) \quad (\xi - \zeta_\alpha; X) \sim_{(R_2 \times R_1)} (\eta; Z).$$

Since (3.1) is a morphism of Manin pairs, axiom (m1) of Definition 2.5 implies that

$$(A.3) \quad \xi - \zeta_\alpha \in \mathfrak{h}.$$

Thus Eq. (A.2) implies that $\eta \in \mathfrak{h}_{\mathfrak{c}}$, establishing the first claim.

Next, suppose $\eta = 0$, then Eq. (A.2) implies that $\xi - \zeta_{\alpha} \in \mathfrak{c}^{\perp}$ in addition to Eq. (A.3). That is $\xi - \zeta_{\alpha} \in \mathfrak{c}^{\perp} \cap \mathfrak{h}$, and hence Eqs. (2.8) and (A.1) imply that X is tangent to the $\mathfrak{c}^{\perp} \cap \mathfrak{h}$. Therefore $Z = q_M(X) = 0$, establishing the second claim. \square

Lemma A.2. *Under the assumptions of Theorem 3.1, the Courant relation*

$$R_2 \times R_1 : (\mathfrak{d} \times N) \times \overline{\mathbb{T}M} \dashrightarrow (\mathfrak{d}_{\mathfrak{c}} \times N_{\mathfrak{c},S}) \times \overline{\mathbb{T}M_{\mathfrak{c},S}}$$

composes cleanly with the Dirac structure $R \subseteq (\mathfrak{d} \times N) \times \overline{\mathbb{T}M}$.

Moreover, the composition

$$R_{\mathfrak{c},S} := R_2 \circ R \circ R_1^{\top}$$

is a well defined subbundle of $(\mathfrak{d}_{\mathfrak{c}} \times N_{\mathfrak{c},S}) \times \overline{\mathbb{T}M_{\mathfrak{c},S}}$.

Proof. We begin by proving that the composition $(R_2 \times R_1) \circ R$ is clean. For this, it is sufficient to show that

- (1) the rank of the intersections $\ker(R_2 \times R_1)^{\perp} \cap R$ and $\ker(R_2 \times R_1) \cap R$ are constant, and
- (2) the composition of the underlying relation of vector bundle bases,

$$(\mathrm{gr}(q_N \times q_M) \circ \mathrm{gr}(i_N \times i_M)^{\top}) \circ \mathrm{gr}(\mu)$$

is clean.

We now show that the rank of $\ker(R_2 \times R_1) \cap R$ is constant. We claim that the sequence

$$(A.4) \quad 0 \rightarrow \mathfrak{h} \cap \mathfrak{c}^{\perp} \xrightarrow{\xi \mapsto (\xi, \rho_R(\xi))} \ker(R_2 \times R_1) \cap R \xrightarrow{(\xi, X + \alpha) \mapsto \alpha} \mathrm{ann}(T\mu^{-1}S) \rightarrow 0$$

is exact, where ρ_R is defined in Eq. (2.8).

First, the second map is surjective: for any $\alpha \in \mathrm{ann}(T\mu^{-1}S)$, let $\tilde{\alpha} \in \mathrm{ann}(TS)$ be chosen so that $\mu^* \tilde{\alpha} = \alpha$. Since S is \mathfrak{c} invariant,

$$\zeta_{\alpha} := \mathbf{a}^* \tilde{\alpha} \in \mathfrak{c}^{\perp}.$$

Moreover, since R is supported on the graph of μ ,

$$(\zeta_{\alpha}, \alpha) = \mathbf{a}^*(-\tilde{\alpha}, \mu^* \tilde{\alpha}) \in \ker(R_2 \times R_1) \cap R.$$

Next, we prove exactness at $\ker(R_2 \times R_1) \cap R$. Suppose $(\xi, X) \in \ker(R_2 \times R_1) \cap R$. Since Eq. (3.1) is a morphism of Manin pairs, $\xi \in \mathfrak{h}$ and $X = \rho_R(\xi - \zeta_{\alpha})$. Since $\xi \in \ker(R_2) = \mathfrak{c}^{\perp}$, we conclude that $\xi \in \mathfrak{h} \cap \mathfrak{c}^{\perp}$.

This shows that the sequence Eq. (A.4) is exact. Since $\mathfrak{h} \cap \mathfrak{c}^{\perp}$ and $\mathrm{ann}(T\mu^{-1}S)$ are both of constant rank, so is $\ker(R_2 \times R_1) \cap R$. Consequently $(\ker(R_2 \times R_1) \cap R)^{\perp} = \ker(R_2 \times R_1)^{\perp} + R$ is also of constant rank, and thus so is $\ker(R_2 \times R_1)^{\perp} \cap R$.

Next, we need to show that the following composition of relations is clean:

$$\begin{array}{ccc}
 M & \xrightarrow{\mu} & N \\
 i_M \uparrow & & \uparrow i_N \\
 \tilde{S} & & S \\
 q_M \downarrow & & \downarrow q_N \\
 M_{\mathfrak{c},S} & \xrightarrow{\tilde{\mu}} & N_{\mathfrak{c},S}
 \end{array}$$

where, for brevity, we have introduced the notation $\tilde{S} := \mu^{-1}(S)$. Since $\text{gr}(\mu)$ intersects $S \times M$ cleanly, the composition $\text{gr}(\mu|_{\tilde{S}}) = \text{gr}(i_N)^\top \circ \text{gr}(\mu \circ i_M)$ is clean

$$\begin{array}{ccc}
 M & \xrightarrow{\mu} & N \\
 i_M \uparrow & & \uparrow i_N \\
 \tilde{S} & \xrightarrow{\mu|_{\tilde{S}}} & S \\
 q_M \downarrow & & \downarrow q_N \\
 M_{\mathfrak{c},S} & \xrightarrow{\tilde{\mu}} & N_{\mathfrak{c},S}
 \end{array}$$

Next,

$$\text{gr}(q_N \circ \mu|_{\tilde{S}}) \times \text{gr}(q_M)^\top \subseteq N_{\mathfrak{c},S} \times \tilde{S} \times \tilde{S} \times M_{\mathfrak{c},S}$$

intersects $N_{\mathfrak{c},S} \times \tilde{S} \times M_{\mathfrak{c},S}$ transversely, since q_M and $q_N \circ \mu|_{\tilde{S}}$ are both maps. Moreover, since $M_{\mathfrak{c},S}$ is the set of $\mathfrak{h} \cap \mathfrak{c}^\perp$ orbits in \tilde{S} , while $N_{\mathfrak{c},S}$ is the set of \mathfrak{c}^\perp orbits in S , the projection

$$\left(\text{gr}(q_N \circ \mu|_{\tilde{S}}) \times \text{gr}(q_M)^\top \right) \cap \left(N_{\mathfrak{c},S} \times \tilde{S} \times M_{\mathfrak{c},S} \right) \rightarrow \text{gr}(\tilde{\mu})$$

is a surjective submersion. Thus, by definition, $\text{gr}(q_N \circ \mu|_{\tilde{S}})$ composes cleanly with $\text{gr}(q_M)^\top$.

It follows that $R_2 \times R_1$ composes cleanly with R .

Finally, since $\{(\xi; \rho_R(\xi)) \mid \xi \in \mathfrak{h} \cap \mathfrak{c}^\perp\} \subseteq R$, it follows that R is $\mathfrak{h} \cap \mathfrak{c}^\perp$ invariant. Hence $R_{\mathfrak{c},S} := R_2 \circ R \circ R_1^\top$ is a well defined subbundle of

$$(\mathfrak{d}_{\mathfrak{c}} \times D_{\mathfrak{c}}/G_{\mathfrak{c}}) \times \overline{\mathbb{T}Q}.$$

□

We are now ready to prove Theorems 3.1 and 3.2.

Proof of Theorem 3.1. Since $N_{\mathfrak{c},S}$ is the space of \mathfrak{c}^\perp -orbits of S , while $M_{\mathfrak{c},S}$ is the space of $\mathfrak{c}^\perp \cap \mathfrak{h}$ orbits of $\mu^{-1}(S)$, the \mathfrak{h} -equivariant map

$$\mu : \mu^{-1}(S) \rightarrow S$$

descends to a define a unique map

$$\tilde{\mu} : M_{\mathfrak{c},S} \rightarrow N_{\mathfrak{c},S}.$$

The composition $R_{\mathfrak{c},S} := R_2 \circ R \circ R_1^\top$ is supported on the graph of $\tilde{\mu}$. Thus Lemma A.2 and Proposition 2.1 show that

$$R_{\mathfrak{c},S} : \mathbb{T}M_{\mathfrak{c},S} \dashrightarrow \mathfrak{d}_{\mathfrak{c}} \times N_{\mathfrak{c},S}$$

is a Courant morphism.

Finally, Lemma A.1 proves that (3.2) satisfies the defining conditions for a morphism of Manin pairs. \square

Proof of Theorem 3.2. We need to show that Eq. (3.2) is a exact morphism of Manin pairs. We do this by first showing that $\mathfrak{d}_{\mathfrak{c}} \times N_{\mathfrak{c},S}$ is an exact Courant algebroid along the image of $\mu(M) \cap S$, and next by showing that the anchor maps $R_{\mathfrak{c},S}$ surjectively onto $T\text{gr}(\mu_{\mathfrak{c},S} : M_{\mathfrak{c},S} \rightarrow N_{\mathfrak{c},S})$.

The fact that $\mathfrak{d}_{\mathfrak{c}} \times N_{\mathfrak{c},S}$ is exact follows from [10, Theorem 3.3], but we include a proof here anyways. We must show that

$$(A.5) \quad 0 \rightarrow T^*N_{\mathfrak{c},S} \xrightarrow{\mathfrak{a}^*} \mathfrak{d}_{\mathfrak{c}} \times N_{\mathfrak{c},S} \xrightarrow{\mathfrak{a}} TN_{\mathfrak{c},S} \rightarrow 0$$

is an exact sequence. By assumption, \mathfrak{c} acts transitively on S . Thus $\mathfrak{d}_{\mathfrak{c}} := \mathfrak{c}/\mathfrak{c}^\perp$ acts transitively on $N_{\mathfrak{c},S} := S/\mathfrak{c}^\perp$. It follows that the sequence (A.5) is exact at $TN_{\mathfrak{c},S}$, and hence, by duality, also at $T^*N_{\mathfrak{c},S}$.

Next, Eq. (A.5) is exact at $\mathfrak{d}_{\mathfrak{c}} \times N_{\mathfrak{c},S}$ if and only if $\ker(\mathfrak{a})$ is isotropic. This, in turn, holds if and only if $\mathfrak{c} \cap (\ker(\mathfrak{a}_{\mathfrak{d}}) + \mathfrak{c}^\perp)$ is isotropic, where $\mathfrak{a}_{\mathfrak{d}}$ denotes the anchor map for $\mathfrak{d} \times N$. But

$$(\mathfrak{c} \cap (\ker(\mathfrak{a}_{\mathfrak{d}}) + \mathfrak{c}^\perp))^\perp = \mathfrak{c}^\perp + \text{Im}(\mathfrak{a}_{\mathfrak{d}}^*) \cap \mathfrak{c}^\perp = \mathfrak{c} \cap (\text{Im}(\mathfrak{a}_{\mathfrak{d}}^*) + \mathfrak{c}^\perp).$$

Therefore, (A.5) is exact at $TN_{\mathfrak{c},S}$ if $\text{Im}(\mathfrak{a}_{\mathfrak{d}}^*) = \ker(\mathfrak{a}_{\mathfrak{d}})$, which holds whenever $\mathfrak{d} \times N$ is exact (as it is along $\mu(M)$).

Next we need to show that the anchor maps $R_{\mathfrak{c},S}$ surjectively onto $T\text{gr}(\mu_{\mathfrak{c},S} : M_{\mathfrak{c},S} \rightarrow N_{\mathfrak{c},S})$. More precisely, for any $Z \in TM_{\mathfrak{c},S}$, we must show there exists $\eta \in \mathfrak{d}_{\mathfrak{c}}$ and $\gamma \in T^*M_{\mathfrak{c},S}$ such that

$$(\eta, Z + \gamma) \in R_{\mathfrak{c},S}.$$

Let $X \in T\mu^{-1}(S)$ be chosen so that it maps to Z under the quotient map $q_M : \mu^{-1}(S) \rightarrow M_{\mathfrak{c},S}$. Since Eq. (3.1) is exact, there exists $\alpha \in T^*M$ and $\xi \in \mathfrak{d}$ such that $(\xi, X + \alpha) \in R$. Now R is supported on the graph of μ , so $\mathfrak{a}_{\mathfrak{d}}(\xi) = d\mu(X) \in TS$. Since \mathfrak{c} acts transitively on S , we must have $\xi \in \mathfrak{c} + \ker(\mathfrak{a}_{\mathfrak{d}})$. Since $\mathfrak{d} \times N$ is exact, $\ker(\mathfrak{a}_{\mathfrak{d}}) = \text{Im}(\mathfrak{a}_{\mathfrak{d}}^*)$. Thus there exists $\xi' \in \mathfrak{c}$ and $\beta \in T^*N$ such that $\xi = \xi' + \mathfrak{a}_{\mathfrak{d}}^*\beta$. Since R is supported on the graph of μ , $(\mathfrak{a}_{\mathfrak{d}}^*\beta, \mu^*\beta) \in R$, and thus

$$(\xi', X + \alpha - \mu^*\beta) \in R.$$

Since R is Lagrangian, pairing this element with $(\zeta, \rho_R(\zeta)) \in R$, where $\zeta \in \mathfrak{h} \cap \mathfrak{c}^\perp$, we see that

$$0 = \langle \xi', \zeta \rangle = \langle \alpha - \mu^* \beta, \rho_R(\zeta) \rangle.$$

Since $\zeta \in \mathfrak{h} \cap \mathfrak{c}^\perp$ was arbitrary, this shows that $\alpha - \mu^* \beta = q_M^* \gamma$ for some $\gamma \in T^*M_{\mathfrak{c},S}$.

Thus, we have shown that

$$(\xi' + \mathfrak{c}^\perp, Z + \gamma) \in R_{\mathfrak{c},S},$$

where $\xi' + \mathfrak{c}^\perp$ is the image of ξ' under the quotient map $\mathfrak{c} \rightarrow \mathfrak{c}/\mathfrak{c}^\perp$. Since $Z \in TM_{\mathfrak{c},S}$ was arbitrary, we may conclude that Eq. (3.2) is exact. \square

Proof of Proposition 3.1. First we show that the 2-form $\omega_\theta := i_M^* \omega - \mu^* \langle \theta, \vartheta_s - \frac{1}{2} s \circ \mathbf{a}(\theta) \rangle \in \Omega^2(\mu^{-1}(S))$ is $\mathfrak{h} \cap \mathfrak{c}^\perp$ -invariant and basic. To show invariance, note that (by definition) $(\xi, \rho_R(\xi)) \in R$ for any $\xi \in \mathfrak{h}$, and thus R is \mathfrak{h} -invariant. Since $s : TN \rightarrow \mathfrak{d} \times N$ is an \mathfrak{d} -invariant splitting, it follows that $R_{\mu,\omega}$ and hence $\omega \in \Omega^2(M)$ is \mathfrak{h} -invariant. Additionally, s, \mathbf{a} and ϑ_s are \mathfrak{d} -equivariant, while θ is \mathfrak{c}^\perp -equivariant. Hence $\langle \theta, \vartheta_s - \frac{1}{2} s \circ \mathbf{a}(\theta) \rangle$ is \mathfrak{c}^\perp -invariant. It follows that the sum, ω_θ is $\mathfrak{h} \cap \mathfrak{c}^\perp$ -invariant.

The isomorphism $\mathfrak{d} \times N \rightarrow \mathbb{T}_\gamma N$ is given by $\xi \rightarrow \mathbf{a}(\xi) + s^*(\xi)$, for all $\xi \in \mathfrak{d}$. Thus, for $\xi, \eta \in \mathfrak{d}$, we have

$$\langle \xi, \eta \rangle = \langle \mathbf{a}(\xi) + s^*(\xi), \mathbf{a}(\eta) + s^*(\eta) \rangle = \langle s \circ \mathbf{a}(\xi), \eta \rangle + \langle \xi, s \circ \mathbf{a}(\eta) \rangle.$$

Thus, since \mathfrak{c}^\perp is coisotropic, the assignment $\xi, \eta \rightarrow \langle \xi, s \circ \mathbf{a} \eta \rangle$ defines a skew-symmetric form on \mathfrak{c}^\perp .

Now suppose $\xi \in \mathfrak{c}^\perp$, then

$$\begin{aligned} & \iota_{\mathbf{a}(\xi)} \langle \theta, \vartheta_s - \frac{1}{2} s \circ \mathbf{a}(\theta) \rangle \\ &= \langle \xi, \vartheta_s - \frac{1}{2} s \circ \mathbf{a}(\theta) \rangle - \langle \theta, s \circ \mathbf{a}(\xi) - \frac{1}{2} s \circ \mathbf{a}(\xi) \rangle \\ &= s^* \xi - \frac{1}{2} \langle \xi, s \circ \mathbf{a}(\theta) \rangle - \langle \theta, s \circ \mathbf{a}(\xi) - \frac{1}{2} s \circ \mathbf{a}(\xi) \rangle \\ &= s^* \xi, \end{aligned} \tag{A.6}$$

where the last line follows from the skew-symmetry of the assignment $\xi, \eta \rightarrow \langle \xi, s \circ \mathbf{a}(\eta) \rangle$. Now, for $\xi \in \mathfrak{h}$, we have $\mu_* \rho_R(\xi) = \mathbf{a}(\xi)$, and thus Eq. (3.7) implies that

$$(A.7) \quad \rho_R(\xi) - \iota_{\rho_R(\xi)} \omega + \iota_{\rho_R(\xi)} \mu^* \langle \theta, \vartheta_s - \frac{1}{2} s \circ \mathbf{a}(\theta) \rangle \sim_{R_{\mu,\omega}} \mathbf{a}(\xi) + s^*(\xi), \quad \xi \in \mathfrak{h} \cap \mathfrak{c}^\perp.$$

Since Eq. (3.8) is a morphism of Manin pairs, and since $\mathbf{a}(\xi) + s^*(\xi) \in E$ for any $\xi \in \mathfrak{h}$, it follows that the left hand side of Eq. (A.7) lies in TM . That is, $\iota_{\rho_R(\xi)} \omega_\theta = 0$ for any $\xi \in \mathfrak{h} \cap \mathfrak{c}^\perp$. We conclude that there is a unique 2-form $\tilde{\omega}_\theta \in \Omega^2(M_{\mathfrak{c},S})$ such that $q_M^* \tilde{\omega}_\theta = \omega_\theta$.

We define the Courant relation

$$R_{\mathfrak{c},s,\theta} := \mathbf{gr}(\mathbf{a} \oplus \tilde{s}_\theta^*) \circ R_{\mathfrak{c}} \circ \mathbf{gr}(\mathbf{a} \oplus s^*)^\top : \mathbb{T}_\gamma N \dashrightarrow \mathbb{T}_{\gamma_\theta} N_{\mathfrak{c},S},$$

so that

$$\mathbf{gr}(\mathbf{a} \oplus \tilde{s}_\theta^*) \circ R_{\mathfrak{c},S} = R_{\mathfrak{c},s,\theta} \circ R_{\mu,\omega} \circ \mathbf{gr}(i_M) \circ \mathbf{gr}(q_M)^\top.$$

Now the definition of s_θ shows that for any $X \in \mathfrak{X}(N_{\mathfrak{c},S})$, we have

$$X^h + \iota_{X^h} \langle \theta, \vartheta_s \rangle \sim_{R_{\mathfrak{c},s,\theta}} X.$$

Since X^h is horizontal with respect to θ and $X = (q_N)_* X^h$, we also have

$$X^h + \iota_{X^h} \langle \theta, \vartheta_s - \frac{1}{2} s \circ \mathbf{a}(\theta) \rangle \sim_{R_{\mathfrak{c},s,\theta}} (q_N)_* X^h.$$

On the other hand, for $\xi \in \mathfrak{c}^\perp$, we have $\mathbf{a}(\xi) + s^*(\xi) \sim_{R_{\mathfrak{c},s,\theta}} 0$, so Eq. (A.6) shows that

$$\mathbf{a}(\xi) + \iota_{\mathbf{a}(\xi)} \langle \theta, \vartheta_s - \frac{1}{2} s \circ \mathbf{a}(\theta) \rangle \sim_{R_{\mathfrak{c},s,\theta}} (q_N)_* \mathbf{a}(\xi).$$

Therefore, for any $X \in TS$,

$$X + \iota_X \langle \theta, \vartheta_s - \frac{1}{2} s \circ \mathbf{a}(\theta) \rangle \sim_{R_{\mathfrak{c},s,\theta}} (q_N)_* X,$$

since this relation holds for both horizontal and vertical vector fields on the bundle $q_N : S \rightarrow N_{\mathfrak{c},S}$. This implies $R_{\mathfrak{c},s,\theta} \circ R_{\mu,\omega} \circ \text{gr}(i_M) = R_{\mu,\omega_\theta}$. Finally, since $\omega_\theta = q_M^* \tilde{\omega}_\theta$, we conclude that

$$R_{\mathfrak{c},s,\theta} \circ R_{\mu,\omega} \circ \text{gr}(i_M) \circ \text{gr}(q_M)^\top = R_{\tilde{\mu},\tilde{\omega}_\theta},$$

which proves the first part of the proposition.

Now suppose that $s(TS) \subseteq \mathfrak{c}$, then $s^*(\mathfrak{c}^\perp) \subseteq \text{ann}(TS)$, so $\vartheta_s = 0$ and $s^*\theta = 0$. Thus $\langle \theta, \vartheta_s - \frac{1}{2} s \circ \mathbf{a}(\theta) \rangle = 0$, and $\tilde{\omega}_\theta$ is independent of θ . This concludes the proof of the proposition. \square

A.0.2. Proof of the commutativity of partial reduction.

Proof of Proposition 3.2. By symmetry, it is sufficient to prove the first equality in Eq. (3.13). The key fact is that when $(\mathfrak{c}_1 \cap \mathfrak{c}_2, S_1 \cap S_2)$ is reductive data then $\mathfrak{c}_1 \cap \mathfrak{c}_2$ is coisotropic. As a result,

$$(A.8) \quad \mathfrak{c}_1^\perp \subseteq (\mathfrak{c}_1 \cap \mathfrak{c}_2)^\perp \subseteq \mathfrak{c}_1 \cap \mathfrak{c}_2 \subseteq \mathfrak{c}_2.$$

Thus,

$$\mathfrak{c}_{2,1} = (\mathfrak{c}_1 \cap \mathfrak{c}_2) / (\mathfrak{c}_1^\perp \cap \mathfrak{c}_2) = (\mathfrak{c}_1 \cap \mathfrak{c}_2) / \mathfrak{c}_1^\perp,$$

and

$$\mathfrak{c}_{2,1}^\perp = (\mathfrak{c}_1^\perp + \mathfrak{c}_2^\perp) / \mathfrak{c}_1^\perp.$$

Hence $\xi \in \mathfrak{d}$, $\xi' \in \mathfrak{d}_{\mathfrak{c}_1}$ and $\xi'' \in (\mathfrak{d}_{\mathfrak{c}_1})_{\mathfrak{c}_{2,1}}$ satisfy

$$\xi \sim_{R_{\mathfrak{c}_1}} \xi' \sim_{R_{\mathfrak{c}_{2,1}}} \xi''$$

if and only if

$$\xi \in \mathfrak{c}_1, \quad \xi' = \xi + \mathfrak{c}_1^\perp, \quad \xi' \in (\mathfrak{c}_1 \cap \mathfrak{c}_2) + \mathfrak{c}_1^\perp, \text{ and } \xi'' = \xi' + \mathfrak{c}_1^\perp + \mathfrak{c}_2^\perp.$$

Equivalently,

$$\xi \in \mathfrak{c}_1 \cap \mathfrak{c}_2, \quad \xi'' = \xi + (\mathfrak{c}_1 \cap \mathfrak{c}_2)^\perp, \text{ and } \xi' = \xi + \mathfrak{c}_1.$$

So

$$(A.9) \quad R_{\mathfrak{c}_{2,1}} \circ R_{\mathfrak{c}_1} = R_{\mathfrak{c}_1 \cap \mathfrak{c}_2}.$$

Before continuing, we introduce some notation. Let $\mu_{\mathbf{c}_1, S_1} : M_{\mathbf{c}_1, S_1} \rightarrow N_{\mathbf{c}_1, S_1}$ be the function whose graph is the support of $R_{\mathbf{c}_1, S_1}$, let

$$\begin{aligned} i_{N_1} : S_1 &\rightarrow N, & i_{M_1} : \mu^{-1}(S_1) &\rightarrow M, \\ i_N : S_1 \cap S_2 &\rightarrow N, & i_M : \mu^{-1}(S_1 \cap S_2) &\rightarrow M, \\ i_{N_{21}} : S_{2,1} &\rightarrow N_{\mathbf{c}_1, S_1}, & i_{M_{21}} : \mu_{\mathbf{c}_1, S_1}^{-1}(S_{2,1}) &\rightarrow M_{\mathbf{c}_1, S_1} \end{aligned}$$

denote the inclusions, and

$$\begin{aligned} q_{N_1} : S_1 &\rightarrow N_{\mathbf{c}_1, S_1}, & q_{M_1} : \mu^{-1}(S_1) &\rightarrow M_{\mathbf{c}_1, S_1}, \\ q_N : S_1 \cap S_2 &\rightarrow N_{\mathbf{c}_1 \cap \mathbf{c}_2, S_1 \cap S_2}, & q_M : \mu^{-1}(S_1 \cap S_2) &\rightarrow M_{\mathbf{c}_1 \cap \mathbf{c}_2, S_1 \cap S_2}, \\ q_{N_{21}} : S_{2,1} &\rightarrow (N_{\mathbf{c}_1, S_1})_{\mathbf{c}_{21}, S_{21}}, & q_{M_{21}} : \mu_{\mathbf{c}_1, S_1}^{-1}(S_{2,1}) &\rightarrow (M_{\mathbf{c}_1, S_1})_{\mathbf{c}_{21}, S_{21}} \end{aligned}$$

denote the quotient maps.

Now

$$\begin{aligned} (R_{\mathbf{c}_1, S_1})_{\mathbf{c}_{21}, S_{21}} &= \left((R_{\mathbf{c}_{2,1}} \circ R_{\mathbf{c}_1}) \times \text{gr}(q_{N_{21}}) \circ \text{gr}(i_{N_{21}})^\top \circ \text{gr}(q_{N_1}) \circ \text{gr}(i_{N_1})^\top \right) \\ &\quad \circ R \circ (R_{q_{N_{21}}} \circ R_{i_{N_{21}}}^\top \circ R_{q_{N_1}} \circ R_{i_{N_1}}^\top)^\top \end{aligned}$$

Now

$$(A.10) \quad x \sim_{\text{gr}(q_{N_1}) \circ \text{gr}(i_{N_1})^\top} y \sim_{\text{gr}(q_{N_{21}}) \circ \text{gr}(i_{N_{21}})^\top} z$$

if and only if

$$x \in S_1, \quad y = q_{N_1}(x), \quad y \in S_{2,1}, \quad \text{and } z = q_{N_{21}}(y).$$

Equation (A.8) implies that the $\mathbf{c}_1 \cap \mathbf{c}_2$ invariant manifold $S_1 \cap S_2$ is also \mathbf{c}_1^\perp invariant. Since $S_{2,1}$ is the set of \mathbf{c}_1^\perp orbits in $S_1 \cap S_2$, it follows that $y \in S_{2,1}$ if and only if $x \in S_1 \cap S_2$. Thus, Eq. (A.10) holds if and only if

$$x \in S_1 \cap S_2, \quad z = q_{N_{21}} \circ q_{N_1}(x), \quad \text{and } y = q_{N_1}(x).$$

But

$$q_{N_{21}} \circ q_{N_1}|_{S_1 \cap S_2} : S_1 \cap S_2 \rightarrow N_{\mathbf{c}_1 \cap \mathbf{c}_2, S_1 \cap S_2}$$

is the quotient map for the $\mathbf{c}_1^\perp + \mathbf{c}_2^\perp$ action, i.e. $q_{N_{21}} \circ q_{N_1}|_{S_1 \cap S_2} = q_N$. Therefore

$$(A.11) \quad \text{gr}(q_{N_{21}}) \circ \text{gr}(i_{N_{21}})^\top \circ \text{gr}(q_{N_1}) \circ \text{gr}(i_{N_1})^\top = \text{gr}(q_N) \circ \text{gr}(i_N)^\top.$$

A similar calculation shows that

$$\text{gr}(q_{M_{21}}) \circ \text{gr}(i_{M_{21}})^\top \circ \text{gr}(q_{M_1}) \circ \text{gr}(i_{M_1})^\top = \text{gr}(q_M) \circ \text{gr}(i_M)^\top,$$

and since the composition is clean we also have

$$(A.12) \quad R_{q_{N_{21}}} \circ R_{i_{N_{21}}}^\top \circ R_{q_{N_1}} \circ R_{i_{N_1}}^\top = R_{q_M} \circ R_{i_M}^\top$$

Combining Eqs. (A.9), (A.11) and (A.12) shows that

$$(R_{\mathbf{c}_1, S_1})_{\mathbf{c}_{21}, S_{21}} = (R_{\mathbf{c}_1 \cap \mathbf{c}_2} \times (\text{gr}(q_N) \circ \text{gr}(i_N)^\top)) \circ R \circ (R_{q_M} \circ R_{i_M}^\top)^\top = R_{\mathbf{c}_1 \cap \mathbf{c}_2, S_1 \cap S_2}.$$



REFERENCES

- [1] Anton Yu Alekseev, Henrique Bursztyn, and Eckhard Meinrenken. Pure spinors on Lie groups. *Astérisque*, (327):131–199 (2010), 2009.
- [2] Anton Yu Alekseev and Yvette Kosmann-Schwarzbach. Manin pairs and moment maps. *Journal of Differential Geometry*, 56(1):133–165, 2000.
- [3] Anton Yu Alekseev, Yvette Kosmann-Schwarzbach, and Eckhard Meinrenken. Quasi-Poisson manifolds. *Canadian Journal of Mathematics*, 54(1):3–29, 2002.
- [4] Anton Yu Alekseev, Anton Z. Malkin, and Eckhard Meinrenken. Lie group valued moment maps. *Journal of Differential Geometry*, 48(3):445–495, 1998.
- [5] Anton Yu Alekseev and Ping Xu. Derived brackets and Courant algebroids. Available at <http://www.math.psu.edu/ping/anton-final.pdf>, 2002.
- [6] Michael Atiyah and Raoul Bott. The Yang-Mills Equations over Riemann Surfaces. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 308(1505):523–615, March 1983.
- [7] Philip Paul Boalch. Quasi-Hamiltonian geometry of meromorphic connections. *Duke Mathematical Journal*, 139(2):369–405, August 2007.
- [8] Philip Paul Boalch. Through the analytic Halo: fission via irregular singularities. *Université de Grenoble. Annales de l'Institut Fourier*, 59(7):2669–2684, 2009.
- [9] Philip Paul Boalch. Geometry and braiding of Stokes data; Fission and wild character varieties. November 2011. Available at <http://arxiv.org/abs/1111.6228>.
- [10] Henrique Bursztyn, Gil R. Cavalcanti, and Marco Gualtieri. Reduction of Courant algebroids and generalized complex structures. *Advances in Mathematics*, 211(2):726–765, 2007.
- [11] Henrique Bursztyn, David Iglesias Ponte, and Pavol Ševera. Courant morphisms and moment maps. *Mathematical Research Letters*, 16(2):215–232, 2009.
- [12] Theodore James Courant. Dirac manifolds. *Transactions of the American Mathematical Society*, 319(2):631–661, 1990.
- [13] Theodore James Courant and Alan Weinstein. Beyond Poisson structures. In *Action hamiltoniennes de groupes. Troisième théorème de Lie (Lyon, 1986)*, pages 39–49. Hermann, Paris, 1988.
- [14] Vladimir Gershonovich Drinfel'd. Hamiltonian structures on Lie groups, Lie bialgebras and the geometric meaning of classical Yang-Baxter equations. *Doklady Akademii Nauk SSSR*, 268(2):285–287, 1983.
- [15] Vladimir Gershonovich Drinfel'd. *Quasi-Hopf Algebras*. Algebra i Analiz, 1989.
- [16] Vladimir Gershonovich Drinfel'd. On Poisson homogeneous spaces of Poisson-Lie groups. *Theoretical and Mathematical Physics*, 95(2):524–525, May 1993.
- [17] Jurgen Fuchs, Christoph Schweigert, and Alessandro Valentino. Bicategories for boundary conditions and for surface defects in 3-d TFT . March 2012. Available at <http://arxiv.org/pdf/1203.4568.pdf>.
- [18] Lars Hörmander. *The analysis of linear partial differential operators. III*. Classics in Mathematics. Springer, Berlin, 2007. Pseudo-differential operators, Reprint of the 1994 edition.
- [19] David Iglesias Ponte and Ping Xu. Hamiltonian spaces for Manin pairs over manifolds. September 2008. Available at <http://arxiv.org/abs/0809.4070>.
- [20] Anton Kapustin and Natalia Saulina. Surface operators in 3d topological field theory and 2d rational conformal field theory. In *Proc. Sympos. Pure Math.*, pages 175–198. Amer. Math. Soc., Providence, RI, 2011.
- [21] Anton Kapustin and Natalia Saulina. Topological boundary conditions in abelian Chern-Simons theory. *Nuclear Physics. B*, 845(3):393–435, 2011.
- [22] Citrad Klimčík and Thomas Strobl. WZW-Poisson manifolds. *Journal of Geometry and Physics*, 43(4):341–344, 2002.

- [23] David Li-Bland and Eckhard Meinrenken. Courant algebroids and Poisson geometry. *International Mathematics Research Notices*, 2009(11):2106–2145, 2009.
- [24] David Li-Bland and Eckhard Meinrenken. Dirac Lie Groups. pages 1–46, October 2011. Available at <http://arxiv.org/abs/1110.1525>.
- [25] David Li-Bland and Pavol Ševera. Quasi-Hamiltonian groupoids and multiplicative Manin pairs. *International Mathematics Research Notices*, 2011(10):2295–2350, 2011.
- [26] David Li-Bland and Pavol Ševera. Moduli spaces for quilted surfaces and Poisson structures. December 2012. Available at <http://arxiv.org/abs/1212.2097>.
- [27] Zhang-Ju Liu, Alan Weinstein, and Ping Xu. Manin triples for Lie bialgebroids. *Journal of Differential Geometry*, 45(3):547–574, 1997.
- [28] Jiang-Hua Lu. *Multiplicative and affine Poisson structures on Lie groups*. PhD thesis, University of California, Berkeley, 1990.
- [29] Jiang-Hua Lu and Alan Weinstein. Groupoides symplectiques doubles des groupes de Lie-Poisson. *Comptes Rendus de l'Academie des Sciences - Series I: Mathematics*, 309:951–954, 1989.
- [30] Jiang-Hua Lu and Milen Yakimov. Group orbits and regular partitions of Poisson manifolds. *Communications in Mathematical Physics*, 283(3):729–748, 2008.
- [31] Dmitry Roytenberg. *Courant algebroids, derived brackets and even symplectic supermanifolds*. PhD thesis, University of California, Berkeley, 1999.
- [32] Michael A. Semenov-Tian-Shansky. Dressing transformations and Poisson group actions. *Kyoto University. Research Institute for Mathematical Sciences. Publications*, 21(6):1237–1260, 1985.
- [33] Pavol Ševera. Letters to A. Weinstein. Available at <http://sophia.dtp.fmph.uniba.sk/~severa/letters/>.
- [34] Pavol Ševera. (Non-)abelian Kramers-Wannier duality and topological field theory. *Journal of High Energy Physics. A SISSA Journal*, (5):no. 49–10, 2002.
- [35] Pavol Ševera. Some title containing the words “homotopy” and “symplectic”, e.g. this one. In *Travaux mathématiques. Fasc. XVI*, pages 121–137. Univ. Luxemb., Luxembourg, 2005.
- [36] Pavol Ševera. Moduli spaces of flat connections and Morita equivalence of quantum tori. *Documenta Math*, 17:607–625, 2012.
- [37] Pavol Ševera and Alan Weinstein. Poisson geometry with a 3-form background. *Progress of Theoretical Physics. Supplement*, (144):145–154, 2001.
- [38] Vladimir G Turaev. Intersections of loops in two-dimensional manifolds. *Matematicheskii Sbornik. Novaya Seriya*, 106(148)(4):566–588, 1978.
- [39] Kyouzuke Uchino. Remarks on the definition of a Courant algebroid. *Letters in Mathematical Physics*, 60(2):171–175, 2002.
- [40] Katrin Wehrheim and Christopher Thomas Woodward. Quilted Floer cohomology. *Geometry & Topology*, 14(2):833–902, 2010.
- [41] Alan Weinstein. The symplectic structure on moduli space. In *The Floer memorial volume*, pages 627–635. Birkhäuser, Basel, September 1995.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY
E-mail address: libland@math.berkeley.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITÉ DE GENÈVE, GENEVA, SWITZERLAND, ON LEAVE FROM
 DEPT. OF THEORETICAL PHYSICS, FMFI UK, BRATISLAVA, SLOVAKIA
E-mail address: pavol.severa@gmail.com