

The strange (hi)story of particles and waves*

H. Dieter Zeh – www.zeh-hd.de – (March 2013)

Abstract: Attempt of a non-technical but conceptually consistent presentation of the state of the art in quantum theory from a historical perspective. While its first part is written for a general readership, Sect. 4 may be provocative, in particular, to quantum field theorists. I argue that the wave functions of quantum mechanics, in contrast to the wave functionals of QFT, which represent the true quantum states, have to be interpreted as field modes and their combinations that are “occupied” (excited states of their corresponding quantum oscillators).

1. Early History

The conceptual distinction between a discrete or continuous structure of matter (and perhaps other „substances“) goes back at least to the pre-Socratic philosophers. However, their concepts and early ideas were qualitative and speculative. They remained restricted to some general properties, such as symmetries, while the rational and quantitative understanding of a continuum in space or time had to wait two millennia until the development of calculus. Quantitative laws of nature and the concept of a mass point were only invented as part of classical mechanics.

These concepts were first applied to extended “clumps of matter”, such as the heavenly bodies or falling rocks and apples. It was in fact a great surprise to Newton and his contemporaries (about 1680) that such very different objects – or rather their centers of mass – obeyed the same laws of motion.¹ The objects themselves still seemed to consist of continuous matter. However, the new concept of mass points was quite early also applied to the structure of matter, that is, in the sense of an atomism. Already in 1738, Daniel Bernoulli explained the pressure of a gas by the mean kinetic energy of assumed gas particles, but without recognizing its relation to the phenomenon of heat. If one regarded these particles themselves as small hard spheres, though, the question for their internal structure would in principle arise anew.

* Free and extended translation of my unpublished German text “Die sonderbare Geschichte von Teilchen und Wellen” – available on my website since October 2011.

During the same time, Newton's theory of mechanics was also generalized by means of the concept of infinitesimal massive volume elements that can move and change their form and density according to their local interaction with their direct neighbors. This route to continuum mechanics was mainly a mathematical program of calculus that did not require any fundamentally novel physical concepts. The assumption of an unlimited divisibility of matter thus led to a consistent theory. In particular, it gave rise to wave-like propagating density oscillations which explained the phenomenon of sound. So it seemed that the fundamental question for the structure of matter had been answered.

As a byproduct of this "substantial" (or "Laplacian") picture of continuum mechanics, based on the assumption of individually moving infinitesimal elements of matter, also a formally elegant "local" (or "Eulerian") picture could be formulated. In the latter one neglects any information about trajectories of individual pieces of matter in order to consider only its spatial density distribution together with its corresponding current density as the kinematical objects of interest. In modern language, they would be called a scalar and a vector *field*. In spite of this new form, continuum mechanics remains based on the concept of a conserved material substance with its consequence of a continuity equation.

The picture of individually moving elements of a substance would lose any practical meaning, however, if the true elements of matter could move irregularly, as suspected for a gas by Daniel Bernoulli. Since his gas pressure is given by the density of kinetic energy, that is, by the product of the number density of gas particles and their mean kinetic energy, these fundamental elements could still be understood as forming a "chaotic continuum" by considering an appropriately defined simultaneous limit of infinite number density and vanishing size and mass of the particles. This remained a possibility even when chemists began to successfully apply Dalton's and Avogadro's hypotheses about molecular structures from the beginning of the nineteenth century in order to understand the chemical properties of the various substances. Similar to August Bravais's concept of crystal lattices (about 1849), these structures were often regarded as no more than an intuitive tool to describe the internal structure of a multi-component continuum. It was upheld by many even after Maxwell's and Boltzmann's explanation of thermodynamic concepts in terms of molecular kinetics, and in spite of repeated but until then unsuccessful attempts to determine a finite value for Avogadro's or Loschmidt's number. The "energeticists", such as Wilhelm Ostwald, Ernst Mach and initially also Max Planck remained convinced until about 1900 that internal energy, heat and entropy were fundamental fields or continuous substances. Indeed, even after the determination of Loschmidt's number could they have used an argument that formed a severe

2

Loschmidt's number could they have used an argument that formed a severe problem for atomists: Gibbs' paradox of the missing entropy of self-mixing of a gas. Today it is usually countered by referring to the indistinguishability of elementary particles, although the argument requires more, namely the *identity* of states obtained by particle permutations. However, this assumption is in conflict with the substantial picture of particles following individual trajectories, while it would readily apply to the permutation of different wave packets of a *field*. Although we are using quite novel theories today, these conceptual differences do remain relevant (see Sect. 4).

Another object affected by the early dispute about particles and waves is light. According to its appearance, light was traditionally regarded as a "medium" rather than a substance. Nonetheless, and in spite of Huygens' early ideas of light as a wave phenomenon, Newton tried to explain it by means of particles of light, which are supposed to move along their trajectories according to the local refractive index of matter. This proposal was later refuted by various interference experiments, in particular those of Thomas Young in 1802. It remained open, though, what substance (called the ether) did oscillate – even after light had been demonstrated by Heinrich Hertz in 1886 to represent an electromagnetic phenomenon in accordance with Maxwell's equations. The property of these fields to propagate and carry energy gave them a certain material character that seemed to support the world of continua as envisioned by the energeticists. For a long time, Ernst Mach used to reply "Have you seen one?" whenever somebody mentioned atoms to him. Later in this article I will argue that this doubting question may still be appropriate today – even though we *seem* to observe individual atoms as particles.

At the end of the nineteenth century, the successful continuum hypothesis suffered a number of decisive blows. In 1897, J. J. Thomson determined the elementary electric charge; in 1900, Max Planck postulated his radiation quanta with great success; and in 1905, Albert Einstein estimated the value of Loschmidt's number N_L by means of his theory of Brownian motion. Thereafter, even the last energeticists resigned. Einstein then proposed the concept of particles of light (later called photons) – although he regarded them merely as a "heuristic point of view" that he was never ready to fully accept himself. Niels Bohr, in 1913, replaced even the concept of continuous motion by stochastic "jumps" between his discrete electron orbits in the hydrogen atom – in accordance with Planck's and Einstein's ideas about the radiation process. These early ideas led later to the insufficient interpretation of quantum mechanics as no more than some stochastic dynamics for otherwise classical particles.

However, the development soon began to proceed in the opposite direction again.² In 1923, Louis de Broglie inverted Einstein's speculative step from light waves to photons by postulating a wave length $\lambda = c/\nu = h/p$ for the electron, where p is its momentum, in analogy to Planck's relation $E = h\nu$. For him this could only mean that all microscopic objects must consist of both, a particle and a wave, whereby the wave has to serve as a "guiding field" for the particle. This field would thereby have to be more powerful than a conventional field of force, since it has to determine the velocity rather than merely the acceleration; the initial velocity cannot be freely chosen any more in this description that was later further developed and completed by Madelung and by Bohm. It turned out that the assumed guiding wave cannot be local (defined in space), as it must be identical with the entangled wave function to be described in Sect. 3.

2. Wave Mechanics

Inspired by de Broglie's ideas, Schrödinger based his novel wave mechanics of 1926 on the assumption that electrons are *solely* and uniquely described by wave functions (spatial fields, as he first thought). This allowed him to explain the hydrogen spectrum by replacing Bohr's specific electron orbits in the atom by standing waves (energy eigenstates). For a special case, the harmonic oscillator, he was furthermore able to construct narrow "wave packets" that would dynamically behave like particles (see the Figure below for the case of *free* motion). Shortly thereafter, interference phenomena in agreement with de Broglie's wave length were observed by Davisson and Germer for electrons scattered from crystal lattices. According to Schrödinger's equation, a wave function can also penetrate a potential barrier and thus explain "quantum tunneling", as required for the possibility of α -decay. Does this not very convincingly demonstrate that electrons and other "particles" are in reality just wave packets of some fields that are described by Schrödinger's equation?




Figure: A wave packet (real part of an initial Gaussian modulated by a plane wave $e^{2\pi ix/\lambda}$) moving freely according to the Schrödinger equation, depicted at three different times (blue: $t=0$, red: $t=0.04$, yellow: $t=1$ in arbitrary units). When comparing blue and red one recognizes that the packet as a whole moves faster than its waves, while the yellow curve demonstrates a slight dispersion of the packet (in contrast to the mentioned harmonic oscillator). The center of the packet moves according to the group velocity $v = p/m = h/m\lambda$, where the mass m is now just a parameter of the wave equation. For this reason, momentum is in wave mechanics *defined* by h/λ . It can indirectly be measured even for plane waves, which do *not* possess a group velocity, by means of a conservation law for the sum of wave numbers that holds for scattering (“momentum transfer”) with objects which do exist in localized wave packets (such as observable Brownian particles). Already for atomic masses and thermal velocities, the de Broglie wave length is clearly smaller than the radius of a hydrogen atom. So one may construct quite narrow wave packets for their center of mass wave functions. Although the dispersion of the wave packet is reduced with increasing mass m , it becomes non-negligible with growing time. Therefore, one would need a new mechanism that dynamically reduces the “coherence lengths” characterizing the packet in order to retain the appearance of a particle.

A few months before Schrödinger invented his wave mechanics, Heisenberg had already proposed his matrix mechanics. In contrast to Schrödinger, he kept the concept of particles, but introduced certain formal “rules” that were to replace the concept of their trajectories. Later, matrix mechanics turned out to be “equivalent” to Schrödinger’s theory when used to calculate probabilities for closed systems with a well defined Hamiltonian. Together with Born and Jordan, Heisenberg constructed an elegant algebraic framework that could be applied in order to “quantize” mechanical systems in general. This mathematical abstraction was in accordance with Heisenberg’s preferred view of the physical world as ultimately being ruled by Platonic ideas (such as in mathematics) in contrast to materialism. A year later, he complemented this formalism by his uncertainty relations between position and momentum of an electron or similar “conjugate” pairs of variables. This fundamental uncertainty is evidently in conflict with the concept of a particle, while in wave mechanics it would simply be a consequence of the Fourier theorem – without any uncertainty of the wave function itself. It is not surprising that Schrödinger’s intuitive wave mechanics was preferred by most practicing physicists – for a short time even by Heisenberg’s mentor Max Born. For example, Arnold Sommerfeld wrote only a “Wellenmechanischer Ergänzungsband” to his important book on “Atombau und Spektrallinien”.

So everything seemed to look fine if there had not been those incredible stochastic quantum jumps between the different standing waves of an electron in the atom. They appeared absolutely incompatible with Schrödinger’s otherwise successful wave dynamics. Similarly, other initially extended wave functions seemed to “jump” into particle-like local

wave packets during measurements. In a Wilson chamber, one could even observe tracks of droplets that can be regarded as successions of position measurements along particle trajectories. As a consequence, Schrödinger seemed to resign when Max Born, influenced by Wolfgang Pauli, re-interpreted his new probability postulate, that was originally meant to describe jumps between different wave functions, in terms of probabilities for the spontaneous occurrence of genuine particle properties. This interpretation turned out to be very successful (and earned Born a Nobel prize) even though it was never quite honest, because the wave function does *not only* describe probabilities. It also defines observable individual properties of microscopic objects, such as energy or angular momentum – represented by their eigenstates. Similarly, a spinor (that is, a quantum state of spin) is known to describe probabilities for the “occurrence” of other individual spinor states rather than for any classical properties.

The problem was so painful that Heisenberg spoke of the wave function as “a new form of human knowledge as an intermediary kind of reality”, while Bohr introduced his irrational principle of complementarity that required the application of incompatible classical concepts, such as particles and waves, to the same objects. No doubt – this was an ingenious kind of pragmatism, but from there on the quest for a consistent description of Nature was not allowed any more in microscopic physics. Bohr insisted that “there is no microscopic reality” – a conclusion that was often regarded as philosophically very deep. Only few dared to object that “this emperor is naked”, and the new terminology no more than empty words, used to verbally circumvent an inconsistency. The great number of philosophical or formal arguments for its justification is even the more impressive. In particular, it has always remained open when and where precisely the probability interpretation or the Heisenberg cut have to be applied. The Hungarian Eugene Wigner called this situation a “Balkanization of physics” – a traditional (Prussian-Austrian) name for a region without law and order.

3. Entanglement

So one should take a fresh and more complete look at Schrödinger’s wave mechanics. When he first formulated it, he used Hamilton’s partial differential equations as a guiding principle. These equations, the result of a reformulation of classical mechanics, are solved by a function that would describe a whole ensemble of classical trajectories with different initial conditions – sort of a wave function without interference. Hamilton was mainly interested in the elegant mathematical form of this theory rather than in applications. This turned out to be advanta-

geous for Schrödinger. He assumed that Hamilton's equations were no more than a short wave lengths approximation (corresponding to the limit $h \rightarrow 0$) of a fundamental wave theory – similar to the approximation of geometric optics that could be applied to Maxwell's theory in order to obtain trajectories for Newton's hypothetical particles of light. With respect to Heisenberg's particle concept, he later remarked ironically that even Newton's assumption of light particles could have been saved in spite of the observed interference phenomena if one had required some "uncertainty relations" for them. However, the short wave length approximation means only that local parts of an extended wave propagate independently of one another along trajectories – not that they represent particles. These partial waves *can* interfere, that is, they can act coherently if focused in some way.

While light waves propagate in three-dimensional space, Hamilton's waves must according to their construction exist in the configuration space that represents the system under consideration. Therefore, Schrödinger, too, obtained wave functions on configuration spaces of various dimensions. Later this turned out to be the only correct form of wave mechanics. It can also be understood as a consequence of Dirac's general superposition principle, since the superposition of all classical configurations defines a wave function on configuration space. Dirac himself understood his superpositions in Born's sense as probability amplitudes for properties that are formally represented by Heisenberg's "observables", that is, not only for *points* in configuration space. However, if the observables are represented by a dyadic product of their eigenfunction, this leads back to probabilistic relations corresponding to a collapse of the wave function (a projection).

Schrödinger was convinced of a reality in space and time, and he intended to describe the real electron as a field. So he initially restricted himself with great success to single-particle problems (quantized mass points, whose configuration space is isomorphic to space). Consequently, he spoke originally of a " ψ -field". A three-dimensional wave function can also be used to describe scattering problems – either for the center of mass wave function of an object scattered from a potential, or for the relative coordinates of a two-body problem. In such scattering problems, Born's probability interpretation is particularly suggestive because of the subsequent position measurement in a detector. Usually this three-dimensional wave function is meant when one speaks of the *wave-particle dualism*. Unfortunately, many textbooks still describe mainly such spatial wave functions for didactic but very misleading reasons – in particular when presenting the two-slit interference experiment as *the* key to understanding quantum mechanics.

The generalization (or rather the return) to wave functions in configuration space happened almost unnoticed at those times of great confusion (for some physicists even until today). In contrast to fields, even single-particle wave functions do not describe additive charge or energy distributions. Each partial wave of a plane wave, projected out by means of absorbers, for example, describes the full charge and energy, since, in the absence of a potential, energy is defined by the wave length.

Schrödinger took initially great pains to disregard or to re-interpret this general wave equation in configuration space, even though it is precisely its application to oscillating field amplitudes rather than mass points that is able to explain Planck's quanta $h\nu$. The oscillator spectrum $E = nh\nu$ is proportional to the natural numbers, and thus allows one to think of n as a *number* of quanta (later identified with photons). In Schrödinger's wave mechanics, these quantum numbers n are explained by the numbers of nodes of wave functions that have to obey certain boundary conditions. This wave function is here not a spatial function, as assumed in the Figure, but an (in general time-dependent) wave function on the configuration space of field amplitudes. Different field modes can fortunately be treated separately for harmonically coupled oscillators. Although the oscillator spectrum can alternatively be derived from Heisenberg's algebra of observables, nodes of the wave function in this configuration space (which are quite unrelated to the spatial nodes of the fixed field mode) have recently been observed for various photon number eigenstates in an elegant experiment performed by the group of Haroche and Raimond in Paris.³ The importance of this fundamental experiment for the wave-particle debate has as yet not been sufficiently appreciated by the physics community.

The difference between Schrödinger's theory and a classical field theory becomes particularly obvious from the fact that the amplitudes of the classical field now appear as *arguments* in Schrödinger's function. While a general "one-photon wave function" can be understood as a superposition of various classical *field modes* in their first excited quantum state ("occupied once" – with all others being in their ground state), a semi-classical *field* state representing a certain classical mode has to be described as a coherent superposition of many *different* excitations (photon number eigenstates) of this mode. In contrast to the free wave packet shown in the Figure, these "coherent states" (Gaussians again) preserve their shape and size while following the classical motion of an oscillator. In this sense, they imitate a classically oscillating field. These different kinds of states thus can describe both particle numbers or fields – mutually excluded by a Fourier theorem – in one and the same formalism of QFT.

Before trying to study interacting quantum fields, physicists of the end of the twenties investigated the quantum mechanics of many-particle systems, such as molecules or multi-electron atoms. These could often *approximately* be described by assuming different single-particle wave functions for each electron, while the atomic nuclei seemed to possess fixed or slowly moving positions similar to classical objects. For example, this assumption explained the periodic system of the chemical elements. On closer inspection it turned out – at least for small molecules – that *all* particles, including the nuclei, had to be described by one common wave function in their full configuration space. This cannot normally be a product or determinant of single-particle wave functions – a consequence that is called entanglement. Every physics student is using the entanglement between an electron and a proton in the hydrogen atom when writing the wave function as a product of cms and relative coordinates. The simplest nontrivial case, the Helium atom, was successfully studied in great numerical detail by Hylleraas, using variational methods, in a series of papers since 1929. Similarly, in the theory of multi-electron atoms one has to take into account “configuration mixing” as a correction to the independent-particle (Hartree-Fock) approximation. Already Arnold Sommerfeld remarked in his *Wellenmechanischer Ergänzungsband* that “Heisenberg’s method”, which considered only the anti-symmetrization of product wave functions, is insufficient for this purpose. Anti-symmetrization is often misunderstood as entanglement, although it represents no more than a *formal* entanglement between physical properties and meaningless particle numbers that disappears in the formalism of quantum field theory (see Sect. 4). Today, all closed non-relativistic multi-particle systems are accepted as being correctly described by one entangled wave function in their high-dimensional configuration space.

But how can the space of all possible classical configurations, which even has different number of dimensions for different systems, replace three-dimensional space as the new stage for the dynamics of a wave function that might be expected to describe physical reality? If our Universe consisted of N particles (and nothing else), its configuration space would possess $3N$ dimensions. For early quantum physicists – including Schrödinger, of course – this was inconceivable. On the other hand, the concept of a space of possible configurations fits excellently with Born’s probability interpretation for the occurrence of such classical states. Entanglement can then conveniently be understood as describing statistical correlations between measured variables. But only for *measured* variables! In the mentioned case of the Helium atom, for example, entanglement is responsible for the precise individual ground state energy and other observable properties – regardless of any statistical interpretation. This irritating fact is usually simply “overlooked” for convenience. Even in scattering processes one

often needs coherent amplitudes, which can only be described by one entangled wave function for all scattering fragments. Only after Einstein, Podolski and Rosen had clearly shown in 1935 that entanglement can have observable consequences, did Schrödinger regard this property as the deepest interpretational challenge to his theory (although he kept calling it a “statistical correlation”). The three mentioned authors had in fact incorrectly concluded that the wave function can only be a statistical (not a complete or ontic) description, so that further (“hidden”) variables would have to exist.

While it was often thought that these new variables might perhaps never be found in an experiment, it came as a surprise to most physicists when John Bell showed in 1964 that *every* local theory (regardless of whether it consists of particles, fields or anything else) cannot be compatible with certain observable consequences of entangled wave functions. In order to demonstrate this incompatibility, Bell used quite arbitrary local variables (just names for something not yet known but assumed to exist). However, most physicists had by then become so much accustomed to Bohr’s denial of a microscopic reality that they immediately accused Bell for having used a “long refuted assumption”. As the crucial consequences of entangled wave functions have since always been confirmed, physicists disagree deeply about whether this experimental result excludes microscopic locality (in space) or rather reality. For neither those who are ready to accept a non-local wave function as representing reality nor the opponents of any kind of microscopic reality feel particularly disturbed by Bell’s theorem. Actually, these two camps are essentially identical with those preferring either the Schrödinger or the Heisenberg picture, respectively. This seems to be the origin of many misunderstandings between them.

If, for consistency, one assumes Schrödinger’s theory to apply universally, one has to accept the existence of one entangled wave function for the whole universe. Heisenberg and Bohr assumed instead that the wave function “loses its meaning” after a final measurement that concludes an experiment – another way of avoiding unwanted consequences. This “end of the experiment” would remain vaguely defined just as Heisenberg’s cut. However, if one does assume that there is a universal wave function that always evolves according to the Schrödinger equation, this leads to an entirely novel world view that appears unacceptable to most physicists because of its unusual consequences. For example, if one measures a microscopic property that is initially in a wave function with components representing two or more different values, this will give rise to an entangled state for the microscopic system and the pointer position, including – if appropriately generalized – Schrödinger’s infamous cat. To

cure this dilemma, one assumes in von Neumann's orthodox interpretation that the Schrödinger dynamics must be complemented by a stochastic "collapse of the wave function" into a product of narrow wave packets for all macroscopic variables. The same collapse may also be able to re-localize the spreading free wave packet of the Figure. In the Copenhagen interpretation, one would instead conceptually jump from a description in terms of wave functions to one in classical terms. This unsatisfactory situation – obviously related to the mentioned "Balkanization" of physics – is also called the quantum measurement problem.

So if one insists on the universal validity of the Schrödinger equation, one has to explain what an entangled wave function for the microscopic system and an apparatus can possibly mean. That is, one has to discuss what happens when one tries to observe such a state. Toward that end one has to include the observer into this quantum mechanical description.⁴ If he reads off the measurement result, he must himself become part of the entanglement. So he would then simultaneously exist in different states of awareness – similar to the fate of Schrödinger's cat. As Hugh Everett first noticed in 1957, this result is not in conflict with our subjective observation of *one* individual outcome of the measurement, since each "component state" of the observer would register precisely one component of the quantum world (his "relative state", representing one definite outcome). As there are many such component states in their global superposition, the question of which one contains the "true" successor of the experimenter who prepared the experiment has no unique answer. After the measurement, these different versions of the observer are dynamically clearly distinct (see the following remarks about decoherence) – even more so than spatially distinct observers in one interwoven classical world, who can at least communicate. A conceptually novel theory must be expected to require also quite novel kinds of interpretation! Certainly this is a very unusual, but at least a consistent solution of the problem, as it does not need any new physical assumptions. Any attempt to avoid this conclusion would instead require that the validity of quantum theory were limited in some way.

Until very recently one did in fact generally believe that such conceptual or dynamical border lines between micro- and macrophysics do exist – even though they have never been directly confirmed in an experiment. Otherwise it should be possible to observe individual properties of the entangled combination of the microscopic system and its measurement apparatus – similar to the energy or other properties of Hylleraas's whole Helium atom.

However, this bipartite entanglement is not yet complete under realistic conditions. Every macroscopic system must unavoidably, very fast, and in practice irreversibly interact

with its “environment” (the rest of the quantum universe), and thereby uncontrollably extend the entanglement that resulted from a measurement – even before an observer enters the scene. Because of the large number of effective degrees of freedom of a realistic environment, this situation can not be formulated in complete detail, but one may at least understand how a wave function representing different macroscopic pointer positions, for example, would permanently be transformed into an *apparent* ensemble of narrow wave packets that mimic classical states (points in configuration space).⁵ A local observer is unable to observe the resulting global entangled state as a whole. This unavoidable consequence for macroscopic variables, which defines the true border line between microphysics and macrophysics, is called decoherence, since predominantly the phase relation between separate wave packets are locally lost (dislocalized) in this way. Decoherence was the first successful application of entanglement beyond microscopic systems, and it explains also why one seems to observe individual atoms as apparent particles in a Paul trap or tracks in a Wilson chamber.⁶

Still, many leading physicists who are not happy any more with the Copenhagen interpretation prefer to assume some new dynamics (such as a collapse of the wave function) in order to avoid the consequence of Many Worlds. This is definitely a possibility that would also solve the measurement problem under the assumption of a universally valid wave function describing reality. One should keep in mind, though, that *all* as yet observed apparent deviations from the Schrödinger equation, such as quantum jumps or measurements, can be described (and have in some cases even been confirmed experimentally) as decoherence phenomena. If a collapse mechanism does nonetheless exist, it would therefore have to be triggered by decoherence – but this would then not make any difference in practice.

If one of two spatially separated but entangled microscopic systems were measured, their total state would according to a unitary description become entangled with the apparatus, including the latter’s environment. Nothing else so far. An observer, for example at the position of the second system, becomes part of this entanglement only when he receives a signal about the result. He would thereafter exist in various versions in the different world components that have been dynamically separated from one another by the decoherence process. In contrast, a collapse caused by the measurement would have to affect the distant objects instantaneously in order to avoid other weird consequences. If the distant observer measures also the microsystem at his own location (before or after he receives the signal about the first measurement – thus including delayed choice experiments), his state of awareness may then depend on the result of both measurements, that is, it must have split twice, whereby the order

does not matter. But only if one *postulates* statistical weights according to Born's probability measure for the observer's different components or subjective "versions", will he according to the thus defined subjective ensemble very *probably* confirm the frequency correlations that are predicted from quantum theory (and which violate Bell's inequality). These statistical weights are in fact the only conceivable ones that are conserved under further measurements, that is, which factorize into products of probabilities for subsequent measurements. Everett regarded this as a proof of Born's rule,⁷ while his branches can be defined according to the irreversible dislocalization of their coherence.

In this unusual way, *all* those much discussed "absurdities" of quantum theory can be consistently explained. It is, in fact, precisely the way how they were all predicted – except that the chain of unitary interactions is usually interrupted with the last "relevant" measurement that is accompanied by decoherence. So-called quantum teleportation is another example, where one can easily show, using a consistent unitary description, that nothing is teleported that was not prepared in advance at its final position in one of the components of the entangled wave function.⁶ This demonstrates again that entanglement cannot merely describe "information" about possibilities in the future, even though one is free to *assume* that a global collapse with undetermined outcome did occur (or that this outcome came into existence in some other way) together with the first irreversible decoherence process in a measurement. Since such a pragmatic restriction of "quantum reality" to one single component ("our quantum world") represents here a mere convention rather than a physical process, one may even define it to occur superluminally. However, a quantum state described by a wave function is non-local in general, so that "quantum teleportation" does not require any additional spooky action at a distance.

Since the global wave function evolves deterministically, the observed quantum indeterminism cannot represent any objective dynamics. In Everett's interpretation, it is a "subjective" phenomenon related to the multiple future of all potential observers. It is nonetheless objectivized between those versions of different observers who live in one and the same branch or Everett world (including corresponding versions of Wigner's friend). In contrast to the conceptual dualism of the Copenhagen interpretation, classical properties can then also be understood in terms of wave packets as mere phenomena, objectively based on decoherence.

4. Quantum Field Theory

We have seen that the interpretation of quantum theory in terms of a universal wave function admits a consistent (even though quite unusual) description of Nature, but this does not yet bring the strange story of particles and waves to an end. Instead of the expected spatial waves (fields) we were led to wave functions on some high-dimensional “configuration space” (a name that is justified only because of its classical appearance). For a universe consisting of N particles, its configuration space would possess $3N$ dimensions, but we have already seen that for QED (quantum electrodynamics) it must be complemented by the infinite-dimensional configuration space of the Maxwell fields. A factorizing wave functional for its free field modes was sufficient to explain Planck’s quanta in terms of the number of nodes of the corresponding factor wave functions. The spontaneous local appearance of photons as apparent particles can then again be understood by means of a decoherence process in the detector.

However, it is known from the theory of relativistic electrons that they, too, have to be described by a quantized *field*, a consequence that must then also apply to the non-relativistic limit. The relativistic generalization of a one-electron wave function is called the *Dirac field*, and thus regarded as a function on spacetime. It can *not* be generalized to an N -electron field on a $4N$ -dimensional “configuration spacetime”, even though this has occasionally been proposed. In fact, in the Schrödinger picture of QED, the Dirac field is only used to define, by its configuration space, the space on which the corresponding time-dependent wave functionals live. According to the rules of canonical quantization, these wave functionals have then again to obey a generalized Schrödinger equation.^{8,9} This assumption avoids the N -dependence of the relevant configuration spaces for varying numbers of “particles”. Since Schrödinger had originally discovered one-electron wave functions by the same formal quantization procedure, the quantization of the Dirac field is for this purely historical reason called a “second quantization”. As explained above, however, the particle concept, and with it the first quantization, are no more than historical artifacts from a fundamental point of view.[†]

[†] It should be emphasized, though, that the Pauli principle, valid for fermions, may not yet be sufficiently understood. It is true that the individual components of the Dirac equation also obey the Klein-Gordon equation, but the latter’s quantization would again lead to *all* bosonic quantum numbers $n = 0, 1, 2, \dots$. Anti-commuting field operators, which lead to anti-symmetric multi-particle wave functions, were proposed quite *ad hoc* by Jordan and Wigner, and initially appeared artificial even to Dirac. Interpreted consistently, their configuration space would represent a spatial continuum of coupled bits (“empty” or “occupied”) rather than a continuum of coupled oscillators. The n -th excited state of this bit continuum (that is, n

The hamiltonian form of the Dirac equation is unusual as a consequence of its linearization especially insofar as canonical momenta are not defined by time derivatives of the variables any more. Nonetheless, the two occupation numbers 0 and 1 resulting from the *assumption* of anti-commuting field operators are again interpreted as “particle” numbers because of their well-known consequences in the quasi-classical world (“clicks in the counter”) resulting from decoherence. Field modes “occupied” once and their superpositions define what are usually called single-particle wave functions. In contrast to the case of photons, however, one does not observe any superpositions (wave functionals) of *different* electron numbers. This has traditionally been regarded as a fundamental restriction of the superposition principle (a “superselection rule”), but it may again be understood as a consequence of decoherence. For charged particles, the unavoidable Coulomb field assumes the role of an unavoidable environment.¹⁰

For relativistic reasons, *all* known elementary physical objects (still called “elementary particles” because of their phenomenology that results from decoherence) have to be described as quantum fields. The contrast between the first order in time of the Schrödinger equation and the second order of the classical field equations with their negative frequencies opens the door to the concept of “anti-bosons”. (For fermions this relation may take a different form.) Because of the universality of this concept of quantum fields, one also expects a “theory for everything” to exist in the form of a unified quantum field theory. At present, it is no more than the most plausible attempt, though, that the fundamental stage for the universal wave function is given by the configuration space of some fundamental field(s). On the other hand, the general framework of Schrödinger’s quantum theory with its concept of a universal wave function (or Dirac’s universal superposition) obeying unitary dynamics seems as yet to be well justified.

occupied positions) would then represent n point-like “objects”. Because of the dynamical coupling between bit-neighbors, these objects can move, but only *after* their quantization (application of the superposition principle) would this give rise to propagating waves. In contrast, coupled oscillators, which define a field, propagate as waves already classically, and they obey a *classical* superposition principle (in space rather than in configuration space) in addition to the quantum superposition principle that is realized by the field functionals only.

An additional (and possibly related) source of confusion is the fact that relativistic invariance can and need not be manifest in the canonical formalism of the Schrödinger picture. For example, the canonical quantization of the Maxwell field leads to a wave functional $\Psi\{A(\mathbf{x});t\}$, with a vector field A defined at all space-points \mathbf{x} on an arbitrary simultaneity t .

For example, the traditional formulation that one particle is in the state described by the wave function ψ_1 and a second one in the state ψ_2 has to be replaced by the statement that the two *field modes* ψ_1 and ψ_2 are both in their first excited field quantum state. Obviously, a permutation of the two modes is then an identity, and there is only one total state of this kind to be counted in any statistics. This consequence eliminates Gibbs' paradox in a natural way.

It is similarly inappropriate to conclude that *wave functions* can be directly observed in Bose-Einstein condensates (as often claimed). What one does observe in this case is again the multiply "occupied" three-dimensional boson *field mode* – even though most bosons are in general regarded as particles because of their normal appearance in detectors. Instead of the free field modes used for photons, for strongly interacting bosons one may here more appropriately use self-consistent modes in analogy to the selfconsistent Hartree (-Fock) single-fermion wave functions. Both cases lead to an effective non-linear equation – for bosons called the Gross-Pitaevskii equation. In spite of this non-linearity, the quantum states proper are, of course, described by the linear and unitary universal Schrödinger equation, relativistically always understood in the sense of Tomonaga.⁸ Their true wave functions, which are very different from the easily observable field modes, have now also been observed in the form of their Wigner functions in Ref. 3 for various "photon" number eigenstates in a fixed electromagnetic laser mode. The Wigner function is here a partial Fourier transform of the dyadic product $\Psi(q)\Psi^*(q')$, and hence equivalent to the wave function $\Psi(q)$ itself, where q is the amplitude of the field mode. The resulting three-dimensional plot of the Wigner function allows one to clearly recognize the nodes which define "photon" numbers n . Nonetheless, apparent photons can be observed as clicks in the detector or spots on a screen.

Unfortunately, the general theory of interacting fields requires such an enormous effective number of entangled degrees of freedom that it can normally not even approximately be treated in a fundamental way. Instead of consistently applying the successful concepts of quantum mechanics, such as wave functions, to the new variables (field amplitudes), only semi-phenomenological methods of calculation are therefore used for specific purposes of quantum field theory - mostly for solving scattering problems. Their construction and interpretation is based on particle pictures again (for example in Feynman diagrams or in the form of probabilities for the appearance of apparent particles). Freeman Dyson's "equivalence"¹¹ between the use of relativistic field functionals (Tomonaga) or field operators (Feynman) is again essentially the same as that between the Schrödinger and the Heisenberg picture. However, the latter will hardly ever be able even in principle to characterize the complex situation

of steadily growing global entanglement. As relativistic physics is based on the absence of absolute time, the relativistic generalization of the Schrödinger equation can only be given by the Tomonaga equation with its “many-fingered” concept of time – but certainly *not* by the Dirac equation. The time parameter of a relativistic single-particle equation could only represent proper time along the particle trajectory, which does not exist any more in quantum mechanics. Apparent particle lines in Feynman diagrams, on the other hand, are just shorthand for certain field modes (such as plane waves indicated by particle momenta) that appear under an integral for calculating scattering amplitudes. Closed lines (“virtual particles”) then represent entanglement of the corresponding quantum fields.

Similar semi-phenomenological methods as in QFT are also much used in solid state physics, even when solid bodies are regarded as given N -particles systems. For example, they may give rise to effective phonon fields or other “quasi-particles”. Different thermodynamic phases of many-particle systems, which often determine an effective ground state in analogy to a Fock vacuum, can in some cases be explained by a different number of particles (such as electrons in a metal) that contribute to *stable entanglement* in their specific ground state. Analogously, effective fields of QFT that would merely have to be “renormalized” must not be expected to represent its fundamental variables which span the space for the universal wave function. This opens up quite novel possibilities, perhaps even to understand all fermions as quantum consequences of certain topological effects.

The successful phenomenology of the familiar effective fields (“elementary particles”), such as described by the Standard Model, must certainly form the major touchstone for any fundamental theory of the future. At present, it does not seem to offer really convincing hints for the structure of such a theory, while purely mathematical arguments must remain speculative and even questionable as far as there is no general consensus about the meaning of their quantization. Many modern quantum field theorists and mathematical physicists seem to regard these phenomenological models together with the calculational methods and recipes – often applied by using classical field or particle concepts – as the quantum field theory proper. In fact, why should one expect a consistent theory if there is no microscopic reality to be described (as assumed in the Copenhagen interpretation and its variants)?

Our conclusion that the particle aspect is a mere consequence of the decoherence process in the detector may understandably not be of particular interest to a practicing high-energy experimentalist, but it seems to be unknown even to many theoreticians in this field. So they sometimes simply call the enigmatic objects of their research “wavicles” because they

cannot make up their mind between particles and waves. This confusing language represents just another example of Wigner's "Balkanization of physics" (many words instead of many worlds). Waves are in the context of the wave-particle "dualism" usually understood in the sense of spatial waves – not as high-dimensional wave function(al)s. As we have seen, this is quite insufficient for a complete and consistent quantum theory.

If one also allowed for a multiverse of "landscapes" (Tegmark's level 2 of multiverses), which is suggested in many as yet speculative field theories that lead to an inhomogeneous universe on the very large scale, the "subjective selection" of an observer with his frog's perspective had to be roughly characterized by a hierarchy of at least four steps: (1) the selection of one of many component landscapes from their quantum superposition (in the sense of Tegmark's level 3, that is, Everett), (2) the selection of a particular region in this three or higher dimensional landscape (our specific "world" that may be characterized by specific values of certain "constants of nature" – level 2), (3) the selection of one specific complex organism from all of those that may exist in this world plus some "moment of awareness" for it (giving rise to an approximate subjective localization in space and time), and (4) the selection of one of its "quantum versions" (again in the sense of level 3).⁴ Only step 3 can *not* be objectivized in the usual sense of this word, that is, with respect to different observers in the same "world", but at least at this step "our" selection as humans must have used the weak anthropic principle.

5. Quantum gravity

I cannot finish this brief review of quantum theory without having mentioned quantum gravity.¹² Although one cannot hope to observe quanta of gravity in the foreseeable future, the formal quantization of gravity can hardly be avoided for consistency in view of the quantization of all other fields. The spacetime metric of general relativity may be regarded as a "normal" tensor field – in close analogy, in particular, to Maxwell's theory with its vector potential as the dynamical variable. It must then also appear among the arguments of a universal wave function, and thus be entangled with all other fields – as it turns out in an essential way.

The hamiltonian formulation of Einstein's theory was brought into its final form only in 1962 by Arnowitt, Deser and Misner (ADM). They demonstrated that the configuration space of gravity can be understood as consisting of all spatial geometries of three-dimensional space-like hypersurfaces of spacetime that would define arbitrary simultaneities (time coordi-

nates) t . This hamiltonian form is therefore also known as “geometro-dynamics”. Its canonical quantization leads to a Schrödinger equation in the sense of Tomonaga for the wave functional on all these geometries, and it is known as the *Wheeler-DeWitt equation*.

This equation is remarkable insofar as it is of the time-independent form $H\Psi = 0$. The reason for this property is that the spatial metric that occurs as its argument determines proper times (“physical times”) along all time-like curves connecting it with another spatial geometry, while the value of the time coordinate t has no physical meaning. So in spite of this formal timelessness, the Wheeler-DeWitt equation *does* depend on (physical) time by means of the entanglement between matter and geometry that it describes. In general, this physical time is many-fingered (it depends on the progression of the space-like hypersurfaces of spacetime at any space point), but in the case of an exactly homogenous and isotropic Friedmann cosmology, time may be represented by *one* single “finger”: the expansion parameter a . If regarded as a probability amplitude, however, the wave function would define probabilities *for* time – not as a function *of* time.

It is further remarkable that for Friedmann type universes the apparently static Wheeler-DeWitt equation assumes a hyperbolic form in its infinite-dimensional configuration space – again with the time-like variable a . This property is very important, since it allows one to define an initial value problem for the wave functional with respect to this variable – for example at $a = 0$.¹³ Claus Kiefer could even show that one may derive the usual time-dependent Schrödinger equation for matter from the Wheeler-DeWitt equation under a short wave length approximation for the geometric degrees of freedom – in analogy to a Born Oppenheimer approximation (see Ch. 4 of Ref. 5). This result demonstrates once more that the Wheeler-DeWitt equation can only describe a whole Everett multiverse, since each trajectory through the configuration space of spatial geometries would define a classical spacetime. Wave packets following such trajectories according to the WKB approximation are decohered from one another by the matter variables – in analogy to large molecules whose nuclei are observed following quasi-classical orbits.

These remarks about quantum gravity may bring the strange story of particles and waves in principle to an at least preliminary end. While the particle concept has been recognized as a mere illusion, waves exist only as part of a global “wave function” in a very high-dimensional (if not infinite-dimensional) space. Although classical concepts, such as particles and fields, remain essential for our every-day life (including that in physical laboratories), their limited validity must deeply affect our physical world view (cosmology, in particular).

The *observed* quantum world (our “relative state”) can be no more than a very small (but dynamically autonomous in the future) partial wave of the global wave function. We have to accept, however, that the precise structure of a fundamental Hilbert space basis, which is usually assumed to be given by the configuration space of some fundamental fields, remains unknown. Because of the unavoidable entanglement of all fields (“dressing”, renormalization, etc.) one cannot expect the *observed* fields to be related to such fundamental variables in a simple way. Rather, they seem to be only “effective fields” (comparable to phonon fields in solid bodies) – a conclusion that puts in doubt the whole traditional approach to QFT.

References:

¹ E. Segrè, *From Falling Bodies to Radio Waves – Classical Physicists and their Discoveries* (Freeman, NY 1984)

² M. Jammer, *The Conceptual Development of Quantum Mechanics* (McGraw-Hill 1966)

³ S. Deléglise et al., Reconstruction of non-classical cavity field states with snapshots of their decoherence, *Nature* **455**, 510 (2008) – cf. also Fig. 3.14 from Ref. 5

⁴ H. D. Zeh, The role of the observer in the Everett interpretation, arxiv:1211.0196

⁵ E. Joos et al., *Decoherence and the Appearance of a Classical World in Quantum Theory* (Springer 2003), Chs. 1-4

⁶ H. D. Zeh, Quantum Discreteness is an Illusion, *Found. Phys.* **40**, 1476 (2010)

⁷ P. Byrne, *The Many Worlds of Hugh Everett III* (Oxford UP, 2010), Ch. 27

⁸ S. Tomonaga, On a Relativistically Invariant Formulation of the Quantum Theory of Wave Fields, *Progr. Theor. Phys.* **1**, 27 (1946)

⁹ H. D. Zeh, There is no ‘first’ quantization, *Phys. Lett.* **A309**, 329 (2003)

¹⁰ D. Giulini, C. Kiefer, and H. D. Zeh, Symmetries, superselection rules, and decoherence, *Phys. Lett.* **A199**, 291 (1995)

¹¹ F. J. Dyson, The Radiation Theories of Tomonaga, Schwinger and Feynman, *Phys. Rev.* **75**, 486 (1949)

¹² C. Kiefer, *Quantum Gravity*, 2nd edn. (Oxford UP, 2007)

¹³ H. D. Zeh, Emergence of classical time from a universal wave function, *Phys. Lett.* **A116**, 9 (1986); The nature and origin of time-asymmetric spacetime structures, arxiv:1012.4708