# A Fixed Point Theorem and Equilibria of Abstract Economies with w-Upper Semicontinuous Set-Valued Maps

Carlos Hervés-Beloso<sup>1</sup> and Monica Patriche<sup>2</sup>

1 University of Vigo e-mail: cherves@uvigo.es 2 University of Bucharest e-mail: monica.patriche@yahoo.com

The date of receipt and acceptance will be inserted by the editor

Abstract We introduce the notion of w-upper semicontinuous set valued maps

Key words Fixed point theorem, w-upper semicontinuous set valued maps,

2010 Mathematics Subject Classification. 47H10, 91A47, 91A80.

# 1. Introduction

The pioneer work of Nash [11] first proved a theorem of equilibrium existence for games where the player's payoffs are represented by continuous quasi-concave utilities. Arrow and Debreu used the work by Nash to prove the existence of equilibrium in a generalized N-person game or on abstract economy [7] which implies the Walrasian equilibrium existence [2]. These ideas were extended by various authors in several ways. In [16], Shafer and Sonnenschein proved the existence of equilibrium of an economy with finite dimensional commodity space and irreflexive preferences represented as set valued maps with open graph. Yannelis and Prahbakar [22] developed new techniques based on selection theorems and fixed-point theorems. Their main result concerns the existence of equilibrium when the constraint and preference set valued maps have open lower sections. They work within different frameworks (countable infinite number of agents, infinite dimensional strategy spaces).

Borglin and Keiding [3] used new concepts of K.F.-set valued maps and KF-majorized set valued maps for their existence results . The concept of

*Correspondence to*: 1 RGEA, Facultad de Económicas, Universidad de Vigo, Campus Universitario, E-36310 Vigo, Spain

<sup>2</sup> University of Bucharest, Faculty of Mathematics and Computer Science, 14 Academiei Street, 010014 Bucharest, Romania

KF-majorized set valued maps was extended by Yannelis and Prabhakar [22] to L-majorized set valued maps. In [23], Yuan proposed a more general model of abstract economy than the one introduced by Borglin and Keing in [3], in the sense that the constraint mapping was split into two parts A and B. This is due to the "small" constraint set valued map A which could not have enough fixed points even though the "big" constraint set valued map B could.

Most existence theorems of equilibrium deal with preference set valued maps which have lower open sections or are majorized by set valued maps with lower open sections. In the last few years, some existence results were obtained for lower semicontinuous and upper semicontinuous set valued maps. Some recent results concerning upper semicontinuous set valued maps and fixed points can be found in [1], [4], [18], [19], [20], [24]. New results on equilibrium existence in games are given in [10], [12], [13], [17].

In this paper, we define two types of set valued maps: w-upper semicontinuous set valued maps and set valued maps that have e-USS-property. We prove a fixed point theorem for *w*-upper semicontinuous set valued maps. This result is a Wu like result [20] and generalizes the Himmelberg's fixed point theorem in [9]. We use this theorem for proving our first theorem of equilibrium existence for abstract economies having w-upper semicontinuous constraint and preference set valued maps. On the other hand, we use a technique of approximation to prove an equilibrium existence theorem for set valued maps having e-USS-property.

The paper is organized in the following way: Section 2 contains preliminaries and notations. The fixed point theorem is presented in Section 3 and the equilibrium theorems are stated in Section 4.

#### 2. Preliminaries and Notation

Throughout this paper, we shall use the following notations and definitions:

Let A be a subset of a topological space X.  $2^A$  denotes the family of all subsets of A. clA denotes the closure of A in X. If A is a subset of a vector space, coA denotes the convex hull of A. If  $F, G : X \rightrightarrows Y$  are set valued maps, then conv G, cl G,  $G \cap F : X \rightrightarrows Y$  are set valued maps defined by  $(\operatorname{conv} G)(x) = \operatorname{conv} G(x), (\operatorname{cl} G)(x) = \operatorname{cl} G(x) \text{ and } (G \cap F)(x) = G(x) \cap F(x)$ for each  $x \in X$ , respectively. The graph of  $T : X \rightrightarrows Y$  is the set Gr  $(T) = \{(x, y) \in X \times Y \mid y \in T(x)\}.$ 

The set valued map  $\overline{T}$  is defined by  $\overline{T}(x) := \{y \in Y : (x, y) \in cl_{X \times Y} \text{ Gr} T\}$  (the set  $cl_{X \times Y}$  Gr (T) is called the adherence of the graph of T). It is easy to see that  $cl T(x) \subset \overline{T}(x)$  for each  $x \in X$ .

Let X, Y be topological spaces and  $T: X \rightrightarrows Y$  be a set valued map. T is said to be *upper semicontinuous* if for each  $x \in X$  and each open set V in Y with  $T(x) \subset V$ , there exists an open neighborhood U of x in X such that  $T(y) \subset V$  for each  $y \in U$ . T is said to be *almost upper semicontinuous* if for each  $x \in X$  and each open set V in Y with  $T(x) \subset V$ , there exists an open neighborhood U of x in X such that  $T(y) \subset V$  for each  $y \in U$ . **Lemma 2.1**(Lemma 3.2, pag. 94 in [25]) Let X be a topological space, Y be a topological linear space, and let  $S: X \rightrightarrows Y$  be an upper semicontinuous set valued map with compact values. Assume that the set  $C \subset Y$  is closed and  $K \subset Y$  is compact. Then  $T: X \rightrightarrows Y$  defined by  $T(x) = (S(x) + C) \cap K$ for all  $x \in X$  is upper semicontinuous.

Lemma 2.2 is a version of Lemma 1.1 in [21] (for D = Y, we obtain Lemma 1.1 in [21]). For the reader's convenience, we include its proof below. **Lemma 2.2** Let X be a topological space, Y be a nonempty subset of a locally convex topological vector space E and  $T: X \rightrightarrows Y$  be a set valued map. Let  $\beta$  be a basis of neighbourhoods of 0 in E consisting of open absolutely convex symmetric sets. Let D be a compact subset of Y. If for each  $V \in \beta$ , the set valued map  $T^V: X \rightrightarrows Y$  is defined by  $T^V(x) = (T(x) + V) \cap D$  for each  $x \in X$ , then  $\cap_{V \in B} \overline{T^V}(x) \subset \overline{T}(x)$  for every  $x \in X$ .

*Proof* Let x and y be such that  $y \in \bigcap_{V \in B} \overline{T^V}(x)$  and suppose, by way of contradiction, that  $y \notin \overline{T}(x)$ . This means that  $(x, y) \notin cl \ Gr \ T$ , so that there exists an open neighborhood U of x and  $V \in \beta$  such that:

Choose  $W \in \beta$  such that  $W - W \subseteq V$  (e.g.  $W = \frac{1}{2}V$ ). Since  $y \in T^W(x)$ , then  $(x, y) \in cl$  Gr  $T^W$ , so that

$$(U \times (y + W)) \cap \operatorname{Gr} T^W \neq \emptyset.$$

There are some  $x' \in U$  and  $w' \in W$  such that  $(x', y + w') \in \text{Gr } T^W$ , i.e.  $y + w' \in T^W(x')$ . Then,  $y + w' \in D$  and y + w' = y' + w'' for some  $y' \in T(x')$  and  $w'' \in W$ . Hence,  $y' = y + (w' - w'') \in y + (W - W) \subseteq y + V$ , so that  $T(x') \cap (y+V) \neq \emptyset$ . Since  $x' \in U$ , this means that  $(U \times (y+V)) \cap \operatorname{Gr} T \neq \emptyset$ , contradicting (1). 

We introduce the following definitions.

Let X be a topological space, Y be a nonempty subset of a topological vector space E and D be a subset of Y.

**Definition 2.1** The set valued map  $T: X \Rightarrow Y$  is said to be *w*-upper semicontinuous (weakly upper semicontinuous) with respect to the set D if there exists a basis  $\beta$  of open symmetric neighborhoods of 0 in E such that, for each  $V \in \mathcal{B}$ , the set valued map  $T^V$  is upper semicontinuous.

**Definition 2.2** The set valued map  $T: X \rightrightarrows Y$  is said to be *almost w-upper* semicontinuous (almost weakly upper semicontinuous) with respect to the set D if there exists a basis  $\beta$  of open symmetric neighborhoods of 0 in E such that, for each  $V \in \mathfrak{G}$ , the set valued map  $\overline{T^V}$  is upper semicontinuous.

*Example 2.1* Let  $T_1: (0,2) \rightrightarrows (0,2)$  be defined by  $T_1(x) := \begin{cases} (0,1) \text{ if } x \in (0,1]; \\ [1,2) \text{ if } x \in (1,2). \end{cases}$ 

 $T_1$  and  $T_1 \cap \{1\} := \begin{cases} \phi & \text{if } x \in (0,1]; \\ \{1\} & \text{if } x \in (1,2) \end{cases}$  are not upper semicontinuous on (0, 2).

Let  $D := \{1\}$  and let  $V := (-\varepsilon, \varepsilon), \varepsilon > 0$ , be an open symmetric neighbourhood of 0 in IR. Then, it results that

for 
$$\varepsilon > 0$$
,  $T_1(x) + (-\varepsilon, \varepsilon) := \begin{cases} (-\varepsilon, 1+\varepsilon) & \text{if} \quad x \in (0,1];\\ (1-\varepsilon, 2+\varepsilon) & \text{if} \quad x \in (1,2); \end{cases}$   
$$\frac{T_1^V(x) := (T_1(x) + (-\varepsilon, \varepsilon)) \cap \{1\} = \{1\} \text{ for any } x \in (0,2).$$
  
$$\frac{T_1^V(x) = \{1\} \text{ for } x \in (0,2). \end{cases}$$

For each  $V = (-\varepsilon, \varepsilon)$  with  $\varepsilon > 0$ , the set valued maps  $T_1^V$  and  $\overline{T_1^V}$  are upper semicontinuous and  $\overline{T_1^V}$  has nonempty values. We conclude that  $T_1$  is w-upper semicontinuous with respect to D and it is also almost w-upper semicontinuous with respect to D.

We also define the dual w-upper semicontinuity with respect to a compact set.

**Definition 2.3** Let  $T_1, T_2 : X \rightrightarrows Y$  be set valued maps. The pair  $(T_1, T_2)$  is said to be *dual almost w-upper semicontinuous* (dual almost weakly upper semicontinuous) with respect to the set D if there exists a basis  $\beta$  of open symmetric neighborhoods of 0 in E such that, for each  $V \in \beta$ , the set valued map  $\overline{T_{(1,2)}^V} : X \rightrightarrows D$  is lower semicontinuous, where  $T_{(1,2)}^V : X \rightrightarrows D$  is defined by  $T_{(1,2)}^V(x) := (T_1(x) + V) \cap T_2(x) \cap D$  for each  $x \in X$ .

Example 2.2 Let  $D := [1,2], T_1 : (0,2) \Rightarrow [1,4]$  be the set valued map defined by

$$T_1(x) := \begin{cases} [2-x,2], & \text{if } x \in (0,1); \\ \{4\} & \text{if } x = 1; \\ [1,2] & \text{if } x \in (1,2). \end{cases}$$
  
and  $T_2 : (0,2) \Rightarrow [2,3]$  be the set valued map defined by  
 $T_2(x) := \begin{cases} [2,3], & \text{if } x \in (0,1]; \\ \{2\} & \text{if } x \in (1,2); \end{cases}$ .

The set valued map  $T_1$  is not upper semicontinuous on (0, 2).

For  $\varepsilon \in (0,2]$ ,  $(T_1(x) + (-\varepsilon,\varepsilon)) \cap D \cap T_2(x) = \begin{cases} \{2\} \text{ if } x \in (0,1) \cup (1,2); \\ \phi & \text{if } x = 1. \end{cases}$ For  $\varepsilon \in (2,\infty)$ ,  $(T_1(x) + (-\varepsilon,\varepsilon)) \cap D \cap T_2(x) = \{2\}$  for each  $x \in (0,2)$ . Then, we have that for each  $\varepsilon > 0$ ,  $\overline{T_{(1,2)}^V}(x) = \{2\}$  for each  $x \in [0,2]$ 

and the set valued map  $\overline{T^V_{(1,2)}}$  is upper semicontinuous and has nonempty values.

We conclude that the pair  $(T_1, T_2)$  is dual almost w-upper semicontinuous with respect to D.

### 3. A New Fixed Point Theorem

We obtain the following fixed point theorem which generalizes Himmelberg's fixed point theorem in [9]:

**Theorem 3.1** Let I be an index set. For each  $i \in I$ , let  $X_i$  be a nonempty convex subset of a Hausdorff locally convex topological vector space  $E_i$ ,  $D_i$  be a nonempty compact convex subset of  $X_i$  and  $S_i, T_i : X := \prod_{i \in I} X_i \rightrightarrows X_i$ 

be two set valued maps with the following conditions:

1) for each  $x \in X$ ,  $\overline{S}_i(x) \subseteq T_i(x)$ ;

2)  $S_i$  is almost w-upper semicontinuous with respect to  $D_i$  and  $S_i^{V_i}$  is convex nonempty valued for each absolutely convex symmetric neighborhood  $V_i$  of 0 in  $E_i$ .

Then there exists  $x^* \in D := \prod_{i \in I} D_i$  such that  $x_i^* \in T_i(x^*)$  for each  $i \in I$ .

 $\begin{array}{l} Proof \mbox{ Since } D_i \mbox{ is compact}, D := \prod_{i \in I} D_i \mbox{ is also compact in } X. \mbox{ For each } i \in I, \\ \mbox{let } \mbox{$\mathfrak{B}_i$ be a basis of open absolutely convex symmetric neighborhoods of zero in } E_i \mbox{ and let } \mbox{$\mathfrak{B}=\prod_{i \in I} \mbox{$\mathfrak{B}_i$}. \mbox{ For each system of neighborhoods } V = (V_i)_{i \in I} \in \prod_{i \in I} \mbox{$\mathfrak{B}_i$}, \\ \mbox{let's define the set valued maps } S_i^{V_i} : X \rightrightarrows D_i, \mbox{ by } S_i^{V_i}(x) = (S_i(x) + V_i) \cap D_i, \\ x \in X, \ i \in I. \mbox{ By assumption } 2) \mbox{ each } \overline{S_i^{V_i}} \mbox{ is u.s.c with nonempty closed convex values. Let's define } S^V : X \rightrightarrows D \mbox{ by } S^V(x) = \prod_{i \in I} \overline{S_i^{V_i}}(x) \mbox{ for each } \end{array}$ 

 $x \in D$ . The set valued map  $S^V$  is upper semicontinuous with closed convex values. Therefore, according to Himmelberg's fixed point theorem [9], there exists  $x_V^* = \prod_{i \in I} (x_V^*)_i \in D$  such that  $x_V^* \in S^V(x_V^*)$ . It follows that  $(x_V^*)_i \in \overline{V}$ 

 $\overline{S_i^{V_i}}(x_V^*)$  for each  $i \in I$ .

For each  $V = (V_i)_{i \in I} \in \mathcal{B}$ , let's define  $Q_V = \bigcap_{i \in I} \{x \in D : x_i \in \overline{S_i^{V_i}}(x)\}.$ 

 $Q_V$  is nonempty since  $x_V^* \in Q_V$ , then  $Q_V$  is nonempty and closed.

We prove that the family  $\{Q_V : V \in \beta\}$  has the finite intersection property.

Let  $\{V^{(1)}, V^{(2)}, ..., V^{(n)}\}$  be any finite set of  $\beta$  and let  $V^{(k)} = \prod_{i \in I} V_i^{(k)}$ , k = 1, ..., n. For each  $i \in I$ , let  $V_i = \bigcap_{k=1}^n V_i^{(k)}$ , then  $V_i \in \beta_i$ ; thus  $V = \prod_{i \in I} V_i \in \prod_{i \in I} \beta_i$ . Clearly  $Q_V \subseteq \bigcap_{k=1}^n Q_{V^{(k)}}$  so that  $\bigcap_{k=1}^n Q_{V^{(k)}} \neq \emptyset$ .

Since D is compact and the family  $\{Q_V : V \in \beta\}$  has the finite intersection property, we have that  $\cap\{Q_V : V \in \beta\} \neq \emptyset$ . Take any  $x^* \in \cap\{Q_V : V \in \beta\}$ , then for each  $V_i \in \beta_i$ ,  $x_i^* \in \overline{S_i^{V_i}}(x^*)$ . According to Lemma 2.2, we have that  $x_i^* \in \overline{S_i}(x^*)$ , for each  $i \in I$ , therefore  $x_i^* \in T(x^*)$ .  $\Box$ 

If |I| = 1 we get the result below.

**Corollary 3.1** Let X be a nonempty subset of a Hausdorff locally convex topological vector space F, D be a nonempty compact convex subset of X and  $S, T : X \rightrightarrows X$  be two set valued maps with the following conditions:

1) for each  $x \in X$ ,  $\overline{S}(x) \subseteq T(x)$  and  $S(x) \neq \emptyset$ ,

2) S is almost w-upper semicontinuous with respect to D and  $\overline{S^V}$  is convex valued for each open absolutely convex symmetric neighborhood V of 0 in E.

Then, there exists a point  $x^* \in D$  such that  $x^* \in T(x^*)$ .

In the particular case that the set valued map S = T the following result stands.

**Corollary 3.2** Let X be a nonempty subset of a Hausdorff locally convex topological vector space F, D be a nonempty compact convex subset of X and  $T: X \rightrightarrows X$  be an almost w- upper semicontinuous set valued map with respect to D and  $\overline{T^V}$  is convex valued for each open absolutely convex symmetric neighborhood V of 0 in E. Then, there exists a point  $x^* \in D$ such that  $x^* \in \overline{T}(x^*)$ .

### 4. Application in the Equilibrium Theory

## 4.1 The Model of an Abstract Economy

We will consider further Yuan's model of an abstract economy (see [23]). Let I be a nonempty set (the set of agents). For each  $i \in I$ , let  $X_i$  be a non-empty subset of a topological vector space representing the agent's iset of actions and define  $X := \prod_{i \in I} X_i$ ; let  $A_i, B_i : X \rightrightarrows X_i$  be the constraint

set valued maps and  $P_i$  the preference set valued map.

**Definition 4.1** [23] An abstract economy  $\Gamma = (X_i, A_i, P_i, B_i)_{i \in I}$  is defined as a family of ordered quadruples  $(X_i, A_i, P_i, B_i)$ .

**Definition 4.2 [23]** An *equilibrium* for  $\Gamma$  is defined as a point  $x^* \in X$  such that for each  $i \in I$ ,  $x_i^* \in \overline{B}_i(x^*)$  and  $A_i(x^*) \cap P_i(x^*) = \emptyset$ .

Remark 4.1 When, for each  $i \in I$ ,  $A_i(x) = B_i(x)$  for all  $x \in X$ , the abstract economy model coincides with the classical one introduced by Borglin and Keiding in [3]. If in addition,  $\overline{B}_i(x^*) = \operatorname{cl}_{X_i} B_i(x^*)$  for each  $x \in X$ , which is the case where  $B_i$  has a closed graph in  $X \times X_i$ , the definition of equilibrium coincides with the one used by Yannelis and Prabhakar in [22].

Remark 4.2 If the preference set valued map  $P_i$  is defined by using a utility function  $u_i$ , that is  $P_i(x) = \{y \in X_i : u_i(y) > u_i(x_i)\}$ , the irreflexibility condition  $x_i \notin \overline{P_i}(x)$ , which appears among the hypothesis of the existence equilibrium theorems, may fail. A case in which this condition is verified, is when  $P_i$  is an order interval preference. Order interval preferences are studied, for instance, in Chateauneuf [5]. These preference relations  $\prec$  (on X) are representable, if two real valued functions u and v on X exist and are such that:  $x \prec y \Leftrightarrow u(x) < v(y)$ . If a representation of the preference relation  $\prec_i$  exist, we can define the preference set valued map  $P_i$  by  $P_i(x) =$  $\{y \in X_i : v_i(y) > u_i(x_i)\}$  and the condition  $x_i \notin \overline{P_i}(x)$  can be fulfilled.

# 4.2 Examples of Abstract Economies with Two Constraint Set Valued Maps

A first example of an abstract economy with two constraint set valued maps is the one associated to the model proposed by Radner [14] and his followers. This is a model of a pure exchange economy with two periods, present and future, and uncertainty on the state of nature in the future. There is a finite number n of agents and a finite number m of possible future states of nature. Let  $I = \{1, 2, ..., n\}$  be the set of agents and  $\Omega = \{s_1, s_2, ..., s_m\}$  the set of the future states of the nature. Each agent has his own private, and tipically incomplete, information about the future state of nature. For each agent i in I, the initial private information is a partition on  $\Omega$ , induced by

a signal  $\pi_i: \Omega \to Y_i$ . In Radner [14], agents make decisions today without knowing the future state of nature tomorrow. The initial agent's information is kept fixed and their consumption plans need to be made compatible with their information, in the sense that their consumption must be the same in states that they do not distinguish. In a different framework, Radner [15] consider the notion of rational expectation equilibrium. In this model, agents are able to forecast the future equilibrium price. Consequently, their initial information is updated with a signal given by the future equilibrium prices p and a more refined partition of  $\Omega$  is obtained as the joint of the initial information and the information generated by  $\hat{\pi}_i(p): \Omega \to \Delta$ , defined by  $\hat{\pi}_i(p)(s) = p_i(s)$ , where  $\Delta$  be the normalized set of prices. Here, we consider a model in which agents may be able to learn from market signals. These market signal are summarized by equilibrium prices that may not be fully revealing. Without loss of generality, let denote the joint of the initial information and the information generated by  $\hat{\pi}_i$ .

For agent i in I, the consumption plan in the first period will be denoted by  $x_0^i \in IR_+^l$  and in the second period, for each state  $s_j, j = 1, 2, ..., m$ , it will be denoted by  $x_j^i \in IR_+^l$ . A bundle for agent i is  $x^i = (x_0^i, x_1^i, x_2^i, ..., x_m^i)$ . Let  $X_i = IR_+^{ml+1}$ . Each agent has a preference set valued map  $Q'_i : \prod_{i \in I} X_i \rightrightarrows X_i$ 

and an initial endowment  $e^i = (e_0^i, e_1^i, e_2^i, ..., e_m^i) \in X_i$ .

**Definition 4.3** A pure exchange economy with asymmetric information is the family  $\mathcal{E} = (I, \Omega, \widehat{\pi}_i, Q'_i, e^i)_{i \in I}$ .

**Definition 4.4** An allocation for the economy  $\mathcal{E}$  is  $x = (x^i)_{i \in I}$ . The allocation is called *phisically feasible* if  $\sum_{i \in I} x^i \leq \sum_{i \in I} e^i$  and *informationally feasible* for each agent i if  $\widehat{\pi}_i(p)(s) = \widehat{\pi}_i(p)(s')$  implies  $x^i_s = x^i_{s'}$ .

Let  $p_0$  be the price in the first period, for the second period let  $p_j$  be the price in the state j, j = 1, 2, ..., m and let  $p = (p_0, p_1, ..., p_m)$ . Let  $\Delta$  be the normalized set of prices. Without loss of generality, we assume that pbelongs to  $\Delta$ .

The budget set valued map of agent *i* is  $B_i : \Delta \rightrightarrows IR_+^{lm+1}$ , defined by  $B_i(p) = \{x^i \in IR_+^{lm+1} : px^i < pe^i\}.$ 

The information set valued map of agent *i* is  $I_i : \Delta \rightrightarrows IR_+^{lm+1}$ , defined by  $I_i(p) = \{x^i \in IR_+^{lm+1} : x_s^i = x_{s'}^i \text{ if } \widehat{\pi}_i(p)(s) = \widehat{\pi}_i(p)(s')\}.$ 

**Definition 4.5** The pair  $(x^*, p^*) \in IR^{n(lm+1)}_+ \times \Delta$  is an *equilibrium* for the asymmetrically informed economy  $\mathcal{E}$  if

1)  $\sum_{i \in I} (x^*)^i \leq \sum_{i \in I} e^i$ and for each  $i \in I$ , 2)  $(x^*)^i \in \overline{I_i}(p^*) \cap \overline{B_i}(p^*);$ 3)  $y^i \in Q'_i(x^*) \cap I_i(p^*)$  implies that  $y^i \notin B_i(p^*)$ .

Let  $X := \prod_{i \in I} X_i \times \Delta$ , where for  $i \in I$ ,  $X_i = IR_+^{lm+1}$  is the consumption set of agent *i*. Let's define the following set valued maps:

-for each  $i \in I$ ,  $Q_i : X \rightrightarrows X_i$  is the preference set valued map defined by  $Q_i(x, p) = Q'_i(x)$  for each  $(x, p) \in X$ ;

 $\begin{array}{l} -Q_{n+1}:X \rightrightarrows \Delta \text{ is the preference set valued map defined by } Q_{n+1}(x,p) := \\ \{q \in \Delta: q(\sum_{i \in I} (x^i - e^i)) > p(\sum_{i \in I} (x^i - e^i))\} \text{ for each } (x,p) \in X; \end{array}$ 

-for  $i \in I$ ,  $A_i : X \rightrightarrows 2^{X_i}$  is defined by  $A_i(x, p) := \{y^i \in IR^{lm+1}_+ : py^i < pe^i\}$  for each  $(x, p) \in X$ ;

 $\begin{array}{l} -A_{n+1}: X \rightrightarrows \varDelta \text{ is defined by } A_{n+1}(x,p) := \varDelta \text{ for each } (x,p) \in X; \\ \text{-for } i \in I, \ I_i: X \rightrightarrows X_i \text{ is defined by } I_i(x,p) := \{y^i \in IR_+^{lm+1}: y^i_s = y^i_{s'} \text{ if } \widehat{\pi}_i(p)(s) = \widehat{\pi}_i(p)(s')\} \text{ for each } (x,p) \in X; \end{array}$ 

-  $I_{n+1}: X \rightrightarrows \Delta$  is defined by  $I_{n+1}(x,p) := \Delta$  for each  $(x,p) \in X$ ;

**Definition 4.6** The abstract economy associated to the model of the pure exchange economy with asymmetric information is  $\Gamma = (X_i, A_i, P_i, B_i)_{i \in \{1, 2, ..., n+1\}}$ , where:

-for  $i \in I$ ,  $X_i := IR_+^{lm+1}$  is the consumption set of agent i and let  $X := \prod X_i \times \Delta;$ 

 $-P_i : X \Rightarrow X_i \ (i \in I)$  and  $P_{n+1} : X \Rightarrow \Delta$  are the preference set valued maps defined by  $P_i(x,p) = Q_i(x,p) \cap I_i(x,p)$  for each  $(x,p) \in X$  and  $i \in \{1, 2, ..., n+1\}$ ;

 $-A_i: X \rightrightarrows X_i \ (i \in I)$  and  $A_{n+1}: X \rightrightarrows \Delta$  are the constraint set valued maps defined above;

 $-B_i: X \Rightarrow X_i \ (i \in I)$  and  $B_{n+1}: X \Rightarrow \Delta$  are the constraint set valued maps defined by  $B_i(x,p) := A_i(x,p) \cap I_i(x,p)$  for each  $(x,p) \in X$  and  $i \in \{1, 2, ..., n+1\}.$ 

Remark 4.3 We note that  $A_i(x,p) \cap P_i(x,p) \subseteq B_i(x,p)$  for each  $(x,p) \in X$  and for each  $i \in \{1, 2, ..., n+1\}$ .

**Proposition 4.1** An equilibrium for the associated abstract economy  $\Gamma$  is an equilibrium of the economy with asymmetric information E.

Proof Let  $(x^*, p^*)$  be an equilibrium for  $\Gamma$ .

1) For each  $i \in \{1, 2, ..., n\}$ , we have that  $(x^*)^i \in \overline{B_i}(x^*, p^*) = \overline{(A_i \cap I_i)}(x^*, p^*)$ and then, by definition of  $A_i$  and  $I_i$ ,  $(x^*)^i \in \overline{(I_i \cap B_i)}(p^*)$ ;

2)  $p^* \in \overline{B_{n+1}}(x^*, p^*) = \Delta;$ 

3) for each  $i \in \{1, 2, ..., n\}$ , we have that  $A_i(x^*, p^*) \cap P_i(x^*, p^*) = \phi$ , which implies that if  $y^i \in P_i(x^*, p^*) = Q_i(x^*, p^*) \cap I_i(x^*, p^*)$ , then  $y^i \notin A_i(x^*, p^*)$ ; This means that  $y^i \in Q'_i(x^*) \cap I_i(p^*)$  implies that  $y^i \notin B_i(p^*)$ ;

4) we have that  $A_{n+1}(x^*, p^*) \cap P_{n+1}(x^*, p^*) = \phi$ , which is equivalent with  $\{q \in \Delta : q(\sum_{i \in I} ((x^*)^i - e^i)) > p^*(\sum_{i \in I} ((x^*)^i - e^i))\} \cap \Delta = \phi$ . This fact implies that  $q(\sum_{i \in I} ((x^*)^i - e^i)) \leq p^*(\sum_{i \in I} ((x^*)^i - e^i)) \leq 0$  for all  $q \in \Delta$ . If we choose q as a vector of the canonical basis of  $IR^{ml+1}$ , that is  $q_j = 1$  and  $q_i = 0$  for  $i \neq j$ , where  $i, j \in \{1, 2, ..., ml + 1\}$ , we obtain that  $\sum_{i \in I} (x^*)^i \leq \sum_{i \in I} e^i$ .  $\Box$ 

The second example is the abstract economy associated to an exchange economy with two constraint set valued maps, the first one being the budget

set valued map and the second one being the consumption set that depends on prices.

The third example follows the idea of an exchange economy which has, beyond the budget set valued map, a second constraint set valued map  $G_i$ , defined by the delivery conditions as stated in the paper by Correia-da-Silva and Herves-Beloso [6].

Let's assume that the set of the states of nature is  $\Omega = \{1, 2, ..., m\},\$ the future prices are  $p_1, p_2, ..., p_m \in IR^l_+$  and that each agent *i* has a signal  $f_i: \Omega \to Y_i$  such that  $f_i(s) = f_i(s')$  if s and s' are states that cannot be distinguished. The agent *i* chooses a portfolio y(s) in the following way:

 $p_s y(s) \le p_s y(s')$  for all s' such that  $f_i(s') = f_i(s)$ .

The set valued map  $G_i: X \times \Delta \rightrightarrows IR^{lm}$  is defined by  $G_i(x,p) = \{y \in$  $IR^{lm}: p_s y(s) \le p_s y(s')$  for all s' such that  $f_i(s') = f_i(s)$ .

# 4.3 The Existence of Equilibria in Locally Convex Spaces

As an application of the fixed point Theorem 3.1, we have the following result.

**Theorem 4.1** Let  $\Gamma = \{X_i, A_i, B_i, P_i\}_{i \in I}$  be an abstract economy such that for each  $i \in I$ , the following conditions are fulfilled:

1)  $X_i$  is a nonempty convex subset of a Hausdorff locally convex topological vector space  $E_i$  and  $D_i$  is a nonempty compact convex subset of  $X_i;$ 

2) for each  $x \in X := \prod_{i \in I} X_i$ ,  $A_i(x)$  and  $P_i(x)$  are convex,  $B_i(x)$  is nonempty, convex and  $A_i(x) \cap P_i(x) \subset B_i(x)$ ;

3)  $W_i := \{x \in X : A_i(x) \cap P_i(x) \neq \emptyset\}$  is open in X.

4)  $H_i: X \rightrightarrows X_i$  defined by  $H_i(x) := A_i(x) \cap P_i(x)$  for each  $x \in X$  is almost w-upper semicontinuous with respect to  $D_i$  on  $W_i$  and  $\overline{H_i^{V_i}}$  is convex nonempty valued for each open absolutely convex symmetric neighborhood  $V_i$  of 0 in  $E_i$ ;

5)  $B_i: X \rightrightarrows X_i$  is almost w-upper semicontinuous with respect to  $D_i$  and  $\overline{B_i^{V_i}}$  is convex nonempty valued for each open absolutely convex symmetric neighborhood  $V_i$  of 0 in  $E_i$ ;

6) for each  $x \in X$ ,  $x_i \notin \overline{(A_i \cap P_i)}(x)$ ; Then there exists  $x^* \in D = \prod_{i \in I} D_i$  such that  $x_i^* \in \overline{B}_i(x^*)$  and  $(A_i \cap P_i)$ 

 $P_i(x^*) = \emptyset$  for each  $i \in I$ .

*Proof* Let  $i \in I$ . By condition (3) we know that  $W_i$  is open in X.

Let's define  $T_i : X \rightrightarrows X_i$  by  $T_i(x) := \begin{cases} A_i(x) \cap P_i(x), & \text{if } x \in W_i, \\ B_i(x), & \text{if } x \notin W_i \end{cases}$  for each  $x \in X$ .

Then  $T_i: X \rightrightarrows X_i$  is a set valued map with nonempty convex values. We shall prove that  $T_i: X \Rightarrow D_i$  is almost w-upper semicontinuous with respect to  $D_i$ . Let  $\beta_i$  be a basis of open absolutely convex symmetric neighborhoods of 0 in  $E_i$  and let  $\beta = \prod \beta_i$ .

For each  $V = (V_i)_{i \in I} \in \prod_{i \in I} \beta_i$ , for each  $x \in X$ , let for each  $i \in I$   $B^{V_i}(x) := (B_i(x) + V_i) \cap D_i$ ,  $F^{V_i}(x) := ((A_i(x) \cap P_i(x)) + V_i) \cap D_i$  and  $T_i^{V_i}(x)^{"} = \begin{cases} F^{V_i}(x), \text{ if } x \in W_i, \\ B^{V_i}(x), \text{ if } x \notin W_i. \end{cases}$ For each open set  $V_i'$  in  $D_i$ , the set  $\left\{x \in X : \overline{T_i^{V_i}}(x) \subset V_i'\right\} =$   $= \left\{x \in W_i : \overline{F^{V_i}}(x) \subset V_i'\right\} \cup \left\{x \in X \smallsetminus W_i : \overline{B^{V_i}}(x) \subset V_i'\right\}$  $= \left\{x \in W_i : \overline{F^{V_i}}(x) \subset V_i'\right\} \cup \left\{x \in X : \overline{B^{V_i}}(x) \subset V_i'\right\}.$ 

According to condition (4), the set  $\left\{x \in W_i : \overline{F^{V_i}}(x) \subset V'_i\right\}$  is open in X. The set  $\left\{x \in X : \overline{B^{V_i}}(x) \subset V'_i\right\}$  is open in X because  $\overline{B^{V_i}}$  is upper semicontinuous.

Therefore, the set  $\left\{x \in X : \overline{T_i^{V_i}}(x) \subset V_i'\right\}$  is open in X. It shows that  $\overline{T_i^{V_i}}: X \Rightarrow D_i$  is upper semicontinuous. According to Theorem 3.1, there exists  $x^* \in D = \prod_{i \in I} D_i$  such that  $x^* \in \overline{T_i}(x^*)$ , for each  $i \in I$ . By condition (5) we have that  $x_i^* \in \overline{B_i}(x^*)$  and  $(A_i \cap P_i)(x^*) = \emptyset$  for each  $i \in I$ .  $\Box$ 

Example 4.1 Let  $\Gamma = \{X_i, A_i, B_i, P_i\}_{i \in I}$  be an abstract economy, where  $I = \{1, 2, ..., n\}, X_i := [0, 4]$  be a compact convex choice set,  $D_i := [0, 2]$  for each  $i \in I$  and  $X := \prod_{i \in I} X_i$ .

Let the set valued maps  $A_i, B_i, P_i : X \rightrightarrows X_i$  be defined as follows: for each  $(x_1, x_2, ..., x_n) \in X$ ,

$$A_{i}(x) := \begin{cases} [1 - x_{i}, 2] \text{ if } x \in (0, \frac{1}{2})^{n}; \\ [1 - x_{i}, 2) \text{ if } x \in [\frac{1}{2}, 1)^{n}; \\ [3, 4] & \text{if } x = 0; \\ [0, \frac{1}{2}], & \text{otherwise}; \end{cases}$$
$$P_{i}(x) := \begin{cases} [\frac{3}{2}, 2 + x_{i}] \text{ if } x \in [0, 1)^{n}; \\ [1, 2], & \text{otherwise}; \end{cases}$$
$$B_{i}(x) := \begin{cases} [0, 2] \text{ if } x \in [0, 1); \\ [3, 4] & \text{if } x = 0; \\ [0, 2), & \text{otherwise}. \end{cases}$$

The set valued maps  $A_i, B_i, P_i$  are not upper semicontinuous on X.

$$A_i(x) \cap P_i(x) := \begin{cases} [\frac{3}{2}, 2] & \text{if } x \in (0, \frac{1}{2})^n; \\ [\frac{3}{2}, 2) & \text{if } x \in [\frac{1}{2}, 1)^n; \\ \phi, & \text{otherwise.} \end{cases}$$
$$W_i := \{x \in X : A_i(x) \cap P_i(x) \neq \emptyset\} = (0, 1)^n \text{ is open in } X.$$
$$\overline{(A_i \cap P_i)}(x) := \begin{cases} [\frac{3}{2}, 2] & \text{if } x \in [0, 1]^n; \\ \phi, & \text{otherwise.} \end{cases}$$
We notice that for each  $x \in X$ ,  $x_i \notin \overline{(A_i \cap P_i)}(x)$ .

We shall prove that  $B_i$  and  $(A_i \cap P_i)_{W_i}$  are almost w-upper semicontinuous with respect to  $D_i = [0, 2]$ .

On  $W_i$ ,

 $\begin{array}{l} (A_i \cap P_i) \left( x \right) := \begin{cases} \left[ \frac{3}{2}, 2 \right] \text{ if } x \in (0, \frac{1}{2})^n; \\ \left[ \frac{3}{2}, 2 \right] \text{ if } x \in \left[ \frac{1}{2}, 1 \right)^n; \\ (A_i \cap P_i) \left( x \right) + \left( -\varepsilon, \varepsilon \right) = \left( \frac{3}{2} - \varepsilon, 2 + \varepsilon \right) \text{ if } x \in (0, 1)^n; \\ \text{Let } \left( A_i \cap P_i \right)^V \left( x \right) = \left( \left( A_i \cap P_i \right) \left( x \right) + \left( -\varepsilon, \varepsilon \right) \right) \cap [0, 2], \text{ where } V = \left( -\varepsilon, \varepsilon \right). \end{cases}$ Then, if  $\varepsilon \in (0, \frac{3}{2}]$ ,  $(A_i \cap P_i)^V(x) = (\frac{3}{2} - \varepsilon, 2]$  if  $x \in (0, 1)^n$ ; if  $\varepsilon > \frac{3}{2}$ ,  $(A_i \cap P_i)^V(x) = [0, 2]$  if  $x \in (0, 1)^n$ ; Hence, for each  $V = (-\varepsilon, \varepsilon)$ ,  $(A_i \cap P_i)^V_{W_i}$  is upper semicontinuous and

has nonempty values.  $(-\varepsilon \ 2+\varepsilon)$  if  $x \in (0,1)^n$ :

$$B_{i}(x) + (-\varepsilon, \varepsilon) = \begin{cases} (-\varepsilon, 2+\varepsilon) \text{ if } x \in (0,1), \\ (3-\varepsilon, 4+\varepsilon) & \text{if } x = 0; \\ (-\varepsilon, 2+\varepsilon) & \text{otherwise.} \end{cases}$$
Let  $B_{i}^{V}(x) = (B_{i}(x) + (-\varepsilon, \varepsilon)) \cap [0,2]$ , where  $V = (-\varepsilon, \varepsilon)$ .  
Then,  
if  $\varepsilon \in (0,1]$ ,  
 $B_{i}^{V}(x) = \begin{cases} \phi & \text{if } x = 0; \\ [0,2] & \text{otherwise;} \end{cases}$   
if  $\varepsilon \in (1,3], B_{i}^{V}(x) = \begin{cases} [0,2] & \text{if } x \in [0,1)^{n}; \\ (3-\varepsilon,2] & \text{if } x = 0; \\ [0,2], & \text{otherwise.} \end{cases}$ 

and if  $\varepsilon > 3$ ,  $B_i^V(x) = [0, 2]$  if  $x \in X$ .

Then, for each  $V = (-\varepsilon, \varepsilon)$ ,  $\overline{B_i^V}$  is upper semicontinuous and has nonempty values.

Therefore, all hypotheses of Theorem 4.1 are satisfied, so that there exist equilibrium points. For example,  $x^* = \{\frac{3}{2}, \frac{3}{2}, ..., \frac{3}{2}\} \in X$  verifies  $x_i^* \in \overline{B}_i(x^*)$ and  $(A_i \cap P_i)(x^*) = \emptyset$ .

Theorem 4.2 deals with abstract economies which have dual w-upper semicontinuous pairs of set valued maps.

**Theorem 4.2** Let  $\Gamma = \{X_i, A_i, B_i, P_i\}_{i \in I}$  be an abstract economy such that for each  $i \in I$ , the following conditions are fulfilled:

1)  $X_i$  is a nonempty convex subset of a Hausdorff locally convex topological vector space  $E_i$  and  $D_i$  is a nonempty compact convex subset of  $X_i$ ;

2) for each  $x \in X := \prod_{i \in I} X_i$ ,  $P_i(x) \subset D_i$ ,  $A_i(x) \cap P_i(x) \subset B_i(x)$  and  $B_i(x)$  is nonempty;

3) the set  $W_i := \{x \in X : A_i(x) \cap P_i(x) \neq \emptyset\}$  is open in X;

4) the pair  $(A_{i|c|W_i}, P_{i|c|W_i})$  is dual almost w-upper semicontinuous with respect to  $D_i, B_i : X \rightrightarrows X_i$  is almost w-upper semicontinuous with respect to  $D_i$ ;

5) if  $T_{i,V_i} : X \rightrightarrows X_i$  is defined by  $T_{i,V_i}(x) := (A_i(x) + V_i) \cap D_i \cap P_i(x)$  for each  $x \in X$ , then the set valued maps  $\overline{B_i^{V_i}}$  and  $\overline{T_{i,V_i}}$  are nonempty convex valued for each open absolutely convex symmetric neighborhood  $V_i$  of 0 in  $E_i$ ;

 $\begin{array}{l} E_i;\\ 6) \ for \ each \ x \in X \ , \ x_i \notin \overline{P}_i \ (x);\\ Then, \ there \ exists \ x^* \in D := \prod_{i \in I} D_i \ such \ that \ x_i^* \in \overline{B}_i \ (x^*) \ and \ A_i \ (x^*) \cap P_i \ (x^*) = \emptyset \ for \ all \ i \in I. \end{array}$ 

*Proof* For each  $i \in I$ , let  $\beta_i$  denote the family of all open absolutely convex symmetric neighborhoods of zero in  $E_i$  and let  $\beta = \prod_{i \in I} \beta_i$ . For each V =

$$\begin{split} &\prod_{i \in I} V_i \in \prod_{i \in I} \mathbb{B}_i, \text{ for each } i \in I, \text{ let} \\ &B^{V_i}(x) := (B_i(x) + V_i) \cap D_i \text{ for each } x \in X \text{ and} \\ &S_i^{V_i}(x) := \begin{cases} T_{i,V_i}(x), & \text{ if } x \in W_i, \\ B_i^{V_i}(x), & \text{ if } x \notin W_i, \end{cases} \end{split}$$

 $\overline{S_i^{V_i}}$  has closed values. Next, we shall prove that  $\overline{S_i^{V_i}}: X \Rightarrow D_i$  is upper semicontinuous.

For each open set 
$$V'$$
 in  $D_i$ , the set  

$$\left\{x \in X : \overline{S_i^{V_i}}(x) \subset V'\right\} =$$

$$= \left\{x \in W_i : \overline{T_{i,V_i}}(x) \subset V'\right\} \cup \left\{x \in X \smallsetminus W_i : \overline{B_i^{V_i}}(x) \subset V'\right\}$$

$$= \left\{x \in W_i : \overline{T_{i,V_i}}(x) \subset V'\right\} \cup \left\{x \in \underline{X} : \overline{B_i^{V_i}}(x) \subset V'\right\}.$$

We know that the set valued map  $\overline{T_{i,V_i}}(x)|_{W_i}: W_i \Rightarrow D_i$  is upper semicontinuous. The set  $\{x \in W_i : \overline{T_{i,V_i}}(x) \subset V'\}$  is open in X. Since  $\overline{B_i^{V_i}}(x):$  $X \Rightarrow D_i$  is upper semicontinuous, the set  $\{x \in X : \overline{B_i^{V_i}}(x)\} \subset V'$  is open in X and therefore, the set  $\{x \in X : \overline{S_i^{V_i}}(x) \subset V'\}$  is open in X. It proves that  $\overline{S_i^{V_i}}: X \Rightarrow D_i$  is upper semicontinuous. According to Himmelberg's Theorem, applied for the set valued maps  $\overline{S_i^{V_i}}$ , there exists a point  $x_V^* \in D =$  $\prod_{i \in I} D_i$  such that  $(x_V^*)_i \in S_i^{V_i}(x_V^*)$  for each  $i \in I$ . By condition (5), we have that  $(x_{V_i}^*)_i \notin \overline{P_i}(x_V^*)$ , hence,  $(x_V^*)_i \notin \overline{A_i^{V_i}}(x_V^*) \cap \overline{P_i}(x_V^*)$ .

that  $(x_V^*)_i \notin \overline{P_i}(x_V^*)$ , hence,  $(x_V^*)_i \notin \overline{A_i^{V_i}}(x_V^*) \cap \overline{P_i}(x_V^*)$ . We also have that cl Gr  $(T_{i,V_i}) \subseteq$  cl Gr  $(A_i^{V_i}) \cap$ cl Gr  $P_i$ . Then  $\overline{T_{i,V_i}}(x) \subseteq \overline{A_i^{V_i}}(x) \cap \overline{P_i}(x)$  for each  $x \in X$ . It follows that  $(x_V^*)_i \notin \overline{T_{i,V_i}}(x_V^*)$ . Therefore,  $(x_V^*)_i \in \overline{B^{V_i}}(x_V^*)$ .

For each  $V = (V_i)_{i \in I} \in \prod_{i \in I} \beta_i$ , let's define  $Q_V = \bigcap_{i \in I} \{x \in D : x \in \overline{B^{V_i}}(x) \text{ and } A_i(x) \cap P_i(x) = \emptyset \}.$ 

 $Q_V$  is nonempty since  $x_V^* \in Q_V$ , and it is a closed subset of D according to (3). Then,  $Q_V$  is nonempty and compact.

Let  $\beta = \prod_{i \in I} \beta_i$ . We prove that the family  $\{Q_V : V \in \beta\}$  has the finite intersection property.

Let  $\{V^{(1)}, V^{(2)}, ..., V^{(n)}\}$  be any finite set of  $\beta$  and let  $V^{(k)} = \prod_{i \in I} V_i^{(k)}{}_{i \in I},$ k = 1, ..., n. For each  $i \in I$ , let  $V_i = \bigcap_{k=1}^n V_i^{(k)}$ , then  $V_i \in \beta_i$ ; thus  $V \in \prod_{i \in I} \beta_i$ .

Clearly  $Q_V \subset \bigcap_{k=1}^n Q_{V^{(k)}}$  so that  $\bigcap_{k=1}^n Q_{V^{(k)}} \neq \emptyset$ .

Since D is compact and the family  $\{Q_V : V \in \beta\}$  has the finite intersection property, we have that  $\cap \{Q_V : V \in \beta\} \neq \emptyset$ . Take any  $x^* \in \cap \{Q_V : V \in \beta\}$ , then for each  $V \in \beta$ ,

 $x^* \in \bigcap_{i \in I} \left\{ x^* \in D : x_i^* \in \overline{B^{V_i}}(x) \text{ and } A_i(x) \cap P_i(x) = \emptyset \right\}.$ 

Hence,  $x_i^* \in \overline{B^{V_i}}(x^*)$  for each  $V \in \beta$  and for each  $i \in I$ . According to Lemma 2.2, we have that  $x_i^* \in \overline{B_i}(x^*)$  and  $(A_i \cap P_i)(x^*) = \emptyset$  for each  $i \in I$ .  $\Box$ 

We now introduce the following concept, which also generalizes the concept of lower semicontinuous set valued maps.

**Definition 4.7** Let X be a non-empty convex subset of a topological linear space E, Y be a non-empty set in a topological space and  $K \subseteq X \times Y$ .

The set valued map  $T : X \times Y \rightrightarrows X$  has the e-USCS-property (eupper semicontinuous selection property) on K, if for each absolutely convex neighborhood V of zero in E, there exists an upper semicontinuous set valued map with convex values  $S^V : X \times Y \rightrightarrows X$  such that  $S^V(x,y) \subset$ T(x,y) + V and  $x \notin cl S^V(x,y)$  for every  $(x,y) \in K$ .

The following theorem is an equilibrium existence result for economies with constraint set valued maps having e-USCS-property.

**Theorem 4.3** Let  $\Gamma = (X_i, A_i, P_i, B_i)_{i \in I}$  be an abstract economy, where I is a (possibly uncountable) set of agents such that for each  $i \in I$ :

- (1)  $X_i$  is a non-empty compact convex set in a locally convex space  $E_i$ ;
- (2) cl  $B_i$  is upper semicontinuous with non-empty convex values;

(3) the set  $W_i := \{x \in X / (A_i \cap P_i)(x) \neq \emptyset\}$  is open;

(3) cl  $(A_i \cap P_i)$  has the e-USCS-property on  $W_i$ .

Then there exists an equilibrium point  $x^* \in X$  for  $\Gamma$ , i.e., for each  $i \in I, x_i^* \in \overline{B}_i(x^*)$  and  $A_i(x^*) \cap P_i(x^*) = \emptyset$ .

*Proof* For each  $i \in I$ , let  $\mathfrak{B}_i$  denote the family of all open convex neighborhoods of zero in  $E_i$ . Let  $V = (V_i)_{i \in I} \in \prod_{i \in I} \mathfrak{B}_i$ . Since cl  $(A_i \cap P_i)$  has the

e-USCS-property on  $W_i$ , it follows that there exists an upper semicontinuous set valued map  $F_i^{V_i} : X \rightrightarrows X_i$  such that  $F_i^{V_i}(x) \subset \operatorname{cl} (A_i \cap P_i)(x) + V_i$  and  $x_i \notin \operatorname{cl} F_i^{V_i}(x)$  for each  $x \in W_i$ .

Define the set valued map  $T_i^{V_i}: X \rightrightarrows X_i$ , by

$$T_i^{V_i}(x) := \begin{cases} \operatorname{cl} \{F_i^{V_i}(x)\}, & \text{if } x \in W_i, \\ \operatorname{cl} (B_i(x) + V_i) \cap X_i, & \text{if } x \notin W_i; \end{cases}$$
$$B_i^{V_i} : X \rightrightarrows X_i, B_i^{V_i}(x) := \operatorname{cl} (B_i(x) + V_i) \cap X_i = (\operatorname{cl} B_i(x) + \operatorname{cl} V_i) \cap X_i \text{ is pper semicontinuous by Lemma 2.1}$$

upper semicontinuous by Lemma 2.1.

Let U be an open subset of  $X_i$ , then  $U' := \{x \in X \mid T_i^{V_i}(x) \subset U\}$ 

$$= \{ x \in W_i \mid T_i^{V_i}(x) \subset U \} \cup \{ x \in X \setminus W_i \mid T_i^{V_i}(x) \subset U \}$$
$$= \{ x \in W_i \mid \text{cl } F_i^{V_i}(x) \subset U \} \cup \{ x \in X \mid (\text{cl } B_i(x) + \overline{V_i}) \cap X_i \subset U \}$$

U' is an open set, because  $W_i$  is open,  $\left\{x \in W_i \mid \operatorname{cl} F_i^{V_i}(x) \subset U\right\}$  is open since  $\operatorname{cl} F_i^{V_i}(x)$  is an upper semicontinuous map on  $W_i$ . We have also that the set  $\{x \in X \mid (\operatorname{cl} B_i(x) + \operatorname{cl} V_i) \cap X_i \subset U\}$  is open since  $(\operatorname{cl} B_i + \operatorname{cl} V_i) \cap X_i$  is u.s.c. Then  $T_i^{V_i}$  is upper semicontinuous on X and has closed convex values.

Define 
$$T^V: X \rightrightarrows X$$
 by  $T^V(x) := \prod_{i \in I} T_i^{V_i}(x)$  for each  $x \in X$ .

 $T^V$  is an upper semicontinuous set valued map and has also non-empty convex closed values.

Since X is a compact convex set, by Fan's fixed-point theorem [8], there exists  $x_V^* \in X$  such that  $x_V^* \in T^V(x_V^*)$ , i.e., for each  $i \in I$ ,  $(x_V^*)_i \in T_i^{V_i}(x_V^*)$ . If  $x_V^* \in W_i$ ,  $(x_V^*)_i \in Cl F_i^{V_i}(x_V^*)$ , which is a contradiction.

Hence,  $(x_V^*)_i \in cl (B_i(x_V^*) + V_i) \cap X_i$  and  $(A_i \cap P_i)(x_V^*) = \emptyset$ , i.e.  $x_V^* \in Q_V$  where

 $Q_V = \bigcap_{i \in I} \{ x \in X : x_i \in cl \ (B_i(x) + V_i) \cap X_i \text{ and } (A_i \cap P_i)(x) = \emptyset \}.$ 

Since  $W_i$  is open,  $Q_V$  is the intersection of non-empty closed sets, therefore it is non-empty, closed in X.

We prove that the family  $\{Q_V : V \in \prod_{i \in I} \beta_i\}$  has the finite intersection property.

Let  $\{V^{(1)}, V^{(2)}, ..., V^{(n)}\}$  be any finite set of  $\prod_{i \in I} \beta_i$  and let  $V^{(k)} = (V_i^{(k)})_{i \in I}$ , k = 1, ...n. For each  $i \in I$ , let  $V_i = \bigcap_{k=1}^n V_i^{(k)}$ , then  $V_i \in \beta_i$ ; thus  $V = (V_i)_{i \in I} \in \prod_{i \in I} \beta_i$ . Clearly  $Q_V \subset \bigcap_{k=1}^n Q_{V^{(k)}}$  so that  $\bigcap_{k=1}^n Q_{V^{(k)}} \neq \emptyset$ .

Since X is compact and the family  $\{Q_V : V \in \prod_{i \in I} \beta_i\}$  has the finite intersection property, we have that  $\cap \{Q_V : V \in \prod_{i \in I} \beta_i\} \neq \emptyset$ . Take any  $x^* \in$  $\cap \{Q_V : V \in \prod_{i \in I} \beta_i\}$ , then for each  $i \in I$  and each  $V_i \in \beta_i$ ,  $x_i^* \in \operatorname{cl}(B_i(x^*) + V_i) \cap X_i$  and  $(A_i \cap P_i)(x^*) = \emptyset$ ; but then  $x_i^* \in \overline{B}_i(x^*)$  from Lemma 2.2 and  $(A_i \cap P_i)(x^*) = \emptyset$  for each  $i \in I$  so that  $x^*$  is an equilibrium point of  $\Gamma$  in X.  $\square$ 

# 5. Concluding Remarks

We proved a fixed point theorem for the w-upper semicontinuous set valued maps. We also obtained results concerning the existence of equilibria for Yuan's model of an abstract economy without continuity assumptions.

The first author thanks the support by Research Grants ECO2012-38860-C02-02 (Ministerio de Economia y Competitividad), RGEA and 10PXIB300141PR (Xunta de Galicia and FEDER).

# References

### References

- R. P. Agarwal, O'Regan, 2001. A Note on Equilibria for abstract economies, Mathematical and Computer Modelling 34, 331-343.
- 2. K. J. Arrow, G. Debreu, 1954. Existence of an Equilibrium for a Competitive Economy. Econometrica 22, 265-290.
- 3. A. Borglin and H. Keiding, 1976. Existence of equilibrium action and of equilibrium: A note on the "new" existence theorem. J. Math. Econom. 3, 313-316.
- S.-Y. Chang, 2010. Inequalities and Nash equilibria. Nonlinear Analysis: Theory, Methods & Applications, Vol. 73, 9, 2933-2940.
- 5. A. Chateauneuf, 1987. Continuous representation of a preference relation on a connected topological space, Journal of Mathematical Economics 16, 139-146.
- J. Correia-da-Silva and C. Hervés-Beloso, 2012. General equilibrium in economies with uncertain delivery, Economic Theory, 51 3, 729-755.
- G. Debreu, 1952. A social equilibrium existence theorem. Proc. Nat. Acad. Sci. 38, 886-893.
- 8. K. Fan,1961. A generalization of Tychonoff's fixed point theorem, Math. Ann. 142, 305-310.
- C. J. Himmelberg, 1972. Fixed points of compact multifunctions. J. Math. Anal. Appl. 38, 205-207.
- A. A. Kulkarni, V. Shanbhag, Revisiting Generalized Nash Games and Variational Inequalities. Journal of Optimization Theory and Applications, 154 (1), 175-186 (2012)
- 11. J.F. Nash, 1951. Non-cooperative games. Ann. Math. 54, 286-295.
- R. Nessah, G. Tian, Existence of Solution of Minimax Inequalities, Equilibria in Games and Fixed Points Without Convexity and Compactness Assumptions. J.O.T.A., 2012 DOI 10.1007/s10957-012-0176-5
- 13. M. Patriche, Equilibrium in games and competitive economies. The Publishing House of the Romanian Academy (2011)
- 14. R. Radner, 1968. Competitive Equilibrium Under Uncertainty, Econometrica, 36 1, 31-58.
- 15. R. Radner, 1979. Rational Expectations Equilibrium: Generic Existence and the Information Revealed by Prices. Econometrica, 47 *3*, 655-678.
- W. Shafer, H. Sonnenschein, 1975. Equilibrium in abstract economies without ordered preferences. J. Math. Econom. 2, 345-348.
- A. Stefanescu, M. Ferrara, M. V. Stefanescu, Equilibria of the Games in Choice Form. Journal of Optimization Theory and Applications, 155 (3), 1060-1072 (2012)
- K.K. Tan, Z. Wu, 1998. A Note on Abstract Economies with upper semicontinuous correspondence. Appl. Math. Letters. 5 Vol. 11, 21-22.
- L. A. Tuan, G. M. Lee, P. H. Sach, 2010. Upper semicontinuity in a parametric general variational problem and application. Nonlinear Analysis: Theory, Methods & Applications, Vol. 72, 3-4, 1500-1513.
- X. Wu, 1997. A new fixed point theorem and its applications, Proc. Amer. Math. Soc. 125, 1779-1783.
- X. Wu, 20001. Equilibria of lower semicontinuous games, Computer Math. Appl. 42, 13-22.
- N. C. Yannelis and N. D. Prabhakar, 1983. Existence of maximal elements and equilibrium in linear topological spaces. J. Math. Econom. 12, 233-245.

- X. Z. Yuan, 1998. The Study of Minimax inequalities and Applications to Economies and Variational inequalities. Memoirs of the American Society 132, 625.
- X. Z. Yuan, E. Taradfar, 1999. Maximal elements and equilibria of generalized games for U-majorized and condensing correspondences, Internat. J. Math. Sci. 1, vol 22, 179-189.
- 25. X. Zheng, 1997. Approximate selection theorems and their applications. J. Math. An. Appl. 212, 88-97.