# CYCLIC STABILIZERS AND INFINITELY MANY HYPERBOLIC ORBITS FOR PSEUDOGROUPS ON $(\mathbb{C},0)$

#### JULIO C. REBELO & HELENA REIS

ABSTRACT. Consider a pseudogroup on  $(\mathbb{C},0)$  generated by two local diffeomorphisms having analytic conjugacy classes *a priori* fixed in Diff  $(\mathbb{C},0)$ . We show that a generic pseudogroup as above is such that every point has (possibly trivial) cyclic stabilizer. It also follows that these generic groups possess infinitely many hyperbolic orbits. This result possesses several applications to the topology of leaves of foliations and we shall explicitly describe the case of nilpotent foliations associated to Arnold's singularities of type  $A^{2n+1}$ .

Key-words: pseudogroups on  $(\mathbb{C},0)$  - fixed points - cyclic stabilizers

AMS-Classification: 37C85 - 37F75 - 34M35

### 1. Introduction

This paper provides answers to two types of well-known problems about pseudogroups on  $(\mathbb{C},0)$ . For the convenience of the reader, this Introduction will begin with a detailed presentation of them. The first problem has to do with the existence of hyperbolic fixed points for certain elements of pseudogroups generated by local diffeomorphisms fixing  $0 \in \mathbb{C}$ . The reader is reminded that a fixed point p for a local diffeomorphism f between open sets of  $\mathbb{C}$  is said to be hyperbolic if and only if  $||f'(p)|| \neq 1$ . In particular these fixed points are necessarily isolated, i.e. f cannot coincide with the identity on a neighborhood of a hyperbolic fixed point p. The existence of "generic" pseudogroups for which there are infinitely many points  $p_i, p_i \neq 0$  for every i, such that each  $p_i$  is an hyperbolic fixed point for a certain element  $h_i$  belonging to the pseudogroup in question was first singled out by Y. Il'yashenko in his seminal paper [Il-1]. The meaning of these fixed points for the topology of the leaves of the corresponding foliations is explained in [Il-1], [Il-2]. The more recent paper [Sh-R-O] improves Il'yashenko's result on the existence of the fixed points in question. Another result of similar nature was established in [GM-W] and applies to certain pseudogroups obtained through "perturbations" of pseudogroups generated by local diffeomorphisms fixing  $0 \in \mathbb{C}$ (note that these "perturbed" pseudogroups need no longer have the origin as a common fixed point).

Among the most interesting cases where the existence of hyperbolic fixed points have been considered, there is the case of pseudogroups generated by local diffeomorphisms fixing  $0 \in \mathbb{C}$  and such that the associated group of germs at  $0 \in \mathbb{C}$  is not solvable. Here the existence of elements in the corresponding pseudogroup exhibiting hyperbolic fixed points has been known since [Sh], [N] and [BLL-1], see also [Lo] for a comprehensive discussion of these non-solvable pseudogroups. However the question on whether or not these pseudogroups exhibit more than one single orbit of hyperbolic fixed points, at least in the case of "typical" pseudogroups, has remained open. This question deserves to be elaborated further.

Suppose  $p \neq 0$  happens to be a hyperbolic fixed point for some element h of a given pseudogroup. Then the orbit of p is fully constituted by points that are hyperbolic fixed points for suitable conjugates of h in the pseudogroup in question. The problem that is naturally raised in this context consists of deciding whether the pseudogroup contains two or more hyperbolic fixed points whose orbits are mutually disjoint. This question has a natural motivation going back to foliations: in fact, whereas the existence of an "orbit of hyperbolic fixed points" implies the existence of a leaf carrying hyperbolic holonomy, it falls short from ensuring the existence of "many leaves" with hyperbolic holonomy. The existence of these leaves can only be asserted if hyperbolic fixed points with mutually disjoint orbits is guaranteed.

The other problem concerning  $(\mathbb{C},0)$  that will be tackled in this paper has a more typical "generic nature". It concerns the existence of points that are fixed by "more than one element in the pseudogroup". More precisely, if  $p \in \mathbb{C}$  is fixed by some element h, then p is obviously fixed by every iterate of h as well. The question is then about the existence of some  $\overline{h}$  in the pseudogroup fixing p and not (locally) coinciding with an iterate of h. In other words, the question is whether or not every point different from  $0 \in \mathbb{C}$  has cyclic (possibly trivial) stabilizer.

The answers to the above mentioned problems given in this work sit naturally in the continuation of [M-R-R] as will be seen below. Some terminology is however needed before stating our main results. First, given an element  $f \in \text{Diff}(\mathbb{C},0)$ , we shall denote by  $f^j$  its  $j^{\text{th}}$ -iterate for  $j \in \mathbb{Z}$  ( $f^0 = \text{id}$  and  $f^j = (f^{-1})^{|j|}$  for j < 0). This definition has a clear meaning in terms of germs whereas this meaning is less clear in the context of pseudogroups, cf. Section 2 for details.

Now consider Diff  $(\mathbb{C},0)$  equipped with the analytic topology (cf. Section 2) turning Diff  $(\mathbb{C},0)$  into a Baire space. Denote by  $\mathrm{Diff}_{\alpha}(\mathbb{C},0)$  the normal subgroup of Diff  $(\mathbb{C},0)$  consisting of those germs of diffeomorphisms that are tangent to the identity to order  $\alpha \in \mathbb{N}$  (if  $\alpha = 0$  then  $\mathrm{Diff}_{\alpha}(\mathbb{C},0) = \mathrm{Diff}(\mathbb{C},0)$ ). The subgroup  $\mathrm{Diff}_{\alpha}(\mathbb{C},0) \subseteq \mathrm{Diff}(\mathbb{C},0)$  is closed for the analytic topology and, in fact, it is a Baire space in its own right. Next, let  $\mathrm{Diff}_{\alpha}(\mathbb{C},0) \times \mathrm{Diff}_{\alpha}(\mathbb{C},0)$  be endowed with the product analytic topology. Suppose we are given two elements f,g in  $\mathrm{Diff}(\mathbb{C},0)$  and denote by  $G_1$  (resp.  $G_2$ ) the cyclic group generated by f (resp. g). Naturally, the groups  $G_1,G_2$  may or may not be finite and their orders are respectively the orders of the germs f,g which are denoted by f and f. In other words, f (resp. f) is the smallest strictly positive integer for which f in this case, the group f integer does not exist, then we set f is f (resp. f and f in this case, the group f (resp. f and f in this case, the group f (resp. f and f in this case, the group f (resp. f and f in this case, the group f (resp. f and f in this case, the group f (resp. f and f in this case, the group f (resp. f and f in this case, the group f (resp. f and f in this case, the group f (resp. f and f in this case, the group f (resp. f and f in this case, the group f (resp. f and f in this case, the group f in the group f in this case, the group f in this case, the group f in this case, the group f in the group f in this case, the group f in the group f in the group f in this case, the group f in the grou

Now denote by  $F_2$  the free group on two generators a, b and consider the natural evaluation morphism from  $F_2$  to Diff  $(\mathbb{C}, 0)$  consisting of making the substitutions  $a \mapsto f$  and  $b \mapsto g$  (and of interpreting the "concatenation of letters" as composition of germs). Let N be the normal subgroup of  $F_2$  generated by  $\{a^r, b^s\}$ , with the convention that  $a^{\infty} = b^{\infty} = \mathrm{id}$ . The quotient group  $F_2/N$  is isomorphic to the free product  $G_1 * G_2$  of the groups  $G_1, G_2$ . Furthermore, the above mentioned evaluation morphism factors through the quotient  $F_2/N$  so as to induce a homomorphism  $\mathcal{E}$  from  $G_1 * G_2$  to Diff  $(\mathbb{C}, 0)$ .

Another explicit construction for the homomorphism  $\mathcal{E}$  consists of using the fact that every element in the free product  $G_1 * G_2$  is represented by a *unique* reduced word in the letters a, b, where the empty-word represents the identity (cf. Section 2 for further details).

Therefore, the elements of  $G_1 * G_2$  are naturally identified to reduced words W(a, b). With this notation,  $\mathcal{E}(W(a, b))$  is simply the element of Diff  $(\mathbb{C}, 0)$  obtained by substituting  $a \mapsto f$  and  $b \mapsto g$  in the spelling of W(a, b) (where again the "concatenation of letters" becomes composition of germs). In the sequel, the element of Diff  $(\mathbb{C}, 0)$  given by  $\mathcal{E}(W(a, b))$  is going to be denoted by W(f, g).

To state Theorem A, recall that a local diffeomorphism f fixing  $0 \in \mathbb{C}$  is linearizable if and only if it is conjugate to the linear map  $z \mapsto f'(0) z$  by a local holomorphic change of coordinates, where f'(0) stands for the derivative of f at  $0 \in \mathbb{C}$ . This local diffeomorphism is said to have a Cremer point (at  $0 \in \mathbb{C}$ ) if it is not linearizable and if f'(0) has norm 1 but is not a root of unity. Suppose now that we are given local diffeomorphisms f, g such that none of them has a Cremer point at  $0 \in \mathbb{C}$ . If W(a, b) is a non-empty reduced word in a, b (w.r.t  $G_1 * G_2$ ), the following was shown in [M-R-R] (cf. Theorem 3.1): there is a  $G_{\delta}$ -dense  $\mathcal{V} \subset \operatorname{Diff}_{\alpha}(\mathbb{C}, 0) \times \operatorname{Diff}_{\alpha}(\mathbb{C}, 0)$  such that, whenever  $(h_1, h_2) \in \mathcal{U}$ , the element  $W(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)$  of the pseudogroup generated by  $h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2$  does not coincide with the identity on any connected component of its domain of definition. Equivalently, all fixed points of  $W(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)$  are isolated, though not necessarily hyperbolic. Here we shall improve on this result by proving Theorem A below.

In the sequel  $\alpha \in \mathbb{N}$  is fixed as well as local diffeomorphisms f, g. Given local diffeomorphisms  $h_1, h_2$ , we denote by  $\Gamma_{h_1, h_2}$  the pseudogroup generated by  $h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2$  on some local neighborhood of  $0 \in \mathbb{C}$  (to be chosen later).

**Theorem A.** Suppose we are given f, g in  $\mathrm{Diff}_{\alpha}(\mathbb{C}, 0)$  and denote by D an open disc about  $0 \in \mathbb{C}$  where f, g and their inverses are defined. Assume that none of the local diffeomorphisms f, g has a Cremer point at  $0 \in \mathbb{C}$ . Then, there is a  $G_{\delta}$ -dense set  $\mathcal{U} \subset \mathrm{Diff}_{\alpha}(\mathbb{C}, 0) \times \mathrm{Diff}_{\alpha}(\mathbb{C}, 0)$  such that, whenever  $(h_1, h_2)$  lies in  $\mathcal{U}$ , the pseudogroup  $\Gamma_{h_1, h_2}$ generated by  $\tilde{f} = h_1^{-1} \circ f \circ h_1$ ,  $\tilde{g} = h_2^{-1} \circ g \circ h_2$  on D satisfies the following:

- (1) The stabilizer of every point  $p \in D$  is either trivial or cyclic.
- (2) There is a sequence of points  $\{Q_i\}$ ,  $Q_i \neq 0$  for every  $i \in \mathbb{N}^*$ , converging to  $0 \in \mathbb{C}$  and such that every  $Q_n$  is a hyperbolic fixed point of some element  $W_i(\tilde{f}, \tilde{g}) \in \Gamma_{h_1, h_2}$ . Furthermore the orbits under  $\Gamma_{h_1, h_2}$  of  $Q_{n_1}$ ,  $Q_{n_2}$  are disjoint provided that  $n_1 \neq n_2$ .

Note that, in the statement of Theorem A, the analytic conjugacy classes of f and g in Diff  $(\mathbb{C},0)$  are supposed to be fixed. This condition is naturally imposed by the use of the Krull topology in classical problems about singular foliations, cf. below.

The assumption that neither f nor g has a Cremer point at  $0 \in \mathbb{C}$  is indeed necessary for the statement of Theorem A to hold, cf. Section 2. On the other hand, the theorem holds equally well for pseudogroups generated by every finite collection of local diffeomorphisms. Some comments on possible improvements of Theorem A also deserve to be included here. Essentially these improvements concern extensions "generic/general" for the corresponding statement. As to item (2), it is conceivable that the existence of "more than one" orbit of hyperbolic fixed points may be verified for every non-solvable pseudogroup of Diff ( $\mathbb{C}$ , 0). On the other hand, item (1) is unlikely to be a general phenomenon and we believe that only a "generic" affirmative answer can be expected. A more optimistic perspective might suggest that an affirmative answer can still hold true for "open and dense sets" though this already seems a bit unlikely to happen. In any event, in order to make further progress in this

type of questions, it seems clear that the next step is to investigate the possible existence of "Closing lemmas" for the mentioned pseudogroups.

Theorem A has a few consequences on the topology of the leaves of a foliation on a neighborhood of an invariant curve and/or a neighborhood of the singular point since in most well-known problems this information can be codified into the holonomy pseudogroup associated to the invariant curve (or to the reduction of the singular point). An important class of applications stems from classical problems abut deciding the topology of leaves for a (local) "generic" foliation, where "generic" is understood in terms of dense sets for the Krull topology (see [LF], [M-M], [M-R-R], [M-S]). These applications are by now standard so that they will not be detailed in this paper (apart from stating Corollary B). On the other hand, for the convenience of non-experts, let us explain why the use of Krull topology forces us to fix the conjugacy classes of the initial local diffeomorphisms. For this, suppose that  $\mathcal{F}$  is the singular foliation associated to the local orbits of a holomorphic vector field X defined on a neighborhood of  $(0,0) \in \mathbb{C}^2$ . Let  $\widetilde{\mathcal{F}}$  denote the transform of  $\mathcal{F}$  by a suitable birational transformation so that  $\widetilde{\mathcal{F}}$  is defined on a neighborhood of an invariant divisor. Suppose that the singularities of  $\widetilde{\mathcal{F}}$  are all hyperbolic (an assumption frequently satisfied). Note that, in the present context, a singular point p of  $\widetilde{\mathcal{F}}$  is said to be hyperbolic if  $\widetilde{\mathcal{F}}$  can locally be given by a holomorphic vector field Y whose linear part at p has non-zero eigenvalues  $\lambda_1, \lambda_2$ verifying  $\lambda_1/\lambda_2 \in \mathbb{C} \setminus \mathbb{R}$ . A classical problem in differential equations/singularity theory considers dynamical/topological properties of foliations  $\mathcal{F}$  as before that are satisfied by Krull-dense sets of foliations. In practice, this means that we should look for properties that are verified by foliations  $\mathcal{F}'$  given by vector fields X' having the same Taylor series as X up to an arbitrarily fixed order. Since the invariant divisor of  $\mathcal{F}$  as well as the position of its singular points and the values of corresponding eigenvalues are determined by some finite jet of X, these data cannot be changed in the construction of the "perturbed" foliation. Thus the holonomy maps arising from corresponding singular points of  $\widetilde{\mathcal{F}}$  and of the "perturbed foliation" are necessarily conjugate to each other since they are both linearizable in view of Poincaré theorem. In other words, while constructing foliations "near" to  $\mathcal{F}, \widetilde{\mathcal{F}}$ , the analytic conjugacy classes of the mentioned holonomy maps are necessarily fixed.

In general, the transverse structure of a (singular) holomorphic foliation is described through the pseudogroup associated to the holonomy with respect to an invariant curve. The existence of hyperbolic fixed points, for example, is strictly related to the presence of hyperbolic limit cycles for the foliation in question. Though we have mentioned hyperbolic singularities to illustrate the interest of having fixed analytic conjugacy classes in the statement of Theorem A, we shall explicitly state an application of this theorem in a rather "orthogonal" setting where singularities are far from hyperbolic. This setting corresponds to the much studied case of nilpotent foliations associated to Arnold  $A^{2n+1}$  singularities.

Recall that a nilpotent foliation about  $(0,0) \in \mathbb{C}^2$  is the singular foliation associated to the local orbits of a (local) holomorphic vector field X having an isolated singularity at  $(0,0) \in \mathbb{C}^2$  where the linear part of X at (0,0) is nilpotent (different from zero). Consider then nilpotent foliations possessing a unique separatrix which is given by a cusp of the form  $\{y^2 + x^{2n+1} = 0\}$ , i.e. the separatrix is an analytic curve locally equivalent to the mentioned cusp. Foliations in this class are called nilpotent foliation  $\mathcal{F}$  of type  $A^{2n+1}$ . In [M-R-R], it

was proved that a generic nilpotent foliation  $\mathcal{F}$  of type  $A^{2n+1}$  possesses only countably many non-simply connected leaves.

Let  $X \in \mathfrak{X}_{(\mathbb{C}^2,0)}$  be a holomorphic vector field with an isolated singularity at the origin and defining a germ of nilpotent foliation  $\mathcal{F}$  of type  $A^{2n+1}$ . By relying on the construction carried out in Section 5 of [M-R-R], Theorem A yields:

Corollary B (Cusps). For every (sufficiently large)  $N \in \mathbb{N}$ , there exists a vector field  $X' \in \mathfrak{X}_{(\mathbb{C}^2,0)}$  defining a germ of a foliation  $\mathcal{F}'$  and satisfying the following conditions:

- (1)  $J_0^N X' = J_0^N X$  (i.e. the vector field X, X' are tangent to order N at the origin).
- (2) The foliations  $\mathcal{F}$  and  $\mathcal{F}'$  have S as a common separatrix.
- (3) There exists a fundamental system of open neighborhoods  $\{U_n\}_{n\in\mathbb{N}}$  of S, inside a closed ball  $\bar{B}(0,R)$ , such that for all  $n\in\mathbb{N}$ , the leaves of the restriction of  $\mathcal{F}'$  to  $U_n\setminus S$  are simply connected except for a countable set of them.
- (4) The countable set mentioned above is, in fact, infinite.
- (5) Every leaf of the restriction of  $\mathcal{F}'$  to  $U_n \setminus S$  is either simply-connected or topologically equivalent to a cylinder.

Compared to the analogous statement in [M-R-R], the improvements made in this paper lies in items (4) and (5). The remaining items make up the previous result in [M-R-R] and depend solely on the fact that, for a "generic choice" of  $(h_1, h_2) \in \operatorname{Diff}_{\alpha}(\mathbb{C}, 0) \times \operatorname{Diff}_{\alpha}(\mathbb{C}, 0)$ , every element in the resulting pseudogroup  $\Gamma_{h_1,h_2}$  coinciding with the identity on a nonempty open set must coincide with the identity on all of its domain of definition. Naturally items (4) and (5) are implied by our Theorem A.

The proof of Theorem A naturally starts from the above mentioned result for the pseudogroup  $\Gamma_{h_1,h_2}$ . However it requires a rather different type of analysis which is vaguely reminiscent from the classical Kupka-Smale theorems or, more precisely, with the part of its statement asserting that "generic" diffeomorphisms have only hyperbolic periodic points, cf. for example [K-H]. In closing this Introduction, let us give a brief outline of the structure of the proof of Theorem A. Compared to Kupka-Smale statement, the first difference is naturally the fact that we are dealing with a pseudogroup of local diffeomorphisms rather than with a single globally defined diffeomorphism. This requires us to pay special attention to domains of definitions as well as to the boundary behavior of maps. In Section 2, an appropriate setting to handle these pseudogroups will be worked out in detail.

The main analogy with Kupka-Smale theorem appears at the level of "stability" of hyperbolic fixed points. The idea of "stability" for hyperbolic fixed points is materialized by saying that once a (local) diffeomorphism possessing a (unique) hyperbolic fixed point is perturbed, the new diffeomorphism will also possess a (unique) hyperbolic fixed point which, in addition, is "near" the initial fixed point. In our holomorphic context, there is no need to worry about fixed points being hyperbolic since the desired "stability" property is verified by "multiplicity one" fixed points of (local) holomorphic diffeomorphisms. In fact, this type of "stability" follows at once from the Argument principle (or Rouché theorem). It is this application of the Argument principle that, ultimately, will allow us to keep track of the number of fixed points that may arise when a diffeomorphism having only isolated fixed points is perturbed. This material is carefully developed in Section 3 which also contains the proof of Theorem A modulo Proposition 3.6 whose proof, at this point, will essentially

be reduced to the construction of certain types of perturbations for pairs of local diffeomorphisms  $(h_1, h_2) \in \operatorname{Diff}_{\alpha}(\mathbb{C}, 0) \times \operatorname{Diff}_{\alpha}(\mathbb{C}, 0)$ . These constructions will then be detailed in Sections 4 and 5. In particular, Section 4 begins with an outline of the structure of the proof of Proposition 3.6 itself. As it is to be expected, our "perturbations" have no analogue in the Kupka-Smale context, not only due to the holomorphic nature of our problem, but mainly due to the fact that the conjugacy classes for the initial diffeomorphisms f, g have to be fixed during all the procedure.

**Acknowledgments**. We are very grateful to J.-F. Mattei for several discussions about the content of this paper as well as for having encouraged us to work on this problem.

The authors thank the IMPA and the UMI-CNRS for their hospitality during the preparation of this paper. The first author also thanks CNPq-Brazil for financial support. The second author thanks FCT for financial support through CMUP and through the project PTDC/MAT/103319/2008.

#### 2. General set up

In the sequel, let  $\mathrm{Diff}\,(\mathbb{C},0)$  stand for the group of local holomorphic diffeomorphisms fixing  $0 \in \mathbb{C}$  whereas  $\mathrm{Diff}_{\alpha}(\mathbb{C},0)$  stands for its normal subgroup consisting of elements tangent to the identity to order  $\alpha$  ( $\alpha=0$  corresponds to  $\mathrm{Diff}\,(\mathbb{C},0)$  itself). Also, let  $\mathrm{Hol}\,(\mathbb{C},0)$  denote the space of (germs of) holomorphic functions defined about  $0 \in \mathbb{C}$ . Clearly  $\mathrm{Diff}\,(\mathbb{C},0) \subset \mathrm{Hol}\,(\mathbb{C},0)$  and an element  $f \in \mathrm{Hol}\,(\mathbb{C},0)$  belongs to  $\mathrm{Diff}\,(\mathbb{C},0)$  if and only if  $f'(0) \neq 0$ .

Let us equip both Diff  $(\mathbb{C}, 0)$  and  $\operatorname{Hol}(\mathbb{C}, 0)$  with the so-called analytic topology (or  $C^{\omega}$ -topology) that was first considered by Takens [T] in the context of real analytic diffeomorphisms of an analytic manifold who also observed that it inherits of the Baire property. The definition of the analytic topology can naturally be adapted to  $\operatorname{Hol}(\mathbb{C}, 0)$  and it was shown in [M-R-R] that it turns  $\operatorname{Hol}(\mathbb{C}, 0)$  into a complete metric space. The metric is defined as follows. Suppose that f, g in  $\operatorname{Hol}(\mathbb{C}, 0)$  are given and consider the holomorphic function f - g which is defined in a neighborhood of  $0 \in \mathbb{C}$ . Denote by  $c_1x + c_2x^2 + \cdots$  the Taylor series of f - g at  $0 \in \mathbb{C}$ . The metric  $d_A$  inducing the analytic topology in  $\operatorname{Hol}(\mathbb{C}, 0)$  is then given by

$$d_A(f,g) = \sup_{k \in \mathbb{N}} ||c_k||^{1/k}.$$

Being a complete metric space,  $\operatorname{Hol}(\mathbb{C},0)$  has the Baire property. Since  $\operatorname{Diff}(\mathbb{C},0)$  is clearly an open and dense subset of  $\operatorname{Hol}(\mathbb{C},0)$ , we recover the fact that  $\operatorname{Diff}(\mathbb{C},0)$  has the Baire property as well. Furthermore, if  $\alpha \geq 1$ ,  $\operatorname{Diff}_{\alpha}(\mathbb{C},0)$  is a complete metric space in its own right and hence it also possesses the Baire property.

Let f, g be given elements in Diff  $(\mathbb{C}, 0)$  and denote by  $G_1, G_2$  the cyclic group generated by f, g, respectively. Fix a sufficiently small open disc D about  $0 \in \mathbb{C}$  where f, g and their inverses are defined. To begin the discussion concerning Theorem A a few notions need to be recalled. Also, while in the statement of Theorem A and in the discussion below, we shall restrict our attention to pseudogroups generated by two local diffeomorphisms f, g, all the arguments immediately carry over to pseudogroups generated by a finite number of local diffeomorphisms.

Let us begin by recalling the formal notion of a reduced word in two letters. Let  $f, g \in \text{Diff}(\mathbb{C}, 0)$  be two holomorphic diffeomorphisms fixing the origin of  $\mathbb{C}$  and assume that they are both different from the identity. Denote by r (resp. s) the order of f (resp. g), namely

- (1)  $\vartheta_i$  takes on the values  $\{a, b\}$ .
- (2) If  $\vartheta_{i_0}$  takes on the value a (resp. b) then  $\vartheta_{i_0-1}$  and  $\vartheta_{i_0+1}$  take on the value b (resp. a) provided that  $\vartheta_{i_0-1}$  and  $\vartheta_{i_0+1}$  are defined.
- (3) If  $\vartheta_i$  takes on the value a, then  $r_i$  takes values in the set  $\{1, \ldots, r-1\}$  provided that  $r < \infty$ . If  $r = \infty$ , then  $r_i$  takes values in  $\mathbb{Z}^*$  (it is understood that, for  $r_i < 0$ ,  $a^{r_i}$  means  $(a^{-1})^{|r_i|}$ ).
- (4) Similarly, if  $\vartheta_i$  takes on the value b, then  $r_i$  takes values in the set  $\{1, \ldots, s-1\}$  provided that  $s < \infty$ . If  $s = \infty$ , then  $r_i$  takes values in  $\mathbb{Z}^*$  (where  $b^{r_i}$  means  $(b^{-1})^{|r_i|}$  whenever  $r_i < 0$ ).

With the previous notations, let us consider the free product  $G_1 * G_2$  between  $G_1$  and  $G_2$ . In terms of presentation, this group is isomorphic to the group defined by  $\{a, b; a^{\mathsf{r}} = b^{\mathsf{s}} = \mathrm{id}\}$ , where the relation  $a^{\mathsf{r}} = \mathrm{id}$  (resp.  $b^{\mathsf{s}} = \mathrm{id}$ ) is understood to be void if  $\mathsf{r} = \infty$  (resp.  $\mathsf{s} = \infty$ ). In the sequel we shall use this presentation to refer to the free product  $G_1 * G_2$ . It should be noticed that every element in  $\{a, b; a^{\mathsf{r}} = b^{\mathsf{s}} = \mathrm{id}\}$  is represented by a unique reduced word W(a, b) (where the neutral element corresponds to the empty word).

The fundamental object involved in the subsequent discussion is the notion of pseudogroup generated by local diffeomorphisms f and g about  $0 \in \mathbb{C}$ . For local diffeomorphisms f, g as above, consider an open disc D about  $0 \in \mathbb{C}$  where  $f, f^{-1}, g, g^{-1}$  are all defined and one-to-one. We want to consider the pseudogroup  $\Gamma = \Gamma(f, g, D)$  generated by  $f, f^{-1}, g, g^{-1}$  on D (in the sequel this pseudogroup will be referred to as being generated by f, g and their inverses or simply by f, g, when no confusion is possible). An accurate definition of this pseudogroup is required for the subsequent discussion. Consider a fixed word  $W(a, b) = \vartheta_l^{r_l} * \cdots * \vartheta_1^{r_1}$ . In the sequel, unless otherwise stated, every word is supposed to be reduced with respect to the group  $\{a,b; a^r = b^s = \mathrm{id}\}$ , with the previously defined conventions about r, s. In some cases, we shall also consider the spelling of W(a,b) arising from splitting the components  $\vartheta_i^{r_i}$ . More precisely, when both r or s equals  $\infty$ , since the exponents  $r_j$  can be negative, the mentioned splitting of the components  $\vartheta_i^{r_i}$  takes on the form  $\theta_s * \cdots * \theta_1$ , where:

- $\theta_j$  takes on one of the values  $a, b, a^{-1}, b^{-1}$
- if  $\theta_j$  takes on the value a (resp.  $a^{-1}$ ) then, whenever defined, neither  $\theta_{j-1}$  nor  $\theta_{j+1}$  takes on the value  $a^{-1}$  (resp. a). A similar rule applies to  $b, b^{-1}$ .

In the cases where both  $\mathbf{r}, \mathbf{s} < \infty$ ,  $\theta_j$  takes only on the values a, b and every sequence  $\theta_i, \theta_{i+1}, \ldots$  of " $\theta_i$ " with the same value is contained in the split of some  $\vartheta_j^{r_j}$  in the natural sense. Adaptations to the mixed cases  $\mathbf{r} < \infty$ ,  $\mathbf{s} = \infty$  or  $\mathbf{r} = \infty$ ,  $\mathbf{s} < \infty$  are straightforward and left to the reader. In any event, we obtain  $s = \sum_{i=1}^l |r_i|$ . Now, consider the corresponding local diffeomorphism W(f,g) written under the form  $H_s \circ \cdots \circ H_1$  where each  $H_i, i \in \{1,\ldots,s\}$ , belongs to the set  $\{f^{\pm 1}, g^{\pm 1}\}$ . In other words,  $H_i$  replaces  $\theta_i$  by means of the substitutions  $f^{\pm 1} \mapsto a^{\pm 1}, g^{\pm 1} \mapsto b^{\pm 1}$ . The domain of definition of  $W(f,g) = H_s \circ \cdots \circ H_1$  as an element

of  $\Gamma$  can be introduced by recursively defining the domains of definitions of each element  $H_i \circ \cdots \circ H_1$  of  $\Gamma$ ,  $i = 1, \ldots, s$ , as follows:

- The domain of definition of  $H_1$  is all of D and, for every  $z \in D$ ,  $H_1(z)$  is defined in the obvious way.
- Suppose that the domain of definition  $\operatorname{Dom}_{H_i \circ \cdots \circ H_1}$  of  $H_i \circ \cdots \circ H_1$  is already known and that the image  $H_i \circ \cdots \circ H_1(z)$  of  $z \in \operatorname{Dom}_{H_i \circ \cdots \circ H_1}$  under  $H_i \circ \cdots \circ H_1$  is also defined. Then the domain of definition  $\operatorname{Dom}_{H_{i+1} \circ \cdots \circ H_1}$  of  $H_{i+1} \circ \cdots \circ H_1$  is obtained by setting

$$\operatorname{Dom}_{H_{i+1}\circ\cdots\circ H_1} = \{z \in \operatorname{Dom}_{H_i\circ\cdots\circ H_1} \ ; \ H_i \circ \cdots \circ H_1(z) \in D\}.$$

In particular,  $\text{Dom}_{H_{i+1}\circ\cdots\circ H_1}\subseteq \text{Dom}_{H_i\circ\cdots\circ H_1}$  and hence the domain of definition of every element in  $\Gamma$  is naturally contained in D. Besides, for  $z\in \text{Dom}_{H_{i+1}\circ\cdots\circ H_1}$ , the value of  $H_{i+1}\circ\cdots\circ H_1(z)$  is defined by setting  $H_{i+1}\circ\cdots\circ H_1(z)=H_{i+1}\circ[H_i\circ\cdots\circ H_1](z)$ .

Let f, g be local diffeomorphisms as above and consider now a disc D whose closure D is contained in a larger (open) disc where f, g and their inverses are well-defined one-to-one maps. This disc D will be fixed in the sequel and its boundary is going to be denoted by  $\partial D$ . With the previously defined notations, suppose we are given a word  $W(a, b) = \emptyset_l^{r_l} * \cdots * \emptyset_1^{r_1} = \theta_s * \cdots * \theta_1$  which, as always, is supposed to be non-empty and reduced w.r.t. the group  $\{a, b; a^r = b^s = \text{id}\}$ . Given local diffeomorphisms  $h_1, h_2 \in \text{Diff}_{\alpha}(\mathbb{C}, 0)$ , denote by  $\tilde{f}, \tilde{g}$  the local diffeomorphisms given by  $\tilde{f} = h_1^{-1} \circ f \circ h_1$  and  $\tilde{g} = h_2^{-1} \circ g \circ h_2$ . In terms of pseudogroups generated by f, g and by  $\tilde{f}, \tilde{g}$ , note that the domains of definition of the corresponding elements W(f, g) and  $W(\tilde{f}, \tilde{g})$  may drastically differ. To make sense of all these pseudogroups on the fixed disc D, or on the closed disc  $\overline{D}$ , we need the following definition.

- **Definition 2.1.** (1) The domain of definition of  $h_1$  is defined as follows: let  $\rho$  be the radius of the maximal open disc about  $0 \in \mathbb{C}$  in which  $h_1$  is defined and one-to-one. Then the open domain of definition of  $h_1$  is defined to be the open disc of radius  $9\rho/10$ . The closed domain of definition of  $h_1$  will also be considered and this will be nothing but the closed disc of radius  $9\rho/10$ . Analogous definitions apply to each of the local diffeomorphisms:  $h_1^{-1}$ ,  $h_2$ ,  $h_2^{-1}$ .
  - (2) The domain of definition of  $\tilde{f} = h_1^{-1} \circ f \circ h_1$  consists of those points p verifying all the following conditions: p belongs to the open domain of definition of  $h_1$ ,  $h_1(p)$  belongs to the domain of definition of f, i.e. to D. Besides  $f \circ h_1(p)$  must belong to the open domain of definition of  $h_1^{-1}$ . Analogous considerations apply to the domain of definition of  $\tilde{g} = h_2^{-1} \circ g \circ h_2$  and to  $\tilde{f}^{-1}$ ,  $\tilde{g}^{-1}$ .
  - (3) Finally, considering the pseudogroup generated by  $\tilde{f}$ ,  $\tilde{g}$  on the open disc D, the domain of definition of the element  $W(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2) = W(\tilde{f}, \tilde{g})$  is obtained according to the above given general definitions concerning pseudogroups, with  $\tilde{f}$ ,  $\tilde{g}$  in the place of f, g.
  - (4) The (closed) domain of definition of  $\tilde{f} = h_1^{-1} \circ f \circ h_1$ ,  $\tilde{g} = h_1^{-1} \circ g \circ h_1$  on  $\overline{D}$  is defined according to item (2) by using closed domains of definition for f, g as well as for  $h_1, h_1^{-1}, h_2, h_2^{-1}$ . Finally the pseudogroup generated by  $\tilde{f}, \tilde{g}$  on the closed disc  $\overline{D}$  is such that the domain of definition of a general element  $W(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2) =$

 $W(\tilde{f}, \tilde{g})$  is obtained by starting with closed domains of definition for  $\tilde{f}, \tilde{g}$  and following the general pseudogroup rules.

Once  $(h_1, h_2) \in \operatorname{Diff}_{\alpha}(\mathbb{C}, 0) \times \operatorname{Diff}_{\alpha}(\mathbb{C}, 0)$  is chosen, the pseudogroup generated by  $\tilde{f} = h_1^{-1} \circ f \circ h_1$  and by  $\tilde{g} = h_1^{-1} \circ g \circ h_1$  on the open disc D is going to be denoted by  $\Gamma_{h_1,h_2}$ . Consider now an element  $W(\tilde{f}, \tilde{g})$  of  $\Gamma_{h_1,h_2}$  and denote by  $\operatorname{Dom}_W(D)$  its domain of definition. The local diffeomorphism  $W(\tilde{f}, \tilde{g})$  may also be considered as an element of the pseudogroup generated by  $\tilde{f}, \tilde{g}$  on the closed disc  $\overline{D}$  and, in this case, it has a new domain of definition  $\operatorname{Dom}_W(\overline{D})$ . Clearly the topological closure  $\overline{\operatorname{Dom}_W(D)}$  of  $\operatorname{Dom}_W(D)$  is contained in  $\operatorname{Dom}_W(\overline{D})$  whereas, in principle, these two sets may be distinct. However, in what follows, we shall primarily be interested in the pseudogroup  $\Gamma_{h_1,h_2}$  and, when studying its elements, it will be important to consider their local behaviors around points in  $\overline{\operatorname{Dom}_W(D)}$ . Thus possible points lying in  $\operatorname{Dom}_W(\overline{D}) \setminus \overline{\operatorname{Dom}_W(D)}$  will play no role in the subsequent discussion. In view of this and to abridge notations, by a small abuse of language we shall consider that elements in the pseudogroup generated by  $\tilde{f} = h_1^{-1} \circ f \circ h_1$  and by  $\tilde{g} = h_1^{-1} \circ g \circ h_1$  on the closed disc  $\overline{D}$  are defined only on  $\overline{\operatorname{Dom}_W(D)}$ . In other words, we shall write  $\operatorname{Dom}_W(\overline{D})$  to denote the domain of definition in question and, throughout the paper,  $\operatorname{Dom}_W(\overline{D})$  will be assumed to coincide with the closure of  $\operatorname{Dom}_W(D)$ .

The pseudogroup generated by  $\tilde{f}, \tilde{g}$  on the (closed) disc  $\overline{D}$  will be denoted by  $\overline{\Gamma}_{h_1,h_2}$ . Since  $0 \in \mathbb{C}$  is fixed by f, g, we conclude that every word has a non-empty domain of definition as element of both  $\Gamma_{h_1,h_2}$  and  $\overline{\Gamma}_{h_1,h_2}$ . Furthermore, since non-constant holomorphic maps are open maps, the domain of definition of every element  $W(\tilde{f}, \tilde{g})$  in  $\Gamma_{h_1,h_2}$  is necessarily an open set. It may, however, be disconnected.

The above given definition of  $\overline{\Gamma}_{h_1,h_2}$  involves a technical issue playing a significant role in the forthcoming sections which is singled out by the following lemma.

**Lemma 2.2.** Let  $W(\tilde{f}, \tilde{g})$  be an element in  $\overline{\Gamma}_{h_1,h_2}$  whose domain of definition is denoted by  $\mathrm{Dom}_W(\overline{D})$ . Suppose that U is a connected component of  $\mathrm{Dom}_W(\overline{D})$ . Then U is a closed set and, moreover, the holomorphic map  $W(\tilde{f}, \tilde{g}) : U \to \mathbb{C}$  possesses a holomorphic extension to some open neighborhood of U.

*Proof.* The fact that U is closed follows again from the fact that the maps  $\tilde{f}, \tilde{g}$  are holomorphic and hence open (since non-constant). The existence of the desired holomorphic extension follows from the fact that f, g have holomorphic extensions to a neighborhood of  $\overline{D}$  along with the construction of closed domains of definition for  $h_1, h_1^{-1}, h_2, h_2^{-1}$  given in Definition 2.1. The lemma is proved.

Considering the pseudogroups as above, there is already a point to be made about the iterates  $f^j$ ,  $g^j$ , of f, g, which should themselves be understood as elements of suitable pseudogroups defined on some open neighborhood of  $0 \in \mathbb{C}$ . More generally, given  $F \in \text{Diff}(\mathbb{C}, 0)$  and fixed a neighborhood D of  $0 \in \mathbb{C}$  where F is defined, the notation  $F^j$ , where  $j \in \mathbb{Z}^*$ , refers to the element  $F^j$  viewed as an element of the pseudogroup generated by F on D. Now, by combining Poincaré theorem on the existence of linearizing coordinates about hyperbolic fixed points to the well-known topological dynamics of diffeomorphisms tangent to the identity (see for example [C-G]), the following lemma immediately follows.

**Lemma 2.3.** Suppose that  $F \in \text{Diff}(\mathbb{C},0)$  is a local diffeomorphism that does not have a Cremer point at  $0 \in \mathbb{C}$ . Then there is a small disc  $B(\varepsilon)$  about  $0 \in \mathbb{C}$  where, for every  $j \in \mathbb{Z}^*$ ,  $F^j$  has no fixed point unless  $F^j$  coincides with the identity on all of its domain of definition. In particular, if  $F^j$ ,  $j \in \mathbb{Z}$ , coincides with the identity on some connected component of its domain of definition, then it coincides with the identity on all of its domain of definition.  $\square$ 

In closing this section, let us point out that Lemma 2.3 is no longer valid if F is allowed to have a Cremer point at  $0 \in \mathbb{C}$ . Also, as it was the case in [M-R-R], the conclusions of our Theorem A no longer holds if f, g are allowed to have Cremer points. This is a consequence of a construction due to Perez-Marco, cf. [Y] of local diffeomorphisms  $F \in \text{Diff}(\mathbb{C}, 0)$  exhibiting a Cremer point at  $0 \in \mathbb{C}$  and satisfying the following conditions:

- There exists a sequence of points  $\{q_i\}$  accumulating on  $0 \in \mathbb{C}$  along with a sequence of periods  $\{n_i\}$ ,  $n_i \neq 0$ , going to infinity such that  $F^{n_i}(q_i) = q_i$  for every  $i \in \mathbb{N}$ .
- The dynamics of  $F^{n_i}$  about its fixed point  $q_i$  may arbitrarily be fixed: in particular, it can be chosen so that  $F^{n_i}$  coincides with the identity on some (very small) neighborhood of  $q_i$ .

## 3. Proof of Theorem A

Let  $f, g \in \text{Diff}(\mathbb{C}, 0)$  and  $D \subset \mathbb{C}$  be as in Section 2. Fixed  $\alpha \in \mathbb{N}$  and given  $(h_1, h_2) \in \text{Diff}_{\alpha}(\mathbb{C}, 0) \times \text{Diff}_{\alpha}(\mathbb{C}, 0)$ , consider the pseudogroups  $\Gamma_{h_1, h_2}$  and  $\overline{\Gamma}_{h_1, h_2}$  introduced in Section 2. Given a point  $p \in D$ , consider all those elements  $W(\tilde{f}, \tilde{g})$  in  $\Gamma_{h_1, h_2}$  verifying the following:

- The point p belongs to the domain of definition of  $W(\tilde{f}, \tilde{g})$  (as element of  $\Gamma_{h_1,h_2}$ ).
- We have  $W(\tilde{f}, \tilde{g})(p) = p$ .

The germs at p of all elements  $W(\tilde{f}, \tilde{g})$  in  $\Gamma_{h_1,h_2}$  satisfying the mentioned conditions form a group named the stabilizer of p in  $\Gamma_{h_1,h_2}$ . The stabilizer of p is said to be trivial if this group is reduced to the identity.

As mentioned, Theorem A is a natural continuation of the results obtained in [M-R-R]. In fact, to begin the approach to Theorem A, we remind the reader that the main result of [M-R-R] can be summarized as follows.

**Theorem 3.1.** [M-R-R] Suppose that f, g and D as above are fixed. Then there exists a  $G_{\delta}$ -dense set  $\mathcal{V} \subset \operatorname{Diff}_{\alpha}(\mathbb{C}, 0) \times \operatorname{Diff}_{\alpha}(\mathbb{C}, 0)$  such that, whenever  $(h_1, h_2) \in \mathcal{V}$ , for every non-empty reduced word W(a, b) the corresponding element  $W(\tilde{f}, \tilde{g})$  of  $\Gamma_{h_1, h_2}$  does not coincide with the identity on any connected component of its domain of definition.

In the rest of this section we shall prove Theorem A modulo a more technical statement, namely Proposition 3.6, whose proof will be deferred to Sections 4 and 5.

Let us begin with Proposition 3.2 below which plays a significant role in everything that follows. This proposition, as well as the argument used in its proof, is already important for the discussion conducted in this section although its full strength will only be needed in the next section.

As always, let W(a, b) be a fixed non-empty reduced word. Given  $(h_1, h_2) \in \operatorname{Diff}_{\alpha}(\mathbb{C}, 0) \times \operatorname{Diff}_{\alpha}(\mathbb{C}, 0)$ , consider the pseudogroup  $\Gamma_{h_1, h_2}$  (resp.  $\overline{\Gamma}_{h_1, h_2}$ ) generated by  $\tilde{f} = h_1^{-1} \circ f \circ h_1$  and by  $\tilde{g} = h_2^{-1} \circ g \circ h_2$  on D (resp.  $\overline{D}$ ). Finally recall that both f, g are supposed not to have Cremer points at  $0 \in \mathbb{C}$ . The reader is also reminded that, according to our conventions, the

domain of definition  $\mathrm{Dom}_W(\overline{D})$  of an element  $W(\tilde{f}, \tilde{g}) \in \overline{\Gamma}_{h_1,h_2}$  is the closure of its domain of definition  $\mathrm{Dom}_W(D)$  when viewed as element of  $\Gamma_{h_1,h_2}$ .

**Proposition 3.2.** There is an open and dense set  $\mathfrak{U}_W \subset \operatorname{Diff}_{\alpha}(\mathbb{C},0) \times \operatorname{Diff}_{\alpha}(\mathbb{C},0)$  such that, whenever  $(h_1,h_2) \in \mathfrak{U}_W$ , the element  $W(\tilde{f},\tilde{g})$  of  $\overline{\Gamma}_{h_1,h_2}$  has only finitely many fixed points in its (closed) domain of definition  $\operatorname{Dom}_W(\overline{D})$ .

Proof. Naturally the set  $\mathfrak{U}_W$  is defined as consisting of those pairs  $(h_1, h_2) \in \operatorname{Diff}_{\alpha}(\mathbb{C}, 0) \times \operatorname{Diff}_{\alpha}(\mathbb{C}, 0)$  for which the statement of our proposition holds. Let us begin by showing that  $\mathfrak{U}_W$  is dense. For this, it suffices to check that every pair  $(h_1, h_2)$  lying in the  $G_{\delta}$ -dense set  $\mathcal{V}$  provided by Theorem 3.1 gives rise to a pseudogroup  $\overline{\Gamma}_{h_1,h_2}$  whose elements, apart from the identity, have only finitely many fixed points in their (closed) domains of definition.

Consider then  $W(\tilde{f}, \tilde{g})$  where  $h_1$ ,  $h_2$  are as above and denote by  $\mathrm{Dom}_W(\overline{D})$  its (closed) domain of definition. By construction,  $\mathrm{Dom}_W(\overline{D})$  is the closure of the domain of definition  $\mathrm{Dom}_W(D)$  of  $W(\tilde{f}, \tilde{g})$  viewed as element of  $\Gamma_{h_1,h_2}$ . Suppose for a contradiction that  $W(\tilde{f}, \tilde{g})$  possesses infinitely many fixed points in  $\mathrm{Dom}_W(\overline{D})$ . Since  $\mathrm{Dom}_W(\overline{D})$  is closed, there is  $P \in \mathrm{Dom}_W(\overline{D})$  that is the limit of a non-trivial sequence  $\{p_i\} \subset \mathrm{Dom}_W(\overline{D})$  consisting of fixed points of  $W(\tilde{f}, \tilde{g})$ . However, by virtue of Lemma 2.2,  $W(\tilde{f}, \tilde{g})$  admits a holomorphic extension to some open neighborhood  $V \subset \mathbb{C}$  of P. Besides  $W(\tilde{f}, \tilde{g})$  must coincide with the identity on V since P is accumulated by a non-trivial sequence  $\{p_i\}$  of fixed points of  $W(\tilde{f}, \tilde{g})$ . On the other hand, the open set V must intersect the open set  $\mathrm{Dom}_W(D)$  non-trivially since P lies in the closure of  $\mathrm{Dom}_W(D)$ . Therefore  $W(\tilde{f}, \tilde{g})$  coincides with the identity on some connected component of  $\mathrm{Dom}_W(D)$  what is impossible since  $(h_1, h_2) \in \mathcal{V}$ . The resulting contradiction ensures that  $\mathfrak{U}_W$  is dense in  $\mathrm{Diff}_{\alpha}(\mathbb{C}, 0) \times \mathrm{Diff}_{\alpha}(\mathbb{C}, 0)$ .

It remains to show that  $\mathfrak{U}_W$  is also open in the analytic topology. For this consider an element  $(h_1, h_2) \in \mathfrak{U}_W$  and the corresponding element  $W(\tilde{f}, \tilde{g})$  of  $\overline{\Gamma}_{h_1, h_2}$ . We need to check that every pair  $(\overline{h}_1, \overline{h}_2) \in \mathrm{Diff}_{\alpha}(\mathbb{C}, 0) \times \mathrm{Diff}_{\alpha}(\mathbb{C}, 0)$  sufficiently close to  $(h_1, h_2)$  must belong to  $\mathfrak{U}_W$  as well.

Since  $(h_1, h_2)$  lies in  $\mathfrak{U}_W$ ,  $W(\tilde{f}, \tilde{g})$  has only finitely many fixed points in its (closed) domain of definition  $\mathrm{Dom}_W(\overline{D})$ . These points will be denoted by  $p_1, \ldots, p_n$ . Now we consider a small open neighborhood  $U^{\epsilon}$  of  $\mathrm{Dom}_W(\overline{D})$  such that  $W(\tilde{f}, \tilde{g})$  still possesses a holomorphic extension to a neighborhood of the closure  $\overline{U}^{\epsilon}$  of  $U^{\epsilon}$ , cf. Lemma 2.2. Modulo choosing  $U^{\epsilon}$  very small, the points  $p_1, \ldots, p_n$  are isolated points also for the set of fixed points of  $W(\tilde{f}, \tilde{g})$  on the closed set  $\overline{U}^{\epsilon}$ . In fact, we can suppose that the latter set of fixed points coincide with  $p_1, \ldots, p_n$ , i.e.  $W(\tilde{f}, \tilde{g})$  has no additional fixed point in  $\overline{U}^{\epsilon}$ . In particular none of these fixed points lies in the boundary of  $\overline{U}^{\epsilon}$ .

Next, for each  $i=1,\ldots,n$ , let  $B_i(\delta)$  denote the disc of radius  $\delta>0$  about  $p_i$ . If  $\delta$  is chosen sufficiently small, then  $B_i(\delta)\subset U^\epsilon$  and  $B_i(\delta)\cap B_j(\delta)=\emptyset$  provided that  $i\neq j$ . Let now K be the compact set given by  $K=\overline{U}^\epsilon\setminus\bigcup_{i=1}^nB_i(\delta)$ . Since K is compact and  $W(\tilde{f},\tilde{g})$  has no fixed point in K, it follows the existence of  $\tau>0$  such that  $\|W(\tilde{f},\tilde{g})(z)-z\|\geq \tau>0$  for every  $z\in K$ .

Finally consider a sequence  $\overline{h}_{1,j}$  (resp.  $\overline{h}_{2,j}$ ) of elements in  $\mathrm{Diff}_{\alpha}(\mathbb{C},0)$  converging to  $h_1$  (resp.  $h_2$ ) in the analytic topology. Set  $\overline{f}_j = \overline{h}_{1,j}^{-1} \circ f \circ \overline{h}_{1,j}$  and  $\overline{g}_j = \overline{h}_{2,j}^{-1} \circ g \circ \overline{h}_{2,j}$  and

consider the corresponding element  $W(\overline{f}_j, \overline{g}_j)$  of the pseudogroup generated by  $\overline{f}_j$ ,  $\overline{g}_j$  on D. The proposition is now reduced to the following claim:

Claim. For sufficiently large j,  $W(f_i, \overline{g}_i)$  has only finitely many fixed points in its domain of definition.

*Proof of the Claim.* Since convergence in the analytic topology ensures both convergence of domains of definition as well as uniform convergence in these domains, modulo choosing jvery large, the local diffeomorphism  $W(f_i, \overline{g}_i)$  satisfies the conditions below.

- (1) The domain of definition of  $W(\overline{f}_j, \overline{g}_j)$  as element of the pseudogroup generated by  $\overline{f}_j$ ,  $\overline{g}_j$  on D is contained in  $U^{\epsilon} \subset \overline{U}^{\epsilon}$ . Besides  $W(\overline{f}_j, \overline{g}_j)$  possesses a holomorphic extension to a neighborhood of  $\overline{U}^{\epsilon}$ .
- (2) For every  $z \in K \subset \overline{U}^{\epsilon}$ , we have  $\|W(\overline{f}, \overline{g})(z) z\| \ge \tau/2 > 0$ . In particular,  $W(\overline{f}_i, \overline{g}_i)$ has no fixed point in K.

To conclude the proof of the claim, it suffices to check that  $W(\overline{f}_j, \overline{g}_j)$  can have only finitely many fixed points in  $B_i(\delta)$ , for every  $i=1,\ldots,n$ . This is however an immediate consequence of the Argument principle. Indeed, the uniform convergence of  $W(\overline{f}_i, \overline{g}_i)$  to  $W(f, \tilde{g})$  on (a neighborhood of  $B_i(\delta)$ ) implies, by Cauchy formula and the Argument principle, that the sum of the zeros of  $W(f_i, \overline{g}_i)(z) - z$  on  $B_i(\delta)$  counted with multiplicity equals the multiplicity of  $p_i$  as zero of  $W(f, \tilde{g})(z) - z$  since the values of the corresponding integrals must agree by uniform convergence of the integrand. The claim is proved. 

The proof of Proposition 3.2 is over.

For posterior reference, it is useful to explicitly state some by-products of the proof of Proposition 3.2. First we have:

Corollary 3.3. Consider a non-empty reduced word W(a,b). Suppose that  $(h_1,h_2) \in$  $\operatorname{Diff}_{\alpha}(\mathbb{C},0) \times \operatorname{Diff}_{\alpha}(\mathbb{C},0)$  is such that the element  $W(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2) = W(\tilde{f},\tilde{g})$  has only finitely many fixed points in its (open) domain of definition  $\operatorname{Dom}_W(D)$  when viewed as element of  $\Gamma_{h_1,h_2}$ . Then  $W(\tilde{f},\tilde{g})$  also has finitely many fixed points in its (closed) domain of definition  $Dom_W(\overline{D})$  when viewed as element of  $\overline{\Gamma}_{h_1,h_2}$ .

*Proof.* Since  $W(\tilde{f}, \tilde{g})$  has only finitely many fixed points in  $Dom_W(D)$ , it follows that  $W(\tilde{f}, \tilde{g})$ does not coincide with the identity on any connected component of the open set  $Dom_W(D)$ . Now the second paragraph in the proof of Proposition 3.2 can be repeated word-by-word to establish the desired statement.

In the above discussion, it is understood that the fixed points in question can be counted with their multiplicities. On the other hand, if  $(h_1, h_2)$  leads to an element  $W(\tilde{f}, \tilde{g})$  having only finitely many fixed points in its (closed) domain of definition  $Dom_W(D)$ , then the proof of Proposition 3.2 also establishes that every pair  $(h_1, h_2) \in \mathrm{Diff}_{\alpha}(\mathbb{C}, 0) \times \mathrm{Diff}_{\alpha}(\mathbb{C}, 0)$ sufficiently close to  $(h_1, h_2)$  gives rise to a new pseudogroup  $\Gamma_{\overline{h_1}, \overline{h_2}}$  on D whose corresponding element  $W(\overline{f}, \overline{g})$  has only finitely many fixed points in its domain of definition. For reference,

**Lemma 3.4.** Suppose that a reduced word W(a,b) is fixed. Let  $\mathfrak{V}_W \subset \mathrm{Diff}_{\alpha}(\mathbb{C},0) \times$  $\operatorname{Diff}_{\alpha}(\mathbb{C},0)$  denote the set of pairs  $(h_1,h_2)$  giving rise to a pseudogroup  $\Gamma_{h_1,h_2}$  for which the element  $W(\tilde{f}, \tilde{g})$  possesses only finitely many fixed points in its closed domain of definition. Then  $\mathfrak{V}_W$  is open.

Furthermore, if  $(h_1, h_2) \in \mathfrak{V}$  is "perturbed" to  $(\overline{h}_1, \overline{h}_2)$  (necessarily in  $\mathfrak{V}$ ), then the fixed points of  $W(\overline{h}_1, \overline{h}_2)$  are close to the fixed points of  $W(\tilde{f}, \tilde{g})$ .

Note that, when it comes to the assumption of Lemma 3.4, Corollary 3.3 ensures that the mentioned assumption is, indeed, equivalent to saying that  $W(\tilde{f}, \tilde{g})$  has finitely many fixed points in its open domain of definition  $\mathrm{Dom}_W(D)$ .

Two reduced words  $W_1(a, b)$ ,  $W_2(a, b)$  are said to be commensurable if there is a word  $W_{1-2}(a, b)$  such that both  $W_1(a, b)$ ,  $W_2(a, b)$  are obtained by concatenating finitely many "copies" of  $W_{1-2}(a, b)$ . This means that the local diffeomorphism  $W_1(f, g)$  (resp.  $W_2(f, g)$ ) is a finite power of the local diffeomorphism  $W_{1-2}(f, g)$  for every pair  $f, g \in \text{Diff}(\mathbb{C}, 0)$ . If there is not such word  $W_{1-2}(a, b)$ , then  $W_1(a, b)$ ,  $W_2(a, b)$  are said to be incommensurable. Let  $\mathcal{I}$  denote the collection of pairs  $(W_1(a, b), W_2(a, b))$  of incommensurable words. Clearly  $\mathcal{I}$  is a countable set.

Given a reduced word W(a, b) and a pair  $(h_1, h_2) \in \mathfrak{U}_W \subset \operatorname{Diff}_{\alpha}(\mathbb{C}, 0) \times \operatorname{Diff}_{\alpha}(\mathbb{C}, 0)$ , denote by  $\operatorname{Fix}(W(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2))$  the set of fixed points of  $W(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2) = W(\tilde{f}, \tilde{g})$  in its (closed) domain of definition  $\operatorname{Dom}_W(\overline{D})$  (i.e.  $W(\tilde{f}, \tilde{g})$  is viewed as element of  $\overline{\Gamma}_{h_1,h_2}$ ). Let us now choose two incommensurable words  $W_i(a,b)$  and  $W_j(a,b)$  so that the pair  $(W_i(a,b),W_j(a,b))$  defines an element of  $\mathcal{I}$ . Consider the set  $\mathcal{U}_{i,j} \subset \operatorname{Diff}_{\alpha}(\mathbb{C},0) \times \operatorname{Diff}_{\alpha}(\mathbb{C},0)$  consisting of those pairs  $(h_1,h_2)$  of local diffeomorphisms fulfilling the following two conditions:

- Both sets Fix  $(W_i(\tilde{f}, \tilde{g}))$  and Fix  $(W_i(\tilde{f}, \tilde{g}))$  are finite.
- Fix  $(W_i(\tilde{f}, \tilde{g})) \cap$  Fix  $(W_j(\tilde{f}, \tilde{g})) = \{0\} \subset \mathbb{C}$ .
- The multiplicity of  $0 \in \mathbb{C}$  as fixed point of  $W_i(\tilde{f}, \tilde{g})$  and of  $W_j(\tilde{f}, \tilde{g})$  does not change by perturbation.

Note that the first item above means that  $(h_1, h_2)$  lies in the intersection  $\mathfrak{V}_i \cap \mathfrak{V}_j$  which is open owing to Lemma 3.4. Next, in view of the third item, no additional common fixed point can "bifurcate" from  $0 \in \mathbb{C}$  by perturbing  $(h_1, h_2)$  due to the usual Argument principle. Note also that the condition in the third item is always verified when  $0 \in \mathbb{C}$  has multiplicity 1 as fixed point of  $W_i(\tilde{f}, \tilde{g}), W_j(\tilde{f}, \tilde{g})$ . Since, apart from the  $0 \in \mathbb{C}$ ,  $W_i(\tilde{f}, \tilde{g}), W_j(\tilde{f}, \tilde{g})$  share no fixed point, it follows from the second part of the statement of Lemma 3.4 that Fix  $(W_i(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)) \cap \text{Fix}(W_j(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)) = \{0\} \subset \mathbb{C}$  for every pair  $(\overline{h}_1, \overline{h}_2)$  sufficiently close to  $(h_1, h_2)$ .

Summarizing what precedes, we have proved the following.

**Lemma 3.5.** The set  $\mathcal{U}_{i,j} \subset \operatorname{Diff}_{\alpha}(\mathbb{C},0) \times \operatorname{Diff}_{\alpha}(\mathbb{C},0)$  is open for every pair of incommensurable words  $W_i(a,b), W_j(a,b)$ .

On the other hand, we shall also prove:

**Proposition 3.6.** The set  $\mathcal{U}_{i,j} \subset \mathrm{Diff}_{\alpha}(\mathbb{C},0) \times \mathrm{Diff}_{\alpha}(\mathbb{C},0)$  is dense for every pair of incommensurable words  $W_i(a,b)$ ,  $W_i(a,b)$ .

As soon as Proposition 3.6 is established, we are able to prove Theorem A. Since the proof of Proposition 3.6 is long and technical, we shall first derive Theorem A deferring to the next section the proof of Proposition 3.6.

Proof of Theorem A. Fixed a pair  $(W_i(a,b),W_j(a,b))$  in  $\mathcal{I}$ , i.e. a pair of incommensurable words, consider the above defined set  $\mathcal{U}_{i,j}$ . According to Lemma 3.5 and to Proposition 3.6, the set  $\mathcal{U}_{i,j}$  is open and dense in  $\mathrm{Diff}_{\alpha}(\mathbb{C},0) \times \mathrm{Diff}_{\alpha}(\mathbb{C},0)$ . Let us then define

$$\mathcal{U} = igcap_{(W_i(a,b),W_j(a,b)) \in \mathcal{I}} \mathcal{U}_{i,j} \,.$$

Clearly  $\mathcal{U}$  is a  $G_{\delta}$ -dense subset of  $\mathrm{Diff}_{\alpha}(\mathbb{C},0) \times \mathrm{Diff}_{\alpha}(\mathbb{C},0)$ . By construction of  $\mathcal{U}$ , it also clear that the stabilizer of every point  $z \neq 0 \in D$  is either cyclic or trivial. Therefore, the proof of Theorem A is reduced to show the existence of a sequence of points  $\{Q_n\}$ ,  $Q_n \neq 0$  for every  $n \in \mathbb{N}$ , with the properties indicated in the statement of Theorem A.

For this, note that  $\mathcal{U}$  is contained in the set  $\mathcal{V}$  provided by Theorem 3.1. In fact, for every reduced word W(a,b), the element  $W(\tilde{f},\tilde{g})$  has only isolated fixed points in  $\mathrm{Dom}_W(D)$  so long  $(h_1,h_2)\in\mathcal{U}$ . Therefore  $W(\tilde{f},\tilde{g})$  cannot coincide with the identity on any connected component of its domain of definition  $\mathrm{Dom}_W(D)$ . It follows, in particular, that the germ of  $\Gamma_{h_1,h_2}$  at  $0\in\mathbb{C}$  is not solvable, there are points in D that are hyperbolic fixed points for certain elements of  $\Gamma_{h_1,h_2}$ , cf. [BLL-1]. Furthermore, the multipliers of these fixed points can a priori be fixed in a dense set of  $\mathbb{C}$ . However, inasmuch there are infinitely many points whose stabilizers contain a hyperbolic element, it may happen that all these points are contained in a single orbit of the pseudogroup  $\Gamma_{h_1,h_2}$ .

To complete the proof of the theorem, we proceed as follows. Let  $Q_1$  be a hyperbolic fixed point of some element  $W_1(\tilde{f}, \tilde{g}) \in \Gamma_{h_1,h_2}$ . In particular the stabilizer of  $Q_1$  is not trivial and hence it must be cyclic. Thus we can assume that  $W_1(\tilde{f}, \tilde{g})$  is the generator of the stabilizer of  $Q_1$ . As mentioned above, the orbit of  $Q_1$  by  $\Gamma_{h_1,h_2}$  is constituted by points whose stabilizers contain some hyperbolic element, namely a certain conjugate of  $W_1(\tilde{f}, \tilde{g})$ . Consider for each point  $\gamma.Q_1$  in the  $Q_1$ -orbit the multipliers of elements in the stabilizer of  $\gamma.Q_1$ . The collection of all multipliers obtained from points in the  $Q_1$ -orbit is then denoted by  $M_1$ . Now note that  $M_1$  is a discrete subset of  $\mathbb{C}$ , indeed,  $M_1$  is nothing but a cyclic subgroup of  $\mathbb{C}^*$  generated by the derivative of  $W_1(\tilde{f}, \tilde{g})$  at  $Q_1$ . Thus, after [BLL-2], there must exist another hyperbolic fixed point  $Q_2$  whose multiplier lies away from a neighborhood of  $M_1$  in  $\mathbb{C}$ . In particular, the orbits of  $Q_1$  and  $Q_2$  must be disjoint. However, the preceding argument applies again to ensure that the orbit of  $Q_2$  yields another discrete set of multipliers  $M_2 \subset \mathbb{C}$ . The construction can then be continued to yield infinitely many hyperbolic fixed points with pairwise disjoint orbits. Theorem A is proved.

Corollary B is an immediate consequence of what precedes.

Proof of Corollary B. The construction detailed in Section 5 of [M-R-R] allows us to translate information on the topology of the leaves of the corresponding foliations into dynamical properties of the pseudogroup  $\Gamma_{h_1,h_2}$ , and conversely. By means of this connection, the items (4) and (5) of Corollary B turn out to be implied by Theorem A. The remaining items were already established in [M-R-R]. An alternative possibility is to resort to the general statements of [M-M]. In any event the proof of Corollary B is completed.

## 4. Proof of Proposition 3.6

The rest of the paper is entirely devoted to the proof of Proposition 3.6. This proof will be accomplished in this section whereas the proof of a more technical lemma on the existence of suitable perturbations will be supplied only in the last section of this article.

Let us start by explaining the strategy for proving Proposition 3.6. Fix two reduced incommensurable words  $W_i(a,b)$  and  $W_j(a,b)$  and suppose we are given a pair  $(h_1,h_2) \in \operatorname{Diff}_{\alpha}(\mathbb{C},0) \times \operatorname{Diff}_{\alpha}(\mathbb{C},0)$ . We need to find  $(\overline{h}_1,\overline{h}_2) \in \mathcal{U}_{i,j}$  arbitrarily close to  $(h_1,h_2)$ . The existence of the desired pair  $(\overline{h}_1,\overline{h}_2)$  will be shown by successively approximating  $(h_1,h_2)$  by elements in  $\operatorname{Diff}_{\alpha}(\mathbb{C},0) \times \operatorname{Diff}_{\alpha}(\mathbb{C},0)$  that will fulfil "more and more" the conditions needed to belong to  $\mathcal{U}_{i,j}$ . This will be done so that, after finitely many steps, a pair  $(\overline{h}_1,\overline{h}_2) \in \mathcal{U}_{i,j}$  will be found in a given  $\varepsilon$ -neighborhood of  $(h_1,h_2)$ .

This goes as follows. First, by using the "denseness part" of the statement of Proposition 3.2 applied to both words  $W_i(a,b)$ ,  $W_j(a,b)$ , we see that arbitrarily close to  $(h_1,h_2)$  there is  $(\overline{h}_1,\overline{h}_2)$  leading to a pseudogroup  $\overline{\Gamma}_{\overline{h}_1,\overline{h}_2}$  whose elements  $W_i(\overline{h}_1^{-1}\circ f\circ \overline{h}_1,\overline{h}_2^{-1}\circ g\circ \overline{h}_2)$  and  $W_j(\overline{h}_1^{-1}\circ f\circ \overline{h}_1,\overline{h}_2^{-1}\circ g\circ \overline{h}_2)$  possess only finitely many fixed points in their closed domains of definitions  $\mathrm{Dom}_{W_i}(\overline{D})$ ,  $\mathrm{Dom}_{W_j}(\overline{D})$ . In other words, to abridge notations, we can assume without loss of generality that the initial local diffeomorphisms are already such that the words  $W_i(h_1^{-1}\circ f\circ h_1,\,h_2^{-1}\circ g\circ h_2)=W_i(\tilde{f},\tilde{g})$  and  $W_j(h_1^{-1}\circ f\circ h_1,\,h_2^{-1}\circ g\circ h_2)=W_j(\tilde{f},\tilde{g})$  have only finitely many fixed points as elements of  $\overline{\Gamma}_{h_1,h_2}$ .

Denote then by  $\operatorname{Fix}(W_i(\tilde{f},\tilde{g}))$  (resp.  $\operatorname{Fix}(W_j(\tilde{f},\tilde{g}))$ ) the set of fixed points of  $W_i(\tilde{f},\tilde{g})$  (resp.  $W_j(\tilde{f},\tilde{g})$ ) in its (closed) domain of definition as element of  $\overline{\Gamma}_{h_1,h_2}$ . These sets are both finite. The proof of Proposition 3.6 is essentially reduced to checking that  $(h_1,h_2)$  can be approximated by a pair  $(\overline{h}_1,\overline{h}_2)$  yielding corresponding local diffeomorphisms  $W_i(\overline{h}_1^{-1}\circ f\circ \overline{h}_1,\overline{h}_2^{-1}\circ g\circ \overline{h}_2)=W_j(\overline{f},\overline{g})$  having no common fixed point other than  $0\in\mathbb{C}$ . The condition that the multiplicity of  $0\in\mathbb{C}$  as fixed point of  $W_i(\overline{f},\overline{g}),W_j(\overline{f},\overline{g})$  should not change under perturbations is a minor one and it will be dealt with below. Consider then the problem of showing that  $W_i(\tilde{f},\tilde{g}),W_j(\tilde{f},\tilde{g})$  can be perturbed so as not to have common fixed point other than  $0\in\mathbb{C}$ . Note that the construction of the desired perturbations is a problem that is naturally localized at the mentioned common fixed points. To explain this assertion and clarify the rest of our strategy to approach Proposition 3.6, suppose for example that the initial pair  $(h_1,h_2)$  is such that  $p\neq 0$  is the only common fixed point for  $W_i(\tilde{f},\tilde{g}),W_j(\tilde{f},\tilde{g})$  away from  $0\in\mathbb{C}$ . Consider then a small disc  $B(\delta)$  about p. Also the (closed) domain of definition of  $W_i(\tilde{f},\tilde{g})$  (resp.  $W_j(\tilde{f},\tilde{g})$ ) is going to be denoted by  $\operatorname{Dom}_{W_i}(\overline{D})$  (resp.  $\operatorname{Dom}_{W_j}(\overline{D})$ ). According to Lemma 2.2 (cf. Corollary 3.3 and Proposition 3.2), we can choose a (closed) neighborhood  $\overline{U}_i^e$  of  $\operatorname{Dom}_{W_i}(\overline{D})$  (resp.  $\overline{U}_j^e$  of  $\operatorname{Dom}_{W_i}(\overline{D})$ ) where the following holds:

- $W_i(\tilde{f}, \tilde{g})$  (resp.  $W_j(\tilde{f}, \tilde{g})$ ) has a holomorphic extension to some neighborhood of  $\overline{U}_i^{\epsilon}$  (resp.  $\overline{U}_i^{\epsilon}$ ).
- The fixed points of  $W_i(\tilde{f}, \tilde{g})$  (resp.  $W_j(\tilde{f}, \tilde{g})$ ) in  $\overline{U}_i^{\epsilon}$  (resp.  $\overline{U}_j^{\epsilon}$ ) are all contained in  $\mathrm{Dom}_{W_i}(\overline{D})$  (resp.  $\mathrm{Dom}_{W_i}(\overline{D})$ ).

Now consider the compact set  $\overline{U}_{j}^{\epsilon} \cap [\overline{U}_{i}^{\epsilon} \setminus B(\delta)]$  and note that for every point  $z \in \overline{U}_{j}^{\epsilon} \cap [\overline{U}_{i}^{\epsilon} \setminus B(\delta)]$ , we have  $||W_i(\tilde{f},\tilde{g})(z)-W_i(\tilde{f},\tilde{g})(z)||\neq 0$ . Since this set is compact, it follows the existence of some  $\tau > 0$  such that we actually have  $||W_i(\tilde{f}, \tilde{g})(z) - W_j(\tilde{f}, \tilde{g})(z)|| \geq \tau > 0$  for every  $z \in \overline{U}_i^{\epsilon} \cap [\overline{U}_i^{\epsilon} \setminus B(\delta)]$ . Next the reader is reminded that convergence in the analytic topology implies convergence of domains of definition as well as uniform convergence of maps on  $\underline{\text{the}}$ corresponding domains. Thus, recalling that  $\mathrm{Dom}_{W_i}(\overline{D})$  is contained in the interior of  $\overline{U}_i^{\mathfrak{c}}$ , by taking  $(\overline{h}_1, \overline{h}_2)$  very close to  $(h_1, h_2)$  the following holds:

- (i) The closed domain of definition of  $W_i(\overline{f}, \overline{g})$  (resp.  $W_i(\overline{f}, \overline{g})$ ) is contained in  $\overline{U}_i^{\epsilon}$  (resp.  $\overline{U}_{i}^{\epsilon}$ ).
- (11) The local diffeomorphism  $W_i(\overline{f}, \overline{g})$  (resp.  $W_i(\overline{f}, \overline{g})$  possesses a holomorphic extension to a neighborhood of  $\overline{U}_i^{\epsilon}$  (resp.  $\overline{U}_i^{\epsilon}$ ).

It follows from (i) and (ii) that  $W_i(\overline{f}, \overline{g})(z) - W_j(\overline{f}, \overline{g})(z)$  is defined for every  $z \in \overline{U}_j^{\epsilon} \cap [\overline{U}_i^{\epsilon} \setminus \overline{U}_j^{\epsilon}]$  $B(\delta)$ ]. Next, by uniform convergence, it also follows that  $||W_i(\overline{f}, \overline{g})(z) - W_j(\overline{f}, \overline{g})(z)|| \ge$  $\tau/2 > 0$  for every  $z \in \overline{U}_j^{\epsilon} \cap [\overline{U}_i^{\epsilon} \setminus B(\delta)]$ . Therefore we conclude that the all possible common fixed points of  $W_i(\overline{f}, \overline{g})$  and  $W_i(\overline{f}, \overline{g})$  lie in  $B(\delta)$ . Thus, the proof Proposition 3.6 is essentially reduced to an analysis of  $W_i(\tilde{f}, \tilde{g})$  and of  $W_i(\tilde{f}, \tilde{g})$  on a neighborhood of their common fixed points. Namely we need to show that these common fixed points can be split by arbitrarily small perturbations of the initial local diffeomorphisms  $(h_1, h_2)$ . In the sequel, we shall provide full detail for this construction.

Summarizing what precedes, we can assume that both  $W_i(\overline{f}, \overline{g})$  and  $W_i(\overline{f}, \overline{g})$  have only finitely many fixed points in their closed domains of definition denoted respectively by  $\mathrm{Dom}_{W_i}(\overline{D})$  and  $\mathrm{Dom}_{W_i}(\overline{D})$ . Moreover, we can choose a neighborhood  $U_i^{\epsilon}$  of  $\mathrm{Dom}_{W_i}(\overline{D})$ (resp.  $U_j^{\epsilon}$  of  $\mathrm{Dom}_{W_j}(\overline{D})$ ) such that  $W_i(\tilde{f}, \tilde{g})$  (resp.  $W_j(\tilde{f}, \tilde{g})$ ) has a holomorphic extension to a neighborhood of  $\overline{U}_i^{\epsilon}$  (resp.  $\overline{U}_j^{\epsilon}$ ). Furthermore the fixed points of  $W_i(\tilde{f}, \tilde{g})$  (resp.  $W_j(\tilde{f}, \tilde{g})$ ) in  $\overline{U}_i^{\epsilon}$  (resp.  $\overline{U}_j^{\epsilon}$ ) are all contained in  $\mathrm{Dom}_{W_i}(\overline{D})$  (resp.  $\mathrm{Dom}_{W_j}(\overline{D})$ ).

Recalling that our purpose is to find  $(\overline{h}_1, \overline{h}_2) \in \mathcal{U}_{i,j}$  arbitrarily close to  $(h_1, h_2)$ , we begin with the following lemma.

**Lemma 4.1.** To construct  $(\overline{h}_1, \overline{h}_2)$ , we can assume that the multiplicity associated to each fixed point of both  $W_i(\tilde{f}, \tilde{g})$  and  $W_j(\tilde{f}, \tilde{g})$  does not change under perturbations.

*Proof.* Consider the case of  $W_i(\tilde{f}, \tilde{g})$ . Denote by  $P_1, \ldots, P_s$  its fixed points. Suppose first that  $W_i(\tilde{f}, \tilde{g})$  has no fixed point in the boundary of its domain of definition. Let  $B_k(\delta)$  be a small ball about  $P_k$ , k = 1, ..., s containing no other fixed point of  $W_i(\tilde{f}, \tilde{g})$ . Also denote by  $N_k$  the multiplicity of  $P_k$  as fixed point of  $W_i(\tilde{f}, \tilde{g})$ . By continuity, if  $(\overline{h}_1, \overline{h}_2)$  is sufficiently close to  $(h_1, h_2)$ , then we have:

- (1) The (closed) domain of definition of  $W_i(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$  is contained in  $\overline{U}_i^{\epsilon}$ . (2) The diffeomorphism  $W_i(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$  has no fixed point in  $\overline{U}_i^{\epsilon} \setminus \bigcup_{k=1}^s B_k(\delta)$ .

Also, fixed k, the number of fixed points of  $W_i(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$  in  $B_k(\delta)$  counted with their multiplicities is precisely  $N_k$ , as it follows again from the Argument principle. In particular, the multiplicity of a fixed point may decrease, but never increase, under perturbations. Thus, if there are arbitrarily small perturbations for which each point  $P_k$  splits into  $N_k$  fixed points with multiplicity 1, then the statement becomes immediate: we consider a first perturbation  $(\overline{h}_1, \overline{h}_2)$  such that all fixed points become of multiplicity 1 and then, we only need to approximate  $(\overline{h}_1, \overline{h}_2)$  by elements  $(\overline{h}_1, \overline{h}_2)$  as above. Since, in this case, every fixed point of  $W_i(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$  has multiplicity 1, this multiplicity will not change by perturbations as already seen. The general case follows from this argument, we consider the least multiplicity that can be achieved for each fixed point of  $W_i(\tilde{f}, \tilde{g})$  by perturbing  $(h_1, h_2)$ . Being given by local minima, these multiplicities cannot further decrease under perturbations. On the other hand, the previous general argument shows that they cannot increase either. Therefore they must remain constant what proves the lemma in this case.

Finally the argument when  $W_i(\tilde{f}, \tilde{g})$  possesses fixed points in the boundary of its domain of definition is essentially the same. These points are in finite number. If by an arbitrarily small perturbation some of them fall in the open domain and others fall away from the closed domain, then the situation is reduced to the preceding case. Otherwise there are points that remain in the boundary of the domain of definition of  $W_i(\tilde{f}, \tilde{g})$  for every sufficient small perturbation of  $(h_1, h_2)$ . The same argument above can then be applied to these fixed points.

Owing to Lemma 4.1, we can assume without loss of generality that, in addition, the multiplicities of the fixed points of  $W_i(\tilde{f}, \tilde{g})$  and of  $W_j(\tilde{f}, \tilde{g})$  do not change under perturbation of  $(h_1, h_2)$ . Recalling that  $W_i(\tilde{f}, \tilde{g})$  and  $W_j(\tilde{f}, \tilde{g})$  have only isolated fixed points, the next lemma establishes that a common fixed point for  $W_i(\tilde{f}, \tilde{g})$  and  $W_j(\tilde{f}, \tilde{g})$  can always be destroyed by arbitrarily small perturbations provided that  $W_i(a, b)$ ,  $W_j(a, b)$  are two incommensurable words. More precisely:

**Lemma 4.2.** Consider two incommensurable words  $W_i(a, b)$ ,  $W_j(a, b)$  as above along with a given pair  $(h_1, h_2) \in \operatorname{Diff}_{\alpha}(\mathbb{C}, 0) \times \operatorname{Diff}_{\alpha}(\mathbb{C}, 0)$ . Suppose that  $q \neq 0$  is a common fixed point for  $W_i(\tilde{f}, \tilde{g})$  and  $W_j(\tilde{f}, \tilde{g})$  (viewed as elements of  $\Gamma_{h_1,h_2}$ ). Let  $B(\delta)$  be a small disc about q containing no other fixed point of  $W_i(\tilde{f}, \tilde{g})$  and  $W_j(\tilde{f}, \tilde{g})$ . Then, arbitrarily close to  $(h_1, h_2)$ , there is  $(\overline{h}_1, \overline{h}_2)$  such that  $W_i(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$  and  $W_j(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$  have no common fixed point in  $B(\delta)$ .

After the preceding discussion, Lemma 4.2 is the main technical result needed for the proof of Proposition 3.6. We shall close this section with the proof of Proposition 3.6. The next section will be devoted to the constructions leading to the proof of Lemma 4.2

Proof of Proposition 3.6. The argument is now clear. Again denote by  $P_1, \ldots, P_s$  the fixed points of, say,  $W_i(\tilde{f}, \tilde{g})$ . As explained in the proof of Lemma 4.1, modulo constructing a first perturbation of  $(h_1, h_2)$ , we can assume that  $W_i(\tilde{f}, \tilde{g})$  has no fixed points in the boundary of its domain of definition, unless the fixed points lying in this boundary remain in it for every sufficiently small perturbation of  $(h_1, h_2)$ . In the rest of the discussion, both possibilities will be treated together since there is no essential difference between them.

Consider the above fixed neighborhood  $U_i^{\epsilon}$  of the closed domain of definition of  $W_i(\tilde{f}, \tilde{g})$ . Let  $\delta > 0$  very small be fixed. For every  $k = 1, \ldots, s$ , denote by  $B_k(\delta)$  a small disc about the fixed point  $P_k$ . The choice of  $\delta$  is dictated by the fact that the distance between any pair of these discs must be strictly positive and by the fact that they should all be contained in a compact part of  $U_i^{\epsilon}$ . By construction,  $W_i(\tilde{f}, \tilde{g})$  has no additional fixed point in  $U_i^{\epsilon}$ . Thus, if  $(\overline{h}_1, \overline{h}_2)$  is close enough to  $(h_1, h_2)$ , it follows that  $W_i(\overline{f}, \overline{g}) = W_i(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$  satisfies the following conditions:

- The closed domain of definition of  $W_i(\overline{f}, \overline{g})$  is contained in  $U_i^{\epsilon}$ .
- $W_i(\overline{f}, \overline{g})$  has no fixed point in  $\overline{U}_i^{\epsilon} \setminus \bigcup_{k=1}^s B_k(\delta)$ .

Analogous conclusions hold for  $W_j(\overline{f}, \overline{g}) = W_j(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$ . In the sequel all perturbations will be chosen small enough to guarantee that the final "perturbed" diffeomorphisms still satisfy the above conditions for  $W_i(\overline{f}, \overline{g})$  and for  $W_j(\overline{f}, \overline{g})$ .

In view of what precedes, and given that the multiplicities of  $P_1, \ldots, P_s$  do not change under perturbations, the number s of fixed points of  $W_i(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)$  will not change under sufficiently small perturbations of  $(h_1, h_2)$  (owing again to the Argument principle).

Consider again the above defined discs  $B_k(\delta)$ ,  $k=1,\ldots,s$ . Recall that every sufficiently small perturbation  $(\tilde{h}_1,\tilde{h}_2)$  of  $(h_1,h_2)$  leads to a new local diffeomorphism  $W_i(\tilde{h}_1^{-1}\circ f\circ \tilde{h}_1,\tilde{h}_2^{-1}\circ g\circ \tilde{h}_2)$  having exactly one fixed point in each small disc  $B_k(\delta)$  and no fixed point in the complement of  $\bigcup_{k=1}^s B_k(\delta)$  in the closed domain of definition of  $W_i(\tilde{h}_1^{-1}\circ f\circ \tilde{h}_1,\tilde{h}_2^{-1}\circ g\circ \tilde{h}_2)$  itself. Starting from k=1, suppose that  $P_1$  is a common fixed point for  $W_i(\overline{f},\overline{g})$  and  $W_j(\overline{f},\overline{g})$ . Then Lemma 4.2 allows us to find an arbitrarily small perturbation  $(\overline{h}_{1,1},\overline{h}_{2,1})$  of  $(h_1,h_2)$  such that  $W_i(\overline{h}_{1,1}^{-1}\circ f\circ \overline{h}_{1,1},\overline{h}_{2,1}^{-1}\circ g\circ \overline{h}_{2,1})$  and  $W_j(\overline{h}_{1,1}^{-1}\circ f\circ \overline{h}_{1,1},\overline{h}_{2,1}^{-1}\circ g\circ \overline{h}_{2,1})$  have no longer a common fixed point in  $B_1(\delta)$ . Furthermore, if  $(\overline{h}_{1,1},\overline{h}_{2,1})$  is close enough to  $(h_1,h_2)$ , then  $W_i(\overline{h}_{1,1}^{-1}\circ f\circ \overline{h}_{1,1},\overline{h}_{2,1}^{-1}\circ g\circ \overline{h}_{2,1})$  still satisfies the previous conditions regarding its own fixed points, which will now be denoted by  $P_{1,1},\ldots,P_{s,1}$ . In particular, each  $P_{k,1}$  lies in  $B_k(\delta)$ . By construction  $P_{1,1}$  is not a common fixed point for  $W_i(\overline{h}_{1,1}^{-1}\circ f\circ \overline{h}_{1,1},\overline{h}_{2,1}^{-1}\circ g\circ \overline{h}_{2,1})$  and  $W_j(\overline{h}_{1,1}^{-1}\circ f\circ \overline{h}_{1,1},\overline{h}_{2,1}^{-1}\circ g\circ \overline{h}_{2,1})$ . Thus, if no point  $P_{k,1}$ ,  $k=2,\ldots,s$  turns out to be fixed by  $W_j(\overline{h}_{1,1}^{-1}\circ f\circ \overline{h}_{1,1},\overline{h}_{2,1}^{-1}\circ g\circ \overline{h}_{2,1})$  then the statement is proved.

Thus let us suppose that  $P_{2,1}$  is fixed also by  $W_j(\overline{h}_{1,1}^{-1} \circ f \circ \overline{h}_{1,1}, \overline{h}_{2,1}^{-1} \circ g \circ \overline{h}_{2,1})$ . By using again Lemma 4.2 we can find a new perturbation  $(\overline{h}_{1,2}, \overline{h}_{2,2})$  of  $(\overline{h}_{1,1}, \overline{h}_{2,1})$  so that  $W_i(\overline{h}_{1,2}^{-1} \circ f \circ \overline{h}_{1,2}, \overline{h}_{2,2}^{-1} \circ g \circ \overline{h}_{2,2})$  and  $W_j(\overline{h}_{1,2}^{-1} \circ f \circ \overline{h}_{1,2}, \overline{h}_{2,2}^{-1} \circ g \circ \overline{h}_{2,2})$  have no longer a common fixed point in  $B_2(\delta)$ . Furthermore, modulo choosing this perturbation sufficiently small, the following conditions can again be ensured:

- No common fixed point for  $W_i(\overline{h}_{1,2}^{-1} \circ f \circ \overline{h}_{1,2}, \overline{h}_{2,2}^{-1} \circ g \circ \overline{h}_{2,2})$  and  $W_j(\overline{h}_{1,2}^{-1} \circ f \circ \overline{h}_{1,2}, \overline{h}_{2,2}^{-1} \circ g \circ \overline{h}_{2,2})$  is produced in  $B_1(\delta)$ .
- $g \circ \overline{h}_{2,2}$ ) is produced in  $B_1(\delta)$ . •  $W_i(\overline{h}_{1,1}^{-1} \circ f \circ \overline{h}_{1,1}, \overline{h}_{2,1}^{-1} \circ g \circ \overline{h}_{2,1})$  still has exactly s fixed points, denoted by  $P_{1,2}, \ldots, P_{s,2}$ . Besides, for every  $k = 1, \ldots, s$ , the fixed point  $P_{k,2}$  belongs to the disc  $B_k(\delta)$ .

In particular, after this second perturbation,  $W_i(\overline{h}_{1,2}^{-1} \circ f \circ \overline{h}_{1,2}, \overline{h}_{2,2}^{-1} \circ g \circ \overline{h}_{2,2})$  and  $W_j(\overline{h}_{1,2}^{-1} \circ f \circ \overline{h}_{1,2}, \overline{h}_{2,2}^{-1} \circ g \circ \overline{h}_{2,2})$  can have at most s-2 common fixed points. By inductively continuing this argument, we shall eventually obtain a perturbation  $(\overline{h}_1, \overline{h}_2) = (\overline{h}_{1,s}, \overline{h}_{2,s})$  of  $(h_1, h_2)$  satisfying the condition required in the statement. The proof of Proposition 3.6 is over.

## 5. Constructing analytic perturbations: proof of Lemma 4.2

In this last section we shall introduce some perturbation techniques leading to the proof of Lemma 4.2. First, we assume that the conditions used in the previous section still hold in the present context. This means that  $W_i(f, \tilde{g})$  and  $W_i(f, \tilde{g})$  have only isolated fixed points and, furthermore, that the multiplicity of these fixed points do not change under perturbations of  $(h_1, h_2)$ .

The approach to the proof of Lemma 4.2 begins with some simple reductions in the statement. First it will be proved that, if  $q \in \mathbb{C}$ ,  $q \neq 0$ , is a fixed point for  $W_i(\tilde{f}, \tilde{g})$ , then we can perturb  $(h_1, h_2)$  into  $(\overline{h}_1, \overline{h}_2)$  so that q is no longer fixed by  $W_j(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$ . Assuming that q is a common fixed point for both  $W_i(\tilde{f}, \tilde{g})$  and  $W_j(\tilde{f}, \tilde{g})$ , the proof of Lemma 4.2 amounts to checking that such perturbation can be applied to, say,  $W_i(\tilde{f}, \tilde{g})$  at q while keeping the point q fixed by  $W_i(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$ . As already explained, since the multiplicity of q as fixed point of  $W_i(\tilde{f}, \tilde{g})$  does not change under perturbations of  $(h_1, h_2)$ , it follows that for every pair  $(\overline{h}_1, \overline{h}_2)$  sufficiently close to  $(h_1, h_2)$ ,  $W_i(\overline{h}_1^{-1} \circ f \circ f)$  $\overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$  will still have a unique fixed point on a fixed neighborhood of q. Thus, if q remains fixed by the perturbed diffeomorphism  $W_i(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$ , it follows that this diffeomorphism cannot have additional fixed points in the neighborhood in question. In other words,  $W_i(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$  and  $W_j(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$  have no common fixed point in  $B(\delta)$  as desired.

Let us begin by making accurate the first statement above.

**Lemma 5.1.** Consider the element  $W(\tilde{f}, \tilde{g})$  viewed as belonging to the pseudogroup  $\Gamma_{h_1,h_2}$ . Suppose that  $q \neq 0$  lies in the (open) domain of definition of  $W(\tilde{f}, \tilde{g})$ . Suppose also that  $W(\tilde{f}, \tilde{g})(q) = q$ . Then, there is  $(h_{1,*}, h_{2,*}) \in \mathrm{Diff}_{\alpha}(\mathbb{C}, 0) \times \mathrm{Diff}_{\alpha}(\mathbb{C}, 0)$  arbitrarily close to  $(h_1, h_2)$  and such that the following holds:

- (a) q lies in the (open) domain of definition of  $W(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$  viewed as element of the pseudogroup generated by  $h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*}$  on D. (b)  $W(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})(q) \neq q$ .

*Proof.* Condition (a) is always satisfied provided that  $(h_{1,*}, h_{2,*})$  is very close to  $(h_1, h_2)$ . Thus we only need to prove that, arbitrarily close to  $(h_1, h_2)$ , there is  $(h_{1,*}, h_{2,*}) \in \mathrm{Diff}_{\alpha}(\mathbb{C}, 0) \times$  $\operatorname{Diff}_{\alpha}(\mathbb{C},0)$  satisfying condition (b).

Consider the spelling of W(a,b) under the form  $W(a,b) = \vartheta_l^{r_l} * \cdots * \vartheta_1^{r_1}$ . The proof of the existence of  $(h_{1,*}, h_{2,*})$  satisfying condition (b) and arbitrarily close to  $(h_1, h_2)$  is going to be carried out by induction on l. Suppose first that l equals to 1. In this case, the statement follows at once from the fact that f does not have a Cremer point at  $0 \in \mathbb{C}$ , cf. Lemma 2.3.

By inducting on the length of the words, the proposition can be assumed to hold for words of length  $1, \ldots, l-1$ . We need to show that it also holds for words of length l. First consider the itinerary  $q = q_0, \ldots, q_{l-1}, q_l$  of q under  $W(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2) = W(\tilde{f}, \tilde{g}).$ By assumption we have  $q = q_0 = q_l$ . The induction assumption allows us to suppose that the points  $q_0, \ldots, q_{l-1}$  are pairwise distinct. Indeed, given  $0 \le k_1 < k_2 < l$ , we have that  $q_{k_2} = W'(f, \tilde{g})(q_{k_1})$  where W'(a, b) is a word whose length is at most l-1. Thus, by the induction assumption,  $(h_1, h_2)$  can be perturbed into  $(h_{1,*}, h_{2,*}) \in \mathrm{Diff}_{\alpha}(\mathbb{C}, 0) \times \mathrm{Diff}_{\alpha}(\mathbb{C}, 0)$ 

so as to satisfy  $q_{k_2} = W'(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})(q_{k_1}) \neq q_{k_1}$ . Since, once obtained, the condition  $q_{k_2} = W'(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})(q_{k_1}) \neq q_{k_1}$  is open, the fact that there are only finitely many words W'(a,b) that need to be considered allows us to construct a first perturbation  $(\tilde{h}_{1,*}, \tilde{h}_{2,*}) \in \operatorname{Diff}_{\alpha}(\mathbb{C}, 0) \times \operatorname{Diff}_{\alpha}(\mathbb{C}, 0)$  of  $(h_1, h_2)$  so that the itinerary of  $q = q_0$  by  $W(\tilde{h}_{1,*}^{-1} \circ f \circ \tilde{h}_{1,*}, \tilde{h}_{2,*}^{-1} \circ g \circ \tilde{h}_{2,*})$  satisfies the required condition. In other words, we can assume without loss of generality that the itinerary  $q = q_0, \ldots, q_{l-1}, q_l$  of q under  $W(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)$  is such that the points  $q_0, \ldots, q_{l-1}$  are pairwise distinct.

Let us now construct pairs of local diffeomorphisms  $(h_{1,*}, h_{2,*}) \in \operatorname{Diff}_{\alpha}(\mathbb{C}, 0) \times \operatorname{Diff}_{\alpha}(\mathbb{C}, 0)$  arbitrarily close to  $(h_1, h_2)$  and such that  $W(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})(q) \neq q$ . First, since  $W(a, b) = \vartheta_l^{r_l} * \cdots * \vartheta_1^{r_1}$ , with  $l \geq 2$ , we shall assume that  $\vartheta_1$  takes on the value a (with  $r_1 > 0$ ) and that  $\vartheta_l$  takes on the value b (with  $r_l > 0$ ). Note that, if the word W(a, b) is such that  $\vartheta_1, \vartheta_l$  takes on the same value (a or b), then W(f, g) is conjugate to a word of smaller length and the desired conclusion can immediately be derived.

Let P be a polynomial such that  $P(q_0) = \cdots = P(q_{l-2}) = 0$  and  $P(q_{l-1}) \neq 0$ . Since  $\vartheta_l$  takes on the value b, we set

$$h_{1,t} = h_1$$
 and  $h_{2,t} = h_2 + tz^{\alpha+1}P$ 

where  $t \in [0,1]$ . Clearly  $h_{2,t}$  converges to  $h_2$  in the analytic topology when  $t \to 0$  and  $h_{2,t} \in \text{Diff}_{\alpha}(\mathbb{C},0)$  for every  $t \in [0,1]$ . Therefore, to conclude the proof, it suffices to show that  $W(h_{1,t}^{-1} \circ f \circ h_{1,t}, h_{2,t}^{-1} \circ g \circ h_{2,t})(q) \neq q$  for arbitrarily small t > 0 (strictly). As already observed, for t sufficiently small q belongs to the domain of definition of  $W(h_{1,t}^{-1} \circ f \circ h_{1,t}, h_{2,t}^{-1} \circ g \circ h_{2,t})$  viewed as an element of the pseudogroup generated on the open disc D by  $h_{1,t}^{-1} \circ f \circ h_{1,t}, h_{2,t}^{-1} \circ g \circ h_{2,t}$ . The corresponding itinerary is going to be denoted by  $q = q_{0,t}, \ldots, q_{l-2,t}, q_{l-1,t}$  and  $q_{l,t}^l = h_{2,t}^{-1} \circ g^{r_l} \circ h_{2,t}(q_{l-1,t})$ . By construction, it follows that  $q_k = q_{k,t}$  for  $k = 0, 1, \ldots, l-1$ . However,  $h_{2,t}(q_{l-1}) = h_{2,t}(q_{l-1,t}) \neq h_2(q_{l-1,t})$ . Now the assumption concerning the injective character of both  $h_2^{\pm 1}, g^{\pm 1}$  on the domains in question implies that  $q = q_0 = q_l = W(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)(q_0) \neq W(h_{1,t}^{-1} \circ f \circ h_{1,t}, h_{2,t}^{-1} \circ g \circ h_{2,t})(q_0)$  for every t > 0 sufficiently small. The lemma is proved.

As already been mentioned, our strategy consists of showing that perturbations as in Lemma 5.1 can be applied to  $W_j(\tilde{f}, \tilde{g})$  while keeping q as a fixed point of  $W_i(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$ . The rest of the material goes in this direction.

Let us write  $W_i(a,b) = \vartheta_l^{r_l} * \cdots * \vartheta_1^{r_1}$  and  $W_j(a,b) = \vartheta_m^{s_m} * \cdots * \vartheta_1^{s_1}$ . Note that  $\vartheta_1$  on  $W_i$  does not necessarily take on the same value of  $\vartheta_1$  on  $W_j$ . Nonetheless, although the same notation is used for simplicity, throughout the text, each time we refer to  $\vartheta_1$  it will be explicitly mentioned if we are considering  $\vartheta_1$  on  $W_i$  or in  $W_j$ . Modulo re-labeling these two words, we always assume that  $m \leq l$ . The construction of the required perturbation  $(\overline{h}_1, \overline{h}_2)$  will be carried out by induction on m, i.e. on the length of the shorter word. To initialize the induction, note that for every pair  $(h_1, h_2) \in \operatorname{Diff}_{\alpha}(\mathbb{C}, 0) \times \operatorname{Diff}_{\alpha}(\mathbb{C}, 0)$ , the map  $W_j(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)$  has no isolated fixed points provided that m = 1 as it follows from the fact that none of the local diffeomorphisms f, g has a Cremer point at the origin, cf. Lemma 2.3. The statement is then immediately true regardless of the value of l. Therefore, by means of the induction, we assume that the existence of the desired perturbation  $(\overline{h}_1, \overline{h}_2)$  was already established for  $1, \ldots, m-1$  and every  $l \in \mathbb{N}$ . All we need to prove is the existence

of  $(\overline{h}_1, \overline{h}_2)$  arbitrarily close to  $(h_1, h_2)$  such that q remains fixed by exactly one of the words  $W_i(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$  and  $W_j(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$ .

**Lemma 5.2.** We can assume that  $W_i(a,b)$  (resp.  $W_j(a,b)$ ) is not a power of some (third) word  $W_3(a,b)$  verifying  $W_3(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)(q) = q$ .

Proof. Consider first  $W_i(a,b)$  and suppose it is a power of  $W_3(a,b)$  verifying  $W_3(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)(q) = q$ . In this situation, to establish Lemma 4.2, it suffices to work with  $W_3(a,b)$  and  $W_j(a,b)$ . In fact, if the statement is verified for these two latter words, the "new" fixed point  $\overline{q}$  of  $W_3(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$  is automatically a fixed point for  $W_i(\overline{h}_1^{-1} \circ f \circ \overline{h}_1, \overline{h}_2^{-1} \circ g \circ \overline{h}_2)$  in  $B(\delta)$ .

The case of  $W_j(a,b)$  is automatic: if  $W_j(a,b)$  is a power of  $W_3(a,b)$  verifying  $W_3(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)(q) = q$ , then the induction assumption ensures the existence of an arbitrarily small perturbation  $(h_{1,*}, h_{2,*})$  such that  $W_i(\overline{h_1}^{-1} \circ f \circ \overline{h_1}, \overline{h_2}^{-1} \circ g \circ \overline{h_2})(q) \neq q$  while  $W_3(\overline{h_1}^{-1} \circ f \circ \overline{h_1}, \overline{h_2}^{-1} \circ g \circ \overline{h_2})(q) = q$ . Clearly q must still be the unique fixed point of  $W_j(\overline{h_1}^{-1} \circ f \circ \overline{h_1}, \overline{h_2}^{-1} \circ g \circ \overline{h_2})$  in  $B(\delta)$  so that the lemma follows.  $\square$ 

Next consider the itinerary  $q=q_0^1,\ldots,q_l^1=q_0^1$  of q under  $W_i(\tilde{f},\tilde{g})$  and the itinerary  $q=q_0^2,\ldots,q_m^2=q_0^2$  of q under  $W_j(\tilde{f},\tilde{g})$ .

**Lemma 5.3.** We can suppose, without loss of generality, that  $q_k^1 \neq q_0^1$  for k = 1, ..., l-1 and that  $q_k^2 \neq q_0^2$  for k = 1, ..., m-1.

*Proof.* It follows from the same argument of the proof of Lemma 5.2. In fact, if this condition is not verified, the word  $W_i(a,b)$  (resp.  $W_j(a,b)$ ) can be split into shorter words and it will be enough to work with these latter words to eventually arrive to a situation where the condition in question is satisfied.

Nonetheless, with the preceding notations, Lemma 5.1 does not allow us to suppose that the points  $q = q_0^1, \ldots, q_{l-1}^1$  are pairwise disjoint since, when carrying out the perturbations described in this lemma, it may happen that the condition  $q_0^1 = q_l^1$  becomes no longer fulfilled. Analogous considerations apply to the itinerary  $q = q_0^2, \ldots, q_m^2 = q_0^2$ . For this reason, we shall need a version of Lemma 5.1 adapted to the present setting. However, to state this result, some new terminology is needed.

As always all words W(a, b) are supposed to be non-empty and reduced. A word W(a, b) of length l is said to be a conjugate of type 1 of a shorter word if there are words  $W_1(a, b)$  and  $W_2(a, b)$  such that the spelling of W(a, b) has the form  $[W_1(a, b)]^{-1} * W_2(a, b) * W_1(a, b)$ , where the concatenation "\*" leads to no simplification. This last assumption implies that the length of  $W_2(a, b)$  plus twice the length of  $W_1(a, b)$  equals the length of W(a, b), i.e. it equals l. Therefore, a word W(a, b) of length l is a conjugate of type 1 of a shorter word if the usual spelling  $W(a, b) = \vartheta_l^{r_l} * \cdots * \vartheta_1^{r_1}$  of W(a, b) has the form

$$[\vartheta_{l}^{r_{l}}*\cdots*\vartheta_{l-s_{0}+1}^{r_{l-s_{0}+1}}]*[\vartheta_{l-s_{0}}^{r_{l-s_{0}}}*\cdots*\vartheta_{s_{0}+1}^{r_{s_{0}+1}}]*[\vartheta_{s_{0}}^{r_{s_{0}}}*\cdots*\vartheta_{1}^{r_{1}}]$$

where  $[\vartheta_l^{r_l} * \cdots * \vartheta_{l-s_0+1}^{r_{l-s_0+1}}]$  represents the inverse of  $[\vartheta_{s_0}^{r_{s_0}} * \cdots * \vartheta_1^{r_1}]$ . Naturally this construction does not affect the domain of definition of W(a,b) as element of the corresponding

pseudogroup. The word  $W_2(a, b)$  is then a (shorter) conjugate of W(a, b). Naturally this definition does not unequivocally characterize  $W_1(a, b)$ ,  $W_2(a, b)$ . However it does characterize the *minimal conjugate of type* 1  $W_2(a, b)$  of W(a, b) which corresponds to  $W_1(a, b)$ ,  $W_2(a, b)$  as above such that  $W_2(a, b)$  is not a conjugate of type 1 of a shorter word.

Let now  $W(a,b) = [W_1(a,b)]^{-1} * W_2(a,b) * W_1(a,b)$  where  $W_2$  is the minimal conjugate of type 1 of W(a,b). Setting  $W_2(a,b) = \vartheta_{l-s_0}^{r_{l-s_0}} * \cdots * \vartheta_{s_0+1}^{r_{s_0+1}}$ , the minimal conjugate  $W_4(a,b)$  of W(a,b) is defined as follows:

- (1) If  $\vartheta_{l-s_0}$  and  $\vartheta_{s_0+1}$  take on different values (negative exponents allowed), then the minimal conjugate  $W_4(a,b)$  of W(a,b) (or equivalently of  $W_2(a,b)$ ) is  $W_2(a,b)$  itself.
- (2) If both  $\vartheta_{l-s_0}$  and  $\vartheta_{s_0+1}$  take on the same value, say a, then the minimal conjugate  $W_4(a,b)$  of W(a,b) (or equivalently of  $W_2(a,b)$ ) is given by

$$\vartheta_{l-s_0}^{r_{l-s_0}-r_{s_0+1}} * \cdots * \vartheta_{s_0+2}^{r_{s_0+2}}$$

In particular, a word  $W(a,b) = \vartheta_l^{r_l} * \cdots * \vartheta_1^{r_1}$  coinciding with its own minimal conjugate must be such that  $\vartheta_1$  and  $\vartheta_l$  take on different values. The preceding definition is coherent in the sense that  $\vartheta_{s_0+2}$  and  $\vartheta_{l-s_0}$  takes on different values. Indeed,  $r_{l-s_0} - r_{s_0+1} \neq 0$  since  $W_2(a,b)$  is the minimal conjugate of type 1 of W(a,b).

Remark 5.4. Although in the definition of conjugate of type 1 the concatenation leads to no simplification, the same does not necessarily occurs with the definition of minimal conjugate. For example, consider the reduced word  $W(a,b) = a^{-1}ba^2$ . This word is not a conjugate of type 1 of a shorter word. In fact, albeit W(a,b) admits the spelling  $W(a,b) = a^{-1} * (ba) * a$ , this spelling leads to a simplification. Nonetheless, the minimal conjugate  $W_4(a,b)$  of W(a,b) does not coincide with W(a,b) itself. First, note that  $\vartheta_1$  and  $\vartheta_3$  take on the same value, namely a. Since  $s_0$  (in the definition of the minimal conjugate of type 1) is equal to zero, the length of the minimal conjugate should be equal to  $l - s_0 - (s_0 + 2) + 1$ , i.e. equal to two. Indeed, the minimal conjugate of W(a,b) coincides with the word ab since W(a,b) can be written under the form  $W(a,b) = a^{-2} * (ab) * a^2$ , where a simplification on the concatenation "\*" can be used. This type of simplification is unique and it occurs only once.

With the previous notations, a version of Lemma 5.1 adapted to the present setting is as follows.

**Lemma 5.5.** Let W(a,b) be a word of length l and consider a point q in the domain of definition of  $W(\tilde{f},\tilde{g})=W(h_1^{-1}\circ f\circ h_1,\,h_2^{-1}\circ g\circ h_2)$  viewed as element of  $\Gamma_{h_1,h_2}$ . Suppose that  $W(\tilde{f},\tilde{g})$  has only isolated fixed points and that the multiplicity of each of these fixed points does not change under perturbations of  $(h_1,h_2)$ . Denote by  $q=q_0,\ldots,q_l$  the itinerary of q under l and assume that  $q_0 \notin \{q_1,\ldots,q_{l-1}\}$ . Then arbitrarily close to  $(h_1,h_2)$ , there is  $(\tilde{h}_1,\tilde{h}_2)$  such that the itinerary  $q=\tilde{q}_0,\tilde{q}_1,\ldots,\tilde{q}_l$  of q under  $W(\tilde{h}_1^{-1}\circ f\circ \tilde{h}_1,\tilde{h}_2^{-1}\circ g\circ \tilde{h}_2)$  satisfies the following conditions:

- (1) Suppose  $q_0 \neq q_l$ . Then  $\tilde{q}_l = q_l$  and the points  $q = q_0, \tilde{q}_1, \dots, \tilde{q}_{l-1}$  are pairwise distinct.
- (2) Suppose  $q_0 = q_l$  and W(a,b) is not a conjugate of a shorter word. Then again  $\tilde{q}_l = q_l = q_0$  and the points  $q = q_0, \tilde{q}_1, \dots, \tilde{q}_{l-1}$  are pairwise distinct.
- (3) Suppose  $q_0 = q_l$  and  $W(a,b) = [W_3(a,b)]^{-1} * W_4(a,b) * W_3(a,b)$ , where  $W_4(a,b)$  represents the minimal conjugate of W(a,b). Denote the length of  $W_3(a,b)$  (resp.  $W_4(a,b)$ ) by  $S_0$  (resp. U). If the natural assumption that  $Q_{s_0}$  is different from all the

points  $q_{s_0+1}, \ldots, q_{s_0+l'-1}$  is added, then the points  $q = \tilde{q}_0, \tilde{q}_1, \ldots, \tilde{q}_{s_0+l'-1}$  are pairwise distinct.

First let us make some comments concerning item (3) of Lemma 5.5. Consider words W(a,b),  $W_3(a,b)$  and  $W_4(a,b)$  as in item (3) of the previous lemma. Denote by l (resp.  $s_0$ , l') the length of the word W(a,b) (resp.  $W_3(a,b)$ ,  $W_4(a,b)$ ). Two cases may occur.

- (a) The minimal conjugate  $W_4(a, b)$  coincides with the minimal conjugate of type 1 of W(a, b). In this case  $l' + 2s_0 = l$ , i.e. the concatenation "\*" leads to no simplification in the spelling of  $[W_3(a, b)]^{-1} * W_4(a, b) * W_3(a, b)$ .
- (b) The minimal conjugate  $W_4(a, b)$  does not coincide with the minimal conjugate of type 1 of W(a, b). In this case  $l' + 2s_0 = l + 1$ , i.e. the concatenation "\*" leads to a (unique) simplification in the spelling of  $[W_3(a, b)]^{-1} * W_4(a, b) * W_3(a, b)$ .

Next, consider the itinerary  $q=q_0,\ldots,q_l$  of q under  $W(\tilde{f},\tilde{g})$  and the itinerary  $q=q'_0,\ldots,q'_p$  of q under  $([W_3]^{-1}*W_4*W_3)(\tilde{f},\tilde{g})$ , where p=l or l+1 according to we are in case (a) or in case (b). In the first case, the itinerary  $q_0,\ldots,q_l$  coincides with the itinerary  $q'_0,\ldots,q'_p$ . In the second case, these itineraries satisfy  $q'_k=q_k$  for  $1\leq k\leq s_0+l'-1$  and  $q'_{k+1}=q_k$  for  $s_0+l'\leq k\leq l$ . In other words, the two itineraries coincide up to the point  $q'_{s_0+l'}$ . This point is, in fact, "fictitious" for the itinerary in question in the sense that  $[W_3(a,b)]^{-1}*W_4(a,b)*W_3(a,b)$  is not written in a reduced way and itineraries should be defined only in this case. In fact, with the above notations, taking  $W_3=\vartheta_{s_0}^{r_{i_0}}*\cdots*\vartheta_1^{r_1}$  and  $W_4=\vartheta_{l'+s_0}^{r_{l'+s_0}}*\cdots*\vartheta_{s_0+1}^{r_{s_0+1}}$ , we have that  $\vartheta_{l'+s_0}$  and  $\vartheta_{s_0}^{-1}$  take on same same value if  $W_4$  does not coincide with the minimal conjugate of type 1 of W.

Item (3) of Lemma 5.5 ensures that the elements  $q = \tilde{q}_0, \tilde{q}_1, \dots, \tilde{q}_{s_0+l'-1}$  in the itinerary of q under  $W(\tilde{h}_1^{-1} \circ f \circ \tilde{h}_1, \tilde{h}_2^{-1} \circ g \circ \tilde{h}_2)$  are pairwise distinct. The "fictitious" point is not included in this list and this will play a role in the proof of Lemma 4.2.

To not interrupt the discussion, let us first conclude the proof of Lemma 4.2 before proving Lemma 5.5.

*Proof of Lemma 4.2.* Let us begin by summarizing what have already been mentioned concerning the proof of this lemma.

Let  $W_i(a,b) = \vartheta_l^{r_l} * \cdots * \vartheta_1^{r_1}$  and  $W_j(a,b) = \vartheta_m^{s_m} * \cdots * \vartheta_1^{s_1}$  and assume that  $m \leq l$ . The construction of the required perturbation  $(\overline{h}_1,\overline{h}_2)$  will be carried out by induction on m. It has already been mentioned that the statement is true for m=1. Assume that the existence of the desired perturbation  $(\overline{h}_1,\overline{h}_2)$  was already established for  $1,\ldots,m-1$  and every  $l \in \mathbb{N}$ . Let us now construct  $(\overline{h}_1,\overline{h}_2)$  arbitrarily close to  $(h_1,h_2)$  and such that q remains fixed by exactly one of the elements  $W_i(\overline{h}_1,\overline{h}_2)$  and  $W_j(\overline{h}_1,\overline{h}_2)$ . The result will immediately follow in this case.

Let  $W_4(a,b)$  denote the minimal conjugate of  $W_i(a,b)$  and set

$$W_i(a,b) = [W_3(a,b)]^{-1} * W_4(a,b) * W_3(a,b),$$

where  $W_3(a, b)$  is empty if  $W_i(a, b)$  coincides with its minimal conjugate. The length of the minimal conjugate  $W_4(a, b)$  is going to be denoted by l'. If l' < l, then  $s_0$  will denote the length of the word  $W_3(a, b)$ . The induction assumption allows us to suppose that  $l' \ge m$ , otherwise we can conjugate the whole group by  $W_3(a, b)$  and the problem will be reduced to eliminate the corresponding common fixed point between  $W_4(\tilde{f}, \tilde{g})$  and another word  $\overline{W}(\tilde{f}, \tilde{g})$ 

which which is conjugate to  $W_j(\tilde{f}, \tilde{g})$ . Since the length of  $W_4(a, b)$  is l' < m the induction assumption implies that the common fixed point in question can effectively be eliminated. Similarly it is clear that the word  $W_j(a, b)$ , of length m, is not a conjugate of a shorter word since otherwise the statement results immediately.

In view of the preceding, in the sequel we always have  $l' \geq m$ . The proof is divided in three cases. First we will assume that l' is strictly greater than m, i.e. l' > m. When l' = m two further cases need to be considered, according to whether or not  $W_3(a, b)$  is void, i.e. according to  $s_0 \geq 1$  or  $s_0 = 0$ .

Case 1. Suppose first that l'>m strictly. This case will be handled with a useful general observation. By using Lemma 5.5, the points  $q=q_0^1,\ldots,q_{s_0+l'-1}^1$  can be supposed pairwise distinct. Also, it is clear that the points  $q_{s_0+1}^1,\ldots,q_{s_0+l'-1}^1$  appears exactly once in the (full) itinerary of q under  $W_i(\tilde{f},\tilde{g})$ . Since l'>m and  $q_0^2=q_m^2=q_0^1\not\in\{q_{s_0+1}^1,\ldots,q_{s_0+l'-1}^1\}$ , it follows the existence of a point  $q_N^1$ ,  $i_0< N< k_0+l'$ , which does not belong to the itinerary of q under  $W_j(h_1^{-1}\circ f\circ h_1,h_2^{-1}\circ g\circ h_2)$ , i.e. to the set  $\{q_0^2,\ldots,q_m^2\}$ . Without loss of generality, the value of  $\vartheta_{m+1}$  can be supposed to be a. In this case, we shall consider perturbations  $h_{1,t}$  of  $h_1$  given by

(1) 
$$h_{1,t}(z) = h_1(z) + tz^{\alpha+1}P(z)$$

where P is a polynomial vanishing at all points  $q_k^1$ ,  $k \neq N$  and  $k \in \{0, \ldots, l\}$ , and over all points  $q_k^2$ ,  $k \in \{0, \ldots, m\}$ . Yet, P verifies  $P(q_N^1) \neq 0$ . Let then  $h_{1,*} = h_{1,t}$ , for some sufficiently small t > 0, and  $h_{2,*} = h_2$ . Clearly  $W_j(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})(q) = q$ . It remains to check that  $W_i(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})(q) \neq q$ . Setting  $W_i(a,b) = W_B(a,b) * W_A(a,b)$  where  $W_A(a,b) = \vartheta_{N+1}^{r_{N+1}} * \cdots * \vartheta_1^{r_1}$  and  $W_B(a,b) = \vartheta_l^{r_l} * \cdots * \vartheta_{N+2}^{r_{N+2}}$  (being  $W_B(a,b)$  possibly empty), by construction, we have:  $W_B(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})(q_0^1) \neq q_{N+1}^1$ . Since all the maps involved are one-to-one, we conclude that  $W_i(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})(q_0^1) \neq q_{N+1}^1$ . Since all the statement is proved in this first case.

Note that in the construction above it was implicitly used the fact that  $q_N^1$ , the element in the itinerary of q under  $W_i(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)$  which does not belong to the itinerary of q under  $W_j(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)$ , does not correspond to the above mentioned "fictitious" point. In fact, if this were the case, then both  $\vartheta_{l'+s_0}$  in  $W_4$  and  $\vartheta_{s_0}^{-1}$  in  $W_3^{-1}$  would take on same value and, therefore, the effect of P would be void.

Case 2. Suppose now that l'=m. The preceding argument can then easily be adapted to handle the case where  $s_0 \geq 1$ . The main difference between the present case and the above discussion lies in the fact that  $q_N^i$ , the element in the itinerary of q under  $W_i(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)$ , which does not belong to the itinerary of q under  $W_j(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)$ , may appear "twice" in the (full) itinerary of q under  $W_i(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)$ . In other words, N belongs to  $\{1, \ldots, s_0 + l' - 1\}$  and not necessarily to  $\{s_0 + 1, \ldots, s_0 + l' - 1\}$  as in the previous case. Let us present the adaptations required to establish the result in this context.

If  $q_N^1$  appears only once in the full itinerary of q, then the proof follows as above. So, let us assume that  $q_N^1$  appears twice and that the value of  $\vartheta_{N+1}$  is a. Let us first consider perturbations  $h_{1,t}$  of  $h_1$  given by

$$h_{1,t}(z) = h_1(z) + tz^{\alpha+1}P(z)$$

where P is a polynomial vanishing at all points  $q_k^1$ ,  $k \neq N$  and  $k \in \{0, ..., l\}$  and over all points  $q_k^2$ ,  $k \in \{0, ..., m\}$ . Furthermore  $P(q_N^1)$  should be different from zero. Let then  $h_{1,*} = h_{1,t}$ , for some sufficiently small t > 0, and  $h_{2,*} = h_2$ . Clearly  $W_j(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})(q) = q$ . Nonetheless we cannot yet ensure that  $W_i(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})(q) \neq q$ . To begin with, let us describe when q is a fixed point for  $W_i(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$ .

We are assuming that  $q_N^1$  appears twice in the itinerary of q so that  $q_N^1$  coincides with  $q_{l-N}^1$ . Without loss of generality, the value of  $\vartheta_{N+1}$  can be supposed to be a. If  $\vartheta_{l-N+1}$  takes on value b then the argument presented in the previous case ensures that q is not a fixed point for  $W_i(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$ . So, it can be assumed that  $\vartheta_{l-N+1}$  also takes on value a. It can easily be checked that q is a fixed point for  $W_i(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$  if and only if

(2) 
$$h_{1,t} \circ \vartheta_{l-N}^{r_{l-N}} \circ \cdots \circ \vartheta_{N+1}^{r_{N+1}}(q_N^1) = h_1(q_N^1)$$

where  $\vartheta_k$  is substituted by  $h_{1,t}^{-1} \circ f \circ h_{1,t}$  (resp.  $h_{2,t}^{-1} \circ g \circ h_{2,t}$ ) if  $\vartheta_k$  takes on value a (resp. b). Therefore, to make sure that q is not a fixed point for  $W_i(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$ , we should guarantee that  $h_{1,t}^{-1} \circ h_1(q_N^1) \neq \vartheta_{l-N}^{r_{l-N}} \circ \cdots \circ \vartheta_{N+1}^{r_{N+1}}(q_N^1)$ . Let us assume that  $h_{1,t}^{-1} \circ h_1(q_N^1) = \vartheta_{l-N}^{r_{l-N}} \circ \cdots \circ \vartheta_{N+1}^{r_{N+1}}(q_N^1)$ . Let Q be a polynomial vanishing at all points  $q_k^1, k \in \{0, \ldots, l\}$  and over all points  $q_k^2, k \in \{0, \ldots, m\}$ . Let also Q verify  $Q(h_{1,t}^{-1} \circ h_1(q_N^1)) \neq 0$ . Now, since t is arbitrary, it can be chosen so that  $h_{1,t}^{-1} \circ h_1(q_N^1) \notin \{q_0^1, \ldots, q_l^1\} \cup \{q_0^2, \ldots, q_m^2\}$ . Consider the perturbations of  $h_1$  given by

$$\overline{h}_{1,t} = h_{1,t} + tz^{\alpha+1}Q(z).$$

Let  $\overline{h}_{1,*} = \overline{h}_{1,t}$ , for some sufficiently small t > 0, and  $\overline{h}_{2,*} = h_2$ . Clearly  $W_j(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})(q) = q$ . Moreover, since  $Q(h_{1,t}^{-1} \circ h_1(q_N^1)) \neq 0$ , Condition (2) is no longer verified for  $(\overline{h}_{1,*}, \overline{h}_{2,*})$  and, therefore,  $W_i(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})(q) \neq q$ . The statement is proved in this case.

Case 3. Summarizing what precedes, it only remains to deal with words  $W_i(a,b)$ ,  $W_j(a,b)$  of same length (i.e. l=m) and none of them being a conjugate of a shorter word. In particular, both words  $W_i(a,b)$ ,  $W_j(a,b)$  are on the same "footing" concerning the argument below. Next note that, since none of them is conjugate to a shorter word, Lemma 5.5 allows us to suppose without loss of generality that the points  $q=q_0^1,\ldots,q_{l-1}^1$  (resp.  $q=q_0^2,\ldots,q_{m-1}^2$ ) are pairwise distinct. In particular, unless these two itineraries coincide up to relabeling the points, there exists again  $q_N^1$ , 0 < N < l, which does not belong to the set  $\{q_0^2,\ldots,q_m^2\}$ . Thus the preceding argument can still be employed to settle the lemma.

Finally, we only have to deal with the case where the itineraries of q under  $W_i(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)$  and under  $W_j(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)$  coincide up to relabeling the points. The discussion splits in three subcases.

Case 3a. Suppose that in the word  $W_i(a,b)$ ,  $\vartheta_1$  takes on the value a whereas, in the word  $W_j(a,b)$ ,  $\vartheta_1$  takes on the value b (or the other way around). Consider then perturbations  $h_{1,t}$  of  $h_1$  having the form (1), where, this time,  $P(q) \neq 0$  and P vanishes over all the remaining points of the common itinerary  $q = q_0^1, \ldots, q_{l-1}^1$ . Again we set  $h_{1,*} = h_{1,t}$ , for small t > 0, and  $h_{2,*} = h_2$ . Since  $q_k^2 \neq q_0$  for every  $k = 1, \ldots, k-1$ , it follows that  $W_j(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})(q) = q$ . On the other hand, the same argument employed

above shows that  $W_i(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)(q) \neq q$  and finishes the proof of the proposition in the present case.

Case 3b. Suppose that in both words  $W_i(a,b)$ ,  $W_j(a,b)$ ,  $\vartheta_1$  takes on the same value, say a, but we have  $r_1 \neq s_1$  (where the exponents are allowed to be negative). Since  $0 \in \mathbb{C}$  is not a Cremer point for f, it follows that  $q_1^1 \neq q_1^2$ . However  $q_1^2$  must belong to the itinerary of q under  $W_i(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)$ . Let then  $q_N^1 = q_1^2$  where  $N \in \{2, \ldots, m-1\}$  (recall that l = m). If N < m-1 consider the word  $W^{(1)}(a,b) = W_B * W_A$  where  $W_B(a,b) = a^{-s_1}$  and  $W_A(a,b) = \vartheta_m^{r_m} * \cdots * \vartheta_1^{r_1}$ , where  $\vartheta_k$  takes on the same value taken in the spelling of  $W_i(a,b)$ , for  $k=1,\ldots,N$ . It follows that  $W^{(1)}(a,b)$  is a word of length less than m satisfying  $W^{(1)}(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2)(q) = q$ . Hence, by induction assumption, we can destroy the fact that q is a common fixed point for  $W^{(1)}(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$  and  $W_{i_2}(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$ .

Naturally we can assume that  $W_i(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*}), W_j(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$  still have a common fixed point in  $B(\delta)$ , otherwise the statement is established. Let us still denote by q this common fixed point. We claim that now the itineraries of q under  $W_i(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$  and under  $W_j(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$  do not coincide, even though these points are relabeled. In fact, denote by  $q = \tilde{q}_0^1, \ldots, \tilde{q}_m^1$  (resp.  $q = \tilde{q}_0^2, \ldots, \tilde{q}_m^2$ ) the itinerary of q under  $W_i(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$  (resp.  $W_j(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$ ). The perturbation  $h_{1,*}$  can be chosen so that

$$||q_k^1 - \tilde{q}_1^1|| < \tau/4$$
 and  $||q_k^2 - \tilde{q}_1^2|| < \tau/4$ 

for all  $k=1,\ldots,m$ , where  $\tau$  denotes a lower bound for the distance between two distinct points on each itinerary of q under  $W_i(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$ ,  $W_j(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$ . Assume for a contradiction that the itineraries  $q=\tilde{q}_0^1,\ldots,\tilde{q}_m^1$  and  $q=\tilde{q}_0^2,\ldots,\tilde{q}_m^2$  coincide up to relabeling the points. The estimate above guarantees that  $\tilde{q}_1^2$  again coincides with  $\tilde{q}_N^2$ . Therefore q is a common fixed point for  $W^{(1)}(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$  and  $W_j(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$ , contradicting the choice of  $h_{1,*}$ . Thus, we have just proved that  $\tilde{q}_N^1 \notin \{\tilde{q}_0^2,\ldots,\tilde{q}_m^2\}$ . Therefore the same arguments employed in the previous cases can now be brought to settle the lemma in the present case as well.

Let us finally consider the case N=m-1. We are going to show that the induction assumption can also be used to eliminate this situation as well. This goes as follows. Set  $W^{(1)}(a,b)=W_B*W_A$  where  $W_A(a,b)=a^{r_1-s_1}$  and  $W_B(a,b)=\vartheta_m^{r_m}*\cdots*\vartheta_2^{r_2}$ . Then  $W^{(1)}(a,b)$  is a word of length m-1 such that  $W^{(1)}(h_1^{-1}\circ f\circ h_1,\,h_2^{-1}\circ g\circ h_2)(q_1^2)=q_1^2$ . On the other hand  $q_1^2$  is also fixed by  $W^{(2)}(h_1^{-1}\circ f\circ h_1,\,h_2^{-1}\circ g\circ h_2)$  where  $W^{(2)}(a,b)$  is the word of length m obtained from  $W_j$  through conjugation under  $a^{-s_1}$ , i.e.  $W^{(2)}(a,b)=a^{s_1}*W_j(a,b)*a^{-s_1}$ . Since one of this words has length strictly smaller then m, the induction assumption allows us to eliminate the common fixed point  $q_1^2$ . Therefore, if  $W_i(h_{1,*}^{-1}\circ f\circ h_{1,*},\,h_{2,*}^{-1}\circ g\circ h_{2,*})$  and  $W_j(h_{1,*}^{-1}\circ f\circ h_{1,*},\,h_{2,*}^{-1}\circ g\circ h_{2,*})$  still have a common fixed point, again we claim that the itineraries of q under  $W^{(1)}(h_{1,*}^{-1}\circ f\circ h_{1,*},\,h_{2,*}^{-1}\circ g\circ h_{2,*})$  and under  $W_j(h_{1,*}^{-1}\circ f\circ h_{1,*},\,h_{2,*}^{-1}\circ g\circ h_{2,*})$ . Now the same argument can again be applied to eliminate this common fixed point. The lemma is proved in subcase 3b.

Case 3c. Suppose that in both words  $W_i(a, b)$ ,  $W_j(a, b)$ ,  $\vartheta_1$  takes on the same value, say a, and that  $r_1 = s_1$ . In this case we have  $q_1^1 = q_1^2$ . This case amounts to consider the new

words  $\widetilde{W}_i(a,b)$  and  $\widetilde{W}_j(a,b)$  of length m and given respectively by  $a^{r_1} * \vartheta_m^{r_m} * \cdots * \vartheta_2^{r_2}$  and  $a^{r_1} * \vartheta_m^{s_m} * \cdots * \vartheta_2^{s_2}$ . Clearly  $q_1^1 = q_1^2$  is a common fixed point for these words. Note that  $\vartheta_2$  takes on the same value b. However unless  $r_2 = s_2$ , this common fixed point can be destroyed by means of the preceding arguments. Otherwise, we shall have  $q_2^1 = q_2^2$  and we may move on to the new words spelled out respectively as  $b^{r_2} * a^{r_1} * \vartheta_m^{r_m} * \cdots * \vartheta_3^{r_3}$  and as  $b^{r_2} * a^{r_1} * \vartheta_m^{s_m} * \cdots * \vartheta_3^{s_3}$  which have  $q_1^2 = q_2^2$  as common fixed points. Continuing in this way, since  $W_i(a,b), W_j(a,b)$  are incommensurable, at some step or order N we must have  $r_N \neq s_N$ . At this moment the common fixed point can be destroyed and the rest of the statement will quickly follows. This ends the proof of Lemma 4.2.

To finish the paper, the last step is to supply the proof of Lemma 5.5.

Proof of Lemma 5.5. The statement can naturally be proved by inducting on the length l of W(a,b). To begin with, recall that W(a,b) is spelled out as  $W(a,b) = \vartheta_l^{r_l} * \cdots * \vartheta_1^{r_1}$ . Consider the itinerary  $q = q_0, \ldots, q_l$  of q under  $W(\tilde{f}, \tilde{g})$ . Let us first consider the situations described in items (1) or (2). Then the statement amounts to showing that, modulo perturbing  $(h_1, h_2)$  into certain  $(\tilde{h}_1, \tilde{h}_2)$ , arbitrarily close to  $(h_1, h_2)$ , we obtain an itinerary  $q = \tilde{q}_0, \ldots, \tilde{q}_l$  such that the points  $q = \tilde{q}_0, \ldots, \tilde{q}_{l-1}$  are pairwise distinct. The idea is to use a perturbation similar to the one employed in Lemma 5.1 to make the mentioned points pairwise distinct and then to "locally correct" it by adding a new localized perturbation that will ensure the points  $q_0, q_l$  do not move at the final situation.

Let then  $\epsilon > 0$  be fixed. We shall look for  $(\tilde{h}_1, \tilde{h}_2)$   $\epsilon$ -close to  $(h_1, h_2)$  satisfying the required conditions where, by saying that  $(\tilde{h}_1, \tilde{h}_2)$  is  $\epsilon$ -close to  $(h_1, h_2)$ , it is meant that  $\max\{d_A(\tilde{h}_1, h_1), d_A(\tilde{h}_2, h_2)\} < \epsilon$ .

To fix notations, assume also that  $\vartheta_1$  takes on the value a since the other possibility is totally analogous (throughout the discussion negative exponents are allowed). Let us first suppose that  $q = q_0 \neq q_l$ . Because  $q_0$  is supposed to be different from each of the points  $q_1, \ldots, q_{l-1}$ , cf. Lemma 5.3, in addition to  $q_l$ , there is  $\tau > 0$  so that, for every  $k \in \{1, \ldots, l\}$  we have  $|q_0 - q_k| > 2\tau$ . In other words, the distances of the mentioned points  $q_k$  to  $q_0$  is bounded from below by a positive constant.

Now consider the word  $\overline{W}(a,b)$  which is nothing but W(a,b) spelled out in the "reverse order and with exponents of opposite signs" (i.e.  $\overline{W}(a,b)$  is the inverse element of W(a,b)). In particular,  $q_l, \ldots, q_0$  becomes the itinerary of  $q_l$  under  $\overline{W}(h_1^{-1} \circ f \circ h_1, h_2^{-1} \circ g \circ h_2) = \overline{W}(\tilde{f}, \tilde{g})$  whose "final point" is  $q_0$ . By successively applying Lemma 5.1, we can find an arbitrarily small perturbation  $(h_{1,*}, h_{2,*})$  of  $(h_1, h_2)$  so that the itinerary  $q_l = q'_l, q'_{l-1}, \ldots, q'_0$  of  $q_l$  under  $\overline{W}(h_{1,*}^{-1} \circ f \circ h_{1,*}, \frac{1}{2,*} \circ g \circ h_2, *)$  is constituted by pairwise distinct points. In addition, we can assume that  $(h_{1,*}, h_{2,*})$  is  $\epsilon/2$ -close to  $(h_1, h_2)$  whereas, for every  $k = 0, \ldots, l$ , the distance  $|q_k - q'_k|$  is arbitrarily small. In particular we have

$$(3) |q_0 - q_0'| < \tau.$$

Modulo choosing  $(h_{1,*}, h_{2,*})$  closer to  $(h_1, h_2)$ , we are going to construct a new perturbation  $\tilde{h}_1$  of  $h_{1,*}$ , verifying  $d_A(\tilde{h}_1, h_{1,*}) < \epsilon/2$ , and such that  $(\tilde{h}_1, \tilde{h}_2) = (\tilde{h}_1, h_{2,*})$  satisfies the required conditions. To construct  $\tilde{h}_1$  consider the elementary Lagrange interpolation consisting of a Polynomial P (of degree at most l) satisfying the following conditions:

(1) 
$$P(q_1') = \cdots = P(q_l') = 0.$$

(2) 
$$P(q_0) = h_{1,*}(q'_0) - h_{1,*}(q_0).$$

Thanks to Estimate (3), it follows that all coefficients of P converge to 0 provided that all the distances  $|q_0 - q_0'|$  do so. Note that Estimate (3) guarantees that  $q_0'$  does not coincide with any one of the elements  $q_1', \ldots, q_l'$ . Since the degree of P is uniformly bounded, it follows that P converges in the analytic topology for the null function. In particular, modulo taking  $(h_{1,*}, h_{2,*})$  sufficiently close to  $(h_1, h_2)$  in the analytic topology, we conclude that  $\tilde{h}_1 = h_{1,*} + P$  is  $\epsilon/2$ -close to  $h_{1,*}$  and, hence,  $\epsilon$ -close to  $h_1$ . Finally it is clear that the itinerary of  $q_l = q_l'$  under  $\overline{W}(\tilde{h}_1^{-1} \circ f \circ \tilde{h}_1, \tilde{h}_2^{-1} \circ g \circ \tilde{h}_2)$  is simply  $q_l = q_l', q_{l-1}', \ldots, q_1', q_0$ . The lemma is proved in the first case.

Consider now the case where  $q_0 = q_l$ . The word W(a, b) is first supposed not to be conjugate to a shorter word. Recalling that  $\vartheta_1$  takes on the value a, we conclude that  $\vartheta_l$  takes on the value b. The proof then amounts to noticing that the same argument above applies: a first perturbation  $(h_{1,*}, h_{2,*})$  is constructed so that the itinerary  $q_l = q'_l, q'_{l-1}, \ldots, q'_0$  of  $q_l$  under  $\overline{W}(h_{1,*}^{-1} \circ f \circ h_{1,*}, h_{2,*}^{-1} \circ g \circ h_{2,*})$  is constituted by pairwise distinct points. To make  $q'_0$  to coincide with  $q_0 = q_l$ , just note that  $q'_{l-1}$  is not affected by the replacing of  $h_{1,*}$  by  $\tilde{h}_1$ , since  $\vartheta_l$  takes on the value b. This settles the second item in the statement of the lemma.

Finally let us consider the case where W(a, b) is conjugate to a shorter word. Set  $W(a, b) = [W_3(a, b)]^{-1} * W_4(a, b) * W_3(a, b)$  where  $W_4(a, b)$  is the minimal conjugate of W(a, b). This case is nothing but a blend of item (1), applied to  $W_3(a, b)$  and of item (2) applied to  $W_4(a, b)$ , details are left to the reader.

#### References

- [BLL-1] M, Belliart, I. Liousse & F. Loray, Sur l'existence de points fixes attractifs pour les sous-groupes de Aut  $(\mathbb{C}, 0)$ , C. R. Acad. Sci. Paris, **324**, Sér. I, (1997), 443-446.
- [BLL-2] M, BELLIART, I. LIOUSSE & F. LORAY, The generic differential equation  $dw/dz = P_n(w,z)/Q_n(w,z)$  on  $\mathbb{C}P(2)$  carries no interesting transverse structure, *Ergod. Th. & Dynam. Sys.*, **21**, (2001), 1599-1607.
- [C-G] L. CARLESON & T. GAMELIN, Complex Dynamics, Springer-Verlag, New York, (1993).
- [GM-W] X. Gomez-Mont & B. Wirtz, On fixed points of conformal pseudogroups, Bull. Braz. Math. Soc., 26, 2, (1995), 201-209.
- [II-1] Yu. Il'yashenko, Topology of phase portraits of analytic differential equations on a complex projective plane, *Trudy Sem. Petrovsk.*, **4**, (1978), 83-136.
- [II-2] Yu. Il'Yashenko, Global and local aspects of the theory of complex differential equations, *Proc. Int. Cong. Math. Helsinki*, Acad. Scient. Fennica, **2**, (1978), 821-826.
- [K-H] A. KATOK & B. HASSELBLATT, Introduction to the Modern Theory of Dynamical Systems, Encyclopedia of Mathematics and its Applications, Volume 54, Cambridge University Press, (1997).
- [LF] L. LE FLOCH, Rigidité générique des feuilletages Ann. Sc. de l'ENS, Sér. 4, 31, 6, (1998), 765-785.
- [Lo] F. LORAY, Pseudo-groupe d'une singularité de feuilletage holomorphe en dimension deux, available from hal.archives-ouvertes.fr/hal-00016434, (2005).
- [M-M] D. MARIN & J.-F. MATTEI, Incompressibilité des feuilles de germes de feuilletages holomorphes singuliers, Ann. Scient. de l'ENS, 4-Série, 41, (2008), 855-903.
- [M-R-R] J.-F. MATTEI, J.C. REBELO & H. REIS, Generic pseudogroups on  $(\mathbb{C},0)$  and the topology of leaves, to appear in Compositio Mathematica.
- [M-S] J.-F. MATTEI & E. SALEM, Modules formels locaux de feuilletages holomorphes, available from arXiv:math/0402256v1, (2004), 89 pages.
- [N] I. NAKAI, Separatrizes for non solvable dynamics on (C, 0), Ann. Inst. Fourier, 44, (1994), 569-599.

- [Sh] A. Shcherbakov, On the density of an orbit of a pseudogroup of conformal mappings and a generalization of the Hudai-Verenov theorem, *Vestinik Movskovskogo Universiteta Mathematika*, **31**, 4, (1982), 10-15.
- [Sh-R-O] A. SHCHERBAKOV, E. ROSALES-GONZLEZ & L. ORTIZ-BOBADILLA, Countable set of limit cycles for the equation  $dw/dz = P_n(z, w)/Q_n(z, w)$ , J. Dynam. Control Systems, 4, 4, (1998), 539-581.
- [T] F. Takens, A nonstabilizable jet of a singularity of a vector field: the analytic case, Algebraic and differential topology global differential geometry, Teubner-Text Math., 70, Teubner, Leipzig, (1984), 288-305.
- [Y] J.-C. Yoccoz, Centralisateurs et conjugaison différentiable des difféomorphismes du cercle. Petits diviseurs en dimension 1, *Astérisque*, **231**, (1995), 89-242.

# Julio Rebelo

Institut de Mathématiques de Toulouse 118 Route de Narbonne F-31062 Toulouse, FRANCE. rebelo@math.univ-toulouse.fr

# HELENA REIS

Centro de Matemática da Universidade do Porto, Faculdade de Economia da Universidade do Porto, Portugal hreis@fep.up.pt