

# COHOMOLOGICAL CHARACTERIZATION OF HYPERQUADRICS OF ODD DIMENSIONS IN CHARACTERISTIC TWO

KATSUHISA FURUKAWA

ABSTRACT. We consider characterizations of projective varieties in terms of their tangents. S. Mori established the characterization of projective spaces in arbitrary characteristic by ampleness of tangent bundles. J. Wahl characterized projective spaces in characteristic zero by cohomological condition of tangent bundles; in addition, he remarked that a counter-example in characteristic two is constructed from odd-dimensional hyperquadrics  $Q_{2n-1}$  with  $n > 1$ . This is caused by existence of a common point in  $\mathbb{P}^{2n}$  which every embedded tangent space to the quadric contains. In general, a projective variety in  $\mathbb{P}^N$  is said to be strange if its embedded tangent spaces admit such a common point in  $\mathbb{P}^N$ . A non-linear smooth projective curve is strange if and only if it is a conic in characteristic two (E. Lluís, P. Samuel). S. Kleiman and R. Piene showed that a non-linear smooth hypersurface in  $\mathbb{P}^N$  is strange if and only if it is a quadric of odd-dimension in characteristic two. In this paper, we investigate complete intersections, and prove that, a non-linear smooth complete intersection in  $\mathbb{P}^N$  is strange if and only if it is a quadric in  $\mathbb{P}^N$  of odd dimension in characteristic two; these conditions are also equivalent to non-vanishing of 0-cohomology of  $(-1)$ -twist of the tangent bundle.

## 1. INTRODUCTION

S. Mori [8] established the characterization of projective spaces in characteristic  $p \geq 0$  by ampleness of tangent bundles. The work has motivated approaches via tangential properties to characterizing several projective varieties. J. Wahl [12] characterized projective spaces in  $p = 0$  by cohomological condition of tangent bundles; in addition, he remarked that a counter-example in  $p = 2$  is constructed from odd-dimensional hyperquadrics  $Q_{2n-1}$  with  $n > 1$  [12, p. 316]. This is caused by existence of a common point  $v \in \mathbb{P}^{2n}$  which every embedded tangent space to the quadric contains (Remark 2.8). In general, a projective variety  $X \subset \mathbb{P}^N$  is said to be *strange* if its embedded tangent spaces admit such a common point  $v \in \mathbb{P}^N$  (see §2.1, for more details).

It is classically known that a non-linear smooth projective curve  $X \subset \mathbb{P}^N$  is strange if and only if  $X$  is a conic in  $\mathbb{P}^N$  in  $p = 2$  (E. Lluís [6], P. Samuel [11]; see also [3, IV, Theorem 3.9]). In higher-dimensions, S. Kleiman and R. Piene [5, Theorem 7] focused on hypersurfaces, and showed that a non-linear smooth

---

*Date:* February 2, 2014.

*2000 Mathematics Subject Classification.* Primary 14F10; Secondary 14N05.

*Key words and phrases.* strange variety, characterization of hyperquadrics.

hypersurface  $X \subset \mathbb{P}^N$  is strange if and only if  $X$  is a quadric in  $\mathbb{P}^N$  of odd dimension in  $p = 2$ .

In this paper, we investigate whether a smooth complete intersection variety other than quadrics can be strange, and answer it negatively by examining a parameter space of strange complete intersections. In consequence, we have:

**Theorem 1.1.** *Let  $X$  be a smooth projective variety which is a non-linear complete intersection in  $\mathbb{P}^N$ , and let  $\mathcal{O}_X(1) := \mathcal{O}_{\mathbb{P}^N}(1)|_X$ . Then the following are equivalent:*

- (a)  $X$  is strange in  $\mathbb{P}^N$ ,
- (b)  $X$  is a quadric in  $\mathbb{P}^N$  of odd dimension in  $p = 2$ .

Moreover, in the case where  $\dim(X) \geq 2$ , the conditions (a) and (b) are equivalent to

- (c)  $H^0(X, T_X(-1)) \neq 0$ .

This paper is organized as follows: In §2.1, we study some basic properties of strangeness. In §2.2, considering relation between strangeness and cohomology, we show the equivalence “(a)  $\Leftrightarrow$  (c)” (for the exceptional case  $\dim(X) = 1$ , see Remark 2.12). In §2.3, we analyze defining polynomials of strange varieties. We say that an  $(N - r)$ -dimensional projective variety in  $\mathbb{P}^N$  is an  $(e^1, \dots, e^r)$ -complete intersection if it is scheme-theoretically equal to an intersection of  $r$  hypersurfaces of degrees  $e^1, \dots, e^r$ , where  $e^1, \dots, e^r$  are  $r$  positive integers. In (3) of §3, we define an irreducible parameter space  $\mathcal{H}^v$  of  $(e^1, \dots, e^r)$ -complete intersections being strange for  $v$ . In order to show the implication “(a)  $\Rightarrow$  (b)”, it is essential to consider the case where all  $e^k > 1$  and where  $\mathcal{H}^v$  is *not* equal to the parameter space of quadrics of odd dimensions in  $p = 2$ . In §3.1, we construct an  $(e^1, \dots, e^r)$ -complete intersection variety  $X_0 \subset \mathbb{P}^N$  having an isolated singular point  $\alpha \neq v$ . In §3.2, we take the incidence scheme  $I \subset \mathcal{H}^v \times \mathbb{P}^N$  parameterizing pairs of strange complete intersections and their singular points, and in addition, take an irreducible component  $\Lambda \subset I$  whose subset parameterizes the orbit of the pair  $(X_0, \alpha)$  under automorphisms of  $\mathbb{P}^N$  with fixed point  $v$ . Calculating the dimension of  $\Lambda$  and using the existence of  $(X_0, \alpha)$ , we show that the projection  $\Lambda \rightarrow \mathcal{H}^v$  is surjective in the case. This means that every  $(e^1, \dots, e^r)$ -complete intersection variety belonging to  $\mathcal{H}^v$  is singular (Theorem 3.8), yielding Theorem 1.1.

## 2. COHOMOLOGY AND DEFINING POLYNOMIALS

**2.1. Preliminary.** Let  $X \subset \mathbb{P}^N$  be a projective variety over an algebraically closed field  $K$  of characteristic  $p \geq 0$ . We say that  $X$  is *strange for a point*  $v \in \mathbb{P}^N$  if  $v \in \mathbb{T}_x X$  for any smooth point  $x \in X$ , where  $\mathbb{T}_x X \subset \mathbb{P}^N$  is the embedded tangent space to  $X$  at  $x$ . We simply say that  $X$  is *strange* in  $\mathbb{P}^N$  if  $X$  is strange for some point of  $\mathbb{P}^N$ .

Let  $(z_0 : z_1 : \dots : z_N)$  be the homogeneous coordinates on  $\mathbb{P}^N$ . We denote by  $f_{z_j} := \partial f / \partial z_j$  for a homogeneous polynomial  $f \in K[z_0, z_1, \dots, z_N]$ .

**Proposition 2.1.** *Let  $X \subset \mathbb{P}^N$  be a hypersurface defined by a homogeneous polynomial  $f$ , and let  $v = (1 : 0 : \cdots : 0) \in \mathbb{P}^N$ . Then  $X$  is strange for  $v$  if and only if  $f_{z_0}$  is the zero polynomial.*

*Proof.* Let  $a \in X$  be a smooth point, where recall that  $\mathbb{T}_a X = (f_{z_0}(a)z_0 + \cdots + f_{z_N}(a)z_N = 0) \subset \mathbb{P}^N$ . Then  $v \in \mathbb{T}_a X$  if and only if  $f_{z_0}(a) = 0$ . Hence  $X$  is strange for  $v$  if and only if  $f_{z_0}|_X = 0$ . Here, the latter condition  $f_{z_0}|_X = 0$  means that  $f_{z_0}$  is contained in the ideal  $(f) \subset K[z_0, \dots, z_N]$ , and then we have  $f_{z_0} = 0$  because of  $\deg(f_{z_0}) < \deg(f)$ .  $\square$

Note that, in the case where  $X \subset \mathbb{P}^N$  is a degenerate subvariety, i.e.,  $X$  is contained in an  $m$ -dimensional linear subvariety  $L$  of  $\mathbb{P}^N$  with  $m < N$ ,  $X$  is strange in  $\mathbb{P}^N$  if and only if  $X$  is strange in  $L \simeq \mathbb{P}^m$ .

**Example 2.2.** Let  $X \subset \mathbb{P}^N$  be a smooth quadric, i.e., a smooth projective variety of degree 2 in  $\mathbb{P}^N$ . Then  $X$  is strange if and only if  $\dim(X)$  is odd and  $p = 2$ . The reason is as follows: It is sufficient to consider the case where  $X$  is non-degenerate; thus we set  $N = \dim(X) + 1$ . Let  $f$  be the defining equation of  $X$ . Choosing suitable coordinates  $(z_0 : z_1 : \cdots : z_N)$  on  $\mathbb{P}^N$ , we can assume that

$$f = \begin{cases} z_0^2 + z_1 z_2 + z_3 z_4 + \cdots + z_{N-1} z_N & \text{if } \dim(X) \text{ is odd,} \\ z_0 z_1 + z_2 z_3 + \cdots + z_{N-1} z_N & \text{if } \dim(X) \text{ is even.} \end{cases}$$

(a) Assume that  $\dim(X)$  is odd and  $p = 2$ . Then  $f_{z_0} = 2z_0 = 0$ ; hence it follows from Proposition 2.1 that  $X$  is strange for  $v = (1 : 0 : \cdots : 0)$ .

(b) Assume that  $\dim(X)$  is even or  $p \neq 2$ . Let us consider the Gauss map  $\gamma : X \rightarrow (\mathbb{P}^N)^\vee$  sending  $a \mapsto \mathbb{T}_a X$ , where  $(\mathbb{P}^N)^\vee = \mathbb{G}(N-1, \mathbb{P}^N)$  is the space of hyperplanes of  $\mathbb{P}^N$ . Indeed, we can describe  $\gamma$  by

$$a \mapsto (f_{z_0}(a) : \cdots : f_{z_N}(a)).$$

Then, by assumption,  $\gamma$  is isomorphic with  $\gamma(\mathcal{O}(1)) = \mathcal{O}(1)$ , and  $\gamma(X)$  is also a smooth quadric hypersurface in  $(\mathbb{P}^N)^\vee$ . We denote by  $u^* \subset (\mathbb{P}^N)^\vee$  the set of hyperplanes containing a point  $u \in \mathbb{P}^N$ . Then  $u^*$  is a hyperplane of  $(\mathbb{P}^N)^\vee$ . If  $X$  is strange for some  $u \in \mathbb{P}^N$ , then we have  $\gamma(X) \subset u^*$ , a contradiction. Hence  $X$  is not strange.

We say that a projective variety  $X \subset \mathbb{P}^N$  is a *cone* with vertex  $v \in \mathbb{P}^N$  if the line  $\overline{xv}$  is contained in  $X$  for any  $x \in X$ . If  $X$  is a cone with vertex  $v$ , then  $X$  is strange for  $v$ .

**Remark 2.3.** In  $p = 0$ , if  $X$  is strange for  $v$ , then  $X$  is a cone with vertex  $v$ . The reason is as follows: Let  $\pi_v : \mathbb{P}^N \setminus \{v\} \rightarrow \mathbb{P}^{N-1}$  be the linear projection from  $v$ , and let  $d_x \pi_v : T_x \mathbb{P}^N \rightarrow T_{\pi_v(x)} \mathbb{P}^{N-1}$  be the linear map between Zariski tangent spaces. For any smooth point  $x \in X$  with  $x \neq v$ , since  $T_x \overline{xv} \subset T_x X \cap \ker(d_x \pi_v)$ , we have  $\dim(d_x \pi_v(T_x X)) = \dim(X) - 1$ . It follows from  $p = 0$  that  $\dim(\pi_v(X \setminus \{v\})) = \dim(X) - 1$ ; hence  $X$  is a cone with vertex  $v$ .

**Remark 2.4.** Let  $X \subset \mathbb{P}^N$  be a projective variety being strange for a point  $v$ . Then we immediately have the following properties:

(a) If  $X$  is smooth and  $L \subset \mathbb{P}^N$  is a hyperplane not containing  $v$ , then  $X \cap L$  is smooth. This is because, for each point  $x \in X \cap L$ , it follows from  $v \in \mathbb{T}_x X$  that  $\mathbb{T}_x X \not\subset L$ ; hence  $X \cap L$  is smooth at  $x$  and  $\mathbb{T}_x(X \cap L) = \mathbb{T}_x(X) \cap L$ .

(b) Let  $\pi_z : \mathbb{P}^N \setminus \{z\} \rightarrow \mathbb{P}^{N-1}$  be a linear projection from a point  $z \in \mathbb{P}^N$  with  $z \neq v$ . Then the image  $Y \subset \mathbb{P}^{N-1}$  of  $X$  under  $\pi_z$  is strange for  $\pi_z(v)$ . This is because,  $\mathbb{T}_{\pi_z(x)} Y$  contains  $\pi_z(\mathbb{T}_x X)$  for a general point  $x \in X$ .

**2.2. Section of 0-cohomology of  $(-1)$ -twist of a tangent bundle.** Let  $X \subset \mathbb{P}^N$  be a smooth quasi-projective variety, and let  $\mathcal{O}_X(1) := \mathcal{O}_{\mathbb{P}^N}(1)|_X$ . We identify  $\mathbb{P}^N$  with  $(H^0(\mathbb{P}^N, \mathcal{O}(1))^\vee \setminus 0)/(K \setminus 0)$ , the projectivization of the dual vector space of  $H^0(\mathbb{P}^N, \mathcal{O}(1))$ . Let

$$\hat{v} \subset H^0(\mathbb{P}^N, \mathcal{O}(1))^\vee$$

be the one-dimensional vector subspace corresponding to  $v \in \mathbb{P}^N$ . Considering the Euler sequence  $0 \rightarrow \mathcal{O}_{\mathbb{P}^N} \rightarrow H^0(\mathbb{P}^N, \mathcal{O}(1))^\vee \otimes \mathcal{O}_{\mathbb{P}^N}(1) \xrightarrow{\xi} T_{\mathbb{P}^N} \rightarrow 0$ , we can define a composite homomorphism  $s_v$  of bundles on  $\mathbb{P}^N$  by

$$s_v : \hat{v} \otimes \mathcal{O}_{\mathbb{P}^N}(1) \hookrightarrow H^0(\mathbb{P}^N, \mathcal{O}(1))^\vee \otimes \mathcal{O}_{\mathbb{P}^N}(1) \xrightarrow{\xi} T_{\mathbb{P}^N}.$$

**Proposition 2.5.** *Let  $X \subset \mathbb{P}^N$  be a smooth quasi-projective variety. Then  $X$  is strange for  $v$  if and only if  $s_v|_X$  factors through  $T_X \subset T_{\mathbb{P}^N}|_X$ . Hence, in this case,  $s_v|_X$  gives a nonzero section of  $H^0(X, T_X(-1)) \simeq \text{Hom}_{\mathcal{O}_X}(\mathcal{O}_X(1), T_X)$ .*

To show this, we consider  $\mathcal{P}_X^1 = \mathcal{P}_X^1(\mathcal{O}_X(1))$ , the bundle of principal parts of  $\mathcal{O}_X(1)$  of first order, which gives an exact sequence

$$0 \rightarrow \Omega_X^1 \rightarrow \mathcal{P}_X^1 \otimes \mathcal{O}_X(-1) \rightarrow \mathcal{O}_X \rightarrow 0$$

(see [10, Remark (6.4)]). Taking the dual of this, we have the following commutative diagram with exact rows and columns:

$$(1) \quad \begin{array}{ccccccc} & & & 0 & & 0 & \\ & & & \downarrow & & \downarrow & \\ 0 & \longrightarrow & \mathcal{O}_X & \longrightarrow & \mathcal{P}_X^1 \otimes \mathcal{O}_X(1) & \longrightarrow & T_X \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ & & \mathcal{O}_X & \longrightarrow & H^0(\mathbb{P}^N, \mathcal{O}(1))^\vee \otimes \mathcal{O}_X(1) & \xrightarrow{\xi|_X} & T_{\mathbb{P}^N}|_X \longrightarrow 0 \\ & & & & \downarrow & & \downarrow \\ & & & & N_{X/\mathbb{P}^N} & \xlongequal{\quad\quad\quad} & N_{X/\mathbb{P}^N} \\ & & & & \downarrow & & \downarrow \\ & & & & 0 & & 0. \end{array}$$

**Remark 2.6.** For a smooth point  $x \in X$ , the projectivization of  $\mathcal{P}_X^{1\vee} \otimes k(x) \subset H^0(\mathbb{P}^N, \mathcal{O}(1))^\vee$  corresponds to the embedded tangent space  $\mathbb{T}_x X \subset \mathbb{P}^N$ . The reason is as follows. Let  $r = \text{codim}(X, \mathbb{P}^N)$ , and let  $f^1, \dots, f^r$  be  $r$  homogeneous polynomials which locally define  $X$  around the smooth point  $x$ . Then the linear map  $H^0(\mathbb{P}^N, \mathcal{O}(1))^\vee \rightarrow N_{X/\mathbb{P}^N} \otimes k(x)$  induced from the middle column of the above diagram (1) is represented by the matrix  $[f_{z_i}^k(x)]_{1 \leq k \leq r, 0 \leq i \leq N}$ , and hence its kernel is the zero set of  $r$  linear polynomials  $\sum_i f_{z_i}^1(x) \cdot z_i, \dots, \sum_i f_{z_i}^r(x) \cdot z_i$ . This implies the assertion.

From Remark 2.6, we find that  $\hat{v} \subset \mathcal{P}_X^{1\vee} \otimes k(x)$  if and only if  $v \in \mathbb{T}_x X$ . In particular, the following holds:

**Lemma 2.7.** *The subbundle  $\hat{v} \otimes \mathcal{O}_X \subset H^0(\mathbb{P}^N, \mathcal{O}(1)) \otimes \mathcal{O}_X$  is contained in  $\mathcal{P}_X^{1\vee}$  if and only if  $X$  is strange for  $v$ .*

*Proof of Proposition 2.5.* If  $X$  is strange for  $v$ , then it follows from Lemma 2.7 that  $s_v|_X$  is equal to the composite map,  $\hat{v} \otimes \mathcal{O}_X(1) \hookrightarrow \mathcal{P}_X^{1\vee} \otimes \mathcal{O}_X(1) \rightarrow T_X$ . If  $X$  is not strange for  $v$ , then  $\hat{v} \not\subset \mathcal{P}_X^{1\vee} \otimes k(x)$  for some  $x$ , and then the image of  $s_v(x) : \hat{v} \rightarrow T_x \mathbb{P}^N$  is not contained in  $T_x X$ .  $\square$

**Remark 2.8.** If a smooth projective variety  $X$  is strange and is not isomorphic to a projective space, then it follows from Proposition 2.5 that  $X$  gives a counter-example in  $p > 0$  of the statement of Wahl's cohomological characterization of projective spaces. For example, smooth quadrics in  $p = 2$  whose dimensions are odd and  $\geq 3$  (Example 2.2). (One dimensional quadrics, i.e., conics, are still isomorphic to  $\mathbb{P}^1$ .)

In addition, we can restate strangeness of tangents as a cohomological condition, as follows:

**Corollary 2.9.** *Let  $X \subset \mathbb{P}^N$  be a smooth projective variety with  $\mathcal{O}_X(1) := \mathcal{O}_{\mathbb{P}^N}(1)|_X$ , and denote by  $\mathcal{P}_X^1 = \mathcal{P}_X^1(\mathcal{O}_X(1))$  the bundle of principal parts of  $\mathcal{O}_X(1)$  of first order. Then  $X$  is strange in  $\mathbb{P}^N$  if and only if  $H^0(X, \mathcal{P}_X^{1\vee}) \neq 0$ .*

*Proof.* The ‘‘only if’’ part follows immediately from Lemma 2.7. To show the ‘‘if’’ part, we consider the case where there exists a nonzero section  $s \in H^0(X, \mathcal{P}_X^{1\vee})$ . By the middle column of (1),  $s$  is regarded as a nonzero section of  $H^0(\mathbb{P}^N, \mathcal{O}(1))^\vee$ , which gives a point  $v \in \mathbb{P}^N$  such that  $K \cdot s = \hat{v}$ . Then  $X$  is strange for  $v$ .  $\square$

Now, let us consider the case where  $X$  satisfies  $H^1(X, \mathcal{O}_X(-1)) = 0$ .

**Proposition 2.10.** *Let  $X \subset \mathbb{P}^N$  be a smooth projective variety, and assume that  $H^1(X, \mathcal{O}_X(-1)) = 0$ . Then  $X$  is strange in  $\mathbb{P}^N$  if and only if  $H^0(X, T_X(-1)) \neq 0$ .*

*Proof.* The ‘‘only if’’ part was shown in Proposition 2.5. Let us consider the ‘‘if’’ part. By the first row of (1), it follows from  $H^1(X, \mathcal{O}_X(-1)) = 0$  that

$$H^0(X, \mathcal{P}_X^{1\vee}) \rightarrow H^0(X, T_X(-1))$$

is surjective. Hence the assertion follows from Corollary 2.9.  $\square$

**Lemma 2.11.** *Let  $X = X^1 \cap \cdots \cap X^r \subset \mathbb{P}^N$  be a complete intersection of  $r$  hypersurfaces  $X^1, \dots, X^r$  of degrees  $e^1, \dots, e^r$ . Then  $H^j(X, \mathcal{O}_X(i)) = 0$  for  $0 < j < N - r$  and all  $i \in \mathbb{Z}$ .*

*Proof.* Since  $H^j(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(i)) = 0$  for  $0 < j < N$  and all  $i$ , it follows from the exact sequence  $0 \rightarrow \mathcal{O}_{\mathbb{P}^N}(-e^1) \rightarrow \mathcal{O}_{\mathbb{P}^N} \rightarrow \mathcal{O}_{X^1} \rightarrow 0$  that  $H^j(X^1, \mathcal{O}_{X^1}(i)) = 0$  for  $0 < j < N - 1$  and all  $i$ . Similarly, it follows from the exact sequence  $0 \rightarrow \mathcal{O}_{X^1}(-e^2) \rightarrow \mathcal{O}_{X^1} \rightarrow \mathcal{O}_{X^1 \cap X^2} \rightarrow 0$  that  $H^j(X^1 \cap X^2, \mathcal{O}_{X^1 \cap X^2}(i)) = 0$  for  $0 < j < N - 2$  and all  $i$ . Inductively, we have  $H^j(X, \mathcal{O}_X(-i)) = 0$  for  $0 < j < N - r$  and all  $i$ .  $\square$

**Remark 2.12.** Let  $X \subset \mathbb{P}^N$  be a smooth complete intersection of dimension  $\geq 2$ . Then  $X$  satisfies the statement of Proposition 2.10, because of Lemma 2.11. Note that the case where  $\dim(X) = 1$  is exceptional: Assume that  $X$  is a smooth  $(e^1, \dots, e^{N-1})$ -complete intersection curve in  $\mathbb{P}^N$ . Then  $T_X = \mathcal{O}_X(N + 1 - \sum_{1 \leq k \leq N-1} e^k)$ , which implies that  $H^0(X, T_X(-1)) \neq 0$  if and only if  $X$  is a conic or line (in any  $p \geq 0$ ). We recall that a smooth conic in  $p \neq 2$  is not strange.

**2.3. Defining polynomials of a strange complete intersection variety.** Let  $e^1, \dots, e^r$  be  $r$  integers greater than 1. We recall that an  $(N - r)$ -dimensional projective variety  $X \subset \mathbb{P}^N$  is an  $(e^1, \dots, e^r)$ -complete intersection if  $X$  is scheme-theoretically equal to an intersection of  $r$  hypersurfaces of degrees  $e^1, \dots, e^r$ , i.e., a minimal set of generators of the defining homogeneous ideal  $I_X \subset K[z_0, z_1, \dots, z_N]$  of  $X$  consists of  $r$  homogeneous polynomials of degrees  $e^1, \dots, e^r$ . We generalize Proposition 2.1, as follows:

**Proposition 2.13.** *Let  $X \subset \mathbb{P}^N$  be an  $(e^1, \dots, e^r)$ -complete intersection variety which is strange for a point  $v = (1 : 0 : \cdots : 0)$ . Then  $I_X$  is generated by  $r$  homogeneous polynomials  $f^1, \dots, f^r$  of degrees  $e^1, \dots, e^r$  such that  $f_{z_0}^k$  is the zero polynomial for  $1 \leq k \leq r$ .*

To prove the above statement, we first show the following lemma:

**Lemma 2.14.** *Let  $X \subset \mathbb{P}^N$  be a projective variety which is strange for  $v = (1 : 0 : \cdots : 0)$ . For a homogeneous polynomial  $g \in I_X$  of degree  $e$ , there exists a homogeneous polynomial  $\tilde{g} \in I_X$  of degree  $e$ , such that  $\tilde{g}_{z_0} = 0$  and the two ideals  $(g, z_0), (\tilde{g}, z_0) \subset K[z_0, z_1, \dots, z_N]$  coincide.*

*Proof.* For a homogeneous polynomial  $g \in I_X$ , we have  $g_{z_0} \in I_X$ . The reason is as follows: There is nothing to prove if  $g$  is the zero polynomial. Let  $g$  be nonzero, and denote by  $Y := (g = 0) \subset \mathbb{P}^N$ , the hypersurface defined by  $g$ . For any  $a \in X$ , it follows that  $v \in \mathbb{T}_a X \subset \mathbb{T}_a Y$ ; then, since  $v = (1 : 0 : \cdots : 0)$  and  $\mathbb{T}_a Y = (g_{z_0}(a)z_0 + \cdots + g_{z_N}(a)z_N = 0)$ , we have  $g_{z_0}(a) = 0$ . Hence  $g_{z_0} \in I_X$ .

Applying the above argument inductively, we have  $\partial^j g / \partial z_0^j \in I_X$  for any  $j > 0$ . Now let

$$\tilde{g} := g + \sum_{j=1}^{p-1} \frac{(-1)^j z_0^j}{j!} \cdot \frac{\partial^j g}{\partial z_0^j},$$

which is contained in  $I_X$  and satisfies that  $(g, z_0) = (\tilde{g}, z_0)$ . In addition, we have  $\tilde{g}_{z_0} = 0$ , because of

$$\begin{aligned} \frac{\partial}{\partial z_0} \left( \sum_{j=1}^{p-1} \frac{(-1)^j z_0^j}{j!} \cdot \frac{\partial^j g}{\partial z_0^j} \right) &= \sum_{j=1}^{p-1} \frac{(-1)^j z_0^{j-1}}{(j-1)!} \cdot \frac{\partial^j g}{\partial z_0^j} + \sum_{j=1}^{p-1} \frac{(-1)^j z_0^j}{j!} \cdot \frac{\partial^{j+1} g}{\partial z_0^{j+1}} \\ &= -\frac{\partial g}{\partial z_0} + \frac{(-1)^{p-1} z_0^{p-1}}{(p-1)!} \cdot \frac{\partial^p g}{\partial z_0^p} = -\frac{\partial g}{\partial z_0}, \end{aligned}$$

where  $\partial^p g / \partial z_0^p = 0$  since  $p$  is the characteristic of the ground field.  $\square$

**Remark 2.15.** The operation making  $\tilde{g}$  with  $\tilde{g}_{z_0} = 0$  has appeared in an algorithm of derivation kernel computation (see [2, p. 27]; for positive characteristic, see [9]).

*Proof of Proposition 2.13.* We can assume  $N - r \geq 1$ . Let  $I_X$  be generated by homogeneous polynomials  $f^1, \dots, f^r$  of degrees  $e^1, \dots, e^r$ . From Lemma 2.14, for each  $1 \leq k \leq r$ , we have a homogeneous polynomial  $\tilde{f}^k$  of degree  $e^k$  satisfying that  $\tilde{f}^k \in I_X$ ,  $\tilde{f}_{z_0}^k = 0$ , and  $(f^k, z_0) = (\tilde{f}^k, z_0)$ . We set  $\tilde{X} \subset \mathbb{P}^N$  to be the zero set of  $\tilde{f}^1, \dots, \tilde{f}^r$ . Then we have  $X \subset \tilde{X}$  and  $X \cap (z_0 = 0) = \tilde{X} \cap (z_0 = 0)$  for the hyperplane  $(z_0 = 0) \subset \mathbb{P}^N$ .

Let  $V_1, \dots, V_m$  be the irreducible components of  $\tilde{X}$ , where  $V_i$  is of dimension  $\geq N - r \geq 1$  for each  $1 \leq i \leq m$ . Since  $X$  is strange for  $v$  and  $v \notin (z_0 = 0)$ , we have  $X \not\subset (z_0 = 0)$ ; in particular,  $\dim(X \cap (z_0 = 0)) = N - r - 1$ . Since  $X \cap (z_0 = 0) = \bigcup_i (V_i \cap (z_0 = 0))$ , we have  $\dim(V_i \cap (z_0 = 0)) = N - r - 1$ , and hence  $\dim(V_i) = N - r$ . In particular,  $X$  coincides with some  $V_i$ . Since  $\deg(X) = \deg(\tilde{X})$ , we have  $X = \tilde{X}$ .  $\square$

**Corollary 2.16.** *Let  $X \subset \mathbb{P}^N$  be an  $(e^1, \dots, e^r)$ -complete intersection variety. Assume that  $X$  is strange for a point  $v$  and assume that  $e^k < p$  for any  $k$  with  $1 \leq k \leq r$ . Then  $X$  is a cone with vertex  $v$ .*

*Proof.* Changing coordinates, we can assume that  $v = (1 : 0 : \dots : 0) \in \mathbb{P}^N$ . Then it follows from Proposition 2.13 that  $X$  is defined by homogeneous polynomials  $f^1, \dots, f^r$  of degrees  $e^1, \dots, e^r$  such that  $f_{z_0}^k = 0$  for  $1 \leq k \leq r$ . If  $f^k \notin K[z_1, \dots, z_N]$ , then the inequality  $e^k < p$  implies  $f_{z_0}^k \neq 0$ , a contradiction. Hence  $f^k \in K[z_1, \dots, z_N]$  for any  $k$ , which means that  $X$  is a cone with vertex  $v$ .  $\square$

### 3. PARAMETER SPACE OF STRANGE COMPLETE INTERSECTIONS

We fix some notation. Let  $\mathcal{H}_e = |\mathcal{O}_{\mathbb{P}^N}(e)|$  be the projectivization of  $H^0(\mathbb{P}^N, \mathcal{O}(e))$ , whose general members parameterize hypersurfaces of  $\mathbb{P}^N$  of degree  $e$ . For a

point  $v \in \mathbb{P}^N$ , we denote by  $\mathcal{H}_e^v$  the subset of  $\mathcal{H}_e$  whose general members parameterize hypersurfaces being strange for  $v$ . Changing homogeneous coordinates  $(z_0 : z_1 : \cdots : z_N)$  on  $\mathbb{P}^N$ , we can assume

$$v = (1 : 0 : \cdots : 0).$$

Then we have  $\mathcal{H}_e^v = \{f \in \mathcal{H}_e \mid f_{z_0} = 0\}$  due to Proposition 2.1. In particular,  $\mathcal{H}_e^v$  is regarded as a linear subvariety of  $\mathcal{H}_e$ , since it is equal to the projectivization of the kernel of the linear map  $H^0(\mathbb{P}^N, \mathcal{O}(e)) \rightarrow H^0(\mathbb{P}^N, \mathcal{O}(e-1)) : f \mapsto f_{z_0}$ .

Let  $e^1, \dots, e^r$  be  $r$  integers greater than 1. We denote by

$$(2) \quad \mathcal{H} := \mathcal{H}_{e^1} \times \cdots \times \mathcal{H}_{e^r},$$

whose general members parameterize  $(N-r)$ -dimensional  $(e^1, \dots, e^r)$ -complete intersections in  $\mathbb{P}^N$ . Let us consider

$$(3) \quad \mathcal{H}^v := \mathcal{H}_{e^1}^v \times \cdots \times \mathcal{H}_{e^r}^v,$$

whose general members parameterize complete intersections being strange for  $v$ . Indeed, we have:

$$(4) \quad \mathcal{H}^v = \{(f^1, \dots, f^r) \in \mathcal{H} \mid f_{z_0}^1 = \cdots = f_{z_0}^r = 0\}.$$

Note that  $\mathcal{H}^v$  is irreducible, since so is each  $\mathcal{H}_e^v$ .

**Remark 3.1.** Let  $X \subset \mathbb{P}^N$  be an  $(e^1, \dots, e^r)$ -complete intersection variety being strange for  $v$ . Then, from Proposition 2.13, we can find homogeneous polynomials  $(f^1, \dots, f^r) \in \mathcal{H}^v$  whose zero set is equal to  $X$ .

**3.1. Strange varieties with isolated singular points.** We construct a member of  $\mathcal{H}^v$  which defines a complete intersection variety  $X_0$  having an isolated singular point  $\alpha \neq v$ . Later, the pair  $(X_0, \alpha)$  will play an essential role.

For a member  $(f^1, \dots, f^r) \in \mathcal{H}^v$  and for  $a \in \mathbb{P}^N$ , we set

$$(5) \quad D((f^k), a) = D((f^k), a; \mathbb{P}^N) := \begin{bmatrix} f_{z_1}^1(a) & \cdots & f_{z_N}^1(a) \\ \vdots & & \vdots \\ f_{z_1}^r(a) & \cdots & f_{z_N}^r(a) \end{bmatrix},$$

where we need not consider  $f_{z_0}^k$ 's since these are zero as in (4). The complete intersection variety  $X \subset \mathbb{P}^N$  defined by  $(f^1, \dots, f^r)$  is singular at  $a \in \mathbb{P}^N$  if and only if  $f^1(a) = \cdots = f^r(a) = 0$  and  $\text{rk } D((f^k), a) < r$ .

**Proposition 3.2.** *Assume that  $\mathcal{H}$  is not equal to the parameter space of quadric hypersurfaces in  $\mathbb{P}^N$  of odd dimensions in  $p = 2$ . Assume that  $p > 0$  and  $e^k \geq p$  for some  $k$ . Then there exists a complete intersection variety  $X_0 \subset \mathbb{P}^N$  defined by a member of  $\mathcal{H}^v$  such that  $\text{Sing } X_0 \neq \{v\}$  and  $0 < \#(\text{Sing } X_0) < \infty$ . In particular, there exists an isolated singular point  $\alpha$  of  $X_0$  such that  $\alpha \neq v$ .*

**Remark 3.3** (Hypersurfaces). Let us consider the case where  $r = 1$ , i.e.,  $\mathcal{H}$  is equal to  $\mathcal{H}_e$ , the parameter space of hypersurfaces in  $\mathbb{P}^N$  of degree  $e$ .

(a) Suppose that  $\mathcal{H} = \mathcal{H}_2$  in  $p = 2$ , and suppose that  $N$  is even (i.e., hypersurfaces are of odd dimensions). Then every member of  $\mathcal{H}_2^v$  does *not* satisfy the statement “ $\text{Sing } X_0 \neq \{v\}$  and  $0 < \#(\text{Sing } X_0) < \infty$ ”. This is because, if a quadric  $X \subset \mathbb{P}^N$  is defined by a member of  $\mathcal{H}^v$  and is singular at a point  $\alpha \neq v$ , then we have  $\dim(\text{Sing } X) \geq 1$ , as follows: Let  $\pi : \mathbb{P}^N \setminus \{\alpha\} \rightarrow \mathbb{P}^{N-1}$  be the linear projection from  $\alpha$ . Then  $Y = \pi(X) \subset \mathbb{P}^{N-1}$  is an  $(N-2)$ -dimensional quadric, and  $X = \overline{\pi^{-1}(Y)}$  (it is a cone with vertex  $\alpha$ ). Since  $X$  is strange for  $v$ ,  $Y$  is strange for  $\pi(v)$  as in Remark 2.4(b). From Example 2.2,  $Y$  is not smooth. Hence, for  $w \in \text{Sing}(Y)$ , the line  $\overline{\pi^{-1}(w)}$  is contained in  $\text{Sing}(X)$ .

(b) Suppose that  $\mathcal{H} = \mathcal{H}_2$  in  $p = 2$ , and suppose that  $N$  is odd. Then  $\mathcal{H}_2^v$  satisfies the statement, as follows: We take  $L \subset \mathbb{P}^N$  to be a hyperplane containing  $v$ , and take  $Y$  to be an  $(N-2)$ -dimensional smooth quadric in  $L$  which is strange for  $v$ . For a point  $\alpha \notin L$ , we set  $X_0 = \text{Cone}_\alpha(Y)$ , the cone over  $Y$  with vertex  $\alpha$ . Then it follows that  $\text{Sing}(X_0) = \{\alpha\}$  with  $\alpha \neq v$ .

(c) For  $e \geq 3$  and  $e \geq p > 0$ , we can construct a member of  $\mathcal{H}_e^v$  satisfying the statement of Proposition 3.2, as follows. Suppose that  $p \mid e$ . Then we set  $X_0 \subset \mathbb{P}^N$  to be the hypersurface defined by

$$f = z_N z_{N-1}^{e-1} + z_{N-1} z_{N-2}^{e-1} + \cdots + z_2 z_1^{e-1} + z_0^e.$$

Then  $X_0$  is strange for  $v$  because of  $f_{z_0} = 0$ . On the other hand,  $D(f, z) = [f_{z_1}(z) \ \cdots \ f_{z_N}(z)]$  is equal to

$$[(e-1)z_1^{e-2}z_2 \quad z_1^{e-1} + (e-1)z_2^{e-2}z_3 \quad \cdots \quad z_{N-2}^{e-1} + (e-1)z_N z_{N-1}^{e-2} \quad z_{N-1}^{e-1}].$$

Then  $\text{Sing}(X_0) = \{(0 : \cdots : 0 : 1)\}$  holds, as follows: We have “ $\supset$ ” immediately. To show “ $\subset$ ”, we take  $z \in \text{Sing}(X_0)$ . Since  $D(f, z) = 0$ , by the first polynomial from the right of the above description of  $D(f, z)$ , we have  $z_{N-1} = 0$ . By the second polynomial from the right, we have  $z_{N-2} = 0$  because of  $e-2 \geq 1$ . Similarly, we have  $z_1 = \cdots = z_{N-1} = 0$ . Since  $f(z) = 0$ , we have  $z_0 = 0$ . Therefore  $z = (0 : \cdots : 0 : 1)$ .

Suppose that  $e > p$  and  $p \nmid e$ . Then we set  $X_0 \subset \mathbb{P}^N$  to be the hypersurface defined by

$$f = z_0^p z_1^{e-p} + z_2^e + \cdots + z_N^e.$$

Then  $X_0$  is strange for  $v$ . In addition,  $D(f, z)$  is equal to

$$[(e-p)z_0^p z_1^{e-p-1} \quad e z_2^{e-1} \quad \cdots \quad e z_N^{e-1}].$$

Thus  $\text{Sing}(X_0) = \{(1 : 0 : 0 : \cdots : 0), (0 : 1 : 0 : \cdots : 0)\}$  if  $e > p + 1$ , and  $\text{Sing}(X_0) = \{(0 : 1 : 0 : \cdots : 0)\}$  if  $e = p + 1$ .

*Proof of Proposition 3.2.* We have already shown the assertion in the case where  $\mathcal{H}$  is the space of hypersurfaces, as above. Thus we assume that  $r \geq 2$ . Without loss of generality, we can assume that  $e_1 \geq p$ . Let us take  $Z \subset \mathbb{P}^{N-1}$  to be an  $(N-r)$ -dimensional complete intersection variety defined by  $r-1$

homogeneous polynomials  $f^2, \dots, f^r \in K[z_1, \dots, z_N]$  of degrees  $e^2, \dots, e^r$ , such that  $0 < \#(\text{Sing}(Z)) < \infty$  (for example, a cone over a smooth variety). By Bertini's theorem, we can choose a hypersurface  $Y^1 \subset \mathbb{P}^{N-1}$  defined by a homogeneous polynomial  $f^1 \in K[z_1, \dots, z_N]$  of degree  $e^1$  such that the intersection  $Y := Y^1 \cap Z \subset \mathbb{P}^{N-1}$  is smooth, where  $Y^1 \cap \text{Sing}(Z) = \emptyset$ . Note that Bertini's theorem for subvarieties of  $\mathbb{P}^N$  in any  $p \geq 0$  is deduced from [3, II, Theorem 8.18] by using  $d$ -uple embeddings.

Let  $\beta = (\beta_1 : \dots : \beta_N) \in \text{Sing}(Z)$ . Then we have

$$f^1(\beta) \neq 0 \text{ and } \text{rk } D((f^2, \dots, f^r), \beta; \mathbb{P}^{N-1}) < r - 1.$$

Here, without loss of generality, we can assume that  $\beta_1 \neq 0$ . We regard  $Y$  as a subvariety of  $\mathbb{P}^N$  contained in the hyperplane  $(z_0 = 0) \simeq \mathbb{P}^{N-1}$ .

Now, let us define a complete intersection variety

$$X_0 := (g^1 = f^2 = \dots = f^r = 0) \subset \mathbb{P}^N,$$

where

$$g^1 := f^1 - \frac{f^1(\beta)}{\beta_1^{e^1-p}} \cdot z_0^p z_1^{e^1-p} \in K[z_0, z_1, \dots, z_N].$$

Then  $X_0$  is strange for  $v$  because of  $g^1_{z_0} = 0$ . In addition, we have  $\dim(\text{Sing}(X_0)) < 1$ , as follows: By definition of  $g^1$ , the intersection  $X_0 \cap (z_0 = 0)$  is equal to the smooth variety  $Y$ . Then  $X_0$  is smooth at any  $x \in Y$ . (This deduced from [7, Theorem. 14.2], as follows: Let  $\ell$  be a homogeneous linear polynomial satisfying  $\ell(x) \neq 0$ . Then, since  $\mathcal{O}_{Y,x} = \mathcal{O}_{X_0,x}/(z_0/\ell)$  is a regular local ring, the  $r+1$  functions  $z_0/\ell, g^1/\ell, f^2/\ell, \dots, f^r/\ell$  give a subset of a regular system of parameters of  $\mathcal{O}_{\mathbb{P}^N,x}$ . Hence  $\mathcal{O}_{X_0,x} = \mathcal{O}_{\mathbb{P}^N,x}/(g^1/\ell, f^2/\ell, \dots, f^r/\ell)$  is also regular.) Thus  $\text{Sing}(X_0)$  does not intersect with  $(z_0 = 0)$ , which implies that  $\text{Sing}(X_0)$  must be of dimension  $< 1$ .

Next, we set  $\alpha := (1 : \beta_1 : \dots : \beta_N) \in \mathbb{P}^N$ , where  $\alpha \neq v$ . We have  $\alpha \in \text{Sing}(X_0)$ , as follows: Since  $\beta \in Z$ , we have  $f^2(\alpha) = \dots = f^r(\alpha) = 0$ . By definition of  $g^1$ , it follows that  $g^1(\alpha) = 0$ . Hence  $\alpha \in X_0$ . Since  $\text{rk } D((f^2, \dots, f^r), \beta; \mathbb{P}^{N-1}) < r - 1$ , we have  $\text{rk } D((g^1, f^2, \dots, f^r), \alpha; \mathbb{P}^N) < r$ . Hence  $\alpha \in \text{Sing}(X_0)$ .  $\square$

**3.2. Irreducible components of the incidence scheme.** Let us consider

$$I = \left\{ ((f^k), a) \in \mathcal{H}^v \times \mathbb{P}^N \left| \begin{array}{l} f^1(a) = \dots = f^r(a) = 0 \\ \text{and } \text{rk } D((f^k), a) < r \end{array} \right. \right\},$$

the incidence scheme whose general members parameterize pairs  $(X, a)$  such that  $X$  is a complete intersection variety defined by  $(f^k) \in \mathcal{H}^v$  and that  $a$  is a singular point of  $X$ . Let  $\text{pr}_1 : I \rightarrow \mathcal{H}^v$  and  $\text{pr}_2 : I \rightarrow \mathbb{P}^N$  be the first and second projections. Here, the image  $\text{pr}_1(I) \subset \mathcal{H}^v$  is the parameter space of singular complete intersections.

For a member  $(f^1, \dots, f^r) \in \mathcal{H}^v$ , we set

$$D'((f^k), a) := \begin{bmatrix} f_{z_1}^1(a) & \cdots & f_{z_{N-1}}^1(a) \\ \vdots & & \vdots \\ f_{z_1}^r(a) & \cdots & f_{z_{N-1}}^r(a) \end{bmatrix}$$

(i.e., we remove  $f_{z_N}^k$ 's from  $D((f^k), a)$  defined in (5)), and set

$$I' := \left\{ ((f^k), a) \in \mathcal{H}^v \times \mathbb{P}^N \left| \begin{array}{l} f^1(a) = \cdots = f^r(a) = 0 \\ \text{and } \text{rk } D'((f^k), a) < r \end{array} \right. \right\},$$

where we have  $I \subset I'$ . We will investigate irreducible components of  $I'$  and  $I$ .

**Lemma 3.4.** *Let  $\Lambda'$  be an irreducible component of  $I'$ . If  $\text{pr}_2(\Lambda') = \mathbb{P}^N$ , then  $\Lambda'$  is of dimension  $\geq \dim(H^v)$ .*

**Remark 3.5.** Let  $M_k = M_k(r, n)$  be the set of  $(x_{i,j})_{1 \leq i \leq r, 1 \leq j \leq n} \in \mathbb{A}^{rn}$  such that

$$\text{rk} \begin{bmatrix} x_{1,1} & \cdots & x_{1,n} \\ \vdots & & \vdots \\ x_{r,1} & \cdots & x_{r,n} \end{bmatrix} \leq k.$$

Then  $M_k$  is an irreducible subvariety of codimension  $(r-k)(n-k)$  in  $\mathbb{A}^{rn}$  (see [1, p. 67, II, §2, Prop.]).

**Remark 3.6.** Let  $a = (a_0 : \cdots : a_N) \in \mathbb{P}^N$  be a point with  $a_N \neq 0$ . Let  $\hat{\mathcal{H}}_e^v \subset H^0(\mathbb{P}^N, \mathcal{O}(e))$  be the affine subvariety corresponding to  $\mathcal{H}_e^v \subset \mathcal{H}_e$ . We consider the morphism

$$\Phi_e : \hat{\mathcal{H}}_e^v \rightarrow \mathbb{A}^{N-1} : f \mapsto (f_{z_1}(a), \dots, f_{z_{N-1}}(a)).$$

Then  $\Phi_e$  is surjective. This is because, for  $b = (b_1, \dots, b_{N-1}) \in \mathbb{A}^{N-1}$ , we have

$$\Phi_e(b_1/a_N^{e-1} \cdot z_1 z_N^{e-1} + \cdots + b_{N-1}/a_N^{e-1} \cdot z_{N-1} z_N^{e-1}) = b.$$

*Proof of Lemma 3.4.* Let  $a = (a_0 : \cdots : a_N) \in \mathbb{P}^N$  be general. Then we can assume  $a_N \neq 0$ . Let us calculate the codimension of the intersection  $I' \cap (\mathcal{H}^v \times \{a\})$ , which is equal to the fiber of  $\text{pr}_2 : I' \rightarrow \mathbb{P}^N$  at  $a$ . First, we set

$$\Phi = \bigoplus_{1 \leq k \leq r} \Phi_{e^k} : \hat{\mathcal{H}}^v \rightarrow \mathbb{A}^{r(N-1)} : (f^k) \mapsto D'((f^k), a),$$

where  $\hat{\mathcal{H}}^v := \bigoplus \hat{\mathcal{H}}_{e^k}^v \subset \bigoplus H^0(\mathbb{P}^N, \mathcal{O}(e^k))$  is the affine subvariety corresponding to  $\mathcal{H}^v \subset \mathcal{H}$ . From Remark 3.6,  $\Phi$  is surjective. Moreover,  $\Phi$  is a smooth morphism, since it is regarded as a linear map of vector spaces. Hence  $\Phi^{-1}(M_{r-1}) \subset \hat{\mathcal{H}}^v$  is of codimension  $N - r$ , where  $M_{r-1} = M_{r-1}(r, N-1) \subset \mathbb{A}^{r(N-1)}$  is the subvariety defined in Remark 3.5. Then each irreducible component of

$$(6) \quad \Phi^{-1}(M_{r-1}) \cap F_a \subset \hat{\mathcal{H}}^v$$

is of codimension  $\leq N$ , where  $F_a := \{(f^k) \in \mathcal{H}^v \mid f^1(a) = \cdots = f^r(a) = 0\}$ . Now, the projective variety in  $\mathcal{H}^v$  corresponding to the affine variety (6) can be identified with

$$I' \cap (\mathcal{H}^v \times \{a\}) \subset (\mathcal{H}^v \times \{a\}).$$

Hence each irreducible component of  $I' \cap (\mathcal{H}^v \times \{a\})$  is of codimension  $\leq N$ .

Since  $\Lambda' \rightarrow \mathbb{P}^N$  is surjective and since  $a \in \mathbb{P}^N$  is general, there exists an irreducible component  $W$  of  $I' \cap (\mathcal{H}^v \times \{a\})$  such that  $W$  coincides with an irreducible component of  $\Lambda' \cap (\mathcal{H}^v \times \{a\})$ . The reason is as follows. Let  $U = \Lambda' \setminus \bigcup_{i=1}^m V_i$ , where  $V_1, \dots, V_m$  are the irreducible components of  $I'$  with  $V_i \neq \Lambda'$ . Since  $U \rightarrow \mathbb{P}^N$  is dominant, we can take an irreducible component  $W$  of  $I' \cap (\mathcal{H}^v \times \{a\})$  such that  $W \cap U \neq \emptyset$ . Then  $W$  is contained in the fiber  $\Lambda' \cap (\mathcal{H}^v \times \{a\})$ , and is indeed an irreducible component of this fiber.

In addition, since  $a$  is general, we have

$$\text{codim}(\Lambda', \mathcal{H}^v \times \mathbb{P}^N) = \text{codim}(W, \mathcal{H}^v \times \{a\})$$

(see [3, II, Ex. 3.22]). Hence  $\text{codim}(\Lambda', \mathcal{H}^v \times \mathbb{P}^N) \leq N$ , which implies the assertion.  $\square$

**Lemma 3.7.** *Let  $\Lambda' \subset I'$  be an irreducible subset. If  $\text{pr}_2(\Lambda') \not\subset (z_N = 0)$ , then  $\Lambda' \subset I$ .*

*Proof.* Let  $((f^k), a) \in \Lambda'$  be a general member. Since  $\text{pr}_2(\Lambda') \not\subset (z_N = 0)$ ,  $a = (a_0 : \cdots : a_N)$  satisfies  $a_N \neq 0$ . Then, since  $f_{z_0}^k = 0$ , it follows from Euler's formula  $\sum_{j=0}^N a_j f_{z_j}^k(a) = e^k f^k(a) = 0$  that

$$f_{z_N}^k(a) = -(a_1/a_N \cdot f_{z_1}^k(a) + \cdots + a_{N-1}/a_N \cdot f_{z_{N-1}}^k(a))$$

holds for  $1 \leq k \leq r$ . Thus the last column vector of the matrix  $D((f^k), a)$  defined in (5) can be written as a linear combination of the other column vectors. Then, since  $D'((f^k), a)$  is of rank  $< r$ , so is  $D((f^k), a)$ . It follows that  $((f^k), a) \in I$ .  $\square$

**Theorem 3.8.** *Let  $v \in \mathbb{P}^N$  be a point, and let  $e^1, \dots, e^r$  be  $r$  integers greater than 1. Let  $\mathcal{H}^v = \mathcal{H}_{e^1}^{v_1} \times \cdots \times \mathcal{H}_{e^r}^{v_r}$  be the parameter space defined in (3). Assume that the space  $\mathcal{H}$  defined in (2) is not equal to the parameter space of quadric hypersurfaces of odd dimensions in  $p = 2$ . Then  $\text{pr}_1(I) = \mathcal{H}^v$ , which means that every complete intersection variety defined by a member of  $\mathcal{H}^v$  is singular.*

*Proof.* If  $p = 0$  or  $e^k < p$  for any  $k$ , then it follows from Remark 2.3 and Corollary 2.16 that every  $X$  defined by a member of  $\mathcal{H}^v$  is a cone with vertex  $v$ , in particular, is singular.

Now we assume that  $p > 0$  and  $e^k \geq p$  for some  $k$ . From Proposition 3.2, we can take  $((f^k), \alpha) \in \mathcal{H}^v \times \mathbb{P}^N$  such that  $\alpha \neq v$  is an isolated singular point of the complete intersection variety  $X_0 \subset \mathbb{P}^N$  defined by  $(f^k)$ .

We denote by  $\text{PGL}(\mathbb{P}^N; v) \subset \text{PGL}(\mathbb{P}^N)$  the group of automorphisms  $\sigma$  of  $\mathbb{P}^N$  such that  $\sigma(v) = v$ . Let us consider the subset of  $I$  parameterizing pairs

$(\sigma(X_0), \sigma(\alpha))$  with  $\sigma \in \mathrm{PGL}(\mathbb{P}^N; v)$ , which is actually given by the orbit of  $((f^k), \alpha)$  in  $I$ ,

$$\{ (((\sigma^{-1})^* f^k), \sigma(\alpha)) \in I \mid \sigma \in \mathrm{PGL}(\mathbb{P}^N; v) \},$$

where  $(\sigma^{-1})^* f^k(z) := f^k(\sigma^{-1}(z))$ . Now we take an irreducible component  $\Lambda$  of  $I$  containing the above orbit. Then  $\mathrm{pr}_2(\Lambda) = \mathbb{P}^N$ . Here we have

$$(7) \quad \dim(\Lambda) = \dim(\mathrm{pr}_1(\Lambda)).$$

The reason is as follows: The fiber of  $\mathrm{pr}_1 : I \rightarrow \mathcal{H}^v$  at  $(f^k)$  is equal to  $\mathrm{Sing}(X_0)$ . Since  $\alpha$  is isolated, the set  $\{\alpha\}$  is an irreducible component of  $\mathrm{Sing}(X_0)$ . Since each irreducible component of a fiber of  $\Lambda \rightarrow \mathrm{pr}_1(\Lambda)$  must be of dimension  $\geq \dim(\Lambda) - \dim(\mathrm{pr}_1(\Lambda))$ , the equality  $\dim(\Lambda) = \dim(\mathrm{pr}_1(\Lambda))$  holds.

Let  $\Lambda' \subset I'$  be an irreducible component of  $I'$  such that  $\Lambda \subset \Lambda'$ . From Lemma 3.4,  $\dim(\Lambda') \geq \dim(\mathcal{H}^v)$ . From Lemma 3.7, we have  $\Lambda = \Lambda'$ . From (7), we have  $\dim(\mathrm{pr}_1(\Lambda)) \geq \dim(\mathcal{H}^v)$ , and hence  $\mathrm{pr}_1(\Lambda) = \mathcal{H}^v$ .  $\square$

*Proof of Theorem 1.1.* The equivalence “(a)  $\Leftrightarrow$  (c)” follows immediately from Proposition 2.10 and Remark 2.12. The implication “(b)  $\Rightarrow$  (a)” follows as in Example 2.2. Now we show the implication “(a)  $\Rightarrow$  (b)”. Let  $X \subset \mathbb{P}^N$  be a smooth  $(e^1, \dots, e^r)$ -complete intersection variety, and assume that  $X$  is strange for a point  $v \in \mathbb{P}^N$ . It is sufficient to consider the case where  $X$  is non-degenerate. Then  $e^k > 1$  for any  $k$ . As in Remark 3.1, it follows from Proposition 2.13 that  $X$  is defined by a member of  $(f^k) \in \mathcal{H}^v$ . Since  $X$  is smooth, it follows from Theorem 3.8 that  $X$  must be a quadric of odd dimension in  $p = 2$ .  $\square$

*Acknowledgments.* The author would like to thank Hajime Kaji for helpful discussions and suggestions. The author also would like to thank Satoru Fukasawa and Yasunari Nagai for their valuable comments. The author was partially supported by JSPS KAKENHI Grant Number 25800030.

## REFERENCES

- [1] E. Arbarello, M. Cornalba, P. A. Griffiths, J. Harris, *Geometry of Algebraic Curves*, Volume I. Springer 1985.
- [2] A. van den Essen, *Polynomial automorphisms and the Jacobian conjecture*, Progress in Mathematics, vol. **190**, Birkhauser Verlag, Basel, 2000.
- [3] R. Hartshorne, *Algebraic Geometry*, Graduate Texts in Math. **52**, Springer, New York, 1977.
- [4] A. Holme and E. Lluís, Strange varieties, Appendix to “Embeddings, projective invariants and classifications”, *Monografias del Instituto de Matemáticas, Univ. Nac. Autónoma de México*. Vol. 7, 1979, pp. 93–122.
- [5] S. L. Kleiman and R. Piene, On the inseparability of the Gauss map, “Enumerative Algebraic Geometry (Proceedings of the 1989 Zeuthen Symposium)”, *Contemp. Math.* **123**, Amer. Math. Soc., Providence, 1991, pp. 107–129.
- [6] E. Lluís: Variedades algebraicas con ciertas condiciones en sus tangentes, *Bol. Soc. Mat. Mexicana* (2) **7** (1962), 47–56.

- [7] H. Matsumura, Commutative ring theory, Cambridge Stud. Adv. Math., vol. **8**, Cambridge University Press, Cambridge, 1989.
- [8] S. Mori, Projective manifolds with ample tangent bundles, Ann. Math. (2) **110**(3) (1979), 593–606.
- [9] S. Okuda, Kernels of derivations in positive characteristic, Hiroshima Math. J. **34** (2004), 1–19
- [10] R. Piene, Numerical characters of a curve in projective n-space. Real and complex singularities, Proc. Conf. Oslo, 1976 (P. Holm, Ed.), pp. 475–495, Groningen, Sijthoff & Nordhoff, 1977.
- [11] P. Samuel: Lectures on old and new results on algebraic curves, Tata Inst. Fund. Res. Lectures on Math. **36**, Tata Inst. Fund. Res., Bombay 1966.
- [12] J. M. Wahl, A cohomological characterization of  $P^n$ , Invent. Math, **72** (1983), 315–322.

*E-mail address:* [katu@toki.waseda.jp](mailto:katu@toki.waseda.jp)

*URL:* <http://www.aoni.waseda.jp/katu/index.html>

DEPARTMENT OF MATHEMATICS, SCHOOL OF FUNDAMENTAL SCIENCE AND ENGINEERING, WASEDA UNIVERSITY, OHKUBO 3-4-1, SHINJUKU, TOKYO, 169-8555, JAPAN