

# ON CREMONA TRANSFORMATIONS OF $\mathbb{P}^3$ WITH ALL POSSIBLE BIDEGREES

IVAN PAN

**ABSTRACT.** For every ordered pair  $(d, e)$  of integer numbers  $d, e \geq 2$ , such that  $\sqrt{d} \leq e \leq d$ , we construct a birational map  $\mathbb{P}^3 \dashrightarrow \mathbb{P}^3$  defined by homogeneous polynomials of degree  $d$  whose inverse map is defined by homogeneous polynomials of degree  $e$ .

## 1. INTRODUCTION

The aim of this note is to correct a mistake in the proof of Theorem [Pa2000-2, Théorème. 2.2]. The proof of that theorem depends on the example [Pa2000-2, Exemple 2.1] which is wrong.

We propose an explicit construction of Cremona transformations of  $\mathbb{P}^3$  (see §2, especially Lemma 2) which, together with their inverse maps, provide all possible bidegrees (Theorem 3 and Corollary 4).

**Acknowledge** We would like to thank Igor Dolgachev for pointing out a mistake in [Pa2000-2, Exemple 2.1].

## 2. MAIN CONSTRUCTION AND RESULTS

Let  $\mathbb{P}^3$  be the projective space over an algebraically closed field  $k$  of characteristic zero; we fix homogeneous coordinates  $w, x, y, z$  on  $\mathbb{P}^3$ .

We recall that a Cremona transformation of  $\mathbb{P}^3$  is a birational map  $F : \mathbb{P}^3 \dashrightarrow \mathbb{P}^3$ . We say  $F$  has *bidegree*  $(d, e)$  when  $F$  and its inverse  $F^{-1}$  are defined by homogeneous polynomials, without non trivial common factors, of degrees  $d$  and  $e$  respectively; notice that in this case  $F^{-1}$  has bidegree  $(e, d)$ . If  $V \subset \mathbb{P}^3$  is a dense open set over which  $F^{-1}$  is defined and injective and  $L \subset \mathbb{P}^3$  is a line with  $L \cap V \neq \emptyset$ , then  $e$  is the degree of the closure of  $F^{-1}(L \cap V)$ ; one deduces that  $\sqrt{d} \leq e \leq d$  (see for example [Pa2000-2, §1]).

If  $X \subset \mathbb{P}^2$  is a curve and  $p \in \mathbb{P}^2$  we denote by  $\text{mult}_p(X)$  the multiplicity of  $X$  at  $p$ . If  $S, S' \subset \mathbb{P}^3$  are surfaces and  $C \subset S \cap S'$  is an irreducible component, we denote by  $\text{mult}_C(S, S')$  the intersection multiplicity of  $S$  and  $S'$  along  $C$ .

---

<sup>1</sup>Partially supported by the *Agencia Nacional de Investigadores* of Uruguay

Consider a rational map  $T : \mathbb{P}^3 \dashrightarrow \mathbb{P}^3$  defined by

$$T = (g : qt_1 : qt_2 : qt_3),$$

where  $t_1, t_2, t_3 \in k[x, y, z]$  are homogeneous of degree  $r$ , without non trivial common factors, and  $g, q \in k[w, x, y, z]$  are homogeneous of degrees  $d, d - 1$ , with  $d \geq r \geq 1$  and  $g$  irreducible. We know that  $T$  is birational if  $\tau := (t_1 : t_2 : t_3) : \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$  is birational and  $g, q$  vanish at  $o = (1 : 0 : 0 : 0)$  with orders  $d - 1$  and  $\geq d - r - 1$ , respectively (see [Pa2000-1, Proposition 2.2]).

On the other hand, consider  $2r - 1$  points  $p_0, p_1, \dots, p_{2r-2}$  in  $\mathbb{P}^2$ ,  $r \geq 2$ , satisfying the following condition:

*There exist curves  $X_r, Y_{r-1} \subset \mathbb{P}^2$  of degrees  $r, r - 1$ , respectively, with  $X_r$  irreducible, such that  $\text{mult}_{p_0}(X_r) = r - 1$ ,  $\text{mult}_{p_0}(Y_{r-1}) \geq r - 2$  and  $p_i \in X_r \cap Y_{r-1}$  for  $i = 1, \dots, 2r - 2$ .* (I)

Hence *loc. cit.* also implies there exists a plane Cremona transformation defined by polynomials of degree  $r$  with a point of multiplicity  $r - 1$  at  $p_0$  and passing through  $p_1, \dots, p_{2r-2}$  with multiplicity 1: indeed, if we consider  $p_0 = (1 : 0 : 0)$  and take polynomials  $t_1$  and  $f$ , of degrees  $r$  and  $r - 1$ , defining  $X_r$  and  $Y_{r-1}$  respectively, then  $(t_1 : yf : zf) : \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$  is a Cremona transformation as required; such a transformation is said to be *associated* to the points  $p_0, p_1, \dots, p_{2r-2}$ .

*Remark 1.* The transformations satisfying the condition (I) are general cases of the so-called *de Jonquières transformations* (see [dJo1864] or [Alb2000, Def. 2.6.10]). We note that the Enriques criterion [Alb2000, Thm. 5.1.1] may be used to prove that a set of  $2r - 2$  points  $p_0, p_1, \dots, p_{2r-2}$  with assigned multiplicities  $r - 1, 1, \dots, 1$ , and satisfying the condition (I), defines a de Jonquières transformation.

Set  $r = d$  and take an irreducible homogeneous polynomial  $g = wA(x, y, z) + B(x, y, z)$  of degree  $d$ ; that is,  $g \in k - \{0\}$  in the considerations above. Denote by  $T_{g,\tau}$  the Cremona transformation defined by

$$T_{g,\tau} = (g : t_1 : t_2 : t_3), \tag{1}$$

where  $\tau = (t_1 : t_2 : t_3)$  is associated to  $2d - 1$  points satisfying the condition (I).

We have

**Lemma 2.** *Let  $d \geq 2$  be an integer number. Then*

- (a) *there exist  $g$  and  $\tau$  such that  $T_{g,\tau}$  has bidegree  $(d, 2d - 1 - m)$ , for  $0 \leq m \leq d - 1$ .*
- (b) *there exist  $g$  and  $\tau$  such that  $T_{g,\tau}$  has bidegree  $(d, d^2 - \ell^2 - m)$ , for  $0 \leq \ell < d - 1$  and  $0 \leq m \leq 2d - 2$ .*

*Proof.* We identify  $\mathbb{P}^2$  with the plane  $\{w = 0\} \subset \mathbb{P}^3$  and consider a point  $p_0 \in \mathbb{P}^2$ . Without loss of generality, we may suppose  $p_0 = (0 : 1 : 0 : 0)$ . We recall  $o = (1 : 0 : 0 : 0)$ .

In order to prove (a) we first choose  $g \in k[w, x, y, z]$  to be a homogeneous polynomial which vanishes along the line  $op_0$  with order  $d - 1$  and is general with respect to this condition. In other words, one has  $g = wA + B$  with

$$A = A_{d-1}(y, z), B = xB_{d-1}(y, z) + B_d(y, z),$$

where  $A_i, B_i \in k[y, z]$  are general homogeneous polynomials of degree  $i$ . Hence  $A = 0$  defines a union of  $d - 1$  distinct lines in  $\mathbb{P}^2$  passing through  $p_0$  and  $B = 0$  defines an irreducible curve of degree  $d$  with an ordinary singular point of multiplicity  $d - 1$  at  $p_0$ .

Notice that, by construction, in the open set  $\mathbb{P}^2 - \{p_0\}$  the curves  $A = 0$  and  $B = 0$  intersect at  $d(d - 1) - (d - 1)^2 = d - 1$  points; in particular, if  $m \leq d - 1$ , there exist  $m$  points  $p_1, \dots, p_m \in \mathbb{P}^2$  satisfying  $A(p_i) = B(p_i) = 0$  for  $1 \leq i \leq m$ . We consider  $m$  such points and choose  $2d - 1 - m$  points  $p_{m+1}, \dots, p_{2d-2} \in \mathbb{P}^2$  with  $A(p_j) \neq 0$  and  $B(p_j) = 0$ , for all  $j = m + 1, \dots, 2d - 2$ , such that  $p_0, p_1, \dots, p_{2d-2}$  satisfy (I). Let  $\tau$  be a plane Cremona transformation associated to these  $2d - 1$  points.

Now we consider a Cremona transformation  $T_{g,\tau} : \mathbb{P}^3 \dashrightarrow \mathbb{P}^3$  as in (1). A general member in the linear system defining  $T_{g,\tau}$  is an irreducible surface of degree  $d$ ,  $S$  say, with equation of the form

$$ag + a_1t_1 + a_2t_2 + a_3t_3 = 0,$$

where  $a, a_1, a_2, a_3 \in k$  are general. Therefore  $S$  admits an ordinary singularity of multiplicity  $d - 1$  at the generic point of (the line)  $op_0$  and is smooth at the generic point of  $op_i$  for  $1 \leq i \leq m$ . If  $S'$  is another general member of that linear system, then there exists an irreducible rational curve  $\Gamma$  of degree  $e = \deg(T_{g,\tau}^{-1})$  such that the intersection scheme  $S \cap S'$  is supported on

$$\Gamma \cup (\cup_{i=0}^m op_i).$$

We have

$$\text{mult}_\Gamma(S, S') = 1, \text{mult}_{op_0}(S, S') = (d - 1)^2, \text{mult}_{op_i}(S, S') = 1, i = 1, \dots, m,$$

hence  $e = d^2 - (d - 1)^2 - m = 2d - 1 - m$ , which proves the assertion (a).

To prove (b) we proceed analogously. This time we choose  $g = wA + B$  with

$$A = \sum_{i=\ell}^{d-1} x^{d-1-i} A_i(y, z), B = \sum_{j=\ell}^d x^{d-j} B_j(y, z),$$

where  $A_i, B_i \in k[y, z]$  are general homogeneous polynomials of degree  $i$ . Since  $\ell \leq d - 2$  there exist points  $p_1, \dots, p_{2d-2} \in \mathbb{P}^2$  such that  $A(p_i) = B(p_i) = 0$  for  $1 \leq i \leq m$  and  $A(p_j) \neq 0, B(p_j) = 0$  for  $j = m + 1, \dots, 2d - 2$ : indeed, in the open set  $\mathbb{P}^2 - \{p_0\}$ , the curves  $A = 0$  and  $B = 0$  intersect at  $d(d - 1) - \ell^2 \geq d(d - 1) - (d - 2)^2 = 3d - 4$  points. Thus we can define  $\tau$  as before and obtain the assertion (b).

□

**Theorem 3.** *There exist Cremona transformations of bidegree  $(d, e)$  for  $d \leq e \leq d^2$ .*

*Proof.* From the part (a) of Lemma 2 we deduce that there exist Cremona transformations of bidegrees  $(d, e)$  for  $d \leq e \leq 2d - 1$ .

Now we use the part (b) of Lemma 2. Suppose  $\ell < d - 1$  and think of  $e = d^2 - \ell^2 - m$  as a function  $e(\ell, m)$  depending on  $\ell, m$ ; to complete the proof it suffices to show that the image of that function contains  $\{2d, 2d + 1, \dots, d^2\}$ .

We note that  $e(d - 2, 2d - 2) = 2d - 2$  and  $e(0, 0) = d^2$ ; in other words, the part (b) of Lemma 2 implies that there exist Cremona transformations of bidegrees  $(d, 2d - 2)$  and  $(d, d^2)$ . On the other hand  $e(\ell, 0) - e(\ell - 1, 2d - 2) = 2(d - \ell) - 1 > 0$ . Since  $e(\ell, m)$  decreases with respect to  $m$ , we easily obtain the result. □

For  $d = 2$  the theorem above asserts that there exist Cremona transformations of bidegrees  $(2, 2), (2, 3), (2, 4)$ ; analogously for  $d = 3$  and bidegrees  $(3, 3), (3, 4), \dots, (3, 9)$ , and so on. By symmetry we deduce

**Corollary 4.** *There exist Cremona transformations of bidegrees  $(d, e)$  with  $\sqrt{d} \leq e \leq d^2$ .*

*Remark 5.* The inequality  $\sqrt{d} \leq e \leq d^2$  is the unique obstruction to the degree for the inverse of a Cremona transformation of degree  $d$  in  $\mathbb{P}^3$ .

## REFERENCES

- [Alb2000] M. Alberich-Carramiñana, *Geometry of the plan Cremona maps*, Lectures Notes in Math. 1769, Springer (2000).
- [dJo1864] E. de Jonquières, *Mémoire sur les figures isographiques et sur un mode uniforme de génération des courbes à courbure d'un ordre quelconque au moyen de deux faisceaux correspondants de droites*, Nouvelles annales de mathématiques 2e série, tome 3 (1864), p. 97-111.
- [Pa2000-1] I. Pan, *Les transformations de Cremona stellaires*, Proc. of AMS, **129**, N. 5 (2000), pp.1257-1262.
- [Pa2000-2] I. Pan, *Sur les multidegrés des transformations de Cremona*, C. R. Acad. Sci. Paris, t. 330, Série I, pp. 297-300, 2000.

*E-mail address:* ivan@cmat.edu.uy