

On radiation reaction and the Abraham-Lorentz-Dirac equation

A. Cabo Montes de Oca*** and Jorge Castineiras*

* *Programa de Pós-Graduação em Física (PPGF) da Universidade Federal do Pará (UFPA), Av. Augusto Correa, No. 01, Campus Básico do Guamá, Belém, Pará, Brasil and*

** *Departamento de Física Teórica, Instituto de Cibernética Matemática y Física (ICIMAF), Calle E, No. 309, entre 13 y 15, Vedado, La Habana, Cuba*

It is underlined that the Lienard-Wiechert solutions indicate that after the external force is instantly removed from a small charged particle, the field in its close neighborhood becomes a Lorentz boosted Coulomb field. It suggests that the force of the self-field on the particle should instantaneously vanish after a sudden removal of the external force. A minimal modification of Abraham-Lorentz-Dirac equation is searched seeking to implement this property. A term assuring this behaviour is added to the equation by maintaining Lorentz covariance and vanishing scalar product with the four-velocity. The simple Diracs constant force example does not show runaway acceleration.

A. Cabo E-Mail: cabo@icimaf.cu,
J. Castineiras E-Mail: jcastin@ufpa.br

The correctness of the Abraham-Lorentz-Dirac (ALD) equations for the point particles interacting with their own fields had been a recurrent issue in searching for a consistent formulation of Classical Electrodynamics [1–10]. The formulation of the equations, already from the start, presented theoretical difficulties which had been discussed along decades in the extensive literature devoted to theme. The presence of solutions with values growing without limit (runaway behavior) or pre-accelerated motions in advance to the applied forces, were the basic un-physical properties exhibited by the ALD equations [8, 9].

A relevant advance in the analysis of these difficulties was done in Ref. [8]. In this work, it was argued that the ALD equations can't be derived in a completely exact way from the integral equations. This happens in the coupled system formed by the charged body having the electric charge concentrated in its surface and interacting with its electromagnetic field, when the force has a non analytic behavior. Moreover, the author was able to derive a correction to the ALD equations which should be introduced in order to consider suddenly changing forces. The change corresponded to multiplying the reaction force of the field on the particle by a proper-time dependent function $\eta(\tau)$ which vanish at the proper time value at which the forces is started to be applied and attains a unit value in a small time interval of the order of the time required by light to travel the radius of the considered structured body studied in Ref. [8]. This change was argued to allow eliminating both: the runaway and the pre-accelerated types of solutions. Thus, in general terms, it can be concluded that the ALD equations are not exact consequences of the integral equations of motions for the coupled system describing the structured particle investigated in Ref. [8]. It can be also remarked that, since the size a of the system considered in the work is arbitrary, the elimination of the un-physical nature of the solutions is possible for whatever arbitrary small value of a is chosen. This indicates that the solution might be valid also in the point like limit of the particle structure.

Although, the described study constitute a relevant step in the understanding of the problem, the wrote modification of the ALD equations seems somewhat restricted by the particular process of applying a force showing a non analytic proper time dependence given by a sudden jump from zero to a finite value at a given instant. Henceforth, it makes sense to investigate the possibility of formulating the point particle equations of motion in alternative form exhibiting causality and absence of runaway solutions.

In this work we intend to advance a formulation of the equation of motion in a form directly showing causality. The discussion, is suspected that could implement the main physical elements leading to the elimination of the runaway and pre-accelerated solutions which were discovered in Ref. [8]. The basic idea motivating the search is the following observation: when an applied force is suddenly removed from acting over a point particle, the Lienard-Wiechert solutions for the self-field surrounding it, shows that after an arbitrarily short time interval τ passed, the field solution within a sufficiently small neighborhood of the particle becomes rigorously given by a Lorentz boosted Coulomb field. But, a Lorentz boosted Coulomb field is not expected to exert any force on the particle generating it. This known simple property strongly suggests, that the force applied by the self-field on the particle should also instantaneously vanish after the sudden removal of the force. Therefore, since the ALD equations are widely expected to be valid for continuously varying forces, a minimal modifications of them is here searched in order to implement the instantaneously vanishing of the acceleration after a sudden removal of the applied force.

The proposal for the modified equation of motion is simple: to add a term of the same order of the Schott and Larmor one being jointly determined by the dynamics and by the external force. The dynamic part of the new terms

is simply linear in the four-acceleration and the element defined by the forces is a Lorentz invariant function of the proper time. It is completely defined by an assumed piecewise continuous external force as a function of the proper time. The linear in the acceleration dependence of the added contribution assures that its scalar product with the four velocity vanishes, which is a consistency condition of the relativistic invariant ALD equations. In this letter, in first place the modified equation is presented and the satisfaction of the equations argued for solutions being linear motions in the intervals of vanishing forces, and given by motions solving the ALD equations in the region for which the forces are finite. Next, the equations are solved for the linear motion in which a constant force in the observer's frame is applied during a finite time interval. This is one simple motion discussed by Dirac in its classical article. For this case he encountered for the first time the situation in which the acceleration (due to its continuity required by its second derivative appearing in the equations), after a sudden elimination of the applied force remained non vanishing and more drastically, it did not decrease to a vanishing value continuously, but increased indefinitely. The presented solution eliminates this problem because the new term adds a Dirac's delta contribution in the proper time, which just cancels the also Delta like terms appearing with the acceleration instantaneously jump to its zero value. That is, the modification done simply allows the discontinuous transition between the accelerated motion solution of the ALD equation (just existing before the instant in which the force is eliminated) to the also solving the ALD equation, uniform motion state. One point of interest to remark is that the presented alternative equation seems to be, at least, coherent with the picture of the electromagnetic propagation as described by the Lienard-Wiechert potentials. These basic equations for the interaction of this field with charged objects, indicate that when a particle is instantly acquiring a state of uniform motion, at the exactly the same instant, a wavefront start propagating from the point in which the particles is. Inside the wavefront the field is the one associated to a moving charge with constant velocity. Therefore, the self forces acting on the particle in this interval should be zero. The modified equations, since they predict vanishing of the acceleration after removing the forces, is at least compatible with this property and solves the difficulty with the runaway solution after the force suddenly vanishes [11]. The discussion presented also suggests a way for the discretization of a general trajectory produced by a continuous in the proper time external force, in a way which could probably eliminate the instabilities that currently beset the numerical solution of the ADL equation. The idea is that by a discretization of the movement in infinitesimal time intervals in which the forces periodically vanishes, and modifying the ALD equations in the described way, the developing of the numerical instabilities of the usual ALD equation (being valid in the intervals in which the force is not vanishing) could be eliminated. These questions are expected to be studied in extensions of the work to be considered elsewhere.

The letter proceeds as follows. Firstly the modified ALD equations and its general solution are presented. Next, the constant force case is discussed

Let us consider a general motion of a particle P along a space-time trajectory C defined by a curve $x(\tau) = (x_0^0(\tau), x^1(\tau), x^2(\tau), x^3(\tau))$ and parameterized by the proper time τ . The metric signature to be used will be $(1, -1, -1, -1)$. The modified ALD equations are proposed in the specific form

$$\begin{aligned} f^\mu(\tau) &= m a^\mu(\tau) - \kappa \frac{da^\mu(\tau)}{d\tau} - \kappa a^\nu(\tau) a_\nu(\tau) u^\mu(\tau) + \kappa a^\mu(\tau) \sigma(\tau), \\ u^\mu(\tau) &= \frac{dx^\mu(\tau)}{d\tau}, \\ a^\mu(\tau) &= \frac{du^\mu(\tau)}{d\tau}, \end{aligned}$$

where the proper time function $\sigma(\tau)$ will be defined in what follows. The equations satisfies the relativistic condition consisting in that the scalar product of the four-velocity with them should vanish. This follows from the form of the new added term, since

$$\kappa u^\mu(\tau) a_\mu(\tau) \sigma(\tau) = 0.$$

The scalar function of the proper time $\sigma(\tau)$ will be chosen to be determined by the force. Let us assume that the force is a piecewise continuous function of the proper time τ , and define the sequences of proper times values (s_n, q_n) , where s_n is the set of increasing with the integer value of n , values of the proper time when the forces discontinuously turns its values from zero to a finite one. The values of q_n indicate the instants at which the force is removed. An instant before q_n the force is finite and an instant after, it vanishes. Further, in the pair (s_n, q_n) , the time s_n is assumed to be one at which the force is "connected" for afterwards being "disconnected" only at the subsequent

instant q_n . A first step in the definition of $\sigma(\tau)$ is to define the function $\beta(\tau)$ in the form

$$\beta_\epsilon(\tau) = \begin{cases} 0 & \tau \in (q_{n-1}, s_n) \\ \frac{\tau - s_n}{\epsilon} & \tau \in (s_n, s_n + \epsilon) \\ 1 & \tau \in (s_n + \epsilon, q_n - \epsilon) \\ 1 - \frac{\tau - (q_n - \epsilon)}{\epsilon} & \tau \in (q_n - \epsilon, q_n) \\ 0 & \tau \in (q_n, s_{n+1}) \end{cases}$$

$$\tau \in (q_{n-1}, s_{n+1}), \quad \epsilon > 0.$$

Note for all value of n the function becomes defined at all times. The small ϵ parameter appearing is introduced in order that the derivative over the proper time of the function leads to Dirac delta functions with their support defined in the internal part of the intervals (s_n, q_n) . That means that the delta functions under integration will give the values of the limits at the boundaries of the function when taken from the interior points of the intervals (s_n, q_n) .

The function $\sigma(\tau)$ is defined as the proper time derivative of $\beta(\tau)$ when the limit $\epsilon \rightarrow 0$

$$\begin{aligned} \sigma(\tau) &= \lim_{\epsilon \rightarrow 0} \frac{d\beta_\epsilon(\tau)}{d\tau} \\ &= \sum_{n=1} (\delta^{(+)}(\tau - s_n) - \delta^{(-)}(\tau - q_n)), \end{aligned}$$

and the plus and minus signs as superindices in $\delta^{(+)}(\tau - s_n)$ and $\delta^{(-)}(\tau - q_n)$, indicate that the supports of the Delta functions are at the interior of the intervals (s_n, q_n) . This completes the definition of the modified ALD equations.

Lets us present an explicit constructions of solutions of these equations. For this purpose consider the following definitions. Assume that the acceleration of the proposed solutions is only non vanishing in all intervals (s_n, q_n) . That is

$$a^\mu(\tau) = \sum_{n=1} \theta(\tau - s_n) \theta^{(-)}(q_n - \tau) a_n^\mu(\tau),$$

where $a_n^\mu(\tau)$ is the acceleration of a solution of the ALD equations in the considered interval, which satisfies initial conditions expressing the continuity of the coordinates, velocity and acceleration at all the points s_n , that is

$$\begin{aligned} f^\mu(\tau) &= m a_{s_n}^\mu(\tau) - \kappa \frac{da_{s_n}^\mu(\tau)}{d\tau} - \kappa a_{s_n}^\nu(\tau) a_{s_n \nu}(\tau) u_{s_n}^\mu(\tau), \\ \tau &\in (s_n, q_n), \\ x_{s_n}^\mu(s_n^+) &= x_{q_n}^\mu(s_n^-), \\ u_{s_n}^\mu(s_n^+) &= u_{q_n}^\mu(s_n^-), \\ a_{s_n}^\mu(s_n^+) &= 0. \end{aligned}$$

In the interior points of the intervals (q_n, s_{n+1}) , since the accelerations are vanishing the motion will be assumed to be uniform ones. The coordinates and velocities, indicated with the subindex q_n , take the expressions

$$\begin{aligned} x_{q_n}^\mu(\tau) &= \sum_{n=1} \theta(\tau - q_n) \theta(s_{n+1} - \tau) (x_n^\mu(q_n) + u_n^\mu(q_n)(\tau - q_n)), \\ u_{q_n}(\tau) &= \sum_{n=1} \theta^{(+)}(\tau - q_n) \theta^{(-)}(s_{n+1} - \tau) u_n^\mu(q_n), \\ a_{q_n}^\mu(\tau) &= 0. \end{aligned}$$

The being constructed solution for the coordinates in all the proper time axis is written as the sum of the two expressions defined above. The coordinate and their derivatives are continuous at the values of the proper times thanks to their definition in the above equations.

$$\begin{aligned} x^\mu(\tau) &= \sum_{n=1} (x_{q_n}^\mu(\tau) \theta(\tau - q_n) \theta(s_{n+1} - \tau) + \\ &\quad x_{s_n}^\mu(\tau) \theta(\tau - s_n) \theta(q_n - \tau)). \end{aligned}$$

It can be noted that the ALD equations are satisfied almost everywhere, possibly except at the discrete set of points s_n, q_n . Further, since the coordinates and velocities are defined in a continuous way at those points, but also

the acceleration vanishes at the right of the points s_n (due to the chosen initial conditions after the force is connected) the new added term vanishes at these points and the proposed equation of motion is also satisfied there. Thus, the equations could be not be obeyed only at the ending points of the intervals q_n . However, although at those points there is a discontinuity in the acceleration, the evaluation of the proposed solution in the modified ALD equations in the sense of the generalized functions leads to

$$\begin{aligned}
& -f^\mu(\tau) + m a^\mu(\tau) - \kappa \frac{da^\mu(\tau)}{d\tau} - m a^\nu(\tau) a_\nu(\tau) u^\mu(\tau) + \kappa a^\mu(\tau) \sigma(\tau) \\
& = \kappa \sum_{n=1} a_{s_n}^\mu(q_n) \delta(\tau - q_n) - \kappa \sum_{n=1} a_{s_n}^\mu(q_n) \delta^{(-)}(\tau - q_n) \\
& = 0.
\end{aligned}$$

Therefore the defined function satisfies the modified equations of motion. The general set of equations in terms of the coordinates can be written in the form

$$\begin{aligned}
f^\mu(\tau) &= m \frac{d^2 x^\mu(\tau)}{d\tau^2} - \kappa \frac{d^3 x^\mu(\tau)}{d\tau^3} - \kappa \frac{d^2 x^\nu(\tau)}{d\tau^2} \frac{d^2 x_\nu(\tau)}{d\tau^2} \frac{dx^\mu(\tau)}{d\tau} + \kappa \frac{d^2 x^\mu(\tau)}{d\tau^2} \sigma(\tau), \\
x^\mu(\tau_0) &= x_0^\mu, \\
\frac{dx^\mu(\tau_0)}{d\tau} &= u^\mu(\tau_0) = u_0^\mu, \\
\frac{d^2 x^\mu(\tau_0)}{d\tau^2} &= a^\mu(\tau_0) = 0,
\end{aligned}$$

in which initial conditions were fixed at given point internal to the intervals of vanishing acceleration.

At this point it seems helpful to recall that the sudden vanishing of the acceleration at the points q_n was motivated by the Lienard-Wiechert field solution for a moving charge. Just at the moment at which a particle acquires a uniform motion, the field around it starts to become the Coulomb solution for a uniformly moving charge. However, on the contrary, if a charge is initially in uniform motion, and a force is suddenly applied, the radiation reaction, which is accepted to be defined by the time derivative of the acceleration, would be infinite, if the particle is assumed to instantly get a finite value for the acceleration. Thus, the vanishing value of the acceleration fixed here just after the instant of kind s_n seems also a natural physical choice.

Let us present here the solution for the simple case of the one dimensional motion under a constant force which acts during a finite proper time interval. The analysis of this problem led Dirac, as discussed in his classic work on the theme, to the difficulties of the ALD equations in connection with the runaway solutions. Consider the particle at the beginning being at the origin of coordinates and that it moves with four velocity u_0^μ , at that starting instant, taken as the origin of the proper time $\tau = 0$. If the motion is in the x^1 direction, the four-force and four-velocity have the expressions

$$\begin{aligned}
u^\mu(\tau) &= \gamma(1, v, 0, 0), \\
\gamma &= \frac{1}{\sqrt{1 - v^2}}, \\
f^\mu(\tau) &= (F v, F, 0, 0), \\
u^\mu(\tau) f_\mu(\tau) &= 0, \\
u^\mu(\tau) u_\mu(\tau) &= 1.
\end{aligned}$$

The modified equations takes the explicit form

$$\begin{aligned}
F v \theta(\tau) \theta(T - \tau) &= m \frac{d^2 x^0(\tau)}{d\tau^2} - \kappa \frac{d^3 x^0(\tau)}{d\tau^3} - \kappa \left(\frac{d^2 x^0(\tau)}{d\tau^2} \frac{d^2 x_0(\tau)}{d\tau^2} - \frac{d^2 x^1(\tau)}{d\tau^2} \frac{d^2 x^1(\tau)}{d\tau^2} \right) \frac{dx^0(\tau)}{d\tau} + \kappa \frac{d^2 x^0(\tau)}{d\tau^2} \sigma(\tau), \\
F \theta(\tau) \theta(T - \tau) &= m \frac{d^2 x^1(\tau)}{d\tau^2} - \kappa \frac{d^3 x^1(\tau)}{d\tau^3} - \kappa \left(\frac{d^2 x^0(\tau)}{d\tau^2} \frac{d^2 x_0(\tau)}{d\tau^2} - \frac{d^2 x^1(\tau)}{d\tau^2} \frac{d^2 x^1(\tau)}{d\tau^2} \right) \frac{dx^1(\tau)}{d\tau} + \kappa \frac{d^2 x^1(\tau)}{d\tau^2} \sigma(\tau), \\
x^0(0) &= x^0(0) = 0, \\
\frac{dx^0(0)}{d\tau} &= \gamma, \quad \frac{dx^1(0)}{d\tau} = \gamma v, \\
\frac{d^2 x^0(0)}{d\tau^2} &= 0, \quad \frac{d^2 x^1(0)}{d\tau^2} = 0.
\end{aligned}$$

The parameters and initial conditions fixed were $\kappa = 1, m = 10, f = 10, v = 0.5$ and the proper time interval in which the force acts was selected as $T = 0.1$. The solution for the acceleration $\frac{d^2 x^1(\tau)}{d\tau^2}$ of the equations is illustrated in figure 1. The basic point to note is that the solution in the interval of proper time in which the force acts, is the same as the one for the ALD equation. However, when the force is removed the acceleration instantly vanishes. Thus, the only function of the added term in the ALD equation has been simply to facilitate the system to suddenly attain its uniform motion state, also satisfying the ALD equation. In the case of the ALD equation, discussed by Dirac in the reference [5], the acceleration, after the force is eliminated, is not vanishing due to the continuity of the acceleration, and more troublesome, it tends to grow without limit. The application of this particular solution to developing a discretization of a general motion, searching for avoiding the numerical development of instabilities leading to runaway solutions will be considered in extensions of this work.




FIG. 1: The figure shows the solution of the equations under study for the important case of a force which is suddenly applied remains constant during a time interval $T = 0.1$ and afterwards is also suddenly removed. The parameters of the equation fixed were $\kappa = 1, m = 10, f = 10, v = 0.5$

In this sense, the present discussion seems to be close connected with the analysis done in Ref. [8], which underlines, that in the zones of the evolution in which the forces have a non analytic behavior, the ALD equations can not be consistently derived and alternative equations should be obeyed. Therefore, the added term could represent an allowed modification of the dynamics, being able to solve the basic difficulties of the ALD equations, and which is determined by the physically plausible condition that the self-force should vanish instantaneously after the sudden (non-analytical) removal of the external force. This reasonable nature of this last condition is strongly supported by the fact that exactly after the instant of removal of the force, the wave-like propagation of the field, suggests that it starts to become exactly given by a boosted Coulomb field in sufficiently small neighborhood of the particle.

The solution of equations for more complex situations will be explored in the extension of the work. Of particular interest seems its application in improving numerical calculations of the solutions of the ALD equations, which had been systematically plagued with instabilities due to the existence of the nonphysical runaway solutions [8]. One specific problem will be the application to the one dimensional motion of a particle under the action of a repulsive Coulomb potential. In this task the runaway instabilities appear and their elimination had not been attained [12].

The authors would like to deeply acknowledge the helpful comments on the work received from Prof. Danilo Villarroel (Chile) and Prof. Luis Carlos Bassalo Crispino (UFPA). One of the authors (A.C) strongly acknowledges the support received from the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) of Brazil and the Postgraduation Programme in Physics (PPGF) of the Federal University of Pará at Belém, Pará (Brazil), in which this work was done, in the context of a CAPES External Visiting Professor Fellowship. The support also received by (A.C.) from the Caribbean Network on Quantum Mechanics, Particles and Fields (Net-35) of the ICTP Office of External Activities (OEA), the "Proyecto Nacional de Ciencias Básicas" (PNCB) of CITMA, Cuba is also very much acknowledged.

[1] H. A. Lorentz, *The theory of electrons*, Leipzig: Teubner, 1909 (2nd edition, 1916).

[2] M. Abraham, *Theorie der Elektrizitat, V. 2: Elektromagnetische, Theorie der Strahlung*, Leipzig: Teubner, 1905.

[3] H. Poincare, *On the dynamics of the electron*, Rendiconti del Circolo Matematico de Palermo 21, 129-176, 1906.

- [4] G. A. Schott, *Electromagnetic radiation*, Cambridge University Press, 1912.
- [5] P. A. M. Dirac, Proc. Roy. Soc. A 167, 148 (1938).
- [6] C. Teitelboim, D. Villarroel and Ch. G. van Weerth, Rev. Nuov. Cim. 3, 1, 1980.
- [7] F. Rohrlich, *Classical charged particles*, Redwood City, CA: Addison-Wesley, 1990.
- [8] A. Yaghjian, *Relativistic dynamics of a charged particle*, Lecture notes in Physics, m11, Springer Verlag, New York, Berlin, Heidelberg, 1992.
- [9] H. Spohn, *Dynamics of charged particles and their radiation field*, Cambridge University Press, Cambridge, 2004.
- [10] L. D. Landau, and E. M. Lifshitz, *The classical theory of fields*, Addison-Wesley, London 1959.
- [11] E. M. Purcell, *Electricidade e Magnetismo, Curso de Física de Berkeley*, V. 2, Edgar Blucher, Sao Paulo 1973.
- [12] , D. Vogt and P. S. Letelier, Gen. Rel. Grav. 35, 2261, 2003.