# **REGULARITY OF KÄHLER-RICCI FLOW**

\*GANG TIAN AND \*\*ZHENLEI ZHANG

ABSTRACT. In this short note we announce a regularity theorem for Kähler-Ricci flow on a compact Fano manifold (Kähler manifold with positive first Chern class) and its application to the limiting behavior of Kähler-Ricci flow on Fano 3-manifolds. Moreover, we also present a partial  $C^0$  estimate to the Kähler-Ricci flow under the regularity assumption, which extends previous works on Kähler-Einstein metrics and shrinking Kähler-Ricci solitons (cf. [17], [11], [20], [15]). The detailed proof will appear in [23].

#### 1. INTRODUCTION

Let M be a Fano *n*-manifold and  $g_0$  be any Kähler metric with Kähler class  $2\pi c_1(M)$ . Consider the normalized Kähler-Ricci flow:

(1.1) 
$$\frac{\partial g}{\partial t} = g - \operatorname{Ric}(g), \quad g(0) = g_0.$$

It was proved in [1] that (1.1) has a global solution g(t) for  $t \ge 0$ . The main problem is to understand the limit of g(t) as t tends to  $\infty$ .

By Perelman's non-collapsing result [12], there is a uniform constant  $\kappa = \kappa(g(0)) > 0$  satisfying:

(1.2) 
$$\operatorname{vol}_{g(t)}(B_{g(t)}(x,r)) \ge \kappa r^{2n}, \quad \forall t \ge 0, r \le 1.$$

Since the volume stays the same along the Kähler-Ricci flow, this implies that for any sequence  $t_i \to \infty$ , by taking a subsequence if necessary,  $(M, g(t_i))$  converge to a limiting length space  $(M_{\infty}, d)$  in the Gromov-Harsdorff topology:

(1.3) 
$$(M, g(t_i)) \xrightarrow{a_{GH}} (M_{\infty}, d)$$

The question is the regularity of  $M_{\infty}$ . A desirable picture is given in the following folklore conjecture <sup>1</sup>.

**Conjecture 1.1** ([18], also see [9]). (M, g(t)) converges (at least along a subsequence) to a shrinking Kähler-Ricci soliton with mild singularities.

<sup>\*</sup>The first author is supported in part by a National Science Foundation grant.

<sup>\*\*</sup>The second author is supported by a grant of Beijing Municipal Commission of Education.

<sup>&</sup>lt;sup>1</sup>It is often referred as the Hamilton-Tian conjecture (see [18]).

Here, "mild singularities" may be understood in two ways: (i) A singular set of codimension at least 4, and (ii) a singular set of a normal variety. By extending the partial  $C^0$ -estimate conjecture [19] to Kähler-Ricci flow, one can show that these two approaches are actually equivalent (see Section 3 or [23]).

As pointed out by the second named author, this conjecture implies the Yau-Tian-Donaldson conjecture, in the case of Fano manifolds. The conjecture states that if a Fano manifold M admits a Kähler-Einstein metrics if it is K-stable. Recently, solutions were provided for this conjecture in the case of Fano manifolds ([20], also see [6, 7, 8]).

### 2. Regularity of Kähler-Ricci flow

Let M be a Fano *n*-manifold and g(t) a normalized Kähler-Ricci flow in the Kähler class  $2\pi c_1(M)$ . Let  $(M_{\infty}, d)$  be a sequence limit of the Kähler-Ricci flow as phrased in (1.3). The main regularity result is the following theorem:

**Theorem 2.1.** Suppose that for some uniform p > n and  $\Lambda < \infty$ ,

(2.1) 
$$\int_{M} |Ric(g(t))|^{p} dv_{g(t)} \leq \Lambda.$$

Then the limit  $M_{\infty}$  is smooth outside a closed subset S of (real) codimension  $\geq 4$ and d is induced by a smooth Kähler-Ricci soliton  $g_{\infty}$  on  $M_{\infty} \setminus S$ . Moreover,  $g(t_i)$ converge to  $g_{\infty}$  in the  $C^{\infty}$ -topology outside S.<sup>2</sup>

The proof of the theorem relies on Perelman's Pseudolocality theorem [12] of Ricci flow and a regularity theory for manifolds with  $L^p$  bounded Ricci curvature (*p* bigger than half dimension) and uniformly local volume noncollapsing condition as in (1.2). This is a generalization of the regularity theory of Cheeger-Colding [2, 3, 4] and Cheeger-Colding-Tian [5]. The proof can be carried out following the lines given in these papers under the framework established by Petersen-Wei [13, 14] on the geometry of manifolds with integral bounded Ricci curvature. Note that due to an example of Yang [25], the uniformly volume noncollapsing condition (1.2) can not be replaced by a lower bound of total volume or local volume of metric balls of a definite size.

We shall show in [23] that there is a uniform  $L^4$  bound on the Ricci curvature along the Kähler-Ricci flow on any Fano manifold. Therefore, by the above regularity result, we have

**Corollary 2.2.** Conjecture 1.1, i.e., the Hamilton-Tian conjecture, holds for dimension  $n \leq 3$ .

In the case of Del-Pezzo surfaces, Conjecture 1.1 follows from [24] and [10].

 $<sup>^{2}</sup>$ The convergence with these properties is usually referred as the convergence in the Cheeger-Gromov topology, see [17] for instance.

## 3. Partial $C^0$ estimate of Kähler-Ricci flow

The partial  $C^0$  estimate of Kähler-Einstein manifolds plays the key role in Tian's program to resolve the Yau-Tian-Donaldson conjecture, see [17], [18], [19], [11] and [20] for examples. An extension of the partial  $C^0$  estimate to shrinking Kähler-Ricci solitons was given in [15]. These works are based on the compactness of Cheeger-Colding-Tian [5] and its generalizations to solitons by [22]. We shall generalize these to the Kähler-Ricci flow on Fano manifolds in [23] under the regularity assumption of the limit  $M_{\infty}$ .

Let u(t) denote the Ricci potentials of the Kähler-Ricci flow g(t) which satisfy

(3.1) 
$$Ric(g(t)) + \partial \partial u(t) = g(t).$$

The Hermitian metrics  $\tilde{g}(t) = e^{-\frac{1}{n}u(t)}g(t)$  have  $\omega(t)$ , the Kähler forms of g(t), as their Chern curvature forms. Let H(t) be the induced metric on  $K_M^{-l}$ , the *l*-th power of the anti-canonical bundle  $(l \ge 1)$ , and D be the Chern connection of H(t) on  $K_M^{-l}$ .

Let  $\nabla$  and  $\overline{\nabla}$  denote the (1,0) and (0,1) part of the Levi-Civita connection respectively. Then, at any time t, we have the Bochner type formula for  $\sigma \in H^0(M, K_M^{-l})$ 

$$(3.2) \qquad \triangle |\nabla\sigma|^2 = |\nabla\nabla\sigma|^2 + |\bar{\nabla}\nabla\sigma|^2 - ((n+2)l-1)|\nabla\sigma|^2 - \langle\partial\bar{\partial}u(\nabla\sigma,\cdot),\nabla\sigma\rangle$$

and the Weitzenböch type formulas for  $\xi \in C^{\infty}(M, T^{1,0}M \otimes K_M^{-l})$ 

where  $\Delta_{\bar{\partial}}$  is the Hodge Laplacian of  $\bar{\partial}$ . By [26] there is a uniform bound of Sobolev constant under the Kähler-Ricci flow. Then apply the Moser iteration one can prove the gradient estimate to  $\sigma \in H^0(M, K_M^{-l})$  and  $L^2$  estimate to solutions  $\bar{\partial}\vartheta =$  $\xi \in C^{\infty}(M, T^{1,0}M \otimes K_M^{-l})$ ; see Lemmas 4.1 and 5.4 of [20]. The gradient estimate implies dim  $H^0(M, K_M^{-l}) \leq N_l$  uniformly at any time t. We remark that Perelman's  $C^1$  estimate to u(t) [16] will be used in the iteration arguments to cancel the bad terms containing  $\partial \bar{\partial} u(t)$ .

Now, let  $\{s_{t,l,i}\}_{i=1}^{N_{t,l}}$  be an orthonormal basis of  $H^0(M, K_M^{-l})$  with respect to the  $L^2$  norm defined by H(t) and Riemannian volume form, and put

(3.5) 
$$\rho_{t,l}(x) = \sum_{i=1}^{N_{t,l}} |s_{t,l,i}|_H^2(x), \quad \forall x \in M.$$

By using arguments similar to those in [11] or [20], we can prove

**Theorem 3.1** (Partial  $C^0$  estimate). If  $(M, g(t_i)) \xrightarrow{d_{GH}} (M_\infty, g_\infty)$  as phrased in Theorem 2.1, then the partial  $C^0$  estimate

(3.6) 
$$\inf_{t_j} \inf_{x \in M} \rho_{t_j,l}(x) > 0$$

holds for a sequence of  $l \to \infty$ .

A direct corollary of this is to refine the regularity in Theorem 2.1.

**Theorem 3.2.** Suppose  $(M, g(t_i)) \xrightarrow{d_{GH}} (M_{\infty}, g_{\infty})$  as phrased in Theorem 2.1. Then  $M_{\infty}$  is a normal projective variety and S is a subvariety of complex codimension at least 2.

Finally, let us indicate how to deduce the Yau-Tian-Donaldson conjecture from the Hamilton-Tian conjecture. Suppose M is K-stable as defined in [18]. Then, under the Kähler-Ricci flow g(t), we get a shrinking Kähler-Ricci soliton. This, together with the uniqueness theorem on shrinking solitons, we can conclude that the Lie algebra of holomorphic vector fields on  $M_{\infty}$  is reductive. Then the K-stability implies the vanishing of Futaki invariant of  $M_{\infty}$ , consequently, the limit  $(M_{\infty}, g_{\infty})$ is Kähler-Einstein. If  $M_{\infty}$  is not biholomorphic to M, then the eigenspaces of the first eigenvalues of  $-\Delta_{g(t)} + g^{i\bar{j}}(t)\partial_i u(t)\partial_{\bar{j}}$  will converge to a subspace of potential functions on  $M_{\infty}$  whose complex gradients are nontrivial holomorphic vector fields, cf. [27]. These vector fields induce the required degeneration of M to  $M_{\infty}$ , with vanishing Futaki invariants. This gives a contradiction to the K-stability of M. So we have

**Theorem 3.3.** Suppose M is K-stable. If  $(M, g(t_i)) \xrightarrow{d_{GH}} (M_{\infty}, g_{\infty})$  as phrased in Theorem 2.1, then M coincides with  $M_{\infty}$  and admits a Kähler-Einstein  $g_{\infty}$ .

In view of the regularity of low dimensional Kähler-Ricci flow in §2 we have

**Corollary 3.4.** The Yau-Tian-Donaldson conjecture holds for dimension  $n \leq 3$ .

#### References

- H.D. Cao, Deformation of Kähler metrics to Kähler-Einstein metrics on compact Kähler manifolds, Invent. Math., 81 (1985), 359-372.
- [2] J. Cheeger and T. H. Colding, Lower bounds on the Ricci curvature and the almost rigidity of warped products, Annal. Math., 144 (1996), 189-237.
- [3] J. Cheeger and T. H. Colding, On the structure of spaces with Ricci curvature bounded below I, J. Diff. Geom., 46 (1997), 406-480.
- [4] J. Cheeger and T. H. Colding, On the structure of spaces with Ricci curvature bounded below II, J. Diff. Geom.,
- [5] J. Cheeger, T. H. Colding and G. Tian, On the singularities of spaces with bounded Ricci curvature, Geom. Funct. Anal., 12 (2002), 873-914.
- [6] X.X. Chen, S. Donaldson and S. Sun, Kahler-Einstein metrics on Fano manifolds, I: approximation of metrics with cone singularities, arXiv:1211.4566
- [7] X.X. Chen, S. Donaldson and S. Sun, Kahler-Einstein metrics on Fano manifolds, II: limits with cone angle less than 2π, arXiv:1212.4714
- [8] X.X. Chen, S. Donaldson and S. Sun, Kahler-Einstein metrics on Fano manifolds, III: limits as cone angle approaches 2 and completion of the main proof, arXiv:1302.0282
- [9] X.X. Chen and G. Tian, Ricci flow on Kähler-Einstein manifolds, Duke Math. Jour., 131 (2006), 17-73.

- [10] X.X. Chen and B. Wang, Space of Ricci flows I, Comm. Pure Appl. Math., 65 (2012), 1399-1457.
- S. Donaldson and S. Sun, Gromov-Hausdorff limits of Kähler manifolds and algebraic geometry, arXiv:1206.2609v1
- [12] G. Perelman, The entropy formula for the Ricci flow and its geometric applications, arXiv:math.DG/0211159.
- [13] P. Petersen and G.F. Wei, Relative volume comparison with integral curvature bounds, Geom. Funct. Anal., 7 (1997), 1031-1045.
- [14] P. Petersen and G.F. Wei, Analysis and geometry on manifolds with integral Ricci curvature bounds. II, Trans. AMS., 353 (2001), 457-478.
- [15] D. H. Phong, J. Song and J. Sturm, Degeneration of Kähler-Ricci solitons on Fano manifolds, arXiv:1211.5849v1
- [16] N. Sesum and G. Tian, Bounding scalar curvature and diameter along the Kähler-Ricci flow (after Perelman), J. Inst. Math. Jussiu, 7 (2008), 575-587.
- [17] G. Tian, On Calabi's conjecture for complex surfaces with positive first Chern class, Invent. Math., 101 (1990), 101-172.
- [18] G. Tian, Kähler-Einstein metrics with positive scalar curvature, Invent. Math., 130 (1997), 1-39.
- [19] G. Tian, Existence of Einstein metrics on Fano manifolds, Metric and Differential Geometry, Prog. Math., 297 (2012), 119-159.
- [20] G. Tian, K-stability and Kähler-Einstein metrics, arXiv:1211.4669
- [21] G. Tian, S.J. Zhang, Z.L. Zhang and X.H. Zhu, Supremun of Perelman's entropy and Kähler-Ricci flow on a Fano manifold, arXiv:1107.4018.
- [22] G. Tian and Z.L. Zhang, Degeneration of Kähler-Ricci solitons, Intern. Math. Res. Notices, 2012, 957-985.
- [23] G. Tian and Z.L. Zhang, Long-time limits of the Kähler-Ricci flow, preprint.
- [24] G. Tian and X.H. Zhu Convergence of K?hler-Ricci flow. J. Amer. Math. Soc. 20 (2007), 675-699.
- [25] D. Yang, Convergence of Riemannian manifolds with integral buonds on curvature I, Ann. scient. Éc. Norm. Sup., 25 (1992), 77-105.
- [26] Q. S. Zhang, A uniform Sobolev inequality under Ricci flow, Int. Math. Res. Not., 2007, 1-12.
- [27] Z.L. Zhang, Kähler-Ricci flow on Fano manifolds with vanished Futaki invariants, Math. Res. Lett., 18 (2011), 969-982.

BICMR, PEKING UNIVERSITY, YIHEYUAN ROAD 5, BEIJING 100871, CHINA

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON NJ 08544, USA

*E-mail address*: gtian@math.pku.edu.cn

School of Mathematics, Capital Normal University, Xisanhuan North Road 105, Beijing 100048, China

E-mail address: zhleigo@yahoo.com.cn