

MEASURE CONTRACTION PROPERTIES OF SASAKIAN MANIFOLDS

PAUL W. Y. LEE, CHENGBO LI, AND IGOR ZELENKO

ABSTRACT. Measure contraction properties are generalizations of the notion of Ricci curvature lower bounds in Riemannian geometry to more general metric measure spaces. In this paper, we give sufficient conditions for a Sasakian manifold equipped with a natural sub-Riemannian distance to satisfy these properties. Moreover, the sufficient conditions are defined by the Tanaka-Webster curvature. This generalizes the earlier work in [2] for the three dimensional case and in [14] for the Heisenberg group. To obtain our results we use the intrinsic Jacobi equations along sub-Riemannian extremals, coming from the theory of canonical moving frames for curves in Lagrangian Grassmannians [17, 18]. The crucial new tool here is a certain decoupling of the corresponding matrix Riccati equation. It is also worth pointing out that our measure contraction properties are sharp: the corresponding inequalities become equalities for the corresponding homogeneous models in the considered class of sub-Riemannian structures.

1. INTRODUCTION

In recent years, there are lots of efforts in generalizing the notion of Ricci curvature lower bounds in Riemannian geometry and its consequences to more general metric measure spaces. One of them is the work of [20, 21, 26, 27] where the notion of curvature-dimension conditions was introduced. These conditions are generalizations of Ricci curvature lower bounds to length spaces equipped with a measure (length spaces are metric spaces on which the notion of geodesics is defined). In [24], it was shown that the curvature-dimension conditions coincide

Date: February 28, 2019.

2010 Mathematics Subject Classification. 53C17, 53D10, 70G45, 34C10, 53C25, 53C55.

Key words and phrases. Sub-Riemannian metrics, Measure Contraction properties, contact manifolds, Jacobi equations and Jacobi curves, curves in Lagrangian Grassmannians, symplectic invariants, conjugate points, Matrix Riccati equation, Bonnet-Myers theorems.

The first author's research was supported by the Research Grant Council of Hong Kong (RGC Ref. No. CUHK404512). The second author was supported in part by the National Natural Science Foundation of China (Grant No. 11201330).

with the pre-existing notion of Ricci curvature lower bounds in the case of Finsler manifolds.

On the contrary, it was shown in [14] that the curvature-dimension conditions defined using the theory of optimal transportation are not satisfied on the Heisenberg group, the simplest sub-Riemannian manifold (Note however that a type of curvature-dimension conditions were defined in [6, 7] using a sub-Riemannian version of the Bochner formula). It was also shown in [14] that the Heisenberg group satisfies another generalization of Ricci curvature lower bounds to length spaces called measure contraction properties [26, 27, 23].

Measure contraction properties are essentially defined by the rate of contraction of volume along geodesics inspired by the classical Bishop volume comparison theorem. In the Riemannian case, the measure contraction property $\mathcal{MCP}(k, n)$ is equivalent to the conditions that the Ricci curvature is bounded below by k and the dimension is bounded above by n . In [14], it was shown that the left-invariant sub-Riemannian metric on the Heisenberg group of dimension $2n + 1$ satisfies the condition $\mathcal{MCP}(0, 2n + 3)$. Such sub-Riemannian metrics can be regarded as the flat one among all sub-Riemannian metrics on Sasakian manifolds of the same dimension.

The next natural task is to study the measure contraction property for general (curved) sub-Riemannian metrics on Sasakian manifolds and, in particular, to understand what differential invariants of such metrics are important for their measure contraction property. A natural way of doing this is to analyze the Jacobi equation along a sub-Riemannian extremal. In order to write the Jacobi equation intrinsically one needs first to construct a connection canonically associated with a geometric structure. The construction of such connection is known in several classical cases such as the Levi-Civita connection for Riemannian metrics, the Tanaka-Webster connection for a special class of sub-Riemannian contact metrics associated with CR structures ([28, 30]) or its generalization, the Tanno connection, to more general class of sub-Riemannian contact metrics associated with (non-integrable) almost CR structures ([29]). However, these constructions use specific properties of the geometric structures under consideration and it is not clear how to generalize them to more general sub-Riemannian structures.

An alternative approach for obtaining intrinsic Jacobi equation *without preliminary construction of a canonical connection* was initiated in [1] and further developed in [5, 17, 18]. In this approach one replaces the Jacobi equation along an extremal by a special and a priori intrinsic curve of Lagrangian subspaces in a linear symplectic space, i.e. a curve

in a Lagrangian Grassmannian. This curve is defined up to the natural action of the linear symplectic group. It contains all information about the space of Jacobi fields along the extremal and therefore it is called the *Jacobi curve of the extremal*.

By analogy with the classical Frenet-Serret frame for a curve in an Euclidean space, one can construct a bundle of canonical moving symplectic frames for a curve in a Lagrangian Grassmannian satisfying very general assumptions [17, 18]. The structure equations for these moving frames can be considered as the intrinsic Jacobi equations, while the nontrivial entries in the matrices of these structure equations give the invariants of the original geometric structures. These invariants can be used in principal for obtaining various comparison type results including the measure contraction properties. Although the construction of these invariants is algorithmic, to express them explicitly in terms of the original geometric structure is not an easy task already for sub-Riemannian contact case ([2, 19, 9]). Besides, in contrast to the Riemannian case, the level sets of a sub-Riemannian Hamiltonian are not compact. Therefore, to control the bounds for the symplectic invariants of the Jacobi curves along extremals, additional assumptions has to be imposed.

For the first time the scheme based on the geometry of Jacobi curves was used in the study of the measure contraction properties in [2] (see also [13, 10, 3] for closely related results), where general three dimensional contact sub-Riemannian manifolds were treated. In particular, it was shown there that the measure contraction properties $\mathcal{MCP}(0, 5)$ are characterized by the Tanaka-Webster sectional curvature of the planes which are the fibers of the contact distribution. More precisely, a sub-Riemannian metric naturally associated with a three-dimensional Sasakian manifold satisfies $\mathcal{MCP}(0, 5)$ if and only if the corresponding Tanaka-Webster section curvature is bounded below by 0.

Moreover, the generalized measure contraction properties $\mathcal{MCP}(k; 2, 3)$ were defined there. It was shown that a three-dimensional Sasakian manifold satisfies this condition if and only if the aforementioned Tanaka-Webster sectional curvature is bounded below by k .

In this paper, we generalize the results in [14, 2] to Sasakian manifolds equipped with a natural sub-Riemannian structure. We introduce new generalized measure contraction properties $\mathcal{MCP}(k_1, k_2; N - 1, N)$ and discuss when a Sasakian manifold equipped with the sub-Riemannian structure mentioned above satisfies them.

To obtain these results we analyze the matrix Riccati equation associated with the structure equation for the canonical moving frame of the Jacobi curves (Lemma 4.1 and section 5 below) on the basis of

comparison theorems for matrix Riccati equations ([25, 16]) and use the expressions for the symplectic invariants of the Jacobi curves (see section 5 below) based on the calculation in [9].

The key new observation, which was first appeared in an earlier version of this paper, is the decoupling of these equations after taking the traces of appropriate blocks (as in the proof of Lemma 5.3 below): the coupled equations after taking the traces yields to decoupled inequalities, which leads to the desired estimates, involving the Tanaka-Webster curvature of the manifold. Surprisingly, the passage from equations to inequalities does not affect the sharpness of the estimates: our measure contraction properties for the considered class of sub-Riemannian structures cannot be improved because the corresponding inequalities turn to equalities for the corresponding homogeneous models (see Corollaries 2.5 and 2.6 below).

Next, we state the condition $\mathcal{MCP}(k_1, k_2; N-1, N)$ and some consequences of the main results in more detail. Let M be a sub-Riemannian manifold (see Section 2 for a discussion on some basic notions in sub-Riemannian geometry). For simplicity, we assume that M satisfies the following property: given any point x_0 in M , there is a set of Lebesgue measure zero such that any point outside the set is connected to x_0 by a unique length minimizing sub-Riemannian geodesic. By the result in [8], this is satisfied by all contact sub-Riemannian manifolds.

Let $t \mapsto \varphi_t(x)$ be the unique geodesic starting from x and ending at x_0 . This defines a 1-parameter family of Borel maps. Let d be the sub-Riemannian distance and let μ be a Borel measure. The following is the original measure contraction property studied in [26, 27, 23]:

A metric measure space (M, d, μ) satisfies $\mathcal{MCP}(0, N)$ if

$$\mu(\varphi_t(U)) \geq (1-t)^N \mu(U)$$

for each point x_0 and each Borel set U .

Next, we introduce the new generalized measure contraction properties $\mathcal{MCP}(k_1, k_2; N-1, N)$: A metric measure space (M, d, μ) satisfies $\mathcal{MCP}(k_1, k_2; N-1, N)$ if, for each point x_0 and each Borel set U ,

$$\mu(\varphi_t(U)) \geq \int_U \frac{(1-t)^{N+2} \mathcal{M}_1(k_1 d^2(x, x_0), t) \mathcal{M}_2^{N-3}(k_2 d^2(x, x_0), t)}{\mathcal{M}_1(k_1 d^2(x, x_0), 0) \mathcal{M}_2^{N-3}(k_2 d^2(x, x_0), 0)} d\mu(x),$$

where $\mathcal{D}(k, t) = \sqrt{|k|}(1-t)$,

$$\mathcal{M}_1(k, t) = \begin{cases} \frac{2-2\cos(\mathcal{D}(k,t))-\mathcal{D}(k,t)\sin(\mathcal{D}(k,t))}{\mathcal{D}(k,t)^4} & \text{if } k > 0 \\ \frac{1}{12} & \text{if } k = 0 \\ \frac{2-2\cosh(\mathcal{D}(k,t))+\mathcal{D}(k,t)\sinh(\mathcal{D}(k,t))}{\mathcal{D}(k,t)^4} & \text{if } k < 0, \end{cases}$$

and

$$\mathcal{M}_2(k, t) = \begin{cases} \frac{\sin(\mathcal{D}(k, t))}{\mathcal{D}(k, t)} & \text{if } k > 0 \\ 1 & \text{if } k = 0 \\ \frac{\sinh(\mathcal{D}(k, t))}{\mathcal{D}(k, t)} & \text{if } k < 0. \end{cases}$$

Note, in particular, that $\mathcal{MCP}(0, 0; N-1, N)$ is the same as $\mathcal{MCP}(0, N+2)$. If $k_1 \geq 0$ and $k_2 \geq 0$, then $\mathcal{MCP}(k_1, k_2; N-1, N)$ implies $\mathcal{MCP}(0, N+2)$. The reason for the notations in the conditions $\mathcal{MCP}(k_1, k_2; N-1, N)$ is clarified by Theorem 1.1 below.

Next, we state a simple consequence of the main results. For this, let M be a Sasakian manifold of dimension $2n+1$ equipped with a contact form α_0 and a Riemannian metric $\langle \cdot, \cdot \rangle$ (see the next section for a detail discussion of Sasakian manifold and the corresponding sub-Riemannian structure). Let μ be the corresponding Riemannian volume form and let d_{CC} be the sub-Riemannian distance. Here the distribution is defined by $\mathcal{D} = \ker \alpha_0$ and the sub-Riemannian metric is given by the restriction of the Riemannian one on \mathcal{D} .

Theorem 1.1. *Assume that the Tanaka-Webster curvature \mathbf{Rm}^* of the Sasakian manifold M satisfies*

$$\langle \mathbf{Rm}^*(Jv, v)v, Jv \rangle \geq k_1$$

and

$$\sum_{i=1}^{2n-2} \langle \mathbf{Rm}^*(w_i, v)v, w_i \rangle \geq (2n-2)k_2.$$

for all unit tangent vectors v in \mathcal{D} and for all orthonormal frame in $\mathcal{D} \cap \{v, Jv\}^\perp$. Then the metric measure space (M, d_{CC}, μ) satisfies $\mathcal{MCP}(k_1, k_2; 2n, 2n+1)$, where d is the sub-Riemannian distance of M .

In particular, we recover the following result in [14].

Theorem 1.2. [14] *The Heisenberg group of dimension $2n+1$ equipped with the standard sub-Riemannian distance d_{CC} and the Lebesgue measure μ satisfies $\mathcal{MCP}(0, 0; 2n, 2n+1) = \mathcal{MCP}(0, 2n+3)$.*

We also have

Theorem 1.3. *The complex Hopf fibration equipped with the above sub-Riemannian distance d_{CC} and measure μ satisfies the condition $\mathcal{MCP}(4, 1; 2n, 2n+1)$. In particular, it satisfies $\mathcal{MCP}(0, 2n+3)$.*

We also remark that the estimates for the proof of Theorem 1.2 and 1.3 are sharp (see Corollary 2.5 and 2.6 for more detail).

Note that the condition $\mathcal{MCP}(0, N)$ implies the volume doubling property of μ and a local Poincaré inequality (see [26, 27, 23, 2], see also [6, 7] for an alternative approach to the following results).

Corollary 1.1. (*Doubling*) *Assume that the assumption of Theorem 1.1 holds. Then there is a constant $C > 0$ such that*

$$\mu(B_x(2R)) \leq C\mu(B_x(R))$$

for all x in M and all $R > 0$, where $B_x(R)$ is the sub-Riemannian ball of radius R centered at x .

Corollary 1.2. (*Poincaré inequality*) *Assume that the assumption of Theorem 1.1 holds. Then there is a constant $C > 0$ such that*

$$\begin{aligned} & \frac{1}{\text{vol}(B_x(R))} \int_{B_x(R)} |f(x) - \langle f \rangle_{B_x(R)}| d\text{vol}(x) \\ & \leq C \frac{R}{\text{vol}(B_x(2R))} \int_{B_x(2R)} |\nabla_{\text{hor}} f| d\text{vol}(x), \end{aligned}$$

where $\langle f \rangle_{B_x(R)} = \frac{1}{\text{vol}(B_x(R))} \int_{B_x(R)} f(x) d\text{vol}(x)$ and $\nabla_{\text{hor}} f$ is the horizontal gradient of f which is the projection of the Riemannian gradient onto the distribution \mathcal{D} .

By restriction the Riemannian Hessian of a function f to the distribution \mathcal{D} and taking the trace, we obtain the sub-Laplacian $\Delta_{\text{hor}} f$ of f . By combining the result in [11], we also obtain

Corollary 1.3. (*Harnack inequality*) *Assume that the assumption of Theorem 1.1 holds. Then there is a constant $C > 0$ such that any positive solution to the equation $\Delta_{\text{hor}} f = 0$ on $B_x(R)$ satisfies*

$$\sup_{B_x(R/2)} f \leq C \inf_{B_x(R/2)} f.$$

Corollary 1.4. (*Liouville theorem*) *Assume that the assumption of Theorem 1.1 holds. Then any non-negative solution to the equation $\Delta_{\text{hor}} f = 0$ on M is a constant.*

In the earlier versions of this paper, we proved the same results for contact sub-Riemannian manifolds with a transversal symmetry. While in the present version, we use the calculations in [9] appeared later than the earlier version of this paper to rewrite the result in the Sasakian case with essentially the same proof. The new calculations are in some sense more direct and the statements of results are simpler. We also remark that, unlike the Riemannian case, Bishop volume comparison theorem and measure contraction properties are very different in the

sub-Riemannian case. This is because any sub-Riemannian ball contains cut points.

In the next section, some basic notions in sub-Riemannian geometry will be recalled and the main results of the paper will be stated. The rest of the sections will be devoted to the proof of the main results.

2. SUB-RIEMANNIAN STRUCTURES ON SASAKIAN MANIFOLDS

In this section, we recall various notions on sub-Riemannian structures of Sasakian manifolds which are needed. A sub-Riemannian manifold is a triple $(M, \mathcal{D}, \langle \cdot, \cdot \rangle)$, where M is a manifold of dimension N , \mathcal{D} is a sub-bundle of the tangent bundle TM , and $\langle \cdot, \cdot \rangle$ is a smoothly varying inner product defined on \mathcal{D} . The sub-bundle \mathcal{D} and the inner product $\langle \cdot, \cdot \rangle$ are commonly known as a distribution and a sub-Riemannian metric, respectively. A curve $\gamma(\cdot)$ is horizontal if $\dot{\gamma}(t)$ is contained in \mathcal{D} for almost every t . The length $l(\gamma)$ of a horizontal curve γ can be defined as in the Riemannian case:

$$l(\gamma) = \int_0^1 |\dot{\gamma}(t)| dt.$$

Assume that the distribution \mathcal{D} satisfies the following bracket generating or Hörmander condition: the sections of \mathcal{D} and their iterated Lie brackets span each tangent space. Under this assumption and that the manifold M is connected, the Chow-Rashevskii Theorem (see [22]) guarantees that any two given points on the manifold M can be connected by a horizontal curve. Therefore, we can define the sub-Riemannian (or Carnot-Carathéodory) distance d_{CC} as

$$(2.1) \quad d_{CC}(x_0, x_1) = \inf_{\gamma \in \Gamma} l(\gamma),$$

where the infimum is taken over the set Γ of all horizontal paths $\gamma : [0, 1] \rightarrow M$ which connect x_0 with x_1 : $\gamma(0) = x_0$ and $\gamma(1) = x_1$. The minimizers of (2.1) are called length minimizing geodesics (or simply geodesics). As in the Riemannian case, reparametrizations of a geodesic is also a geodesic. Therefore, we assume that all geodesics have constant speed.

In this paper, we will focus on Sasakian manifolds equipped with a natural sub-Riemannian structure. First, we recall that a distribution \mathcal{D} is contact if it is given by the kernel of a 1-form α_0 , called a contact form, defined by the condition that the restriction of $d\alpha_0$ to \mathcal{D} is non-degenerate. A Sasakian manifold M is a contact manifold equipped with a contact form α_0 , a vector field v_0 , a $(1, 1)$ -tensor J ,

and a Riemannian metric $\langle \cdot, \cdot \rangle$ such that

$$\begin{aligned}\alpha_0(v_0) &= 1, \\ J^2v &= -v + \alpha_0(v)v_0, \\ d\alpha_0(v, w) &= \langle v, Jw \rangle,\end{aligned}$$

and the Nijenhuis tensor

$$[J, J](v, w) := J^2[v, w] + [Jv, Jw] - J[Jv, w] - J[v, Jw]$$

satisfies

$$[J, J] = -d\alpha_0(v, w)v_0.$$

The sub-Riemannian metric is defined simply by restricting the Riemannian one on the distribution $\mathcal{D} := \ker \alpha_0$.

The Tanaka connection ∇^* is defined by

$$\nabla_X^* Y = \nabla_X Y + \frac{1}{2}\alpha_0(X)JY - \alpha_0(Y)\nabla_X v_0 + \nabla_X \alpha_0(Y)v_0$$

and the corresponding curvature \mathbf{Rm}^* is called Tanaka-Webster curvature

$$\mathbf{Rm}^*(X, Y)Z = \nabla_X^* \nabla_Y^* Z - \nabla_Y^* \nabla_X^* Z - \nabla_{[X, Y]}^* Z.$$

The simplest example of Sasakian manifold is the Heisenberg group. In this case, the manifold M is the $(2n+1)$ -dimensional Euclidean space \mathbb{R}^{2n+1} with coordinates $x_1, \dots, x_n, y_1, \dots, y_n, z$. The vector field v_0 is $v_0 = \partial_z$, and the contact form α_0 is $\alpha_0 = dz - \frac{1}{2}\sum_{i=1}^n x_i dy_i + \frac{1}{2}\sum_{i=1}^n y_i dx_i$. Let

$$X_i = \partial_{x_i} - \frac{1}{2}y_i \partial_z, \quad Y_i = \partial_{y_i} + \frac{1}{2}x_i \partial_z,$$

where $i = 1, \dots, n$. The Riemannian metric is defined in such a way that $\{X_1, \dots, X_n, Y_1, \dots, Y_n, \partial_z\}$ is orthonormal and the tensor J satisfies $J(X_i) = Y_i$, $J(Y_i) = -X_i$, $J(\partial_z) = 0$. Finally, the Tanaka-Webster curvature $\mathbf{Rm}^* = 0$ on \mathcal{D} in this case. Remark that the Riemannian volume in this case coincides with the $(2n+1)$ -dimensional Lebesgue measure \mathcal{L}^{2n+1} .

Corollary 2.5. *The Heisenberg group satisfies*

$$\mathcal{L}^{2n+1}(\varphi_t(U)) = \int_U \frac{(1-t)\mathcal{M}_1(\mathbf{k}_1(x), t)\mathcal{M}_2^{2n-2}(\mathbf{k}_2(x), t)}{\mathcal{M}_1(\mathbf{k}_1(x), 0)\mathcal{M}_2^{2n-2}(\mathbf{k}_2(x), 0)} d\mathcal{L}^{2n+1}(x)$$

for any Borel set U , where

$$\begin{aligned}\mathbf{k}_1(x) &= \mathbf{f}(x)^2(v_0 \mathbf{f}(x))^2, & \mathbf{k}_2(x) &= \frac{\mathbf{f}(x)^2(v_0 \mathbf{f}(x))^2}{4}, \\ \mathbf{f}(\mathbf{x}) &= -\frac{1}{2}d_{CC}^2(\mathbf{x}, \mathbf{x}_0).\end{aligned}$$

In particular, the metric measure space $(\mathbb{H}, d_{CC}, \mathcal{L}^{2n+1})$ satisfies the condition $\mathcal{MCP}(0, 0; 2n, 2n + 1) = \mathcal{MCP}(0, 2n + 3)$.

Another example which is relevant for us is the complex Hopf fibration. In this case, the manifold M is $S^{2n+1} = \{z \in \mathbb{C}^{n+1} \mid |z| = 1\}$. The vector field v_0 is given by $v_0 = 2 \sum_{i=1}^n (-y_i \partial_{x_i} + x_i \partial_{y_i})$ and the 1-form is $\alpha_0 = \frac{1}{2} \sum_{i=1}^n (x_i dy_i - y_i dx_i)$. The Riemannian metric coincides with the Euclidean one on \mathcal{D} and J coincides with standard complex structure on \mathcal{D} . The Tanaka-Webster curvature \mathbf{Rm}^* , in this case, satisfies

$$\langle Rm^*(JX, X)X, JX \rangle = 4|X|^4, \quad \langle Rm^*(Y, X)X, Y \rangle = |X|^2|Y|^2$$

for all X in \mathcal{D} and Y in $\{X, JX\}^\perp \cap \mathcal{D}$. Let μ be the Riemannian volume.

Corollary 2.6. *The complex Hopf fibration satisfies*

$$\mu(\varphi_t(U)) = \int_U \frac{(1-t)\mathcal{M}_1(\mathbf{k}_1(x), t)\mathcal{M}_2^{2n-2}(\mathbf{k}_2(x), t)}{\mathcal{M}_1(\mathbf{k}_1(x), 0)\mathcal{M}_2^{2n-2}(\mathbf{k}_2(x), 0)} d\mu(x)$$

for any Borel set U , where

$$\mathbf{k}_1(x) = \mathfrak{f}(x)^2(4 + (v_0\mathfrak{f}(x))^2), \quad \mathbf{k}_2(x) = \frac{\mathfrak{f}(x)^2}{4}(4 + (v_0\mathfrak{f}(x))^2),$$

$$\mathfrak{f}(\mathbf{x}) = -\frac{1}{2}d_{CC}^2(\mathbf{x}, \mathbf{x}_0).$$

In particular, the metric measure space (M, d_{CC}, μ) satisfies the condition $\mathcal{MCP}(4, 1; 2n, 2n + 1)$ and hence $\mathcal{MCP}(0, 2n + 3)$.

Note that Corollary 2.5 and 2.6 show that the main result is sharp in the sense that all the inequalities in Theorem 1.1, including both assumptions and conclusions, are equality in the case of the Heisenberg group and the complex Hopf fibration.

3. SUB-RIEMANNIAN GEODESIC FLOWS AND CURVATURE ON SASAKIAN MANIFOLDS

In this section, we recall the definition of the sub-Riemannian geodesic flow and its connections with the contraction of measures appeared in [2, 3].

As in the Riemannian case, the (constant speed) minimizers of (2.1) can be found by minimizing the following kinetic energy functional

$$(3.2) \quad \inf_{\gamma \in \Gamma} \int_0^1 \frac{1}{2} |\dot{\gamma}(t)|^2 dt.$$

In the Riemannian case, the minimizers of (3.2) are given by the geodesic equation. In the sub-Riemannian case, the minimization problem

in (3.2) becomes a constrained minimization problem and it is more convenient to look at the geodesic flow from the Hamiltonian point of view in this case. For this, let $\mathbf{H} : T^*M \rightarrow \mathbb{R}$ be the Hamiltonian defined by the Legendre transform:

$$\mathbf{H}(\mathbf{x}, \mathbf{p}) = \sup_{\mathbf{v} \in \mathcal{D}} \left(\mathbf{p}(\mathbf{v}) - \frac{1}{2}|\mathbf{v}|^2 \right).$$

This Hamiltonian, in turn, defines a Hamiltonian vector field $\vec{\mathbf{H}}$ on the cotangent bundle T^*M which is a sub-Riemannian analogue of the geodesic equation. It is given, in the local coordinates $(x_1, \dots, x_N, p_1, \dots, p_N)$, by

$$\vec{\mathbf{H}} = \sum_{i=1}^N (\mathbf{H}_{p_i} \partial_{x_i} - \mathbf{H}_{x_i} \partial_{p_i}).$$

We assume, through out this paper, that the vector field $\vec{\mathbf{H}}$ defines a complete flow which is denoted by $e^{t\vec{\mathbf{H}}}$. In the Riemannian case, the minimizers of (3.2) are given by the projection of the trajectories of $e^{t\vec{\mathbf{H}}}$ to the manifold M . In the sub-Riemannian case, minimizers obtained this way are called normal geodesics and they do not give all the minimizers of (3.2) in general (see [22] for more detailed discussions on this). On the other hand, all minimizers of (3.2) are normal if the distribution \mathcal{D} is contact (see [22]).

Next, we discuss an analogue of the Jacobi equation in the above Hamiltonian setting. For this, let ω be the canonical symplectic form of the cotangent bundle T^*M . In local coordinates $(x_1, \dots, x_N, p_1, \dots, p_N)$, ω is given by

$$\omega = \sum_{i=1}^N dp_i \wedge dx_i.$$

Let $\pi : T^*M \rightarrow M$ be the canonical projection and let \mathbf{ver} the vertical sub-bundle of the cotangent bundle T^*M defined by

$$\mathbf{ver}_{(\mathbf{x}, \mathbf{p})} = \{v \in T_{(\mathbf{x}, \mathbf{p})}T^*M \mid \pi_*(v) = 0\}.$$

Recall that a n -dimensional subspace of a symplectic vector space is Lagrangian if the symplectic form vanishes when restricted to the subspace. Each vertical space $\mathbf{ver}_{(\mathbf{x}, \mathbf{p})}$ is a Lagrangian subspace of the symplectic vector space $T_{(\mathbf{x}, \mathbf{p})}T^*M$. Since the flow $e^{t\vec{\mathbf{H}}}$ preserves the symplectic form ω , it also sends a Lagrangian subspace to another Lagrangian one. Therefore, the following also forms a one-parameter family of Lagrangian subspaces contained in $T_{(\mathbf{x}, \mathbf{p})}T^*M$

$$(3.3) \quad \mathfrak{J}_{(\mathbf{x}, \mathbf{p})}(t) = e_*^{-t\vec{\mathbf{H}}}(\mathbf{ver}_{e^{t\vec{\mathbf{H}}}(\mathbf{x}, \mathbf{p})}).$$

This family defines a curve in the Lagrangian Grassmannian (the space of Lagrangian subspaces) of $T_{(\mathbf{x}, \mathbf{p})}T^*M$ and it is called the Jacobi curve at (\mathbf{x}, \mathbf{p}) of the flow $e^{t\tilde{\mathbf{H}}}$.

Assume that the distribution is contact. Then we have the following particular case of the results in [18, 19].

Theorem 3.1. *Assume that the distribution \mathcal{D} is contact. Then there exists a one-parameter family of bases*

$$\begin{aligned} E(t) &= (E_1(t), \dots, E_{2n+1}(t))^T \\ F(t) &= (F_1(t), \dots, F_{2n+1}(t))^T \end{aligned}$$

of the symplectic vector space $T_{(\mathbf{x}, \mathbf{p})}T^*M$ such that the followings hold for any t :

- (1) $\mathfrak{J}_{(\mathbf{x}, \mathbf{p})}(t) = \text{span}\{E_1(t), \dots, E_{2n+1}(t)\}$,
- (2) $\text{span}\{F_1(t), \dots, F_{2n+1}(t)\}$ is a family of Lagrangian subspaces,
- (3) $\omega(F_i(t), E_j(t)) = \delta_{ij}$,
- (4) $\omega(F_i(t), F_j(t)) = \omega(E_i(t), E_j(t)) = 0$,
- (5) $\dot{E}(t) = C_1E(t) + C_2F(t)$,
- (6) $\dot{F}(t) = -R(t)E(t) - C_1^T F(t)$,

where $R(t)$ is a symmetric matrix, C_1 and C_2 are $(2n+1) \times (2n+1)$ matrices defined by

- (1) $\tilde{C}_1 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ is a 2×2 matrix,
- (2) $\tilde{C}_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ is a 2×2 matrix,
- (3) $C_1 = \begin{pmatrix} \tilde{C}_1 & O \\ O & O \end{pmatrix}$
- (4) $C_2 = \begin{pmatrix} \tilde{C}_2 & O \\ O & I \end{pmatrix}$.

Moreover, a moving frame $\tilde{E}_1(t), \dots, \tilde{E}_{2n+1}(t), \tilde{F}_1(t), \dots, \tilde{F}_{2n+1}(t)$ satisfies conditions (1)-(6) above if and only if

$$(3.4) \quad (\tilde{E}_1(t), \tilde{E}_2(t), \tilde{F}_1(t), \tilde{F}_2(t)) = \pm(E_1(t), E_2(t), F_1(t), F_2(t))$$

and there exists a constant orthogonal matrix U of size $(2n-1) \times (2n-1)$ (independent of time t) such that

$$(3.5) \quad \tilde{E}_i(t) = \sum_{j=3}^{2n+1} U_{i-2, j-2} E_j(t) \quad \text{and} \quad \tilde{F}_i(t) = \sum_{j=3}^{2n+1} U_{i-2, j-2} F_j(t)$$

for all time $t \geq 0$.

We call any frame $(E(t), F(t))$ in Theorem 3.1 a canonical frame at the point (\mathbf{x}, \mathbf{p}) and call the equations in (5) and (6) of Theorem 3.1 the structural equation of the Jacobi curve (3.3). Note that the conditions (3) - (4) means that the canonical frame is a family of symplectic bases.

Let $\mathcal{R} : \mathbf{ver} \rightarrow \mathbf{ver}$ be the operator defined by

$$\mathcal{R}(E_i(0)) = \sum_{j=1}^{2n+1} R_{ij}(0)E_j(0).$$

Under the following identification, we can also consider \mathcal{R} as an operator on TM

$$v \in T_{\mathbf{x}}M \mapsto \alpha(\cdot) = \langle v, \cdot \rangle \in T_{\mathbf{x}}^*M \mapsto \left. \frac{d}{dt}(\mathbf{p} + t\alpha) \right|_{t=0} \in \mathbf{ver}_{(\mathbf{x}, \mathbf{p})}.$$

Let $\mathbf{ver}_1 = \text{span}\{E_1(0)\}$, $\mathbf{ver}_2 = \text{span}\{E_2(0)\}$, and

$$\mathbf{ver}_3 = \text{span}\{E_3(0), \dots, E_{2n+1}(0)\}.$$

The proofs of following results can be found in [9].

Theorem 3.2. *Under the above identification of $\mathbf{ver}_{(\mathbf{x}, \mathbf{p})}$ and $T_{\mathbf{x}}M$, we have the followings*

- (1) $\mathbf{ver}_1 = \mathbb{R}v_0$,
- (2) $\mathbf{ver}_2 = \mathbb{R}J\mathbf{p}^h$,
- (3) $\mathbf{ver}_3 = \mathbb{R}(\mathbf{p}^h + \mathbf{p}(v_0)v_0) \oplus \{v \in T_{\mathbf{x}}M \mid \langle v, \mathbf{p}^h \rangle = \langle v, J\mathbf{p}^h \rangle = \langle v, v_0 \rangle = 0\}$,

where \mathbf{p}^h is the tangent vector in the distribution \mathcal{D} defined by $\mathbf{p}(\cdot) = \langle \mathbf{p}^h, \cdot \rangle$.

Theorem 3.3. *Assume that the manifold is Sasakian. Then, under the above identifications, \mathcal{R} is given by*

- (1) $\mathcal{R}(v) = 0$ for all v in \mathbf{ver}_1 ,
- (2) $\mathcal{R}(v) = \mathbf{Rm}^*(J\mathbf{p}^h, \mathbf{p}^h)\mathbf{p}^h + p(v_0)^2 J\mathbf{p}^h$ for all v in \mathbf{ver}_2 ,
- (3) $\mathcal{R}(\mathbf{p}^h + p(v_0)v_0) = 0$,
- (4) $\mathcal{R}(v)_{\mathbf{ver}_3} = \mathbf{Rm}^*(v^h, \mathbf{p}^h)\mathbf{p}^h + \frac{1}{4}p(v_0)^2 v^h$ for all v in \mathbf{ver}_3 satisfying $\langle v^h, \mathbf{p}^h \rangle = 0$,

where \mathbf{p}^h and v^h are the tangent vectors in the distribution \mathcal{D} defined by $\mathbf{p}(\cdot) = \langle \mathbf{p}^h, \cdot \rangle$ and $\langle v, \cdot \rangle = \langle v^h, \cdot \rangle$, respectively.

4. MEASURE CONTRACTION AND MATRIX RICCATI EQUATION

In this section, we discuss the connections between measure contraction properties and the matrix Riccati equations. For this, let us fix a point \mathbf{x}_0 in M and let $\mathfrak{f}(\mathbf{x}) = -\frac{1}{2}d_{CC}^2(\mathbf{x}, \mathbf{x}_0)$. By the result of [8], \mathfrak{f} is locally semi-concave in $M - \{x_0\}$. In particular, it is differentiable almost

everywhere and we can define the family of Borel maps $\varphi_t : M \rightarrow M$ by $\varphi_t(\mathbf{x}) = \pi(e^{t\tilde{\mathbf{H}}}(d\mathbf{f}_{\mathbf{x}}))$, where $0 \leq t \leq 1$. Note that $t \mapsto \varphi_t(\mathbf{x})$ is a minimizing geodesic between the points \mathbf{x} and \mathbf{x}_0 (see for instance [2]).

Let μ be the Riemannian volume form. By the result in [12], the measures $(\varphi_t)_*\mu$ are absolutely continuous with respect to μ for all time t in the interval $[0, 1)$. If $(\varphi_t)_*\mu = \rho_t\mu$, then the following equation holds on a set of full measure where \mathbf{f} is twice differentiable:

$$\rho_t(\varphi_t(\mathbf{x})) \det((d\varphi_t)_{\mathbf{x}}) = 1$$

and the determinant is computed with respect to frames of the above mentioned Riemannian structure. Moreover, the map φ_t is invertible for all t in $[0, 1)$ and so we have

$$(4.6) \quad \mu(\varphi_t(U)) = \int_U \frac{1}{\rho_t(\varphi_t(\mathbf{x}))} d\mu(\mathbf{x}) = \int_U \det((d\varphi_t)_{\mathbf{x}}) d\text{vol}(\mathbf{x}).$$

Therefore, in order to prove the main results and the measure contraction properties, it remains to estimate $\det((d\varphi_t)_{\mathbf{x}})$ which can be done using the canonical frame mentioned above. The explanations on this will occupy the rest of this section.

Let \mathbf{x} be a point where the function \mathbf{f} is twice differentiable and let $(E(t), F(t))$ be a canonical frame at the point $(\mathbf{x}, d\mathbf{f}_{\mathbf{x}})$. Let $\varsigma_i = \pi_*(F_i(0))$ be the projection of the frame $F(0)$ onto the tangent bundle TM . Let $dd\mathbf{f}$ be the differential of the map $\mathbf{x} \mapsto d\mathbf{f}_{\mathbf{x}}$ which pushes the above frame on $T_{\mathbf{x}}M$ to a tuple of vectors in $T_{(\mathbf{x}, d\mathbf{f})}T^*M$. Therefore, we can let $A(t)$ and $B(t)$ be the matrices defined by

$$(4.7) \quad dd\mathbf{f}(\varsigma) = A(t)E(t) + B(t)F(t),$$

where $\varsigma = (\varsigma_1, \dots, \varsigma_{2n+1})^T$ and $dd\mathbf{f}(\varsigma)$ is the column obtained by applying $dd\mathbf{f}$ to each entries of ς .

Lemma 4.1. *Let $S(t) = B(t)^{-1}A(t)$. Then*

$$\mu(\varphi_t(U)) = \int_U e^{-\int_0^t \text{tr}(S(\tau)C_2) d\tau} d\mu(\mathbf{x}).$$

Moreover, $S(t)$ satisfies the following matrix Riccati equation

$$\dot{S}(t) - S(t)C_2S(t) + C_1^T S(t) + S(t)C_1 - R(t) = 0, \quad \lim_{t \rightarrow 1} S(t)^{-1} = 0.$$

Proof of Lemma 4.1. By (4.7) and the definition of φ_t , we have

$$d\varphi_t(\varsigma) = B(t)(\pi_* e^{t\tilde{\mathbf{H}}} F(t)).$$

Note that $\tau \mapsto e_*^{t\tilde{\mathbf{H}}}F(t+\tau)$ is a canonical frame at $e^{t\tilde{\mathbf{H}}}(\mathbf{x}, d\mathbf{f})$. Therefore, by [9, Theorem 4.3], we have

$$\begin{aligned} |\nabla_{\text{hor}}\mathbf{f}(\mathbf{x})| |\det(d\varphi_t)| &= |\text{vol}(d\varphi_t(\zeta))| \\ &= |\det(B(t))\text{vol}(\pi_*e_*^{t\tilde{\mathbf{H}}}F(t))| = |\det(B(t))| |\nabla_{\text{hor}}\mathbf{f}(\mathbf{x})|. \end{aligned}$$

Here $\nabla_{\text{hor}}\mathbf{f}$ denotes the horizontal gradient of \mathbf{f} defined by $d\mathbf{f}(v) = \langle \nabla_{\text{hor}}\mathbf{f}, v \rangle$, where v is any vector in the distribution \mathcal{D} .

By combining this with (4.6), we obtain

$$(4.8) \quad \mu(\varphi_t(U)) = \int_U |\det(B(t))| d\text{vol}(\mathbf{x}).$$

On the other hand, by differentiating (4.7) with respect to time t and using the structural equation, we obtain

$$(4.9) \quad \dot{A}(t) + A(t)C_1 - B(t)R_t = 0, \quad \dot{B}(t) + A(t)C_2 - B(t)C_1^T = 0.$$

Therefore,

$$\frac{d}{dt} \det(B(t)) = \det(B(t)) \mathbf{tr}(B(t)^{-1} \dot{B}(t)) = -\det(B(t)) \mathbf{tr}(S(t)C_2).$$

By setting $t = 0$ and apply π_* on each side of (4.7), we have $B(0) = I$. Therefore, we obtain

$$\det(B(t)) = e^{-\int_0^t \mathbf{tr}(S(\tau)C_2) d\tau}.$$

By combining this with (4.8), we obtain the first assertion.

Since $\varphi_1(\mathbf{x}) = \mathbf{x}_0$ for all \mathbf{x} , we have $d\varphi_1 = 0$ and so $B(1) = 0$. By (4.9) and the definition of $S(t)$, we have

$$\dot{S}(t) - S(t)C_2S(t) + C_1^T S(t) + S(t)C_1 - R(t) = 0, \quad \lim_{t \rightarrow 1} S(t)^{-1} = 0$$

as claimed. \square

5. ESTIMATES FOR SOLUTIONS OF MATRIX RICCATI EQUATIONS

According to Lemma 4.1, we need to estimate the term $\mathbf{tr}(S(t)C_2)$. In this section, we provide two such estimates which lead to the main results.

Throughout this section, we assume that the matrix $R(t)$ is of the form

$$R(t) = \begin{pmatrix} 0 & 0 & O_{2n-2} & 0 \\ 0 & R_{bb}(t) & R_{cb}(t) & 0 \\ O_{2n-2}^T & R_{cb}(t)^T & R_{cc}(t) & O_{2n-2}^T \\ 0 & 0 & O_{2n-2} & 0 \end{pmatrix},$$

where O_{2n-2} is the zero matrix of size $1 \times (2n - 2)$.

The following is a consequence of the result in [25].

Lemma 5.2. *Assume that the curvature $R(t)$ satisfies*

$$\begin{pmatrix} R_{bb}(t) & R_{cb}(t) \\ R_{cb}(t)^T & R_{cc}(t) \end{pmatrix} \geq \begin{pmatrix} \mathfrak{k}_1 & 0 \\ 0 & \mathfrak{k}_2 I_{2n-2} \end{pmatrix},$$

where \mathfrak{k}_1 and \mathfrak{k}_2 are two constants. Then

$$e^{-\int_0^t \text{tr}(C_2 S(\tau)) d\tau} \geq (1-t)^{2n+3} \frac{\mathcal{M}_1(\mathfrak{k}_1, t) \mathcal{M}_2^{2n-2}(\mathfrak{k}_2, t)}{\mathcal{M}_1(\mathfrak{k}_1, 0) \mathcal{M}_2^{2n-2}(\mathfrak{k}_2, 0)},$$

where

$$\begin{aligned} \mathcal{D}(k, t) &= \sqrt{|k|}(1-t), \\ \mathcal{M}_1(k, t) &= \begin{cases} \frac{2-2\cos(\mathcal{D}(k, t)) - \mathcal{D}(k, t) \sin(\mathcal{D}(k, t))}{\mathcal{D}(k, t)^4} & \text{if } k > 0 \\ \frac{1}{12} & \text{if } k = 0 \\ \frac{2-2\cosh(\mathcal{D}(k, t)) + \mathcal{D}(k, t) \sinh(\mathcal{D}(k, t))}{\mathcal{D}(k, t)^4} & \text{if } k < 0, \end{cases} \\ \mathcal{M}_2(k, t) &= \begin{cases} \frac{\sin(\mathcal{D}(k, t))}{\mathcal{D}(k, t)} & \text{if } k > 0 \\ 1 & \text{if } k = 0 \\ \frac{\sinh(\mathcal{D}(k, t))}{\mathcal{D}(k, t)} & \text{if } k < 0. \end{cases} \end{aligned}$$

Moreover, equality holds if

$$\begin{pmatrix} R_{bb}(t) & R_{cb}(t) \\ R_{cb}(t)^T & R_{cc}(t) \end{pmatrix} = \begin{pmatrix} \mathfrak{k}_1 & 0 \\ 0 & \mathfrak{k}_2 I \end{pmatrix},$$

Proof. We only prove the case when both constants \mathfrak{k}_1 and \mathfrak{k}_2 are positive. The proofs for other cases are similar and are therefore omitted. Recall that $S(t)$ satisfies

$$\dot{S}(t) - S(t)C_2S(t) + C_1^T S(t) + S(t)C_1 - R(t) = 0, \quad \lim_{t \rightarrow 1} S(t)^{-1} = 0.$$

Let $\Gamma(t)$ be a solution of the following

$$\dot{\Gamma}(t) - \Gamma(t)C_2\Gamma(t) + C_1^T \Gamma(t) + \Gamma(t)C_1 - K = 0, \quad \lim_{t \rightarrow 1} \Gamma(t)^{-1} = 0,$$

where $K = \begin{pmatrix} \mathfrak{k}_1 & 0 \\ 0 & \mathfrak{k}_2 I \end{pmatrix}$.

By the result of [25], $S(t) \leq \Gamma(t)$. Therefore, $\text{tr}(C_2 S(t)) \leq \text{tr}(C_2 \Gamma(t))$. On the other hand, $\Gamma(t)$ can be computed using the result in [16] and it follows that

$$\begin{aligned} \text{tr}(C_2 S(t)) &\leq \frac{\sqrt{\mathfrak{k}_1}(\sin(\mathcal{D}(\mathfrak{k}_1, t)) - \mathcal{D}(\mathfrak{k}_1, t) \cos(\mathcal{D}(\mathfrak{k}_1, t)))}{2 - 2\cos(\mathcal{D}(\mathfrak{k}_1, t)) - \mathcal{D}(\mathfrak{k}_1, t) \sin(\mathcal{D}(\mathfrak{k}_1, t))} \\ &\quad + (2n-2)\sqrt{\mathfrak{k}_2} \cot(\mathcal{D}(\mathfrak{k}_2, t)) + \frac{1}{1-t}. \end{aligned}$$

The assertion follows from integrating the above inequality. \square

Next, we consider the case where the assumptions are weaker than those in Lemma 5.2.

Lemma 5.3. *Assume that the curvature $R(t)$ satisfies $R_{bb}(t) \geq \mathfrak{k}_1$ and $\mathbf{tr}(R_{cc}(t)) \geq (2n-2)\mathfrak{k}_2$ for some constants \mathfrak{k}_1 and \mathfrak{k}_2 . Then*

$$e^{-\int_0^t \mathbf{tr}(C_2 S(\tau)) d\tau} \geq (1-t)^{2n+3} \frac{\mathcal{M}_1(\mathfrak{k}_1, t) \mathcal{M}_2^{2n-2}(\mathfrak{k}_2, t)}{\mathcal{M}_1(\mathfrak{k}_1, 0) \mathcal{M}_2^{2n-2}(\mathfrak{k}_2, 0)}.$$

Proof. Once again, we only prove the case when both constants \mathfrak{k}_1 and \mathfrak{k}_2 are positive. Let us write

$$S(t) = \begin{pmatrix} S_1(t) & S_2(t) & S_3(t) \\ S_2(t)^T & S_4(t) & S_5(t) \\ S_3(t)^T & S_5(t)^T & S_6(t) \end{pmatrix},$$

where $S_1(t)$ is a 2×2 matrix and $S_6(t)$ is 1×1 . Then

$$(5.10) \quad \begin{aligned} & \dot{S}_1(t) - S_1(t) \tilde{C}_2 S_1(t) - S_2(t) S_2(t)^T \\ & - S_3(t) S_3(t)^T + \tilde{C}_1^T S_1(t) + S_1(t) \tilde{C}_1 - R_1(t) = 0, \\ & \dot{S}_4(t) - S_4(t)^2 - S_5(t) S_5(t)^T - S_2(t)^T \tilde{C}_2 S_2(t) - R_{cc}(t) = 0, \\ & \dot{S}_6(t) - S_6(t)^2 - S_5(t)^T S_5(t) - S_3(t)^T \tilde{C}_2 S_3(t) = 0, \end{aligned}$$

where $\tilde{C}_1 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, $\tilde{C}_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$, and $R_1(t) = \begin{pmatrix} 0 & 0 \\ 0 & R_{bb}(t) \end{pmatrix}$.

By the same argument as in Lemma 5.2, we have

$$(5.11) \quad \mathbf{tr}(\tilde{C}_2 S_1(t)) \leq \frac{\sqrt{\mathfrak{k}_1} (\sin(\mathcal{D}(\mathfrak{k}_1, t)) - \mathcal{D}(\mathfrak{k}_1, t) \cos(\mathcal{D}(\mathfrak{k}_1, t)))}{2 - 2 \cos(\mathcal{D}(\mathfrak{k}_1, t)) - \mathcal{D}(\mathfrak{k}_1, t) \sin(\mathcal{D}(\mathfrak{k}_1, t))}.$$

For the term $S_4(t)$, we can take the trace and obtain

$$\frac{d}{dt} \mathbf{tr}(S_4(t)) \geq \frac{1}{2n-2} \mathbf{tr}(S_4(t))^2 + (2n-2)\mathfrak{k}_2.$$

Therefore, an argument as in Lemma 5.2 again gives

$$(5.12) \quad \mathbf{tr} S_4(t) \leq \sqrt{|\mathfrak{k}_2|} (2n-2) \cot(\mathcal{D}(\mathfrak{k}_2, t)).$$

Finally, for the term $S_6(t)$, we also have

$$\dot{S}_6(t) \geq S_6(t)^2.$$

Therefore,

$$S_6(t) \leq \frac{1}{1-t}.$$

By combining this with (5.11) and (5.12), we obtain

$$\begin{aligned} \mathbf{tr}(C_2S(t)) &\leq \sqrt{|\mathfrak{k}_2|}(2n-2)\cot(\mathcal{D}(\mathfrak{k}_2, t)) + \frac{1}{1-t} \\ &\quad + \frac{\sqrt{|\mathfrak{k}_1|}(\sin(\mathcal{D}(\mathfrak{k}_1, t)) - \mathcal{D}(\mathfrak{k}_1, t)\cos(\mathcal{D}(\mathfrak{k}_1, t)))}{2 - 2\cos(\mathcal{D}(\mathfrak{k}_1, t)) - \mathcal{D}(\mathfrak{k}_1, t)\sin(\mathcal{D}(\mathfrak{k}_1, t))}. \end{aligned}$$

The rest follows as in Lemma 5.2. \square

Finally, we finish the proof of the main result. Note that Corollary 2.5 and 2.6 are consequences of the proof of Theorem 1.1 and equality case of Lemma 5.2.

Proof of Theorem 1.1. If $E(t), F(t)$ is a canonical frame at the point $(\mathbf{x}, d\mathfrak{f}_{\mathbf{x}})$ in the cotangent bundle T^*M , then

$$t \mapsto (e_*^{\tau\bar{\mathbf{H}}}(E(t+\tau)), e_*^{\tau\bar{\mathbf{H}}}(F(t+\tau)))$$

is a canonical frame at the point $e^{\tau\bar{\mathbf{H}}}(\mathbf{x}, d\mathfrak{f}_{\mathbf{x}})$. It follows from this that $R(t)$ is the matrix representation of the operator $\mathfrak{R}_{e^{t\bar{\mathbf{H}}}(\mathbf{x}, d\mathfrak{f}_{\mathbf{x}})}$ with respect to the frame $e_*^{\tau\bar{\mathbf{H}}}(E(\tau))$.

Since v_0 is a symmetry, the function $u_0(\mathbf{x}, \mathbf{p}) = \mathbf{p}(v_0(\mathbf{x}))$ is constant along the flow $e^{t\bar{\mathbf{H}}}$ (see for instance [22]). Therefore, by the assumptions and Theorem 3.3, $R_{bb}(t)$ and $\mathbf{tr}(R_{cc}(t))$ are bounded below by $k_b|\nabla_{\mathbf{hor}}\mathfrak{f}|^2 + u_0^2(\mathbf{x}, d\mathfrak{f})$ and $k_c(2n-2)|\nabla_{\mathbf{hor}}\mathfrak{f}|^2 + \frac{u_0^2(\mathbf{x}, d\mathfrak{f})}{4}$, respectively. Therefore, the assumptions of Lemma 5.3 are satisfied. By combining this with Lemma 4.1, the result follows. \square

REFERENCES

- [1] A.A. Agrachev, R.V. Gamkrelidze: Feedback-invariant optimal control theory - I. Regular extremals, *J. Dynamical and Control Systems*, **3**, No. 3, 343-389 (1997).
- [2] A. Agrachev, P. Lee: Generalized Ricci curvature bounds for three dimensional contact subriemannian manifolds, to appear in *Math Ann.*, arXiv: 0903.2550 (2009), 31pp.
- [3] A. Agrachev, P. Lee: Bishop and Laplacian comparison theorems on three dimensional contact subriemannian manifolds with symmetry, to appear in *J. Geom. Anal.*, arXiv: 1105.2206 (2011), 25pp.
- [4] A. Agrachev, Y. Sachkov: Control theory from the geometric viewpoint. *Encyclopaedia of Mathematical Sciences*, 87. Control Theory and Optimization, II. Springer-Verlag, Berlin, 2004.
- [5] A. Agrachev, I. Zelenko: Geometry of Jacobi curves. I, *J. Dynamical and Control systems*, **8**, No. 1, 93-140 (2002).
- [6] F. Baudoin, N. Garofalo: Generalized Bochner formulas and Ricci lower bounds for sub-Riemannian manifolds of rank two, preprint, arXiv:0904.1623.

- [7] F. Baudoin, N. Garofalo: Curvature-dimension inequalities and Ricci lower bounds for sub-Riemannian manifolds with transverse symmetries, preprint, arXiv:1101.3590.
- [8] P. Cannarsa, L. Rifford: Semiconcavity results for optimal control problems admitting no singular minimizing controls. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 25 (2008), no. 4, 773–802.
- [9] P.W.Y. Lee, C. Li: Bishop and Laplacian comparison theorems on Sasakian manifolds, preprint, arXiv: 1310.5322 (2013), 38pp.
- [10] S. Chanillo, P. Yang: Isoperimetric inequalities & volume comparison theorems on CR manifolds. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* 8 (2009), no. 2, 279-307.
- [11] T. Coulhon, I. Holopainen, L. Saloff-Coste: Harnack inequality and hyperbolicity for subelliptic p -Laplacians with applications to Picard type theorems. *Geom. Funct. Anal.* 11 (2001), no. 6, 1139-1191.
- [12] A. Figalli, L. Rifford: Mass Transportation on sub-Riemannian Manifolds. *Geom. Funct. Anal.* 20 (2010), no. 1, 124-159.
- [13] K. Hughen: The geometry of sub-riemannian three-manifolds, Ph.D. Dissertation 1995.
- [14] N. Juillet: Geometric Inequalities and Generalized Ricci Bounds in the Heisenberg Group, *Int. Math. Res. Not. IMRN* 2009, no. 13, 2347-2373.
- [15] P.W.Y. Lee: Displacement interpolations from a Hamiltonian point of view, *J. Func. Anal.* 265 (12), 3163-3203
- [16] J.J. Levin: On the matrix Riccati equation. *Proc. Amer. Math. Soc.* 10 (1959) 519–524.
- [17] C. Li, I. Zelenko: Parametrized curves in Lagrange Grassmannians. *C.R. Acad. Sci. Paris, Ser. I, Vol. 345, Issue 11* (2007) 647–652.
- [18] C. Li, I. Zelenko: Differential geometry of curves in Lagrange Grassmannians with given Young diagram, *Differ. Geom. Appl.*, 27 (2009) 723-742.
- [19] C.Li, I. Zelenko: Jacobi equations and comparison theorems for corank 1 sub-Riemannian structures with symmetries, *J. Geom. Phys.* 61 (2011) 781-807.
- [20] J. Lott, C. Villani: Ricci curvature for metric-measure spaces via optimal transport, *Ann. of Math. (2)*, in press
- [21] J. Lott, C. Villani: Weak curvature conditions and functional inequalities. *J. Funct. Anal.* 245 (2007), no. 1, 311–333
- [22] R. Montgomery: A tour of subriemannian geometries, their geodesics and applications. *Mathematical Surveys and Monographs*, 91. American Mathematical Society, Providence, RI, 2002
- [23] S. Ohta: On the measure contraction property of metric measure spaces. *Comment. Math. Helv.* 82 (2007), no. 4, 805–828
- [24] S. Ohta: Finsler interpolation inequalities, *Calc. Var. Partial Differential Equations* 36 (2009), 211-249
- [25] H.L. Royden: Comparison theorems for the matrix Riccati equation. *Comm. Pure Appl. Math.* 41 (1988), no. 5, 739–746.
- [26] K.T. Sturm: On the geometry of metric measure spaces. *Acta Math.* 196, no.1, 65-131 (2006)
- [27] K.T. Sturm: On the geometry of metric measure spaces II. *Acta Math.* 196, no. 1, 133-177 (2006)

- [28] N. Tanaka: A differential geometric study on strongly pseudo-convex manifold, Kinokunya Book Store Co., Ltd., Kyoto, 1975.
- [29] S. Tanno: Variational problems on contact Riemannian manifolds, Trans. Amer. Math. Soc. 314 n. 1 (1989), 349379.
- [30] S.M. Webster: Pseudo-Hermitian structures on a real hypersurface. J. Differential Geometry, 13, 25-41 (1978).

ROOM 216, LADY SHAW BUILDING, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG

E-mail address: wylee@math.cuhk.edu.hk

DEPARTMENT OF MATHEMATICS, TIANJIN UNIVERSITY, TIANJIN, 300072, P.R.CHINA

E-mail address: chengboli@tju.edu.cn

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843-3368, USA

E-mail address: zelenko@math.tamu.edu