

POWER SERIES DETERMINED BY AN EXPERIMENT ON THE UNIT INTERVAL

Predrag M. Rajković^{1*}; Sladjana D. Marinković²
Miomir S. Stanković³

¹Department of Mathematics, Faculty of Mechanical Engineering

²Department of Mathematics, Faculty of Electronic Engineering

³Department of Mathematics, Faculty of Occupational Safety

University of Niš, Serbia

June 19, 2018

Abstract

We consider the linear combinations of elements of two sequences: the first one a priori given nonnegative sequence and the second random sequence from the unit interval. We investigate the expected value of the smallest natural number such that the value of these linear combinations exceed a positive number. After very clear geometrical conclusions, we find the function which expresses the expected value. Here, we recognize a few known results like the special cases.

MSC 2010: 34K60, 60G50, 33B10.

Keywords: Experiment, expected value, probability, exponential function.

1 Introduction

The well known number $e \approx 2.71818\dots$ appeared like an expected value in a trial problem by Putnam [1] in 1958. This wonderful, riddle-like problem motivated others to solve it in various ways [4]. Even more, B. Čurgus and R.I. Jewetts [2] have considered like a function of upper bound which should be exceed. In that manner, they have got the exponential function.

In this paper, we will expose our generalization of this problem. Namely, consider the next experiment.

Procedure.

Let $a = \{a_k\}_{k \in \mathbb{N}}$ be a given positive nondecreasing sequence, i.e.

$$0 < a_1 \leq a_2 \leq a_3 \leq \dots \leq a_k \leq a_{k+1} \leq \dots \quad (1)$$

*The corresponding author, *E-mail*: pedja.rajk@masfak.ni.ac.rs

To any random sequence $x = \{x_n\}$ ($x_n \in [0, 1], \forall n \in \mathbb{N}$), we can join the finite sequence of weighted partial sums

$$S_{k,m}(a, x) = a_k x_k + a_2 x_2 + \cdots + a_m x_m.$$

Take $t > 0$.

Step 1. Take a random number x_1 from the interval $[0, 1]$ and denote $n = 1$.

Step 2. If $S_{1,n}(x, a) > t$, then memorize the value n and stop.

Step 3. If $S_{1,n}(x, a) \leq t$, then increase n into $n + 1$, take the next random number x_n from $[0, 1]$ and return to Step 2.

Repeat this procedure enough times. What is the expected value for n ?

We are looking for the expectation

$$f(t, a) = E\{n \in \mathbb{N} : S_{1,n-1}(x, a) \leq t < S_{1,n}(x, a)\}. \quad (2)$$

Denote with $p_m(t)$ the following function

$$p_m = \mu(P_m), \quad \text{where } P_m = \{x \in [0, 1]^m : S_{1,m}(a, x) \leq t\}. \quad (3)$$

Here, μ is the Euclidean measure in \mathbb{R}^m .

The expected value for n is

$$f(t, a) = \sum_{m=0}^{\infty} m(p_{m-1} - p_m).$$

If it is fulfilled

$$\lim_{m \rightarrow \infty} m p_m = 0, \quad (4)$$

then

$$f(t, a) = \sum_{n=0}^{\infty} p_n. \quad (5)$$

Also, we will use notation

$$(t)_+ = \begin{cases} t, & t > 0, \\ 0, & t \leq 0, \end{cases} \quad f_+(t) = \begin{cases} f(t), & f(t) > 0, \\ 0, & f(t) \leq 0. \end{cases}$$

2 The geometrical approach

Here, we will expose cases which can be illustrated by one, two and three dimensional figures.

We accept $p_0 = 1$.

In one dimensional space, there are 2 cases:

(1.1) For $0 \leq t \leq a_1$, i.e., when is valid $t/a_1 \leq 1$, the border point is $X_1(t/a_1)$.

Hence $P_1 = X_1 O$, and the measure is

$$p_1 = \mu(P_1) = \frac{t}{a_1}.$$

(1.2) For $a_1 < t$, it is $P_1 = [0, 1]$, and $p_1 = 1$.

In two dimensional space, the line $a_1x_1 + a_2x_2 = t$ passes through the points $X_1(t/a_1, 0)$, $X_2(0, t/a_2)$.

(2.1) For $0 \leq t \leq a_1$, the whole segment X_2X_1 belongs to the square $[0, 1]^2$. Hence P_2 is the interior of the triangle X_2X_1O and the measure is

$$p_2 = \mu(X_2X_1O) = \int_0^{t/a_1} dx_1 \int_0^{(t-a_1x_1)/a_2} dx_2 = \frac{1}{2} \frac{t}{a_1} \frac{t}{a_2}.$$

(2.2) Let $a_1 < t \leq a_2$. This condition guarantees that the points $O(0, 0)$ and $S(1, 1)$ are from the different sides of the line $a_1x_1 + a_2x_2 = t$. That is why it exists the cutting segment of this line and the unit square. Since $t - a_1 > 0$, or equivalently $1 < t/a_1$, the point X_1 is outside the unit square, like on the Figure 3a. Hence

$$p_2 = \frac{t^2 - (t - a_1)^2}{2a_1a_2}.$$

(2.3) Let $a_2 < t \leq a_1 + a_2$. In that case (see Figure 3b), the segment X_1X_2 contains the points $B(1, (t - a_1)/a_2)$ and $C((t - a_2)/a_1, 1)$. This line divides the square $[0, 1]^2$ into 2 pieces. Hence

$$P_2 = \overline{OS_{1,0}BCS_{0,1}} = \overline{X_2X_1O} \setminus (\overline{X_1S_{1,0}B} \cup \overline{X_2CS_{0,1}}),$$

wherefrom

$$p_2 = \mu(P_2) = \frac{1}{2} \frac{t}{a_1} \frac{t}{a_2} - \frac{1}{2} \left(\frac{t}{a_1} - 1 \right) \frac{t - a_1}{a_2} - \frac{1}{2} \left(\frac{t}{a_2} - 1 \right) \frac{t - a_2}{a_1},$$

i.e.

$$p_2 = \frac{t^2 - (t - a_1)^2 - (t - a_2)^2}{2a_1a_2}.$$

We can unify (2.1-3) into one formula:

$$p_2(t) = \frac{1}{2a_1a_2} \left(t^2 - \sum_{k=1}^2 (t - a_k)_+^2 \right) \quad (0 \leq t \leq a_1 + a_2).$$

(2.4) For $a_1 + a_2 < t$, it is valid

$$a_1x_1 + a_2x_2 \leq a_1 + a_2 < t \quad (\forall x_1, x_2 \in [0, 1])$$

Hence $P_2 = [0, 1]^2$, and $p_2 = 1$.

For arbitrary dimensional space, the hyper-plane $a_1x_1 + \dots + a_mx_m = t$ passes through the points

$$X_1(t/a_1, 0, 0, \dots, 0), X_2(0, t/a_2, 0, \dots, 0), \dots, X_m(0, 0, \dots, 0, t/a_m).$$

For $0 \leq t \leq a_1$, the measure of the m -dimensional solid between this hyper-plane and coordinate hyper-planes is

$$\mu(X_m \dots X_1O) = \int_0^{t/a_1} dx_1 \int_0^{t-a_1x_1/a_2} dx_2 \dots \int_0^{\frac{t-a_1x_1-\dots-a_{m-1}x_{m-1}}{a_m}} dx_m,$$




Figure 1: Two dimensional case: a) $1 \leq t/a_1 \wedge t/a_2 \leq 1$ b) $1 \leq t/a_1, t/a_2$

i.e.,

$$\mu(X_m \dots X_1 O) = \frac{t^m}{m! \prod_{j=1}^m a_j}.$$

Applying the formula (5), we get

$$f_0(t; a_1) = \sum_{m=0}^{\infty} \frac{t^m}{m! \prod_{j=1}^m a_j} \quad (t \in [0, a_1]). \quad (6)$$

3 The analytical approach

Let us denote by

$$f_n(t; a_k) = \sum_{i=n}^{\infty} \frac{t^i}{i! \prod_{j=k}^{i+k-1} a_j} \quad (t \in [0, a_k])(n \in \mathbb{N}_0; k \in \mathbb{N}). \quad (7)$$

We accept $f_n(t; a_k) \equiv 0$ ($t < 0$).

It is important to note that the functions $f_n(t; a_k)$ are defined for every $t > 0$ according to D'Alembert criteria and the fact $\inf\{a_i\} = a_1 > 0$.

Lemma 3.1 *The next relations are true for all $k, n \in \mathbb{N}$:*

$$\begin{aligned} f_n(t; a_k) &= \frac{t^n}{n! \prod_{i=k}^{n+k-1} a_i} + f_{n+1}(t; a_k), \\ f'_0(t; a_k) &= \frac{1}{a_k} f_0(t; a_{k+1}), \\ f'_n(t; a_k) &= \frac{1}{a_k} f_{n-1}(t; a_{k+1}) \\ f_0(t; a_{k+n}) &= f_0^{(n)}(t; a_k) \prod_{i=k}^{k+n-1} a_i \\ f_0(t; a_{n+1}) &= f_n^{(n)}(t; a_1) \prod_{i=1}^n a_i. \end{aligned} \quad (8)$$

The function $f(t, a)$ is known on $(0, a_1)$. It is

$$f(t, a) = f_0(t; a_1).$$

Lemma 3.2 *It is valid*

$$f(t, a) = 1 + f_{1,+}(t; a_1) - f_{1,+}(t - a_1; a_1) \quad (a_1 < t \leq a_2).$$

Proof. If we take $a_1 < t \leq a_2$, and a number $x_1 \in [0, 1]$, then

$$f(t - a_1x_1, a) = E\{m \in \mathbb{N} : S_{2,m-1} \leq t - a_1x_1 < S_{2,m}\} = f_{0,+}(t - a_1x_1; a_2).$$

Including the average of $f_{0,+}(t - a_1x_1; a_2)$ when $x_1 \in [0, 1]$, we have

$$f(t, a) = 1 + \int_0^1 f_{0,+}(t - a_1x_1; a_2) dx_1 \quad (a_1 < t \leq a_2). \quad (9)$$

By change $z = t - a_1x_1$, and the fact $f_{0,+}(z; a_2) = a_1f'_{1,+}(z; a_1)$, yield

$$f(t, a) = 1 + \frac{1}{a_1} \int_{t-a_1}^t f_{0,+}(z; a_2) dz = 1 + \int_{t-a_1}^t f'_{1,+}(z; a_1) dz,$$

what finishes the proof. \square

Lemma 3.3 *If $a_2 < t \leq a_3$, then*

$$f(t, a) = 2 + f_{2,+}(t; a_1) - \sum_{k=1}^2 f_{2,+}(t - a_k; a_1) + f_{2,+}(t - a_1 - a_2; a_1).$$

Proof. If we take two numbers $x_1, x_2 \in [0, 1]$, then

$$f(t - a_1x_1 - a_2x_2, a) = E\{m \in \mathbb{N} : S_{3,m-1} \leq t - a_1x_1 - a_2x_2 < S_{3,m}\},$$

i.e.,

$$f(t - a_1x_1 - a_2x_2, a) = f_{0,+}(t - a_1x_1 - a_2x_2; a_3).$$

Including the average of $f_{0,+}(t - a_1x_1 - a_2x_2; a_3)$ when $x_1, x_2 \in [0, 1]$, we have

$$f(t, a) = 2 + \int_0^1 dx_1 \int_0^1 f_{0,+}(t - a_1x_1 - a_2x_2; a_3) dx_2 \quad (a_1 + a_2 < t \leq a_3). \quad (10)$$

By change $z = t - a_1x_1 - a_2x_2$, we yield

$$f(t, a) = 2 + \frac{1}{a_2} \int_0^1 dx_1 \int_{t-a_1x_1-a_2}^{t-a_1x_1} f_{0,+}(z; a_3) dz.$$

Since

$$f_{0,+}(z; a_3) = a_2f'_{1,+}(z; a_2) = a_1a_2f''_{2,+}(z; a_1),$$

we have

$$f(t, a) = 2 + \int_0^1 f'_{2,+}(t - a_1 x_1; a_1) dx_1 - \int_0^1 f'_{2,+}(t - a_2 - a_1 x_1; a_1) dx_1.$$

Applying the Lemma 3.2, we finish the proof. \square

Notice that the conclusions from the previous lemmas can be written in the following forms:

$$f(t, a) = 1 + \sum_{k=0}^1 (-1)^k \sum_{0 \leq i_0 < i_1 \leq 1} f_{1,+} \left(t - \sum_{j=1}^k a_{i_j}; a_1 \right) \quad (a_1 < t \leq a_2)$$

$$f(t, a) = 2 + \sum_{k=0}^2 (-1)^k \sum_{0 \leq i_0 < i_1 < i_2 \leq 2} f_{2,+} \left(t - \sum_{j=1}^k a_{i_j}; a_1 \right) \quad (a_2 < t \leq a_3)$$

Theorem 3.4 *If $a_n < t < a_{n+1}$, it is valid*

$$f(t, a) = n + \sum_{k=0}^n (-1)^k \sum_{0 \leq i_0 < \dots < i_k \leq k} f_{n,+} \left(t - \sum_{j=1}^k a_{i_j}; a_1 \right) \quad (11)$$

Proof. We will prove by mathematical induction. For $n = 1$ it is proven in Lemma 3.2. Also, for $n = 2$ it is proven in Lemma 3.3.

Let us suppose that the formula (11) is true for n . If we take numbers $x_1, x_2, \dots, x_{n+1} \in [0, 1]$, then

$$f(t - S_{1,n+1}, a) = E\{m \in \mathbb{N} : S_{n+2,m-1} \leq t - S_{1,n+1} < S_{n+2,m}\},$$

i.e.,

$$f(t - S_{1,n+1}, a) = f_{0,+}(t - S_{1,n+1}; a_{n+2}) \quad (n + 1 < m).$$

Including the average of $f_{0,+}(t - S_{1,n+1}; a_{n+2})$ when $x_1, x_2, \dots, x_{n+1} \in [0, 1]$, we have

$$f(t, a) = n+1 + \int_0^1 dx_1 \cdots \int_0^1 dx_n \int_0^1 f_{0,+}(t - a_1 x_1 - \dots - a_{n+1} x_{n+1}; a_{n+2}) dx_{n+1},$$

By change $z = t - a_1 x_1 - \dots - a_{n+1} x_{n+1}$, we yield

$$f(t, a) = n + 1 + \frac{1}{a_{n+1}} \int_0^1 dx_1 \cdots \int_0^1 dx_n \int_{t - a_1 x_1 - \dots - a_n x_n - a_{n+1}}^{t - a_1 x_1 - \dots - a_n x_n} f_{0,+}(z; a_{n+2}) dz,$$

Applying the relations (8), we have

$$f_{0,+}(z; a_{n+2}) = f_{n+1,+}^{(n+1)}(z; a_1) \prod_{i=1}^{n+1} a_i,$$

wherefrom

$$f(t, a) = n + 1 + \prod_{i=1}^n a_i \left(\int_0^1 dx_1 \cdots \int_0^1 dx_{n-1} \int_0^1 f_{n+1,+}^{(n)}(t - a_1 x_1 - \cdots - a_n x_n; a_1) dx_n - \int_0^1 dx_1 \cdots \int_0^1 dx_{n-1} \int_0^1 f_{n+1,+}^{(n)}(t - a_{n+1} - a_1 x_1 - \cdots - a_n x_n; a_1) dx_n \right).$$

The integrals are average values for the functions $f_{n+1,+}(t - a_1 x_1 - \cdots - a_n x_n; a_1)$ and $f_{n+1,+}(t - a_{n+1} - a_1 x_1 - \cdots - a_n x_n; a_1)$ when $x_1, x_2, \dots, x_n \in [0, 1]$. That is why we can apply inductual assumption. Hence

$$f(t, a) = n + 1 + \left(n - 1 + \sum_{k=0}^{n-1} (-1)^k \sum_{0 \leq i_0 < \cdots < i_k \leq k} f_{n+1,+} \left(t - \sum_{j=1}^k a_{i_j}; a_1 \right) \right) - \left(n - 1 + \sum_{k=0}^{n-1} (-1)^k \sum_{0 \leq i_0 < \cdots < i_k \leq k} f_{n+1,+} \left(t - a_{n+1} - \sum_{j=1}^k a_{i_j}; a_1 \right) \right).$$

By simplifying, we get the formula (11) for $n + 1$, what finishes the proof. \square

4 Examples

We prepared the programs in the software *Mathematica* to test our conclusions. We used the function `RandomReal[0, 1, WorkingPrecision->32]`. Blue points are provided on the next figures in this way every after 500 random chosen numbers.

4.1 The Laguerre-type exponential function

The *Laguerre-type exponential function* were introduced by G. Dattoli [3]:

$$e_s(t) = \sum_{n=0}^{\infty} \frac{t^n}{(n!)^{s+1}} \quad (0 \leq t \leq 1) (s \in \mathbb{N}_0).$$

From the previous consideration, we can establish the next statement.

Corollary 4.1 *When $a_k = k^s$ ($k \in \mathbb{N}$) ($s \geq 0$), it is valid*

$$f(t, a) = \begin{cases} e_s(t), & (0 \leq t \leq 1) \\ e_s(t) - e_s(t-1) + 1, & (1 < t \leq 2^s). \end{cases}$$

In general, in the formula (11) appears the next fuction

$$f_{n,+}(t) = e_s(t) - \sum_{k=0}^{n-1} \frac{t^k}{(k!)^{s+1}}.$$

Remark 4.1 The case $s = 0$ was examined in [2]. Since $a = \{1\}$, we easily find that $f(t, a) = e^t$ for $0 \leq t \leq 1$, but our approach can not give $f(t, a)$ for $t > 1$.




Figure 2: Dattoli cases: a) $a_k = k$

b) $a_k = k^2$.

4.2 The Laguerre-type q -exponential function

Another theoretically interesting example is the case

$$a_k = 1 - q^k \quad (0 \leq q < 1) \quad (k \in \mathbb{N}).$$

It is leading to a *Laguerre type q -exponential function*[5]

$$e_{1,q}(t) = \sum_{n=0}^{\infty} \frac{t^n}{n!(q; q)_n} \quad (0 < t \leq 1 - q).$$

Here

$$(q; q)_0 = 1, \quad (q; q)_n = \prod_{k=1}^n (1 - q^k) \quad (n \in \mathbb{N}).$$




Figure 3: Case with q -numbers $a_k = 1 - q^k$ ($q = 0.4$).

Acknowledgements. This paper is supported by the Ministry of Science and Technological Development of the Republic Serbia, projects No 174011 and No 44006.

References

- [1] L.E. Bush, *The William Putnam mathematical competition*, Amer. Math. Monthly **68** (1961) 18–33.
- [2] B. Ćurgus, R.I. Jewett, *An unexpected limit of expected values*, Expo. Math. **25** (2007) 1–20.
- [3] G. Dattoli, M.X. He, P.E. Ricci, *Eigenfunctions of Laguerre-Type Operators and Generalized Evolution Problems*, Mathematical and Computer Modelling **42** (2005) 1263–1268.
- [4] H.S. Shultz, *An expected value problem*, Two-Year College Math. J. **10** (1979) 179.
- [5] M.S. Stanković, S.D. Marinković, P.M. Rajković, *The deformed exponential functions of two variables in the context of various statistical mechanics*, Applied Mathematics and Computation **218** (2011) 2439–2448.