LOG-MAJORIZATION OF THE MODULI OF THE EIGENVALUES OF A MATRIX POLYNOMIAL BY TROPICAL ROOTS

MARIANNE AKIAN, STÉPHANE GAUBERT, AND MEISAM SHARIFY

ABSTRACT. We show that the sequence of moduli of the eigenvalues of a matrix polynomial is log-majorized, up to universal constants, by a sequence of "tropical roots" depending only on the norms of the matrix coefficients. These tropical roots are the non-differentiability points of an auxiliary tropical polynomial, or equivalently, the opposites of the slopes of its Newton polygon. This extends to the case of matrix polynomials some bounds obtained by Hadamard, Ostrowski and Pólya for the roots of scalar polynomials. We also obtain new bounds in the scalar case, which are accurate for "fewnomials" or when the tropical roots are well separated.

1. Introduction

Let $p(x) = \sum_{j=0}^{n} a_j x^j$, $a_j \in \mathbb{C}$ be a polynomial of degree n in a complex variable x. Let ζ_1, \ldots, ζ_n denote the roots of p(x) arranged by non-decreasing modulus (i.e., $|\zeta_1| \leq \ldots \leq |\zeta_n|$). We associate with p the tropical polynomial t p(x), defined for all nonnegative numbers x by

$$t p(x) := \max_{0 \leqslant j \leqslant n} |a_j| x^j .$$

The tropical roots of tp, $\alpha_1, \ldots, \alpha_n$, ordered by non-decreasing value (i.e., $\alpha_1 \leq \ldots \leq \alpha_n$), are defined as the non-differentiability points of the function tp, counted with certain multiplicities. They coincide with the exponential of the opposite of the slopes of the edges of a Newton polygon, defined by Hadamard [19] and Ostrowski [30, 31] as the upper boundary of the convex hull of the set of points $\{(j, \log |a_j|) \mid 0 \leq j \leq n\}$. The logarithms of these roots were called the *inclinaisons numériques* by Ostrowski. One interest of these roots is that they can be easily computed (linear number of arithmetic operations and comparisons). See Section 3.1 below for details.

Hadamard was probably the first to prove a log-majorization type inequality for the modulus of the roots of a scalar polynomial by using what we call today the

1

²⁰¹⁰ Mathematics Subject Classification. 15A22,15A80,15A18,47J10.

Key words and phrases. Matrix polynomial, Tropical algebra, Majorization of eigenvalues, Tropical roots, Roots of polynomial, Bound of Pólya.

The first and second authors were partially supported by PGMO, a joint program of EDF and FMJH (Fondation Mathématique Jacques Hadamard).

The last two authors were partly supported by the Arpege programme of the French National Agency of Research (ANR), project "ASOPT", number ANR-08-SEGI-005.

This work was performed when the third author was with INRIA Saclay–Île-de-France and CMAP, Ecole polytechnique.

tropical roots. His result (page 201 of [19], third inequality) can be restated as follows in tropical terms:

(1)
$$\frac{|\zeta_1\zeta_2\dots\zeta_k|}{\alpha_1\cdots\alpha_k}\geqslant \frac{1}{k+1}.$$

This bound, proved in passing in a memoir devoted to the Riemann zeta function, remained apparently not so well known. In particular, the special case $|\zeta_1|/\alpha_1 \ge 1/2$ is equivalent to the homogeneous form of the classical bound of Cauchy, established later on by Fujiwara [16], and a weaker inequality, with α_1^k at the denominator instead of $\alpha_1 \cdots \alpha_k$, appeared later on in the work of Specht [37].

Ostrowski proved several bounds on the roots of a polynomial in his work on the method of Graeffe [30, 31], in which he used again the Newton polygon considered by Hadamard. In particular, he obtained the following upper bound (see [30, \S 7]),

(2)
$$\frac{|\zeta_1\zeta_2\cdots\zeta_k|}{\alpha_1\cdots\alpha_k} \leqslant \binom{n}{k} ,$$

which can be thought of as a generalization of a "reverse" of the Cauchy inequality due to Birkhoff [11] (corresponding to the case k = 1 in (2)). He also gave a different proof of a variant of (1), with the constant 1/(2k) instead of 1/(k+1), and reported a private communication of Pólya, leading to a tighter constant

(3)
$$\frac{|\zeta_1\zeta_2\cdots\zeta_k|}{\alpha_1\cdots\alpha_k}\geqslant \frac{1}{\sqrt{\mathcal{E}(k)(k+1)}},$$

with

(4)
$$\mathcal{E}(k) := \left(\frac{k+1}{k}\right)^k < e .$$

In this paper, we generalize the bounds of Hadamard, Ostrowski, and Pólya, to the case of a matrix polynomial

(5)
$$P(\lambda) = A_0 + A_1 \lambda + \dots + A_d \lambda^d, \quad A_j \in \mathbb{C}^{n \times n}, \quad 0 \leqslant j \leqslant d.$$

We now associate with the matrix polynomial P the tropical polynomial

(6)
$$t p(x) := \max_{0 \le j \le d} ||A_j|| x^j ,$$

where $\|\cdot\|$ is a norm on the space of matrices, and show that the moduli of the roots $\zeta_1, \ldots, \zeta_{nd}$ of P can still be controlled in terms of the tropical roots $\alpha_1, \ldots, \alpha_n$ of t p. Our results give in particular bounds on the ratios $|\zeta_1, \ldots, \zeta_{nk}|/(\alpha_1, \ldots, \alpha_k)^n$, which extend and refine the above bounds. In particular, in Theorem 4.4, we extend the lower bound (3) of Pólya to the matrix polynomial case, and in Theorem 6.1, we extend the upper bound (2) of Ostrowski. Note that, for the lower bounds to be similar to the one of Pólya and in some sense independent of the dimension of the matrices, the norm $\|\cdot\|$ must satisfy assumptions described in Section 2.2, which hold in particular for the normalized Frobenius norm.

We also obtain other lower bounds that are new even in the case of scalar polynomials. In particular, in Theorem 4.1, we obtain a lower bound which may be tighter for "fewnomials". In Theorems 4.5 and 4.6, we obtain general lower bounds, which extend the bound of Pólya and its extension to the matrix case, and which may be much tighter when the tropical roots are sufficiently separated. Then, all together our results show that the tropical roots give tight estimates of the moduli of the

eigenvalues if the tropical roots are sufficiently separated and if certain matrices are sufficiently well conditioned.

The results of the present paper combine ideas from max-plus algebra and tropical geometry, and numerical linear algebra. In [5, 1, 2, 4], Akian, Bapat, and Gaubert studied the eigenvalues and eigenvectors of matrices and matrix polynomials whose entries (the A_i) are functions, for instance Puiseux series, of a (perturbation) parameter. It is shown there that the leading exponents of the Puiseux series representing the different eigenvalues (resp. eigenvectors) coincide, under some genericity conditions, with the "tropical eigenvalues" (resp. eigenvectors) of the tropical matrix polynomial with entries equal to the leading exponents of the entries of the initial matrix polynomial. This can be interpreted in the light of tropical geometry (see [39, 23, 34]), using the notion of non-archimedean amoeba [13] with respect to the usual non-archimedean valuation on the field of Puiseux series (taking the leading exponent). Since amoebas with respect to the archimedean valuation $z \mapsto \log |z|$ on the field of complex numbers \mathbb{C} can be approximated by non-archimedean amoebas [32, 33], tropical eigenvalues are expected to provide approximations of the log of moduli of the classical eigenvalues. The above bounds of Hadamard, Ostrowski, and Pólya can be interpreted as an approximation result of this nature, for roots of univariate scalar valued polynomials. In the case of a matrix with non-negative coefficients, Friedland [15] established a bound for its spectral radius (or its Perron eigenvalue), which can be interpreted as a bound of the maximal eigenvalue of A by the maximal tropical eigenvalue of the valuation of A [3]. Similar bounds for the other eigenvalues have been established by Akian, Gaubert and Marchesini in [6].

Here, we replaced the valuation on \mathbb{C} by the "valuation" $A \mapsto \log \|A\|$ on the ring of $n \times n$ matrices over \mathbb{C} , which leads to (6). The tropical roots obtained in this way are easy to compute and we shall see they provide a good approximation of the moduli of the eigenvalues under reasonable assumptions. The idea of using the norm instead of the usual valuation was inspired by several works in numerical linear algebra, suggesting that the information of the norms is relevant. For instance, Higham and Tisseur [21] extended to matrix polynomials the bound of Cauchy (related to the special case k=1 in the Hadamard-Ostrowski-Pólya inequality), by using the norms of $A_0^{-1}A_j$ and $A_d^{-1}A_j$. Fan, Lin and Van Dooren [14] introduced a scaling based on the norms of the matrix coefficients of a matrix quadratic polynomial.

The interest of tropical roots is not limited to theoretical bounds, they can be used to perform scalings allowing one to improve the accuracy of the numerical computation of the eigenvalues of a matrix polynomial. In [17], the tropical polynomial of (6) was initially introduced to refine the results of [14]. It was shown there (see also [35]) that the tropical roots of tp can be used to perform scalings allowing one to improve the backward stability of eigenvalue computations for a matrix polynomial P. In the special case of quadratic pencils, this was confirmed by a work of Hamarling, Munro, and Tisseur [20], who implemented the tropical scaling as an option of quadeig, and gave theoretical estimates showing that the tropical scaling does reduce the backward error under some conditions. We shall explain this scaling in more details in Section 8. We also explain why the tropical scaling helps to compute the eigenvalues of a matrix polynomial, as soon as all

the ratios $\zeta_{n(k-1)+l}/\alpha_k$ with $k=1,\ldots,d$ and $\ell=1,\ldots,n$ are close to 1. We also illustrate this behavior numerically.

Some bounds on the modulus of the eigenvalues of P, involving the tropical roots of tp, also appeared in [17] in the case in which d=2, and in [35] for the smallest and largest tropical roots when $d \ge 2$, which may be seen as particular cases of the lower bounds of the present paper.

Let us finally point out some further related works. Bini used in [9] what we call the tropical roots (from the Newton polygon technique) to initialize the Aberth method of computation of the roots of a scalar polynomial. Also, Malajovich and Zubelli applied Ostrowski's analysis to effective root solving [27]. Finally, Bini, Noferini, and Sharify recently proved in [10] some location results by using the tropical roots of (6) for the eigenvalues of a specific class of matrix polynomials such that the coefficient matrices are unitary up to some constants. They also generalized the classical Pellet's theorem to the case of matrix polynomials, which involve the norms of the $A_k^{-1}A_j$, except in the specific above class, where they involve the norms of the A_j . They also relate it to the Newton polygon method, thus to the tropical roots. Moreover, Melman [28] proved a different generalization and some variations of Pellet's theorem, which are in general less costly since they involve the norms of the A_j and A_k^{-1} only (we need to compute O(d) norms instead of $O(d^2)$, each of these computations requiring at least an inversion or a matrix product). Also a recent work by Noferini, Sharify, Tisseur [29] improved these results. Note that the information obtained by Pellet type methods is complementary and incomparable to the one obtained by the present log-majorization inequalities, see Remark 6.4.

The paper is organized as follows. We first recall a variation on Jensen formula due to Landau (Section 2.1) and extend it to the matrix case (Section 2.3), exploiting some technical results on matrix norms (Section 2.2). In Section 3.1, we recall the construction of the Newton polygon and of tropical roots. Then, we derive in Section 3.2 a general estimate for the modulus of the product of K smallest eigenvalues of a matrix polynomial in terms of tropical roots. This leads to various explicit lower bounds, stated in Section 4, which extend the Hadamard-Ostrowski-Polyá inequality to the matrix case. These bounds are proved in Section 5.1 in the scalar case, and then in Section 5.2 in general. An upper bound (reverse inequality) is stated in Section 6.1 and proved in Section 6.2. In Section 7 we provide examples showing the tightness of the lower bounds. In Section 8, we explain the tropical scaling and the interest of the above bounds in this context.

- 2. Estimates of the eigenvalues of matrix polynomials using norms of matrices
- 2.1. An inequality of Landau. An ingredient of our results is a bound on the modulus of the product of the k smallest roots of a polynomial in terms of its coefficients. It is a consequence of Jensen's formula, derived by Landau in [25], building on an earlier observation of Lindelöf [26].

Lemma 2.1 ([25]). Let ζ_1, \ldots, ζ_d be the roots of a polynomial $p(x) = \sum_{j=0}^d a_j x^j$, arranged by non-decreasing modulus, and assume that $a_0 \neq 0$. For all $1 \leq k \leq d$, we have

(7)
$$\log |\zeta_1 \dots \zeta_k| \geqslant -\inf_{r>0} \frac{1}{2} \log \left(\sum_{j=0}^d \frac{|a_j|^2}{|a_0|^2} r^{2(j-k)} \right) .$$

We include the short proof, as its idea will be used in the extension to the matrix case.

Proof. The formula of Jensen [24] shows that if ζ_1, \ldots, ζ_k are the roots of p(z) in the closed disk of $\mathbb C$ of radius r, counted with multiplicities, then

$$\log|\zeta_1 \dots \zeta_k| = k \log r + \log|p(0)| - \frac{1}{2\pi} \int_0^{2\pi} \log|p(re^{i\theta})| d\theta.$$

It follows that, for all r > 0, and k = 1, ..., d:

(8)
$$\log |\zeta_1 \dots \zeta_k| \ge k \log r + \log |p(0)| - \frac{1}{2\pi} \int_0^{2\pi} \log |p(re^{i\theta})| d\theta.$$

Using the comparison between the geometric and the L^2 mean, together with Parseval's identity, we get

$$\begin{split} &\frac{1}{2\pi} \int_0^{2\pi} \log |p(re^{i\theta})| d\theta \leqslant \frac{1}{2} \log \left(\frac{1}{2\pi} \int_0^{2\pi} |p(re^{i\theta})|^2 d\theta\right) \\ &= &\frac{1}{2} \log \left(\frac{1}{2\pi} \int_0^{2\pi} |\sum_{j=0}^d a_j r^j e^{ij\theta}|^2 d\theta\right) = \frac{1}{2} \log \left(\sum_{j=0}^d |a_j|^2 r^{2j}\right) \;. \end{split}$$

Gathering this inequality with (8) yields

$$\log |\zeta_1 \dots \zeta_k| \geqslant k \log r + \log |a_0| - \frac{1}{2} \log \left(\sum_{j=0}^d |a_j|^2 r^{2j} \right).$$

Since this holds for all r > 0, this shows the inequality of the lemma.

- 2.2. Preliminary results on matrix norms. In order to generalize Lemma 2.1 to matrix polynomials, and to derive effective lower bounds for eigenvalues, we need to introduce technical assumptions on matrix norms, like the following one:
- (A1) $|\det A| \leq ||A||^n$, for all $A \in \mathbb{C}^{n \times n}$.

The normalized Frobenius norm.

$$||A||_{\text{nF}} := \left(\frac{1}{n} \sum_{i,j=1}^{n} |A_{ij}|^2\right)^{\frac{1}{2}}, \quad \forall A \in \mathbb{C}^{n \times n},$$

will be of special interest, as some of our estimates rely on L_2 methods. Therefore, the next assumption will also be considered:

(A2) There exists $Q, Q' \in \mathbb{C}^{n \times n}$, such that $\det Q = \det Q' = 1$ and $\|QAQ'\|_{nF} \le \|A\|$, for all $A \in \mathbb{C}^{n \times n}$.

We next show that Assumption (A2) implies Assumption (A1), and that a number of commonly used norms satisfy Assumption (A1) or Assumption (A2).

For all $p \in [1, \infty]$, and $n \ge 1$, we shall denote by $\|\cdot\|_p$ the ℓ^p norm of \mathbb{C}^n : $\|v\|_p := (\sum_{i=1}^n |v_i|^p)^{1/p}$ for $p < \infty$ and $\|v\|_{\infty} := \max_{i=1,\dots,n} |v_i|$. In particular, $\|\cdot\|_2$ is the Euclidean norm. The norm on the space of matrices $\mathbb{C}^{n \times n}$ induced by the norm $\|\cdot\|_p$ on \mathbb{C}^n will also be denoted by $\|\cdot\|_p$ (the same norm is used for the domain and the range of A). In particular the norm $\|\cdot\|_2$ on $\mathbb{C}^{n \times n}$ is the spectral norm. Moreover, for all $p \in [1,\infty)$, and $n \ge 1$, we shall denote by $\|\cdot\|_{*p}$ the following normalized Schatten p-norm on the space of matrices $\mathbb{C}^{n \times n}$: $\|A\|_{*p} := (\frac{1}{n} \sum_{i=1}^n \sigma_i^p)^{1/p}$, where $\sigma_1, \dots, \sigma_n$ are the singular values of A (the eigenvalues of $\sqrt{A^*A}$, where A^* denotes

the conjugate transpose of A). Then, the normalized Schatten 2-norm coincides with the normalized Frobenius norm. Recall that the (unnormalized) Schatten 1-norm is also called the trace norm or the Ky Fan n-norm. We shall also denote by $\|A\|_{\infty} := \max_{i=1,\dots,n} \sigma_i$ the Schatten ∞ -norm, which coincides with the spectral norm.

Property 2.2. The normalized Frobenius norm satisfies Assumption (A1). Therefore, Assumption (A2) implies Assumption (A1).

Proof. From Hadamard's inequality, we get that $|\det A| \leqslant \|A^{(1)}\|_2 \cdots \|A^{(n)}\|_2$, for all $A \in \mathbb{C}^{n \times n}$, where $A^{(j)}$ denotes the j-th column of A. Then, since the geometric mean is less than or equal to the arithmetic mean (or simply by the concavity of the logarithm), we get that $|\det A|^{2/n} \leqslant \frac{1}{n}(\|A^{(1)}\|_2^2 + \cdots + \|A^{(n)}\|_2^2) = \|A\|_{\mathrm{nF}}^2$, which implies that the normalized Frobenius norm satisfies Assumption (A1). Since for all $Q, Q' \in \mathbb{C}^{n \times n}$ such that $\det Q = \det Q' = 1$, we get $|\det A| = |\det(QAQ')| \leqslant \|QAQ'\|_{\mathrm{nF}}^n$ for all $A \in \mathbb{C}^{n \times n}$, Assumption (A2) implies Assumption (A1).

Property 2.3. For all $p \in [1, \infty]$, the norm $\|\cdot\|_p$ on $\mathbb{C}^{n \times n}$ satisfies (A2).

Proof. Indeed, let us denote by $\|A\|_{p,q}$ the norm induced by the norms $\|\cdot\|_p$ and $\|\cdot\|_q$ on the range and domain \mathbb{C}^n of A respectively, which means that $\|A\|_{p,q} = \max\{\|Av\|_q/\|v\|_p, \ v\in\mathbb{C}^n, v\neq 0\}$. Since (for all $v\in\mathbb{C}^n$) $p\in[1,\infty]\mapsto\|v\|_p\in\mathbb{R}^+$ is a nonincreasing map, we get that $\|A\|_{p,q}\leqslant\|A\|_{p',q'}$ when $p\leqslant p'$ and $q\geqslant q'$. In particular $\|A\|_p=\|A\|_{p,p}\geqslant\|A\|_{2,\infty}$ when $2\leqslant p$, and $\|A\|_p=\|A\|_{p,p}\geqslant\|A\|_{1,2}$ when $p\leqslant 2$. In addition, it is easy to show that $\|A\|_{2,\infty}=\max\{\|(A^*)^{(i)}\|_2, i=1,\ldots,n\}$. Since the last expression is greater or equal to $\|A\|_{\mathrm{nF}}$, we deduce that $\|A\|_p\geqslant\|A\|_{2,\infty}\geqslant\|A\|_{\mathrm{nF}}$ for all $A\in\mathbb{C}^{n\times n}$ and $2\leqslant p\leqslant \infty$. Similarly, $\|A\|_{1,2}=\max\{\|A^{(i)}\|_2, i=1,\ldots n\}\geqslant\|A\|_{\mathrm{nF}}$, hence $\|A\|_p\geqslant\|A\|_{1,2}\geqslant\|A\|_{\mathrm{nF}}$ for all $A\in\mathbb{C}^{n\times n}$ and $1\leqslant p\leqslant 2$. This shows that $\|\cdot\|_p$ satisfies (A2) with Q=Q'=I the identity matrix.

This result implies that all norms induced by the norm $||v||_d = ||Qv||_p$ in the domain of A and the norm $||v||_r = ||Q'v||_p$ in the range of A with the same p, but possibly different matrices Q, Q' such that $\det Q = \det Q' = 1$, satisfy (A2).

Property 2.4. If $\|\cdot\|$ is the norm on $\mathbb{C}^{n\times n}$ induced by a norm on \mathbb{C}^n (the same for the domain and the range of matrices), then it satisfies (A1).

Proof. Since det A is the product of the eigenvalues of A counted with multiplicities, we get that $|\det A| \leq \rho(A)^n$, where $\rho(A)$ is the spectral radius of A. Since $\rho(A) \leq ||A||$ for all $A \in \mathbb{C}^{n \times n}$ and all induced norms $||\cdot||$ on the space of matrices, we get the result.

Property 2.5. The normalized Schatten p-norm $\|\cdot\|_{*p}$ on $\mathbb{C}^{n\times n}$ satisfies (A1), for all $p\in[1,\infty]$. It satisfies (A2) if and only if $p\geqslant 2$. Moreover, for $p\in[1,2)$, the least constant η such that $\eta\|\cdot\|_{*p}$ satisfies (A2) is given by $\eta=n^{1/p-1/2}>1$.

Proof. Since $|\det A|$ is the product of the singular values of A counted with multiplicities, we get that $|\det A|^p = \sigma_1^p \cdots \sigma_n^p$, and using that the geometric mean is less than or equal to the arithmetic mean, we obtain that $|\det A|^{p/n} \leqslant \frac{1}{n}(\sigma_1^p + \cdots + \sigma_n^p) = ||A||_{*p}^p$, which implies that the normalized Schatten p-norm satisfies Assumption (A1).

We have that $p \mapsto ||A||_{*p}$ is a nondecreasing map. Hence, when $p \in [2, \infty]$, $||A||_{*p} \geqslant ||A||_{*2} = ||A||_{nF}$, for all $A \in \mathbb{C}^{n \times n}$, which implies that $||\cdot||_{*p}$ satisfies (A2) when $p \ge 2$. Let $\eta > 0$ be the least constant such that $\eta \| \cdot \|_{*p}$ satisfies (A2) and let us show that $\eta = n^{1/p-1/2}$ when p < 2. This will implies in particular that $\eta > 1$ hence $\|\cdot\|_{*p}$ does not satisfy (A2) for p < 2, which will finishes the proof of the equivalence " $\|\cdot\|_{*p}$ satisfies (A2) if and only if $p \ge 2$ ". Since $n^{1/p}\|\cdot\|_{*p}$ is the unnormalized Schatten norm, which is nonincreasing with respect to p, we get that, for all $p \in [1,2)$, and $A \in \mathbb{C}^{n \times n}$, $n^{1/2} ||A||_{nF} \leq n^{1/p} ||A||_{*p}$, which implies that $\eta \leqslant n^{1/p-1/2}$. Let us fix $p, Q, Q' \in \mathbb{C}^{n \times n}$ such that $\det Q = \det Q' = 1$ and $\|QAQ'\|_{nF} \leqslant \eta \|A\|_{*p}$, for all $A \in \mathbb{C}^{n \times n}$. For all $i = 1, \ldots, n$, let us consider the matrix A whose entries are all zero but the entry ii equal to 1. Then, the singular values of A are all zero except one which is equal to 1, so that $||A||_{*p}$ $(1/n)^{1/p}$. Moreover $QAQ' = Q^{(i)}Q'_{(i)}$, where $Q'_{(i)}$ denotes the *i*-th raw of Q'. Hence $\|QAQ'\|_{nF} = (1/n)^{1/2} \|Q^{(i)}\|_2 \|(Q'^*)^{(i)}\|_2$. From $\|QAQ'\|_{nF} \leqslant \eta \|A\|_{*p}$, we deduce that $\|Q^{(i)}\|_2 \|(Q'^*)^{(i)}\|_2 \leqslant \eta (1/n)^{1/p-1/2}$. Since $\det Q = \det Q' = 1$, we get using Hadamard's inequality, $1 \leq \|Q^{(1)}\|_2 \|(Q'^*)^{(1)}\|_2 \cdots \|Q^{(n)}\|_2 \|(Q'^*)^{(n)}\|_2 \leq (\eta(1/n)^{1/p-1/2})^n$, which shows that $\eta \geq n^{1/p-1/2}$. We deduce that $\eta = n^{1/p-1/2} > 1$ for p > 2, which completes the proof.

2.3. Generalization of the inequality of Landau to matrix polynomials. The following generalization of Lemma 2.1 will be a key tool to establish lower bounds for the eigenvalues of matrix polynomials.

Lemma 2.6. Consider the matrix polynomial P with degree d defined in (5), and let $\zeta_1, \ldots, \zeta_{nd}$ denote its eigenvalues, arranged by non-decreasing modulus. Assume that $\|\cdot\|$ is any norm on the space of matrices satisfying (A1), that $\det A_0 \neq 0$ and let $c = \frac{|\det A_0|}{\|A_0\|^n}$. Then, for $K \in \{1, \ldots, nd\}$, we have

(9)
$$\log |\zeta_1 \dots \zeta_K| \ge \log c - n \inf_{r>0} \log \left(\sum_{j=0}^d \frac{\|A_j\|}{\|A_0\|} r^{j-\frac{K}{n}} \right) .$$

When $\|\cdot\|$ satisfies (A2), the previous bound can be improved as follows

(10)
$$\log |\zeta_1 \dots \zeta_K| \geqslant \log c - n \inf_{r>0} \frac{1}{2} \log \left(\sum_{j=0}^d \frac{\|A_j\|^2}{\|A_0\|^2} r^{2(j-\frac{K}{n})} \right) .$$

Proof. From Inequality (8) applied to $\tilde{p}(\lambda) = \det P(\lambda)$, we get, for all r > 0,

(11)
$$\log |\zeta_1 \dots \zeta_K| \geqslant K \log r + \log |\det A_0| - \frac{1}{2\pi} \int_0^{2\pi} \log |\det P(re^{i\theta})| d\theta.$$

Since $\|\cdot\|$ satisfies (A1), we have

(12)
$$|\det P(re^{i\theta})| \le ||P(re^{i\theta})||^n = ||\sum_{j=0}^d A_j (re^{i\theta})^j||^n \le (\sum_{j=0}^d ||A_j||r^j)^n$$
,

for all $\theta \in [0, 2\pi]$. Gathering (12) with (11), we obtain (9).

Assume now that $\|\cdot\|$ satisfies (A2) with some matrices Q, Q' such that $\det Q = \det Q' = 1$, and let us prove (10). Since from Property 2.2, the normalized Frobenius norm satisfies (A1), we get that $|\det P(re^{i\theta})| = |\det(QP(re^{i\theta})Q')| \leq$

 $||QP(re^{i\theta})Q'||_{nF}^n$. Now using the comparison between geometric and L^2 means, we deduce

$$\frac{1}{2\pi} \int_0^{2\pi} \log|\det P(re^{i\theta})| d\theta \leqslant n \frac{1}{2\pi} \int_0^{2\pi} \log||QP(re^{i\theta})Q'||_{nF} d\theta$$

$$\leqslant \frac{n}{2} \log\left(\frac{1}{2\pi} \int_0^{2\pi} ||QP(re^{i\theta})Q'||_{nF}^2 d\theta\right) .$$
(13)

From the formula of the normalized Frobenius norm, we get by applying Parseval's identity to each coordinate $(QP(re^{i\theta})Q')_{\ell m}$

$$\frac{1}{2\pi} \int_0^{2\pi} \|QP(re^{i\theta})Q'\|_{\mathrm{nF}}^2 d\theta = \frac{1}{n} \sum_{\ell,m=1}^n \left(\frac{1}{2\pi} \int_0^{2\pi} |(QP(re^{i\theta})Q')_{\ell,m}|^2 d\theta \right) \\
= \frac{1}{n} \sum_{\ell,m=1}^n \left(\sum_{j=0}^d (|(QA_jQ')_{\ell,m}|r^j)^2 \right) = \sum_{j=0}^d (\|QA_jQ'\|_{\mathrm{nF}}^2 r^{2j}) \leqslant \sum_{j=0}^d (\|A_j\|^2 r^{2j}) .$$

Gathering this inequality with Inequalities (13) and (11), we obtain (10).

3. Tropical polynomials and numerical Newton Polygons

3.1. Preliminary results on tropical roots. We recall here basic results on tropical polynomials of one variable. See for instance [8, 39, 23] for more background on tropical polynomials from different perspectives.

Let \mathbb{R}_{\max} denotes the set $\mathbb{R} \cup \{-\infty\}$. A (max-plus) tropical polynomial f is a function of a variable $x \in \mathbb{R}_{\max}$ of the form

(14)
$$f(x) = \max_{0 \le j \le d} (f_j + jx) ,$$

where d is an integer, and f_0, \ldots, f_d are given elements of \mathbb{R}_{\max} . We say that f is of degree d if $f_d \neq -\infty$. We shall assume that at least one of the coefficients f_0, \ldots, f_d is finite (i.e., that f is not the tropical "zero polynomial"). Then, f is a real valued convex function, piecewise affine, with integer slopes.

Cuninghame-Green and Meijer showed [12] that the analogue of the fundamental theorem of algebra holds in the tropical setting, i.e., f(x) can be written uniquely as

$$f(x) = f_d + \sum_{j=1}^{d} \max(x, \alpha_j) ,$$

where $\alpha_1 \leqslant \cdots \leqslant \alpha_d \in \mathbb{R}_{\max}$. The numbers $\alpha_1, \ldots, \alpha_d$ are called the *tropical roots*. The finite tropical roots can be checked to be the points at which the maximum in the expression (14) of f(x) is attained at least twice, whereas $-\infty$ arises as a tropical root if $f_0 = -\infty$. The *multiplicity* of a root α is defined as the cardinality of the set $\{j \in \{1, \ldots, d\} \mid \alpha_j = \alpha\}$.

The multiplicity of a finite root α can be checked to coincide with the variation of the derivative of the map f at point α , whereas the multiplicity of the root $-\infty$ is given by $\inf\{j \mid f_j \neq -\infty\}$ or by the slope of the map f at $-\infty$. The notion of tropical roots is an elementary special case of the notion of tropical variety which has arisen recently in tropical geometry [23].

The tropical roots can be computed by the following variant of the classical Newton polygon construction. Define the Newton polygon $\Delta(f)$ of f to be the

upper boundary of the convex hull of the region

$$\{(j,\lambda) \in \mathbb{N} \times \mathbb{R} \mid \lambda \leqslant f_j, \ 0 \leqslant j \leqslant d\} \subset \mathbb{R}^2$$
.

The latter region is the hypograph of the map $\tilde{f}: \mathbb{R} \to \mathbb{R} \cup \{-\infty\}$ such that $\tilde{f}(j) = f_j$ for $j \in \{0, \ldots, d\}$ and $\tilde{f}(x) = -\infty$ otherwise. Hence, the Newton polygon coincides with the graph of the concave hull of \tilde{f} . Let us denote by \hat{f} this concave hull and by \hat{f}_j its value at an integer point j. Then the Newton polygon of f consists of (linear) segments relying (or passing through) the points $(j, \hat{f}_j), j \in \{0, \ldots, d\}$.

The following result relies on standard Legendre-Fenchel duality. A proof can be found in [2, Proposition 2.10] (for min-plus polynomials instead of max-plus polynomials).

Proposition 3.1. There is a bijection between the set of finite tropical roots of f and the set of segments of the Newton polygon $\Delta(f)$: the tropical root corresponding to a segment is the opposite of its slope, and the multiplicity of this root is the length of this segment (measured by the difference of the abscissæ of its endpoints).

Remark 3.2. The Graham scan algorithm [18] allows one to compute the convex hull of a finite set of points by making O(n) arithmetical operations and comparisons, provided that the given set of points is already arranged by increasing abscissæ, which is the case in the present setting. It follows that the tropical roots, counted with multiplicities, can be computed in linear time (see also [17, Proposition 1]).


In particular, the maximal tropical root is given by

$$\alpha_d = \max_{0 \le j \le d} \frac{f_j - f_d}{d - j} .$$

Example 3.3. Consider

$$f(x) = \max(0, 1+x, 6+2x, 4+4x, 9+8x, 5+10x, 1+16x)$$
.

The graph of f and the Newton polygon of f are shown in Figure 1. The tropical roots are -3, -1/2, and 1, with respective multiplicities 2, 6 and 8.






FIGURE 1. Graph of the tropical polynomial function f on left, and of its Newton polygon on right.

The notion of root also applies with trivial changes to the "max-times" model of the tropical structure, in which polynomial functions now have the form

$$t p(x) = \max_{0 \le j \le d} a_j x^j ,$$

where a_0, \ldots, a_d are nonnegative numbers, and the variable x now takes nonnegative values. Then, the *tropical roots* of tp(x) are, by definition, the exponentials of the tropical roots of its log-exp transformation $f(x) := \log t p(\exp(x)) = \max_{0 \le j \le d} (\log a_j + jx)$.

In the sequel we shall consider max-times polynomials associated with usual scalar or matrix polynomials. We shall need the following result which follows from the above definitions and properties.

Proposition 3.4. Let $tp(x) = \max_{0 \le j \le d} a_j x^j$ be a max-times polynomial of degree d, with $a_j \ge 0$, $j = 0, \dots d$. Assume that $a_0 \ne 0$. Let $0 < \alpha_1 \le \dots \le \alpha_d$ denote the tropical roots of tp arranged in non-decreasing order. Then,

$$a_j \leqslant a_0 \prod_{\ell=1}^{j} \alpha_{\ell}^{-1}, \quad j = 0, \dots d.$$

Moreover, let k be the abscissa of a vertex of the Newton polygon of p, then $a_k > 0$ and

$$a_j \leqslant a_k \prod_{\ell=k+1}^j \alpha_\ell^{-1}, \quad for \ j = k, \dots d,$$

 $a_j \leqslant a_k \prod_{\ell=j+1}^k \alpha_\ell, \quad for \ j = 0, \dots k.$

Proof. Let $f(x) := \max_{0 \leqslant j \leqslant d} (\log a_j + jx)$. By definition $\log \alpha_j, j = 1, \ldots d$, are the tropical roots of f and by the assumption $a_0 \neq 0$, they are all finite. Let $f_j = \log a_j$, for $j = 0, \ldots, d$ be the coefficients of the tropical polynomial f. From the above observations, the Newton polygon of f coincides with the graph of the concave hull \hat{f} of the map $\tilde{f} : \mathbb{R} \to \mathbb{R} \cup \{-\infty\}$ such that $\tilde{f}(j) = f_j$ for $j \in \{0, \ldots, d\}$ and $\tilde{f}(x) = -\infty$ otherwise. Then the Newton polygon of f consists of (linear) segments relying (or passing through) the points $(j, \hat{f}_j), j \in \{0, \ldots, d\}$. By Proposition 3.1, the finite tropical roots are the opposites of the slopes of \hat{f} and their multiplicities are the lengths of the segments where \hat{f} has this slope. This means that $\log \alpha_j = \hat{f}_{j-1} - \hat{f}_j$, hence $\hat{f}_j = \hat{f}_0 - \sum_{\ell=1}^j \log \alpha_\ell$, and using that \hat{f} is above the map \tilde{f} , and that both maps coincide at the boundary point j = 0, we get the first inequality of the proposition. If now k is the abscissa of a vertex of the Newton polygon of f, then f is an exposed point of the hypograph of f, which implies that f is an exposed point of the hypograph of f, which implies that two last inequalities of the proposition.

3.2. A general lower bound using tropical roots. Using the same method as in the proof of Pólya's inequality reproduced in [30], we derive from Landau's inequality the following result which involves now the tropical roots α_k instead of the moduli $|a_k|$ of the coefficients of p.

Proposition 3.5. Let ζ_1, \ldots, ζ_d be the roots of a univariate scalar polynomial p of degree d, $p(x) = \sum_{j=0}^{d} a_j x^j$, arranged by non-decreasing modulus, and assume that $a_0 \neq 0$. Let $\alpha_1, \ldots, \alpha_d$ denote the tropical roots of the associated tropical polynomial $t p(x) = \max_{0 \leq j \leq d} |a_j| x^j$, arranged in non-decreasing order. Then, for

all $k \in \{1, \ldots, d\}$, the inequality

$$\frac{|\zeta_1 \dots \zeta_k|}{\alpha_1 \dots \alpha_k} \geqslant L_k \ ,$$

holds with

(16)
$$(L_k)^{-2} = \inf_{\xi > 0} \left(\sum_{j=0, a_j \neq 0}^d \beta_{k,j}^2 \xi^{j-k} \right) ,$$

and

(17)
$$\beta_{k,j} := \begin{cases} \prod_{\ell=j+1}^{k-1} \left(\frac{\alpha_{\ell}}{\alpha_{k}}\right) & \text{if } j < k-1 \\ \prod_{\ell=k+1}^{j} \left(\frac{\alpha_{k}}{\alpha_{\ell}}\right) & \text{if } j > k \\ 1 & \text{if } j = k-1 \text{ or } k \end{cases},$$

so that in particular $\beta_{k,j} \leq 1$ for all $k, j = 0, \ldots, d$.

Proof. By applying the change of variable $r = \alpha_k \sqrt{\xi}$ in the inequality of Lemma 2.1, we get

$$\log |\zeta_{1} \dots \zeta_{k}| \geq \sup_{\xi > 0} \left(-\frac{1}{2} \log \left(\sum_{j=0}^{d} \frac{|a_{j}|^{2}}{|a_{0}|^{2}} (\alpha_{k}^{2} \xi)^{j-k} \right) \right)$$

$$= \log(\alpha_{1} \dots \alpha_{k}) + \sup_{\xi > 0} \left(-\frac{1}{2} \log \left(\prod_{\ell=1}^{k} \alpha_{\ell}^{2} \left(\sum_{j=0}^{d} \frac{|a_{j}|^{2}}{|a_{0}|^{2}} (\alpha_{k}^{2} \xi)^{j-k} \right) \right) \right).$$

Applying Proposition 3.4 to the max-times polynomial t p, we get $|a_j| \leq |a_0| \prod_{\ell=1}^j \alpha_\ell^{-1}$ for all $j = 0, \ldots, d$, which with the above inequality yields (15) with

$$(L_k)^{-2} := \inf_{\xi > 0} \left(\sum_{j=0, a_j \neq 0}^d \left(\frac{(\prod_{\ell=1}^k \alpha_\ell) (\prod_{\ell=1}^j \alpha_\ell^{-1})}{\alpha_k^{k-j}} \right)^2 \xi^{j-k} \right)$$

which can be written in the form (16) with $\beta_{k,j}$ as in (17). Moreover, since α_j is nondecreasing with respect to j, we get that all the $\beta_{k,j}$ are less than or equal to 1.

The following is a matrix version of Proposition 3.5. It is proved along the same lines. For any real number x, $\lceil x \rceil$ will denote the least integer $\geqslant x$: $\lceil x \rceil - 1 < x \leqslant \lceil x \rceil$.

Proposition 3.6. Let $A_0, \ldots, A_d, P, \zeta_1, \ldots, \zeta_{nd}, \|\cdot\|$, c, be as in the first part of Lemma 2.6, and let $\alpha_1, \ldots, \alpha_d$ be the tropical roots of the tropical polynomial of (6), arranged in non-decreasing order. For $K \in \{1, \ldots, nd\}$, denote $\bar{\alpha}_K = \alpha_{\lceil \frac{K}{n} \rceil}$. Then, for all $K \in \{1, \ldots, nd\}$, the inequality

(18)
$$\frac{|\zeta_1 \dots \zeta_K|}{\bar{\alpha}_1 \dots \bar{\alpha}_K} \geqslant c(L_{\frac{K}{n}})^n ,$$

holds with L_k , $0 < k \le d$, such that:

(19)
$$(L_k)^{-1} = \inf_{\xi > 0} \left(\sum_{j=0, A_j \neq 0}^d \beta_{\lceil k \rceil, j} \, \xi^{j-k} \right) ,$$

with β defined as in (17).

Moreover, when $\|\cdot\|$ satisfies (A2), the constant L_k in (18) can be replaced by the greater one:

(20)
$$(L_k)^{-2} = \inf_{\xi > 0} \left(\sum_{j=0, A_j \neq 0}^d (\beta_{\lceil k \rceil, j})^2 \xi^{j-k} \right) .$$

Note that $\bar{\alpha}_1, \dots, \bar{\alpha}_K$ are the tropical roots of the tropical polynomial $(tp)^n$ arranged in non-decreasing order. Hence the above result compares the eigenvalues of P with the tropical roots of $(tp)^n$.

Remark 3.7. In the above proof, the only ingredient to deduce the result of Proposition 3.5 from that of Lemma 2.1 is the inequality $|a_j| \leq |a_0| \prod_{\ell=1}^j \alpha_\ell^{-1}$ for all $j=0,\ldots,d$. By Proposition 3.4, this inequality is an equality for any abscissa j of a vertex of the Newton polygon of p. Hence, if all the tropical roots of the tropical polynomial of (6) are simple (which means that all $k=0,\ldots,d$ are abscissæ of a vertex of the Newton polygon of p), the inequalities of Proposition 3.5 and Lemma 2.1 are equivalent. The same is true for the inequalities of Proposition genlowercormatrix and Lemma 2.6.

4. Explicit lower bounds in terms of tropical roots

We now derive from Proposition 3.6 explicit lower bounds for the product of the K smallest eigenvalues of a matrix polynomial. These new bounds can be easily computed (in O(d) time, either in floating point or exact arithmetics, except for the ones of Corollary 4.8 which can be computed in $O(d^2)$ time), as soon as c and the tropical roots are given. Recall that the tropical roots themselves can be computed in O(d) time as soon as the coefficients of the tropical polynomial, that is the norms of the coefficient matrices, are given (see Remark 3.2). Different bounds can be given, depending on the information available on the matrix polynomial. Our first bound is useful for "fewnomials" (polynomials with few non-zero coefficients).

Theorem 4.1 (Bounds involving the number of nonzero coefficients). Consider the matrix polynomial P with degree d defined in (5), and let $\zeta_1, \ldots, \zeta_{nd}$ denote its eigenvalues, arranged by non-decreasing modulus. Assume that $\|\cdot\|$ is any norm on the space of matrices satisfying (A1), that $\det A_0 \neq 0$ and let $c = \frac{|\det A_0|}{\|A_0\|^n}$. Let $\alpha_1, \ldots, \alpha_d$ be the tropical roots of the tropical polynomial of (6), arranged in non-decreasing order, and for $K \in \{1, \ldots, nd\}$, denote $\bar{\alpha}_K = \alpha_{\lceil \frac{K}{n} \rceil}$. Also let mon P denote the number of nonzero monomials of P. Then, for all $K \in \{1, \ldots, nd\}$, we have

(21)
$$\frac{|\zeta_1 \dots \zeta_K|}{\bar{\alpha}_1 \dots \bar{\alpha}_K} \geqslant c(L_{\frac{K}{n}})^n ,$$

in particular, for all $1 \leq k \leq d$, we have

(22)
$$\frac{|\zeta_1 \dots \zeta_{nk}|}{(\alpha_1 \dots \alpha_k)^n} \geqslant c(L_k)^n.$$

with, for $0 < k \le d$,

$$(23) L_k = \frac{1}{\operatorname{mon} P} .$$

Moreover, when $\|\cdot\|$ satisfies (A2), the constant L_k can be replaced by the greater one:

$$(24) L_k = \frac{1}{\sqrt{\text{mon } P}} .$$

The proof of all the results of this section will be given in Section 5.

Remark 4.2. When n=1 (thus the matrices A_0, \ldots, A_d are scalars), any norm is proportional to the normalized Frobenius norm which is nothing but the modulus map $|\cdot|$, satisfies (A2), and for which c=1. Therefore the tropical roots of tp are the same for all norms, the best possible inequality (21) is

(25)
$$\frac{|\zeta_1 \dots \zeta_k|}{\alpha_1 \dots \alpha_k} \geqslant L_k ,$$

and this inequality holds with L_k as in (24).

Remark 4.3. If a norm $\|\cdot\|$ on the space of matrices satisfies (A1) but not (A2), we can still obtain a bound of the form (21,24) by changing the constant c. Indeed, by the equivalence between norms on $\mathbb{C}^{n\times n}$, for any norm $\|\cdot\|$ on $\mathbb{C}^{n\times n}$, there exists a constant $\eta > 0$ (which depends on n) such that $||A||_{nF} \leqslant \eta ||A||$ for all $A \in \mathbb{C}^{n \times n}$. Then, the norm N obtained by multiplying $\|\cdot\|$ by η satisfies (A2), hence (21,24). There, the constant L_k is given by (24), thus independent of the norm, whereas the constant c and the tropical roots α_j are computed with respect to the norm N. In particular $c = \frac{|\det A_0|}{(\eta ||A_0||)^n} = c_0/\eta^n$, where c_0 is the constant c associated with the norm $\|\cdot\|$. Moreover, if t p denotes the tropical polynomial associated with the norm $\|\cdot\|$, then the tropical polynomial associated with the norm N is equal to $\eta t p$, and its tropical roots are equal to the ones of tp. Hence, we deduce that (21,24)holds for $\|\cdot\|$ with c/η^n instead of c. However, if $\|\cdot\|$ satisfies (A1) but not (A2), then $\eta > 1$, so that the inequality derived from (21,24) with c/η^n instead of c may be weaker than (21.23): this is indeed the case if and only if $n > \sqrt{\text{mon } P}$. The same type of conclusions can be obtained for the lower bounds that are stated in the next theorems.

The following theorem provides a lower bound generalizing the lower bound of Pólya to the matrix case (since, as said in Remark 4.2, when n=1, the modulus map is a norm satisfying (A2), and for which Inequality (21) reduces to (25)). Up to the constant c, the following bounds are independent of the coefficients of the matrix polynomial P.

Theorem 4.4 (Universal bound). Let $A_0, \ldots, A_d, P, \zeta_1, \ldots, \zeta_{nd}, \|\cdot\|, c$, and $\alpha_1, \ldots, \alpha_d, \bar{\alpha}_1, \ldots, \bar{\alpha}_{nd}$ be as in the first part of Theorem 4.1. Then, for all $K \in \{1, \ldots, nd\}$, Inequality (21) holds with $L_k, 0 < k \le d$, defined as follows:

$$(26) L_k = \frac{1}{\mathcal{E}(k)(k+1)} ,$$

where \mathcal{E} is defined as in (4). Moreover, when $\|\cdot\|$ satisfies (A2), Inequality (21) holds with the greater constant:

(27)
$$L_k = \frac{1}{\sqrt{\mathcal{E}(k)(k+1)}}.$$

The lower bound of Pólya, and its matrix version given above, are tight only for k small. By symmetry, one can obtain a tight lower bound when k is close to d (this was already noted by Ostrowski in the scalar case [30]). Theorem 4.5 below will allow us to obtain a tight lower bound when k lies "in the middle" of the interval [0,d], by using the comparison between the tropical roots. Unlike the lower bound of Pólya, this theorem gives a bound which is not anymore independent of the coefficients of the polynomial P, although it depends only on a small information, namely a ratio measuring the *separation* between some tropical roots. Thus, the bound will involve a coefficient $\mathcal{U}(k,\delta)$ depending both on the index k and on the parameter δ (the ratio). It will be specially useful in situations in which δ is small. This coefficient $\mathcal{U}(k,\delta)$ is defined as follows:

(28)
$$\mathcal{U}(k,\delta) := \mathcal{E}(k) \left(\frac{1+\delta+\sqrt{k^2(1-\delta)^2+4\delta}}{1-\delta} \right)$$
, for $k \geqslant 0$ and $0 \leqslant \delta \leqslant 1$,

where $\mathcal{E}(k)$ is defined by (4) for k > 0, and $\mathcal{E}(0) := 1$, and with the convention that $1/0 = +\infty$, so that $\mathcal{U}(k,1) = +\infty$. It is easy to check that

(29)
$$k + \frac{1+\delta}{1-\delta} \le \mathcal{U}(k,\delta) < e\left(k + \frac{1+\sqrt{\delta}}{1-\sqrt{\delta}}\right).$$

The following asymptotic regime should be kept in mind:

$$\mathcal{U}(k,\delta) \sim \mathcal{E}(k)(k+1) < e(k+1), \qquad \delta \to 0$$

Theorem 4.5 (Master lower bound). Let $A_0, \ldots, A_d, P, \zeta_1, \ldots, \zeta_{nd}, \|\cdot\|, c$, and $\alpha_1, \ldots, \alpha_d, \bar{\alpha}_1, \ldots, \bar{\alpha}_{nd}$ be as in the first part of Theorem 4.1, and denote $\alpha_0 = 0$ and $\alpha_{d+1} = +\infty$.

Let $K \in \{1, ..., nd\}$, k^-, k^+ be integers such that $0 \le k^- < \lceil \frac{K}{n} \rceil \le k^+ - 1 \le d$, and denote $\delta_- := \alpha_{k^-} / \alpha_{\lceil \frac{K}{n} \rceil} \le 1$ and $\delta_+ := \alpha_{\lceil \frac{K}{n} \rceil} / \alpha_{k^+} \le 1$. Then, Inequality (21) holds with L_k , $k^- < k \le k^+ - 1$, defined as follows:

(30)
$$L_k := \max(L_k^{\pm}), \quad L_k^+ := \frac{1}{\mathcal{U}(k^+ - k - 1, \delta_+)}, \ L_k^- := \frac{1}{\mathcal{U}(k - k^-, \delta_-)},$$

with the convention that $1/\infty = 0$. Moreover, when $\|\cdot\|$ satisfies (A2), the constant L_k of (21) can be replaced by the greater constant L_k^* :

$$(31) \ L_k^* := \max(L_k^{\pm *}), \quad L_k^{+*} := \frac{1}{\sqrt{\mathcal{U}(k^+ - k - 1, \delta_+^2)}}, \ L_k^{-*} := \frac{1}{\sqrt{\mathcal{U}(k - k^-, \delta_-^2)}} \ .$$

Theorem 4.5 generalizes Theorem 4.4, and thus the lower bound of Pólya. Indeed, taking $k^- = 0$, $k^+ = d + 1$, and using that $\mathcal{U}(k,0) = \mathcal{E}(k)(k+1)$, we get that the constants $L_k^- \leq L_k$ and $L_k^{-*} \leq L_k^*$ of Theorem 4.5 are exactly the constants L_k of (26) and (27) of Theorem 4.4 respectively. Moreover, Theorem 4.5 is already new in the scalar case (n = 1).

Note that when $\delta_{-}=1$ in Theorem 4.5, $\mathcal{U}(k-k^{-},\delta_{-})=+\infty$, so that $L_{k}^{-}=0$ and $L_{k}=L_{k}^{+}$. Similarly when $\delta_{+}=1$, we get $L_{k}^{+}=0$. Hence, if $\delta_{-}=1=\delta_{+}$, we get $L_{k}=0$ so that (21) does not provide any information, although it is true. However, if for instance $\delta_{-}=1$ and $\delta_{+}<1$, we get that $L_{k}=L_{k}^{+}>0$, which gives a positive lower bound in (21).

Applying Theorem 4.5 to the particular case when $\frac{K}{n} = k = k^- + 1 = k^+ - 1$ leads to the following formula for the constants of (30) and (31):

(32)
$$L_k := \max(L_k^{\pm}), \quad L_k^+ := \frac{1 - \sqrt{\delta_+}}{1 + \sqrt{\delta_+}}, \ L_k^- := \frac{1 - \delta_-}{4(1 + \delta_-)},$$

and

(33)
$$L_k^* := \max(L_k^{\pm *}), \quad L_k^{+*} := \sqrt{\frac{1 - \delta_+}{1 + \delta_+}}, \ L_k^{-*} := \frac{1}{2} \sqrt{\frac{1 - \delta_-^2}{1 + \delta_-^2}} \ .$$

In this special case, we obtain the following stronger lower bounds.

Theorem 4.6. Let us use the notations of Theorem 4.5, and assume that $\frac{K}{n} = k = k^- + 1 = k^+ - 1$. Then, the statements of Theorem 4.5 hold with Inequality (22) instead of (21) and the constants L_k and L_k^* (given in (30) and (31), or (32) and (33)) replaced respectively by the greater constants L_k^{\sharp} and $L_k^{\sharp\sharp}$ given by:

(34)
$$L_k^{\sharp} := \frac{1 - \delta_{-} \delta_{+}}{(1 + \sqrt{\delta_{+}})^2} ,$$

and

(35)
$$L_k^{*\sharp} := \frac{\sqrt{1 - \delta_-^2 \delta_+^2}}{1 + \delta_+} .$$

We next indicate how the indices k^+ and k^- should be chosen, for each $K \in \{1, \ldots, nd\}$, in order to get the best lower bound $L_{\underline{K}}$.

Let us use the notations of Theorem 4.5, so that $\alpha_1, \ldots, \alpha_d$ are the tropical roots of the tropical polynomial tp of (6), $\alpha_0 = 0$ and $\alpha_{d+1} = +\infty$. Let $k_0 = 0, k_1, \ldots, k_q = d$ be the sequence of abscissæ of the vertices of the Newton polygon of tp(x), as shown in Figure 2. For $j = 1, \ldots, q$, we have, $\alpha_{k_{j-1}+1} = \cdots = \alpha_{k_j} < \alpha_{k_j+1}$. We also denote by

(36)
$$\delta_j = \frac{\alpha_{k_j}}{\alpha_{k_j+1}} ,$$

for $j=0,\ldots,q$, the parameters measuring the *separation* between the tropical roots, in particular $\delta_0=\delta_q=0$.




FIGURE 2. Newton polygon corresponding to t p(x).

Proposition 4.7. Each of the constants L_k and L_k^* appearing in Theorem 4.5, is maximized by choosing $k^- = k_r$ for some $0 \le r \le q$ such that $k_r < k$ and $k^+ = k_s + 1$ for some $0 \le s \le q$ such that $k_s \ge k$.

This proposition shows that, to apply Theorem 4.5, we may always require k^- and k^+-1 to be abscissæ of vertices of the Newton polygon. Optimizing the choice of k^\pm , we readily arrive at the following corollary.

Corollary 4.8. Let us use the notations of Theorem 4.5, and let $k_0 = 0, k_1, \ldots, k_q = d$ be the sequence of abscissæ of the vertices of the Newton polygon of tp(x), as shown in Figure 2. Then, the statements of Theorem 4.5 hold with the constants L_k , L_k^{\pm} , L_k^{\star} and $L_k^{\pm \star}$ (given in (30) and (31)) replaced respectively by the following optimal ones:

(37a)
$$L_k^{opt} := \max(L_k^{\pm, opt}),$$

(37b)
$$L_k^{+,opt} := \max_{j: k_j \geqslant k} \frac{1}{\mathcal{U}(k_j - k, \frac{\alpha_{\lceil k \rceil}}{\alpha_{k_j + 1}})},$$

(37c)
$$L_k^{-,opt} := \max_{j: k_{j-1} < k} \frac{1}{\mathcal{U}(k - k_{j-1}, \frac{\alpha_{k_{j-1}}}{\alpha_{\lceil k \rceil}})} ,$$

and

(38a)
$$L_k^{*,opt} := \max(L_k^{\pm *,opt}),$$

(38b)
$$L_k^{+*,opt} := \max_{j: k_j \geqslant k} \frac{1}{\sqrt{\mathcal{U}(k_j - k, (\frac{\alpha_{\lceil k \rceil}}{\alpha_{k_j + 1}})^2)}},$$

(38c)
$$L_k^{-*,opt} := \max_{j: k_{j-1} < k} \frac{1}{\sqrt{\mathcal{U}(k - k_{j-1}, (\frac{\alpha_{k_{j-1}}}{\alpha_{\lceil k \rceil}})^2)}}.$$

However, a simpler choice of k^{\pm} consists in taking the nearest vertices of the Newton polygon in (37) and (38), which lead to the following corollary.

Corollary 4.9. Let us use the notations of Theorem 4.5, let $k_0 = 0, k_1, \ldots, k_q = d$ be the sequence of abscissæ of the vertices of the Newton polygon of tp(x), as shown in Figure 2, and let $\delta_0, \ldots, \delta_q$ be defined by (36). For $K \in \{1, \ldots, nd\}$ and $k = \frac{K}{n}$, let us consider the unique $j \in \{1, \ldots, q\}$ such that $k_{j-1} < k \le k_j$, so that $\alpha_{\lceil k \rceil} = \alpha_{k_{j-1}+1} = \alpha_{k_j}$. Then, the statements of Theorem 4.5 hold with the constants L_k , L_k^{\pm} , L_k^{*} and $L_k^{\pm *}$ (given in (30) and (31)) replaced respectively by the following ones:

(39a)
$$L_k^{prox} := \max(L_k^{\pm, prox}),$$

(39b)
$$L_k^{+,prox} := \frac{1}{\mathcal{U}(k_i - k, \delta_i)}, \quad L_k^{-,prox} := \frac{1}{\mathcal{U}(k - k_{i-1}, \delta_{i-1})},$$

and

$$(40a) L_k^{*,prox} := \max(L_k^{\pm *,prox}),$$

(40b)
$$L_k^{+*,prox} := \frac{1}{\sqrt{\mathcal{U}(k_j - k, \delta_j^2)}}, \quad L_k^{-*,prox} := \frac{1}{\sqrt{\mathcal{U}(k - k_{j-1}, \delta_{j-1}^2)}}$$
.

Moreover, in the particular case where $k = k_i$, we have

$$L_k^{prox} \geqslant L_k^{+,prox} = \frac{1 - \sqrt{\delta_j}}{1 + \sqrt{\delta_j}}, \qquad L_k^{*,prox} \geqslant L_k^{+*,prox} = \sqrt{\frac{1 - \delta_j}{1 + \delta_j}} \ .$$

Remark 4.10. When all the ratios $\delta_1, \ldots, \delta_{q-1}$ are small, the maxima in (37) and (38) are attained by taking j as in Corollary 4.9, that is $L_k^{\text{opt}} = L_k^{\text{prox}}$ and $L_k^{*,\text{opt}} = L_k^{*,\text{prox}}$ for all $0 < k \le d$.

Remark 4.11. Since $\mathcal{U}(k,\delta)$ is increasing in k and δ , the maximizing j in the definition of $L_k^{+*,\mathrm{opt}}$ arises from a compromise between keeping $k_j \geqslant k$ close to k and δ_j small. In particular, when k belongs to an edge of the Newton polygon such that several consecutive edges have almost the same slope than this edge, the maximizing j may be the one corresponding to the first vertex at which the slope changes significantly, i.e., the first one such that δ_j is small. Similar considerations apply to $L_k^{-*,\mathrm{opt}}$

5. Proof of the lower bounds

5.1. Proof of the lower bounds for the roots of scalar polynomials. In this section, we prove the main lower bounds of Section 4 for scalar polynomials, since the arguments are more transparent in this case. The generalization to the matrix case will be given in the next section.

From Remark 4.2, in the scalar case, all lower bounds reduce to (25) with the constant L_k obtained under Assumption (A2), that is (24) in Theorem 4.1, (27) in Theorem 4.4, (31) in Theorem 4.5, and (35) in Theorem 4.6.

Proof of the scalar version of Theorem 4.4. We note first that Pólya's inequality is an immediate consequence of Proposition 3.5. Indeed, using the property that all the $\beta_{k,j}$ are less than or equal to 1, we obtain:

$$(L_k)^{-2} \leqslant \sum_{j=0}^d \xi^{j-k} \leqslant \sum_{j=0}^\infty \xi^{j-k} = \frac{1}{\xi^k (1-\xi)}$$
,

for all $\xi > 0$. The minimum of the right hand side of the previous inequality for $0 < \xi < 1$, is attained for $\xi = k/(k+1)$, from which we deduce (3), which is also the scalar version of Theorem 4.4.

We can now deduce similarly, from Proposition 3.5, the scalar versions of the lower bounds of Theorems 4.1, 4.5, and 4.6. Since in the scalar case, we are reduced to show (25) for some constants L_k , and that this inequality is precisely the statement of Proposition 3.5, we only need to show that these constants L_k are lower bounds of the constant L_k of (16).

Proof of the scalar version of Theorem 4.1. Using the property that $\beta_{k,j} \leq 1$ for all k, j, we obtain that the constant L_k of (16) satisfies $(L_k)^{-2} \leq \sum_{j=0, a_j \neq 0}^d \xi^{j-k}$ for all $\xi > 0$. When $\xi = 1$, the right hand side of this inequality is equal to the number of non-zero coefficients of p, which shows that the constant L_k of (16) is lower bounded by the constant L_k of (24), which implies (25) with this lower bound L_k .

Proof of the scalar version of Theorem 4.5. Assume that $0 \le k^- < k \le k^+ - 1 \le d$ are integers. Denote $\delta_- = \alpha_{k^-}/\alpha_k$ and $\delta_+ = \alpha_k/\alpha_{k^+}$. Then, for all $1 \le \ell \le k$, $\alpha_\ell \le \alpha_k$, and for all $1 \le \ell \le k^-$, $\alpha_\ell \le \alpha_{k^-} = \alpha_k \delta_-$. This implies that $\beta_{k,j} \le \delta_-^{k^- - j}$, for

all $j \leq k^-$. Similarly, for all $\ell \geq k$, $\alpha_{\ell} \geq \alpha_k$, and for all $\ell \geq k^+$, $\alpha_{\ell} \geq \alpha_{k^+} = \alpha_k/\delta_+$, hence, for all $j \geq k^+$, $\beta_{k,j} \leq \delta_+^{j-k^++1}$. Since we also have $\beta_{k,j} \leq 1$ for all j,k, we obtain that the constant L_k of (16) satisfies, for all $\delta_-^2 < \xi < \delta_+^{-2}$, $\xi \neq 1$,

$$(L_k)^{-2} \leqslant \left(\sum_{j=0}^{k^-} \delta_-^{2(k^- - j)} \xi^{j-k} \right) + \left(\sum_{j=k^- + 1}^{k^+ - 1} \xi^{j-k} \right) + \left(\sum_{j=k^+}^{d} \delta_+^{2(j-k^+ + 1)} \xi^{j-k} \right)$$

$$\leqslant \xi^{k^- - k + 1} \frac{1}{\xi - \delta^2} + \frac{\xi^{k^- - k + 1} - \xi^{k^+ - k}}{1 - \xi} + \xi^{k^+ - k - 1} \frac{\delta_+^2 \xi}{1 - \delta_+^2 \xi} .$$

The later inequality can be written as

(41)
$$(L_k)^{-2} \leqslant g(\xi), \quad \forall \delta_-^2 < \xi < \delta_+^{-2}, \ \xi \neq 1,$$

with

$$g(\xi) := g_{-}(\xi) + g_{+}(\xi) ,$$

$$g_{-}(\xi) := \xi^{k^{-}-k+1} \left(\frac{1}{\xi - \delta_{-}^{2}} + \frac{1}{1 - \xi} \right) ,$$

$$g_{+}(\xi) := \xi^{k^{+}-k-1} \left(\frac{1}{\xi - 1} + \frac{1}{1 - \delta_{+}^{2} \xi} \right) .$$

Note that (41) also holds when $k^- = 0$ or $k^+ = d + 1$, since then $\delta_- = 0$ or $\delta_+ = 0$ respectively. When $\delta_- = 1$ and $\delta_+ < 1$, the conditions on ξ in (41) are equivalent to $1 < \xi < \delta_+^{-2}$, whereas when $\delta_- = \delta_+ = 1$, these conditions are never satisfied, but in this case the constant L_k of (31) is equal to 0 so there is nothing to prove.

The functions g_- and g_+ satisfy $g_-(\xi) = g_{k-k-,\delta_-^2}(\xi^{-1})$ and $g_+(\xi) = g_{k+-k-1,\delta_+^2}(\xi)$ where for $k \ge 0$ and $0 \le \delta \le 1$, $g_{k,\delta}$ is defined as a function of $\xi \in (0,\delta^{-1}) \setminus \{1\}$ by:

$$g_{k,\delta}(\xi) := \xi^k \left(\frac{1}{\xi - 1} + \frac{1}{1 - \delta \xi} \right) = \xi^{k+1} \frac{1 - \delta}{(\xi - 1)(1 - \delta \xi)}$$
.

We have $g_{k,\delta}(\xi) \leq 0$ for $0 < \xi < 1$, hence $g_{-}(\xi) \leq 0$ for all $\xi > 1$, and $g_{+}(\xi) \leq 0$ for all $\xi < 1$. When $0 \leq \delta < 1$, the minimum of $g_{k,\delta}$ on $(1, \delta^{-1})$ is attained at

$$\xi_{k,\delta} := \begin{cases} \frac{k(\delta+1) - \sqrt{k^2(\delta+1)^2 - 4\delta(k^2 - 1)}}{2(k-1)\delta} & \text{when } k \neq 1, \ \delta \neq 0 \\ \frac{k+1}{k(1+\delta)} & \text{otherwisewhen } \delta = 0 \end{cases}.$$

The last formula gives $\xi_{k,\delta} = \infty$ when $\delta = 0$ and k = 0, which is the point of infimum of $g_{k,\delta}$ for $1 < \xi < \infty = \delta^{-1}$, since $g_{k,\delta}$ is decreasing. It also gives $\xi_{k,1} = 1$ for all $k \ge 0$. Hence extending $g_{k,\delta}$, g_- and g_+ by $+\infty$ at point 1, we get that the infimum of $g_{k,\delta}$ on $(1,\delta^{-1})$ equals $g_{k,\delta}(\xi_{k,\delta})$, and denoting $\xi_- = (\xi_{k-k^-,\delta^-_-})^{-1}$ and $\xi_+ = \xi_{k^+-k^-1,\delta^2_+}$, we obtain

(42)
$$(L_k)^{-2} \leqslant \min(g(\xi_-), g(\xi_+)) \leqslant \min(g_-(\xi_-), g_+(\xi_+)) .$$

In order to simplify this bound, we need to find good estimates of $\xi_{k,\delta}$. When $k \neq 1$ and $\delta \neq 0$, we have:

$$\xi_{k,\delta} = \frac{k(\delta+1) - \sqrt{k^2(1-\delta)^2 + 4\delta}}{2(k-1)\delta} = \frac{2(k+1)}{k(\delta+1) + \sqrt{k^2(1-\delta)^2 + 4\delta}} \ .$$

Using the property that $\delta \geqslant 0$ in the last formula, we get that $\xi_{k,\delta} \leqslant \frac{k+1}{k} = \xi_{k,0}$. Moreover, this inequality also holds for k = 1. In particular $(\xi_{k,\delta})^k \leqslant \mathcal{E}(k)$, for all $k \geqslant 0$ and $\delta \geqslant 0$ (taking the convention that $\xi^0 = 1$ for all $\xi \in (1, \infty]$). We also have, for $k \geqslant 0$ and $0 \leqslant \delta < 1$,

$$\frac{1}{\xi_{k,\delta} - 1} + \frac{1}{1 - \delta \xi_{k,\delta}} = \frac{1 + \delta + \sqrt{k^2 (1 - \delta)^2 + 4\delta}}{1 - \delta} \leqslant k + \frac{1 + \sqrt{\delta}}{1 - \sqrt{\delta}}.$$

This yields

$$g_{k,\delta}(\xi_{k,\delta}) \leqslant \mathcal{U}(k,\delta) \quad \forall k \geqslant 0, \ 0 \leqslant \delta \leqslant 1,$$

with \mathcal{U} as in (28). Indeed, this inequality holds for $\delta < 1$ by the above arguments. It also holds for $\delta = 1$ since $\mathcal{U}(k,1) = +\infty$. This implies in particular that $g_+(\xi_+) \leq \mathcal{U}(k^+ - k - 1, \delta_+^2)$ and $g_-(\xi_-) \leq \mathcal{U}(k - k^-, \delta_-^2)$. Combining these two inequalities with (42), we obtain that the constant L_k of (16) is lower bounded by the constant L_k^* of (31), which implies (25) with this lower bound L_k^* instead of L_k .

Remark 5.1. In the previous proof, obtaining the minimum of g instead of g_- and g_+ would have led to a better lower bound for L_k than the one of Theorem 4.5. However, such a bound is more difficult to estimate for general values of k^+-k^- . We can use for instance the first inequality in (42), which will give better estimates in some particular cases. For instance, when k=0, we get that $g_{k,\delta}(\xi_{k,\delta})=\mathcal{U}(k,\delta)=(1+\sqrt{\delta})/(1-\sqrt{\delta})$. In particular, when $k=k^+-1$, the bound $(L_k)^{-2} \leq g_+(\xi_+)$ gives the lower bound L_k^+ of (33), with $\delta=\delta_+$. However (42) gives the slightly better bound:

$$(L_k)^{-2} \leqslant g(\xi_+) = \frac{1+\delta_+}{1-\delta_+} - \frac{1-\delta_-^2}{(1-\delta_+\delta_-^2)(1-\delta_+)} \delta_+^{k^+-k^-}.$$

Proof of the scalar version of Theorem 4.6. Let us use the notation of the previous proof. When $k=k^+-1=k^-+1$, Inequality (41) is true for all $\xi\in(\delta_-^2,\delta_+^{-2})$ since g can be well defined at 1 (by continuity for instance). When $\delta_-\delta_+<1$, the interval $(\delta_-^2,\delta_+^{-2})$ is nonempty, the map g is convex there and its minimum is achieved at the point $\xi=(1+\delta_-^2\delta_+)/(\delta_+(1+\delta_+))$. Then (41) yields $(L_k)^{-2}\leqslant(1+\delta_+)^2/(1-\delta_-^2\delta_+^2)$, which implies that the constant L_k of (16) is lower bounded by the constant $L_k^{*\sharp}$ of (35), which implies (25) with this lower bound $L_k^{*\sharp}$ instead of L_k . Moreover, since the minimum of g is less or equal to the right hand side of (42), the proof of the scalar version of Theorem 4.5 implies that $L_k^{*\sharp}\leqslant L_k^*$.

5.2. Proofs of the lower bounds for the eigenvalues of matrix polynomials. In this section, we prove the lower bounds which were stated in Section 4 for matrix polynomials, and proved in Section 5.1 in the special scalar case.

Proof of Theorems 4.1, 4.4, 4.5, and 4.6. In Proposition 3.6, the formula (20) of L_k coincides with (16), except that the condition $a_j \neq 0$ is replaced by $A_j \neq 0$ and $\beta_{k,j}$ is replaced by $\beta_{\lceil k \rceil,j}$. For $0 < k \le d$, and for integers k^- and k^+ , the inequality $0 \le k^- < k \le k^+ - 1 \le d$ is equivalent to $0 \le k^- < \lceil k \rceil \le k^+ - 1 \le d$. Then, for all $0 < k \le d$, the inequalities obtained for $\beta_{\lceil k \rceil,j}$ in the proofs of Theorems 4.1, 4.5, and 4.6 remain true and depend only on k^- and k^+ and not on $\lceil k \rceil$. Hence, the lower bounds of the constant L_k of (16) obtained in these proofs remain true for the constant L_k of (20). Combined with the assertion that Inequality (21) holds with this constant, they now provide the lower bounds of Theorems 4.1, 4.5, and 4.6 respectively, in the case where $\|\cdot\|$ satisfies (A2), that is Inequality (21) with L_k replaced by the constant L_k of (24), the constant L_k^* of (31), and the constant L_k^* of (35), respectively.

In the case of a norm satisfying only (A1), the formula (20) of L_k is replaced by (19), which means that everything behave as if L_k were replaced by its square and the numbers α_k were replaced by their square roots in (20). Hence the assertions of Theorems 4.1, 4.5, and 4.6 in the case of Assumption (A2), are still true under Assumption (A1), up to this transformation of the constant L_k of (24), the constant L_k^* of (31), and the constant L_k^* of (35), respectively.

The constant L_k of (24) does not depend on the values of the constants α_k , hence the above transformation leads to the square of L_k , which is exactly the constant L_k of (23), which finishes the proof of Theorem 4.1. The constant L_k^* of (31) is a function of $\delta_- = \frac{\alpha_{k^-}}{\alpha_{\lceil k \rceil}}$ and $\delta_+ = \frac{\alpha_{\lceil k \rceil}}{\alpha_{k^+}}$. Since replacing all numbers α_i , $i=0,\ldots,d+1$, by their square roots, reduces to replace the numbers δ_- and δ_+ by their square roots too, the above transformation of the constant L_k^* of (31) consists in taking its square and replacing δ_- and δ_+ by their square roots, which leads to the constant L_k of (30), which finishes the proof of Theorem 4.5. Similarly, the above transformation of the constant L_k^* of (35) leads to the constant L_k^* of (34), which finishes the proof of Theorem 4.6.

Now Theorem 4.4 is an immediate corollary of Theorem 4.5. Indeed as said in Section 4, taking $k^- = 0$, $k^+ = d+1$ in Theorem 4.5, and using $L_k^- \leq L_k$, $L_k^{-*} \leq L_k^*$ and $\mathcal{U}(k,0) = \mathcal{E}(k)(k+1)$, we get Theorem 4.4.

Corollaries 4.8 and 4.9 are straightforward consequences of Theorem 4.5 and Proposition 4.7. We only need to prove the latter proposition.

Proof of Proposition 4.7. Let us use the notations of Theorem 4.5, and let $k_0 = 0, k_1, \ldots, k_q = d$ be the sequence of abscissæ of the vertices of the Newton polygon of $\operatorname{t} p(x)$, as shown in Figure 2. Then, the ratio δ_- does not change when k^- moves inside an interval $[k_{r-1}+1,k_r]$. Since $\mathcal{U}(\cdot,\delta)$ is a nondecreasing function, in order to maximize L_k^- or L_k^{-*} and thus L_k , with δ_- constant (and δ_+ and k^+ constant), we need to minimize $k-k^-$, which implies that $k^-=k_r$ for some $0\leqslant r\leqslant q$ such that $k_r< k$. Similarly, $k^+=k_s+1$ for some $0\leqslant s\leqslant q$ such that $k_s\geqslant k$.

Remark 5.2. All the lower bounds in Theorems 4.1, 4.4, 4.5, and 4.6 for K=nk are equivalent to inequalities of the form

(43)
$$(|\zeta_1 \dots \zeta_{nk}|^{-1}|\det A_0|)^{1/n}L_k \leqslant ||A_0||\alpha_1^{-1} \dots \alpha_k^{-1} ...$$

By definition of the tropical roots, $\|A_0\|\alpha_1^{-1}\dots\alpha_k^{-1}=\exp\hat{p}_k$, where \hat{p}_k is the value in k of the concave hull of the map $j\in\{0,\dots,d\}\mapsto\log\|A_j\|$. Hence, if $\|\cdot\|$ and $\|\cdot\|'$ are two norms on the space of matrices such that $\|A\|\leqslant\|A\|'$ for all $A\in\mathbb{C}^{n\times n}$, then the right hand side in (43) is smaller for $\|\cdot\|$ than for $\|\cdot\|'$. Moreover, in (23), (24), (26), and (27), L_k depends only on P or k and n, but not on the norm $\|\cdot\|$, so that (43) is necessarily a tighter inequality for $\|\cdot\|$ than for $\|\cdot\|'$. In particular if (A2) holds with Q and Q', then the lower bounds (24) and (27) for $\|\cdot\|$ are weaker than the corresponding ones for $A\mapsto\|QAQ'\|_{\mathrm{nF}}$. For the lower bounds (30), (31), (34), and (35) of Theorems 4.5 and 4.6, and the ones of Corollaries 4.8 and 4.9, based on the separation between the tropical roots, the comparison is not so simple, because changing the norm changes the separation between the tropical roots.

6.1. Statement of the upper bound and corollary. The previous results provide lower bounds for the product of the K smallest eigenvalues, in terms of tropical roots. We next state a reverse inequality.

Theorem 6.1 (Upper bound). Let $\|\cdot\|$ be any norm on the space of matrices $\mathbb{C}^{n\times n}$. For all $i=1,\ldots,n$, we denote by $A^{(i)}$ the i-th column of A, by η_i the least positive constant such that $\|A^{(i)}\|_2 \leq \eta_i \|A\|$ for all $A \in \mathbb{C}^{n\times n}$ where $\|\cdot\|_2$ is the Euclidean norm of \mathbb{C}^n , and we set $\eta:=\eta_1\cdots\eta_n$. Let A_0,\ldots,A_d , $P,\zeta_1,\ldots,\zeta_{nd}$, $c,\mathsf{t}p,\alpha_1,\ldots,\alpha_d$, be as in Theorem 4.4, let $k_0=0,k_1,\ldots,k_q=d$ be the sequence of all abscissæ of the vertices of the Newton polygon of $\mathsf{t}p(x)$, and define δ_0,\ldots,δ_q as in (36). Denote by $C_{n,d,k}$ the number of maps $\phi:\{1,\ldots,n\}\to\{0,\ldots,d\}$ such that $\sum_{j=1}^n \phi(j)=k$, so that $C_{n,d,k} \leq (\min(d,k,nd-k)+1)^{n-1}$.

Then, for every $j = 1, \ldots, q$, if

(44)
$$c_j := \frac{|\det A_{k_j}|}{\|A_{k_j}\|^n} - (C_{n,d,nk_j} - 1)\eta \delta_j > 0 ,$$

we have

$$\frac{|\zeta_1 \dots \zeta_{nk_j}|}{(\alpha_1 \dots \alpha_{k_j})^n} \leqslant \frac{c}{c_j} \binom{nd}{nk_j} .$$

Using the results of Section 4 and Theorem 6.1, we are able to bound from above and below the ratio between the modulus of an eigenvalue and its corresponding tropical root, as follows.

Corollary 6.2. Let us use the notations of Theorems 4.1 and 6.1, and assume that (21) holds for all $K \in \{1, ..., nd\}$. Then, under the assumptions of Theorem 6.1, we have, for all $1 \le j \le q$, such that $c_j > 0$ and $c_{j-1} > 0$, and all integer K such that $k_{j-1} < \frac{K}{r} \le k_j$

$$(45) \qquad \frac{c_{j-1}}{\binom{nd}{nk_{j-1}}} (L_{k_{j-1} + \frac{1}{n}})^n \leqslant \frac{|\zeta_{nk_{j-1} + 1}|}{\alpha_{k_j}} \leqslant \frac{|\zeta_K|}{\alpha_{k_j}} \leqslant \frac{|\zeta_{nk_j}|}{\alpha_{k_j}} \leqslant \frac{\binom{nd}{nk_j}}{c_j} (L_{k_j - \frac{1}{n}})^{-n} .$$

In particular, if $n \ge 2$ and (A2) holds, we have:

$$(46) \qquad \frac{c_{j-1}}{\binom{nd}{nk_{j-1}}\mathcal{U}(\frac{1}{n},\delta_{j-1}^2)^{n/2}} \leqslant \frac{|\zeta_K|}{\alpha_{k_j}} \leqslant \frac{\binom{nd}{nk_j}\mathcal{U}(\frac{1}{n},\delta_j^2)^{n/2}}{c_j} .$$

Note that the bounds in the above corollary do not depend on the constant c, except if j=1 (in which case $c_{j-1}=c_0=c$). Hence, by an argument of continuity, the assumption that $\det A_0 \neq 0$, which is present there (since the notations and assumptions of Theorem 6.1 include the ones of the first part of Theorem 4.1), can be dispensed with, except when j=1.

Remark 6.3. Since $1 \geqslant \frac{|\det A_{k_j}|}{\|A_{k_j}\|^n} \geqslant (\kappa(A_{k_j}))^{-n}$, where κ denotes the condition number of a matrix with respect to the norm $\|\cdot\|$, we get that Condition (44) of Theorem 6.1 holds if the matrix A_{k_j} is nonsingular and has a sufficiently small condition number, and if the tropical roots α_{k_j} and $\alpha_{k_{j+1}}$ are sufficiently separated, so that $\delta_j = \alpha_{k_j}/\alpha_{k_{j+1}} \ll 1$. Similarly, the tightness of the bounds in Corollary 6.2 depends on the parameters δ_{j-1}, δ_j and on the condition number of the matrices $A_{k_{j-1}}, A_{k_j}$.

Remark 6.4. The bounds of Corollary 6.2 require the δ_j to be very small. In this case, a different approach would be to use the extensions of Pellet's theorem to the case of matrix polynomials [10, 28, 29].

Recall that in the scalar case, Pellet's theorem shows that for $k \notin \{0,d\}$, if the polynomial $q(x) = |a_k| x^k - \sum_{i \neq k, \ 0 \leqslant i \leqslant d} |a_i| x^i$ has precisely two real positive roots r(a) < R(a) (we know by Descartes' rule of sign it has 0 or 2 real positive roots), then p has precisely k roots in the region $|x| \leqslant r(a)$ and d-k roots in the region $|x| \geqslant R(a)$. Moreover, in [9], it is shown that in that case, k is the abscissa of a vertex of the Newton polygon of p, and that tropical roots of p are outside the interval $(\log r(a), \log R(a))$. If Pellet's theorem can be applied to every abscissa k of a vertex of the Newton polygon of p, then each moduli of a root of p can be estimated using the corresponding tropical root. This is in particular possible if $\delta_j < 1/9$ for all $j = 1, \ldots, q$, by a result of Gaubert and Sharify (see [35, Theorem 3.3.3]).

Using an extension of Pellet's theorem to the case of matrix polynomials, one may obtain a similar localisation of the eigenvalues when all the δ_j and condition numbers of the matrices A_{k_j} are small enough, see for instance [10] for the case where the coefficient matrices are unitary up to some constants, and [29] for the general case.

However, it may happen that Pellet's theorem cannot be applied for any vertex of the Newton polygon except the first and last one. In this case, the log-majorization type inequalities of Section 4 do provide a better information. Note that there is some duality between Pellet type estimates and log-majorization estimates: the former hold conditionnally, and give exclusion annuli, whereas the latter hold inconditionnally, and allow one to infer inclusion regions for the eigenvalues. For instance, it does not seem that tight constants like the one in (3) are achievable by Pellet type approaches.

6.2. **Proof of the upper bound.** To prove Theorem 6.1, we first prove the following lemma which provides a lower bound for the modulus of the coefficients of the polynomial $\det(P(\lambda))$.

Lemma 6.5. Let $A_0, \ldots, A_d, P, \|\cdot\|, \eta, \alpha_1, \ldots, \alpha_d, q, \delta_0, \ldots, \delta_q$, and $C_{n,d,k}$ be as in Theorem 6.1, and denote by \tilde{p} the polynomial:

(47)
$$\tilde{p}(\lambda) = \det P(\lambda) = \sum_{l=0}^{nd} \tilde{p}_l \lambda^l .$$

Then, for $j = 0, \ldots, q$, we have $||A_{k_i}|| > 0$ and

$$|\tilde{p}_{nk_i}| \geqslant |\det A_{k_i}| - (C_{n,d,nk_i} - 1)\eta ||A_{k_i}||^n \delta_i$$
.

Proof. Let $k = 0, \ldots, d$. Using the multilinearity of the determinant we get

$$\tilde{p}(\lambda) = \det \sum_{j=0}^{d} A_j \lambda^j = \sum_{\phi} \det(A_{\phi(1)}^{(1)}, A_{\phi(2)}^{(2)}, \dots, A_{\phi(n)}^{(n)}) \lambda^{\sum_{m=1}^{n} \phi(m)} ,$$

where the sum is taken over all maps $\phi: \{1, \ldots, n\} \to \{0, \ldots, d\}$. Denoting by Φ_k the set of all such maps that satisfy $\sum_{m=1}^{n} \phi(m) = nk$, we obtain that the nk-th

coefficient of the polynomial \tilde{p} is equal to:

$$\tilde{p}_{nk} = \sum_{\phi \in \Phi_k} \det(A_{\phi(1)}^{(1)}, A_{\phi(2)}^{(2)}, \dots, A_{\phi(n)}^{(n)})
= \det A_k + \sum_{\phi \in \Phi_k, \phi \neq k} \det(A_{\phi(1)}^{(1)}, A_{\phi(2)}^{(2)}, \dots, A_{\phi(n)}^{(n)}) .$$

Using Hadamard's inequality together with the definition of η yields

$$|\det(A_{\phi(1)}^{(1)},A_{\phi(2)}^{(2)},\dots,A_{\phi(n)}^{(n)})| \leqslant \|A_{\phi(1)}^{(1)}\|_2\dots\|A_{\phi(n)}^{(n)}\|_2 \leqslant \eta \|A_{\phi(1)}\|\dots\|A_{\phi(n)}\|.$$

Assume now that $k=k_j$ for some $j=0,\ldots,q$. Then, k is the abscissa of a vertex of the Newton polygon of the tropical polynomial tp defined in (6). By Proposition 3.4 applied to tp, we get that $\|A_k\| > 0$, $\|A_m\| \le \|A_k\| \prod_{\ell=m+1}^k \alpha_\ell \le \|A_k\| \alpha_k^{k-m}$ for all $m \le k$ and $\|A_m\| \le \|A_k\| \prod_{\ell=k+1}^m \alpha_\ell^{-1} \le \|A_k\| \alpha_{k+1}^{k-m} \le \|A_k\| \alpha_k^{k-m} \delta_j^{m-k}$ for all m > k. Hence

$$||A_{\phi(1)}|| \dots ||A_{\phi(n)}|| \leqslant ||A_k||^n \alpha_k^{\sum_{m=1}^n (k-\phi(m))} \delta_j^{\sum_{m=1,\phi(m)>k}^n (\phi(m)-k)}$$
$$= ||A_k||^n \delta_j^{\sum_{m=1,\phi(m)>k}^n (\phi(m)-k)},$$

when $\sum_{m=1}^{n} \phi(m) = nk$. When in addition $\phi \not\equiv k$, there exists $m = 1, \ldots, n$ such that $\phi(m) > k$, thus $\sum_{m=1, \phi(m) > k}^{n} (\phi(m) - k) \geqslant 1$, which yields

$$||A_{\phi(1)}|| \dots ||A_{\phi(n)}|| \le ||A_k||^n \delta_j$$
.

From all the above inequalities, we deduce that

$$|\tilde{p}_{nk}| \geqslant |\det A_k| - \sum_{\phi \in \Phi_k, \ \phi \neq k} \eta ||A_k||^n \delta_j$$
.

Since by definition $C_{n,d,nk}$ is the cardinality of Φ_k , and there exists exactly one element of Φ_k such that $\phi \equiv k$, we obtain the inequality of the lemma for $k = k_j$. \square

Proof of Theorem 6.1. Consider the polynomial \tilde{p} of (47) and let $\gamma_1, \ldots, \gamma_{nd}$ be the tropical roots of the tropical polynomial $t \, \tilde{p}(x) = \max_{0 \leq l \leq nd} |\tilde{p}_l| x^l$ arranged in non-decreasing order. Let $j = 0, \ldots, q$ and denote $k = k_j$. From (2), we have

$$(48) |\zeta_1 \dots \zeta_{nk}| \leqslant \binom{nd}{nk} \gamma_1 \dots \gamma_{nk} ,$$

By the first part of Proposition 3.4 applied to the tropical polynomial $t \tilde{p}$, we get $|\tilde{p}_{nk}|\gamma_1 \dots \gamma_{nk} \leq |\tilde{p}_0|$. Applying the result of Lemma 6.5, and using the assumption of Theorem 6.1 on c_j , we get that $\tilde{p}_{nk} \geq c_j ||A_k||^n > 0$. Since $\tilde{p}_0 = |\det A_0|$, we deduce that $\gamma_1 \dots \gamma_{nk} \leq |\det A_0|/(c_j ||A_k||^n)$. Gathering this with (48), we get

$$|\zeta_1 \dots \zeta_{nk}| \leqslant \frac{\binom{nd}{nk}}{c_j} \frac{|\det A_0|}{\|A_0\|^n} \frac{\|A_0\|^n}{\|A_k\|^n}.$$

Since $k=k_j$ is the abscissa of a vertex of the Newton polygon of the tropical polynomial tp, we get from Proposition 3.4 applied to tp that $\frac{\|A_0\|}{\|A_k\|} = \alpha_1 \dots \alpha_k$, hence the previous inequality shows Theorem 6.1.

7. Numerical examples illustrating the lower bounds

In this section, we illustrate the bounds by numerical examples. We shall see in particular that the lower bounds stated in Section 4 can be tight when the tropical roots are well separated and when the input matrices are well conditioned, according to Remark 6.3. All the numerical results of the present paper were obtained using Scilab 4.1.2. Note that similar results are obtained using Matlab, specially the polyeig function.

7.1. Illustration of the lower bounds for the roots of scalar polynomials.

In the following examples, for each scalar polynomial, using the notations of Section 4, we shall show in tables or plots the following quantities with respect to $k=1,\ldots,d$ for comparison: - the ratio $\frac{|\zeta_1...\zeta_k|}{\alpha_1...\alpha_k}$, denoted RATIO; - the lower bound of Pólya given in (3), which coincides with the constant L_k of (27), denoted Pólya; - the lower bound L_k of (24) in Theorem 4.1, denoted FEWNOM; - the lower bound $L_k^{*,\text{prox}}$ of (40) in Corollary 4.9, based on the separation between the tropical roots, denoted SEPAR. Also, in tables, the abscissæ of the vertices of the Newton polygon of tp are indicated by the symbol *.

Example 7.1. Consider the following scalar polynomial

(49)
$$p(x) = 1 - \exp(1)x - \exp(6)x^2 + \exp(4)x^4 + \exp(9)x^8 + \exp(5)x^{10} + \exp(1)x^{16}$$
.

The log-exp transformation of its tropical polynomial tp is the tropical polynomial f of Example 3.3. The graph of f and the associated Newton polygon were shown in Figure 1. Hence, the tropical roots of tp are the exponentials of -3, -1/2, and 1, with multiplicities 2, 6 and 8, respectively. Table 1 shows the ratios and lower bounds, as explained at the beginning of the section.

k	RATIO	FEWNOM	PÓLYA	SEPAR
1	0.93475	0.37796	0.5	0.5
2*	1.00034	"	0.3849	0.92102
3	0.98782	"	0.32476	0.49664
4	0.97926	"	0.28622	0.38360
5	0.98348	"	0.25880	0.32403
6	0.98771	"	0.23802	0.37508
7	0.99383	"	0.22156	0.47570
8*	1.00000	"	0.20810	0.79696
9	0.98811	"	0.19683	0.47570
10	0.97636	"	0.18721	0.37508
11	0.96476	"	0.17888	0.31917
12	0.95329	"	0.17158	0.28622
13	0.96476	"	0.16510	0.32476
14	0.97636	"	0.15930	0.3849
15	0.98811	"	0.15407	0.5
16*	1	"	0.14933	1

Table 1. Comparison of the lower bounds for the scalar polynomial p of (49).

Example 7.2. Consider now the following scalar polynomial

$$p(x) = 1 - \exp(3)x - \exp(6)x^2 - \exp(7)x^4 - \exp(8)x^6 + \exp(9)x^8$$

$$(50) \qquad + \exp(7)x^{10} - \exp(4)x^{13} - \exp(3)x^{14} + \exp(1)x^{16} .$$

The log-exp transformation of its tropical polynomial $t\,p$ is again the tropical polynomial f of Example 3.3, but now there are some points of the graph of the map

 $k \mapsto \log |p_k|$, where p_k is the kth coefficient of p, that are on the edges of its Newton polygon, that is its concave hull. Table 2 shows the ratios and lower bounds for p. In Figure 3, we plot in the same graph the values of the ratios and lower bounds

k	RATIO	FEWNOM	PÓLYA	SEPAR
1	0.61759	0.31623	0.5	0.5
2*	0.98684	"	0.3849	0.92102
3	0.84948	"	0.32476	0.49664
4	0.73124	"	0.28622	0.38360
5	0.63280	"	0.25880	0.32403
6	0.54760	"	0.23802	0.37508
7	0.72118	"	0.22156	0.47570
8*	0.95684	"	0.20810	0.79696
9	0.76959	"	0.19683	0.47570
10	0.61899	"	0.18721	0.37508
11	0.57724	"	0.17888	0.31917
12	0.53832	"	0.17158	0.28622
13	0.62117	"	0.16510	0.32476
14	0.71677	"	0.15930	0.3849
15	0.84662	"	0.15407	0.5
16*	1	"	0.14933	1

Table 2. Comparison of the lower bounds for the scalar polynomial p of (50).

for the scalar polynomials p of (49) and (50). This indicates that the lower bound L_k of Corollary 4.9, based on the separation between tropical roots, may become tighter when the polynomial has non zero coefficients between the vertices of the Newton polygon.




FIGURE 3. Plot of RATIO as a function of k for the scalar polynomials of (49) (blue circles) and (50) (red squares), together with the lower bound FEWNOM (blue and red lines resp.), PÓLYA (plus signs) and SEPAR (stars signs).

7.2. Illustration of lower bounds for the eigenvalues of matrix polynomials. Note that to compute the eigenvalues of the following examples we had to use the tropical scaling algorithm that we recall in the next section.

In the following examples, we are comparing the value of the ratio $\frac{|\zeta_1...\zeta_{nk}|}{(\alpha_1...\alpha_k)^n}$, k = 1, ..., d, with the lower bounds (21) or (22) of Section 4 for some matrix polynomials P. In view of Remark 5.2, we computed only the lower bounds obtained under (A2) for the normalized Frobenius norm $\|\cdot\|_{nF}$, and the lower bounds obtained under (A1) for the normalized Schatten 1-norm (or normalized trace norm) $\|\cdot\|_{*1}$ (which satisfies (A1) but not (A2), see Property 2.5).

Then, using the notations of Section 4, we shall show in tables or plots the following quantities with respect to $k=1,\ldots,d$ for comparison: - the ratio $\frac{|\zeta_1\ldots\zeta_{nk}|}{(\alpha_1\ldots\alpha_k)^n}$, denoted RATIO; - the universal lower bound generalizing the lower bound of Pólya to the matrix case, given by cL_k^n (see (22)) with L_k of (26) or (27), denoted G-PÓLYA; - the lower bound involving the number of monomials, given by cL_k^n with L_k of (23) or (24), denoted FEWNOM; - the lower bound based on the separation between the tropical roots, given by $c(L_k^{\text{prox}})^n$ or $c(L_k^{*,\text{prox}})^n$ with L_k^{prox} and $L_k^{*,\text{prox}}$ of (39) and (40) respectively, denoted SEPAR. Also, in tables, the abscissæ of the vertices of the Newton polygon of t p are indicated by the symbol *.

Example 7.3. Consider the following matrix polynomial

$$P_1(\lambda) = 10^{-7}U_0 + 10^2\lambda^2U_2 + 10^7\lambda^4U_4 + 10\lambda^7U_7 + 10^{-8}\lambda^9U_9$$

where all the matrices U_j , $j \in \{0, 2, 4, 7, 9\}$, are unitary of dimension 3. Here, we shall only consider the normalized Frobenius norm since for a unitary matrix U, we have $||U||_{nF} = ||U||_{*1} = 1$, so the tropical polynomial associated with P_1 are the same for both norms, and the lower bounds under (A1) are the same, and are thus weaker than the lower bounds under (A2) for the normalized Frobenius norm. The Newton polygon of the tropical polynomial corresponding to P_1 (for the normalized Frobenius norm) is shown in Figure 4, and its tropical roots are $10^{-9/2}$, $10^{-5/2}$, 10^2 ,




FIGURE 4. The Newton polygon of the tropical polynomial corresponding to P_1 .

and $10^{9/2}$ with multiplicities 2, 2, 3, and 2, respectively. Figure 5 shows the lower bounds for all values of $1 \le k \le 9$, and the maximum of the ratios for a sample of 1000 random choices of the unitary matrices U_j . The ratios slightly change for different random choices of the unitary matrices U_j , but with a difference smaller than $6 \, 10^{-4}$. Note that when $1 \le k \le 9$ increases, the generalized lower bound of Pólya decreases, and should be replaced by its symmetrized version with k replaced by d - k. However the bounds using the separation between tropical roots are the

best ones, and they are tight at the vertices of the Newton polygon of the tropical polynomial associated with P_1 .

_			, , , , , , , , , ,	
k	RATIO	FEWNOM	G-PÓLYA	SEPAR
1	1.00015	0.08944	0.125	0.125
2*	1.00029	"	0.05702	0.97044
3	1.00015	"	0.03425	0.125
4*	1	"	0.02345	0.99991
5	1.00000	"	0.01733	0.125
6	1.00000	"	0.01348	0.12500
7*	1	"	0.01088	0.99056
8	1	"	0.00901	0.125
9*	1	"	0.00763	1




FIGURE 5. Comparison of the bounds for P_1 using the normalized Frobenius norm.

Example 7.4. In the following example, we perturb the previous polynomial by adding coefficients to P_1 , in such a way that the Newton polygon of the tropical polynomial associated with P_1 does not change, either for the normalized Frobenius norm or at least for the normalized Schatten 1-norm. Here we fix the unitary matrices to be either the identity matrix or its opposite, and we consider the following matrices:

$$B = b \begin{pmatrix} 10^{-15} & 1 & 1 \\ 0 & 1 & 1 \\ 1 & 2 & 3 \end{pmatrix}, \quad C = \frac{\sqrt{3}}{14} \begin{pmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 3 & 6 & 9 \end{pmatrix}$$

where b is chosen so that $||B||_{nF} = 1$, and C is also such that $||C||_{nF} = 1$. The matrix C is singular, since it has rank 1, and it satisfies $||C||_{*1} = 1/\sqrt{3}$. The matrix B is ill-conditioned, so that $||B||_{*1} < ||B||_{nF} = 1$ too.

We shall consider the two polynomials:

$$\begin{split} P_2(\lambda) = & 10^{-7} U_0 + 10^{-5/2} \lambda C_1 + 10^2 \lambda^2 U_2 + 10^{9/2} \lambda^3 C_1 + 10^7 \lambda^4 U_4 + 10^5 \lambda^5 C_1 \\ & + 10^3 \lambda^6 B_1 + 10 \lambda^7 U_7 + 10^{-7/2} \lambda^8 B_1 + 10^{-8} \lambda^9 U_9 \\ P_3(\lambda) = & 10^{-7} U_0 + 10^{-5/2} \lambda C_2 + 10^2 \lambda^2 U_2 + 10^{9/2} \lambda^3 C_2 + 10^7 \lambda^4 U_4 + 10^5 \lambda^5 C_2 \\ & + 10^3 \lambda^6 B_2 + 10 \lambda^7 U_7 + 10^{-7/2} \lambda^8 B_2 + 10^{-8} \lambda^9 U_9 \ , \end{split}$$

with $U_0 = U_4 = U_7 = -I$, $U_2 = U_9 = I$, $B_1 = B$, $C_1 = C$, $B_2 = \|B\|_{*1}^{-1}B$, $C_2 = \sqrt{3}C$. Since for any unitary matrix U with dimension 3, $\|U\|_{*1} = \|U\|_{\rm nF} = 1$, and since for any complex matrix A of dimension 3, we have $\|A\|_{*1} \leq \|A\|_{\rm nF}$, the tropical polynomial associated with P_2 for either the normalized Frobenius norm or the Schatten 1-norm coincides with the one associated with P_1 . Hence, as for P_1 , the bounds for P_2 with the normalized Schatten 1-norm are necessarily weaker than the ones with the normalized Frobenius norm. Moreover, all the bounds presented in Table 5 for P_1 remain the same for P_2 except the bounds based on the number of nonzero coefficients. We present all these bounds in Figure 6 together with the new ratios. Since these ratios are different from the ones of Table 5, one can see that the lower bounds based on the separation between tropical roots may be tight.

k	RATIO	FEWNOM	G-PÓLYA	SEPAR
1	0.45628	0.03162	0.125	0.125
2*	1.03298	"	0.05702	0.97044
3	0.45627	"	0.03425	0.125
4*	1.00009	"	0.02345	0.99991
5	0.39258	"	0.01733	0.125
6	0.39258	"	0.01348	0.12500
7*	1.00867	"	0.01088	0.99056
8	0.47428	"	0.00901	0.125
9*	1	"	0.00763	1




FIGURE 6. Comparison of the bounds for P_2 using the normalized Frobenius norm.

Let us consider now the polynomial P_3 . The normalized Schatten 1-norm of the coefficients of P_3 coincide with the normalized Frobenius norm of the coefficients of P_2 , hence the tropical polynomial associated with P_3 for the normalized Schatten 1-norm coincides with the one associated with P_1 for the normalized Frobenius norm, so that the tropical roots are the same. However, since the Schatten 1-norm satisfies (A1) but not (A2), we can only get the bounds based on (A1) which are necessarily lower than the one presented above for P_1 with the normalized Frobenius norm. Finally, the tropical polynomial associated with P_3 for the normalized Frobenius norm differs from the one associated with P_1 , so that the tropical roots and the bounds differ too. All tropical roots of this new tropical polynomial have multiplicity 1, which means that all indices are abscissæ of vertices of its Newton polygon. We present in Figure 7, the ratios and the lower bounds for both normalized Frobenius and Schatten 1-norms. Here the ratios and lower bounds for the normalized Schatten 1-norm are lower than the ones for the normalized Frobenius norm. However, one can see that the results for the normalized Frobenius norm are still better than the ones for the normalized Schatten 1-norm.


Example 7.5. In order to obtain a polynomial for which the lower bounds for the normalized Schatten 1-norm are better than the ones for the normalized Frobenius norm, one need to take matrices A_i such that $||A_i||_{\mathrm{nF}}/||A_i||_{*1}$ is as large as possible. This means that the dimension n is large and that some of the singular values of A_i are zero, or at least that the ratio between the maximal and minimal singular values is large, which implies that A_i is singular or at least has a large condition number. Moreover some parameters of the lower bounds need to be smaller than $||A_i||_{\mathrm{nF}}/||A_i||_{*1}$, which is possible for instance for the lower bounds based on the separation between tropical roots, when these tropical roots have a small multiplicity and are well separated. We shall show here such examples.

We consider a matrix polynomial with degree 3 and dimension n = 10:

$$P_4(\lambda) = A_0 + A_1 \lambda + A_2 \lambda^2 + A_3 \lambda^3$$

where all the matrices A_i , i = 0, ..., 3 are diagonal matrices: $A_i = \text{diag}(d_i)$ with $d_i \in \mathbb{C}^n$. We choose $A_0 = A_3 = I$, so that c = 1, $\alpha_1 \alpha_2 \alpha_3 = 1$ (for norms such that ||I|| = 1) and for k = 3 the ratio $\frac{|\zeta_1...\zeta_{nk}|}{(\alpha_1...\alpha_k)^n}$ is equal to 1, which is equal to the lower bound based on separation of tropical roots. We choose the diagonal elements of

	1	norm. Frob	enius norm	ı	norm. Schatten 1-norm				
k	RATIO	FEWNOM	G-PÓLYA	SEPAR	RATIO	FEWNOM	G-PÓLYA	SEPAR	
1	1.57223	0.03162	0.125	0.35355	0.30258	0.001	0.01563	0.01563	
2*	1.11466	"	0.05702	0.91391	1.11466	"	0.00325	0.54771	
3	1.57223	"	0.03425	0.35355	0.30258	"	0.00117	0.01562	
4*	1.00028	"	0.02345	0.99972	1.00028	"	0.00055	0.96682	
5	1.45041	"	0.01733	0.18345	0.27913	"	0.00030	0.01562	
6	1.01082	"	0.01348	0.05989	0.27913	"	0.00018	0.01533	
7*	1.02104	"	0.01088	0.97788	1.02104	"	0.00012	0.71337	
8	1.26686	"	0.00901	0.25721	0.34983	"	0.00008	0.01563	
9*	1	"	0.00763	1	1	"	0.00006	1	






FIGURE 7. Comparison of the bounds for P_3 using the normalized Frobenius (on left) and Schatten 1-norms (on right). The symbol * corresponds to the abscissæ of the vertices of the Newton polygon of the tropical polynomial associated with the normalized Frobenius norm.

the matrices A_1, A_2 as follows:

$$d_1 := \frac{a_1}{1 + b/n} (1, \dots, 1, 1 + b) , \quad d_2 := \frac{a_2}{1 + b/n} (1 + b, 1, \dots, 1) ,$$

with $a_1, a_2, b > 0$. Then

$$\begin{split} \|A_1\|_{*1} &= \frac{\|d_1\|_1}{n} = a_1 \ , \quad \|A_2\|_{*1} = \frac{\|d_2\|_1}{n} = a_2 \ , \\ \|A_1\|_{\mathrm{nF}} &= \frac{\|d_1\|_2}{\sqrt{n}} = a_1 f_b \ , \quad \|A_2\|_{\mathrm{nF}} = \frac{\|d_2\|_2}{\sqrt{n}} = a_2 f_b \ , \end{split}$$

with

$$f_b := \frac{\sqrt{1 + \frac{2b + b^2}{n}}}{1 + b/n} > 1$$
.

The constant f_b is large when b is large, that is when the condition numbers of A_1 and A_2 are large.

When $a_1 = a_2 > 1$, the tropical polynomial associated with P_4 for the normalized Schatten 1-norm (resp. the normalized Frobenius norm) has 3 different tropical roots equal to $1/a_1$, 1 and a_1 (resp. $1/(a_1f_b)$, 1 and a_1f_b). We present in Tables 3 and 4, the ratios and the lower bounds for both normalized Frobenius and Schatten 1-norms, when $a_1 = 10^4$, and b = 2 and b = 10 respectively. When b = 10, the matrices have a large condition number, so that the ratios $\frac{|\zeta_1...\zeta_{nk}|}{(\alpha_1...\alpha_k)^n}$ for k = 1 or 2 are very large. When either b = 2 or b = 10, we see that the lower bounds based on the separation between the tropical roots are the best ones, and that they are

nearer the ratios $(\frac{|\zeta_1...\zeta_{nk}|}{(\alpha_1...\alpha_k)^n})$ in the case of the Schatten 1-norm than in the case of the Frobenius norm, although they are far in the case where b=10.

	norm. Frobenius norm				r	orm. Schat	ten 1-norm	ı
k	RATIO	FEWNOM	G-PÓLYA	SEPAR	RATIO	FEWNOM	G-PÓLYA	SEPAR
1*	6.30697	0.00098	0.00098	0.99911	2.06667	9.5D-07	9.5D-07	0.81873
2*	6.30697	0.00098	0.00007	0.99911	2.06667	9.5D-07	5.1D-09	0.81873
3*	1	0.00098	0.00001	1	1	9.5D-07	1.7D-10	1

TABLE 3. Comparison of the bounds for P_4 with $a_1 = a_2 = 10^4$ and b = 2.

		norm. Frob	enius norm		norm. Schatten 1-norm			
k	RATIO	FEWNOM	G-PÓLYA	SEPAR	RATIO	FEWNOM	G-PÓLYA	SEPAR
1*	33882.7	0.00098	0.00098	0.99945	93.4462	9.5D-07	9.5D-07	0.81873
2*	33882.7	0.00098	0.00007	0.99945	93.4462	9.5D-07	5.1D-09	0.81873
3*	1	0.00098	0.00001	1	1	9.5D-07	1.7D-10	1

Table 4. Comparison of the bounds for P_4 with $a_1 = a_2 = 10^4$ and b = 10.

	norm. Frobenius norm				n	orm. Schat	ten 1-norm	
k	RATIO	FEWNOM	G-PÓLYA	SEPAR	RATIO	FEWNOM	G-PÓLYA	SEPAR
1*	1.1D+21	0.00098	0.00098	0.99968	1.2D+16	9.5D-07	9.5D-07	0.81873
2*	1.1D+21	0.00098	0.00007	0.99968	1.2D+16	9.5D-07	5.1D-09	0.81873
3*	1	0.00098	0.00001	1	1	9.5D-07	1.7D-10	1

Table 5. Comparison of the bounds for P_4 with $a_1 = a_2 = 10^4$ and $b = 10^3$.

When $a_1 > 1$ and $a_2 = \sqrt{a_1/f_b}$ the tropical polynomial associated with P_4 for the normalized Schatten 1-norm (resp. the normalized Frobenius norm) has only 2 different tropical roots equal to $1/a_1$ and $\sqrt{a_1}$ (resp. $1/(a_1f_b)$, 1 and $\sqrt{a_1f_b}$) with respective multiplicities 1 and 2. We present in Tables 6 and 7, the ratios and the lower bounds for both normalized Frobenius and Schatten 1-norms, when $a_1 = 10^4$, and b = 2 and b = 10 respectively. When b = 10, the matrices have a large condition number, so that for k = 1 or 2 the distances (ratios) between the ratio $\frac{|\zeta_1...\zeta_{nk}|}{(\alpha_1...\alpha_k)^n}$ and its lower bounds are very large, although for k = 2 this ratio is smaller than in the case where $a_1 = a_2 = 10^4$ above. When either b = 2 or b = 10, we see that the lower bounds based on the separation between the tropical roots are the best ones. For k = 2 which is the abscissa of a vertex of the Newton polygon, the lower bound obtained in the case of the Schatten 1-norm is nearer the ratio $\frac{|\zeta_1...\zeta_{nk}|}{(\alpha_1...\alpha_k)^n}$ than the one obtained in the case of the Frobenius norm, whereas for k = 3 the contrary holds.

8. The tropical scaling and its relation with the bounds

In the examples of the previous section, we considered matrix polynomials the coefficients of which have different order of magnitude. Hence, their eigenvalues need to be computed by using some scaling, otherwise the relative error of their approximation may become very large. We used here the "tropical scaling" introduced in [17], that we next recall. We shall see that in most of the examples of

		norm. Frob	enius norm		norm. Schatten 1-norm			
k	RATIO	FEWNOM	G-PÓLYA	SEPAR	RATIO	FEWNOM	G-PÓLYA	SEPAR
1*	6.29864	0.00098	0.00098	0.99999	2.06394	9.5D-07	9.5D-07	0.98020
2	0.68418	0.00098	0.00007	0.00098	0.39165	9.5D-07	5.1D-09	9.5D-07
3*	1	0.00098	0.00001	1	1	9.5D-07	1.7D-10	1

TABLE 6. Comparison of the bounds for P_4 with $a_1 = 10^4$, $a_2 = \sqrt{a_1/f_b}$, and b = 2.

	norm. Frobenius norm					norm. Schat	ten 1-norn	1
k	RATIO	FEWNOM	G-PÓLYA	SEPAR	RATIO	FEWNOM	G-PÓLYA	SEPAR
1*	33754.9	0.00098	0.00098	1.00000	93.0935	9.5D-07	9.5D-07	0.98020
2	9.86557	0.00098	0.00007	0.00098	0.5181	9.5D-07	5.1D-09	9.5D-07
3*	1	0.00098	0.00001	1	1	9.5D-07	1.7D-10	1

TABLE 7. Comparison of the bounds for P_4 with $a_1 = 10^4$, $a_2 = \sqrt{a_1/f_b}$, and b = 10.

Section 7, it improves the relative error, by comparison with a usual method, and that its efficiency is related with the closeness of the ratios $\frac{|\zeta_1...\zeta_{nk}|}{(\alpha_1...\alpha_k)^n}$ to 1.

8.1. **Tropical scaling.** Consider the matrix polynomial $P(\lambda) = \sum_{j=0}^{d} A_j \lambda^j$ with degree d (as in (5)). Denote $\operatorname{t} p(x) = \max_{0 \leq j \leq d} \|A_j\| x^j$ the tropical polynomial associated to P (as in (6)) and $\alpha_1 \leq \ldots \leq \alpha_d$ the tropical roots of $\operatorname{t} p$. Let also $k_0 = 0, k_1, \ldots, k_q = d$ be the sequence of abscissæ of the vertices of the Newton polygon of $\operatorname{t} p(x)$, as shown in Figure 2.

It has been observed numerically in [17], and proved in the special case of quadratic pencils, that when the coefficient matrices are well conditioned, the tropical roots are good approximations of the moduli of the eigenvalues. Corollary 6.2 and Remark 6.3 prove the same result for general matrix polynomials of any degree. So we expect to get $(k_i - k_{i-1})n$ eigenvalues of P with a modulus of order α_{k_i} , for $i = 1, \ldots, q$.

Given the order of magnitude α of an eigenvalue ζ of P, one can apply the following scaling with respect to α : $\lambda = \alpha \mu$ and

(51)
$$\widetilde{P}(\mu) := (\operatorname{t} p(\alpha))^{-1} P(\alpha \mu) = \sum_{j=0}^{d} \widetilde{A}_{j} \mu^{j} .$$

The eigenvalue ζ of P corresponds to the eigenvalue $\xi = \alpha^{-1}\zeta$ of \widetilde{P} . If ζ is of order of magnitude α , we get that ξ is of order 1, so that we expect that the numerical computation of ξ will behaves well, and better than the numerical computation of ζ .

The tropical scaling algorithm, already proposed in [17], consists in doing a different scaling for every expected order of magnitude of the eigenvalues as follows. Let us call C&QZ, the algorithm consisting in computing the eigenvalues (and eigenvectors) of the matrix polynomial P by applying QZ algorithm after a companion linearization L of P. For every tropical root $\alpha = \alpha_{k_i}$, $i = 1, \ldots, q$, we compute the eigenvalues of the scaled matrix polynomial \tilde{P} in (51) using C&QZ algorithm, and we multiply them by α_{k_i} . We get nd candidate eigenvalues of P. Then, we keep among these candidate eigenvalues only the $n(k_i - k_{i-1})$ ones having the closest modulus to α_{k_i} , and we say that these candidate eigenvalues are assigned to α_{k_i} .

The other eigenvalues returned by the C&QZ algorithm are discarded. Note that we measure the distance between two moduli by the absolute value of the difference of the log of these moduli. Finally, the whole spectrum is obtained as the union of the different groups of $n(k_i - k_{i-1})$ candidate eigenvalues assigned to the different tropical roots, for $1 \le i \le q$. If the moduli of the values of some groups interleave, meaning that for i < j, a candidate eigenvalue λ assigned to a tropical root α_{k_i} is such that $|\lambda| > |\lambda'|$ for some candidate eigenvalue λ' assigned to a tropical root α_{k_i} , the algorithm returns "fail" (the assignment is inconsistent).

8.2. Condition numbers and backward errors with or without tropical scalings. Let us now give some new insight of the interest of the tropical scaling, and its relation with the bounds of Sections 4 and 6.1.

Let us consider the condition numbers and backward errors of eigenvalues defined with respect to the relative errors (and left and right eigenvectors) as in [38, Theorems 4 and 5]. Recall that the accuracy of the computation of an eigenvalue can be evaluated by the product of its condition number and of its backward error. In our case, this accuracy can be computed either by using the condition number κ_P and the backward error η_P with respect to P or the condition number κ_L and the backward error η_L with respect to L.

Since the above scaling does not affect the condition number of the eigenvalue with respect to the matrix polynomial [20], and since QZ algorithm is backward stable, the accuracy of C&QZ on a given eigenvalue will be improved after scaling as soon as η_P/η_L decreases or κ_L/κ_P decreases (here η_P , κ_P , η_L , κ_L are respectively the backward error and the condition number of the eigenvalue with respect to the matrix polynomial and with respect to its linearization). The last property writes $\kappa_{\widetilde{L}}/\kappa_{\widetilde{P}} < \kappa_L/\kappa_P$, where $\kappa_{\widetilde{P}}$ is the condition number of the eigenvalue $\xi = \alpha^{-1}\zeta$ with respect to \widetilde{P} , and $\kappa_{\widetilde{L}}$ is the same with respect to the linearization of \widetilde{L} of \widetilde{P} .

The results of [22] and [20] show that, for quadratic pencils, κ_L/κ_P is lower bounded and that under some conditions, it is also upper bounded when using the scaling of [14] and the tropical scaling respectively. This means that in that cases these scalings are almost minimizing the ratio κ_L/κ_P (among all scalings) hence are almost maximizing the efficiency of C&QZ. These results have been generalized and improved in [7] and [36], for the companion linearization and several linearizations of matrix polynomials respectively. In particular, using the property that the scaled polynomial \widetilde{P} of (51) satisfies $\max_{0 \le j \le d} \|\widetilde{A}_j\| = 1$, it is shown that the following holds (for the companion linearization):

$$\frac{1}{\sqrt{d}} \leqslant \kappa_L/\kappa_P \ , \quad \kappa_{\widetilde{L}}/\kappa_{\widetilde{P}} \leqslant C_d \max(|\xi|, 1/|\xi|)^d \ ,$$

where C_d is a constant depending only on d. From these two inequalities, we see that the above scaling is almost maximizing the efficiency of C&QZ as soon as $\xi = \alpha^{-1}\zeta$ is close to 1 or equivalently α is a good order of magnitude of ζ . Hence the tropical scaling algorithm described in previous section leads to almost the best efficiency of C&QZ as soon as all the ratios $\zeta_{n(k-1)+l}/\alpha_k$ with $k=1,\ldots,d$ and $\ell=1,\ldots,n$ are close to 1. Several numerical examples are also given in [7] and [36].

8.3. Condition numbers and backward errors in examples. In the following we compare on the matrix polynomials of Section 7.2, the numerical accuracy of the tropical scaling algorithm to the one of the standard C&QZ algorithm. As said

before this accuracy can be measured either using κ_L or η_P , or the ratios κ_L/κ_P or η_P/η_L . Note that the condition numbers are computed using the result of the computation of the eigenvalues and eigenvectors, so only approximate condition numbers will be obtained. Actually, without scaling some of the eigenvalues will be aberrant. Moreover, the numerical values of κ_P and η_L may change after scaling, although they should not change theoretically.

In Figure 8, we plot the condition numbers and backward errors for the polynomial P_1 of Example 7.4. The number of eigenvalues is 27, but they moduli are close to the 4 tropical eigenvalues as shown in Figure 5. So, we see only 4 points in each graph corresponding to the computations using the tropical scaling. However, in the graphs corresponding to the computations without scaling, we see an additional point with approximate modulus 10^{-3} corresponding to aberrant values (there are no eigenvalues with this order of magnitude). We see also that the condition number of the linearization and the backward error of the matrix polynomial are much improved by using the tropical scaling, in particular for the smallest eigenvalues.

Similar conclusions are obtained for the polynomials P_2 and P_3 of Example 7.4.




FIGURE 8. Eigenvalues condition numbers and backward errors for P_1 of Example 7.3





FIGURE 9. Eigenvalues condition numbers and backward errors for P_2 of Example 7.4







FIGURE 10. Eigenvalues condition numbers and backward errors for P_3 of Example 7.4 without scaling or with scaling with respect to Schatten 1-norm

We finally consider the polynomial P_4 of Example 7.5 with $a_1 = a_2 = 10^4$. In this case, we can compute the eigenvalues using the diagonal structure of the matrices, for verification. We saw that the tightness of the lower bound decreases when b increases. The efficiency of the tropical scaling also decreases with b. Compare the results for b = 10 and b = 100 in Figures 11 and 12, respectively. When $b = 10^3$, the tropical scaling fails, leading to an error for the logarithm of the modulus greater than 1.99.







FIGURE 11. Eigenvalues condition numbers and backward errors for P_4 with b=10, without scaling or with scaling with respect to Frobenius norm

References

- M. Akian, R. Bapat, and S. Gaubert. Perturbation of eigenvalues of matrix pencils and optimal assignment problem. C. R. Acad. Sci. Paris, Série I, 339:103–108, 2004. Also arXiv:math.SP/0402438.
- [2] M. Akian, R. Bapat, and S. Gaubert. Min-plus methods in eigenvalue perturbation theory and generalised Lidskii-Vishik-Ljusternik theorem. arXiv:math.SP/0402090, 2005.
- [3] M. Akian, R. Bapat, and S. Gaubert. Max-plus algebras. In L. Hogben, editor, Handbook of Linear Algebra (Discrete Mathematics and Its Applications), volume 39. Chapman & Hall/CRC, 2006. Chapter 25.






FIGURE 12. Eigenvalues condition numbers and backward errors for P_4 with b = 100, without scaling or with scaling with respect to Frobenius norm

- [4] M. Akian, R. Bapat, and S. Gaubert. Non-archimedean valuations of eigenvalues of matrix polynomials. *Linear Algebra Appl.*, 498:592–627, 2016.
- [5] M. Akian, R. B. Bapat, and S. Gaubert. Asymptotics of the Perron eigenvalue and eigenvector using max algebra. C. R. Acad. Sci. Paris, Sér. I 327:927–932, 1998.
- [6] M. Akian, S. Gaubert, and A. Marchesini. Tropical bounds for eigenvalues of matrices. *Linear Algebra and its Applications*, January 2014. See also arXiv:1309.7319.
- [7] M. Akian, S. Gaubert, A. Marchesini, and F. Tisseur. Tropical diagonal scaling of matrix pencils, 2015. In preparation.
- [8] F. Baccelli, G. Cohen, G.J. Olsder, and J.P. Quadrat. Synchronization and Linearity. Wiley, 1992.
- [9] D. A. Bini. Numerical computation of polynomial zeros by means of Aberth's method. Numerical Algorithms, 13:179–200, 1996.
- [10] D.A. Bini, V. Noferini, and M. Sharify. Locating the Eigenvalues of Matrix Polynomials. SIAM J. Matrix Anal. Appl., 34(4):1708–1727, 2013.
- [11] G. D. Birkhoff. An elementary double inequality for the roots of an algebraic equation having greatest absolute value. Bull. Amer. Math. Soc., 21(10):494–495, 1915.
- [12] R. A. Cuninghame-Green and P. F. J. Meijer. An algebra for piecewise-linear minimax problems. Discrete Appl. Math., 2(4):267–294, 1980.
- [13] M. Einsiedler, M. Kapranov, and D. Lind. Non-Archimedean amoebas and tropical varieties. J. Reine Angew. Math., 601:139–157, 2006.
- [14] H.-Y. Fan, W-W. Lin, and P. Van Dooren. Normwise scaling of second order polynomial matrices. SIAM J. Matrix Anal. Appl., 26(1):252–256, 2004.
- [15] S. Friedland. Limit eigenvalues of nonnegative matrices. Linear Algebra Appl., 74:173–178, 1986.
- [16] M. Fujiwara. Über die obere schranke des absoluten betrages der wurzeln einer algebraischen gleichung. Tohoku Math. J., 1(10):167–171, 1916.
- [17] S. Gaubert and M. Sharify. Tropical scaling of polynomial matrices. In Rafael Bru and Sergio Romero-Vivó, editors, Proceedings of the third Multidisciplinary International Symposium on Positive Systems: Theory and Applications (POSTA 09), volume 389 of LNCIS, pages 291–303, Valencia, Spain, 2009. Springer. See also arXiv:0905.0121.
- [18] R. L. Graham. An efficient algorithm for determining the convex hull of a finite planar set. Inf. Proc. Lett., 1(4):132–133, 1972.
- [19] J. Hadamard. Étude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann. Journal de Mathématiques Pures et Appliquées, 4ième série, 9:171– 216, 1893.
- [20] S. Hammarling, C. J. Munro, and F. Tisseur. An algorithm for the complete solution of quadratic eigenvalue problems. ACM Transactions on Mathematical Software (TOMS), 39(3), 2013.

- [21] N. J. Higham and F. Tisseur. Bounds for eigenvalues of matrix polynomials. *Linear Algebra Appl.*, 358:5–22, 2003. Special issue on accurate solution of eigenvalue problems (Hagen, 2000).
- [22] Nicholas J. Higham, D. Steven Mackey, and Françoise Tisseur. The conditioning of linearizations of matrix polynomials. SIAM J. Matrix Anal. Appl., 28(4):1005–1028, 2006.
- [23] I. Itenberg, G. Mikhalkin, and E. Shustin. Tropical algebraic geometry. Oberwolfach seminars. Birkhäuser, 2007.
- [24] J. Jensen. Sur un nouvel et important théorème de la théorie des fonctions. Acta Mathematica, 22:359–364, 1899.
- [25] E. Landau. Sur quelques théorèmes de m. petrovitch relatifs aux zéros des fonctions analytiques. Bulletin de la S. M. F., 33:251–261, 1905.
- [26] E. L. Lindelöf. Mémoire sur la théorie des fonctions entières de genre fini. Acta Societatis Scientiarum Fennicae, XXXI, 1902.
- [27] G. Malajovich and J. P. Zubelli. Tangent Graeffe iteration. Numer. Math., 89(4):749–782, 2001.
- [28] A. Melman. Generalization and variations of Pellet's theorem for matrix polynomials. *Linear Algebra Appl.*, 439(5):1550–1567, 2013.
- [29] V. Noferini, M. Sharify, and F. Tisseur. Tropical roots as approximations to eigenvalues of matrix polynomials. SIAM J. Matrix Anal. Appl., 36(1):138–157, 2015.
- [30] A. Ostrowski. Recherches sur la méthode de Graeffe et les zéros des polynomes et des séries de Laurent. Acta Math., 72:99–155, 1940.
- [31] A. Ostrowski. Recherches sur la méthode de Graeffe et les zéros des polynomes et des séries de Laurent. Chapitres III et IV. Acta Math., 72:157–257, 1940.
- [32] M. Passare and H. Rullgård. Amoebas, Monge-Ampère measures, and triangulations of the Newton polytope. Duke Math. J., 121(3):481–507, 2004.
- [33] K. Purbhoo. A Nullstellensatz for amoebas. Duke Math. J., 141(3):407-445, 2008.
- [34] J. Richter-Gebert, B. Sturmfels, and T. Theobald. First steps in tropical geometry. In *Idempotent mathematics and mathematical physics*, volume 377 of *Contemp. Math.*, pages 289–317. Amer. Math. Soc., Providence, RI, 2005.
- [35] M. Sharify. Scaling Algorithms and Tropical Methods in Numerical Matrix Analysis: Application to the Optimal Assignment Problem and to the Accurate Computation of Eigenvalues. PhD thesis, Ecole Polytechnique, France, 2011.
- [36] M. Sharify and F. Tisseur. Effect of tropical scaling on linearizations of matrix polynomials: backward error and conditioning. t-mims-eprint, inst-MIMS, inst-MIMS:adr, 2014. In preparation.
- [37] W. Specht. Zur theorie der algebraischen gleichungen. Jahr. DMV, 48:142–145, 1938.
- [38] F. Tisseur. Backward error and condition of polynomial eigenvalue problems. Linear Algebra Appl., 309:339–361, 2000.
- [39] O. Viro. Dequantization of real algebraic geometry on logarithmic paper. In European Congress of Mathematics, Vol. I (Barcelona, 2000), volume 201 of Progr. Math., pages 135–146. Birkhäuser, Basel, 2001.

MARIANNE AKIAN, INRIA AND CMAP, ÉCOLE POLYTECHNIQUE, UMR 7641, CNRS. ADDRESS: CMAP, ÉCOLE POLYTECHNIQUE, ROUTE DE SACLAY, 91128 PALAISEAU CEDEX, FRANCE. E-mail address: Marianne.Akian@inria.fr

STÉPHANE GAUBERT, INRIA AND CMAP, ÉCOLE POLYTECHNIQUE, UMR 7641, CNRS. ADDRESS: CMAP, ÉCOLE POLYTECHNIQUE, ROUTE DE SACLAY, 91128 PALAISEAU CEDEX, FRANCE. E-mail address: Stephane.Gaubert@inria.fr

MEISAM SHARIFY, SCHOOL OF MATHEMATICS, INSTITUTE FOR RESEARCH IN FUNDAMENTAL SCIENCES (IPM), P.O. BOX:19395-5746, TEHRAN, IRAN.

E-mail address: meisam.sharify@ipm.ir