

arXiv:1304.3731v1 [math.CV] 12 Apr 2013

A Note on the Hypercomplex Riemann-Cauchy Like Relations for Quaternions and Laplace Equations.

J. A. P.F. Maranhão¹ and M. F. Borges²

¹Department of Mathematics
Federal University of Maranhão - São Luís-MA
65085-580, Maranhão, BRAZIL
josemarao7@hotmail.com

²UNESP - São Paulo State University
S.J. Rio Preto Campus
15054-000, São José do Rio Preto, BRAZIL
borges@ibilce.unesp.br

June 20, 2018

Abstract

In this note it is worked out a new set of Laplace-Like equations for quaternions through Riemann-Cauchy hypercomplex relations obtained earlier [1]. As in the theory of functions of a complex variable, it is expected that this new set of Laplace-Like equations might be applied to a large number of Physical problems, providing new insights in the Classical Theory Fields.

AMS Subject Classification: 30G99, 30E99

Key Words and Phrases:Quaternions, Laplace's Equations, Riemann-Cauchy Relations

1 Cauchy-Riemann Equations (Functions of one complex variable).

In order to fix ideas will be considered theorem that relates the partial derivatives for the case of a function $f(z)$ of a complex variable $f(z) = u(x, y) + iv(x, y)$ [2], which here will be called simply Riemann-Cauchy conditions. These relations say that the first order partial derivatives of functions $u(x, y)$ and $v(x, y)$ satisfy relations according to the following theorem:

Theorem 1. *Is $f(z) = u(x, y) + iv(x, y)$ a function defined and continues in a neighborhood of the point $z = x + yi$ and differentiable at z . Then the partial derivatives of the first order of $u(x, y)$ and $v(x, y)$ exist and satisfy the relations:*

$$\frac{\partial u(x, y)}{\partial x} = \frac{\partial v(x, y)}{\partial y} \quad (1)$$

$$\frac{\partial u(x, y)}{\partial y} = -\frac{\partial v(x, y)}{\partial x} \quad (2)$$

Thus, if $f(z)$ is analytic in a domain Γ , its partial derivatives exist and satisfy the set of relations (1) and (2) over all point in Γ . Moreover, with the above functions class C^2 using Schwartz's Theorem for partial derivatives immediately follows the following equations:

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \quad (3)$$

$$\frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2} = 0 \quad (4)$$

the above equations are called Laplace's equations.

2 Cauchy-Riemann conditions for quaternionic functions.

Now the conditions are presented as the Riemann-Cauchy like relations for quaternionic functions. It follows the Theorem [1]:

Theorem 2. For any pair points a and b and any path joining them simply connect subdomain of the four-dimensional space, the integral $\int_a^b f dq$ is independent from the given path if and only if there is a function $F = F_1 + F_2i + F_3j + F_4k$ such that $\int_a^b f dq = F(a) - F(b)$, and satisfying the following relations:

$$\frac{\partial F_1}{\partial x_1} = \frac{\partial F_2}{\partial x_2} = \frac{\partial F_3}{\partial x_3} = \frac{\partial F_4}{\partial x_4} \quad (5)$$

$$\frac{\partial F_2}{\partial x_1} = -\frac{\partial F_1}{\partial x_2} = -\frac{\partial F_3}{\partial x_4} = \frac{\partial F_4}{\partial x_3} \quad (6)$$

$$\frac{\partial F_3}{\partial x_1} = -\frac{\partial F_1}{\partial x_3} = -\frac{\partial F_2}{\partial x_4} = \frac{\partial F_4}{\partial x_2} \quad (7)$$

$$\frac{\partial F_4}{\partial x_1} = \frac{\partial F_1}{\partial x_4} = -\frac{\partial F_2}{\partial x_3} = -\frac{\partial F_3}{\partial x_2} \quad (8)$$

Proof. The proof of this theorem can be analyzed in greater detail in [1]. \square

3 The Laplace's Equations.

In this section we show that a new set of hypercomplex Laplace equations may be generated in four dimensions. Through its use of Riemann-Cauchy like relations [1] Therefore, the functions that make up the quaternionic function, depend on x_1, x_2, x_3 and x_4 and are supposed to be of class C^2 and thus the theorem is valid Schwartz.

The first step to obtain the Laplace equation is the derivation of equations (5), (6), (7) and (8) over x_1, x_2, x_3 and x_4 will be done as follows: Deriving the conditions of equation (5), we have:

$$\begin{aligned} \frac{\partial^2 F_1}{\partial x_1^2} &= \frac{\partial^2 F_2}{\partial x_1 \partial x_2} = \frac{\partial^2 F_3}{\partial x_1 \partial x_3} = \frac{\partial^2 F_4}{\partial x_1 \partial x_4} \\ \frac{\partial^2 F_1}{\partial x_1 \partial x_2} &= \frac{\partial^2 F_2}{\partial x_2^2} = \frac{\partial^2 F_3}{\partial x_2 \partial x_3} = \frac{\partial^2 F_4}{\partial x_2 \partial x_4} \\ \frac{\partial^2 F_1}{\partial x_3 \partial x_1} &= \frac{\partial^2 F_2}{\partial x_3 \partial x_2} = \frac{\partial^2 F_3}{\partial x_3^2} = \frac{\partial^2 F_4}{\partial x_4 \partial x_3} \\ \frac{\partial^2 F_1}{\partial x_1 \partial x_4} &= \frac{\partial^2 F_2}{\partial x_4 \partial x_2} = \frac{\partial^2 F_3}{\partial x_4 \partial x_3} = \frac{\partial^2 F_4}{\partial x_4^2}. \end{aligned} \quad (9)$$

Deriving the conditions of equation (6), we have:

$$\begin{aligned}
\frac{\partial^2 F_2}{\partial x_1^2} &= -\frac{\partial^2 F_1}{\partial x_1 \partial x_2} = -\frac{\partial^2 F_3}{\partial x_1 \partial x_4} = \frac{\partial^2 F_4}{\partial x_1 \partial x_3} \\
\frac{\partial^2 F_2}{\partial x_1 \partial x_2} &= -\frac{\partial^2 F_1}{\partial x_2^2} = -\frac{\partial^2 F_3}{\partial x_2 \partial x_4} = \frac{\partial^2 F_4}{\partial x_3 \partial x_2} \\
\frac{\partial x_3 \partial x_1}{\partial^2 F_2} &= -\frac{\partial x_3 \partial x_2}{\partial^2 F_1} = -\frac{\partial x_3 \partial x_4}{\partial^2 F_3} = \frac{\partial x_3^2}{\partial^2 F_4} \\
\frac{\partial x_4 \partial x_1}{\partial^2 F_2} &= -\frac{\partial x_4 \partial x_2}{\partial^2 F_1} = -\frac{\partial x_4^2}{\partial^2 F_3} = \frac{\partial x_4 \partial x_3}{\partial^2 F_4}.
\end{aligned} \tag{10}$$

Deriving the conditions of equation (7), we obtain that:

$$\begin{aligned}
\frac{\partial^2 F_3}{\partial x_1^2} &= -\frac{\partial^2 F_1}{\partial x_1 \partial x_3} = -\frac{\partial^2 F_2}{\partial x_1 \partial x_4} = \frac{\partial^2 F_4}{\partial x_1 \partial x_2} \\
\frac{\partial x_1 \partial x_2}{\partial^2 F_3} &= -\frac{\partial x_2 \partial x_3}{\partial^2 F_1} = -\frac{\partial x_2 \partial x_4}{\partial^2 F_2} = \frac{\partial x_2^2}{\partial^2 F_4} \\
\frac{\partial x_3 \partial x_1}{\partial^2 F_3} &= -\frac{\partial x_3^2}{\partial^2 F_1} = -\frac{\partial x_4 \partial x_3}{\partial^2 F_2} = \frac{\partial x_3 \partial x_2}{\partial^2 F_4} \\
\frac{\partial x_1 \partial x_4}{\partial^2 F_3} &= -\frac{\partial x_4 \partial x_3}{\partial^2 F_1} = -\frac{\partial x_4^2}{\partial^2 F_2} = \frac{\partial x_4 \partial x_2}{\partial^2 F_4}.
\end{aligned} \tag{11}$$

Deriving the conditions of equation (8), we have:

$$\begin{aligned}
\frac{\partial^2 F_4}{\partial x_1^2} &= \frac{\partial^2 F_1}{\partial x_1 \partial x_4} = -\frac{\partial^2 F_2}{\partial x_1 \partial x_3} = -\frac{\partial^2 F_3}{\partial x_1 \partial x_2} \\
\frac{\partial x_1 \partial x_2}{\partial^2 F_4} &= \frac{\partial x_2 \partial x_4}{\partial^2 F_1} = -\frac{\partial x_2 \partial x_3}{\partial^2 F_2} = -\frac{\partial x_2^2}{\partial^2 F_3} \\
\frac{\partial x_3 \partial x_1}{\partial^2 F_4} &= \frac{\partial x_3 \partial x_4}{\partial^2 F_1} = -\frac{\partial x_3^2}{\partial^2 F_2} = -\frac{\partial x_3 \partial x_2}{\partial^2 F_3} \\
\frac{\partial x_1 \partial x_4}{\partial^2 F_4} &= \frac{\partial x_4^2}{\partial^2 F_1} = -\frac{\partial x_4 \partial x_3}{\partial^2 F_2} = -\frac{\partial x_4 \partial x_2}{\partial^2 F_3}.
\end{aligned} \tag{12}$$

In correlating those derived groups of partial derivatives (9), (10), (11) and (12), immediately following the Laplace Equations:

$$\frac{\partial F_1}{\partial x_1^2} + \frac{\partial F_1}{\partial x_2^2} + \frac{\partial F_1}{\partial x_3^2} + \frac{\partial F_1}{\partial x_4^2} = 0 \tag{13}$$

$$\frac{\partial F_2}{\partial x_1^2} + \frac{\partial F_2}{\partial x_2^2} + \frac{\partial F_2}{\partial x_3^2} + \frac{\partial F_2}{\partial x_4^2} = 0 \tag{14}$$

$$\frac{\partial F_3}{\partial x_1^2} + \frac{\partial F_3}{\partial x_2^2} + \frac{\partial F_3}{\partial x_3^2} + \frac{\partial F_3}{\partial x_4^2} = 0 \quad (15)$$

and

$$\frac{\partial F_4}{\partial x_1^2} + \frac{\partial F_4}{\partial x_2^2} + \frac{\partial F_4}{\partial x_3^2} + \frac{\partial F_4}{\partial x_4^2} = 0 \quad (16)$$

Therefore, in a more simplified manner, the set of equations appears below:

$$\begin{aligned} \Delta f_1 &= 0 \\ \Delta f_2 &= 0 \\ \Delta f_3 &= 0 \\ \Delta f_4 &= 0 \end{aligned} \quad (17)$$

4 Conclusion

In this note it is shown the feasibility of obtaining the equations of Laplace through the Cauchy-Riemann conditions for quaternions. This fact will allow the relationship between equations that can explain many physical phenomena. You can also use the above equations as a way of stating a theorem for harmonic functions for quaternions that satisfy the conditions of Cauchy results. It is worth mentioning the importance of [1] established relations used in this work.

References

- [1] Machado, J.M, Borges, M.F. New Remarks on the Differentiability of Hypercomplex Functions, In: *International Journal of Pure and Applied Mathematics*, v. 8 n. 1 (2002), 85-101.
- [2] Kodaira, Kunihiko; *Complex Analysis*, Cambridge Studies in Advanced Mathematics; **Cambridge University Press Cambridge**, (2007) 406pp. .