Localization formulas about two Killing vector fields

Xu Chen *

Abstract

In this article, we will discuss the smooth $(X_M + \sqrt{-1}Y_M)$ -invariant forms on M and to establish a localization formulas. As an application, we get a localization formulas for characteristic numbers.

The localization theorem for equivariant differential forms was obtained by Berline and Vergne(see [2]). They discuss on the zero points of a Killing vector field, the localization formula expresses the integral of an equivariantly closed differential form as an integral over the set of zeros of the Killing vector field. Now, We will discuss the equivariant cohomology about two Killing vector fields and to establish a localization formulas.

Let M be a smooth closed oriented manifold. Let G be a compact Lie group acting smoothly on M, and let \mathfrak{g} be its Lie algebra. Let g^{TM} be a G-invariant metric on TM. If $X, Y \in \mathfrak{g}$, let X_M, Y_M be the corresponding smooth vector field on M. If $X, Y \in \mathfrak{g}$, then X_M, Y_M are Killing vector field. Here we will introduce the equivariant cohomology about two Killing vector fields.

1 Equivariant cohomology by two Killing vector fields

First, let us review the definition of equivariant cohomology by a Killing vector field. Let $\Omega^*(M)$ be the space of smooth differential forms on M, the de Rham complex is $(\Omega^*(M), d)$. Let L_{X_M} be the Lie derivative of X_M on $\Omega^*(M)$, i_{X_M} be the interior multiplication induced by the contraction of X_M .

Set

$$d_X = d + i_{X_M}$$

then $d_X^2 = L_{X_M}$ by the following Cartan formula

$$L_{X_M} = [d, i_{X_M}]$$

Let

$$\Omega_X^*(M) = \{\omega \in \Omega^*(M) : L_{X_M}\omega = 0\}$$

be the space of smooth X_M -invariant forms on M. Then $d_X^2 \omega = 0$, when $\omega \in \Omega_X^*(M)$. It is a complex $(\Omega_X^*(M), d_X)$. The corresponding cohomology group

$$H_X^*(M) = \frac{\operatorname{Kerd}_X|_{\Omega^*_X(M)}}{\operatorname{Imd}_X|_{\Omega^*_X(M)}}$$

^{*}Email: xiaorenwu08@163.com. ChongQing, China.

is called the equivariant cohomology associated with X. If a form ω has $d_X \omega = 0$, then ω called d_X -closed form.

Then we will to definite a new complex by two Killing vector field. If $X, Y \in \mathfrak{g}$, let X_M, Y_M be the corresponding smooth vector field on M.

We know

$$L_{X_M} + \sqrt{-1}L_{Y_M}$$

be the operator on $\Omega^*(M) \otimes_{\mathbb{R}} \mathbb{C}$. Set

$$i_{X_M + \sqrt{-1}Y_M} \doteq i_{X_M} + \sqrt{-1}i_{Y_M}$$

be the interior multiplication induced by the contraction of $X_M + \sqrt{-1}Y_M$. It is also a operator on $\Omega^*(M) \otimes_{\mathbb{R}} \mathbb{C}$.

Set

$$d_{X+\sqrt{-1}Y} = d + i_{X_M+\sqrt{-1}Y_M}.$$

Lemma 1. If $X, Y \in \mathfrak{g}$, let X_M, Y_M be the corresponding smooth vector field on M; then

$$d_{X+\sqrt{-1}Y}^2 = L_{X_M} + \sqrt{-1}L_{Y_M}$$

Proof.

$$(d + i_{X_M + \sqrt{-1}Y_M})^2 = (d + i_{X_M} + \sqrt{-1}i_{Y_M})(d + i_{X_M} + \sqrt{-1}i_{Y_M})$$

= $d^2 + di_{X_M} + i_{X_M}d + \sqrt{-1}di_{Y_M} + \sqrt{-1}i_{Y_M}d + (i_{X_M} + \sqrt{-1}i_{Y_M})^2$
= $L_{X_M} + \sqrt{-1}L_{Y_M}$

Let

$$\Omega^*_{X_M + \sqrt{-1}Y_M}(M) = \{ \omega \in \Omega^*(M) \otimes_{\mathbb{R}} \mathbb{C} : (L_{X_M} + \sqrt{-1}L_{Y_M})\omega = 0 \}$$

be the space of smooth $(X_M + \sqrt{-1}Y_M)$ -invariant forms on M. Then we get a complex $(\Omega^*_{X_M + \sqrt{-1}Y_M}(M), d_{X + \sqrt{-1}Y})$. We call a form ω is $d_{X + \sqrt{-1}Y}$ -closed if $d_{X + \sqrt{-1}Y}\omega = 0$ (this is first discussed by Bimsut, see [3]). The corresponding cohomology group

$$H_{X+\sqrt{-1}Y}^{*}(M) = \frac{\operatorname{Ker} d_{X+\sqrt{-1}Y}|_{\Omega_{X+\sqrt{-1}Y}^{*}(M)}}{\operatorname{Im} d_{X+\sqrt{-1}Y}|_{\Omega_{X+\sqrt{-1}Y}^{*}(M)}}$$

is called the equivariant cohomology associated with K.

2 The set of zero points

Lemma 2. If $X, Y \in \mathfrak{g}$, let X_M, Y_M be the corresponding smooth vector field on M, X', Y' be the 1-form on M which is dual to X_M, Y_M by the metric g^{TM} , then

$$L_{X_M}Y' + L_{Y_M}X' = 0.$$

If $[X_M, Y_M] = 0$, then

$$L_{X_M}Y' = L_{Y_M}X' = 0.$$

Proof. Because

$$(L_{X_M}\omega)(Z) = X_M(\omega(Z)) - \omega([X_M, Z])$$

here $Z \in \Gamma(TM)$, So we get

$$(L_{X_M}Y')(Z) = X_M < Y_M, Z > - < [X_M, Z], Y_M >$$
$$(L_{Y_M}X')(Z) = Y_M < X_M, Z > - < [Y_M, Z], X_M >$$

Because X_M, Y_M are Killing vector fields, so (see [6])

$$X_M < Y_M, Z > = < L_{X_M} Y_M, Z > + < Y_M, L_{X_M} Z >$$

= < [X_M, Y_M], Z > + < Y_M, [X_M, Z] >

$$Y_M < X_M, Z > = < L_{Y_M} X_M, Z > + < X_M, L_{Y_M} Z >$$

= < [Y_M, X_M], Z > + < X_M, [Y_M, Z] >

then we get

$$(L_{X_M}Y' + L_{Y_M}X')(Z) = < [X_M, Y_M], Z > + < [Y_M, X_M], Z > = 0.$$

If $[X_M, Y_M] = 0$, we have

$$(L_{X_M}Y')(Z) = \langle [X_M, Y_M], Z \rangle + \langle Y_M, [X_M, Z] \rangle - \langle [X_M, Z], Y_M \rangle = 0.$$

$$(L_{Y_M}X')(Z) = \langle [Y_M, X_M], Z \rangle + \langle X_M, [Y_M, Z] \rangle - \langle [Y_M, Z], X_M \rangle = 0.$$

Lemma 3. If $X, Y \in \mathfrak{g}$, let X_M, Y_M be the corresponding smooth vector field on M, X', Y' be the 1-form on M which is dual to X_M, Y_M by the metric g^{TM} , then

$$d_{X+\sqrt{-1}Y}(X'+\sqrt{-1}Y')$$

is the $d_{X+\sqrt{-1}Y}$ -closed form.

Proof.

$$\begin{aligned} d_{X+\sqrt{-1}Y}^2(X'+\sqrt{-1}Y') &= d_{X+\sqrt{-1}Y}(d(X'+\sqrt{-1}Y')+i_{X_M+\sqrt{-1}Y_M}(X'+\sqrt{-1}Y')) \\ &= di_{X_M+\sqrt{-1}Y_M}(X'+\sqrt{-1}Y')+i_{X_M+\sqrt{-1}Y_M}d(X'+\sqrt{-1}Y') \\ &= L_{X_M}X'-L_{Y_M}Y'+\sqrt{-1}(L_{X_M}Y'+L_{Y_M}X') \\ &= 0 \end{aligned}$$

So $d_{X+\sqrt{-1}Y}(X'+\sqrt{-1}Y')$ is the $d_{X+\sqrt{-1}Y}$ -closed form.

Lemma 4. For any $\eta \in H^*_{X+\sqrt{-1}Y}(M)$ and $s \ge 0$, we have

$$\int_{M} \eta = \int_{M} \exp\{-s(d_{X+\sqrt{-1}Y}(X'+\sqrt{-1}Y'))\}\eta$$

Proof. Because

$$\frac{\partial}{\partial s} \int_{M} \exp\{-s(d_{X+\sqrt{-1}Y}(X'+\sqrt{-1}Y'))\}\eta \\ = -\int_{M} (d_{X+\sqrt{-1}Y}(X'+\sqrt{-1}Y')) \exp\{-s(d_{X+\sqrt{-1}Y}(X'+\sqrt{-1}Y'))\}\eta$$

and by assumption we have

$$d_{X+\sqrt{-1}Y}\eta = 0$$
$$d_{X+\sqrt{-1}Y}\exp\{-s(d_{X+\sqrt{-1}Y}(X'+\sqrt{-1}Y'))\} = 0$$

So we get

$$(d_{X+\sqrt{-1}Y}(X'+\sqrt{-1}Y')) \exp\{-s(d_{X+\sqrt{-1}Y}(X'+\sqrt{-1}Y'))\}\eta$$

= $d_{X+\sqrt{-1}Y}[(X'+\sqrt{-1}Y') \exp\{-s(d_{X+\sqrt{-1}Y}(X'+\sqrt{-1}Y'))\}\eta]$

and by Stokes formula we have

$$\frac{\partial}{\partial s} \int_{M} \exp\{-s(d_{X+\sqrt{-1}Y}(X^{'}+\sqrt{-1}Y^{'}))\}\eta = 0$$

Then we get

$$\int_{M} \eta = \int_{M} \exp\{-s(d_{X+\sqrt{-1}Y}(X'+\sqrt{-1}Y'))\}\eta$$

We have

$$d_{X+\sqrt{-1}Y}(X'+\sqrt{-1}Y') = d(X'+\sqrt{-1}Y') + \langle X_M + \sqrt{-1}Y_M, X_M + \sqrt{-1}Y_M \rangle$$

and

$$\langle X_M + \sqrt{-1}Y_M, X_M + \sqrt{-1}Y_M \rangle = |X_M|^2 - |Y_M|^2 + 2\sqrt{-1}\langle X_M, Y_M \rangle$$

 Set

$$M_0 = \{ x \in M \mid \langle X_M(x) + \sqrt{-1}Y_M(x), X_M(x) + \sqrt{-1}Y_M(x) \rangle = 0 \}.$$

For simplicity, we assume that M_0 is the connected submanifold of M, dim $M_0 < \dim M$, and \mathcal{N} is the normal bundle of M_0 about M. The set M_0 is first discussed by H.Jacobowitz (see [4]).

Lemma 5. For any $\eta \in H^*_{X+\sqrt{-1}Y}(M)$, the top form of η is exact outside M_0 . *Proof.* By lemma 3., we have on $M \setminus M_0$

$$\eta = d_{X+\sqrt{-1}Y} \left(\frac{(X' + \sqrt{-1}Y') \land \eta}{d_{X+\sqrt{-1}Y}(X' + \sqrt{-1}Y')} \right)$$

If $\eta_{[n]}$ is the top form of η , then we get

$$\eta_{[n]} = d \left(\frac{(X' + \sqrt{-1}Y') \land \eta}{d_{X + \sqrt{-1}Y}(X' + \sqrt{-1}Y')} \right)_{[n-1]}$$

on $M \setminus M_0$.

Corollery 1. For any $\eta \in H^*_{X+\sqrt{-1}Y}(M)$, if $M_0 = \emptyset$, then $\int_M \eta = 0$. *Proof.* By the Stokes formula and Lemma 5., we get the result.

Localization formula on $d_{X+\sqrt{-1}Y}$ -closed form 3

Set E is a G-equivariant vector bundle, if ∇^E is a connection on E which commutes with the action of G on $\Omega(M, E)$, we see that

$$[\nabla^E, L_X^E] = 0$$

for all $X \in \mathfrak{g}$. Then we can get a moment map by

$$\mu^E(X) = L_X^E - [\nabla^E, i_X] = L_X^E - \nabla^E_X$$

We known that if y be the tautological section of the bundle $\pi^* E$ over E, then the vertical component of X_E may be identified with $-\mu^E(X)y$ (see [1] proposition 7.6). For the normal bundle \mathcal{N} of M_0 , the vector field $X^{\mathcal{N}}$ and $Y^{\mathcal{N}}$ are vertical and are given at the point $(x, y) \in$ $M_0 \times \mathcal{N}_x$ by the vectors $-\mu^{\mathcal{N}}(X)y, -\mu^{\mathcal{N}}(Y)y \in \mathcal{N}_x$. If *E* is the tangent bundle *TM* and ∇^{TM} is Levi-Civita connection, then we have

$$\mu^{TM}(X)Y = L_XY - \nabla^{TM}_XY = -\nabla^{TM}_YX$$

We known that for any Killing vector field X, $\mu^{TM}(X)$ as linear endomorphisms of TM is skew-symmetric, $-\mu^{TM}(X)$ annihilates the tangent bundle TM_0 and induces a skewsymmetric automorphism of the normal bundle $\mathcal{N}(\text{see }[5] \text{ chapter II, proposition } 2.2 \text{ and}$ theorem 5.3). The restriction of $\mu^{TM}(X)$ to \mathcal{N} coincides with the moment endomorphism $\mu^{\mathcal{N}}(X).$

Now we construct a one-form α on \mathcal{N} :

$$Z \in \Gamma(T\mathcal{N}) \to \alpha(Z) = \langle -\mu^{\mathcal{N}}(X)y, \nabla_Z^{\mathcal{N}}y \rangle + \sqrt{-1} \langle -\mu^{\mathcal{N}}(Y)y, \nabla_Z^{\mathcal{N}}y \rangle$$

Let $Z_1, Z_2 \in \Gamma(T\mathcal{N})$, we known $d\alpha(Z_1, Z_2) = Z_1\alpha(Z_2) - Z_2\alpha(Z_1) - \alpha([Z_1, Z_2])$, so

$$d\alpha(Z_1, Z_2) = \langle -\nabla_{Z_1}^{\mathcal{N}} \mu^{\mathcal{N}}(X)y, \nabla_{Z_2}^{\mathcal{N}}y \rangle - \langle -\nabla_{Z_2}^{\mathcal{N}} \mu^{\mathcal{N}}(X)y, \nabla_{Z_1}^{\mathcal{N}}y \rangle + \sqrt{-1} \langle -\nabla_{Z_1}^{\mathcal{N}} \mu^{\mathcal{N}}(Y)y, \nabla_{Z_2}^{\mathcal{N}}y \rangle - \sqrt{-1} \langle -\nabla_{Z_2}^{\mathcal{N}} \mu^{\mathcal{N}}(Y)y, \nabla_{Z_1}^{\mathcal{N}}y \rangle + \langle -\mu^{\mathcal{N}}(X)y, R^{\mathcal{N}}(Z_1, Z_2)y \rangle + \sqrt{-1} \langle -\mu^{\mathcal{N}}(Y)y, R^{\mathcal{N}}(Z_1, Z_2)y \rangle$$

Recall that $\nabla^{\mathcal{N}}$ is invariant under L_X for all $X \in \mathfrak{g}$, so that $[\nabla^{\mathcal{N}}, \mu^{\mathcal{N}}(X)] = 0, [\nabla^{\mathcal{N}}, \mu^{\mathcal{N}}(Y)] =$ 0. And by X, Y are Killing vector field, we have $d\alpha$ equals

$$2\langle -(\mu^{\mathcal{N}}(X) + \sqrt{-1}\mu^{\mathcal{N}}(Y)) \cdot, \cdot \rangle + \langle -\mu^{\mathcal{N}}(X)y - \sqrt{-1}\mu^{\mathcal{N}}(Y)y, R^{\mathcal{N}}y \rangle$$

And by $|X_{\mathcal{N}}|^2 = \langle \mu^{\mathcal{N}}(X)y, \mu^{\mathcal{N}}(X)y \rangle, |Y_{\mathcal{N}}|^2 = \langle \mu^{\mathcal{N}}(Y)y, \mu^{\mathcal{N}}(Y)y \rangle$. So We can get

$$d_{X_{\mathcal{N}}+\sqrt{-1}Y_{\mathcal{N}}}(X'_{\mathcal{N}}+\sqrt{-1}Y'_{\mathcal{N}}) = -2\langle (\mu^{\mathcal{N}}(X)+\sqrt{-1}\mu^{\mathcal{N}}(Y))\cdot,\cdot\rangle + \langle -\mu^{\mathcal{N}}(X)y-\sqrt{-1}\mu^{\mathcal{N}}(Y)y, -\mu^{\mathcal{N}}(X)y-\sqrt{-1}\mu^{\mathcal{N}}(Y)y + R^{\mathcal{N}}y\rangle$$

Theorem 1. Let M be a smooth closed oriented manifold, G be a compact Lie group acting smoothly on M. For any $\eta \in H^*_{X+\sqrt{-1}Y}(M)$, $[X_M, Y_M] = 0$, the following identity hold:

$$\int_{M} \eta = \int_{M_0} \frac{\eta}{\Pr\left[\frac{-\mu^{\mathcal{N}}(\mathbf{X}) - \sqrt{-1}\mu^{\mathcal{N}}(\mathbf{Y}) + \mathbf{R}^{\mathcal{N}}}{2\pi}\right]}$$

Proof. Set $s = \frac{1}{2t}$, so by Lemma 4. we get

$$\int_{M} \eta = \int_{M} \exp\{-\frac{1}{2t}(d_{X+\sqrt{-1}Y}(X'+\sqrt{-1}Y'))\}\eta$$

Let V is a neighborhood of M_0 in \mathcal{N} . We identify a tubular neighborhood of M_0 in M with V. Set $V' \subset V$. When $t \to 0$, because $\langle X_M(x) + \sqrt{-1}Y_M(x), X_M(x) + \sqrt{-1}Y_M(x) \rangle \neq 0$ out of M_0 , so we have

$$\int_{M} \exp\{-\frac{1}{2t}(d_{X+\sqrt{-1}Y}(X'+\sqrt{-1}Y'))\}\eta \sim \int_{V'} \exp\{-\frac{1}{2t}(d_{X+\sqrt{-1}Y}(X'+\sqrt{-1}Y'))\}\eta$$

Because

$$\int_{V'} \exp\{-\frac{1}{2t}(d_{X+\sqrt{-1}Y}(X'+\sqrt{-1}Y'))\}\eta = \int_{V'} \exp\{-\frac{1}{2t}(d_{X,V}+\sqrt{-1}Y_{N}(X'_{N}+\sqrt{-1}Y'_{N}))\}\eta$$

then

$$\begin{split} \int_{V'} \exp\{-\frac{1}{2t}(d_{X+\sqrt{-1}Y}(X'+\sqrt{-1}Y'))\}\eta = \\ \int_{V'} \exp\{\frac{1}{t}\langle(\mu^{\mathcal{N}}(X)+\sqrt{-1}\mu^{\mathcal{N}}(Y))\cdot,\cdot\rangle + \frac{1}{2t}\langle\mu^{\mathcal{N}}(X)y+\sqrt{-1}\mu^{\mathcal{N}}(Y)y,R^{\mathcal{N}}y\rangle\}\eta \\ + \int_{V'} \exp\{-\frac{1}{2t}\langle-\mu^{\mathcal{N}}(X)y-\sqrt{-1}\mu^{\mathcal{N}}(Y)y,-\mu^{\mathcal{N}}(X)y-\sqrt{-1}\mu^{\mathcal{N}}(Y)y\rangle\}\eta \end{split}$$

By making the change of variables $y = \sqrt{ty}$, we find that the above formula is equal to

$$t^{n} \int_{V'} \exp\{\frac{1}{t} \langle (\mu^{\mathcal{N}}(X) + \sqrt{-1}\mu^{\mathcal{N}}(Y)) \cdot, \cdot \rangle + \frac{1}{2} \langle \mu^{\mathcal{N}}(X)y + \sqrt{-1}\mu^{\mathcal{N}}(Y)y, R^{\mathcal{N}}y \rangle \} \eta$$
$$+ \int_{V'} \exp\{-\frac{1}{2} \langle -\mu^{\mathcal{N}}(X)y - \sqrt{-1}\mu^{\mathcal{N}}(Y)y, -\mu^{\mathcal{N}}(X)y - \sqrt{-1}\mu^{\mathcal{N}}(Y)y \rangle \} \eta_{\sqrt{t}y}$$

we known that

$$\frac{\left(\frac{\langle (\mu^{\mathcal{N}}(\mathbf{X})+\sqrt{-1}\mu^{\mathcal{N}}(\mathbf{Y}))\cdot,\cdot\rangle}{t}\right)^n}{n!} = \left(\operatorname{Pf}(\mu^{\mathcal{N}}(\mathbf{X})+\sqrt{-1}\mu^{\mathcal{N}}(\mathbf{Y}))\right) \mathrm{dy}$$

here dy is the volume form of the submanifold M_0 , let n be the dimension of M_0 , then we get

$$= \int_{V'} \exp\{\frac{1}{2} \langle \mu^{\mathcal{N}}(X)y + \sqrt{-1}\mu^{\mathcal{N}}(Y)y, R^{\mathcal{N}}y \rangle\} \eta [\det(\mu^{\mathcal{N}}(X) + \sqrt{-1}\mu^{\mathcal{N}}(Y))]^{\frac{1}{2}} dy_1 \wedge \dots \wedge dy_n$$
$$+ \int_{V'} \exp\{-\frac{1}{2} \langle -\mu^{\mathcal{N}}(X)y - \sqrt{-1}\mu^{\mathcal{N}}(Y)y, -\mu^{\mathcal{N}}(X)y - \sqrt{-1}\mu^{\mathcal{N}}(Y)y \rangle\} \eta$$

Because by $[X_M, Y_M] = 0$ we have $[\mu^{TM}(X), \mu^{TM}(Y)] = 0$. And by $-\mu^{\mathcal{N}}(X) - \sqrt{-1}\mu^{\mathcal{N}}(Y)$, $R^{\mathcal{N}}$ are skew-symmetric, so we get

$$= \int_{V'} \exp\{-\frac{1}{2}\langle -\mu^{\mathcal{N}}(X)y - \sqrt{-1}\mu^{\mathcal{N}}(Y)y, -\mu^{\mathcal{N}}(X)y - \sqrt{-1}\mu^{\mathcal{N}}(Y)y + R^{\mathcal{N}}y\rangle\}dy_1 \wedge \dots \wedge dy_n$$
$$\cdot [\det(\mu^{\mathcal{N}}(X) + \sqrt{-1}\mu^{\mathcal{N}}(Y))]^{\frac{1}{2}}\eta$$

$$\begin{aligned} (L_{X_M} + \sqrt{-1}L_{Y_M})\widetilde{\nabla}^{TM}\omega &= (L_{X_M} + \sqrt{-1}L_{Y_M})(\nabla^{TM} + i_{X_M + \sqrt{-1}Y_M})\omega \\ &= (\nabla^{TM} + i_{X_M + \sqrt{-1}Y_M})(L_{X_M} + \sqrt{-1}L_{Y_M})\omega \\ &= 0 \end{aligned}$$

So we get $\widetilde{\nabla}^{TM}\omega \in \Omega^*_{X_M + \sqrt{-1}Y_M}(M, TM).$
We will also denote the restriction of $\widetilde{\nabla}^{TM}$ to $\Omega^* = -(M, TM)$ by $\widetilde{\nabla}^{TM}$

We will also denote the restriction of $\widetilde{\nabla}^{TM}$ to $\Omega^*_{X_M + \sqrt{-1}Y_M}(M, TM)$ by $\widetilde{\nabla}^{TM}$.

= 0

The equivariant curvature \widetilde{R}^{TM} of the equivariant connection $\widetilde{\nabla}^{TM}$ is defined by the formula(see [1])

$$\widetilde{R}^{TM} = (\widetilde{\nabla}^{TM})^2 - L_{X_M} - \sqrt{-1}L_{Y_M}$$

$$= \int_{M_0} (2\pi)^n [\det(\mu^{\mathcal{N}}(X) + \sqrt{-1}\mu^{\mathcal{N}}(Y))]^{-\frac{1}{2}} [\det(-\mu^{\mathcal{N}}(X) - \sqrt{-1}\mu^{\mathcal{N}}(Y) + R^{\mathcal{N}})]^{-\frac{1}{2}} \\ \cdot [\det(\mu^{\mathcal{N}}(X) + \sqrt{-1}\mu^{\mathcal{N}}(Y))]^{\frac{1}{2}} \eta \\ = \int_{M_0} (2\pi)^n [\det(-\mu^{\mathcal{N}}(X) - \sqrt{-1}\mu^{\mathcal{N}}(Y) + R^{\mathcal{N}})]^{-\frac{1}{2}} \eta \\ = \int_{M_0} \frac{\eta}{\Pr[\frac{-\mu^{\mathcal{N}}(X) - \sqrt{-1}\mu^{\mathcal{N}}(Y) + R^{\mathcal{N}}}{2\pi}]}$$

By theorem 1., we can get the localization formulas of Berline and Vergne(see [1] or [2]).

Corollery 2 (N.Berline and M.Vergne). Let M be a smooth closed oriented manifold, G be a compact Lie group acting smoothly on M. For any $\eta \in H^*_X(M)$, the following identity hold:

$$\int_{M} \eta = \int_{M_0} \frac{\eta}{\Pr\left[\frac{-\mu^{\mathcal{N}}(\mathbf{X}) + \mathbf{R}^{\mathcal{N}}}{2\pi}\right]}$$

Proof. By theorem 1., we set Y = 0, then we get the result.

Localization formulas for characteristic numbers 4

rv. G be a compact Lie group acting smoothly on M and \mathfrak{g} be its Lie algebra. Let g^{TM} be a G-invariant Riemannian metric on TM, ∇^{TM} is the Levi-Civita connection associated to g^{TM} . Here ∇^{TM} is a *G*-invariant connection, we see that $[\nabla^{TM}, L_{X_M}] = 0$ for all $X \in \mathfrak{g}$.

The equivariant connection $\widetilde{\nabla}^{TM}$ is the operator on $\Omega^*(M, TM)$ corresponding to a Ginvariant connection ∇^{TM} is defined by the formula

$$\tilde{\nabla}^{TM} = \nabla^{TM} + i_{X_M + \sqrt{-1}Y_M}$$

here X_M, Y_M be the smooth vector field on M corresponded to $X, Y \in \mathfrak{g}$.

Lemma 6. The operator $\widetilde{\nabla}^{TM}$ preserves the space $\Omega^*_{X_M+\sqrt{-1}Y_M}(M,TM)$ which is the space of smooth $(X_M + \sqrt{-1}Y_M)$ -invariant forms with values in TM.

Proof. Let $\omega \in \Omega^*_{X_M + \sqrt{-1}Y_M}(M)$, then we have

Let
$$M$$
 be an even dimensional compact oriented manifold without bounda

$$\Lambda_M + \sqrt{-11}M/(\Lambda_M + \sqrt{-1}M)$$

 $(L_{X,i} + \sqrt{-1}L_{Y,i})\omega$

It is the element of $\Omega^*_{X_M + \sqrt{-1}Y_M}(M, End(TM))$. We see that

$$\widetilde{R}^{TM} = (\nabla^{TM} + i_{X_M + \sqrt{-1}Y_M})^2 - L_{X_M} - \sqrt{-1}L_{Y_M} = R^{TM} + [\nabla^{TM}, i_{X_M + \sqrt{-1}Y_M}] - L_{X_M} - \sqrt{-1}L_{Y_M} = R^{TM} - \mu^{TM}(X) - \sqrt{-1}\mu^{TM}(Y)$$

Lemma 7. The equivariant curvature \widetilde{R}^{TM} satisfies the equivariant Bianchi formula

$$\widetilde{\nabla}^{TM}\widetilde{R}^{TM} = 0$$

Proof. Because

$$\begin{split} [\widetilde{\nabla}^{TM}, \widetilde{R}^{TM}] &= [\widetilde{\nabla}^{TM}, (\widetilde{\nabla}^{TM})^2 - L_{X_M} - \sqrt{-1}L_{Y_M}] \\ &= [\widetilde{\nabla}^{TM}, (\widetilde{\nabla}^{TM})^2] + [\nabla^{TM} + i_{X_M + \sqrt{-1}Y_M}, -L_{X_M} - \sqrt{-1}L_{Y_M}] \\ &= 0 \end{split}$$

Now we to construct the equivariant characteristic forms by \widetilde{R}^{TM} . If f(x) is a polynomial in the indeterminate x, then $f(\widetilde{R}^{TM})$ is an element of $\Omega^*_{X_M+\sqrt{-1}Y_M}(M, End(TM))$. We use the trace map

$$\operatorname{Tr}: \Omega^*_{X_M + \sqrt{-1}Y_M}(M, End(TM)) \to \Omega^*_{X_M + \sqrt{-1}Y_M}(M)$$

to obtain an element of $\Omega^*_{X_M+\sqrt{-1}Y_M}(M)$, which we call an equivariant characteristic form.

Lemma 8. The equivariant differential form $\operatorname{Tr}(f(\widetilde{R}^{TM}))$ is $d_{X_M+\sqrt{-1}Y_M}$ -closed, and its equivariant cohomology class is independent of the choice of the G-invariant connection ∇^{TM} .

Proof. If $\alpha \in \Omega^*_{X_M + \sqrt{-1}Y_M}(M, End(TM))$, because in local $\nabla^{TM} = d + \omega$, we have

$$\begin{aligned} d_{X_M + \sqrt{-1}Y_M} \operatorname{Tr}(\alpha) &= \operatorname{Tr}(d_{X_M + \sqrt{-1}Y_M} \alpha) \\ &= \operatorname{Tr}([d_{X_M + \sqrt{-1}Y_M}, \alpha]) + \operatorname{Tr}([\omega, \alpha]) \\ &= \operatorname{Tr}([\widetilde{\nabla}^{TM}, \alpha]) \end{aligned}$$

then by the equivariant Bianchi identity $\widetilde{\nabla}^{TM}\widetilde{R}^{TM} = 0$, we get

$$d_{X_M + \sqrt{-1}Y_M} \operatorname{Tr}(f(\widetilde{R}^{TM})) = 0.$$

Let ∇_t^{TM} is a one-parameter family of G-invariant connections with equivariant curvature \widetilde{R}_t^{TM} . We have

$$\begin{aligned} \frac{d}{dt} \operatorname{Tr}(f(\widetilde{R}_{t}^{TM})) &= \operatorname{Tr}(\frac{d\widetilde{R}_{t}^{TM}}{dt}f'(\widetilde{R}_{t}^{TM})) \\ &= \operatorname{Tr}(\frac{d(\widetilde{\nabla}_{t}^{TM})^{2}}{dt}f'(\widetilde{R}_{t}^{TM})) \\ &= \operatorname{Tr}([\widetilde{\nabla}_{t}^{TM}, \frac{d\widetilde{\nabla}_{t}^{TM}}{dt}]f'(\widetilde{R}_{t}^{TM})) \\ &= \operatorname{Tr}([\widetilde{\nabla}_{t}^{TM}, \frac{d\widetilde{\nabla}_{t}^{TM}}{dt}f'(\widetilde{R}_{t}^{TM})]) \\ &= d_{X_{M}+\sqrt{-1}Y_{M}}\operatorname{Tr}(\frac{d\widetilde{\nabla}_{t}^{TM}}{dt}f'(\widetilde{R}_{t}^{TM})) \end{aligned}$$

from which we get

$$\operatorname{Tr}(f(\widetilde{R}_1^{TM})) - \operatorname{Tr}(f(\widetilde{R}_0^{TM})) = d_{X_M + \sqrt{-1}Y_M} \int_0^1 \operatorname{Tr}(\frac{d\widetilde{\nabla}_t^{TM}}{dt} f'(\widetilde{R}_t^{TM})) dt$$

so we get the result.

As an application of Theorem 1., we can get the following localization formulas for characteristic numbers

Theorem 2. Let M be an 2*l*-dim compact oriented manifold without boundary, G be a compact Lie group acting smoothly on M and \mathfrak{g} be its Lie algebra. Let $X, Y \in \mathfrak{g}$, and X_M, Y_M be the corresponding smooth vector field on M. M_0 is the submanifold descriped in section 2. If f(x) is a polynomial, then we have

$$\int_{M} \operatorname{Tr}(f(\widetilde{R}^{TM})) = \int_{M_0} \frac{\operatorname{Tr}(f(\widetilde{R}^{TM}))}{\Pr[\frac{-\mu^{\mathcal{N}}(\mathbf{X}) - \sqrt{-1}\mu^{\mathcal{N}}(\mathbf{Y}) + \mathbf{R}^{\mathcal{N}}}{2\pi}]}$$

Proof. By Lemma 8., we have $\text{Tr}(f(\widetilde{R}^{TM}))$ is $d_{X_M + \sqrt{-1}Y_M}$ -closed. And by Theorem 1., we get the result.

We can use this formula to compute these characteristic numbers of M, especially we can use it to Euler characteristic of M. Here we didn't to give the details.

References

- [1] N. Berline, E. Getzler and M. Vergne *Heat Kernels and Dirac Operators*. Germany: Springer-Verlag, 1992.
- [2] N. Berline and M. Vergne. Zéros d'un champ de vecteurs et classes caractéristiques équivariantes. Duke Math. J., 50(2):539C549, 1983.
- [3] J.-M. Bismut. Localization formulas, superconnections, and the index theorem for families. Comm. Math. Phys., 103(1):127C166, 1986.
- [4] H. Jacobowitz. Non-vanishing complex vector fields and the Euler characteristic. Proc. Amer. Math. Soc., 2009, 137: 3163-3165.
- [5] S. Kobayashi. Transformation groups in differential geometry. Springer, New York(1972).
- [6] S. Kobayashi and K. Nomizu. *Foundations of differential geometry*. Vol. I. John Wiley and Sons, New York(1963).