

COMMUTATIVITY PROPERTIES OF QUINN SPECTRA

GERD LAURES AND JAMES E. MCCLURE

ABSTRACT. We give a simple sufficient condition for Quinn’s “bordism-type” spectra to be weakly equivalent to commutative symmetric ring spectra. We also show that the symmetric signature is (up to weak equivalence) a monoidal transformation between symmetric monoidal functors, which implies that the Sullivan-Ranicki orientation of topological bundles is represented by a ring map between commutative symmetric ring spectra. In the course of proving these statements we give a new description of symmetric L theory which may be of independent interest.

1. INTRODUCTION

In [Qui95], Frank Quinn gave a general machine for constructing spectra from “bordism-type theories.” In our paper [LM14] we gave axioms for a structure we call an *ad theory* and showed that when these axioms are satisfied (as they are for all of the standard examples) the Quinn machine can be improved to give a symmetric spectrum \mathbf{M} . We also showed that when the ad theory is multiplicative (that is, when its “target category” is graded monoidal) the symmetric spectrum \mathbf{M} is a symmetric ring spectrum. Finally, we showed that there are monoidal functors to the category of symmetric spectra which represent Poincaré bordism over $B\pi$ (considered as a functor of π) and symmetric L-theory (considered as a functor of a ring R with involution).

In this paper we consider commutativity properties. The reader does not need to have a thorough knowledge of our previous paper; the only sections that are relevant are 3, 6, 7, 9, 10, 17, 18 and 19, and only the definitions and the statements of the results (not the proofs) are used in the present paper.

A “commutative ad theory” is (essentially) an ad theory whose target category is graded symmetric monoidal (the precise definition is given in Section 3). Our first main result is

Date: September 22, 2021.

The second author was partially supported by NSF grants. He thanks the Lord for making his work possible.

Theorem 1.1. *Let \mathbf{M} be the symmetric ring spectrum associated to a commutative ad theory. There is a commutative symmetric ring spectrum \mathbf{M}^{comm} which is weakly equivalent in the category of symmetric ring spectra to \mathbf{M} .*

The proof gives a specific chain of weak equivalences (of length 2) between \mathbf{M} and \mathbf{M}^{comm} .

In particular, Theorem 1.1 shows that the L-theory spectrum of a commutative ring can be realized as a commutative symmetric ring spectrum.¹

For the ad theory ad_{STop} of oriented topological bordism ([LM14, Section 6]), we showed in [LM14, Section 17 and Appendix B] that the underlying spectrum of \mathbf{M}_{STop} is weakly equivalent to the usual Thom spectrum $M\text{STop}$. It is well-known that $M\text{STop}$ is a commutative symmetric ring spectrum, and we have

Theorem 1.2. *$(\mathbf{M}_{\text{STop}})^{\text{comm}}$ and $M\text{STop}$ are weakly equivalent in the category of commutative symmetric ring spectra.*

The proof gives a specific chain of weak equivalences between them.

We also prove a multiplicative property of the symmetric signature. The symmetric signature is a basic tool in surgery theory. In its simplest form, it assigns to an oriented Poincaré complex X an element of the symmetric L-theory of $\pi_1(X)$; this element determines the surgery obstruction up to 8-torsion. Ranicki proved that the symmetric signature of a Cartesian product is the product of the symmetric signatures ([Ran80b, Proposition 8.1(i)]). The symmetric signature gives a map of spectra from Poincaré bordism to L-theory ([KMM, Proposition 7.10]), and we showed in [LM14] that it gives a map of symmetric spectra. In order to investigate the multiplicativity of this map, we give a new (but equivalent) description of the L-spectrum, using “relaxed” algebraic Poincaré complexes (the relation between these and the usual algebraic Poincaré complexes is similar to the relation between Γ -spaces and E_∞ spaces). For a ring with involution R there is an ad theory ad_{rel}^R , and the associated spectrum $\mathbf{M}_{\text{rel}}^R$ is equivalent to the usual L-spectrum. The symmetric signature gives a map sig_{rel} from the Poincaré bordism spectrum (which we denote by $\mathbf{M}_{e,*,1}$; see [LM14, Section 7]) to $\mathbf{M}_{\text{rel}}^R$. We prove that this map is weakly equivalent to a ring map between commutative symmetric ring spectra:

¹Lurie [Lur] has explained another way to prove that the L-theory spectrum of a commutative ring can be realized as a commutative symmetric ring spectrum. The method used in [Lur] does not include other examples we consider such as Poincaré bordism and (in [BLM19]) Witt and IP bordism.

Theorem 1.3. *There are symmetric ring spectra \mathbf{A} and \mathbf{B} , commutative symmetric ring spectra \mathbf{C} and \mathbf{D} , and a commutative diagram*

$$\begin{array}{ccccc} \mathbf{M}_{e,*,1} & \longleftarrow & \mathbf{A} & \longrightarrow & \mathbf{C} \\ \text{sig}_{\text{rel}} \downarrow & & \downarrow & & \downarrow \\ \mathbf{M}_{\text{rel}}^{\mathbb{Z}} & \longleftarrow & \mathbf{B} & \longrightarrow & \mathbf{D} \end{array}$$

in which the horizontal arrows and the right vertical arrow are ring maps and the horizontal arrows are weak equivalences.

In fact \mathbf{C} is $(\mathbf{M}_{e,*,1})^{\text{comm}}$, and \mathbf{D} is weakly equivalent to $(\mathbf{M}_{\text{rel}}^{\mathbb{Z}})^{\text{comm}}$ in the category of commutative symmetric ring spectra (see Remark 18.3).

As far as we are aware, there is no previous result in the literature showing multiplicativity of the symmetric signature at the spectrum level.

In Section 19 we prove a stronger statement, that up to weak equivalence the symmetric signature is a monoidal transformation between symmetric monoidal functors. In [BLM19] we proved the analogous statement about the symmetric signature for Witt and IP bordism, using the methods of the present paper.

Remark 1.4. The *Sullivan-Ranicki orientation* for topological bundles ([Sul05], [MM79], [Ran92, Remark 16.3], [KMM, Section 13.5]) is the following composite in the homotopy category of spectra

$$MSTop \simeq \mathbf{Q}_{\text{STop}} \xrightarrow{\text{sig}} L^{\mathbb{Z}},$$

where \mathbf{Q}_{STop} denotes the Quinn spectrum of oriented topological bordism (which was shown to be equivalent to $MSTop$ in [LM14, Appendix B]). Combining Theorems 1.3 and 1.2 shows that the Sullivan-Ranicki orientation is represented by a ring map of commutative symmetric ring spectra.

The results of the present work were already used amongst others in [BL17] in connection with singularities of Baas-Sullivan type.

Here is an outline of the paper. In Sections 2 and 3 we give the definition of commutative ad theory. The proof of Theorem 1.1 occupies Sections 4–10. We begin in Sections 4 and 5 by giving a multiseisimplicial analogue $\Sigma\mathcal{S}_{\text{ss}}$ of the category of symmetric spectra. We observe that an ad theory gives rise to an object \mathbf{R} of $\Sigma\mathcal{S}_{\text{ss}}$ whose realization is the symmetric spectrum \mathbf{M} mentioned above. Section 6 explains the key idea of the proof, which is to interpolate between the various permutations of the multiplication map by allowing a different order

of multiplication for each cell. In Sections 7 and 8 we use this idea to create a monad in the category $\Sigma\mathcal{S}_{\text{ss}}$ which acts on \mathbf{R} , and in Section 10 (after a brief technical interlude in Section 9) we use a standard rectification argument (as in [May72]) to convert \mathbf{R} with this action to a strictly commutative object of $\Sigma\mathcal{S}_{\text{ss}}$; passage to geometric realization gives \mathbf{M}^{comm} . Next we turn to the proof of Theorem 1.3. In Sections 12–14 we introduce the relaxed symmetric Poincaré ad theory and the corresponding version of the symmetric signature. In Sections 15–17 we create a monad in the category $\Sigma\mathcal{S}_{\text{ss}} \times \Sigma\mathcal{S}_{\text{ss}}$ which acts on the pair $(\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^{\mathbb{Z}})$, and in Section 18 we adapt the argument of Section 10 to prove Theorem 1.3. Section 19 gives the statement of the stronger version of Theorem 1.3 mentioned above, and Section 20 gives the proof. Section 11 gives the proof of Theorem 1.2. There is a property of the smash product in symmetric spectra which is needed to prove the main results of the paper. It is taken care of in Appendix A. Appendix C investigates the functorial properties of the ad theories of relaxed quasi-symmetric complexes.

Acknowledgments. The authors benefited from a workshop on forms of homotopy theory held at the Fields Institute. They would like to thank Matthias Kreck for suggesting the problem to the first author and also Carl-Friedrich Bödigheimer, Jim Davis, Steve Ferry, John Harper, Mike Mandell, Frank Quinn, Andrew Ranicki, John Rognes, Stefan Schwede, Michael Weiss and Bruce Williams for useful hints and helpful discussions. The first author is grateful to the Max Planck Institute in Bonn for its hospitality. The authors also like to thank the first referee for pointing out the need for Appendix A.

2. SOME REDEFINITIONS

One of the ingredients in the definition of ad theory in [LM14] is the target “ \mathbb{Z} -graded category” \mathcal{A} (see [LM14, Definitions 3.3 and 3.10]). For the purposes of that paper, there was no reason to allow morphisms in \mathcal{A} between objects of the same dimension (except for identity maps). For the present paper, we do need such morphisms (see Definition 3.1 below and the proof of Theorem 1.1). We therefore begin by giving modified versions of some of the definitions of [LM14].

Definition 2.1. (cf. Definition 3.3 of [LM14]) A *\mathbb{Z} -graded category* is a small category \mathcal{A} with involution i , together with involution-preserving functors $d : \mathcal{A} \rightarrow \mathbb{Z}$ (called the *dimension function*) and $\emptyset : \mathbb{Z} \rightarrow \mathcal{A}$ such that $d\emptyset$ is equal to the identity functor. Note that if $d(a) > d(b)$ then there are no morphisms from a to b .

The definition of a strict monoidal structure on a \mathbb{Z} -graded category ([LM14, Definition 18.1]) needs no change, provided that one uses the new definition of \mathbb{Z} -graded category.

Next we explain how to modify the specific examples of target categories in [LM14] by adding morphisms which preserve dimension.

For the category $\mathcal{A}_{\text{STop}}$ ([LM14, Example 3.5]) the morphisms between objects of the same dimension are the orientation-preserving homeomorphisms.

For the category $\mathcal{A}_{\pi, \mathbb{Z}, w}$ ([LM14, Definition 7.3]) the morphisms between objects (X, f, ξ) and (X', f', ξ') of the same dimension are the maps $g : X \rightarrow X'$ such that $f' \circ g = f$ and $g_*(\xi) = \xi'$.

We do not need the analogous modification for the category \mathcal{A}^R ([LM14, Definition 9.5]) because we will be using the version in Section 12.

3. COMMUTATIVE AD THEORIES

Definition 3.1. Let \mathcal{A} be a \mathbb{Z} -graded category. A *permutative structure* on \mathcal{A} is a strict monoidal structure (\boxtimes, ε) ([LM14, Definition 18.1]) together with a natural isomorphism

$$\gamma_{x,y} : x \boxtimes y \rightarrow i^{|x||y|} y \boxtimes x$$

such that

- (a) $i\gamma_{x,y} = \gamma_{ix,y} = \gamma_{x,iy}$,
- (b) each of the maps $\gamma_{\emptyset,y}$, $\gamma_{x,\emptyset}$ is the identity map of \emptyset ,
- (c) the composite

$$x \boxtimes y \xrightarrow{\gamma_{x,y}} i^{|x||y|} y \boxtimes x \xrightarrow{i^{|x||y|}(\gamma_{y,x})} x \boxtimes y$$

is the identity.

- (d) $\gamma_{x,\varepsilon}$ is the identity, and
- (e) the diagram

$$\begin{array}{ccc} & x \boxtimes y \boxtimes z & \\ \swarrow 1 \boxtimes \gamma & & \searrow \gamma \\ i^{|y||z|} x \boxtimes z \boxtimes y & \xrightarrow{i^{|y||z|} \gamma \boxtimes 1} & i^{|z|(|x|+|y|)} z \boxtimes x \boxtimes y \end{array}$$

commutes.

Remark 3.2. The analogue of the coherence theorem for symmetric monoidal categories [ML63] holds in this context with essentially the same proof.

Definition 3.3. A *commutative ad theory* is a multiplicative ad theory [LM14, Definitions 3.10 and 18.4], with the extra property that every pre K -ad which is isomorphic to a K -ad is a K -ad, together with a permutative structure on the target category \mathcal{A} .

Examples are ad_C when C is a commutative DGA (see [LM14, Example 3.12]), ad_{STop} (see [LM14, Section 6]), $\text{ad}_{e,*,1}$ (see [LM14, Section 7]), $\text{ad}_{\text{STopFun}}$ (see [LM14, end of Section 8]), ad_{rel}^R when R is commutative (see Section 12 below), ad_{IP} ([BLM19, Section 4]) and ad_{IPFun} ([BLM19, Section 6.1]).

Remark 3.4. The extra property in Definition 3.3 is used in the proof of Lemma 6.3 below.

For later use we record some notation for iterated products.

Definition 3.5. (i) For a permutation $\eta \in \Sigma_j$, let $\epsilon(\eta)$ denote 0 if η is even and 1 if η is odd.

(ii) Let \mathcal{A} be a \mathbb{Z} -graded category with a permutative structure. Let $\eta \in \Sigma_j$. Define a functor

$$\eta_\star : \mathcal{A}^{\times j} \rightarrow \mathcal{A}$$

(where $\mathcal{A}^{\times j}$ is the j -fold Cartesian product) by

$$\eta_\star(x_1, \dots, x_j) = i^{\epsilon(\bar{\eta})}(x_{\eta^{-1}(1)} \boxtimes \cdots \boxtimes x_{\eta^{-1}(j)})$$

where $\bar{\eta}$ is the block permutation that takes blocks $\mathbf{b}_1, \dots, \mathbf{b}_j$ of size $|x_1|, \dots, |x_j|$ into the order $\mathbf{b}_{\eta^{-1}(1)}, \dots, \mathbf{b}_{\eta^{-1}(j)}$.

Remark 3.6. Note that, by Remark 3.2, $\eta_\star(x_1, \dots, x_j)$ is canonically isomorphic to $x_1 \boxtimes \cdots \boxtimes x_j$.

4. MULTISEMISIMPLICIAL SYMMETRIC SPECTRA

In this section we define a category $\Sigma\mathcal{S}_{\text{ss}}$ (the ss stands for “semisimplicial”) which is a multisemisimplicial version of the category $\Sigma\mathcal{S}$ of symmetric spectra. The motivation for the definition is that the sequence R_k in [LM14, Definition 17.2] should give an object of $\Sigma\mathcal{S}_{\text{ss}}$.

Recall that we write Δ_{inj} for the category whose objects are the sets $\{0, \dots, n\}$ and whose morphisms are the monotonically increasing injections.

A based k -fold *multisemisimplicial set* is a contravariant functor from the Cartesian product $(\Delta_{\text{inj}})^{\times k}$ to the category Set_* of based sets. In particular, a based 0-fold multisemisimplicial set is just a based set.

Next note that given a category \mathcal{C} with a left action of a group G one can define a category $G \times \mathcal{C}$ whose objects are those of \mathcal{C} and whose

morphisms are pairs (α, f) with $\alpha \in G$ and f a morphism of \mathcal{C} ; the domain of (α, f) is the domain of f and the target is α^{-1} applied to the target of f . Composition is defined by

$$(\alpha, f) \circ (\beta, g) = (\alpha\beta, \beta^{-1}(f) \circ g)$$

Remark 4.1. (i) \mathcal{C} is imbedded in $G \times \mathcal{C}$ by taking the morphism f of \mathcal{C} to the morphism (e, f) of $G \times \mathcal{C}$, where e is the identity element of G .

(ii) The morphism (α, f) is the composite $(\alpha, \text{id}) \circ (e, f)$.

Definition 4.2. Let Σ_k act on $(\Delta_{\text{inj}}^{\text{op}})^{\times k}$ by permuting the factors (when $k = 0$, Σ_0 is the trivial group). For each subgroup H of Σ_k let $H\text{ss}_k$ be the category of functors from $H \times (\Delta_{\text{inj}}^{\text{op}})^{\times k}$ to Set_* .

By Remark 4.1, an object of $H\text{ss}_k$ can be thought of as a based k -fold multiseisimplicial set with a left ‘‘action’’ of H in which H also acts on the multidegrees.

Definition 4.3. (i) A multiseisimplicial symmetric sequence \mathbf{X} is a sequence X_k , $k \geq 0$, such that X_k is an object of $\Sigma_k\text{ss}_k$.

(ii) A *morphism* of multiseisimplicial symmetric sequences from \mathbf{X} to \mathbf{Y} is a sequence of morphisms $f_k : X_k \rightarrow Y_k$ in $\Sigma_k\text{ss}_k$.

The category of multiseisimplicial symmetric sequences will be denoted by β .

For our next definition, recall [LM14, Definitions 17.2 and 17.3].

Definition 4.4. (i) For each $k \geq 0$ extend the object $R_k = \text{ad}^k(\Delta^\bullet)$ of ss_k to an object of $\Sigma_k\text{ss}_k$ by letting

$$(\alpha, \text{id})_*(F) = i^{\epsilon(\alpha)} \circ F \circ \alpha_\#$$

(where $\alpha \in \Sigma_k$ and $F \in \text{ad}^k(\Delta^n)$).

(ii) Let \mathbf{R} denote the object of β whose k -th term is R_k .

Next we assemble the ingredients needed to define a symmetric monoidal structure on β .

Definition 4.5. Given $A \in \Sigma_k\text{ss}_k$ and $B \in \Sigma_l\text{ss}_l$, define $A \wedge B \in (\Sigma_k \times \Sigma_l)\text{ss}_{k+l}$ by

$$(A \wedge B)_{\mathbf{m}, \mathbf{n}} = A_{\mathbf{m}} \wedge B_{\mathbf{n}}$$

(where \mathbf{m} is a k -fold multi-index and \mathbf{n} is an l -fold multi-index).

Definition 4.6. Given $H \subset G \subset \Sigma_k$, define a functor

$$I_H^G : H\text{ss}_k \rightarrow G\text{ss}_k$$

by letting $I_H^G A$ be the left Kan extension of A along $H \times (\Delta_{\text{inj}}^{\text{op}})^{\times k} \rightarrow G \times (\Delta_{\text{inj}}^{\text{op}})^{\times k}$.

Remark 4.7. For later use we give an explicit description of $I_H^G A$. For each multi-index \mathbf{n} , we have

$$(I_H^G A)_{\mathbf{n}} = \left(\bigvee_{\alpha \in G} A_{\alpha^{-1}(\mathbf{n})} \right) / H$$

where the action of H is defined as follows: if $\beta \in H$ and x is an element in the α -summand then β takes x to the element $(\beta, \text{id})_*(x)$ in the $\alpha\beta^{-1}$ -summand.

Notation 4.8. We denote the equivalence class of an element x in the α -summand of $I_H^G A$ by $[\alpha, x]$; note that $[\alpha, x] = (\alpha, \text{id})_*[e, x]$.

Definition 4.9. Given $\mathbf{X}, \mathbf{Y} \in \mathfrak{B}$, define $\mathbf{X} \otimes \mathbf{Y} \in \mathfrak{B}$ by

$$(\mathbf{X} \otimes \mathbf{Y})_k = \bigvee_{j_1 + j_2 = k} I_{\Sigma_{j_1} \times \Sigma_{j_2}}^{\Sigma_k} (\mathbf{X}_{j_1} \wedge \mathbf{Y}_{j_2}).$$

The proof that \otimes is a symmetric monoidal product is essentially the same as the corresponding proof in [HSS00, Section 2.1]. The symmetry map

$$\tau : \mathbf{X} \otimes \mathbf{Y} \rightarrow \mathbf{Y} \otimes \mathbf{X}$$

is given (as in [HSS00]) by

$$(4.1) \quad \tau([\alpha, x \wedge y]) = [\alpha \rho_{l,k}, y \wedge x]$$

where $x \in (X_k)_{\mathbf{m}}$, $y \in (Y_l)_{\mathbf{n}}$, and $\rho_{l,k}$ is the permutation of $\{1, \dots, k+l\}$ which moves the first l elements to the end and the last k elements to the front.

Next we will give the definition of the category $\Sigma \mathcal{S}_{\text{ss}}$ and its symmetric monoidal product. First we need a sphere object.

Definition 4.10. (i) Let S^1 be the based semisimplicial set that consists of the base point together with a 1-simplex s .

(ii) Let S^k be the object of $\Sigma_k \mathcal{S}_{\text{ss}}$ obtained from $(S^1)^{\wedge k}$ by letting

$$(\alpha, \text{id})_*(s \wedge \cdots \wedge s) = s \wedge \cdots \wedge s$$

(iii) Let \mathbf{S} be the object of \mathfrak{B} whose k -th term is S^k .

It is easy to check that \mathbf{S} is a commutative monoid in \mathfrak{B} .

Definition 4.11. $\Sigma \mathcal{S}_{\text{ss}}$ is the category of modules over \mathbf{S} .

Remark 4.12. One can give a more explicit version of this definition: an object of $\Sigma \mathcal{S}_{\text{ss}}$ consists of an object \mathbf{X} of \mathfrak{B} together with suspension maps

$$\omega : S^1 \wedge X_k \rightarrow X_{k+1}$$

for each k , such that the iterates of the ω 's satisfy appropriate equivariance conditions.

Example 4.13. The object \mathbf{R} of Definition 4.4 can be given suspension maps as follows: with the notation of [LM14, Definition 17.4(i)], define

$$\omega : S^1 \wedge R_k \rightarrow R_{k+1}$$

by

$$\omega(s \wedge F) = \lambda^*(F)$$

(where s is the 1-simplex of S^1 and $F \in \text{ad}^k(\Delta^n)$). The resulting object of $\Sigma\mathcal{S}_{\text{ss}}$ will also be denoted \mathbf{R} .

Definition 4.14. (cf. [HSS00, Definition 2.2.3]) For $\mathbf{X}, \mathbf{Y} \in \Sigma\mathcal{S}_{\text{ss}}$, define the *smash product* $\mathbf{X} \wedge \mathbf{Y}$ to be the coequalizer of the diagram

$$\mathbf{X} \otimes \mathbf{S} \otimes \mathbf{Y} \rightrightarrows \mathbf{X} \otimes \mathbf{Y}$$

where the right action of \mathbf{S} on \mathbf{X} is the composite

$$\mathbf{X} \otimes \mathbf{S} \rightarrow \mathbf{S} \otimes \mathbf{X} \rightarrow \mathbf{X}.$$

The proof that \wedge is a symmetric monoidal product is essentially the same as the corresponding proof in [HSS00, Section 2.2].

5. GEOMETRIC REALIZATION

Let G be a subgroup of Σ_k . By Remark 4.1(i), an object of $G\text{ss}_k$ has an underlying k -fold multisemisimplicial set.

Definition 5.1. The *geometric realization* $|A|$ of an object $A \in G\text{ss}_k$ is the geometric realization of its underlying k -fold multisemisimplicial set.

Definition 5.2. (i) A map in $G\text{ss}_k$ is a *weak equivalence* if it induces a weak equivalence of realizations.

(ii) A map $\mathbf{X} \rightarrow \mathbf{Y}$ in β or in $\Sigma\mathcal{S}_{\text{ss}}$ is a *weak equivalence* if each map $X_k \rightarrow Y_k$ is a weak equivalence.

Proposition 5.3. For $A \in \Sigma_k\text{ss}_k$, the following formula gives a natural left Σ_k action on $|A|$:

$$\alpha([u_1, \dots, u_k, a]) = [u_{\alpha^{-1}(1)}, \dots, u_{\alpha^{-1}(k)}, (\alpha, \text{id})_*(a)];$$

here $(u_1, \dots, u_k) \in \Delta^n$, $a \in A_n$, and $[u_1, \dots, u_k, a]$ denotes the class of (u_1, \dots, u_k, a) in $|A|$. \square

Proposition 5.4. For $H \subset G \subset \Sigma_k$ and $A \in H\text{ss}_k$ there is a natural isomorphism of based G -spaces

$$|I_H^G A| \cong G_+ \wedge_H |A|.$$

Proof. The proof is easy, using Remark 4.7. \square

Corollary 5.5. *Geometric realization induces a symmetric monoidal functor from $\Sigma\mathcal{S}_{\text{ss}}$ to the category of symmetric spectra $\Sigma\mathcal{S}$; in particular, the realization of a (commutative) monoid in $\Sigma\mathcal{S}_{\text{ss}}$ is a (commutative) monoid in $\Sigma\mathcal{S}$. \square*

6. A FAMILY OF MULTIPLICATION MAPS

In this section we begin the proof of Theorem 1.1. From now until the end of Section 10 we fix a \mathbb{Z} -graded permutative category \mathcal{A} and a commutative ad theory with values in \mathcal{A} . Let \mathbf{R} be the object of $\Sigma\mathcal{S}_{\text{ss}}$ constructed from this ad theory as in Example 4.13.

Let \mathbf{M} be the symmetric ring spectrum associated to the ad theory ([LM14, Proposition 17.5 and Theorem 18.5]). By definition, $M_k = |R_k|$. The multiplication of \mathbf{M} is induced by the collection of maps

$$\mu : (R_k)_{\mathbf{m}} \wedge (R_l)_{\mathbf{n}} \rightarrow (R_{k+l})_{\mathbf{m}, \mathbf{n}}$$

defined by

$$(6.1) \quad \mu(F \wedge G)(\sigma_1 \times \sigma_2, o_1 \times o_2) = i^{l \dim \sigma_1} F(\sigma_1, o_1) \boxtimes G(\sigma_2, o_2)$$

(this is well-defined because, by [LM14, Definition 18.1(b)], reversing the orientations o_1 and o_2 does not change the right-hand side). These maps give \mathbf{R} the structure of a monoid in $\Sigma\mathcal{S}_{\text{ss}}$ (the proof is essentially the same as for [LM14, Theorem 18.5]).

In general, even though the ad theory is commutative, \mathbf{R} is not a commutative monoid (this would require the product in the target category \mathcal{A} to be *strictly* graded commutative). Instead we have the following. Recall Definition 3.5 and Notation 4.8.

Lemma 6.1. *Let $m : \mathbf{R} \wedge \mathbf{R} \rightarrow \mathbf{R}$ be the product and let $\eta \in \Sigma_j$. Then the composite*

$$m_\eta : \mathbf{R}^{\wedge j} \xrightarrow{\eta} \mathbf{R}^{\wedge j} \xrightarrow{m} \mathbf{R}$$

is determined by the formula

$$\begin{aligned} m_\eta([e, F_1 \wedge \cdots \wedge F_j])(\sigma_1 \times \cdots \times \sigma_j, o_1 \times \cdots \times o_j) \\ = i^{\epsilon(\zeta)} \eta_\star(F_1(\sigma_1, o_1), \dots, F_j(\sigma_j, o_j)), \end{aligned}$$

where e is the identity element of the relevant symmetric group and ζ is the block permutation that takes blocks $\mathbf{b}_1, \dots, \mathbf{b}_j, \mathbf{c}_1, \dots, \mathbf{c}_j$ of size $\deg F_1, \dots, \deg F_j, \dim \sigma_1, \dots, \dim \sigma_j$ into the order $\mathbf{b}_1, \mathbf{c}_1, \dots, \mathbf{b}_j, \mathbf{c}_j$.

Proof. It suffices to prove this when η is a transposition, and in this case the proof is an easy calculation using Equations (4.1) and (6.1) and [LM14, Definition 17.3]. \square

The key idea in the proof of Theorem 1.1 is that there is a family of operations which can be used to interpolate between the various m_η . To construct this family, we allow a different permutation of the factors for each cell of $\Delta^{\mathbf{n}_1} \times \cdots \times \Delta^{\mathbf{n}_j}$, as explained in our next definition. We begin by defining the operations for pre-ads (see [LM14, Definitions 3.8(i) and 3.10(ii)]).

Definition 6.2. (i) Given a ball complex K , let $U(K)$ denote the set of all cells of K .

(ii) Let k_1, \dots, k_j be non-negative integers and let \mathbf{n}_i be a k_i -fold multi-index for $1 \leq i \leq j$. For any map

$$a : U(\Delta^{\mathbf{n}_1} \times \cdots \times \Delta^{\mathbf{n}_j}) \rightarrow \Sigma_j$$

define

$$a_* : \text{pre}^{k_1}(\Delta^{\mathbf{n}_1}) \times \cdots \times \text{pre}^{k_j}(\Delta^{\mathbf{n}_j}) \rightarrow \text{pre}^{k_1+\cdots+k_j}(\Delta^{(\mathbf{n}_1, \dots, \mathbf{n}_j)})$$

by

$$\begin{aligned} a_*(F_1, \dots, F_j)(\sigma_1 \times \cdots \times \sigma_j, o_1 \times \cdots \times o_j) \\ = i^{\epsilon(\zeta)}(a(\sigma_1 \times \cdots \times \sigma_j))_\star(F_1(\sigma_1, o_1), \dots, F_j(\sigma_j, o_j)), \end{aligned}$$

where ζ is the block permutation that takes blocks $\mathbf{b}_1, \dots, \mathbf{b}_j, \mathbf{c}_1, \dots, \mathbf{c}_j$ of size $k_1, \dots, k_j, \dim \sigma_1, \dots, \dim \sigma_j$ into the order $\mathbf{b}_1, \mathbf{c}_1, \dots, \mathbf{b}_j, \mathbf{c}_j$.

Lemma 6.3. *If $F_i \in \text{ad}^{k_i}(\Delta^{\mathbf{n}_i})$ for $1 \leq i \leq j$ then $a_*(F_1, \dots, F_j) \in \text{ad}^{k_1+\cdots+k_j}(\Delta^{(\mathbf{n}_1, \dots, \mathbf{n}_j)})$.*

Proof. This is immediate from the extra property in Definition 3.3, Remark 3.6, and [LM14, Definition 18.4(b)]. \square

Recalling that $(R_k)_\mathbf{n} = \text{ad}^k(\Delta^\mathbf{n})$ with basepoint at the trivial ad (see [LM14, Definitions 3.8(ii), 3.10(b) and 18.1(c)]), we have now constructed an operation

$$a_* : (R_{k_1})_{\mathbf{n}_1} \wedge \cdots \wedge (R_{k_j})_{\mathbf{n}_j} \rightarrow (R_{k_1+\cdots+k_j})_{\mathbf{n}_1, \dots, \mathbf{n}_j}$$

for each

$$a : U(\Delta^{\mathbf{n}_1} \times \cdots \times \Delta^{\mathbf{n}_j}) \rightarrow \Sigma_j.$$

For later use, we give the relation between a_* and the suspension map $\omega : S^1 \wedge R_k \rightarrow R_{k+1}$.

Definition 6.4. For a ball complex K , let

$$\Pi : U(\Delta^1 \times K) \rightarrow U(K)$$

be the map which takes $\sigma \times \tau$ to τ , where σ is a simplex of Δ^1 and τ is a simplex of K .

Lemma 6.5. *Let s be the 1-simplex of S^1 , let $F_i \in (R_{k_i})_{\mathbf{n}_i}$ for $1 \leq i \leq j$, and let $a : U(\Delta^{\mathbf{n}_1} \times \cdots \times \Delta^{\mathbf{n}_j}) \rightarrow \Sigma_j$. Then*

$$\omega(s \wedge a_*(F_1 \wedge \cdots \wedge F_j)) = (a \circ \Pi)_*(\omega(s \wedge F_1) \wedge \cdots \wedge F_j).$$

Proof. This follows from Lemma 6.1 (because the permuted multiplication commutes with suspension). It can also be proved by a straightforward calculation using Definitions 3.5 and 4.13 and [LM14, Definitions 17.4 and 3.7(ii)]. \square

In the remainder of this section, we show that the action of the operations a_* can be described in a way that begins to resemble the action of an operad; this resemblance will be developed further in the next two sections.

Definition 6.6. (i) For $j, k \geq 0$ define an object $\mathcal{O}(j)_k$ of $\Sigma_k \text{SS}_k$ by

$$(\mathcal{O}(j)_k)_{\mathbf{n}} = \text{Map}(U(\Delta^{\mathbf{n}}), \Sigma_j)_+$$

(where the $+$ denotes a disjoint basepoint); the morphisms in $(\Delta_{\text{inj}}^{\text{op}})^{\times k}$ act in the evident way, and the morphisms of the form (α, id) with $\alpha \in \Sigma_k$ act by permuting the factors in $\Delta^{\mathbf{n}}$.

(ii) For $j \geq 0$ define $\mathcal{O}(j)$ to be the object of β with k -th term $\mathcal{O}(j)_k$.

Definition 6.7. (i) For $A, B \in \Sigma_k \text{SS}_k$, define the *degreewise smash product*

$$A \bar{\wedge} B \in \Sigma_k \text{SS}_k$$

by

$$(A \bar{\wedge} B)_{\mathbf{n}} = A_{\mathbf{n}} \wedge B_{\mathbf{n}},$$

with the diagonal action of Σ_k .

(ii) For $\mathbf{X}, \mathbf{Y} \in \beta$, define $\mathbf{X} \bar{\wedge} \mathbf{Y} \in \beta$ by

$$(\mathbf{X} \bar{\wedge} \mathbf{Y})_k = X_k \bar{\wedge} Y_k.$$

Remark 6.8. The difference between the degreewise smash product $A \bar{\wedge} B$ and the previously defined smash product $A \wedge B$ is that the former is only defined when A and B are k -fold multisemisimplicial sets for the same k , and the result is again a k -fold multisemisimplicial set, whereas $A \wedge B$ is defined when A is k -fold and B is l -fold, and the result is $(k + l)$ -fold.

Our next definition assembles the operations a_* for a given j into a single map.

Definition 6.9. Let $j \geq 0$. Define a map

$$\phi_j : \mathcal{O}(j) \bar{\wedge} \mathbf{R}^{\otimes j} \rightarrow \mathbf{R}$$

in \mathfrak{B} by the formulas

$$\phi_j(a \wedge [e, F_1 \wedge \cdots \wedge F_j]) = a_*(F_1 \wedge \cdots \wedge F_j)$$

(where e denotes the identity element of the relevant symmetric group) and

$$\phi_j(a \wedge [\alpha, F_1 \wedge \cdots \wedge F_j]) = (\alpha, \text{id})_* \phi_j((\alpha^{-1}, \text{id})_* a \wedge [e, F_1 \wedge \cdots \wedge F_j]).$$

Lemma 6.10. *The map ϕ_j induces a map*

$$\psi_j : \mathcal{O}(j) \overline{\wedge} \mathbf{R}^{\wedge j} \rightarrow \mathbf{R}$$

in \mathfrak{B} .

Proof. This is a straightforward calculation using Example 4.13, Definition 4.14, and [LM14, Definition 17.3 and Lemma 18.7]. \square

7. A MONAD IN \mathfrak{B}

In the next section we will show that there is a monad \mathbb{P} in $\Sigma\mathcal{S}_{\text{ss}}$ with the property that the maps ψ_j constructed in Lemma 6.10 give an action of \mathbb{P} on \mathbf{R} . As preparation, in this section we prove the analogous result in \mathfrak{B} ; that is, we show that there is a monad \mathbb{O} in \mathfrak{B} for which the maps ϕ_j of Definition 6.9 give an action of \mathbb{O} on \mathbf{R} .

First we observe that the collection of objects $\mathcal{O}(j)$ has a composition map analogous to that of an operad. Recall that May defines an operad \mathcal{M} in the category of sets with $\mathcal{M}(j) = \Sigma_j$ ([May72, Definition 3.1(i)]). Let $\gamma_{\mathcal{M}}$ denote the composition operation in \mathcal{M} . Also recall Definition 4.9 and Notation 4.8.

Definition 7.1. Given $j_1, \dots, j_i \geq 0$ define a map

$$\gamma : \mathcal{O}(i) \overline{\wedge} (\mathcal{O}(j_1) \otimes \cdots \otimes \mathcal{O}(j_i)) \rightarrow \mathcal{O}(j_1 + \cdots + j_i)$$

in \mathfrak{B} by the formulas

$$\gamma(a \wedge [e, b_1 \wedge \cdots \wedge b_i])(\sigma_1 \times \cdots \times \sigma_i) = \gamma_{\mathcal{M}}(a(\sigma_1 \times \cdots \times \sigma_i), b_1(\sigma_1), \dots, b_i(\sigma_i))$$

(where e is the identity element of the relevant symmetric group) and

$$\gamma(a \wedge [\alpha, b_1 \wedge \cdots \wedge b_i]) = (\alpha, \text{id})_* \gamma((\alpha^{-1}, \text{id})_* a \wedge [e, b_1 \wedge \cdots \wedge b_i]).$$

In order to formulate the associativity property of γ , we note that for $\mathbf{X}_1, \dots, \mathbf{X}_i, \mathbf{Y}_1, \dots, \mathbf{Y}_i \in \mathfrak{B}$ there is a natural map

$$\chi : (\mathbf{X}_1 \overline{\wedge} \mathbf{Y}_1) \otimes \cdots \otimes (\mathbf{X}_i \overline{\wedge} \mathbf{Y}_i) \rightarrow (\mathbf{X}_1 \otimes \cdots \otimes \mathbf{X}_i) \overline{\wedge} (\mathbf{Y}_1 \otimes \cdots \otimes \mathbf{Y}_i)$$

given by

$$\chi([\alpha, x_1 \wedge y_1 \wedge \cdots \wedge x_i \wedge y_i]) = [\alpha, x_1 \wedge \cdots \wedge x_i] \wedge [\alpha, y_1 \wedge \cdots \wedge y_i].$$

Lemma 7.2. *The operation γ has the following associativity property: the composite*

$$\begin{aligned} & \emptyset(i) \overline{\wedge} \left(\left(\emptyset(j_1) \overline{\wedge} (\emptyset(l_{11}) \otimes \cdots \otimes \emptyset(l_{1j_1})) \right) \otimes \cdots \otimes \left(\emptyset(j_i) \overline{\wedge} (\emptyset(l_{i1}) \otimes \cdots \otimes \emptyset(l_{ij_i})) \right) \right) \\ & \xrightarrow{1\overline{\wedge}(\gamma \otimes \cdots \otimes \gamma)} \emptyset(i) \overline{\wedge} \left(\emptyset(l_{11} + \cdots + l_{1j_1}) \otimes \cdots \otimes \emptyset(l_{i1} + \cdots + l_{ij_i}) \right) \\ & \xrightarrow{\gamma} \emptyset(l_{11} + \cdots + l_{ij_i}) \end{aligned}$$

is the same as the composite

$$\begin{aligned} & \emptyset(i) \overline{\wedge} \left(\left(\emptyset(j_1) \overline{\wedge} (\emptyset(l_{11}) \otimes \cdots \otimes \emptyset(l_{1j_1})) \right) \otimes \cdots \otimes \left(\emptyset(j_i) \overline{\wedge} (\emptyset(l_{i1}) \otimes \cdots \otimes \emptyset(l_{ij_i})) \right) \right) \\ & \xrightarrow{1\overline{\wedge} \chi} \emptyset(i) \overline{\wedge} \left(\emptyset(j_1) \otimes \cdots \otimes \emptyset(j_i) \right) \overline{\wedge} \left(\emptyset(l_{11}) \otimes \cdots \otimes \emptyset(l_{i,j_i}) \right) \\ & \xrightarrow{\gamma \overline{\wedge} 1} \emptyset(j_1 + \cdots + j_i) \overline{\wedge} \left(\emptyset(l_{11}) \otimes \cdots \otimes \emptyset(l_{i,j_i}) \right) \\ & \xrightarrow{\gamma} \emptyset(l_{11} + \cdots + l_{ij_i}). \end{aligned}$$

□

To formulate the unital property of γ we first need to consider the unit object for the operation $\overline{\wedge}$.

Definition 7.3. Let \overline{S}_k be the object of $\Sigma_k \text{ss}_k$ which has a copy of S^0 in each multidegree (with each morphism of $\Sigma_k \times (\Delta_{\text{inj}}^{\text{op}})^{\times k}$ acting as the identity of S^0), and let \overline{S} be the object of β with k -th term \overline{S}_k .

Remark 7.4. (i) $\overline{S} \overline{\wedge} \mathbf{X} \cong \mathbf{X}$ for any $\mathbf{X} \in \beta$.

(ii) $\emptyset(0)$ and $\emptyset(1)$ are both equal to \overline{S} .

(iii) \overline{S} is a commutative monoid in β with multiplication

$$m : \overline{S} \otimes \overline{S} \rightarrow \overline{S}$$

given by

$$m([\alpha, s_1 \wedge s_2]) = t,$$

where s_1 and s_2 are any nontrivial simplices and t is the nontrivial simplex in the relevant multidegree.

Lemma 7.5. *The operation γ has the following unital property: the diagrams*

$$\begin{array}{ccc} \emptyset(j) \overline{\wedge} \overline{S}^{\otimes j} & \xrightarrow{=} & \emptyset(j) \overline{\wedge} \emptyset(1)^{\otimes j} \\ 1\overline{\wedge} m \downarrow & & \downarrow \gamma \\ \emptyset(j) \overline{\wedge} \overline{S} & \xrightarrow{\cong} & \emptyset(j) \end{array}$$

and

$$\begin{array}{ccc} \bar{\mathbf{S}} \bar{\wedge} \mathcal{O}(j) & \xrightarrow{\cong} & \mathcal{O}(j) \\ \downarrow = & \nearrow \gamma & \\ \mathcal{O}(1) \bar{\wedge} \mathcal{O}(j) & & \end{array}$$

commute. \square

To complete the analogy between γ and the composition map of an operad, we need an equivariance property.

Definition 7.6. Define a right action of Σ_j on $\mathcal{O}(j)$ by

$$(a\alpha)(\sigma) = a(\sigma) \cdot \alpha,$$

where $a \in \text{Map}(U(\Delta^n), \Sigma_j)_+$, $\sigma \in U(\Delta^n)$, and \cdot is multiplication in Σ_j .

Lemma 7.7. (i) The following diagram commutes for all $\alpha \in \Sigma_i$.

$$\begin{array}{ccc} \mathcal{O}(i) \bar{\wedge} (\mathcal{O}(j_1) \otimes \cdots \otimes \mathcal{O}(j_i)) & \longrightarrow & \mathcal{O}(i) \bar{\wedge} (\mathcal{O}(j_{\alpha^{-1}(1)}) \otimes \cdots \otimes \mathcal{O}(j_{\alpha^{-1}(i)})) \\ \downarrow \alpha \bar{\wedge} (\beta_1 \otimes \cdots \otimes \beta_i) & & \downarrow \gamma \\ & & \mathcal{O}(j_1 + \cdots + j_i) \\ & & \downarrow \gamma_M(\alpha, \beta_1, \dots, \beta_i) \\ \mathcal{O}(i) \bar{\wedge} (\mathcal{O}(j_1) \otimes \cdots \otimes \mathcal{O}(j_i)) & \xrightarrow{\gamma} & \mathcal{O}(j_1 + \cdots + j_i) \end{array}$$

\square

Now we use the data defined so far to construct a monad in the category \mathfrak{B} .

Definition 7.8. (i) For $\mathbf{X} \in \mathfrak{B}$, give $\mathcal{O}(j) \bar{\wedge} \mathbf{X}^{\otimes j}$ the diagonal right Σ_j action.

(ii) Define a functor $\mathbb{O} : \mathfrak{B} \rightarrow \mathfrak{B}$ by

$$\mathbb{O}(\mathbf{X}) = \bigvee_{j \geq 0} (\mathcal{O}(j) \bar{\wedge} \mathbf{X}^{\otimes j}) / \Sigma_j.$$

(iii) Define a natural transformation

$$\iota : \mathbf{X} \rightarrow \mathbb{O}\mathbf{X}$$

to be the composite

$$\mathbf{X} \xrightarrow{\cong} \bar{\mathbf{S}} \bar{\wedge} \mathbf{X} = \mathcal{O}(1) \bar{\wedge} \mathbf{X} \hookrightarrow \mathbb{O}(\mathbf{X}).$$

(iv) Define

$$\mu : \mathbb{O}\mathbb{O}\mathbf{X} \rightarrow \mathbb{O}\mathbf{X}$$

to be the natural transformation induced by the maps

$$\begin{aligned} \emptyset(i) \bar{\wedge} \left((\emptyset(j_1) \bar{\wedge} \mathbf{X}^{\otimes j_1}) \otimes \cdots \otimes (\emptyset(j_i) \bar{\wedge} \mathbf{X}^{\otimes j_i}) \right) \\ \xrightarrow{1\bar{\wedge}\chi} \left(\emptyset(i) \bar{\wedge} (\emptyset(j_1) \otimes \cdots \otimes \emptyset(j_i)) \right) \bar{\wedge} \mathbf{X}^{\otimes(j_1+\cdots+j_i)} \\ \xrightarrow{\gamma\bar{\wedge}1} \emptyset(j_1 + \cdots + j_i) \bar{\wedge} \mathbf{X}^{\otimes(j_1+\cdots+j_i)}. \end{aligned}$$

Proposition 7.9. *The transformations μ and ι define a monad structure on \mathbb{O} .*

Proof. This is immediate from Lemmas 7.2 and 7.5. \square

We conclude this section by giving the action of \mathbb{O} on \mathbf{R} . Observe that the map

$$\phi_j : \emptyset(j) \bar{\wedge} \mathbf{R}^{\otimes j} \rightarrow \mathbf{R}$$

of Definition 6.9 induces a map

$$(7.1) \quad (\emptyset(j) \bar{\wedge} \mathbf{R}^{\otimes j}) / \Sigma_j \rightarrow \mathbf{R}.$$

Definition 7.10. Define

$$\nu : \mathbb{O}\mathbf{R} \rightarrow \mathbf{R}$$

to be the map whose restriction to $(\emptyset(j) \bar{\wedge} \mathbf{X}^{\otimes j}) / \Sigma_j$ is the map (7.1).

Proposition 7.11. *ν is an action of \mathbb{O} on \mathbf{R} .*

Proof. We need to show that the diagrams

$$\begin{array}{ccc} \mathbf{R} & \xrightarrow{\iota} & \mathbb{O}\mathbf{R} \\ & \searrow = & \downarrow \nu \\ & & \mathbf{R} \end{array}$$

and

$$\begin{array}{ccc} \mathbb{O}\mathbb{O}\mathbf{R} & \xrightarrow{\mu} & \mathbb{O}\mathbf{R} \\ \mathbb{O}\nu \downarrow & & \downarrow \nu \\ \mathbb{O}\mathbf{R} & \xrightarrow{\nu} & \mathbf{R} \end{array}$$

commute. The first is obvious and for the second it suffices to check that the composite

$$\begin{aligned} \emptyset(i) \bar{\wedge} \left((\emptyset(j_1) \bar{\wedge} \mathbf{R}^{\otimes j_1}) \otimes \cdots \otimes (\emptyset(j_i) \bar{\wedge} \mathbf{R}^{\otimes j_i}) \right) \\ \xrightarrow{1\bar{\wedge}\chi} \emptyset(i) \bar{\wedge} (\emptyset(j_1) \otimes \cdots \otimes \emptyset(j_i)) \bar{\wedge} \mathbf{R}^{\otimes(j_1+\cdots+j_i)} \\ \xrightarrow{\gamma\bar{\wedge}1} \emptyset(j_1 + \cdots + j_i) \bar{\wedge} \mathbf{R}^{\otimes(j_1+\cdots+j_i)} \xrightarrow{\phi} \mathbf{R} \end{aligned}$$

is the same as the composite

$$\emptyset(i) \overline{\wedge} \left((\emptyset(j_1) \overline{\wedge} \mathbf{R}^{\otimes j_1}) \otimes \cdots \otimes (\emptyset(j_i) \overline{\wedge} \mathbf{R}^{\otimes j_i}) \right) \xrightarrow{1 \overline{\wedge} (\phi \otimes \cdots \otimes \phi)} \emptyset(i) \overline{\wedge} \mathbf{R}^{\otimes i} \xrightarrow{\phi} \mathbf{R}.$$

□

8. A MONAD IN $\Sigma \mathcal{S}_{\text{ss}}$

We begin by giving $\emptyset(j) \overline{\wedge} \mathbf{X}^{\wedge j}$ the structure of a multiseemisimplicial symmetric spectrum when $\mathbf{X} \in \Sigma \mathcal{S}_{\text{ss}}$. The definition is motivated by Lemma 6.5. Recall Definition 6.4.

Definition 8.1. Let $j, k \geq 0$. Let s be the 1-simplex of S^1 . Define

$$\omega : S^1 \wedge (\emptyset(j) \overline{\wedge} \mathbf{X}^{\wedge j})_k \rightarrow (\emptyset(j) \overline{\wedge} \mathbf{X}^{\wedge j})_{k+1}$$

as follows: for $a \in (\emptyset(j)_k)_{\mathbf{n}}$ and $x \in ((X^{\wedge j})_k)_{\mathbf{n}}$, let

$$\omega(s \wedge (a \wedge x)) = (a \circ \Pi) \wedge \omega(s \wedge x).$$

The identification given in Definition 4.14 was omitted in this notation.

Definition 8.2. (i) For $\mathbf{X} \in \Sigma \mathcal{S}_{\text{ss}}$, give $\emptyset(j) \overline{\wedge} \mathbf{X}^{\wedge j}$ the diagonal right Σ_j action.

(ii) Define a functor $\mathbb{P} : \Sigma \mathcal{S}_{\text{ss}} \rightarrow \Sigma \mathcal{S}_{\text{ss}}$ by

$$\mathbb{P}(\mathbf{X}) = \bigvee_{j \geq 0} (\emptyset(j) \overline{\wedge} \mathbf{X}^{\wedge j}) / \Sigma_j.$$

To give \mathbb{P} a monad structure we need

Lemma 8.3. *The composite in Definition 7.8(iv) induces a map*

$$\begin{aligned} \emptyset(i) \overline{\wedge} \left((\emptyset(j_1) \overline{\wedge} \mathbf{X}^{\wedge j_1}) \wedge \cdots \wedge (\emptyset(j_i) \overline{\wedge} \mathbf{X}^{\wedge j_i}) \right) \\ \rightarrow \emptyset(j_1 + \cdots + j_i) \overline{\wedge} \mathbf{X}^{\wedge (j_1 + \cdots + j_i)}. \end{aligned}$$

in $\Sigma \mathcal{S}_{\text{ss}}$.

□

Definition 8.4. (i) Define a natural transformation

$$\iota : \mathbf{X} \rightarrow \mathbb{P}\mathbf{X}$$

to be the composite

$$\mathbf{X} \xrightarrow{\cong} \overline{\mathbf{S}} \overline{\wedge} \mathbf{X} = \emptyset(1) \overline{\wedge} \mathbf{X} \hookrightarrow \mathbb{P}(\mathbf{X}).$$

(ii) Define

$$\mu : \mathbb{P}\mathbb{P}\mathbf{X} \rightarrow \mathbb{P}\mathbf{X}$$

to be the natural transformation induced by the maps constructed in Lemma 8.3.

Proposition 8.5. *The transformations μ and ι define a monad structure on \mathbb{P} .*

Proof. This follows from Proposition 7.9 by passage to quotients. \square

Next we give the action of \mathbb{P} on \mathbf{R} . By Definition 8.1 and Lemma 6.5, the map

$$\psi_j : \mathcal{O}(j) \overline{\wedge} \mathbf{R}^{\wedge j} \rightarrow \mathbf{R}$$

of Lemma 6.10 is a map in $\Sigma\mathcal{S}_{\text{ss}}$. It induces a map

$$(8.1) \quad (\mathcal{O}(j) \overline{\wedge} \mathbf{R}^{\wedge j}) / \Sigma_j \rightarrow \mathbf{R}.$$

in $\Sigma\mathcal{S}_{\text{ss}}$.

Definition 8.6. Define

$$\nu : \mathbb{P}\mathbf{R} \rightarrow \mathbf{R}$$

to be the map whose restriction to $(\mathcal{O}(j) \overline{\wedge} \mathbf{X}^{\wedge j}) / \Sigma_j$ is the map (8.1).

Proposition 8.7. *ν is an action of \mathbb{P} on \mathbf{R} .*

Proof. This follows from Proposition 7.11 by passage to quotients. \square

For use in Section 10 we record a lemma.

Lemma 8.8. (i) *There is a functor Υ from \mathbb{P} algebras to monoids in $\Sigma\mathcal{S}_{\text{ss}}$ (with respect to \wedge) which is the identity on objects.*

(ii) *The geometric realization of $\Upsilon(\mathbf{R})$ is the symmetric ring spectrum \mathbf{M} of [LM14, Theorem 18.5].*

Proof. Part (i). Let \mathbb{A} be the monad

$$\mathbb{A}(\mathbf{X}) = \bigvee_{j \geq 0} \mathbf{X}^{\wedge j}.$$

Then a monoid in $\Sigma\mathcal{S}_{\text{ss}}$ is the same thing as an \mathbb{A} -algebra, so it suffices to give a map of monads from \mathbb{A} to \mathbb{P} .

For each $j, k \geq 0$ and each k -fold multi-index \mathbf{n} , define an element

$$e_{j,k,\mathbf{n}} \in (\mathcal{O}(j)_k)_{\mathbf{n}}$$

to be the constant function $U(\Delta^{\mathbf{n}}) \rightarrow \Sigma_j$ whose value is the identity element of Σ_j . Next define a map

$$\overline{\mathbf{S}} \rightarrow \mathcal{O}(j)$$

by taking the nontrivial simplex of $(\overline{\mathbf{S}}_k)_{\mathbf{n}}$ to $e_{j,k,\mathbf{n}}$.

Now the composite

$$\mathbb{A}(\mathbf{X}) = \bigvee_{j \geq 0} \mathbf{X}^{\wedge j} \cong \bigvee_{j \geq 0} \overline{\mathbf{S}} \overline{\wedge} \mathbf{X}^{\wedge j} \rightarrow \bigvee_{j \geq 0} (\mathcal{O}(j) \overline{\wedge} \mathbf{X}^{\wedge j}) / \Sigma_j = \mathbb{P}(\mathbf{X})$$

is a map of monads.

Part (ii) is an easy consequence of the definitions. \square

9. DEGREEWISE SMASH PRODUCT AND GEOMETRIC REALIZATION

For the proof of Theorem 1.1 we need to know the relation between $\overline{\wedge}$ and geometric realization.

There is a natural map

$$\kappa : |A \overline{\wedge} B| \rightarrow |A| \wedge |B|$$

defined by

$$\kappa([u, x \wedge y]) = [u, x] \wedge [u, y].$$

The analogous map for multisimplicial sets is a homeomorphism, but the situation for multiseisimplicial sets is more delicate.

Definition 9.1. A multiseisimplicial set *has compatible degeneracies* if it is in the image of the forgetful functor from multisimplicial sets to multiseisimplicial sets.

Example 9.2. (i) One can define compatible degeneracies on $\mathcal{O}(j)_k$ for each $j, k \geq 0$ by using the codegeneracy maps between the Δ^n .

(ii) If $\mathbf{X} \in \Sigma \mathcal{S}_{ss}$ and X_k has compatible degeneracies for all k then each $(\mathcal{O}(j) \overline{\wedge} \mathbf{X}^{\wedge j})_k$ has compatible degeneracies.

Proposition 9.3. *If the underlying multiseisimplicial sets of A and B have compatible degeneracies then κ is a weak equivalence.*

Proof. Let \tilde{A} and \tilde{B} be multisimplicial sets whose underlying multiseisimplicial sets are A and B . Then the underlying multiseisimplicial set of the degreewise smash product $\tilde{A} \overline{\wedge} \tilde{B}$ is $A \overline{\wedge} B$. Consider the following commutative diagram, where $| \cdot |$ in the bottom row denotes realization of multisimplicial sets, $\tilde{\kappa}$ is defined analogously to κ , and the vertical maps collapse the degeneracies:

$$\begin{array}{ccc} |A \overline{\wedge} B| & \xrightarrow{\kappa} & |A| \wedge |B| \\ \downarrow & & \downarrow \\ |\tilde{A} \overline{\wedge} \tilde{B}| & \xrightarrow{\tilde{\kappa}} & |\tilde{A}| \wedge |\tilde{B}| \end{array}$$

The map $\tilde{\kappa}$ is a homeomorphism, and the vertical arrows are weak equivalences by the multisimplicial analogue of [Seg74, Lemma A.5], so κ is a weak equivalence. \square

Next we give a sufficient condition for a multiseisimplicial set to have compatible degeneracies. Let D^n denote the semisimplicial set consisting of the nondegenerate simplices of the standard simplicial n -simplex. For a multi-index \mathbf{n} , let $D^{\mathbf{n}}$ denote the k -fold multiseisimplicial set

$$D^{n_1} \times \cdots \times D^{n_k}.$$

Definition 9.4. (i) A *horn* in D^n is a subcomplex E which contains all elements of D^n except for the top-dimensional element and one of its faces.

(ii) A k -fold multiseisimplicial set A satisfies the *multi-Kan condition* if every map from a horn in D^n to A extends to a map $D^n \rightarrow A$.

The following result is proved in [McC].

Proposition 9.5. *If A satisfies the multi-Kan condition then it has compatible degeneracies.*

Our next result is proved in the same way as [LM14, Lemma 15.12] and does not require the ad theory to be commutative.

Proposition 9.6. *For each k , R_k satisfies the multi-Kan condition.*

10. RECTIFICATION

In this section we complete the proof of Theorem 1.1.

First we consider a monad in $\Sigma\mathcal{S}_{ss}$ which is simpler than \mathbb{P} .

Definition 10.1. (i) Define $\mathbb{P}'(\mathbf{X})$ to be $\bigvee_{j \geq 0} \mathbf{X}^{\wedge j} / \Sigma_j$.

(ii) For each $j \geq 0$, let

$$\xi_j : \mathcal{O}(j) \rightarrow \overline{\mathbf{S}}$$

be the map which takes each nontrivial simplex of $\mathcal{O}(j)_k$ to the nontrivial simplex of $\overline{\mathbf{S}}_k$ in the same multidegree. Define a natural transformation

$$\Xi : \mathbb{P} \rightarrow \mathbb{P}'$$

to be the wedge of the composites

$$(\mathcal{O}(j) \overline{\wedge} \mathbf{X}^{\wedge j}) / \Sigma_j \xrightarrow{\xi_j \overline{\wedge} 1} (\overline{\mathbf{S}} \overline{\wedge} \mathbf{X}^{\wedge j}) / \Sigma_j \xrightarrow{\cong} \mathbf{X}^{\wedge j} / \Sigma_j.$$

Proposition 10.2. (i) *An algebra over \mathbb{P}' is the same thing as a commutative monoid in $\Sigma\mathcal{S}_{ss}$.*

(ii) Ξ is a map of monads.

(iii) *Suppose that each X_k has compatible degeneracies (see Definition 9.1). Let \mathbb{P}^q denote the q -th iterate of \mathbb{P} . Then each map*

$$\Xi : \mathbb{P}^q(\mathbf{X}) \rightarrow \mathbb{P}'\mathbb{P}^{q-1}(\mathbf{X})$$

is a weak equivalence.

Parts (i) and (ii) are immediate from the definitions. Part (iii) will be proved at the end of this section.

Proof of Theorem 1.1. We apply the monadic bar construction ([May72, Construction 9.6]) to obtain simplicial objects $B_\bullet(\mathbb{P}, \mathbb{P}, \mathbf{R})$ and $B_\bullet(\mathbb{P}', \mathbb{P}, \mathbf{R})$ in $\Sigma\mathcal{S}_{\text{ss}}$. We write \mathbf{R}_\bullet for the constant simplicial object which is \mathbf{R} in each simplicial degree. There are maps of simplicial \mathbb{P} -algebras

$$(10.1) \quad \mathbf{R}_\bullet \xleftarrow{\varepsilon} B_\bullet(\mathbb{P}, \mathbb{P}, \mathbf{R}) \xrightarrow{\Xi_\bullet} B_\bullet(\mathbb{P}', \mathbb{P}, \mathbf{R}),$$

where ε is induced by the action of \mathbb{P} on \mathbf{R} (see [May72, Lemma 9.2(ii)]). The map ε is a homotopy equivalence of simplicial objects ([May72, Proposition 9.8]) and the map Ξ_\bullet is a weak equivalence in each simplicial degree by Propositions 9.5, 9.6 and 10.2(iii). $B_\bullet(\mathbb{P}', \mathbb{P}, \mathbf{R})$ is a simplicial algebra over \mathbb{P}' , which by Proposition 10.2(i) is the same thing as a simplicial commutative monoid in $\Sigma\mathcal{S}_{\text{ss}}$. Moreover, by Lemma 8.8(i), \mathbf{R}_\bullet and $B_\bullet(\mathbb{P}, \mathbb{P}, \mathbf{R})$ are simplicial monoids, and ε and Ξ_\bullet are maps of simplicial monoids.

The objects of the diagram (10.1) are simplicial objects in $\Sigma\mathcal{S}_{\text{ss}}$. We obtain a diagram

$$(10.2) \quad |\mathbf{R}_\bullet| \xleftarrow{|\varepsilon|} |B_\bullet(\mathbb{P}, \mathbb{P}, \mathbf{R})| \xrightarrow{|\Xi_\bullet|} |B_\bullet(\mathbb{P}', \mathbb{P}, \mathbf{R})|$$

of simplicial objects in $\Sigma\mathcal{S}$ (the category of symmetric spectra) by applying the geometric realization functor $\Sigma\mathcal{S}_{\text{ss}} \rightarrow \Sigma\mathcal{S}$ to the diagram (10.1) in each simplicial degree. The map $|\varepsilon|$ is a homotopy equivalence of simplicial objects and the map $|\Xi_\bullet|$ is a weak equivalence in each simplicial degree. The object $|B_\bullet(\mathbb{P}', \mathbb{P}, \mathbf{R})|$ is a simplicial commutative symmetric ring spectrum, the objects $|\mathbf{R}_\bullet|$ and $|B_\bullet(\mathbb{P}, \mathbb{P}, \mathbf{R})|$ are simplicial symmetric ring spectra, and the maps $|\varepsilon|$ and $|\Xi_\bullet|$ are maps of simplicial symmetric ring spectra.

Finally, we apply geometric realization to the diagram (10.2). We define \mathbf{M}^{comm} to be $||B_\bullet(\mathbb{P}', \mathbb{P}, \mathbf{R})||$. Now we have a diagram

$$(10.3) \quad \mathbf{M} = |\mathbf{R}| \xleftarrow{||\varepsilon||} ||B_\bullet(\mathbb{P}, \mathbb{P}, \mathbf{R})|| \xrightarrow{||\Xi_\bullet||} ||B_\bullet(\mathbb{P}', \mathbb{P}, \mathbf{R})|| = \mathbf{M}^{\text{comm}}$$

in $\Sigma\mathcal{S}$. The map $||\varepsilon||$ is a homotopy equivalence (cf. [May72, Corollary 11.9]) and the map $||\Xi_\bullet||$ is a weak equivalence by [Ree, Theorem E]. \mathbf{M}^{comm} is a commutative symmetric ring spectrum, \mathbf{M} is the symmetric ring spectrum of [LM14, Theorem 18.5] (by Lemma 8.8(ii)), $||B_\bullet(\mathbb{P}, \mathbb{P}, \mathbf{R})||$ is a symmetric ring spectrum, and $||\varepsilon||$ and $||\Xi_\bullet||$ are maps of symmetric ring spectra. \square

We conclude this section with the proof of part (iii) of Proposition 10.2. First we need a lemma (which for later use we state in more generality than we immediately need). Recall that a preorder is a set with a reflexive and transitive relation \leq . Examples are Σ_j , with

every element \leq every other, and $U(K)$ (see Definition 6.2(i)), with \leq induced by inclusions of cells.

Lemma 10.3. *Let P be a preorder with an element which is \geq all other elements, and let $k \geq 0$. Define a k -fold multisemisimplicial set A by*

$$A_{\mathbf{n}} = \text{Map}_{\text{preorder}}(U(\Delta^{\mathbf{n}}), P).$$

Then

- (i) A has compatible degeneracies, and
- (ii) A is weakly equivalent to a point.

Proof. For (i), we can give A compatible degeneracies by using the codegeneracy maps between the $\Delta^{\mathbf{n}}$.

Part (ii). Let \tilde{A} be a multisimplicial set whose underlying multisemisimplicial set is A . Let $d\tilde{A}$ be its diagonal. The multisimplicial analogue of [Seg74, Lemma A.5] implies that $|A|$ is weakly equivalent to $|\tilde{A}|$, and it is well-known that the latter is homeomorphic to $|d\tilde{A}|$. It therefore suffices to show that the simplicial set $d\tilde{A}$ is weakly equivalent to a point.

Let Δ_{simp}^n denote the standard simplicial n simplex and let $\partial\Delta_{\text{simp}}^n$ denote its boundary. Then it suffices by [GJ99, Theorem I.11.2] to show that every map from $\partial\Delta_{\text{simp}}^n$ to $d\tilde{A}$ extends to Δ_{simp}^n .

Let D^n (respectively ∂D^n) be the semisimplicial set consisting of the nondegenerate simplices of Δ_{simp}^n (resp., $\partial\Delta_{\text{simp}}^n$). Since Δ_{simp}^n (resp., $\partial\Delta_{\text{simp}}^n$) is the free simplicial set generated by D^n (respectively ∂D^n), it suffices to show that every semisimplicial map from ∂D^n to $d\tilde{A}$ extends to D^n , and this is obvious from the definition of A . \square

Note that if P is Σ_j with the preorder described above then A_+ is $\tilde{\mathcal{O}}(j)_k$.

Proof of 10.2(iii). We begin with the case $q = 1$, so we want to show that the map $\Xi : \mathbb{P}(\mathbf{X}) \rightarrow \mathbb{P}'(\mathbf{X})$ is a weak equivalence. It suffices to show that the map

$$(\mathcal{O}(j) \bar{\wedge} \mathbf{X}^{\wedge j})/\Sigma_j \rightarrow \mathbf{X}^{\wedge j}/\Sigma_j$$

is a weak equivalence for each j . Proposition A.2 and Remark A.3 show that the Σ_j actions are free away from the basepoint, so it suffices to show that each map

$$\mathcal{O}(j) \bar{\wedge} \mathbf{X}^{\wedge j} \rightarrow \mathbf{X}^{\wedge j}$$

is a weak equivalence. Now the object $(\mathcal{O}(j) \bar{\wedge} \mathbf{X}^{\wedge j})_k$ comes from

$$\bigvee_{k_1 + \dots + k_j = k} \mathcal{O}(j)_k \bar{\wedge} I_{\Sigma_{k_1} \times \dots \times \Sigma_{k_j}}^{\Sigma_k} (\mathbf{X}_{k_1} \wedge \dots \wedge \mathbf{X}_{k_j})$$

and we have

$$\mathcal{O}(j)_k \bar{\wedge} I_{\Sigma_{k_1} \times \cdots \times \Sigma_{k_j}}^{\Sigma_k} (\mathbf{X}_{k_1} \wedge \cdots \wedge \mathbf{X}_{k_j}) \cong I_{\Sigma_{k_1} \times \cdots \times \Sigma_{k_j}}^{\Sigma_k} \left(\mathcal{O}(j)_k \bar{\wedge} (\mathbf{X}_{k_1} \wedge \cdots \wedge \mathbf{X}_{k_j}) \right),$$

so it suffices by Proposition 5.4 to show that each map

$$\mathcal{O}(j)_k \bar{\wedge} (\mathbf{X}_{k_1} \wedge \cdots \wedge \mathbf{X}_{k_j}) \rightarrow \mathbf{X}_{k_1} \wedge \cdots \wedge \mathbf{X}_{k_j}$$

is a weak equivalence, and this follows from Example 9.2, Proposition 9.3, and Lemma 10.3.

The general case follows from the case $q = 1$ and Example 9.2(ii). \square

11. PROOF OF THEOREM 1.2

It is well known that Thom spectra are commutative symmetric ring spectra (see for example [Sch09]; we recall this below). In this section we show that the Thom spectrum $MSTop$ is weakly equivalent, in the category of commutative symmetric ring spectra, to the commutative symmetric ring spectrum $(\mathbf{M}_{S\text{Top}})^{\text{comm}}$ given by Theorem 1.1.

Our first task is to construct the following chain of weak equivalences in the category of symmetric spectra.

$$(11.1) \quad \mathbf{M}_{S\text{Top}} \xleftarrow{f_1} \mathbf{Y} \xrightarrow{f_2} \mathbf{X} \xrightarrow{f_3} MSTop$$

First recall that $MSTop$ has as k -th space the Thom space $T(S\text{Top}(k))$. The Σ_k action on $T(S\text{Top}(k))$ is induced by the conjugation action on $S\text{Top}(k)$.

For the construction of \mathbf{X} we need some facts about multisimplicial sets. Given a space Z and $k \geq 1$, let $S_{\bullet}^{k\text{-multi}}(Z)$ be the k -fold multisimplicial set whose simplices in multidegree \mathbf{n} are the maps $\Delta^{\mathbf{n}} \rightarrow Z$. There is a natural map

$$|S_{\bullet}^{k\text{-multi}}(Z)| \rightarrow Z$$

(where $|\cdot|$ denotes realization of the underlying multisimplicial set) which is a weak equivalence by [And] and the multisimplicial analogue of [Seg74, Lemma A.5]. If Z is a based space, there are natural maps

$$\lambda : \Sigma |S_{\bullet}^{k\text{-multi}}(Z)| \rightarrow |S^1 \wedge S_{\bullet}^{k\text{-multi}}(Z)|$$

and

$$\kappa : S^1 \wedge S_{\bullet}^{k\text{-multi}}(Z) \rightarrow S_{\bullet}^{(k+1)\text{-multi}}(\Sigma Z)$$

defined as follows. Given $t \in [0, 1]$, $u \in \Delta^{\mathbf{n}}$, and $g : \Delta^{\mathbf{n}} \rightarrow Z$, let \bar{t} denote the image of t under the oriented affine homeomorphism $[0, 1] \rightarrow \Delta^1$, and define

$$\lambda(t \wedge [u, g]) = [(\bar{t}, u), s \wedge g],$$

where s is the nontrivial simplex of S^1 . Define

$$\kappa(s \wedge g)(\bar{t}, u) = t \wedge g(u).$$

Then the diagram

$$(11.2) \quad \begin{array}{ccc} \Sigma |S_{\bullet}^{k-\text{multi}}(Z)| & \xrightarrow{\lambda} & |S^1 \wedge S_{\bullet}^{k-\text{multi}}(Z)| \\ \downarrow & & \downarrow |\kappa| \\ \Sigma Z & \longleftarrow & |S_{\bullet}^{(k+1)-\text{multi}}(\Sigma Z)| \end{array}$$

commutes.

Now let $X_k = |S_{\bullet}^{k-\text{multi}}(T(\text{STop}(k)))|$. We define the Σ_k action on X_k as follows. For $\alpha \in \Sigma_k$ and $g : \Delta^n \rightarrow T(\text{STop}(k))$, let $\alpha(\mathbf{n}) = (n_{\alpha^{-1}(1)}, \dots, n_{\alpha^{-1}(k)})$ and let $\alpha(g)$ be the composite

$$\Delta^{\alpha(\mathbf{n})} \xrightarrow{\alpha^{-1}} \Delta^n \xrightarrow{g} T(\text{STop}(k)) \xrightarrow{\alpha} T(\text{STop}(k)).$$

This makes $S_{\bullet}^{k-\text{multi}}(T(\text{STop}(k)))$ an object of $\Sigma_k \text{ss}_k$, and now Proposition 5.3 gives the Σ_k action on X_k . Next define the structure map

$$\Sigma X_k \rightarrow X_{k+1}$$

to be the composite

$$\begin{aligned} & \Sigma |S_{\bullet}^{k-\text{multi}}(T(\text{STop}(k)))| \xrightarrow{\lambda} |S^1 \wedge S_{\bullet}^{k-\text{multi}}(T(\text{STop}(k)))| \\ & \xrightarrow{|\kappa|} |S_{\bullet}^{(k+1)-\text{multi}}(\Sigma T(\text{STop}(k)))| \rightarrow |S_{\bullet}^{(k+1)-\text{multi}}(T(\text{STop}(k+1)))|, \end{aligned}$$

where the last map is induced by the structure map of $M\text{STop}$. Let \mathbf{X} be the symmetric spectrum consisting of the spaces X_k with these structure maps. Define $f_3 : \mathbf{X} \rightarrow M\text{STop}$ to be the sequence of weak equivalences

$$|S_{\bullet}^{k-\text{multi}}(T(\text{STop}(k)))| \rightarrow T(\text{STop}(k)).$$

The commutativity of diagram (11.2) shows that f_3 is a map of symmetric spectra.

Next let $S_{\bullet}^{k-\text{multi}, \heartsuit}(T(\text{STop}(k)))$ be the sub-multisemisimplicial set of $S_{\bullet}^{k-\text{multi}}(T(\text{STop}(k)))$ consisting of maps whose restrictions to each face of Δ^n are transverse to the zero section. Let \mathbf{Y} be the subspectrum of \mathbf{X} with k -th space $|S_{\bullet}^{k-\text{multi}, \heartsuit}(T(\text{STop}(k)))|$, and let $f_2 : \mathbf{Y} \rightarrow \mathbf{X}$ be the inclusion.

Lemma 11.1. *f_2 is a weak equivalence.*

Proof. Since $S_{\bullet}^{k-\text{multi}, \heartsuit}(T(\text{STop}(k)))$ and $S_{\bullet}^{k-\text{multi}}(T(\text{STop}(k)))$ satisfy the multi-Kan condition, they have compatible degeneracies by Proposition 9.5. It therefore suffices to show that the inclusion $S_{\bullet}^{k-\text{multi}, \heartsuit}(T(\text{STop}(k))) \subset S_{\bullet}^{k-\text{multi}}(T(\text{STop}(k)))$ induces a weak equivalence on the diagonal semi-simplicial sets, and this follows from [FQ90,

Section 9.6] and the definition of homotopy groups ([May92, Definition 3.6]). \square

It remains to construct f_1 . Let $S \subset T(\text{STop}(k))$ be the zero section. First we observe that, if $g : \Delta^n \rightarrow T(\text{STop}(k))$ is a map whose restriction to each face is transverse to S , we obtain an element $F \in \text{ad}_{\text{STop}}(\Delta^n)$ by letting $F(\sigma, o)$ be $g^{-1}(S) \cap \sigma$ with the orientation determined by o . This construction gives a map

$$S_{\bullet}^{k-\text{multi}, \uparrow}(T(\text{STop}(k))) \rightarrow (\mathbf{R}_{\text{STop}})_k,$$

in $\Sigma_k \text{ss}_k$, and applying geometric realization gives a Σ_k equivariant map $Y_k \rightarrow (\mathbf{M}_{\text{STop}})_k$; we let f_1 be the sequence of these maps.

Lemma 11.2. *f_1 is a weak equivalence.*

Proof. For a k -fold multisemisimplicial set A , let A' be the semisimplicial set whose n -th set is $A_{0, \dots, 0, n}$. There is an evident map

$$\phi : |A'| \rightarrow |A|.$$

If A is $S_{\bullet}^{k-\text{multi}}(Z)$, then A' is $S_{\bullet}(Z)$, and if A is R_k then A' is the semisimplicial set P_k of [LM14, Definition 15.4(i)], with realization $(\mathbf{Q}_{\text{STop}})_k$ ([LM14, Definitions 15.4(ii) and 15.8]). Now we have a commutative diagram

$$\begin{array}{ccccccc} (\mathbf{M}_{\text{STop}})_k & \xleftarrow{f_1} & Y_k & \xrightarrow{f_2} & X_k & \xrightarrow{f_3} & T(\text{STop}(k)) \\ \uparrow \phi_1 & & \uparrow \phi_2 & & & & \uparrow = \\ (\mathbf{Q}_{\text{STop}})_k & \xleftarrow{g_1} & |S_{\bullet}^{\uparrow}(T(\text{STop}(k)))| & \xrightarrow{g_2} & |S_{\bullet}(T(\text{STop}(k)))| & \xrightarrow{g_3} & T(\text{STop}(k)) \end{array}$$

Here g_3 is the usual weak equivalence, and g_2 is a weak equivalence by [FQ90, Section 9.6] and the definition of homotopy groups ([May92, Definition 3.6]), so ϕ_2 is a weak equivalence. g_1 was shown to be a weak equivalence in [LM14, Appendix B], and ϕ_1 was shown to be a weak equivalence in [LM14, Section 15], so f_1 is a weak equivalence as required. \square

This completes the construction of diagram (11.1).

Next we recall that $M\text{STop}$ is a commutative symmetric ring spectrum with product

$$T(\text{STop}(k)) \wedge T(\text{STop}(l)) \rightarrow T(\text{STop}(k+l)).$$

\mathbf{X} is also a commutative symmetric ring spectrum, with the product

$$\begin{aligned} & |S_{\bullet}^{k-\text{multi}}(T(\text{STop}(k)))| \wedge |S_{\bullet}^{l-\text{multi}}(T(\text{STop}(l)))| \\ & \rightarrow |S_{\bullet}^{(k+l)-\text{multi}}(T(\text{STop}(k)) \wedge T(\text{STop}(l)))| \rightarrow |S_{\bullet}^{(k+l)-\text{multi}}(T(\text{STop}(k+l)))|, \end{aligned}$$

and \mathbf{Y} is a commutative symmetric ring spectrum with the product it inherits from \mathbf{X} . The maps f_2 and f_3 are maps of symmetric ring spectra, so to complete the proof of Theorem 1.2, it suffices to show

Lemma 11.3. $(\mathbf{M}_{\text{STop}})^{\text{comm}}$ and \mathbf{Y} are isomorphic in the homotopy category of commutative symmetric ring spectra.

The proof of Lemma 11.3 is outsourced to Appendix B.

12. RELAXED SYMMETRIC POINCARÉ COMPLEXES

For a commutative ring R with the trivial involution, we would like to apply Theorem 1.1 to obtain a commutative model for the symmetric L-spectrum of R . However, the ad theory ad^R defined in Section 9 of [LM14], with the product defined in [LM14, Definition 9.12], is not commutative. The difficulty is that this product is defined using a noncommutative coproduct

$$\Delta : W \rightarrow W \otimes W.$$

In this section and the next we give an equivalent ad theory which is commutative.

Fix a ring R with involution. Recall the definition of the category \mathcal{D} from [LM14, Definition 9.2(v)].

Definition 12.1. A *relaxed quasi-symmetric complex of dimension n* is a quadruple (C, D, β, φ) , where C is an object of \mathcal{D} , D is an object of \mathcal{D} with a $\mathbb{Z}/2$ action, β is a quasi-isomorphism $C^t \otimes_R C \rightarrow D$ which is also a $\mathbb{Z}/2$ equivariant map, and φ is an element of $D_n^{\mathbb{Z}/2}$.

Example 12.2. (i) If (C, φ) is a quasi-symmetric complex as defined in [LM14, Definition 9.3], then the quadruple $(C, (C^t \otimes_R C)^W, \beta, \varphi)$ is a relaxed quasi-symmetric complex, where $\beta : C^t \otimes_R C \rightarrow (C^t \otimes_R C)^W$ is induced by the augmentation $W \rightarrow \mathbb{Z}$.

(ii) Relaxed quasi-symmetric complexes arise naturally from the construction of the symmetric signature of a Witt space given in [FM]; see [BLM19].

Definition 12.3. We define a category $\mathcal{A}_{\text{rel}}^R$ (the rel stands for relaxed) as follows. The objects of $\mathcal{A}_{\text{rel}}^R$ are the relaxed quasi-symmetric complexes. A morphism $(C, D, \beta, \varphi) \rightarrow (C', D', \beta', \varphi')$ is a pair $(f : C \rightarrow C', g : D \rightarrow D')$, where f and g are R -linear chain maps, g is $\mathbb{Z}/2$ equivariant, $g\beta = \beta'(f \otimes f)$, and (if $\dim \varphi = \dim \varphi'$) $g_*(\varphi) = \varphi'$.

Definition 12.4. A morphism $(f, g) : (C, D, \beta, \varphi) \rightarrow (C', D', \beta', \varphi')$ between objects of the same dimension is a *quasi-isomorphism* if f (and hence also g) is a quasi-isomorphism.

$\mathcal{A}_{\text{rel}}^R$ is a balanced ([LM14, Definition 5.1]) \mathbb{Z} -graded category, where i takes (C, D, β, φ) to $(C, D, \beta, -\varphi)$ and \emptyset_n is the n -dimensional object for which C and D are zero in all degrees.

Remark 12.5. The construction of Example 12.2(i) gives a morphism

$$\mathcal{A}^R \rightarrow \mathcal{A}_{\text{rel}}^R$$

of \mathbb{Z} -graded categories.

Next we must say what the K -ads with values in $\mathcal{A}_{\text{rel}}^R$ are. We need some preliminary definitions and a lemma. For a balanced pre K -ad F we will use the notation

$$F(\sigma, o) = (C_\sigma, D_\sigma, \beta_\sigma, \varphi_{\sigma, o}).$$

Recall [LM14, Definition 9.7].

Definition 12.6. A balanced pre K -ad F is *well-behaved* if C and D are well-behaved.

Next recall [LM14, Definition 12.2].

Lemma 12.7. *Let F be a well-behaved pre K -ad. Then*

- (i) $C^t \otimes_R C$ is well-behaved, and
- (ii) the map

$$(\beta_\sigma)_* : H_*((C^t \otimes_R C)_\sigma / (C^t \otimes_R C)_{\partial\sigma}) \rightarrow H_*(D_\sigma / D_{\partial\sigma})$$

is an isomorphism for each σ .

Proof. For part (i), first recall that (by [LM14, Definitions 9.7(b) and 9.6(ii)]) $C_{\partial\sigma}$ is a summand of C_σ for each σ . Thus, if we let \bar{C}_σ be the quotient $C_\sigma / C_{\partial\sigma}$, we have

$$C_\sigma \cong \bigoplus_{\tau \subset \sigma} \bar{C}_\tau.$$

Part (i) follows easily from this.

For part (ii), first observe that the fact that $C^t \otimes_R C$ and D are well-behaved implies that they are Reedy cofibrant ([Hir03, Definition 15.3.3(2)]). The colim that defines $(C^t \otimes_R C)_{\partial\sigma}$ is a hocolim by [Hir03, Theorem 19.9.1(1) and Proposition 15.10.2(2)], and similarly for $D_{\partial\sigma}$, so the map

$$(\beta_\sigma)_* : H_*((C^t \otimes_R C)_{\partial\sigma}) \rightarrow H_*(D_{\partial\sigma})$$

is an isomorphism by [Hir03, Theorem 19.4.2(1)], and this implies the lemma. \square

Recall [LM14, Example 3.12].

Definition 12.8. A balanced pre K -ad F is *closed* if, for each σ , the map

$$\text{cl}(\sigma) \rightarrow D_\sigma$$

which takes $\langle \tau, o \rangle$ to $\varphi_{\tau, o}$ is a chain map.

Note that if F is closed then $\varphi_{\sigma, o}$ represents an element $[\varphi_{\sigma, o}] \in H_*(D_\sigma/D_{\partial\sigma})$.

Notation 12.9. For a balanced pre K -ad F and a cell σ of K , let

$$j_\sigma : (C^t \otimes_R C)_\sigma / (C^t \otimes_R C)_{\partial\sigma} \rightarrow (C_\sigma / C_{\partial\sigma})^t \otimes_R C_\sigma$$

be the map that takes $[c \otimes c']$ to $[c] \otimes c'$.

The next definition is a little more complicated than the corresponding definition in [LM14, Section 9] in order to satisfy the extra condition in Definition 3.3.

Definition 12.10. (i) A *balanced K -ad* is a pre K -ad F with the following properties:

- (a) it is balanced, well-behaved and closed, and
- (b) for each σ the slant product with $(j_\sigma)_*(\beta_\sigma)_*^{-1}([\varphi_{\sigma, o}])$ is an isomorphism

$$H^*(\text{Hom}_R(C_\sigma, R)) \rightarrow H_{\dim \sigma - \deg F - *} (C_\sigma / C_{\partial\sigma}).$$

(ii) A K -ad is a pre K -ad which is naturally isomorphic to a balanced K -ad.

We write ad_{rel}^R for the set of K ads with values in $\mathcal{A}_{\text{rel}}^R$.

Remark 12.11. The morphism of Remark 12.5 takes ads to ads.

Theorem 12.12. ad_{rel}^R is an ad theory.

Remark 12.13. When R is commutative with the trivial involution, the proof will show that ad_{rel}^R is a commutative ad theory.

For the proof of Theorem 12.12 we need a product operation on ads.

Definition 12.14. (i) For $i = 1, 2$, let R_i be a ring with involution and let $(C^i, D^i, \beta^i, \varphi^i)$ be an object of $\mathcal{A}_{\text{rel}}^{R_i}$. Define

$$(C^1, D^1, \beta^1, \varphi^1) \otimes (C^2, D^2, \beta^2, \varphi^2)$$

to be the following object of $\mathcal{A}_{\text{rel}}^{R_1 \otimes R_2}$:

$$(C^1 \otimes C^2, D^1 \otimes D^2, \gamma, \varphi^1 \otimes \varphi^2),$$

where γ is the composite

$$(C^1 \otimes C^2)^t \otimes_{R_1 \otimes R_2} (C^1 \otimes C^2) \cong ((C^1)^t \otimes_{R_1} C^1) \otimes ((C^2)^t \otimes_{R_2} C^2) \xrightarrow{\beta^1 \otimes \beta^2} D^1 \otimes D^2.$$

(ii) For $i = 1, 2$, suppose given a ball complex K_i and a pre K_i -ad F_i of degree k_i with values in $\mathcal{A}_{\text{rel}}^{R_i}$. Define a pre $(K_1 \times K_2)$ -ad $F_1 \otimes F_2$ with values in $\mathcal{A}_{\text{rel}}^{R_1 \otimes R_2}$ by

$$(F_1 \otimes F_2)(\sigma \times \tau, o_1 \times o_2) = i^{k_2 \dim \sigma} F_1(\sigma, o_1) \otimes F_2(\tau, o_2).$$

Lemma 12.15. *For $i = 1, 2$, suppose given a ball complex K_i and a K_i -ad F_i with values in $\mathcal{A}_{\text{rel}}^{R_i}$. Then $F_1 \otimes F_2$ is a $(K_1 \times K_2)$ -ad. \square*

Proof of 12.12. We only need to verify parts (f) and (g) of [LM14, Definition 3.10].

For part (f), let F be a K' -ad. We may assumed it is balanced. Let

$$F(\sigma, o) = (C_\sigma, D_\sigma, \beta_\sigma, \varphi_{\sigma, o}).$$

We need to define a K -ad E which agrees with F on each residual subcomplex of K . As in the proof of [LM14, Theorem 6.5], we may assume by induction that K is a ball complex structure for the n disk with one n cell τ , and that K' is a subdivision of K which agrees with K on the boundary. We only need to define E on the top cell τ of K . We define $E(\tau, o)$ to be $(C_\tau, D_\tau, \beta_\tau, \varphi_{\tau, o})$, where

- $C_\tau = \text{colim}_{\sigma \in K'} C_\sigma$,
- $D_\tau = \text{colim}_{\sigma \in K'} D_\sigma$,
- $\beta = \text{colim}_{\sigma \in K'} \beta_\sigma$, and
- $\varphi_{\tau, o} = \sum \varphi_{\sigma, o'}$, where (σ, o') runs through the n -dimensional cells of K' with orientations induced by o .

The fact that E satisfies part (a) of Definition 12.10 is a consequence of [LM14, Proposition A.1(ii)]. We will deduce the isomorphism in part (b) of Definition 12.10 from [LM14, Proposition 12.4], and for this we need some facts from [LM14, Section 12].

First recall that for a well-behaved functor

$$B : \text{Cell}^b(K') \rightarrow \mathcal{D},$$

we write

$$\text{Nat}(\text{cl}, B)$$

for the chain complex of natural transformations of graded abelian groups; the differential is given by

$$\partial(\nu) = \partial \circ \nu - (-1)^{|\nu|} \nu \circ \partial.$$

Recall [LM14, Definition 12.3] and also the map Φ defined just before the statement of [LM14, Lemma 12.6]. Consider the diagram

$$\begin{array}{ccc} H_*(\text{Nat}(\text{cl}, D)) & \xrightarrow{\Phi} & H_{*+n}(D_\tau, D_{\partial\tau}) \\ \beta \uparrow & & \beta \uparrow \\ H_*(\text{Nat}(\text{cl}, C^t \otimes_R C)) & \xrightarrow{\Phi} & H_{*+n}((C^t \otimes_R C)_\tau, (C^t \otimes_R C)_{\partial\tau}). \end{array}$$

The horizontal maps are isomorphisms by [LM14, Lemma 12.6], and the right-hand vertical map is an isomorphism by the proof of Lemma 12.7(ii). Hence the left-hand vertical map is an isomorphism.

The collection $\{\varphi_{\sigma,o}\}$ gives a cycle ν in $\text{Nat}(\text{cl}, D)$. Let $\mu \in \text{Nat}(\text{cl}, C^t \otimes_R C)$ be a representative for $\beta^{-1}([\nu])$. Now fix an orientation o for τ . Let $\psi \in C_\tau^t \otimes_R C_\tau$ be $\sum \mu(\langle \sigma, o' \rangle)$, where (σ, o') runs through the n -dimensional cells of K' with orientations induced by o . Then ψ is a representative of $(j_\sigma)_* \beta^{-1}([\varphi_{\tau,o}])$, so it suffices to show that the cap product with ψ is an isomorphism $H^*(\text{Hom}_R(C_\tau, R)) \rightarrow H_{n-\text{deg } F-*}(C_\tau/C_{\partial\tau})$, and this follows from [LM14, Proposition 12.4].

It remains to verify part (g) of [LM14, Definition 3.10]. Let $0, 1, \iota$ denote the three cells of the unit interval I , with their standard orientations. As in the proof of [LM14, Theorem 9.11], it suffices to construct a relaxed symmetric Poincaré I -ad H over \mathbb{Z} which takes both 0 and 1 to the object $(\mathbb{Z}, \mathbb{Z}, \gamma, 1)$, where γ is the isomorphism $\mathbb{Z} \otimes \mathbb{Z} \rightarrow \mathbb{Z}$. The proof of [LM14, Theorem 9.11] gives a symmetric Poincaré I -ad G with $G(0) = G(1) = (\mathbb{Z}, \epsilon)$, where $\epsilon : W \rightarrow \mathbb{Z} \otimes \mathbb{Z}$ is the composite of the augmentation with γ^{-1} . Let us denote the object $G(\iota)$ by (C, φ) . Applying Remark 12.11 to G gives a relaxed symmetric Poincaré I -ad G' with $G'(\iota) = (C, (C \otimes C)^W, \beta, \varphi)$, where β is induced by the augmentation. Let e_0 (resp., e_1) be the inclusion $0 \hookrightarrow \iota$ (resp., $1 \hookrightarrow \iota$) and for $i = 0, 1$ let $g_i = G(e_i) : \mathbb{Z} \rightarrow C$. Then

$$\partial\varphi = (g_1 \otimes g_1) \circ \epsilon - (g_0 \otimes g_0) \circ \epsilon$$

because G is closed. We can therefore construct the required I -ad H from G' by replacing $G'(0)$ and $G'(1)$ by $(\mathbb{Z}, \mathbb{Z}, \gamma, 1)$. \square

13. EQUIVALENCE OF THE SPECTRA ASSOCIATED TO ad^R AND ad_{rel}^R

By Remark 12.11, the morphism

$$\mathcal{A}^R \rightarrow \mathcal{A}_{\text{rel}}^R$$

of Remark 12.5 induces a map of spectra

$$\mathbf{Q}^R \rightarrow \mathbf{Q}_{\text{rel}}^R$$

(see [LM14, Section 15]) and a map of symmetric spectra

$$\mathbf{M}^R \rightarrow \mathbf{M}_{\text{rel}}^R$$

(see [LM14, Section 17]).

Theorem 13.1. *The maps*

$$\mathbf{Q}^R \rightarrow \mathbf{Q}_{\text{rel}}^R$$

and

$$\mathbf{M}^R \rightarrow \mathbf{M}_{\text{rel}}^R.$$

are weak equivalences.

Remark 13.2. The method that will be used to prove Theorem 1.3 can be used to show that $\mathbf{M}^R \rightarrow \mathbf{M}_{\text{rel}}^R$ is weakly equivalent to a map of symmetric ring spectra.

Recall [LM14, Definitions 4.1 and 4.2]. By [LM14, Theorem 16.1, Remark 14.2(i) and Corollary 17.9(iii)], Theorem 13.1 follows from

Proposition 13.3. *The map of bordism groups*

$$\Omega_*^R \rightarrow (\Omega_{\text{rel}}^R)_*$$

is an isomorphism.

The proof of Proposition 13.3 will occupy the rest of this section. The following lemma proves surjectivity. As usual, for a chain complex A with a $\mathbb{Z}/2$ action, we write $A^{h\mathbb{Z}/2}$ for $(A^W)^{\mathbb{Z}/2}$. The augmentation induces a map $A^{\mathbb{Z}/2} \rightarrow A^{h\mathbb{Z}/2}$.

Lemma 13.4. *Let*

$$(C, D, \beta, \varphi)$$

be a relaxed symmetric Poincaré $*$ -ad and let

$$\psi \in (C^t \otimes_R C)^{h\mathbb{Z}/2}$$

represent the image of φ under the map

$$H_*(D^{\mathbb{Z}/2}) \rightarrow H_*(D^{h\mathbb{Z}/2}) \xleftarrow{\cong} H_*((C^t \otimes_R C)^{h\mathbb{Z}/2})$$

(where the isomorphism is induced by β). Then (C, ψ) is a symmetric Poincaré $*$ -ad, and (C, D, β, φ) is bordant to

$$(C, (C^t \otimes_R C)^W, \gamma, \psi),$$

where γ is induced by the augmentation.

For the proof we need another lemma.

Lemma 13.5. *Let (C, D, β, φ) be a relaxed symmetric Poincaré $*$ -ad.*

(i) *If $\psi \in D^{\mathbb{Z}/2}$ is any representative for the homology class $[\varphi] \in H_*(D^{\mathbb{Z}/2})$ then (C, D, β, ψ) is bordant to (C, D, β, φ) .*

(ii) *If*

$$(f, g) : (C, D, \beta, \varphi) \rightarrow (C', D', \beta', \varphi')$$

is a map of $$ -ads of the same dimension for which f (and hence also g) is a quasi-isomorphism then (C, D, β, φ) and $(C', D', \beta', \varphi')$ are bordant.*

The proof of Lemma 13.5 is deferred to the end of the section.

Proof of Lemma 13.4. The fact that (C, ψ) satisfies [LM14, Definition 9.9] (only part (b) is relevant) is immediate from Definition 12.10(b).

To see that (C, D, β, φ) and $(C, (C^t \otimes_R C)^W, \gamma, \psi)$ are bordant, let δ denote the composite

$$C^t \otimes_R C \xrightarrow{\beta} D \rightarrow D^W,$$

let $\omega \in D^{h\mathbb{Z}/2}$ be the image of φ , and let $\omega' \in D^{h\mathbb{Z}/2}$ be the image of ψ under the map $(C \otimes C)^{h\mathbb{Z}/2} \rightarrow D^{h\mathbb{Z}/2}$ induced by β . Part (ii) of Lemma 13.5 shows that (C, D, β, φ) and (C, D^W, δ, ω) are bordant, and also that $(C, (C^t \otimes_R C)^W, \gamma, \psi)$ and $(C, D^W, \delta, \omega')$ are bordant. But $[\omega] = [\omega']$ in $H_*(D^{h\mathbb{Z}/2})$, so the result follows from part (i) of Lemma 13.5. \square

Next we show that the map in Proposition 13.3 is 1-1. So let (C_0, φ_0) and (C_1, φ_1) be symmetric Poincaré $*$ -ads and let F be a relaxed symmetric Poincaré bordism between them. Let $0, 1, \iota$ denote the three cells of the unit interval I , with their standard orientations. Denote the object $F(\iota)$ by (C, D, β, φ) . It suffices to show that there is a symmetric Poincaré I -ad G with

$$(13.1) \quad G(0) = (C_0, \varphi_0), \quad G(1) = (C_1, \varphi_1), \quad G(\iota) = (C, \chi)$$

for an element χ which we will now construct.

φ represents an element

$$[\varphi] \in H_*(D^{\mathbb{Z}/2}, (C_0^t \otimes_R C_0)^{h\mathbb{Z}/2} \oplus (C_1^t \otimes_R C_1)^{h\mathbb{Z}/2}).$$

The map $D \rightarrow D^W$ induced by the augmentation gives a map

$$\begin{aligned} H_*(D^{\mathbb{Z}/2}, (C_0^t \otimes_R C_0)^{h\mathbb{Z}/2} \oplus (C_1^t \otimes_R C_1)^{h\mathbb{Z}/2}) \\ \rightarrow H_*(D^{h\mathbb{Z}/2}, ((C_0^t \otimes_R C_0)^W)^{h\mathbb{Z}/2} \oplus ((C_1^t \otimes_R C_1)^W)^{h\mathbb{Z}/2}); \end{aligned}$$

let x be the image of φ under this map. The map $\beta : C^t \otimes_R C \rightarrow D$ gives an isomorphism

$$\begin{aligned} (\beta^{h\mathbb{Z}/2})_* : H_*((C^t \otimes_R C)^{h\mathbb{Z}/2}, (C_0^t \otimes_R C_0)^{h\mathbb{Z}/2} \oplus (C_1^t \otimes_R C_1)^{h\mathbb{Z}/2}) \\ \rightarrow H_*(D^{h\mathbb{Z}/2}, ((C_0^t \otimes_R C_0)^W)^{h\mathbb{Z}/2} \oplus ((C_1^t \otimes_R C_1)^W)^{h\mathbb{Z}/2}); \end{aligned}$$

let $y = (\beta^{h\mathbb{Z}/2})_*^{-1}(x)$.

Lemma 13.6. *The image of y under the boundary map*

$$\begin{aligned} H_*((C^t \otimes_R C)^{h\mathbb{Z}/2}, (C_0^t \otimes_R C_0)^{h\mathbb{Z}/2} \oplus (C_1^t \otimes_R C_1)^{h\mathbb{Z}/2}) \\ \xrightarrow{\partial} H_{*-1}((C_0^t \otimes_R C_0)^{h\mathbb{Z}/2} \oplus (C_1^t \otimes_R C_1)^{h\mathbb{Z}/2}) \end{aligned}$$

is $-\varphi_0 + \varphi_1$.

Before proving this we conclude the proof of Proposition 13.3. The lemma implies that there is a representative χ of y with

$$(13.2) \quad \partial\chi = -\varphi_0 + \varphi_1.$$

It suffices to show that, with this choice of χ , the symmetric Poincaré pre I -ad G given by equation (13.1) is an ad. Equation (13.2) says that G is closed, and part (b) of [LM14, Definition 9.9] follows from Definition 12.10(b) and the fact that the image of $[\chi]$ under the map

$$\begin{aligned} H_*((C^t \otimes_R C)^{h\mathbb{Z}/2}, (C_0^t \otimes_R C_0)^{h\mathbb{Z}/2} \oplus (C_1^t \otimes_R C_1)^{h\mathbb{Z}/2}) \\ \rightarrow H_*((C^t \otimes_R C)^W, (C_0^t \otimes_R C_0)^W \oplus (C_1^t \otimes_R C_1)^W) \\ \xleftarrow{\cong} H_*(C^t \otimes_R C, (C_0^t \otimes_R C_0) \oplus (C_1^t \otimes_R C_1)) \end{aligned}$$

is the same as the image of $[\varphi]$ under the map

$$\begin{aligned} H_*(D^{\mathbb{Z}/2}, (C_0^t \otimes_R C_0)^{h\mathbb{Z}/2} \oplus (C_1^t \otimes_R C_1)^{h\mathbb{Z}/2}) \\ \rightarrow H_*(D, (C_0^t \otimes_R C_0)^W \oplus (C_1^t \otimes_R C_1)^W) \\ \xrightarrow{\beta_*^{-1}} H_*(C^t \otimes_R C, (C_0^t \otimes_R C_0) \oplus (C_1^t \otimes_R C_1)). \end{aligned}$$

□

Proof of Lemma 13.6. We know that the image of $[\varphi]$ under the boundary map

$$\begin{aligned} H_*(D^{\mathbb{Z}/2}, (C_0^t \otimes_R C_0)^{h\mathbb{Z}/2} \oplus (C_1^t \otimes_R C_1)^{h\mathbb{Z}/2}) \\ \xrightarrow{\partial} H_{*-1}((C_0^t \otimes_R C_0)^{h\mathbb{Z}/2} \oplus (C_1^t \otimes_R C_1)^{h\mathbb{Z}/2}) \end{aligned}$$

is $-\lceil\varphi_0\rceil + \lceil\varphi_1\rceil$, so it suffices to show that for $i = 0, 1$ the maps

$$(C_i^t \otimes_R C_i)^{h\mathbb{Z}/2} \rightarrow ((C_i^t \otimes_R C_i)^W)^{h\mathbb{Z}/2}$$

induced by $D \rightarrow D^W$ and by β give the same map in homology. If we think of these as maps

$$a_i, b_i : ((C_i^t \otimes_R C_i)^W)^{\mathbb{Z}/2} \rightarrow ((C_i^t \otimes_R C_i)^{W \otimes W})^{\mathbb{Z}/2}$$

(with diagonal $\mathbb{Z}/2$ action on $W \otimes W$) then a_i and b_i are induced by the maps

$$e_1, e_2 : W \otimes W \rightarrow W$$

given by the augmentations on the two factors. Now the $\mathbb{Z}/2$ equivariant map

$$\Delta : W \rightarrow W \otimes W$$

of [Ran80a, page 175] has the property that $e_1 \circ \Delta$ and $e_2 \circ \Delta$ are both the identity map, so if

$$d : ((C_i^t \otimes_R C_i)^{W \otimes W})^{\mathbb{Z}/2} \rightarrow ((C_i^t \otimes_R C_i)^W)^{\mathbb{Z}/2}$$

is the map induced by Δ then $d \circ a_i$ and $d \circ b_i$ are both the identity map. But Δ is a $\mathbb{Z}/2$ chain homotopy equivalence, so d is a homology isomorphism and it follows that a_i and b_i induce the same map in homology as required. \square

It remains to prove Lemma 13.5. Let F be the cylinder of (C, D, β, φ) (which was constructed in the last paragraph of the proof of Theorem 12.12). Then $F(0)$ and $F(1)$ are both (C, D, β, φ) . Write

$$F(\iota) = (C_\iota, D_\iota, \beta_\iota, \varphi_\iota)$$

and let

$$(h, k) : (C, D, \beta, \varphi) \rightarrow (C_\iota, D_\iota, \beta_\iota, \varphi_\iota)$$

be the map $F(1) \rightarrow F(\iota)$.

For part (i), the hypothesis gives an element $\rho \in D^{\mathbb{Z}/2}$ with $\partial\rho = \psi - \varphi$. Let $\rho' \in D_\iota^{\mathbb{Z}/2}$ be the image of ρ under $k : D \rightarrow D_\iota$. Define an I -ad G by

$$G(0) = (C, D, \beta, \varphi), \quad G(1) = (C, D, \beta, \psi), \quad G(\iota) = (C_\iota, D_\iota, \beta_\iota, \varphi_\iota + \rho').$$

Then G is the desired bordism.

For part (ii), we first show that (C, D, β, φ) is bordant to $(C, D', \beta_1, \varphi')$, where β_1 is the composite

$$C^t \otimes_R C \xrightarrow{\beta} D \xrightarrow{g} D'$$

The idea is to construct a suitable mapping cylinder. Let D_1 be the pushout of the diagram

$$\begin{array}{ccc} D & \xrightarrow{k} & D_\iota \\ g \downarrow & & \\ & & D' \end{array}$$

Let φ_1 be the image of φ_ι in D_1 and let β_2 be the composite

$$C_\iota^t \otimes_R C_\iota \rightarrow D_\iota \rightarrow D_1.$$

Define an I -ad H by

$$H(0) = (C, D, \beta, \varphi), \quad H(1) = (C, D', \beta_1, \varphi'), \quad H(\iota) = (C_\iota, D_1, \beta_2, \varphi_1).$$

Then H is the desired bordism.

To conclude the proof we show that $(C, D', \beta_1, \varphi')$ is bordant to $(C', D', \beta', \varphi')$. Let C_1 be the pushout of the diagram

$$\begin{array}{ccc} C & \xrightarrow{h} & C_\iota \\ f \downarrow & & \\ & & C' \end{array}$$

Let $(C'_\iota, D'_\iota, \beta'_\iota, \varphi'_\iota)$ be the cylinder of $(C', D', \beta', \varphi')$. Let β_3 be the map

$$C_1^t \otimes_R C_1 \rightarrow D'_\iota.$$

Define an I -ad H_1 by

$$H_1(0) = (C, D', \beta_1, \varphi'), \quad H_1(1) = (C', D', \beta', \varphi'), \quad H_1(\iota) = (C_1, D'_\iota, \beta_3, \varphi'_\iota).$$

Then H_1 is the desired bordism. \square

14. THE SYMMETRIC SIGNATURE REVISITED

Fix a group π , a simply-connected free π -space Z , and a homomorphism $w : \pi \rightarrow \{\pm 1\}$, and recall the symmetric spectrum $\mathbf{M}_{\pi, Z, w}$ ([LM14, Sections 7 and 17]) which represents w -twisted Poincaré bordism over Z/π .

Let R denote the group ring $\mathbb{Z}[\pi]$ with the w -twisted involution ([Ran80b, page 196]).

Let us say that a ball complex is *strict* if each component of the intersection of two cells is a single cell. In [LM14, Section 10] we gave a functor

$$\text{sig} : \mathcal{A}_{\pi, Z, w} \rightarrow \mathcal{A}^R$$

which induces a natural transformation

$$\text{sig} : \text{ad}_{\pi, Z, w}(K) \rightarrow \text{ad}^R(K)$$

for strict ball complexes K .

Remark 14.1. (i) The restriction to strict ball complexes was not mentioned in [LM14] but is necessary, because if K has a cell τ whose boundary is not strict and if F is a K -ad $(X_\sigma, f_\sigma, \xi_{\sigma,o})$ then the map

$$(14.1) \quad \operatorname{colim}_{\sigma \subset \partial\tau} S_*(X_\tau) \rightarrow S_*(X_{\partial\tau})$$

is not a monomorphism (because simplices with support in $\sigma \cap \sigma'$ but not in a cell of $\sigma \cap \sigma'$ will have two representatives in the colimit), and hence $\operatorname{sig} \circ F$ is not well-behaved.

(ii) If K is strict then the map (14.1) has a left inverse for all τ (because its image is the subcomplex of $S_*(X_\tau)$ generated by the simplices that land in some X_σ with $\sigma \subset \partial\tau$, and the left inverse takes each such simplex to a representative for it in the colimit system; all such representatives are identified because K is strict). Hence the map (14.1) is the inclusion of a direct summand, as required for $\operatorname{sig} \circ F$ to be well-behaved.

(iii) The restriction to strict ball complexes does not affect the results about the symmetric signature in [LM14] because the only ball complexes that occur in [LM14, Sections 15, 17–19] are products of simplices, and these are strict.

In this section we give a functor

$$\operatorname{sig}_{\text{rel}} : \mathcal{A}_{\pi, Z, w} \rightarrow \mathcal{A}_{\text{rel}}^R$$

which induces a natural transformation

$$\operatorname{sig}_{\text{rel}} : \operatorname{ad}_{\pi, Z, w}(K) \rightarrow \operatorname{ad}_{\text{rel}}^R(K)$$

for strict ball complexes K .

Let (X, f, ξ) be an object of $\mathcal{A}_{\pi, Z, w}$ ([LM14, Definition 7.3]).

In the special case where π is the trivial group and Z is a point, the definition is easy:

$$\operatorname{sig}_{\text{rel}}(X, f, \xi) = (S_*X, S_*(X \times X), \beta, \varphi),$$

where β is the cross product $S_*X \otimes S_*X \xrightarrow{\times} S_*(X \times X)$ and φ is the image of ξ under the diagonal map.

The definition in the general case is similar. Recall that we write \tilde{X} for the pullback of Z along f and \mathbb{Z}^w for \mathbb{Z} with the right R action determined by w . Also recall [LM14, Definition 7.1].

Definition 14.2. (i) Give $S_*(\tilde{X})$ the left R module structure determined by the action of π on \tilde{X} and give $S_*(\tilde{X} \times \tilde{X})$ and $S_*(\tilde{X}) \otimes S_*(\tilde{X})$ the left R module structures determined by the diagonal actions of π .

(ii) Define

$$\operatorname{sig}_{\text{rel}}(X, f, \xi) = (S_*(\tilde{X}), \mathbb{Z}^w \otimes_R S_*(\tilde{X} \times \tilde{X}), \beta, \varphi),$$

where β is the composite

$$S_*(\tilde{X})^t \otimes_R S_*(\tilde{X}) \cong \mathbb{Z}^w \otimes_R (S_*(\tilde{X}) \otimes S_*(\tilde{X})) \xrightarrow{1 \otimes \times} \mathbb{Z}^w \otimes_R S_*(\tilde{X} \times \tilde{X})$$

and φ is the image of ξ under the map

$$S_*(X, \mathbb{Z}^f) = \mathbb{Z}^w \otimes_R S_*(\tilde{X}) \rightarrow \mathbb{Z}^w \otimes_R S_*(\tilde{X} \times \tilde{X})$$

(where the unmarked arrow is induced by the diagonal map).

Remark 14.3. (i) For set-theoretic reasons one should modify this definition as in [LM14, Section 10]; we leave this to the reader.

(ii) For strict ball complexes, sig_{rel} takes ads to ads, because the composite of the cross product with the Alexander-Whitney map is naturally chain homotopic to the map induced by the diagonal.

Next we compare sig to sig_{rel} .

Let us denote by δ both the map $\mathcal{A}^R \rightarrow \mathcal{A}_{\text{rel}}^R$ of Remark 12.5 and the map $\mathbf{M}^R \rightarrow \mathbf{M}_{\text{rel}}^R$ which it induces.

Proposition 14.4. *The diagram*

$$\begin{array}{ccc} & \mathbf{M}_{\pi, Z, w} & \\ \text{sig} \swarrow & & \searrow \text{sig}_{\text{rel}} \\ \mathbf{M}^R & \xrightarrow{\delta} & \mathbf{M}_{\text{rel}}^R \end{array}$$

commutes in the homotopy category of symmetric spectra.

We will derive this from a more general result. Recall Definition 12.4.

Proposition 14.5. *Let Θ be an ad theory, let R be a ring with involution, and let S_1 and S_2 be morphisms of ad theories $\Theta \rightarrow \text{ad}_{\text{rel}}^R$. Suppose that there is a natural quasi-isomorphism $\nu : S_1 \rightarrow S_2$. Then the maps $\mathbf{M}_{\Theta} \rightarrow \mathbf{M}_{\text{rel}}^R$ induced by S_1 and S_2 are homotopic.*

Proof of Proposition 14.4. . The extended Eilenberg-Zilber map

$$W \otimes S_*(Y \times Z) \rightarrow S_*(Y) \otimes S_*(Z)$$

([FM, proof of Proposition 5.8]) gives a map

$$S_*(Y \times Z) \rightarrow ((S_*(Y) \otimes S_*(Z))^W,$$

and this gives a natural quasi-isomorphism

$$\nu : \text{sig}_{\text{rel}} \rightarrow \delta \circ \text{sig}.$$

□

The rest of this section is devoted to the proof of Proposition 14.5. The basic idea is similar to the proof of Theorem 1.1.

Definition 14.6. Let P be the poset whose two elements are the functors S_1 and S_2 , with $S_1 \leq S_2$.

Recall Definition 6.2(i), and note that $U(K)$ has a poset structure given by inclusions of cells. Our next definition is analogous to Definition 6.2(ii).

Definition 14.7. Let $k \geq 0$, let \mathbf{n} be a k -fold multi-index, and let $F \in \text{pre}_{\pi, Z, w}^k(\Delta^{\mathbf{n}})$. Let

$$b : U(\Delta^{\mathbf{n}}) \rightarrow P$$

be a map of posets.

(i) For an object (σ, o) of $\text{Cell}(\Delta^{\mathbf{n}})$ define the object $b_*(F)(\sigma, o)$ of $\mathcal{A}_{\text{rel}}^R$ to be $b(\sigma)(F(\sigma, o))$.

(ii) For a morphism $f : (\sigma, o) \rightarrow (\sigma', o')$ of $\text{Cell}(\Delta^{\mathbf{n}})$ define the morphism

$$b_*(F)(f) : b_*(F)(\sigma, o) \rightarrow b_*(F)(\sigma', o')$$

to be

$$\begin{cases} S_1(f) & \text{if } b(\sigma', o') = S_1, \\ S_2(f) & \text{if } b(\sigma, o) = S_2, \\ \nu \circ S_1(f) & \text{otherwise.} \end{cases}$$

(iii) Let

$$b_* : \text{pre}_{\pi, Z, w}^k(\Delta^{\mathbf{n}}) \rightarrow (\text{pre}_{\text{rel}}^R)^k(\Delta^{\mathbf{n}})$$

be the map defined by (i) and (ii).

Lemma 14.8. b_* takes ads to ads.

Proof. This follows from the fact that ν is a quasi-isomorphism. \square

Recall that we write \mathbf{R} for the object of $\Sigma\mathcal{S}_{\text{ss}}$ associated to an ad theory (Example 4.13). Then b_* gives a map

$$((\mathbf{R}_{\Theta})_k)_{\mathbf{n}} \rightarrow ((\mathbf{R}_{\text{rel}}^R)_k)_{\mathbf{n}}.$$

Definition 14.9. (i) For $k \geq 0$ define an object \mathcal{P}_k of $\Sigma_k\text{ss}_k$ by

$$(\mathcal{P}_k)_{\mathbf{n}} = \text{Map}_{\text{posets}}(U(\Delta^{\mathbf{n}}), P)_+$$

(where the $+$ denotes a disjoint basepoint); the morphisms in $(\Delta_{\text{inj}}^{\text{op}})^{\times k}$ act in the evident way, and the morphisms of the form (α, id) with $\alpha \in \Sigma_k$ act by permuting the factors in $\Delta^{\mathbf{n}}$.

(ii) Define \mathcal{P} to be the object of \mathfrak{B} with k -th term \mathcal{P}_k .

Next we give $\mathcal{P} \overline{\wedge} \mathbf{R}_{\Theta}$ the structure of a multiseemisimplicial symmetric spectrum (cf. Definition 8.1). Recall Definition 6.4 and let s be the 1-simplex of S^1 . Define

$$\omega : S^1 \wedge (\mathcal{P} \overline{\wedge} \mathbf{R}_{\Theta})_k \rightarrow (\mathcal{P} \overline{\wedge} \mathbf{R}_{\Theta})$$

as follows: for $b \in (\mathcal{P}_k)_{\mathbf{n}}$ and $x \in ((\mathbf{R}_{\Theta})_k)_{\mathbf{n}}$, let

$$\omega(s \wedge (b \wedge x)) = (b \circ \Pi) \wedge \omega(s \wedge x).$$

It follows from the definitions that we obtain a map

$$\beta : \mathcal{P} \overline{\wedge} \mathbf{R}_{\Theta} \rightarrow \mathbf{R}_{\text{rel}}^R$$

in $\Sigma\mathcal{S}_{\text{ss}}$ by

$$\beta(b \wedge F) = b_*(F).$$

For each k and \mathbf{n} , define elements $c_{k,\mathbf{n}}, d_{k,\mathbf{n}} \in (\mathcal{P}_k)_{\mathbf{n}}$ to be the constant functions $U(\Delta^{\mathbf{n}}) \rightarrow P$ whose values are respectively S_1 and S_2 . Then define maps

$$c, d : \overline{\mathbf{S}} \rightarrow \mathcal{P}$$

in \mathfrak{B} by taking the nontrivial simplex of $(\overline{\mathbf{S}}_k)_{\mathbf{n}}$ to $c_{k,\mathbf{n}}$, resp., $d_{k,\mathbf{n}}$. Finally, define maps

$$\mathbf{c}, \mathbf{d} : \mathbf{R}_{\Theta} \rightarrow \mathcal{P} \overline{\wedge} \mathbf{R}_{\Theta}$$

in $\Sigma\mathcal{S}_{\text{ss}}$ by letting \mathbf{c} be the composite

$$\mathbf{R}_{\Theta} \cong \overline{\mathbf{S}} \overline{\wedge} \mathbf{R}_{\Theta} \xrightarrow{c \wedge 1} \mathcal{P} \overline{\wedge} \mathbf{R}_{\Theta}$$

and similarly for \mathbf{d} .

Now $\beta \circ \mathbf{c}$ is the map S_1 and $\beta \circ \mathbf{d}$ is the map S_2 , so to complete the proof of Proposition 14.5 it suffices to show:

Lemma 14.10. *\mathbf{c} and \mathbf{d} are homotopic in $\Sigma\mathcal{S}_{\text{ss}}$.*

Proof of Lemma 14.10. For each $k \geq 0$ and each \mathbf{n} let $e_{k,\mathbf{n}} : (\mathcal{P}_k)_{\mathbf{n}} \rightarrow S^0$ be the map which takes every simplex except the basepoint to the nontrivial element of S^0 , and let

$$e : \mathcal{P} \rightarrow \overline{\mathbf{S}}$$

be the map given by the $e_{k,\mathbf{n}}$. Let

$$\mathbf{e} : \mathcal{P} \overline{\wedge} \mathbf{R}_{\Theta} \rightarrow \mathbf{R}_{\Theta}$$

be the composite

$$\mathcal{P} \overline{\wedge} \mathbf{R}_{\Theta} \xrightarrow{e \wedge 1} \overline{\mathbf{S}} \overline{\wedge} \mathbf{R}_{\Theta} \cong \mathbf{R}_{\Theta}.$$

Then $\mathbf{e} \circ \mathbf{c}$ and $\mathbf{e} \circ \mathbf{d}$ are both equal to the identity. But \mathbf{e} is a weak equivalence by Proposition 9.3 and Lemma 10.3, and the result follows. \square

15. BACKGROUND FOR THE PROOF OF THEOREM 1.3

Notation 15.1. In order to distinguish the product in $\mathcal{A}_{e,*,1}$ from the Cartesian product of categories, we will denote the former by \boxtimes from now on.

We now turn to the proof of Theorem 1.3, which will follow the general outline of the proof of Theorem 1.1. The key ingredient in that proof was the action of the monad \mathbb{P} on \mathbf{R} . That action was constructed from the family of operations given in Definition 6.2(ii), and this family in turn was constructed from the family of functors η_\star given in Definition 3.5(ii). For our present purpose we need the functors η_\star and also a family of functors

$$\mathbf{d}_\blacksquare : \mathcal{A}_1 \times \cdots \times \mathcal{A}_j \rightarrow \mathcal{A}_{\text{rel}}^{\mathbb{Z}},$$

where each \mathcal{A}_i is equal to $\mathcal{A}_{e,*,1}$ or $\mathcal{A}_{\text{rel}}^{\mathbb{Z}}$; these will be built from the symmetric monoidal structures of $\mathcal{A}_{e,*,1}$ and $\mathcal{A}_{\text{rel}}^{\mathbb{Z}}$ and the functor

$$\text{sig}_{\text{rel}} : \mathcal{A}_{e,*,1} \rightarrow \mathcal{A}_{\text{rel}}^{\mathbb{Z}}.$$

It is convenient to represent this situation by a function r from $\{1, \dots, j\}$ to a two element set $\{u, v\}$, with $\mathcal{A}_i = \mathcal{A}_{e,*,1}$ if $r(i) = u$ and $\mathcal{A}_i = \mathcal{A}_{\text{rel}}^{\mathbb{Z}}$ if $r(i) = v$.

Example 15.2. A typical example is the functor

$$\mathcal{A}_{\text{rel}}^{\mathbb{Z}} \times (\mathcal{A}_{e,*,1})^{\times 5} \times \mathcal{A}_{\text{rel}}^{\mathbb{Z}} \rightarrow \mathcal{A}_{\text{rel}}^{\mathbb{Z}}$$

which takes (x_1, \dots, x_7) to

$$i^\epsilon \text{sig}_{\text{rel}}(x_4 \boxtimes x_3) \otimes x_7 \otimes \text{sig}_{\text{rel}}(x_6 \boxtimes x_2 \boxtimes x_5) \otimes x_1,$$

where i^ϵ is the sign that arises from permuting (x_1, \dots, x_7) into the order $(x_4, x_3, x_7, x_6, x_2, x_5, x_1)$. In Definition 15.4(iv) we will represent such a functor by a surjection h which keeps track of which inputs go to which output factors and a permutation η which keeps track of the order in which the inputs to each sig_{rel} factor are multiplied. In the present example h is the surjection $\{1, \dots, 7\} \rightarrow \{1, 2, 3, 4\}$ with

$$h^{-1}(1) = \{3, 4\}, h^{-1}(2) = 7, h^{-1}(3) = \{2, 5, 6\}, h^{-1}(4) = 1$$

and η is the permutation $(256)(34)$.

In order to get the signs right we need a preliminary definition.

Definition 15.3. (i) For totally ordered sets S_1, \dots, S_m , define

$$\prod_{i=1}^n S_i$$

to be the disjoint union with the order relation given as follows: $s < t$ if either $s \in S_i$ and $t \in S_j$ with $i < j$, or $s, t \in S_i$ with $s < t$ in the order of S_i .

(ii) For a surjection

$$h : \{1, \dots, j\} \rightarrow \{1, \dots, m\}$$

define $\theta(h)$ to be the permutation

$$\{1, \dots, j\} \cong h^{-1}(1) \amalg \cdots \amalg h^{-1}(m) \cong \{1, \dots, j\};$$

here the first map restricts to the identity on each $h^{-1}(i)$ and the second is the unique ordered bijection.

In Example 15.2. $\theta(h)$ takes $1, \dots, 7$ respectively to $7, 4, 1, 2, 5, 6, 3$.

Definition 15.4. Let $j \geq 0$ and let $r : \{1, \dots, j\} \rightarrow \{u, v\}$. Let \mathcal{A}_i denote $\mathcal{A}_{e,*,1}$ if $r(i) = u$ and $\mathcal{A}_{\text{rel}}^{\mathbb{Z}}$ if $r(i) = v$.

(i) Let $1 \leq m \leq j$. A surjection

$$h : \{1, \dots, j\} \rightarrow \{1, \dots, m\}$$

is *adapted* to r if r is constant on each set $h^{-1}(i)$ and h is monic on $r^{-1}(v)$.

(ii) Given a surjection

$$h : \{1, \dots, j\} \rightarrow \{1, \dots, m\}$$

which is adapted to r , define

$$h_{\blacklozenge} : \mathcal{A}_1 \times \cdots \times \mathcal{A}_j \rightarrow (\mathcal{A}_{\text{rel}}^{\mathbb{Z}})^{\times m}$$

by

$$h_{\blacklozenge}(x_1, \dots, x_j) = (i^\epsilon y_1, \dots, y_m),$$

where i^ϵ is the sign that arises from putting the objects x_1, \dots, x_j into the order $x_{\theta(h)^{-1}(1)}, \dots, x_{\theta(h)^{-1}(j)}$ and

$$y_i = \begin{cases} \text{sig}_{\text{rel}}(\boxtimes_{l \in h^{-1}(i)} x_l) & \text{if } h^{-1}(i) \subset r^{-1}(u), \\ x_{h^{-1}(i)} & \text{if } h^{-1}(i) \in r^{-1}(v). \end{cases}$$

(iii) A *datum* of type r is a pair

$$(h, \eta),$$

where h is a surjection which is adapted to r and η is an element of Σ_j with the property that $h \circ \eta = h$.

(iv) Given a datum

$$\mathbf{d} = (h, \eta),$$

of type r , define

$$\mathbf{d}_{\blacksquare} : \mathcal{A}_1 \times \cdots \times \mathcal{A}_j \rightarrow \mathcal{A}_{\text{rel}}^{\mathbb{Z}}$$

to be the composite

$$\mathcal{A}_1 \times \cdots \times \mathcal{A}_j \xrightarrow{\eta} \mathcal{A}_{\eta^{-1}(1)} \times \cdots \times \mathcal{A}_{\eta^{-1}(j)} = \mathcal{A}_1 \times \cdots \times \mathcal{A}_j \xrightarrow{h_\diamond} (\mathcal{A}_{\text{rel}}^{\mathbb{Z}})^{\times m} \xrightarrow{\otimes} \mathcal{A}_{\text{rel}}^{\mathbb{Z}},$$

where η permutes the factors with the usual sign.

We also need natural transformations between the functors \mathbf{d}_\blacksquare . First observe that there is a natural transformation from the functor

$$(\mathcal{A}_{e,*,1})^{\times l} \xrightarrow{\text{sig}_{\text{rel}}^{\times l}} (\mathcal{A}_{\text{rel}}^{\mathbb{Z}})^{\times l} \xrightarrow{\otimes} \mathcal{A}_{\text{rel}}^{\mathbb{Z}}$$

to the functor

$$(\mathcal{A}_{e,*,1})^{\times l} \xrightarrow{\boxtimes} \mathcal{A}_{e,*,1} \xrightarrow{\text{sig}_{\text{rel}}} \mathcal{A}_{\text{rel}}^{\mathbb{Z}}$$

given by the maps

$$S_*(X_1) \otimes \cdots \otimes S_*(X_l) \xrightarrow{\otimes} S_*(X_1 \times \cdots \times X_l)$$

and

$$\begin{aligned} S_*(X_1 \times X_1) \otimes \cdots \otimes S_*(X_l \times X_l) &\xrightarrow{\otimes} S_*(X_1 \times X_1 \times \cdots \times X_l \times X_l) \\ &\cong S_*((X_1 \times \cdots \times X_l) \times (X_1 \times \cdots \times X_l)). \end{aligned}$$

Combining this with the symmetric monoidal structures of $\mathcal{A}_{e,*,1}$ and $\mathcal{A}_{\text{rel}}^{\mathbb{Z}}$, we obtain a natural transformation $\mathbf{d}_\blacksquare \rightarrow \mathbf{d}'_\blacksquare$ whenever $\mathbf{d} \leq \mathbf{d}'$, as defined in:

Definition 15.5. For data of type r , define

$$(h, \eta) \leq (h', \eta')$$

if each set $\eta^{-1}(h^{-1}(i))$ is contained in some set $\eta'^{-1}(h'^{-1}(l))$.

Our next definition is analogous to Definition 6.6 (the presence of the letter v in the symbols $P_{r;v}$ and $\mathcal{O}(r;v)$ will be explained in a moment).

Definition 15.6. Let $r : \{1 \dots, j\} \rightarrow \{u, v\}$.

(i) Let $P_{r;v}$ be the preorder whose elements are the data of type r , with the order relation given by Definition 15.5.

(ii) Define an object $\mathcal{O}(r;v)_k$ of $\Sigma_k \text{SS}_k$ by

$$(\mathcal{O}(r;v)_k)_\mathbf{n} = \text{Map}_{\text{preorder}}(U(\Delta^\mathbf{n}), P_{r;v})_+$$

(where the $+$ denotes a disjoint basepoint); the morphisms in $(\Delta_{\text{inj}}^{\text{op}})^{\times k}$ act in the evident way, and the morphisms of the form (α, id) with $\alpha \in \Sigma_k$ act by permuting the factors in $\Delta^\mathbf{n}$.

(iii) Define $\mathcal{O}(r;v)$ to be the object of β with k -th term $\mathcal{O}(r;v)_k$.

Remark 15.7. (i) Given $r : \{1, \dots, j\} \rightarrow \{u, v\}$, let $m = |r^{-1}(v)| + 1$, let

$$h : \{1, \dots, j\} \rightarrow \{1, \dots, m\}$$

be any surjection which is adapted to r , and let e be the identity element of Σ_j . Then the datum (h, e) is \geq every element in $P_{r;v}$.

(ii) Lemma 10.3 shows that each of the objects $\mathcal{O}(r;v)_k$ has compatible degeneracies and is weakly equivalent to a point.

We also need a preorder corresponding to the family of functors

$$\eta_\star : (\mathcal{A}_{e,*,1})^{\times j} \rightarrow \mathcal{A}_{e,*,1} :$$

Notation 15.8. Let $r_u(j)$ (resp., $r_v(j)$) denote the constant function $\{1, \dots, j\} \rightarrow \{u, v\}$ with value u (resp., v).

Definition 15.9. Let $r : \{1, \dots, j\} \rightarrow \{u, v\}$.

(i) If $r = r_u(j)$, let $P_{r;u}$ be the set Σ_j with the preorder in which every element is \leq every other, and let $\mathcal{O}(r;u)_k$ be the object $\mathcal{O}(j)_k$ of Definition 6.6.

(ii) Otherwise let $P_{r;u}$ be the empty set and let $\mathcal{O}(r;u)_k$ be the multisemisimplicial set with a point in every multidegree.

(iii) In either case, let $\mathcal{O}(r;u)$ be the object of \mathfrak{B} with k -th term $\mathcal{O}(r;u)_k$.

In the next section we will use the objects $\mathcal{O}(r;v)$ and $\mathcal{O}(r;u)$ to construct a monad. In preparation for that, we show that the collection of preorders $P_{r;v}$ and $P_{r;u}$ has suitable composition maps. Specifically, we show that it is a colored operad (also called a multicategory) in the category of preorders.

We refer the reader to [EM06, Section2] for the definition of multicategory; we will mostly follow the notation and terminology given there. In our case there are two objects u and v , and we think of a function $r : \{1, \dots, j\} \rightarrow \{u, v\}$ as a sequence of u 's and v 's.

Remark 15.10. Let $r : \{1, \dots, j\} \rightarrow \{u, v\}$ and let \mathcal{A}_i denote $\mathcal{A}_{e,*,1}$ if $r(i) = u$ and $\mathcal{A}_{\text{rel}}^{\mathbb{Z}}$ if $r(i) = v$. Let us write \mathcal{A}_r for the category $\mathcal{A}_1 \times \dots \times \mathcal{A}_j$ and $\mathcal{A}_{r;u}$ (resp., $\mathcal{A}_{r;v}$) for the category of functors $\mathcal{A}_r \rightarrow \mathcal{A}_{e,*,1}$ (resp., $\mathcal{A}_r \rightarrow \mathcal{A}_{\text{rel}}^{\mathbb{Z}}$). Then the objects $\mathcal{A}_{r;u}$ and $\mathcal{A}_{r;v}$ form a multicategory (with the obvious Σ_j actions and composition maps). Moreover, Definitions 3.5(ii) and 15.4(iv) give imbeddings

$$\Phi_{r;u} : P_{r;u} \rightarrow \mathcal{A}_{r;u}$$

and

$$\Phi_{r;v} : P_{r;v} \rightarrow \mathcal{A}_{r;v}.$$

The following definitions are chosen so that these imbeddings preserve the Σ_j actions and the composition operations.

We define the right Σ_j action on the collection of j -morphisms as follows. Let $\alpha \in \Sigma_j$ and $r : \{1, \dots, j\} \rightarrow \{u, v\}$. Define r^α to be the composite

$$\{1, \dots, j\} \xrightarrow{\alpha} \{1, \dots, j\} \xrightarrow{r} \{u, v\}.$$

If $r = r_u(j)$ then $r^\alpha = r$ and the map

$$\alpha : P_{r;u} \rightarrow P_{r^\alpha;u}$$

is the right action of Σ_j on itself. If $(h, \eta) \in P_{r,v}$ define $(h, \eta)\alpha \in P_{r^\alpha;v}$ to be $(h \circ \bar{\alpha}, \bar{\alpha}^{-1} \circ \eta \circ \alpha)$, where $\bar{\alpha} \in \Sigma_j$ is the permutation whose restriction to each $\alpha^{-1}h^{-1}(i)$ is the order-preserving bijection to $h^{-1}(i)$.

We define the composition operation as follows. If the composition involves only $P_{r;u}$'s then it is the composition in the operad \mathcal{M} of [May72, Definition 3.1(i)]. Otherwise let $i, j_1, \dots, j_i \geq 0$, let $r : \{1, \dots, i\} \rightarrow \{u, v\}$, and for $1 \leq l \leq i$ let $r_l : \{1, \dots, j_l\} \rightarrow \{u, v\}$; assume that if $r(l) = u$ then r_l is $r_u(j_l)$. Let $(h, \eta) \in P_{r;v}$ and for $1 \leq l \leq i$ let $x_l \in P_{r_l;r(l)}$. If $r(l) = v$ then x_l has the form (h_l, η_l) , otherwise x_l is an element $\eta_l \in \Sigma_{j_l}$ and we write h_l for the map $\{1, \dots, j_l\} \rightarrow \{1\}$. Define the composition operation Γ by

$$(15.1) \quad \Gamma((h, \eta), x_1, \dots, x_i) = (H, \theta),$$

where H is the multivariable composite $h \circ (h_1, \dots, h_i)$ and θ is the composite $\gamma_{\mathcal{M}}(\eta, \eta_1, \dots, \eta_i)$ in the operad \mathcal{M} of [May72, Definition 3.1(i)].

Proposition 15.11. *With these definitions, the collection of preorders $P_{r;u}$ and $P_{r;v}$ is a multicategory.*

Proof. This is immediate from Remark 15.10. □

16. A MONAD IN $\mathfrak{B} \times \mathfrak{B}$

In this section we construct a monad in $\mathfrak{B} \times \mathfrak{B}$ which acts on the pair $(\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^{\mathbb{Z}})$.

Definition 16.1. Let $j \geq 0$ and let $\mathbf{X}, \mathbf{Y} \in \mathfrak{B}$.

(i) For $\alpha \in \Sigma_j$ and $r : \{1, \dots, j\} \rightarrow \{u, v\}$, define

$$\tilde{\alpha} : \mathcal{O}(r; v) \rightarrow \mathcal{O}(r^\alpha; v)$$

by

$$(\tilde{\alpha}(a))(\sigma) = (a(\sigma))\alpha$$

where $a \in \text{Map}_{\text{preorder}}(U(\Delta^n), P_{r;v})_+$ and $\sigma \in U(\Delta^n)$.

(ii) Define

$$(\mathbf{X}, \mathbf{Y})^{\otimes r} = \mathbf{Z}_1 \otimes \dots \otimes \mathbf{Z}_j,$$

where \mathbf{Z}_i denotes \mathbf{X} if $r(i) = u$ and \mathbf{Y} if $r(i) = v$.

(iii) For $\alpha \in \Sigma_j$ define

$$\bar{\alpha} : \bigvee_r \mathcal{O}(r; v) \bar{\wedge} (\mathbf{X}, \mathbf{Y})^{\otimes r} \rightarrow \bigvee_r \mathcal{O}(r; v) \bar{\wedge} (\mathbf{X}, \mathbf{Y})^{\otimes r}$$

to be the map which takes the r -summand to the r^α -summand by means of the map

$$\mathcal{O}(r; v) \bar{\wedge} (\mathbf{X}, \mathbf{Y})^{\otimes r} \xrightarrow{\bar{\alpha}\bar{\wedge}\alpha} \mathcal{O}(r^\alpha; v) \bar{\wedge} (\mathbf{X}, \mathbf{Y})^{\otimes r^\alpha}.$$

Note that the maps $\bar{\alpha}$ give $\bigvee_r \mathcal{O}(r; v) \bar{\wedge} (\mathbf{X}, \mathbf{Y})^{\otimes r}$ a right Σ_j action. Recall Notation 15.8.

Definition 16.2. (i) Define a functor $\mathbb{O} : \mathfrak{B} \times \mathfrak{B} \rightarrow \mathfrak{B} \times \mathfrak{B}$ by

$$\mathbb{O}(\mathbf{X}, \mathbf{Y}) = (\mathbb{O}_1(\mathbf{X}), \mathbb{O}_2(\mathbf{X}, \mathbf{Y})),$$

where

$$\mathbb{O}_1(\mathbf{X}) = \bigvee_{j \geq 0} (\mathcal{O}(r_u(j); u) \bar{\wedge} \mathbf{X}^{\otimes j}) / \Sigma_j$$

and

$$\mathbb{O}_2(\mathbf{X}, \mathbf{Y}) = \bigvee_{j \geq 0} \bigvee_r (\mathcal{O}(r; v) \bar{\wedge} (\mathbf{X}, \mathbf{Y})^{\otimes r}) / \Sigma_j.$$

(ii) Define a natural transformation

$$\iota : (\mathbf{X}, \mathbf{Y}) \rightarrow \mathbb{O}(\mathbf{X}, \mathbf{Y})$$

to be (ι_1, ι_2) , where ι_1 is the composite

$$\mathbf{X} \xrightarrow{\cong} \bar{\mathbf{S}} \bar{\wedge} \mathbf{X} = \mathcal{O}(r_u(1); u) \bar{\wedge} \mathbf{X} \hookrightarrow \mathbb{O}(\mathbf{X})$$

and ι_2 is the composite

$$\mathbf{Y} \xrightarrow{\cong} \bar{\mathbf{S}} \bar{\wedge} \mathbf{Y} = \mathcal{O}(r_v(1); v) \bar{\wedge} \mathbf{Y} \hookrightarrow \mathbb{O}(\mathbf{Y}).$$

For the structure map $\mu : \mathbb{O}\mathbb{O} \rightarrow \mathbb{O}$ we need a composition operation for the collection of objects $\mathcal{O}(r; u)$ and $\mathcal{O}(r; v)$. Recall Definition 15.3(i) and the map Γ defined in Equation (15.1).

Definition 16.3. Let $i, j_1, \dots, j_i \geq 0$, let $r : \{1, \dots, i\} \rightarrow \{u, v\}$, and for $1 \leq l \leq i$ let $r_l : \{1, \dots, j_l\} \rightarrow \{u, v\}$; assume that if $r(l) = u$ then r_l is $r_u(j_l)$. Let

$$R : \{1, \dots, \sum j_l\} \rightarrow \{u, v\}$$

be the composite

$$\{1, \dots, \sum j_l\} \cong \prod_{l=1}^i \{1, \dots, j_l\} \rightarrow \{u, v\},$$

where the first map is the unique order-preserving bijection and the second restricts on each $\{1, \dots, j_l\}$ to r_l . Define a map

$$\gamma : \mathcal{O}(r; v) \overline{\wedge} (\mathcal{O}(r_1; r(1)) \otimes \cdots \otimes \mathcal{O}(r_i; r(i))) \rightarrow \mathcal{O}(R; v)$$

in \mathfrak{B} by the formulas

$$\gamma(a \wedge [e, b_1 \wedge \dots \wedge b_i])(\sigma_1 \times \cdots \times \sigma_i) = \Gamma(a(\sigma_1 \times \cdots \times \sigma_i), b_1(\sigma_1), \dots, b_i(\sigma_i))$$

(where e is the identity element of the relevant symmetric group) and

$$\gamma(a \wedge [\alpha, b_1 \wedge \dots \wedge b_i]) = (\alpha, \text{id})_* \gamma((\alpha^{-1}, \text{id})_* a \wedge [e, b_1 \wedge \dots \wedge b_i]).$$

This operation satisfies the analogues of Lemmas 7.2, 7.5, and 7.7.

Now we can define

$$\mu : \mathbb{O} \mathbb{O} \rightarrow \mathbb{O}$$

to be (μ_1, μ_2) , where μ_1 is given by Definition 7.8(iv) and μ_2 is defined in a similar way using Definition 16.3.

Proposition 16.4. *The transformations μ and ι define a monad structure on \mathbb{O} . \square*

We conclude this section by giving the action of \mathbb{O} on the pair $(\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^{\mathbb{Z}})$. Recall Remark 15.10.

Definition 16.5. Let k_1, \dots, k_j be non-negative integers and let \mathbf{n}_i be a k_i -fold multi-index for $1 \leq i \leq j$. Let $r : \{1, \dots, j\} \rightarrow \{u, v\}$, and for $1 \leq i \leq j$ let pre_i denote $\text{pre}_{e,*,1}$ if $r(i) = u$ and $\text{pre}_{\text{rel}}^{\mathbb{Z}}$ if $r(i) = v$. For any map of preorders

$$a : U(\Delta^{\mathbf{n}_1} \times \cdots \times \Delta^{\mathbf{n}_j}) \rightarrow P_{r;v}$$

define

$$a_* : \text{pre}_1^{k_1}(\Delta^{\mathbf{n}_1}) \times \cdots \times \text{pre}_j^{k_j}(\Delta^{\mathbf{n}_j}) \rightarrow (\text{pre}_{\text{rel}}^{\mathbb{Z}})^{k_1 + \cdots + k_j}(\Delta^{(\mathbf{n}_1, \dots, \mathbf{n}_j)})$$

by

$$\begin{aligned} a_*(F_1, \dots, F_j)(\sigma_1 \times \cdots \times \sigma_j, o_1 \times \cdots \times o_j) \\ = i^{\epsilon(\zeta)} \Phi_{r;v}(a(\sigma_1 \times \cdots \times \sigma_j))(F_1(\sigma_1, o_1), \dots, F_j(\sigma_j, o_j)), \end{aligned}$$

where ζ is the block permutation that takes blocks $\mathbf{b}_1, \dots, \mathbf{b}_j, \mathbf{c}_1, \dots, \mathbf{c}_j$ of size $k_1, \dots, k_j, \dim \sigma_1, \dots, \dim \sigma_j$ into the order $\mathbf{b}_1, \mathbf{c}_1, \dots, \mathbf{b}_j, \mathbf{c}_j$.

Lemma 16.6. *If $F_i \in \text{ad}_i^{k_i}(\Delta^{\mathbf{n}_i})$ for $1 \leq i \leq j$ then $a_*(F_1, \dots, F_j) \in (\text{ad}_{\text{rel}}^{\mathbb{Z}})^{k_1 + \cdots + k_j}(\Delta^{(\mathbf{n}_1, \dots, \mathbf{n}_j)})$.*

Proof. This is a straightforward consequence of the fact that the natural transformation from the functor

$$(\mathcal{A}_{e,*,1})^{\times l} \xrightarrow{\text{sig}_{\text{rel}}^{\times l}} (\mathcal{A}_{\text{rel}}^{\mathbb{Z}})^{\times l} \xrightarrow{\otimes} \mathcal{A}_{\text{rel}}^{\mathbb{Z}}$$

to the functor

$$(\mathcal{A}_{e,*,1})^{\times l} \xrightarrow{\boxtimes} \mathcal{A}_{e,*,1} \xrightarrow{\text{sig}_{\text{rel}}} \mathcal{A}_{\text{rel}}^{\mathbb{Z}}$$

given by the cross product is a quasi-isomorphism. \square

Definition 16.7. Let $j \geq 0$ and let $r : \{1, \dots, j\} \rightarrow \{u, v\}$. Define a map

$$\phi_r : \mathcal{O}(r; v) \overline{\wedge} (\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^{\mathbb{Z}})^{\otimes r} \rightarrow \mathbf{R}_{\text{rel}}^{\mathbb{Z}}$$

in β by the formulas

$$\phi_r(a \wedge [e, F_1 \wedge \dots \wedge F_j]) = a_*(F_1 \wedge \dots \wedge F_j)$$

(where e denotes the identity element of the relevant symmetric group) and

$$\phi_r(a \wedge [\alpha, F_1 \wedge \dots \wedge F_j]) = (\alpha, \text{id})_* \phi_r((\alpha^{-1}, \text{id})_* a \wedge [e, F_1 \wedge \dots \wedge F_j]).$$

Next observe that the maps ϕ_r induce a map

$$(16.1) \quad \left(\bigvee_r (\mathcal{O}(r; v) \overline{\wedge} (\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^{\mathbb{Z}})^{\otimes r}) \right) / \Sigma_j \rightarrow \mathbf{R}_{\text{rel}}^{\mathbb{Z}}$$

for each $j \geq 0$. We define

$$\nu : \mathbb{O}(\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^{\mathbb{Z}}) \rightarrow (\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^{\mathbb{Z}})$$

to be the pair (ν_1, ν_2) , where ν_1 is given by Definition 7.10 and ν_2 is given by the maps (16.1).

Proposition 16.8. ν is an action of \mathbb{O} on $(\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^{\mathbb{Z}})$.

Proof. This is a straightforward consequence of Remark 15.10. \square

17. A MONAD IN $\Sigma \mathcal{S}_{\text{ss}} \times \Sigma \mathcal{S}_{\text{ss}}$

First we give $\mathcal{O}(r; v) \overline{\wedge} (\mathbf{X}, \mathbf{Y})^{\wedge r}$ the structure of a multiseemisimplicial symmetric spectrum when $\mathbf{X}, \mathbf{Y} \in \beta$. The definition is analogous to Definition 8.1. Recall Definition 6.4.

Definition 17.1. Let $j, k \geq 0$. Let s be the 1-simplex of S^1 . Define

$$\omega : S^1 \wedge (\mathcal{O}(r; v) \overline{\wedge} (\mathbf{X}, \mathbf{Y})^{\wedge r})_k \rightarrow (\mathcal{O}(r; v) \overline{\wedge} (\mathbf{X}, \mathbf{Y})^{\wedge r})_{k+1}$$

as follows: for $a \in (\mathcal{O}(r; v)_k)_{\mathbf{n}}$ and $x \in ((\mathbf{X}, \mathbf{Y})^{\wedge r})_k_{\mathbf{n}}$, let

$$\omega(s \wedge (a \wedge x)) = (a \circ \Pi) \wedge \omega(s \wedge x).$$

Definition 17.2. Define a functor $\mathbb{P} : \mathfrak{B} \times \mathfrak{B} \rightarrow \mathfrak{B} \times \mathfrak{B}$ by

$$\mathbb{P}(\mathbf{X}, \mathbf{Y}) = (\mathbb{P}_1(\mathbf{X}), \mathbb{P}_2(\mathbf{X}, \mathbf{Y})),$$

where

$$\mathbb{P}_1(\mathbf{X}) = \bigvee_{j \geq 0} (\mathcal{O}(r_u(j); u) \overline{\wedge} \mathbf{X}^{\wedge j}) / \Sigma_j$$

and

$$\mathbb{P}_2(\mathbf{X}, \mathbf{Y}) = \bigvee_{j \geq 0} \left(\bigvee_r (\mathcal{O}(r; v) \overline{\wedge} (\mathbf{X}, \mathbf{Y})^{\wedge r}) / \Sigma_j \right).$$

The proof that \mathbb{P} inherits a monad structure and an action on $(\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^{\mathbb{Z}})$ is the same as the corresponding proof in Section 8.

For use in the next section we record a lemma. Let \mathcal{C} be the category whose objects are triples $(\mathbf{X}, \mathbf{Y}, f)$, where \mathbf{X} and \mathbf{Y} are monoids in $\Sigma \mathcal{S}_{\text{ss}}$ and f is a map $\mathbf{X} \rightarrow \mathbf{Y}$ in $\Sigma \mathcal{S}_{\text{ss}}$ which is not required to be a monoid map; the morphisms are commutative diagrams

$$\begin{array}{ccc} \mathbf{X} & \longrightarrow & \mathbf{X}' \\ f \downarrow & & \downarrow f' \\ \mathbf{Y} & \longrightarrow & \mathbf{Y}', \end{array}$$

where the horizontal arrows are monoid maps.

Lemma 17.3. (i) *There is a functor Υ from \mathbb{P} algebras to \mathcal{C} which takes (\mathbf{X}, \mathbf{Y}) to a map $\mathbf{X} \rightarrow \mathbf{Y}$; in particular, \mathbf{X} and \mathbf{Y} have natural monoid structures.*

(ii) $\Upsilon(\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^{\mathbb{Z}})$ *is the map*

$$\text{sig}_{\text{rel}} : \mathbf{R}_{e,*,1} \rightarrow \mathbf{R}_{\text{rel}}^{\mathbb{Z}},$$

where $\mathbf{R}_{e,*,1}$ and $\mathbf{R}_{\text{rel}}^{\mathbb{Z}}$ have the monoid structures given by Lemma 8.8(i).

Proof. Recall Notation 15.8, and let e denote the identity element of Σ_j .

Part (i). The map $f : \mathbf{X} \rightarrow \mathbf{Y}$ is the composite

$$\mathbf{X} \cong \overline{\mathbf{S}} \overline{\wedge} \mathbf{X} = \mathcal{O}(r_u(1); v) \overline{\wedge} \mathbf{X} \hookrightarrow \mathbb{P}_2(\mathbf{X}, \mathbf{Y}) \rightarrow \mathbf{Y}.$$

The monoid structure on \mathbf{X} is given by Lemma 8.8; it remains to give the monoid structure on \mathbf{Y} . It suffices to give an action on \mathbf{Y} of the monad \mathbb{A} defined in the proof of Lemma 8.8, and for this in turn it suffices to give a suitable natural transformation $\mathbb{A} \rightarrow \mathbb{P}_2$.

For each $j \geq 0$ let h_0 be the identity map of $\{1, \dots, j\}$. Then h_0 is adapted to $r_v(j)$, so we obtain an element $(h_0, e) \in P_{r_v(j);v}$.

For each $j, k \geq 0$ and each k -fold multi-index \mathbf{n} , define an element

$$b_{j,k,\mathbf{n}} \in (\mathcal{O}(r_v(j); v)_k)_{\mathbf{n}}$$

to be the constant function $U(\Delta^{\mathbf{n}}) \rightarrow P_{r_v(j);v}$ whose value is (h_0, e) . Next define a map

$$\bar{\mathbf{S}} \rightarrow \mathcal{O}(r_v(j); v)$$

by taking the nontrivial simplex of $(\bar{\mathbf{S}}_k)_{\mathbf{n}}$ to $b_{j,k,\mathbf{n}}$.

Now the composite

$$\begin{aligned} \mathbb{A}(\mathbf{Y}) &= \bigvee_{j \geq 0} \mathbf{Y}^{\wedge j} \cong \bigvee_{j \geq 0} \bar{\mathbf{S}} \bar{\wedge} \mathbf{Y}^{\wedge j} \\ &\rightarrow \bigvee_{j \geq 0} \left(\bigvee_r (\mathcal{O}(r; v) \bar{\wedge} (\mathbf{X}, \mathbf{Y})^{\wedge r}) / \Sigma_j = \mathbb{P}_2(\mathbf{X}, \mathbf{Y}) \right) \end{aligned}$$

is the desired map.

Part (ii) is an easy consequence of the definitions. \square

18. RECTIFICATION

In this section we prove Theorem 1.3. The argument is analogous to that in Section 10.

First we consider a monad in $\Sigma \mathcal{S}_{\text{ss}} \times \Sigma \mathcal{S}_{\text{ss}}$ which is simpler than \mathbb{P} .

Definition 18.1. (i) Define $\mathbb{P}'(\mathbf{X}, \mathbf{Y})$ to be

$$\left(\bigvee_{j \geq 0} \mathbf{X}^{\wedge j} / \Sigma_j, \bigvee_{j \geq 0} \left(\bigvee_r (\mathbf{X}, \mathbf{Y})^{\wedge r} \right) / \Sigma_j \right).$$

(ii) For each $j \geq 0$ and each $r : \{1, \dots, j\} \rightarrow \{u, v\}$, let

$$\xi_j : \mathcal{O}(r_u(j); u) \rightarrow \bar{\mathbf{S}}$$

and

$$\zeta_r : \mathcal{O}(r; v) \rightarrow \bar{\mathbf{S}}$$

be the maps which take each nontrivial simplex of the k -th object to the nontrivial simplex of $\bar{\mathbf{S}}_k$ in the same multidegree. Define a natural transformation

$$\Xi : \mathbb{P} \rightarrow \mathbb{P}'$$

to be the pair (Ξ_1, Ξ_2) , where Ξ_1 is the wedge of the composites

$$(\mathcal{O}(r_u(j); u) \bar{\wedge} \mathbf{X}^{\wedge j}) / \Sigma_j \xrightarrow{\xi_j \bar{\wedge} 1} (\bar{\mathbf{S}} \bar{\wedge} \mathbf{X}^{\wedge j}) / \Sigma_j \xrightarrow{\cong} \mathbf{X}^{\wedge j} / \Sigma_j.$$

and Ξ_2 is the wedge of the composites

$$\begin{aligned} \left(\bigvee_r \mathcal{O}(r; v) \overline{\wedge} (\mathbf{X}, \mathbf{Y})^{\wedge r} \right) / \Sigma_j &\xrightarrow{\vee \zeta_r \overline{\wedge} 1} \left(\bigvee_r \overline{\mathbf{S}} \overline{\wedge} (\mathbf{X}, \mathbf{Y})^{\wedge r} \right) / \Sigma_j \\ &\xrightarrow{\cong} \left(\bigvee_r (\mathbf{X}, \mathbf{Y})^{\wedge r} \right) / \Sigma_j. \end{aligned}$$

Proposition 18.2. (i) *An algebra over \mathbb{P}' is the same thing as a pair of commutative monoids (\mathbf{X}, \mathbf{Y}) in $\Sigma \mathcal{S}_{\text{ss}}$ together with a monoid map $\mathbf{X} \rightarrow \mathbf{Y}$.*

(ii) *Ξ is a map of monads.*

(iii) *Suppose that each X_k and each Y_k has compatible degeneracies (see Definition 9.1). Let \mathbb{P}^q denote the q -th iterate of \mathbb{P} . Then each map*

$$\Xi : \mathbb{P}^q(\mathbf{X}, \mathbf{Y}) \rightarrow \mathbb{P}'\mathbb{P}^{q-1}(\mathbf{X}, \mathbf{Y})$$

is a weak equivalence.

Proof. Part (i). Let (\mathbf{X}, \mathbf{Y}) be an algebra over \mathbb{P}' . The fact that \mathbf{X} and \mathbf{Y} are commutative monoids is immediate from the definitions. The map $f : \mathbf{X} \rightarrow \mathbf{Y}$ is constructed as in the proof of Lemma 17.3(i).

To show that f is a monoid map, we first observe that there are two inclusions of $\mathbf{X}^{\wedge j}$ into $\mathbb{P}'_2\mathbb{P}'(\mathbf{X}, \mathbf{Y})$. Let i_1 be the composite

$$\mathbf{X}^{\wedge j} \hookrightarrow \mathbb{P}'_1(\mathbf{X}) \hookrightarrow \mathbb{P}'_2\mathbb{P}'(\mathbf{X}, \mathbf{Y}),$$

where the second arrow is the inclusion of the summand indexed by $j = 1, r = r_u(1)$. Let i_2 be the composite

$$\mathbf{X}^{\wedge j} \hookrightarrow \mathbb{P}'_2(\mathbf{X}, \mathbf{Y})^{\wedge j} \hookrightarrow \mathbb{P}'_2\mathbb{P}'(\mathbf{X}, \mathbf{Y}),$$

where the first arrow is the j -fold smash of the inclusion of the $r_u(1)$ summand, and the second arrow is the inclusion of the $r_v(j)$ summand.

Consider the commutative diagram

$$\begin{array}{ccccc} \mathbf{X}^{\wedge j} & \xrightarrow{=} & \mathbf{X}^{\wedge j} & & \\ \downarrow & \searrow^{i_1} & \downarrow & & \\ & \mathbb{P}'_2\mathbb{P}'(\mathbf{X}, \mathbf{Y}) & \xrightarrow{\mu} & \mathbb{P}'_2(\mathbf{X}, \mathbf{Y}) & \\ & \mathbb{P}'_2\nu \downarrow & & \downarrow \nu & \\ \mathbf{X} & \longrightarrow & \mathbb{P}'_2(\mathbf{X}, \mathbf{Y}) & \xrightarrow{\nu} & \mathbf{Y}. \end{array}$$

Let H denote the composite of the right-hand vertical arrows. Then the diagram shows that the composite

$$(18.1) \quad \mathbf{X}^{\wedge j} \rightarrow \mathbf{X} \xrightarrow{f} \mathbf{Y}$$

is H .

Next consider the commutative diagram

$$\begin{array}{ccccc}
 \mathbf{X}^{\wedge j} & \xrightarrow{=} & \mathbf{X}^{\wedge j} & & \\
 \downarrow f^{\wedge j} & \searrow i_2 & \downarrow & & \\
 & & \mathbb{P}'_2 \mathbb{P}'(\mathbf{X}, \mathbf{Y}) & \xrightarrow{\mu} & \mathbb{P}'_2(\mathbf{X}, \mathbf{Y}) \\
 & & \downarrow \mathbb{P}'_2 \nu & & \downarrow \nu \\
 \mathbf{Y}^{\wedge j} & \longrightarrow & \mathbb{P}'_2(\mathbf{X}, \mathbf{Y}) & \xrightarrow{\nu} & \mathbf{Y}.
 \end{array}$$

This diagram shows that the composite

$$(18.2) \quad \mathbf{X}^{\wedge j} \xrightarrow{f^{\wedge j}} \mathbf{Y}^{\wedge j} \rightarrow \mathbf{Y}$$

is also H . Therefore the composites (18.1) and (18.2) are equal as required.

Part (ii) is immediate from the definitions, and the proof of part (iii) is the same as for Proposition 10.2(iii) (but using Remark 15.7(ii)). \square

Proof of Theorem 1.3. The proof follows the outline of the proof of Theorem 1.1 (given in Section 10); we refer the reader to that proof for some of the details. We have a diagram of simplicial \mathbb{P} -algebras

$$(\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^{\mathbb{Z}})_{\bullet} \xleftarrow{\varepsilon} B_{\bullet}(\mathbb{P}, \mathbb{P}, (\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^{\mathbb{Z}})) \xrightarrow{\Xi_{\bullet}} B_{\bullet}(\mathbb{P}', \mathbb{P}, (\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^{\mathbb{Z}})).$$

By Lemma 17.3, this gives a diagram

$$(18.3) \quad \begin{array}{ccccc}
 (\mathbf{R}_{e,*,1})_{\bullet} & \xleftarrow{\varepsilon} & B_{\bullet}(\mathbb{P}_1, \mathbb{P}, (\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^{\mathbb{Z}})) & \xrightarrow{(\Xi_1)_{\bullet}} & B_{\bullet}(\mathbb{P}'_1, \mathbb{P}, (\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^{\mathbb{Z}})) \\
 \text{sig}_{\text{rel}} \downarrow & & \downarrow & & \downarrow \\
 (\mathbf{R}_{\text{rel}}^{\mathbb{Z}})_{\bullet} & \xleftarrow{\varepsilon} & B_{\bullet}(\mathbb{P}_2, \mathbb{P}, (\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^{\mathbb{Z}})) & \xrightarrow{(\Xi_2)_{\bullet}} & B_{\bullet}(\mathbb{P}'_2, \mathbb{P}, (\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^{\mathbb{Z}})),
 \end{array}$$

in which all objects are simplicial monoids and all horizontal arrows are monoid maps. By Proposition 18.2(i), the right column is a simplicial monoid map between simplicial commutative monoids. Moreover, each map ε is a homotopy equivalence of simplicial objects, and (using Proposition 18.2(iii)) $(\Xi_1)_{\bullet}$ and $(\Xi_2)_{\bullet}$ are weak equivalences in each simplicial degree.

The objects of the diagram (18.3) are simplicial objects in $\Sigma\mathcal{S}_{\text{ss}}$. We obtain a diagram

$$(18.4) \quad \begin{array}{ccccc} |(\mathbf{R}_{e,*,1})_{\bullet}| & \xleftarrow{|\varepsilon|} & |B_{\bullet}(\mathbb{P}_1, \mathbb{P}, (\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^Z))| & \xrightarrow{|(\Xi_1)_{\bullet}|} & |B_{\bullet}(\mathbb{P}'_1, \mathbb{P}, (\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^Z))| \\ \text{sig}_{\text{rel}} \downarrow & & \downarrow & & \downarrow \\ |(\mathbf{R}_{\text{rel}}^Z)_{\bullet}| & \xleftarrow{|\varepsilon|} & |B_{\bullet}(\mathbb{P}_2, \mathbb{P}, (\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^Z))| & \xrightarrow{|(\Xi_2)_{\bullet}|} & |B_{\bullet}(\mathbb{P}'_2, \mathbb{P}, (\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^Z))| \end{array}$$

of simplicial objects in $\Sigma\mathcal{S}$ (the category of symmetric spectra) by applying the geometric realization functor $\Sigma\mathcal{S}_{\text{ss}} \rightarrow \Sigma\mathcal{S}$ to the diagram (18.3) in each simplicial degree. All objects are simplicial monoids and all horizontal arrows are monoid maps, and the right column is a simplicial monoid map between simplicial commutative monoids. The maps $|\varepsilon|$ are homotopy equivalences of simplicial objects and the maps $|(\Xi_1)_{\bullet}|$ and $|(\Xi_2)_{\bullet}|$ are weak equivalences in each simplicial degree.

Finally, we apply geometric realization to the diagram (18.4). We define \mathbf{A} to be $||B_{\bullet}(\mathbb{P}_1, \mathbb{P}, (\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^Z))||$, \mathbf{B} to be $||B_{\bullet}(\mathbb{P}_2, \mathbb{P}, (\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^Z))||$, \mathbf{C} to be $||B_{\bullet}(\mathbb{P}'_1, \mathbb{P}, (\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^Z))||$, and \mathbf{D} to be $||B_{\bullet}(\mathbb{P}'_2, \mathbb{P}, (\mathbf{R}_{e,*,1}, \mathbf{R}_{\text{rel}}^Z))||$. This gives the diagram of Theorem 1.3. \square

Remark 18.3. The symmetric ring spectrum \mathbf{C} is the same as the symmetric ring spectrum $\mathbf{M}_{e,*,1}^{\text{comm}}$ given by Theorem 1.1. There is a ring map

$$(\mathbf{M}_{\text{rel}}^Z)^{\text{comm}} \rightarrow \mathbf{D}$$

which is a weak equivalence (because there is a commutative diagram whose first row is Diagram (10.3) and whose second row is the second row of the diagram in Theorem 1.3).

19. IMPROVED VERSIONS OF GEOMETRIC AND SYMMETRIC POINCARÉ BORDISM.

In order to state our next theorem we need some background.

Let $\Sigma\mathcal{S}$ denote the category of symmetric spectra.

Recall (from [LM14, Definition 13.2(i) and the second paragraph of Section 19]) the strict monoidal category \mathcal{T} whose objects are the triples (π, Z, w) , where π is a group, Z is a simply-connected free π -space, and w is a homomorphism $\pi \rightarrow \{\pm 1\}$. There is a monoidal functor

$$\mathbf{M}_{\text{geom}} : \mathcal{T} \rightarrow \Sigma\mathcal{S}$$

which takes (π, Z, w) to $\mathbf{M}_{\pi,Z,w}$ ([LM14, Definition 19.1 and Theorem 19.2]).

Let \mathcal{R} be the category of rings with involution. There is a functor

$$\mathbf{M}_{\text{sym}} : \mathcal{R} \rightarrow \Sigma\mathcal{S}$$

which takes R to $\mathbf{M}_{\text{rel}}^R$ (we used the symbol \mathbf{M}_{sym} for a different but related functor in [LM14, Section 19]). The proof of ([LM14, Theorem 19.2]) shows that \mathbf{M}_{sym} is monoidal.

There is a functor $\rho : \mathcal{T} \rightarrow \mathcal{R}$ which takes (π, Z, w) to $\mathbb{Z}[\pi]$ with the w -twisted involution ([LM14, Definition 13.2(ii)]). In Section 14 we constructed a natural transformation

$$\text{sig}_{\text{rel}} : \mathbf{M}_{\text{geom}} \rightarrow \mathbf{M}_{\text{sym}} \circ \rho.$$

Now sig_{rel} is not a monoidal transformation, and \mathbf{M}_{geom} and \mathbf{M}_{sym} are not symmetric monoidal functors (we recall the definitions of monoidal transformation and symmetric monoidal functor below). Our next result shows that there is a monoidal transformation between symmetric monoidal functors which is weakly equivalent to sig_{rel} .

Theorem 19.1. *There are symmetric monoidal functors $\mathbf{P} : \mathcal{T} \rightarrow \Sigma\mathcal{S}$, $\mathbf{L}_{\text{sym}} : \mathcal{R} \rightarrow \Sigma\mathcal{S}$, and a monoidal natural transformation $\text{sig} : \mathbf{P} \rightarrow \mathbf{L}_{\text{sym}} \circ \rho$ such that*

(i) *\mathbf{P} is weakly equivalent as a monoidal functor to \mathbf{M}_{geom} ; specifically, there is a monoidal functor $\mathbf{A} : \mathcal{T} \rightarrow \Sigma\mathcal{S}$ and monoidal weak equivalences*

$$\mathbf{M}_{\text{geom}} \leftarrow \mathbf{A} \rightarrow \mathbf{P}.$$

(ii) *\mathbf{L}_{sym} is weakly equivalent as a monoidal functor to \mathbf{M}_{sym} ; specifically, there is a monoidal functor $\mathbf{B} : \mathcal{R} \rightarrow \Sigma\mathcal{S}$ and monoidal weak equivalences*

$$\mathbf{M}_{\text{sym}} \leftarrow \mathbf{B} \rightarrow \mathbf{L}_{\text{sym}}.$$

(iii) *The natural transformations $\text{sig} : \mathbf{P} \rightarrow \mathbf{L}_{\text{sym}} \circ \rho$ and $\text{sig}_{\text{rel}} : \mathbf{M}_{\text{geom}} \rightarrow \mathbf{M}_{\text{sym}} \circ \rho$ are weakly equivalent; specifically, there is a natural transformation $\mathbf{A} \rightarrow \mathbf{B} \circ \rho$ which makes the following diagram strictly commute*

$$\begin{array}{ccccc} \mathbf{M}_{\text{geom}} & \longleftarrow & \mathbf{A} & \longrightarrow & \mathbf{P} \\ \text{sig}_{\text{rel}} \downarrow & & \downarrow & & \downarrow \text{sig} \\ \mathbf{M}_{\text{sym}} \circ \rho & \longleftarrow & \mathbf{B} \circ \rho & \longrightarrow & \mathbf{L}_{\text{sym}} \circ \rho. \end{array}$$

Remark 19.2. (i) It is perhaps worth mentioning that there is no such thing as a *symmetric* monoidal transformation, just as there is no such thing as a commutative homomorphism between commutative rings.

(ii) Theorem 19.1 implies that $\mathbf{L}_{\text{sym}}(R)$ is a strictly commutative symmetric ring spectrum when R is commutative. Also, $\mathbf{P}(e, *, 1)$ is a strictly commutative symmetric ring spectrum and $\text{sig} : \mathbf{P}(e, *, 1) \rightarrow \mathbf{L}_{\text{sym}}(\mathbb{Z})$ is a map of symmetric ring spectra. This is compatible with

Theorem 1.3: there is a commutative diagram

$$\begin{array}{ccc} \mathbf{C} & \longrightarrow & \mathbf{P}(e, *, 1) \\ \downarrow & & \text{sig} \downarrow \\ \mathbf{D} & \longrightarrow & \mathbf{L}_{\text{sym}}(\mathbb{Z}) \end{array}$$

in which the horizontal arrows are ring maps, and they are weak equivalences by the argument given in Remark 18.3.

(iii) The fact that sig is a monoidal functor is a spectrum-level version of Ranicki's multiplicativity formula for the symmetric signature ([Ran80b, Proposition 8.1(i)]). It seems likely that his multiplicativity formula for the surgery obstruction ([Ran80b, Proposition 8.1(ii)]) can also be given a spectrum-level interpretation.

We recall the definitions of symmetric monoidal functor and monoidal transformation. The theorem says that \mathbf{L}_{sym} (and similarly \mathbf{P}) is a monoidal functor with the additional property that the diagram

$$\begin{array}{ccc} \mathbf{L}_{\text{sym}}(R) \wedge \mathbf{L}_{\text{sym}}(S) & \longrightarrow & \mathbf{L}_{\text{sym}}(R \otimes S) \\ \downarrow & & \downarrow \\ \mathbf{L}_{\text{sym}}(S) \wedge \mathbf{L}_{\text{sym}}(R) & \longrightarrow & \mathbf{L}_{\text{sym}}(S \otimes R) \end{array}$$

strictly commutes. Moreover, sig has the property that the diagrams

$$\begin{array}{ccc} & \mathbf{S} & \\ & \swarrow & \searrow \\ \mathbf{P}(e, *, 1) & \xrightarrow{\text{sig}} & \mathbf{L}_{\text{sym}}(\mathbb{Z}) \end{array}$$

and

$$\begin{array}{ccc} \mathbf{P}(\pi, Z, w) \wedge \mathbf{P}(\pi', Z', w') & \xrightarrow{\text{sig} \wedge \text{sig}} & \mathbf{L}_{\text{sym}}(\mathbb{Z}[\pi]^w) \wedge \mathbf{L}_{\text{sym}}(\mathbb{Z}[\pi']^{w'}) \\ \downarrow & & \downarrow \\ & & \mathbf{L}_{\text{sym}}(\mathbb{Z}[\pi]^w \otimes \mathbb{Z}[\pi']^{w'}) \\ \downarrow & & \downarrow \\ \mathbf{P}(\pi \times \pi', Z \times Z', w \cdot w') & \xrightarrow{\text{sig}} & \mathbf{L}_{\text{sym}}(\mathbb{Z}[\pi \times \pi']^{w \cdot w'}) \end{array}$$

strictly commute.

20. PROOF OF THEOREM 19.1

The proof is a modification of the proof of Theorem 1.3; the main difference is that we need more elaborate notation.

Notation 20.1. (i) For an object x of \mathcal{T} or \mathcal{R} , write \mathcal{A}_x for the corresponding \mathbb{Z} -graded category and \mathbf{R}_x for the associated object of $\Sigma\mathcal{S}_{\text{ss}}$.

(ii) Given a j -tuple (x_1, \dots, x_j) , where each x_i is an object of \mathcal{T} or \mathcal{R} , write

$$[x_1, \dots, x_j]$$

for $y_1 \otimes \dots \otimes y_j$, where y_i is x_i if x_i is an object of \mathcal{R} and $\rho(x_i)$ if x_i is an object of \mathcal{T} .

(iii) Given a j -tuple (f_1, \dots, f_j) , where each f_i is a morphism in \mathcal{T} or \mathcal{R} , write

$$[f_1, \dots, f_j]$$

for $g_1 \otimes \dots \otimes g_j$, where g_i is f_i if f_i is a morphism in \mathcal{R} and $\rho(f_i)$ if f_i is a morphism in \mathcal{T} .

The reader should see Proposition 20.11(i) below for motivation for the following definitions.

Definition 20.2. (i) Let y be an object of \mathcal{T} . An *entity* of type $(r_u(j), y)$ is a $j + 1$ -tuple (x_1, \dots, x_j, f) , where f is a morphism in \mathcal{T} from $x_1 \boxtimes \dots \boxtimes x_j$ to y .

(ii) Let $\mathcal{E}_{r_u(j), y}$ denote the set of entities of type $(r_u(j), y)$.

(iii) Let z be an object of \mathcal{R} and let $r : \{1, \dots, j\} \rightarrow \{u, v\}$ be a function. An *entity* of type (r, z) is a $j + 1$ -tuple (x_1, \dots, x_j, f) , where each x_i is an object of \mathcal{T} or \mathcal{R} and f is a morphism in \mathcal{R} from $[x_1, \dots, x_j]$ to z .

(iv) Let $\mathcal{E}_{r, z}$ denote the set of entities of type (r, z) .

Notation 20.3. (i) Let \mathfrak{S} denote the union of the set of objects of \mathcal{T} and the set of objects of \mathcal{R} .

(ii) Let $\Pi\Sigma\mathcal{S}_{\text{ss}}$ be the infinite product of copies of $\Sigma\mathcal{S}_{\text{ss}}$, indexed over \mathfrak{S} .

We will define a monad in $\Pi\Sigma\mathcal{S}_{\text{ss}}$.

First we need to define the relevant right Σ_j actions. Recall Definition 16.1(i).

Definition 20.4. Let $\{\mathbf{X}_x\}_{x \in \mathfrak{S}}$ be an object of $\Pi\Sigma\mathcal{S}_{\text{ss}}$ and let $j \geq 0$.

(i) Given an object y of \mathcal{T} and $\alpha \in \Sigma_j$, define a map $\bar{\alpha}$ from

$$\bigvee_{(x_1, \dots, x_j, f) \in \mathcal{E}_{r_u(j), y}} \mathcal{O}(r_u(j); u) \bar{\wedge} (\mathbf{X}_{x_1} \wedge \dots \wedge \mathbf{X}_{x_j})$$

to itself to be the map which takes the summand indexed by (x_1, \dots, x_j, f) to the summand indexed by $(x_{\alpha(1)}, \dots, x_{\alpha(j)}, f \circ \alpha)$ by means of the map

$$\mathcal{O}(r_u(j); u) \bar{\wedge} (\mathbf{X}_{x_1} \wedge \dots \wedge \mathbf{X}_{x_j}) \xrightarrow{\alpha \bar{\wedge} \alpha} \mathcal{O}(r_u(j); u) \bar{\wedge} (\mathbf{X}_{x_{\alpha(1)}} \wedge \dots \wedge \mathbf{X}_{x_{\alpha(j)}}).$$

(ii) Given an object z of \mathcal{R} and $\alpha \in \Sigma_j$, define a map $\bar{\alpha}$ from

$$\bigvee_r \bigvee_{(x_1, \dots, x_j, f) \in \mathcal{E}_{(r, z)}} \mathcal{O}(r; v) \bar{\wedge} (\mathbf{X}_{x_1} \wedge \dots \wedge \mathbf{X}_{x_j})$$

to itself to be the map which takes the summand indexed by $(x_1, \dots, x_j, f) \in \mathcal{E}_{(r, z)}$ to the summand indexed by $(x_{\alpha(1)}, \dots, x_{\alpha(j)}, f \circ \alpha) \in \mathcal{E}_{(r^\alpha, z)}$ by means of the map

$$\mathcal{O}(r; v) \bar{\wedge} (\mathbf{X}_{x_1} \wedge \dots \wedge \mathbf{X}_{x_j}) \xrightarrow{\bar{\alpha} \bar{\wedge} \alpha} \mathcal{O}(r^\alpha; v) \bar{\wedge} (\mathbf{X}_{x_{\alpha(1)}} \wedge \dots \wedge \mathbf{X}_{x_{\alpha(j)}}).$$

Note that this definition gives right Σ_j actions on the objects mentioned.

Definition 20.5. Let $\{\mathbf{X}_x\}_{x \in \mathfrak{S}}$ be an object of $\Pi\Sigma\mathcal{S}_{\text{ss}}$.

(i) Given an object y of \mathcal{T} , define

$$\mathbb{P}_y(\{\mathbf{X}_x\}_{x \in \mathfrak{S}})$$

to be

$$\bigvee_{j \geq 0} \left(\bigvee_{(x_1, \dots, x_j, f) \in \mathcal{E}_{r_u(j), y}} \mathcal{O}(r_u(j); u) \bar{\wedge} (\mathbf{X}_{x_1} \wedge \dots \wedge \mathbf{X}_{x_j}) \right) / \Sigma_j.$$

(ii) Given an object z of \mathcal{R} , define

$$\mathbb{P}_z(\{\mathbf{X}_x\}_{x \in \mathfrak{S}})$$

to be

$$\bigvee_{j \geq 0} \left(\bigvee_r \bigvee_{(x_1, \dots, x_j, f) \in \mathcal{E}_{(r, z)}} \mathcal{O}(r; v) \bar{\wedge} (\mathbf{X}_{x_1} \wedge \dots \wedge \mathbf{X}_{x_j}) \right) / \Sigma_j.$$

(iii) Define $\mathbb{P} : \Pi\Sigma\mathcal{S}_{\text{ss}} \rightarrow \Pi\Sigma\mathcal{S}_{\text{ss}}$ to be the functor whose projection on the y factor (where y is an object of \mathcal{T}) is \mathbb{P}_y and whose projection on the z factor (where z is an object of \mathcal{R}) is \mathbb{P}_z .

Definition 20.6. Let $\{\mathbf{X}_x\}_{x \in \mathfrak{S}}$ be an object of $\Pi\Sigma\mathcal{S}_{\text{ss}}$.

(i) For an object y of \mathcal{T} , define

$$\iota_y : \mathbf{X}_y \rightarrow \mathbb{P}_y(\{\mathbf{X}_x\}_{x \in \mathfrak{S}})$$

to be the composite

$$\mathbf{X}_y \cong \bar{\mathbf{S}} \bar{\wedge} \mathbf{X}_y = \mathcal{O}(r_u(1); u) \bar{\wedge} \mathbf{X}_y \hookrightarrow \mathbb{P}_y(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}),$$

where the last map is the inclusion of the summand corresponding to the entity (y, id) .

(ii) For an object z of \mathcal{R} , define

$$\iota_z : \mathbf{X}_z \rightarrow \mathbb{P}_z(\{\mathbf{X}_x\}_{x \in \mathfrak{S}})$$

to be the composite

$$\mathbf{X}_z \cong \overline{\mathbf{S}} \overline{\wedge} \mathbf{X}_z = \mathcal{O}(r_v(1); v) \overline{\wedge} \mathbf{X}_y \hookrightarrow \mathbb{P}_z(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}),$$

where the last map is the inclusion of the summand corresponding to the entity (z, id) .

(iii) Define

$$\iota : \{\mathbf{X}_x\}_{x \in \mathfrak{S}} \rightarrow \mathbb{P}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}})$$

to be the map whose projection on the y factor (where y is an object of \mathcal{T}) is ι_y and whose projection on the z factor (where z is an object of \mathcal{R}) is ι_z .

In order to define the structure map $\mu : \mathbb{P}\mathbb{P} \rightarrow \mathbb{P}$ we need a composition operation on entities. For part (ii) we use Notation 20.1(iii) and the notation of Definition 16.3.

Definition 20.7. Let $i \geq 0$, and for each l with $1 \leq l \leq i$ let $j_l \geq 0$.

(i) Let y be an object of \mathcal{T} and let

$$\mathbf{e} = (x_1, \dots, x_i, f) \in \mathcal{E}_{r_u(i), y}.$$

For each l with $1 \leq l \leq i$ let

$$\mathbf{e}_l = (x_1^{(l)}, \dots, x_{j_l}^{(l)}, f^{(l)}) \in \mathcal{E}_{r_u(j_l), x_l}.$$

Define

$$\mathbf{e} \circ (\mathbf{e}_1, \dots, \mathbf{e}_i) \in \mathcal{E}_{r_u(j_1 + \dots + j_i), y}$$

to be

$$(x_1^{(1)}, \dots, x_{j_i}^{(i)}, g),$$

where g is the composite

$$x_1^{(1)} \boxtimes \dots \boxtimes x_{j_i}^{(i)} \xrightarrow{f^{(1)} \boxtimes \dots \boxtimes f^{(i)}} x_1 \boxtimes \dots \boxtimes x_i \xrightarrow{f} y.$$

(ii) Let z be an object of \mathcal{R} , let $r : \{1, \dots, i\} \rightarrow \{u, v\}$, and let

$$\mathbf{e} = (x_1, \dots, x_i, f) \in \mathcal{E}_{r, z}.$$

For each l with $1 \leq l \leq i$ let $r_l : \{1, \dots, j_l\} \rightarrow \{u, v\}$; assume that if $r(l) = u$ then r_l is $r_u(j_l)$. Let

$$\mathbf{e}_l = (x_1^{(l)}, \dots, x_{j_l}^{(l)}, f^{(l)}) \in \mathcal{E}_{r_l, x_l}.$$

Define

$$\mathbf{e} \circ (\mathbf{e}_1, \dots, \mathbf{e}_i) \in \mathcal{E}_{R, z}$$

to be

$$(x_1^{(1)}, \dots, x_{j_i}^{(i)}, g),$$

where g is the composite

$$[x_1^{(1)}, \dots, x_{j_i}^{(i)}] \xrightarrow{[f^{(1)}, \dots, f^{(i)}]} [x_1, \dots, x_i] \xrightarrow{f} z.$$

Now we can define $\mu : \mathbb{P}\mathbb{P} \rightarrow \mathbb{P}$. We begin with the projection on the y -factor,

$$\mu_y : \mathbb{P}_y\mathbb{P} \rightarrow \mathbb{P}_y,$$

where y is an object of \mathcal{T} . A collection of entities $\mathbf{e}, \mathbf{e}_1, \dots, \mathbf{e}_i$ as in Definition 20.7(i) determines a summand

$$\emptyset(r_u(i); u) \overline{\wedge} \left((\emptyset(r_u(j_1); u) \overline{\wedge} (\mathbf{X}_{x_1^{(1)}} \wedge \dots \wedge \mathbf{X}_{x_{j_1}^{(1)}})) \wedge \dots \right)$$

in $\mathbb{P}_y\mathbb{P}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}})$. We define the restriction of μ_y to this summand to be the map to the summand of $\mathbb{P}_y(\{\mathbf{X}_x\}_{x \in \mathfrak{S}})$ indexed by $\mathbf{e} \circ (\mathbf{e}_1, \dots, \mathbf{e}_i)$ which is induced (after passage to quotients) by the composite

$$\begin{aligned} & \emptyset(r_u(i); u) \overline{\wedge} \left((\emptyset(r_u(j_1); u) \overline{\wedge} (\mathbf{X}_{x_1^{(1)}} \wedge \dots \wedge \mathbf{X}_{x_{j_1}^{(1)}})) \wedge \dots \right) \\ & \rightarrow \left(\emptyset(r_u(i); u) \overline{\wedge} (\emptyset(r_u(j_1); u) \otimes \dots) \right) \overline{\wedge} (\mathbf{X}_{x_1^{(1)}} \otimes \dots \otimes \mathbf{X}_{x_{j_i}^{(i)}}) \\ & \xrightarrow{\gamma \overline{\wedge} 1} \emptyset(r_u(j_1 + \dots + j_i); u) \overline{\wedge} (\mathbf{X}_{x_1^{(1)}} \otimes \dots \otimes \mathbf{X}_{x_{j_i}^{(i)}}). \end{aligned}$$

The projection of μ on the z factor (where z is an object of \mathcal{R}) is defined similarly (using Definition 16.3).

Next we give the action of \mathbb{P} on the object $\{\mathbf{R}_x\}_{x \in \mathfrak{S}}$. Let y be an object of \mathcal{T} and let (x_1, \dots, x_j, f) be an entity of type $(r_u(j), y)$. A slight modification of Definition 6.9 gives a map

$$\emptyset(r_u(j); u) \overline{\wedge} (\mathbf{R}_{x_1} \wedge \dots \wedge \mathbf{R}_{x_j}) \rightarrow \mathbf{R}_{x_1 \boxtimes \dots \boxtimes x_j},$$

and composing with the map induced by f gives a map

$$(20.1) \quad \emptyset(r_u(j); u) \overline{\wedge} (\mathbf{R}_{x_1} \wedge \dots \wedge \mathbf{R}_{x_j}) \rightarrow \mathbf{R}_y.$$

We define

$$\nu_y : \mathbb{P}_y(\{\mathbf{R}_x\}_{x \in \mathfrak{S}}) \rightarrow \mathbf{R}_y$$

to be the map whose restriction to the summand indexed by (x_1, \dots, x_j, f) is the map (20.1). We define

$$\nu_z : \mathbb{P}_z(\{\mathbf{R}_x\}_{x \in \mathfrak{S}}) \rightarrow \mathbf{R}_z$$

similarly when z is an object of \mathcal{R} (using a slight modification of Definition 16.7), and we define

$$\nu : \mathbb{P}(\{\mathbf{R}_x\}_{x \in \mathfrak{S}}) \rightarrow \{\mathbf{R}_x\}_{x \in \mathfrak{S}}$$

to be the map with projections ν_y and ν_z .

Lemma 20.8. ν is an action of \mathbb{P} on $\{\mathbf{R}_x\}_{x \in \mathfrak{S}}$. □

Now we need the analogue of Lemma 17.3. Let \mathcal{C} be the category whose objects are triples $(\mathbf{F}, \mathbf{G}, \mathbf{t})$, where \mathbf{F} is a monoidal functor $\mathcal{T} \rightarrow \Sigma\mathcal{S}_{\text{ss}}$, \mathbf{G} is a monoidal functor $\mathcal{R} \rightarrow \Sigma\mathcal{S}_{\text{ss}}$, and \mathbf{t} is a natural transformation $\mathbf{F} \rightarrow \mathbf{G} \circ \rho$ which is not required to be a monoidal transformation; the morphisms are commutative diagrams

$$\begin{array}{ccc} \mathbf{F} & \longrightarrow & \mathbf{F}' \\ \mathbf{t} \downarrow & & \downarrow \mathbf{t}' \\ \mathbf{G} \circ \rho & \longrightarrow & \mathbf{G}' \circ \rho, \end{array}$$

where the horizontal arrows are monoidal transformations.

Let us write \mathbf{R}_{geom} (resp., \mathbf{R}_{sym}) for the functor $\mathcal{T} \rightarrow \Sigma\mathcal{S}_{\text{ss}}$ (resp., $\mathcal{R} \rightarrow \Sigma\mathcal{S}_{\text{ss}}$) which takes x to \mathbf{R}_x .

Lemma 20.9. (i) *There is a functor Υ from \mathbb{P} algebras to \mathcal{C} which takes $\{\mathbf{X}_x\}_{x \in \mathfrak{S}}$ to a triple $(\mathbf{F}, \mathbf{G}, \mathbf{t})$ with $\mathbf{F}(y) = \mathbf{X}_y$ and $\mathbf{G}(z) = \mathbf{X}_z$.*

(ii) *$\Upsilon(\{\mathbf{R}_x\}_{x \in \mathfrak{S}})$ is the triple*

$$(\mathbf{R}_{\text{geom}}, \mathbf{R}_{\text{sym}}, \text{sig}_{\mathfrak{S}\text{rel}})$$

Proof. Part (i). Let $\{\mathbf{X}_x\}_{x \in \mathfrak{S}}$ be a \mathbb{P} algebra. Define a functor

$$\mathbf{F} : \mathcal{T} \rightarrow \Sigma\mathcal{S}_{\text{ss}}$$

on objects by $\mathbf{F}(y) = \mathbf{X}_y$ and on morphisms by letting $\mathbf{F}(f : y \rightarrow y')$ be the composite

$$\mathbf{X}_y \cong \mathcal{O}(r_u(1); u) \bar{\wedge} \mathbf{X}_y \hookrightarrow \mathbb{P}_{y'}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}) \xrightarrow{\nu_{y'}} \mathbf{X}_{y'},$$

where the unlabeled arrow is the inclusion of the summand indexed by the entity (y, f) . We define

$$\mathbf{G} : \mathcal{R} \rightarrow \Sigma\mathcal{S}_{\text{ss}}$$

similarly. The proof that \mathbf{F} and \mathbf{G} are monoidal functors is similar to the argument, in the proof in Lemma 17.3(i), that \mathbf{X} and \mathbf{Y} are monoids.

It remains to give the natural transformation

$$\mathbf{t} : \mathbf{F} \rightarrow \mathbf{G} \circ \rho.$$

For an object y of \mathcal{T} , let \mathbf{t}_y be the composite

$$\mathbf{X}_y \cong \mathcal{O}(r_u(1); v) \bar{\wedge} \mathbf{X}_y \hookrightarrow \mathbb{P}_{\rho(y)}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}) \xrightarrow{\nu_{\rho(y)}} \mathbf{X}_{\rho(y)},$$

where the unlabeled arrow is the inclusion of the summand indexed by the entity $(y, \text{id}) \in \mathcal{E}_{r_u(1), \rho(y)}$. To show that \mathbf{t} is a natural transformation, let $f : y \rightarrow y'$ be a morphism in \mathcal{T} , and let $z = \rho(y)$, $z' = \rho(y')$.

Let i_1 be the composite

$$\begin{aligned} \mathbf{X}_y \cong \mathcal{O}(r_u(1); u) \overline{\wedge} \mathbf{X}_y &\hookrightarrow \mathbb{P}_{y'}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}) \cong \mathcal{O}(r_u(1); v) \overline{\wedge} \mathbb{P}_{y'}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}) \\ &\hookrightarrow \mathbb{P}_{z'} \mathbb{P}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}), \end{aligned}$$

where the first arrow is the inclusion of the summand indexed by the entity (y, f) and the second is the inclusion of the summand indexed by (y', id) . Let j_1 be the composite

$$\mathbf{X}_{y'} \cong \mathcal{O}(r_u(1); v) \overline{\wedge} \mathbf{X}_{y'} \hookrightarrow \mathbb{P}_{z'}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}),$$

where the inclusion is indexed by (y', id) , and let j_2 be the composite

$$\mathbf{X}_y \cong \mathcal{O}(r_u(1); v) \overline{\wedge} \mathbf{X}_y \hookrightarrow \mathbb{P}_{z'}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}),$$

where the inclusion is indexed by $(y, \rho(f))$.

Consider the commutative diagram

$$\begin{array}{ccccc} \mathbf{X}_y & \xrightarrow{=} & \mathbf{X}_y & & \\ \downarrow \mathbf{F}(f) & \searrow i_1 & & & \downarrow j_2 \\ & \mathbb{P}_{z'} \mathbb{P}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}) & \xrightarrow{\mu} & \mathbb{P}_{z'}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}) & \\ & \mathbb{P}_{z'} \nu \downarrow & & \downarrow \nu & \\ \mathbf{X}_{y'} & \xrightarrow{j_1} & \mathbb{P}_{z'}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}) & \xrightarrow{\nu} & \mathbf{X}_{z'}. \end{array}$$

Let H denote the composite of the right-hand vertical arrows. Then the diagram shows that the composite

$$(20.2) \quad \mathbf{X}_y \xrightarrow{\mathbf{F}(f)} \mathbf{X}_{y'} \xrightarrow{\mathbf{t}} \mathbf{X}_{z'}$$

is H .

Let i_2 be the composite

$$\begin{aligned} \mathbf{X}_y \cong \mathcal{O}(r_u(1); v) \overline{\wedge} \mathbf{X}_y &\hookrightarrow \mathbb{P}_z(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}) \cong \mathcal{O}(r_v(1); v) \overline{\wedge} \mathbb{P}_z(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}) \\ &\hookrightarrow \mathbb{P}_{z'} \mathbb{P}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}), \end{aligned}$$

where the first inclusion is indexed by (y, id) and the second is indexed by $(z, \rho(f))$. Let j_2 be as above and let j_3 be the composite

$$\mathbf{X}_z \cong \mathcal{O}(r_v(1); v) \overline{\wedge} \mathbf{X}_z \hookrightarrow \mathbb{P}_z(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}),$$

where the inclusion is indexed by $(z, \rho(f))$.

Consider the commutative diagram

$$\begin{array}{ccc}
 \mathbf{X}_y & \xrightarrow{=} & \mathbf{X}_y \\
 \downarrow \mathfrak{t} & \searrow i_1 & \downarrow j_2 \\
 & \mathbb{P}_{z'} \mathbb{P}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}) & \xrightarrow{\mu} \mathbb{P}_{z'}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}) \\
 & \downarrow \mathbb{P}_{z'} \nu & \downarrow \nu \\
 \mathbf{X}_z & \xrightarrow{j_3} \mathbb{P}_{z'}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}) & \xrightarrow{\nu} \mathbf{X}_{z'}
 \end{array}$$

This diagram shows that the composite

$$(20.3) \quad \mathbf{X}_y \xrightarrow{\mathfrak{t}} \mathbf{X}_z \xrightarrow{\mathbf{G}(f)} \mathbf{X}_{z'}$$

is also H , so the composites (20.2) and (20.3) are equal as required.

Part (ii) is an easy consequence of the definitions. \square

Finally, we have the analogues of Definition 18.1 and Proposition 18.2.

Definition 20.10. Let $\{\mathbf{X}_x\}_{x \in \mathfrak{S}}$ be an object of $\Pi\Sigma\mathcal{S}_{\text{ss}}$.

(i) Given an object y of \mathcal{T} , define

$$\mathbb{P}'_y(\{\mathbf{X}_x\}_{x \in \mathfrak{S}})$$

to be

$$\bigvee_{j \geq 0} \left(\bigvee_{((x_1, \dots, x_j, f) \in \mathcal{E}_{r_u(j), y}} \mathbf{X}_{x_1} \wedge \cdots \wedge \mathbf{X}_{x_j} \right) / \Sigma_j.$$

(ii) Given an object z of \mathcal{R} , define

$$\mathbb{P}'_z(\{\mathbf{X}_x\}_{x \in \mathfrak{S}})$$

to be

$$\bigvee_{j \geq 0} \left(\bigvee_r \bigvee_{(x_1, \dots, x_j, f) \in \mathcal{E}(r, z)} \mathbf{X}_{x_1} \wedge \cdots \wedge \mathbf{X}_{x_j} \right) / \Sigma_j.$$

(iii) Define $\mathbb{P}' : \Pi\Sigma\mathcal{S}_{\text{ss}} \rightarrow \Pi\Sigma\mathcal{S}_{\text{ss}}$ to be the functor whose projection on the y factor (where y is an object of \mathcal{T}) is \mathbb{P}'_y and whose projection on the z factor (where z is an object of \mathcal{R}) is \mathbb{P}'_z .

A routine modification of Definition 18.1(ii) gives a natural transformation

$$\Xi : \mathbb{P} \rightarrow \mathbb{P}'.$$

Proposition 20.11. (i) *An algebra over \mathbb{P}' is the same thing as a pair of symmetric monoidal functors \mathbf{F} and \mathbf{G} with a monoidal transformation $\mathbf{F} \rightarrow \mathbf{G} \circ \rho$.*

(ii) Ξ is a map of monads.

(iii) Suppose that each $(X_x)_k$ has compatible degeneracies (see Definition 9.1). Let \mathbb{P}^q denote the q -th iterate of \mathbb{P} . Then each map

$$\Xi : \mathbb{P}^q(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}) \rightarrow \mathbb{P}'\mathbb{P}^{q-1}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}})$$

is a weak equivalence.

Proof. Part (i). Let $\{\mathbf{X}_x\}_{x \in \mathfrak{S}}$ be an algebra over \mathbb{P}' . The fact that \mathbf{F} and \mathbf{G} are symmetric monoidal functors is an easy consequence of the definitions. The natural transformation $\mathbf{t} : \mathbf{F} \rightarrow \mathbf{G} \circ \rho$ is constructed as in the proof of Lemma 20.9(i). The proof that \mathbf{t} is monoidal is similar to the proofs of Proposition 18.2(i) and Lemma 20.9(i), using the maps

$$i_1 : \mathbf{X}_{y_1} \wedge \cdots \wedge \mathbf{X}_{y_j} \hookrightarrow \mathbb{P}'_{y_1 \boxtimes \cdots \boxtimes y_j}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}) \hookrightarrow \mathbb{P}'_{\rho(y_1) \boxtimes \cdots \boxtimes y_j} \mathbb{P}'(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}),$$

(where the first inclusion is indexed by $(y_1, \dots, y_j, \text{id})$ and the second by $(y_1 \boxtimes \cdots \boxtimes y_j, \text{id})$), and

$$\begin{aligned} i_2 : \mathbf{X}_{y_1} \wedge \cdots \wedge \mathbf{X}_{y_j} &\hookrightarrow \mathbb{P}'_{\rho(y_1)}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}) \wedge \cdots \wedge \mathbb{P}'_{\rho(y_j)}(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}) \\ &\hookrightarrow \mathbb{P}'_{\rho(y_1) \boxtimes \cdots \boxtimes y_j} \mathbb{P}'(\{\mathbf{X}_x\}_{x \in \mathfrak{S}}), \end{aligned}$$

(where the first map is the smash product of the inclusions indexed by (y_i, id) and the second is indexed by $(\rho(y_1), \dots, \rho(y_j), \rho(y_1) \otimes \cdots \otimes \rho(y_j) \rightarrow \rho(y_1 \boxtimes \cdots \boxtimes y_j))$).

Part (ii) is immediate from the definitions, and the proof of part (iii) is the same as for Proposition 10.2(iii) (but using Remark 15.7(ii)). \square

Now the proof of Theorem 19.1 is the same as the proof of Theorem 1.3 given in Section 18, with only the notation changed.

APPENDIX A. A PROPERTY OF THE SMASH PRODUCT OF SYMMETRIC SPECTRA.

For an object \mathbf{X} of $\Sigma\mathcal{S}_{\text{ss}}$ let

$$\bar{\omega} : X_k \rightarrow X_{k+1}$$

be the map which takes x to $\omega(s \wedge x)$ (where s is the 1-simplex of S^1).

Definition A.1. An object \mathbf{X} of $\Sigma\mathcal{S}_{\text{ss}}$ is *monomorphic* if $\bar{\omega}$ is a monomorphism for every k . It is *strongly monomorphic* if it is monomorphic and has the following property: if $x \in X_k$ and $\alpha \in \Sigma_{k+1}$ with $\alpha(1) \neq 1$, and if $\alpha\bar{\omega}x$ is in the image of $\bar{\omega}$, then there is a $\beta \in \Sigma_k$ such that $\beta^{-1}(1) = \alpha^{-1}(1) - 1$ and βx is in the image of $\bar{\omega}$.

The purpose of this appendix is to prove the following fact, which is used in Section 10:

Proposition A.2. *Let \mathbf{X} be strongly monomorphic and suppose that the only element of X_0 is the basepoint. Then the Σ_j action on $\mathbf{X}^{\wedge j}$ which permutes the factors is free away from the basepoints.*

Remark A.3. (i) It is a straightforward consequence of the definitions that the object \mathbf{R} given in Example 4.13 is strongly monomorphic.

(ii) The analog of Proposition A.2 for simplicial or topological symmetric spectra is also true, with essentially the same proof.

The proof of Proposition A.2 will be given after the proof of our next result, which is the main ingredient in the proof of Proposition A.2.

Proposition A.4. *If $\mathbf{X}_1, \dots, \mathbf{X}_j$ are strongly monomorphic then $\mathbf{X}_1 \wedge \dots \wedge \mathbf{X}_j$ is monomorphic.*

We will give an example at the end of this appendix to show that Propositions A.2 and A.4 both fail if “strongly monomorphic” is replaced by “monomorphic.”

Before we can give the proof of Proposition A.4 we will need quite a bit of background, culminating with Lemma A.15.

If A is a multiseemisimplicial set and $a \in A_{\mathbf{n}}$ for some \mathbf{n} , we will say a is a point of A . If \mathbf{X} is an object of $\Sigma\mathcal{S}_{\text{ss}}$ and x is a point of X_k for some k we will say x is a point of \mathbf{X} and write $|x| = k$.

We will use the fact that (by Remark 4.7) points in the k -fold smash product $\mathbf{X}_1 \wedge \dots \wedge \mathbf{X}_j$ are equivalence classes of symbols of the form

$$\theta \wedge x_1 \wedge \dots \wedge x_j$$

where x_i is a point of \mathbf{X}_i for each i and $\theta \in \Sigma_{|x_1| + \dots + |x_j|}$. Our first task is to describe the equivalence relation \sim explicitly, and for this we need the operations in Definition A.7 below.

Notation A.5. (i) Let $\alpha \in \Sigma_l$, let $L \geq 1$, and let $k \geq L + l - 1$. We will write $\alpha^{[L]}$ for the element of Σ_k which permutes $L, L + 1, \dots, L + l - 1$ in the same way that α permutes $1, \dots, l$.

(ii) Let \mathbf{X} be an object of $\Sigma\mathcal{S}_{\text{ss}}$. Let

$$\psi : \mathbf{X} \wedge \mathbf{S} \rightarrow \mathbf{X}$$

be the right action (that is, $\psi = \omega \circ \tau$, where τ is defined in Equation (4.1)), and let

$$\bar{\psi} : X_k \rightarrow X_{k+1}$$

be the map that takes x to $\psi(x \wedge s)$, where s is the 1-simplex of S^1 . Note that

$$(A.1) \quad \bar{\psi}x = \rho_{1,k}\bar{\omega}x,$$

where $\rho_{1,k}$ is defined after Equation (4.1).

Remark A.6. (i) If x is a point of \mathbf{X} and $\alpha \in \Sigma_{|x|}$ then $\bar{\omega}\alpha x = \alpha^{[2]}\bar{\omega}x$ and $\bar{\psi}\alpha x = \alpha^{[1]}\bar{\psi}x$.

(ii) $\bar{\psi}$ commutes with $\bar{\omega}$.

(iii) Suppose that \mathbf{X} is strongly monomorphic $\Sigma\mathcal{S}_{\text{ss}}$. Using Equation A.1 we see that if x is a point of \mathbf{X} and γ is an element of $\Sigma_{|x|+1}$ for which $\gamma(1) \neq |x| + 1$ and $\gamma\bar{\omega}x$ is in the image of $\bar{\psi}$, then there is a $\delta \in \Sigma_{|x|}$ for which $\delta^{-1}(|x|) = \gamma^{-1}(|x| + 1) - 1$ and δx is in the image of $\bar{\psi}$.

Definition A.7. Let $\mathbf{X}_1, \dots, \mathbf{X}_j$ be monomorphic objects of $\Sigma\mathcal{S}_{\text{ss}}$. Let x_i be a point of \mathbf{X}_i for $1 \leq i \leq j$, and let $\theta \in \Sigma_{|x_1|+\dots+|x_j|}$.

(i) For $1 \leq m \leq j$ and α an element of $\Sigma_{|x_m|}$ other than the identity define

$$A_{m,\alpha}(\theta \wedge x_1 \wedge \dots \wedge x_j) = \theta(\alpha^{[1+\sum_{i<m}|x_i|]})^{-1} \wedge y_1 \wedge \dots \wedge y_j,$$

where

$$y_i = \begin{cases} x_i & \text{if } i \neq m \\ \alpha x_m & \text{if } i = m. \end{cases}$$

(ii) Suppose that $m < j$ and that $x_m = \bar{\psi}z$ for some z (in which case z is uniquely determined since \mathbf{X}_m is monomorphic) define

$$B_m(\theta \wedge x_1 \wedge \dots \wedge x_j) = \theta \wedge y_1 \wedge \dots \wedge y_j,$$

where

$$y_i = \begin{cases} x_i & \text{if } i \neq m, m+1 \\ z & \text{if } i = m \\ \bar{\omega}x_{m+1} & \text{if } i = m+1. \end{cases}$$

(iii) If $m < j$ and $x_{m+1} = \bar{\omega}z$ for some z define

$$C_m(\theta \wedge x_1 \wedge \dots \wedge x_j) = \theta \wedge y_1 \wedge \dots \wedge y_j,$$

where

$$y_i = \begin{cases} x_i & \text{if } i \neq m, m+1 \\ \bar{\psi}x_m & \text{if } i = m \\ z & \text{if } i = m+1 \end{cases}$$

Now we can give an explicit description of the equivalence relation \sim , as follows: $\theta \wedge x_1 \wedge \dots \wedge x_j \sim \kappa \wedge y_1 \wedge \dots \wedge y_j$ if and only if there is a sequence D_1, \dots, D_n of operations of the types given in Definition A.7 with

$$\kappa \wedge y_1 \wedge \dots \wedge y_j = D_n \cdots D_1(\theta \wedge x_1 \wedge \dots \wedge x_j).$$

In this situation we will say that the n -tuple $P = (D_1, \dots, D_n)$ is a *path* from $\theta \wedge x_1 \wedge \dots \wedge x_j$ to $\kappa \wedge y_1 \wedge \dots \wedge y_j$ and we will write

$$P(\theta \wedge x_1 \wedge \dots \wedge x_j) = \kappa \wedge y_1 \wedge \dots \wedge y_j.$$

The *length* of P is the number of A -operations in P plus twice the number of B -operations and twice the number of C -operations.

Notation A.8. If α is the identity element then $A_{m,\alpha}$ will be interpreted as the empty path.

We record some useful calculations in our next two lemmas. We define the *standard path of length $3p$ starting at $\theta \wedge \bar{\omega}x_1 \wedge x_2 \wedge \dots \wedge x_j$* to be the path

$$A_{1,\rho_{1,|x_1|}}, B_1, A_{2,\rho_{1,|x_2|}}, B_2, \dots, A_{p,\rho_{1,|x_p|}}, B_p$$

Note that the B -operations in this sequence are always possible because of Equation (A.1) and that $A_{i,\rho_{1,|x_i|}}$ is interpreted as the empty path when $|x_i| = 0$.

Lemma A.9. *Let P be the standard path of length $3p$ starting at $\theta \wedge \bar{\omega}x_1 \wedge x_2 \wedge \dots \wedge x_j$. Then*

$$P(\theta \wedge \bar{\omega}x_1 \wedge \dots \wedge x_j) = \theta(\rho_{1,|x_1|+\dots+|x_p|}^{[1]})^{-1} \wedge x_1 \wedge \dots \wedge \bar{\omega}x_{p+1} \wedge \dots$$

Proof. This follows by induction on p from Equation A.1 and the equation $\rho_{1,t}^{[s+1]} \rho_{1,s}^{[1]} = \rho_{1,s+t}^{[1]}$ in the group Σ_{s+t+1} . \square

Remark A.10. The operations in Definition A.7 often commute with each other. Specifically, $A_{m,\alpha}$ commutes with $A_{n,\beta}$ for $m \neq n$ and with B_n and C_n for $m \neq n, n-1$; moreover B_m commutes with every B_n and every C_n and C_m commutes with every C_n .

Lemma A.11. *Let P be as in Lemma A.9, and let Q be any path starting at $\theta \wedge \bar{\omega}x_1 \wedge x_2 \wedge \dots \wedge x_j$ of the form*

$$A_{1,\alpha_1}, B_1, A_{2,\alpha_2}, B_2, \dots, A_{p,\alpha_p}, B_p$$

where $\alpha_i(1) = |x_i| + 1$ for each i . For each i let β_i be the element of $\Sigma_{|x_i|}$ with $\beta_i^{[2]} = \rho_{1,|x_i|}^{-1} \alpha_i$ (which exists because $\rho_{1,|x_i|}^{-1} \alpha_i$ takes 1 to 1). Then

(i) the path

$$A_{1,\beta_1^{[2]}}, A_{2,\beta_2}, \dots, A_{p,\beta_p}, P$$

has the same effect on $\theta \wedge \bar{\omega}x_1 \wedge x_2 \wedge \dots \wedge x_j$ as Q , and

(ii) the path

$$P, A_{1,\beta_1}, A_{2,\beta_2}, \dots, A_{p,\beta_p}$$

also has the same effect on $\theta \wedge \bar{\omega}x_1 \wedge x_2 \wedge \dots \wedge x_j$ as Q .

Proof. The proof in each case is by induction on p .

For part (i), the case $p = 1$ is immediate. For $p > 1$, we observe that

$$A_{1,\beta_1^{[2]}}, A_{2,\beta_2}, \dots, A_{p-1,\beta_{p-1}}, P, A_{p,\alpha_p}, B_p$$

has the same effect as

$$A_{1,\beta_1^{[2]}}, A_{2,\beta_2}, \dots, A_{p-1,\beta_{p-1}}, P, A_{p,\beta_p^{[2]}}, A_{p,\rho_{1,|x_p|}}, B_p$$

and by Lemma A.9 this has the same effect as

$$A_{1,\beta_1^{[2]}}, A_{2,\beta_2}, \dots, A_{p,\beta_p}, P, A_{p,\rho_{1,|x_p|}}, B_p$$

as required.

For part (ii), we use the equation $\rho_{1,|x_p|}\beta_p^{[2]} = \beta_p^{[1]}\rho_{1,|x_p|}$. This shows that

$$P, A_{1,\beta_1}, A_{2,\beta_2}, \dots, A_{p-1,\beta_{p-1}}, A_{p,\alpha_p}, B_p$$

has the same effect as

$$P, A_{1,\beta_1}, A_{2,\beta_2}, \dots, A_{p-1,\beta_{p-1}}, A_{p,\rho_{1,|x_p|}}, A_{p,\beta_p^{[1]}}, B_p$$

which has the same effect as

$$P, A_{1,\beta_1}, A_{2,\beta_2}, \dots, A_{p-1,\beta_{p-1}}, A_{p,\rho_{1,|x_p|}}, B_p, A_{p,\beta_p}$$

and by Remark A.10 this has the same effect as

$$P, A_{p,\rho_{1,|x_p|}}, B_p, A_{1,\beta_1}, A_{2,\beta_2}, \dots, A_{p,\beta_p}$$

as required. \square

Notation A.12. Given a symbol $\mathbf{x} = \theta \wedge x_1 \wedge \dots \wedge x_j$, we write $\bar{\omega}\mathbf{x}$ for $\theta^{[2]} \wedge \bar{\omega}x_1 \wedge \dots \wedge x_j$

Definition A.13. Given symbols \mathbf{x} and \mathbf{y} , and a path P from $\bar{\omega}\mathbf{x}$ to $\bar{\omega}\mathbf{y}$, we will say that \mathbf{x} and \mathbf{y} are *P-related*.

With this terminology, Proposition A.4 is true if the following statement is true for all P :

(*) if \mathbf{x} and \mathbf{y} are P -related then $\mathbf{x} \sim \mathbf{y}$.

Before proving Proposition A.4 we need two more lemmas.

Lemma A.14. *Let $\mathbf{x} = \theta \wedge x_1 \wedge \dots \wedge x_j$ and let P be a path starting at \mathbf{x} . Suppose that P consists of a single operation D and that*

- (i) if $D = A_{1,\alpha}$ then $\alpha(1) = 1$, and
- (ii) if $D = B_1$ then $|x_1| \neq 0$.

Then there is a unique \mathbf{y} with $P(\bar{\omega}\mathbf{x}) = \bar{\omega}\mathbf{y}$, and $\mathbf{x} \sim \mathbf{y}$. In particular, Statement () is true for P .*

Proof. If $D = A_{1,\alpha}$ with $\alpha(1) = 1$ then there is a β with $\alpha = \beta^{[2]}$, and

$$P(\bar{\omega}\mathbf{x}) = P(\theta^{[2]} \wedge \bar{\omega}x_1 \wedge \cdots \wedge x_j) = \theta^{[2]}(\beta^{[2]})^{-1} \wedge \bar{\omega}\beta x_1 \wedge \cdots \wedge x_j,$$

so we can let

$$\mathbf{y} = \theta\beta^{-1} \wedge \beta x_1 \wedge \cdots \wedge x_j$$

which is the unique choice since \mathbf{X}_1 is monomorphic. We have $\mathbf{y} = A_{1,\beta}\mathbf{x}$, so $\mathbf{x} \sim \mathbf{y}$.

Next suppose $D = B_1$ and that $|x_1| \neq 0$. Since B_1 can be applied to \mathbf{x} , we must have

$$(A.2) \quad \bar{\omega}x_1 = \bar{\psi}z$$

for some z , and then

$$P(\bar{\omega}\mathbf{x}) = \theta^{[2]} \wedge z \wedge \bar{\omega}x_2 \cdots$$

Combining Equations A.2 and A.1 we have $\rho_{1,|x_1|}\bar{\omega}z = \bar{\omega}x_1$, so since \mathbf{X}_1 is strongly monomorphic there is a $\beta \in \Sigma_{|x_1|}$ with $\beta^{-1}(1) = 1$ and

$$\beta z = \bar{\omega}w$$

for some w . There is a $\tilde{\beta} \in \Sigma_{|x_1|-1}$ with $\beta = \tilde{\beta}^{[2]}$ and we have

$$(A.3) \quad z = (\tilde{\beta}^{[2]})^{-1}\bar{\omega}w = \bar{\omega}\tilde{\beta}^{-1}w.$$

Now let

$$\mathbf{y} = \theta \wedge \tilde{\beta}^{-1}w \wedge \bar{\omega}x_2 \cdots$$

We have $\bar{\omega}\mathbf{y} = P(\bar{\omega}\mathbf{x})$ by Equation A.3. We claim that

$$x_1 = \bar{\psi}\tilde{\beta}^{-1}w$$

Assuming this for the moment, we have $\mathbf{y} = B_1\mathbf{x}$, so $\mathbf{x} \sim \mathbf{y}$ as required. Since \mathbf{X}_1 is monomorphic the claim follows from the equations

$$\bar{\omega}x_1 = \bar{\psi}z = \bar{\psi}\bar{\omega}\tilde{\beta}^{-1}w = \bar{\omega}\bar{\psi}\tilde{\beta}^{-1}w,$$

where we have used Equations A.2 and A.3 and Remark A.6(ii).

The remaining cases are easy. \square

Lemma A.15. *Let Q be a path which can be written as a composite Q_1, Q_2 , where Statement (*) is true for Q_2 and every operation in Q_1 satisfies the hypothesis of Lemma A.14. Then Statement (*) is true for Q .*

Proof. By an iterated application of Lemma A.14, $Q_1(\bar{\omega}\mathbf{x})$ has the form $\bar{\omega}\mathbf{z}$ for some \mathbf{z} , and $\mathbf{x} \sim \mathbf{z}$. But also $\mathbf{z} \sim \mathbf{y}$ since the symbols \mathbf{z} and \mathbf{y} are Q_2 -related. \square

Proof of Proposition A.4. We will prove that Statement (*) holds for all P . So suppose that P is a path of length r from $\bar{\omega}\mathbf{x}$ to $\bar{\omega}\mathbf{y}$.

If $r = 0$ then $\mathbf{x} = \mathbf{y}$ since \mathbf{X}_1 is monomorphic.

Suppose that $r > 0$ and that the result holds for all paths of length $< r$.

Let

$$\mathbf{x} = \theta \wedge x_1 \wedge \cdots \wedge x_j$$

and let

$$\mathbf{y} = \kappa \wedge y_1 \wedge \cdots \wedge y_j.$$

Let p be the largest number for which P begins with a path of the form

$$A_{1,\alpha_1}, B_1, A_{2,\alpha_2}, B_2, \dots, A_{p,\alpha_p}, B_p$$

where $\alpha_i(1) = |x_i| + 1$ for each i . Denote this path by P_1 (p is allowed to be 0 in which case P_1 is the empty path).

Lemmas A.11(i) and A.9 imply that $P \neq P_1$: if $|x_1| = 0$ this is because x_1 cannot equal $\bar{\omega}y_1$, and otherwise it's because $\theta^{[2]}(\beta_1^{[2]})^{-1} \dots (\beta_p^{2+\sum_{i<p}|x_i|})^{-1}(\rho_{1,\sum_{i\leq p}|x_i|}^{[1]})^{-1}$ does not take 1 to 1 and so cannot be equal to $\kappa^{[2]}$. Let D be the next operation in P and let P_2 be the part of P after D . There are three cases.

Case 1. Suppose that $D = A_{m,\gamma}$ for some γ , with $m \neq p + 1$.

If $m > p + 1$ then D commutes with all the earlier operations in P (if any), so P has the same effect as

$$D, P_1, P_2.$$

Since the length of P_1, P_2 is $r - 1$ we are done by Lemma A.15 (taking $Q_1 = D$ in that lemma).

If $m \leq p$, Lemma A.11(ii) shows that P_1, D has the same effect as

$$Q = (A_{1,\alpha'_1}, B_1, A_{2,\alpha'_2}, B_2, \dots, A_{p,\alpha'_p}, B_p)$$

where

$$\alpha'_i = \begin{cases} \alpha_i & \text{if } i \neq m, \\ \rho_{1,|x_m|}^{\gamma^{[2]}} \rho_{1,|x_m|}^{-1} \alpha_m & \text{if } i = m. \end{cases}$$

Then P has the same effect as Q, P_2 , which has length $r - 1$.

Case 2. Suppose that D is B_m or C_m for some m .

If $m > p$ then D commutes with all earlier operations in P (if any), so we are done by Lemma A.15 (taking $Q_1 = D$; note that if $p = 0$ and $|x_1| = 0$ then we cannot have $D = B_1$ because of the way p was chosen).

Let P'_1 be the standard path of length $3p$ starting at $\bar{\omega}\mathbf{x}$ and let Q be the path P'_1, D, P_2 . Lemma A.11(i) says that P_1 has the same effect as

$$A_{1,\beta_1^{[2]}}, A_{2,\beta_2}, \dots, A_{p,\beta_p}, P'_1$$

By Lemma A.14, there is a symbol \mathbf{z} such that the path

$$A_{1,\beta_1^{[2]}}, A_{2,\beta_2}, \dots, A_{p,\beta_p}$$

takes $\bar{\omega}\mathbf{x}$ to $\bar{\omega}\mathbf{z}$ and $\mathbf{x} \sim \mathbf{z}$. Then \mathbf{z} is Q -related to \mathbf{y} , and to complete Case 2 it suffices to show that $\mathbf{z} \sim \mathbf{y}$.

If D is B_m or C_m with $m < p$, Lemma A.9 shows that P'_1, D has the same effect as D, P'_1 . Hence Q has the same effect as D, P'_1, P_2 , and we are done by Lemma A.15 since P'_1, P_2 has length $< r$ (note that if $D = B_1$ in this situation then $|x_1|$ cannot be 0 since it would not be possible to apply B_1 to $P'_1(\mathbf{z})$).

If $D = C_p$ then P'_1, D has the same effect as

$$R = (A_{1,\rho_{1,|x_1|}}, B_1, A_{2,\rho_{1,|x_2|}}, B_2, \dots, A_{p,\rho_{1,|x_p|}})$$

since C_p and B_p are inverses of each other. Then Q has the same effect as R, P_2 , which has length $r - 2$.

If $D = B_p$, Lemma A.9 shows that P'_1, D has the same effect as $D, P'_1, A_{p+1,\gamma}$, where γ is the transposition (12). Hence Q has the same effect as $D, P'_1, A_{p+1,\gamma}, P_2$, and we are done by Lemma A.15 since $P'_1, A_{p+1,\gamma}, P_2$ has length $r - 1$ (this is why operations of type B count for 2 in the definition of length); note that we cannot have $D = B_1$ and $|x_1| = 0$ in this situation since p would be 1 and then P_1 would be B_1 , and it would not be possible to apply the sequence P_1, D to \mathbf{x} .

Case 3. Suppose that $D = A_{p+1,\gamma}$ for some γ .

First suppose $\gamma(1) = 1$, which implies $\gamma = \tilde{\gamma}^{[2]}$ for some $\tilde{\gamma}$. If $p = 0$ we're done by Lemma A.14. If $p > 0$ Lemmas A.11(i) and A.9 show that P_1, D has the same effect as $A_{p+1,\gamma'}, P_1$, so P has the same effect as $A_{p+1,\gamma'}, P_1, P_2$ and we're done by Lemma A.15.

For the rest of Case 3 we suppose that

$$(A.4) \quad \gamma(1) \neq 1.$$

Lemmas A.11(i) and A.9 show that P cannot equal P_1, D : if $|x_1| = 0$ this is because x_1 cannot equal $\bar{\omega}y_1$, and otherwise it's because the permutation

$$\theta^{[2]}(\beta_1^{[2]})^{-1} \dots (\beta_p^{2+\sum_{i < p} |x_i|})^{-1} (\rho_{1,\sum_{i \leq p} |x_i|}^{[1]})^{-1} (\gamma^{[1+\sum_{i \leq p} |x_i|]})^{-1}$$

does not take 1 to 1 and so cannot be equal to $\kappa^{[2]}$. Let E be the next operation in P and let P'_2 be the part of P after E ; then we have

$$(A.5) \quad P = (P_1, D, E, P'_2)$$

There are three cases to consider.

Case 3.1. $E = A_{m,\delta}$ for some δ .

If $m = p + 1$ we can combine D and E into a single operation, which gives a path of length $r - 1$ from \mathbf{x} to \mathbf{y} .

Otherwise we can commute E past D , which reduces to Case 1.

Case 3.2. $E = B_m$ for some m .

If $m \neq p, p + 1$ we can commute E past D , which reduces to Case 2.

If $m = p$ then D, E has the same effect as $B_m, A_{p+1,\gamma^{[2]}}$, which reduces to Case 2.

So suppose $m = p + 1$. Then

$$(A.6) \quad \gamma(1) \neq |x_{p+1}| + 1,$$

because of the way p was chosen. Denote $P_1(\mathbf{x})$ by \mathbf{z} , and let

$$(A.7) \quad \mathbf{z} = \lambda \wedge z_1 \wedge \cdots \wedge z_j.$$

Then

$$(A.8) \quad z_{p+1} = \bar{\omega}x_{p+1}$$

by Lemmas A.11 and A.9.

Because the operation B_{p+1} can be applied to $A_{p+1,\gamma}(\mathbf{z})$, we have that

$$(A.9) \quad \gamma z_{p+1} = \bar{\psi}w$$

for some w , and then

$$(A.10) \quad ED(\mathbf{z}) = \lambda(\gamma^{|z_1|+\cdots+|z_p|+1})^{-1} \wedge \cdots \wedge w \wedge \bar{\omega}z_{p+2} \cdots$$

Equations (A.6), (A.8) and (A.9) allow us to apply Remark A.6(iii) to get a $\delta \in \Sigma_{|x_{p+1}|}$ with the properties that $\delta^{-1}(|x_{p+1}|) = \gamma^{-1}(|x_{p+1}| + 1) - 1$ and

$$(A.11) \quad \delta x_{p+1} = \bar{\psi}v$$

for some v . Let

$$(A.12) \quad \varepsilon = \gamma(\delta^{[2]})^{-1}.$$

Then ε takes $|x_{p+1}| + 1$ to itself, so there is an $\tilde{\varepsilon} \in \Sigma_{|x_{p+1}|}$ with $\varepsilon = \tilde{\varepsilon}^{[1]}$.

We claim that the operation B_{p+1} can be applied to $A_{p+1,\delta^{[2]}}(\mathbf{z})$ and that $A_{p+1,\delta^{[2]}}, B_{p+1}, A_{p+1,\tilde{\varepsilon}}$ has the same effect on \mathbf{z} as D, E . Let us assume this for the moment. If $p = 0$ then (by Equation (A.5)) P has the same effect as $A_{1,\delta^{[2]}}, B_1, A_{1,\tilde{\varepsilon}}, P'_2$; since the length

of $A_{1,\varepsilon}, P'_2$ is $r - 2$ we're done by Lemma A.15 (note that $|x_1| \neq 0$ because of Equation A.4). If $p \neq 0$ then P has the same effect as $P_1, A_{p+1,\delta^{[2]}}, B_{p+1}, A_{p+1,\varepsilon}, P'_2$, which (by Lemmas A.11 and A.9) has the same effect as $A_{p+1,\delta}, P_1, B_{p+1}, A_{p+1,\varepsilon}, P'_2$. Case 2 applies to part of this path after the first operation, so we're done by Lemma A.15.

It remains to verify the claim. By Equations (A.8) and (A.11) and Remark A.6(ii) we have

$$\delta^{[2]}z_{p+1} = \bar{\omega}\delta x_{p+1} = \bar{\omega}\bar{\psi}v = \bar{\psi}\bar{\omega}v,$$

so (using Equation (A.7)) B_{p+1} can be applied to $A_{p+1,\delta^{[2]}}(\mathbf{z})$ and we have

$$\begin{aligned} & A_{p+1,\varepsilon}B_{p+1}A_{p+1,\delta^{[2]}}(\mathbf{z}) \\ &= \lambda((\delta^{[2]})^{-1}(\tilde{\varepsilon}^{[1]})^{-1})^{|z_1|+\dots+|z_p|+1} \wedge \dots \tilde{\varepsilon}\bar{\omega}v \wedge \bar{\omega}z_{p+2} \dots \end{aligned}$$

By Equation (A.10) it suffices to show $\tilde{\varepsilon}^{[1]}\delta^{[2]} = \gamma$ (which follows from Equation A.12) and $w = \tilde{\varepsilon}\bar{\omega}v$. Because \mathbf{X}_{p+1} is monomorphic, for the latter equation it suffices to show $\bar{\psi}w = \bar{\psi}\tilde{\varepsilon}^{[1]}\bar{\omega}v$, and this in turn follows from

$$\begin{aligned} \bar{\psi}w &= \gamma z_{p+1} = \gamma\bar{\omega}x_{p+1} = \varepsilon\delta^{[2]}\bar{\omega}x_{p+1} \\ &= \varepsilon\bar{\omega}\delta x_{p+1} = \varepsilon\bar{\omega}\bar{\psi}v = \varepsilon\bar{\psi}\bar{\omega}v = \bar{\psi}\tilde{\varepsilon}^{[1]}\bar{\omega}v, \end{aligned}$$

where we have used (in this order) Equations (A.9), (A.8), (A.12), Remark A.6(i), Equation A.11, and Remark A.6(ii).

Case 3.3. $E = C_m$ for some m .

If $m \neq p, p+1$ we can commute E past D , which reduces to Case 2.

If $m = p+1$ then D, E has the same effect as $C_m, A_{p+1,\gamma^{[1]}}$, which reduces to Case 2.

So suppose $m = p$ (which implies $p \neq 0$). Let P_0 be $A_{1,\alpha_1}, B_1, A_{2,\alpha_2}, B_2, \dots, A_{p-1,\alpha_p}, B_{p-1}$ if $p > 1$ and the empty path if $p = 1$; note that

$$(A.13) \quad P = (P_0, A_{p,\alpha_p}, B_p, D, E, P'_2).$$

We denote $A_{p,\alpha_p}P_0(\mathbf{x})$ by \mathbf{z} and let

$$(A.14) \quad \mathbf{z} = \lambda \wedge z_1 \wedge \dots \wedge z_j.$$

Because the operation B_p can be applied to \mathbf{z} , we have

$$(A.15) \quad z_p = \bar{\psi}w$$

for some w . Because C_p can be applied to $A_{p+1,\gamma}B_p\mathbf{z}$, we have

$$(A.16) \quad \gamma\bar{\omega}z_{p+1} = \bar{\omega}v$$

for some v , and

$$(A.17) \quad EDB_p(\mathbf{z}) = \lambda(\gamma^{-1})^{[|z_1|+\dots+|z_p|+1]} \wedge \dots \bar{\psi}w \wedge v \dots$$

Since \mathbf{X}_{p+1} is strongly monomorphic, Equations (A.4) and (A.16) imply that there is a $\delta \in \Sigma_{|x_{p+1}|}$ with the properties that $\delta^{-1}(1) = \gamma^{-1}(1) - 1$ and

$$(A.18) \quad \delta z_{p+1} = \bar{\omega}u$$

for some u .

Let ε be the transposition (12). Let

$$(A.19) \quad \eta = \gamma(\delta^{[2]})^{-1}(\varepsilon^{[1]})^{-1}.$$

Then η takes 1 to 1, so there is an $\tilde{\eta} \in \Sigma_{|x_{p+1}|}$ with $\eta = \tilde{\eta}^{[2]}$. We claim that the sequence of operations

$$A_{p+1,\delta}, C_p, A_{p,\varepsilon^{[|z_p|+1]}}, B_p, A_{p+1,\tilde{\eta}}$$

can be applied to \mathbf{z} and that it has the same effect as B_p, D, E . Assuming this for the moment, we see (using Equation (A.13)) that P has the same effect as

$$P_0, A_{p,\alpha_p}, A_{p+1,\delta}, C_p, A_{p,\varepsilon^{[|z_p|+1]}}, B_p, A_{p+1,\tilde{\eta}}, P'_2$$

which (by commuting $A_{p+1,\delta}$ past the earlier operations) has the same effect as

$$A_{p+1,\delta}, P_0, A_{p,\alpha_p}, C_p, A_{p,\varepsilon^{[|z_p|+1]}}, B_p, A_{p+1,\tilde{\eta}}, P'_2$$

which (moving C_p past A_{p,α_p}) has the same effect as

$$A_{p+1,\delta}, P_0, C_p, A_{p,\alpha_p^{[1]}}, A_{p,\varepsilon^{[|z_p|+1]}}, B_p, A_{p+1,\tilde{\eta}}, P'_2$$

which (since C_p commutes with the operations in P_0) has the same effect as

$$A_{p+1,\delta}, C_p, P_0, A_{p,\alpha_p^{[1]}}, A_{p,\varepsilon^{[|z_p|+1]}}, B_p, A_{p+1,\tilde{\eta}}, P'_2$$

The part of this path after the first two operations has length $r - 1$, so we're done by Lemma A.15.

It remains to verify the claim. Equations (A.14), (A.15) and (A.18) give

$$(A.20) \quad A_{p+1,\delta}(\mathbf{z}) = \lambda((\delta^{[2]})^{-1})^{[|z_1|+\dots+|z_p|+1]} \wedge \dots \psi w \wedge \bar{\omega}u \dots$$

so C_p can be applied to $A_{p+1,\delta}(\mathbf{z})$. Now $\varepsilon^{[|z_p|+1]}(\bar{\psi}\bar{\psi}w) = \bar{\psi}\bar{\psi}w$, so Equation (A.20) gives

$$(A.21) \quad \begin{aligned} A_{p,\varepsilon^{[|z_p|+1]}}C_p A_{p+1,\delta}(\mathbf{z}) \\ = \lambda((\delta^{[2]})^{-1}(\varepsilon^{[1]})^{-1})^{[|z_1|+\dots+|z_p|+1]} \wedge \dots \bar{\psi}\bar{\psi}w \wedge u \dots \end{aligned}$$

Then B_p can be applied to this and we have

$$\begin{aligned} A_{p+1, \tilde{\eta}} B_p A_{p, \varepsilon^{|z_p|+1}} C_p A_{p+1, \delta}(\mathbf{z}) \\ = \lambda((\delta^{[2]})^{-1}(\varepsilon^{[1]})^{-1}(\tilde{\eta}^{[2]})^{-1})^{|z_1|+\dots+|z_p|+1} \wedge \dots \bar{\psi} w \wedge \tilde{\eta} \bar{\omega} u \dots \end{aligned}$$

Comparing this to Equation (A.17), we see that it suffices to show $\gamma = \tilde{\eta}^{[2]} \varepsilon^{[1]} \delta^{[2]}$ (which follows from Equation (A.19)) and $\tilde{\eta} \bar{\omega} u = v$. Since \mathbf{X}_{p+1} is monomorphic, the latter equation follows from the equations

$$\bar{\omega} \tilde{\eta} \bar{\omega} u = \eta \bar{\omega} \bar{\omega} u = \gamma \bar{\omega} \delta^{-1} \bar{\omega} u = \gamma \bar{\omega} z_{p+1} = \bar{\omega} v$$

where we have used Remark A.6(i) and Equations A.19, A.18 and A.16.

This completes the proof of Proposition A.4 \square

Proof of Proposition A.2. First we need some notation to distinguish the action of Σ_j on $\mathbf{X}^{\wedge j}$ from the action of Σ_k on the k -th object of $\mathbf{X}^{\wedge j}$: given $\nu \in \Sigma_j$, $\eta \in \Sigma_k$ and a point \mathfrak{x} of $\mathbf{X}^{\wedge j}$ with $|\mathfrak{x}| = k$, we write \mathfrak{x}^ν (resp., $\eta \mathfrak{x}$) for the point obtained by applying ν (resp., η) to \mathfrak{x} .

Now suppose there is a point \mathfrak{x} in $\mathbf{X}^{\wedge j}$ which is not a basepoint and a nontrivial $\nu \in \Sigma_j$ such that $\mathfrak{x}^\nu = \mathfrak{x}$. Let \mathfrak{x} be represented by

$$\mathbf{x} = \theta \wedge x_1 \wedge \dots \wedge x_j$$

and note that $|x_i| > 0$ for all i by our hypothesis on X_0 . Then \mathfrak{x}^ν is represented by

$$\mathbf{y} = \theta \eta^{-1} \wedge x_{\nu^{-1}(1)} \wedge \dots \wedge x_{\nu^{-1}(j)},$$

where η permutes blocks of size $|x_1|, \dots, |x_j|$ in the same way that ν permutes $1, \dots, j$. There must be a path from \mathbf{x} to \mathbf{y} , and it cannot consist entirely of operations of type A, since η is not in $\Sigma_{|x_1|} \times \dots \times \Sigma_{|x_j|}$. Thus some x_i is in the image of $\bar{\omega}$, and this implies (using operations of type A and C) that there is a point \mathfrak{w} of $\mathbf{X}^{\wedge j}$ and a $\zeta \in \Sigma_{|\mathfrak{x}|}$ with $\mathfrak{x} = \zeta \bar{\omega} \mathfrak{w}$. Now ν commutes with both ζ and $\bar{\omega}$, so the fact that $\mathfrak{x}^\nu = \mathfrak{x}$ implies that

$$\zeta \bar{\omega}(\mathfrak{w}^\nu) = \zeta \bar{\omega} \mathfrak{w}.$$

Using Proposition A.4 we see that $\mathfrak{w}^\nu = \mathfrak{w}$, and thus \mathfrak{w} is a nontrivial fixed point of the Σ_j action with $|\mathfrak{w}| < |\mathfrak{x}|$. Continuing in this way would give a nontrivial fixed point in the 0-th object of $\mathbf{X}^{\wedge j}$, which is impossible by our hypothesis on X_0 . \square

We conclude with an example which shows that Propositions A.2 and A.4 fail if “strongly monomorphic” is replaced by “monomorphic.”

Recall Definition 4.10. Let \mathbf{X} be the subobject of \mathbf{S} with $X_0 = *$ and $X_k = S^k$ for $k > 0$.

We will denote the point of $\mathbf{X} \wedge \mathbf{X}$ represented by a symbol $\theta \wedge x \wedge y$ by $[\theta \wedge x \wedge y]$. Let x be the nontrivial simplex in X_1 . Note that \mathbf{X} is

not strongly monomorphic because the transposition (12) takes $\bar{\omega}x$ to itself, so $[(12)\bar{\omega}x]$ is in the image of $\bar{\omega}$, but x is not in the image of $\bar{\omega}$.

Now we claim that

$$\tau([\bar{\omega}x \wedge x]) = [\bar{\omega}x \wedge x],$$

which gives a counterexample for Proposition A.2. We have

$$\begin{aligned} \tau([\bar{\omega}x \wedge x]) &= [\rho_{1,2} \wedge x \wedge \bar{\omega}x] = [(23)(12) \wedge x \wedge \bar{\omega}x] = [(23)(12) \wedge \bar{\psi}x \wedge x] \\ &= [(23) \wedge \bar{\psi}x \wedge x] \text{ because (12) acts trivially on } X_2 \\ &= [(23) \wedge x \wedge \bar{\omega}x] = [x \wedge \bar{\omega}x] = [\bar{\psi}x \wedge x] = [\bar{\omega}x \wedge x]. \end{aligned}$$

Next we observe that $[(12) \wedge x \wedge x] \neq [x \wedge x]$, because there is no nontrivial path beginning at $(12) \wedge x \wedge x$. But we claim that

$$\bar{\omega}([(12) \wedge x \wedge x]) = \bar{\omega}([x \wedge x]),$$

which gives a counterexample for Proposition A.4. We have

$$\begin{aligned} \bar{\omega}([(12) \wedge x \wedge x]) &= [(23) \wedge \bar{\omega}x \wedge x] = [(23) \wedge \bar{\psi}x \wedge x] \\ &= [(23) \wedge x \wedge \bar{\omega}x] = [x \wedge \bar{\omega}x] = [\bar{\psi}x \wedge x] \\ &= [\bar{\omega}x \wedge x] = \bar{\omega}([x \wedge x]). \end{aligned}$$

APPENDIX B. THE PROOF OF LEMMA 11.3

Let \mathbf{W} denote the multiseisimplicial spectrum whose k -th object is $S_{\bullet}^{k-\text{multi}, \bar{\rho}}(T(\text{STop}(k)))$, so that $|\mathbf{W}| = \mathbf{Y}$. We begin by showing that the monad \mathbb{P} of Definition 17.2 acts on the pair $(\mathbf{W}, \mathbf{R}_{\text{STop}})$.

Let us define a \mathbb{Z} -graded category \mathcal{B} as follows. The objects of \mathcal{B} are pairs

$$(g : \Delta^n \rightarrow T(\text{STop}(k)), o),$$

where both \mathbf{n} and k are allowed to vary and o is an orientation of Δ^n ; the grading is given by $d(g, o) = \dim(\Delta^n) - k$. The morphisms are commutative diagrams

$$\begin{array}{ccc} \Delta^n & \xrightarrow{g} & T(\text{STop}(k)) \\ \downarrow \phi & & \downarrow \alpha \\ \Delta^{n'} & \xrightarrow{g'} & T(\text{STop}(k)) \end{array}$$

in which ϕ is a composite of coface maps and permutations of the factors and α is a permutation; we require ϕ to be orientation preserving if the dimensions are equal. \mathcal{B} is a symmetric monoidal \mathbb{Z} -graded category with product \square , where

$$(g, o) \square (g', o')$$

is the pair consisting of the composite

$$\Delta^n \times \Delta^{n'} \xrightarrow{g \times g'} T(\text{STop}(k)) \times T(\text{STop}(k')) \rightarrow T(\text{STop}(k + k'))$$

and the orientation $o \times o'$. The symmetry isomorphism γ is

$$\begin{array}{ccc} \Delta^n \times \Delta^{n'} & \xrightarrow{g \times g'} & T(\text{STop}(k)) \times T(\text{STop}(k')) \longrightarrow T(\text{STop}(k + k')) \\ \downarrow \phi & & \downarrow \alpha \\ \Delta^{n'} \times \Delta^n & \xrightarrow{g' \times g} & T(\text{STop}(k')) \times T(\text{STop}(k)) \longrightarrow T(\text{STop}(k' + k)) \end{array}$$

where ϕ and α are the evident permutations.

In the construction of Definition 15.4, if we replace $\mathcal{A}_{e,*,1}$ by \mathcal{B} , $\mathcal{A}_{\text{rel}}^{\mathbb{Z}}$ by $\mathcal{A}_{\text{STop}}$, \boxtimes by \square and sig_{rel} by Inv we obtain a functor

$$\mathbf{d}_{\blacksquare} : \mathcal{A}_1 \times \cdots \times \mathcal{A}_j \rightarrow \mathcal{A}_{\text{STop}},$$

for each datum \mathbf{d} , where \mathcal{A}_i denotes \mathcal{B} if $r(i) = u$ and $\mathcal{A}_{\text{STop}}$ if $r(i) = v$.

Next we have the analogue of Definition 16.5.

Definition B.1. Let k_1, \dots, k_j be non-negative integers and let \mathbf{n}_i be a k_i -fold multi-index for $1 \leq i \leq j$. Let $r : \{1, \dots, j\} \rightarrow \{u, v\}$, and for $1 \leq i \leq j$ let \mathbf{Z}_i denote \mathbf{W} if $r(i) = u$ and \mathbf{R}_{STop} if $r(i) = v$. Then for each map of preorders

$$a : U(\Delta^{\mathbf{n}_1} \times \cdots \times \Delta^{\mathbf{n}_j}) \rightarrow P_{r,v}$$

we define

$$a_* : ((\mathbf{Z}_1)_{k_1})_{\mathbf{n}_1} \times \cdots \times ((\mathbf{Z}_j)_{k_j})_{\mathbf{n}_j} \rightarrow ((\mathbf{R}_{\text{STop}})_{k_1 + \cdots + k_j})_{(\mathbf{n}_1, \dots, \mathbf{n}_j)}$$

by

$$\begin{aligned} a_*(z_1, \dots, z_j)(\sigma_1 \times \cdots \times \sigma_j, o_1 \times \cdots \times o_j) \\ = i^{\epsilon(\zeta)} a(\sigma_1 \times \cdots \times \sigma_j)_{\blacksquare}(z_1(\sigma_1, o_1), \dots, z_j(\sigma_j, o_j)), \end{aligned}$$

where

- if $r(i) = u$ then $z_i(\sigma_i, o_i)$ denotes $(z_i|_{\sigma_i}, o_i)$, and
- ζ is the block permutation that takes blocks $\mathbf{b}_1, \dots, \mathbf{b}_j, \mathbf{c}_1, \dots, \mathbf{c}_j$ of size $k_1, \dots, k_j, \dim \sigma_1, \dots, \dim \sigma_j$ into the order $\mathbf{b}_1, \mathbf{c}_1, \dots, \mathbf{b}_j, \mathbf{c}_j$.

As in Section 17, this definition leads to a map

$$(B.1) \quad \mathbb{P}_2(\mathbf{W}, \mathbf{R}_{\text{STop}}) \rightarrow \mathbf{R}_{\text{STop}}.$$

Since \mathbf{W} is a commutative multiseisimplicial symmetric ring spectrum, we have a map

$$(B.2) \quad \mathbb{P}_1(\mathbf{W}) \xrightarrow{\Xi_1} \bigvee_{j \geq 0} \mathbf{W}^{\wedge j} / \Sigma_j \rightarrow \mathbf{W},$$

where Ξ_1 is given in Definition 18.1(ii).

The maps (B.1) and (B.2) give the required action of \mathbb{P} on $(\mathbf{W}, \mathbf{R}_{\text{STop}})$. Now the proof of Theorem 1.3 (given in Section 18) gives a map of commutative symmetric ring spectra

$$|B_{\bullet}(\mathbb{P}'_1, \mathbb{P}, (\mathbf{W}, \mathbf{R}_{\text{STop}}))| \rightarrow |B_{\bullet}(\mathbb{P}'_2, \mathbb{P}, (\mathbf{W}, \mathbf{R}_{\text{STop}}))|$$

which is a weak equivalence by Lemma 11.1. As in Remark 18.3, there is a weak equivalence of commutative symmetric ring spectra

$$(\mathbf{M}_{\text{STop}})^{\text{comm}} \rightarrow |B_{\bullet}(\mathbb{P}'_2, \mathbb{P}, (\mathbf{W}, \mathbf{R}_{\text{STop}}))|.$$

To complete the proof, we observe that there is a weak equivalence of commutative symmetric ring spectra

$$|B_{\bullet}(\mathbb{P}'_1, \mathbb{P}, (\mathbf{W}, \mathbf{R}_{\text{STop}}))| = |B_{\bullet}(\mathbb{P}'_1, \mathbb{P}_1, \mathbf{W})| \rightarrow |B_{\bullet}(\mathbb{P}'_1, \mathbb{P}'_1, \mathbf{W})| \rightarrow |\mathbf{W}| = \mathbf{Y},$$

where the first arrow is a weak equivalence by Proposition 18.2(iii) and the second by [May72, Proposition 9.8 and Corollary 11.9].

APPENDIX C. A FUNCTORIAL VERSION OF ad_{rel}^R

In [LM14, Section 13] we explained why ad^R (as defined in [LM14, Section 9], which is the definition we have used in the present paper) is not a functor of R , and how to modify the definition of ad^R to make it a functor (unfortunately, in [LM14] we also denoted the modified version by ad^R ; in this appendix we will be more careful with the notation). Our goal in this appendix is to give a functorial version of ad_{rel}^R , which is needed in [BLM19] (but not in the present paper). The reader should look at [LM14, Section 13] before continuing.

Unfortunately, it seems that we cannot just adapt the method of [LM14, Section 13] to this situation, because ad_{rel}^R isn't even "approximately" functorial: given a ring homomorphism $R \rightarrow S$ and an object (C, D, β, φ) of $\mathcal{A}_{\text{rel}}^R$, there does not seem to be a reasonable way to create an object $(C', D', \beta', \varphi')$ of $\mathcal{A}_{\text{rel}}^S$ from this data (we could let $C' = S \otimes_R C$, but there's no reasonable candidate for D'). So in Subsection C.2 (after some preliminary motivation in Subsection C.1) we give a variant of ad_{rel}^R , which we denote by ad_{Rel}^R , and in Subsection C.3 we show that ad_{Rel}^R is approximately functorial (i.e., functorial up to isomorphism). In Subsection C.4 we show that the ad theories ad^R and ad_{Rel}^R are equivalent, that is, there is a morphism of ad theories from ad_{Rel}^R to ad_{rel}^R which induces an isomorphism of bordism groups. In Subsection C.5 we give an enhanced version of the material in [LM14, Section 13], and in Subsection C.6 we use this to create a variant of ad_{Rel}^R , which we denote by $\text{ad}_{\text{Rel, sch}}^R$. In Subsection C.7 we show (using Subsection C.3) that $\text{ad}_{\text{Rel, sch}}^R$ is a functor of R .

Remark C.1. We could have used ad_{Rel}^R throughout this paper instead of ad_{rel}^R , but that would have added extra complexity with no benefit, since functoriality is not an issue in this paper.

C.1. Background. As motivation for the definition of ad_{Rel}^R , we need

Lemma C.2. *Let M be a right R module and N a left R module, and let $R^{\text{op}} \otimes R$ act on R on the right in the usual way. Then the map*

$$a : R \otimes_{R^{\text{op}} \otimes R} (M \otimes N) \rightarrow M \otimes_R N$$

given by $a(r \otimes m \otimes n) = mr \otimes n$ is an isomorphism.

Proof. This follows immediately from the isomorphisms

$$R \otimes_{R^{\text{op}} \otimes R} (M \otimes N) \cong M \otimes_R R \otimes_R N \cong M \otimes_R N.$$

□

Now fix a ring R with involution.

Definition C.3. Let P and Q be left $R^{\text{op}} \otimes R$ modules. A map

$$b : P \rightarrow Q$$

is *quasi-linear* if $b((r \otimes s)p) = (\bar{s} \otimes \bar{r})b(p)$.

Lemma C.4. *Let $b : P \rightarrow Q$ be a quasi-linear map of left $R^{\text{op}} \otimes R$ modules. Then the map*

$$\hat{b} : R \otimes_{R^{\text{op}} \otimes R} P \rightarrow R \otimes_{R^{\text{op}} \otimes R} Q$$

given by $\hat{b}(r \otimes p) = \bar{r} \otimes b(p)$ is well-defined. □

C.2. The ad theory ad_{Rel}^R . Recall the definition of homotopy finite ([LM14, Definition 9.2(iv)]).

Definition C.5. A *Relaxed quasi-symmetric complex of dimension n* is a quadruple (C, E, γ, ϕ) , where C is a homotopy finite² chain complex over R , E is a homotopy finite chain complex over $R^{\text{op}} \otimes R$ with a $\mathbb{Z}/2$ action for which the generator acts quasi-linearly, γ is a $\mathbb{Z}/2$ equivariant $R^{\text{op}} \otimes R$ -linear quasi-isomorphism $C^t \otimes C \rightarrow E$, and ϕ is an n -dimensional element of $(R \otimes_{R^{\text{op}} \otimes R} E)^{\mathbb{Z}/2}$ (where the $\mathbb{Z}/2$ action is given by Lemma C.4).

For the following example, note that if A is a left $R^{\text{op}} \otimes R$ module which is nonzero in only finitely many dimensions then A^W is (additively) a direct sum of copies of A , and hence the natural map

$$R \otimes_{R^{\text{op}} \otimes R} (A^W) \rightarrow (R \otimes_{R^{\text{op}} \otimes R} A)^W$$

²In [LM14, Section 9] we also required C to be free, but that turns out not to be necessary; see Remark C.18.

is an isomorphism because tensor product preserves direct sums.

Example C.6. If (C, φ) is a quasi-symmetric complex as defined in [LM14, Definition 9.3], and if C is nonzero in only finitely many dimensions, then the quadruple $(C, (C^t \otimes C)^W, \gamma, \phi)$ is a Relaxed quasi-symmetric complex, where $\gamma : C^t \otimes C \rightarrow (C^t \otimes C)^W$ is induced by the augmentation $W \rightarrow \mathbb{Z}$ and ϕ is the image of φ under the composite $((C^t \otimes_R C)^W)^{\mathbb{Z}/2} \cong ((R \otimes_{R^{\text{op}}} \otimes_R (C^t \otimes C))^W)^{\mathbb{Z}/2} \cong (R \otimes_{R^{\text{op}}} \otimes_R ((C^t \otimes C)^W))^{\mathbb{Z}/2}$ (where the first isomorphism is Lemma C.2).

Definition C.7. We define a category $\mathcal{A}_{\text{Rel}}^R$ as follows. The objects of $\mathcal{A}_{\text{Rel}}^R$ are the Relaxed quasi-symmetric complexes. A morphism $(C, E, \gamma, \phi) \rightarrow (C', E', \gamma', \phi')$ is a pair $(f : C \rightarrow C', g : E \rightarrow E')$, where f is an R -linear chain map and g is a $\mathbb{Z}/2$ equivariant $R^{\text{op}} \otimes R$ -linear chain map, such that $g\gamma = \gamma'(f \otimes f)$, and (if $\dim \phi = \dim \phi'$) $(1 \otimes g)_*(\phi) = \phi'$.

$\mathcal{A}_{\text{Rel}}^R$ is a balanced ([LM14, Definition 5.1]) \mathbb{Z} -graded category, where i takes (C, E, γ, ϕ) to $(C, E, \gamma, -\phi)$ and \emptyset_n is the n -dimensional object for which C and E are zero in all degrees.

Remark C.8. There is a morphism $\mathcal{A}_{\text{Rel}}^R \rightarrow \mathcal{A}_{\text{rel}}^R$ of \mathbb{Z} -graded categories which takes (C, E, γ, ϕ) to $(C, R \otimes_{R^{\text{op}}} \otimes_R E, \beta, \phi)$, where β is the composite

$$C^t \otimes_R C \cong R \otimes_{R^{\text{op}}} \otimes_R (C^t \otimes C) \xrightarrow{1 \otimes \gamma} R \otimes_{R^{\text{op}}} \otimes_R E.$$

Remark C.9. Let $\mathcal{A}_{\text{fin}}^R$ be the full subcategory of \mathcal{A}^R consisting of objects (C, φ) with C finite (not just homotopy finite). Let $\mathcal{A}_{\text{Rel,fin}}^R$ be the full subcategory of $\mathcal{A}_{\text{Rel}}^R$ consisting of objects (C, E, γ, ϕ) with C and E finite. The construction of Example C.6 gives a morphism

$$\mathcal{A}_{\text{fin}}^R \rightarrow \mathcal{A}_{\text{Rel,fin}}^R$$

of \mathbb{Z} -graded categories.

Next we must say what the K -ads with values in $\mathcal{A}_{\text{Rel}}^R$ are. For a balanced pre K -ad F we will use the notation

$$F(\sigma, o) = (C_\sigma, E_\sigma, \gamma_\sigma, \phi_{\sigma, o}).$$

Recall [LM14, Definition 9.7].

Definition C.10. A balanced pre K -ad F is *well-behaved* if C and E are well-behaved.

Definition C.11. (i) A *balanced K -ad* is a pre K -ad with the following properties.

- (a) it is balanced, closed, and well-behaved, and
 - (b) the composite of F with the morphism of Remark C.8 satisfies part (i)(b) of Definition 12.10
- (ii) A K -ad is a pre K -ad which is naturally isomorphic to a balanced K -ad.

We write $\text{ad}_{\text{Rel}}^R(K)$ for the set of K -ads with values in $\mathcal{A}_{\text{Rel}}^R$.

Theorem C.12. ad_{Rel}^R is an ad theory.

This follows from the proof of Theorem 12.12 with minor changes.

Remark C.13. The morphisms of Remarks C.8 and C.9 take ads to ads.

C.3. ad_{Rel}^R is **approximately functorial**. The result in this subsection will be used in Subsection C.7.

Definition C.14. Let $h : R \rightarrow S$ be a ring homomorphism. Define a functor

$$h_{\text{Rel}} : \mathcal{A}_{\text{Rel}}^R \rightarrow \mathcal{A}_{\text{Rel}}^S$$

as follows. For an object (C, E, γ, ϕ) of $\mathcal{A}_{\text{Rel}}^R$, let

$$h_{\text{Rel}}(C, E, \gamma, \phi) = (S \otimes_R C, (S^{\text{op}} \otimes S) \otimes_{R^{\text{op}} \otimes R} E, \gamma', \phi'),$$

where γ' is the composite

$$(S \otimes_R C)^t \otimes (S \otimes_R C) \cong (S^{\text{op}} \otimes S) \otimes_{R^{\text{op}} \otimes R} (C^t \otimes C) \xrightarrow{1 \otimes \gamma} (S^{\text{op}} \otimes S) \otimes_{R^{\text{op}} \otimes R} E$$

(which is a quasi-isomorphism by the Künneth spectral sequence [Wei94, Theorem 5.6.4], using the fact that C and E are homotopy finite) and ϕ' is the image of ϕ under the composite

$$(R \otimes_{R^{\text{op}} \otimes R} E)^{\mathbb{Z}/2} \rightarrow (S \otimes_{R^{\text{op}} \otimes R} E)^{\mathbb{Z}/2} \cong (S \otimes_{S^{\text{op}} \otimes S} (S^{\text{op}} \otimes S) \otimes_{R^{\text{op}} \otimes R} E)^{\mathbb{Z}/2}.$$

The reader can check that if $k : S \rightarrow T$ is another ring homomorphism then $(kh)_{\text{Rel}}(C, E, \gamma, \phi)$ is isomorphic to but not equal to $k_{\text{Rel}} h_{\text{Rel}}(C, E, \gamma, \phi)$.

Proposition C.15. h_{Rel} takes ads to ads.

Proof. Let $F \in \text{ad}_{\text{Rel}}^R(K)$. We may assume that F is balanced. Write

$$F(\sigma, o) = (C_\sigma, E_\sigma, \gamma_\sigma, \phi_{\sigma, o}).$$

Let Ψ be the functor of Remark C.8. Let $G = \Psi \circ F$; then G is an element of $\text{ad}_{\text{rel}}^R(K)$. Write

$$G(\sigma, o) = (C_\sigma, D_\sigma, \beta_\sigma, \phi_{\sigma, o}).$$

Let $H = \Psi \circ h_{\text{Rel}} \circ F$; we need to show that H is an ad. It's immediate that H is balanced, well-behaved and closed, so it only remains to show that it satisfies part (ii) of Definition 12.10. Write

$$H(\sigma, o) = (S \otimes_R C_\sigma, D_\sigma^S, \beta_\sigma^S, \psi_{\sigma, o})$$

and fix an oriented cell (σ, o) of K . Recall Notation 12.9 and let

$$\begin{aligned} k_\sigma : ((S \otimes_R C)^t \otimes_S (S \otimes_R C))_\sigma / ((S \otimes_R C)^t \otimes_S (S \otimes_R C))_{\partial\sigma} \\ \rightarrow ((S \otimes_R C_\sigma) / (S \otimes_R C_{\partial\sigma})) \otimes_S (S \otimes_R C_\sigma) \end{aligned}$$

be the analogous map. We need to show that the slant product with $(k_\sigma)_*(\beta_\sigma^S)^{-1}([\psi_{\sigma, o}])$ is an isomorphism

$$H^*(\text{Hom}_S(S \otimes_R C_\sigma, S)) \rightarrow H_{\dim \sigma - \deg F - *} (S \otimes_R C_\sigma / S \otimes_R C_{\partial\sigma}).$$

First we observe that (using the definition of h_{Rel}) the image of $(j_\sigma)_*(\beta_\sigma^S)^{-1}([\phi_{\sigma, o}])$ in $H_*((S \otimes_R C_\sigma) / (S \otimes_R C)_{\partial\sigma}) \otimes_S S \otimes_R C_\sigma$ is $(k_\sigma)_*(\beta_\sigma^S)^{-1}([\psi_{\sigma, o}])$. Now the desired isomorphism follows from our next lemma. □

Lemma C.16. *Let $h : R \rightarrow S$ be a homomorphism of rings with involution. Let A and B be homotopy finite chain complexes over R . Let x be a cycle in $A^t \otimes_R B$ with the property that the slant product with x is an isomorphism*

$$H^*(\text{Hom}_R(B, R)) \rightarrow H_{\dim x - *} A^t.$$

Let y be the image of x under the map

$$A^t \otimes_R B \rightarrow (S \otimes_R A)^t \otimes_S (S \otimes_R B).$$

Then the slant product with y is an isomorphism

$$H^*(\text{Hom}_S(S \otimes_R B, S)) \rightarrow H_{\dim x - *} (S \otimes_R A)^t.$$

Proof of Lemma C.16. By naturality of the slant product, we may assume that A and B are finite. Because B is additively a direct sum of finitely many copies of R , the map

$$\Upsilon : \text{Hom}_R(B, R) \otimes_R S \rightarrow \text{Hom}_S(S \otimes_R B, S)$$

defined by $\Upsilon(f \otimes s)(t \otimes b) = tf(b)s$ is an isomorphism.

Consider the diagram

$$\begin{array}{ccc} H^*(\text{Hom}_S(S \otimes_R B, S)) & \xrightarrow{\text{slant } y} & H_{\dim x - *} (S \otimes_R A)^t \\ \Upsilon \uparrow \cong & & \uparrow \cong \\ H^*(\text{Hom}_R(B, R) \otimes_R S) & \longrightarrow & H_{\dim x - *} (A^t \otimes_R S) \end{array}$$

where the bottom arrow is induced by the chain map $(\setminus x) \otimes 1$. It's straightforward to check that the diagram commutes, and the lower arrow is an isomorphism by the Künneth spectral sequence [Wei94, Theorem 5.6.4]. \square

C.4. Comparison of ad_{Rel}^R and ad_{rel}^R .

Proposition C.17. *The morphism $\text{ad}_{\text{Rel}}^R \rightarrow \text{ad}_{\text{rel}}^R$ induces an isomorphism of bordism groups.*

Proof. Let $\mathcal{A}_{\text{Rel,fin}}^R$ be the full subcategory of $\mathcal{A}_{\text{Rel}}^R$ consisting of objects (C, E, γ, ϕ) for which C is finite. Consider the diagram

$$\begin{array}{ccc} \mathcal{A}_{\text{Rel}}^R & \longrightarrow & \mathcal{A}_{\text{rel}}^R \\ \uparrow a & & \uparrow d \\ \mathcal{A}_{\text{Rel,fin}}^R & & \mathcal{A}^R \\ \uparrow b & \xrightarrow{c} & \\ \mathcal{A}_{\text{fin}}^R & & \end{array}$$

where a is induced by the inclusion of categories, b is given by Remark C.9, c is induced by the inclusion of categories, and d is given by Remark 12.5. This diagram commutes up to natural isomorphism, so it induces a commutative diagram of bordism groups:

$$\begin{array}{ccc} (\Omega_{\text{Rel}}^R)_* & \longrightarrow & (\Omega_{\text{rel}}^R)_* \\ \uparrow \Omega_*^a & & \uparrow \Omega_*^d \\ (\Omega_{\text{Rel,fin}}^R)_* & & \mathcal{A}^R \\ \uparrow \Omega_*^b & \xrightarrow{\Omega_*^c} & \\ (\Omega_{\text{fin}}^R)_* & & \end{array}$$

Ω_*^d is an isomorphism by Proposition 13.3, and the proof of that proposition, with minor modifications, shows that Ω_*^b is an isomorphism.

To see that Ω_*^a is onto, let (C, E, γ, ϕ) represent an element of $\text{ad}_{\text{Rel}}^R(*)$. Since C is homotopy finite, there is a chain homotopy equivalence $f : C' \rightarrow C$ with C' finite. Let $\gamma' = \gamma \circ (f \otimes f)$. Then (C', E, γ', ϕ) is an element of $\text{ad}_{\text{Rel,fin}}^R(*)$, and (f, id) is a morphism from (C', E, γ', ϕ) to (C, E, γ, ϕ) . Let F be the cylinder object of (C, E, γ, ϕ) ([LM14, Definition 3.10(g)]). Let $0, 1, \iota$ denote the three cells of the unit interval

I , with their standard orientations. Replacing $F(0)$ with (C', E, γ', ϕ) gives an I -ad which is a bordism between (C, E, γ, ϕ) and (C', E, γ', ϕ) .

To complete the proof it suffices to show that Ω_*^c is an isomorphism, since this will imply that Ω_*^a is a monomorphism.

To see that Ω_*^c is onto, let (C, φ) represent an element of $\text{ad}^R(*)$. There is a chain homotopy equivalence $f : C \rightarrow C'$ with C' finite. Then $(C', (f \otimes f) \circ \varphi)$ represents an element of $\text{ad}_{\text{fin}}^R(*)$, and (f, ϕ) is a morphism from (C, φ) to $(C', (f \otimes f) \circ \varphi)$. Let F be the cylinder object of $(C', (f \otimes f) \circ \varphi)$. Replacing $F(0)$ with (C, φ) gives a bordism between (C, φ) and $(C', (f \otimes f) \circ \varphi)$.

To see that Ω_*^c is a monomorphism, let $F \in \text{ad}^R(I)$ with $F(0)$ and $F(1)$ in $\mathcal{A}_{\text{fin}}^R$. Write

$$F(0) = (C_0, \varphi_0), \quad F(1) = (C_1, \varphi_1), \quad F(\iota) = (C_\iota, \varphi_\iota).$$

Since C_ι is homotopy finite, there is a chain homotopy equivalence $f : C_\iota \rightarrow B$ with B finite. Let g_0 be the composite $C_0 \rightarrow C_\iota \xrightarrow{f} B$ and similarly for $g_1 : C_1 \rightarrow B$. Let $\text{cl}(I)$ be the cellular chain complex of I and let j_0 (resp., j_1) be the composite $\mathbb{Z} \cong \text{cl}(0) \rightarrow \text{cl}(I)$ (resp., $\mathbb{Z} \cong \text{cl}(1) \rightarrow \text{cl}(I)$), where the second map is the inclusion. Let B' be the colimit

$$\begin{array}{ccccc} & & C_0 & & C_1 & & \\ & \swarrow & & \searrow & \swarrow & \searrow & \\ & j_1 \otimes \text{id} & & g_0 & g_1 & & j_1 \otimes \text{id} \\ & & \text{cl}(I) \otimes C_0 & & B & & \text{cl}(I) \otimes C_1 \end{array}$$

The composites $C_0 \xrightarrow{j_0 \otimes \text{id}} \text{cl}(I) \otimes C_0 \rightarrow B'$ and $C_1 \xrightarrow{j_0 \otimes \text{id}} \text{cl}(I) \otimes C_1 \rightarrow B'$ are strong monomorphisms ([LM14, Definition 9.6]). Let n be the degree of F and let $\psi \in B'_{1-n}$ be the image of $\iota \otimes \varphi_0 + f_*(\phi_\iota) - \iota \otimes \varphi_1$.

Define an I -ad G by

$$G(0) = F(0), \quad G(1) = F(1), \quad G(\iota) = (B', \psi).$$

Then G is the desired bordism. \square

Remark C.18. The proof that Ω_*^c is an isomorphism also shows that the requirement in [LM14, Section 9] that C should be free over R is not needed. That is, if we define ad^R as in [LM14, Section 9] and $(\text{ad}^R)^\dagger$ by requiring only that C be homotopy finite then the forgetful map $\text{ad}^R \rightarrow (\text{ad}^R)^\dagger$ induces an isomorphism of bordism groups.

C.5. Enhanced version of [LM14, Section 13]. In [LM14, Section 13] we gave a model for the category of free R modules which is functorial in R . In this subsection give a similar model for the category of all R

modules. Our terminology and notation will be different from [LM14, Section 13].

We define the category of *schematic free R modules* as follows. An object is a set \mathbb{M} . This should be thought of as representing the free R module generated by \mathbb{M} , which we denote by $R\langle\mathbb{M}\rangle$. We define a map $\mathbb{M} \rightarrow \mathbb{M}'$ to be a map of R -modules $R\langle\mathbb{M}\rangle \rightarrow R\langle\mathbb{M}'\rangle$.

We define the category of *schematic R modules* as follows. An object of this category is a triple $(\mathbb{M}, \mathbb{N}, T)$, where \mathbb{M} and \mathbb{N} are schematic free R modules and T is a map $\mathbb{N} \rightarrow \mathbb{M}$. Such a triple should be thought of as representing the quotient of $R\langle\mathbb{M}\rangle$ by the image of T ; we write $R\langle(\mathbb{M}, \mathbb{N}, T)\rangle$ for this quotient. A map $(\mathbb{M}, \mathbb{N}, T) \rightarrow (\mathbb{M}', \mathbb{N}', T')$ is defined to be an R -module map $R\langle(\mathbb{M}, \mathbb{N}, T)\rangle \rightarrow R\langle(\mathbb{M}', \mathbb{N}', T')\rangle$.

Lemma C.19. *The functor from schematic R modules to R modules which takes $(\mathbb{M}, \mathbb{N}, T)$ to $R\langle(\mathbb{M}, \mathbb{N}, T)\rangle$ is an equivalence of categories.*

Proof. The functor is the identity on morphism sets, so it's only necessary to show that every R module P is isomorphic to one of the form $R\langle(\mathbb{M}, \mathbb{N}, T)\rangle$. Choose an exact sequence $Q_1 \rightarrow Q_2 \rightarrow P \rightarrow 0$ where Q_1 and Q_2 are free, let \mathbb{M} and \mathbb{N} be bases for Q_1 and Q_2 , and let T be the map induced by $Q_1 \rightarrow Q_2$. \square

A *schematic chain complex* \mathbb{C} over R is a sequence of schematic R modules and maps, and we write $R\langle\mathbb{C}\rangle$ for the corresponding sequence of R modules and maps. A map $\mathbb{C} \rightarrow \mathbb{C}'$ of schematic chain complexes is a map of R chain complexes $R\langle\mathbb{C}\rangle \rightarrow R\langle\mathbb{C}'\rangle$.

Let $h : R_1 \rightarrow R_2$ be a homomorphism. For a schematic free R_1 module \mathbb{M} we write $h_{\text{sch}}\mathbb{M}$ for \mathbb{M} thought of as a schematic free R_2 module. There is a canonical isomorphism

$$(C.1) \quad R_2\langle h_{\text{sch}}\mathbb{M}\rangle \cong R_2 \otimes_{R_1} R_1\langle\mathbb{M}\rangle$$

which takes an element m of \mathbb{M} to $1 \otimes m$. For a map $T : \mathbb{M} \rightarrow \mathbb{N}$ we write $h_{\text{sch}}T$ for the map $h_{\text{sch}}\mathbb{M} \rightarrow h_{\text{sch}}\mathbb{N}$ defined by the following diagram.

$$\begin{array}{ccc} R_2\langle h_{\text{sch}}\mathbb{M}\rangle & \xrightarrow{h_{\text{sch}}T} & R_2\langle h_{\text{sch}}\mathbb{N}\rangle \\ \cong \downarrow & & \cong \downarrow \\ R_2 \otimes_{R_1} R_1\langle\mathbb{M}\rangle & \xrightarrow{1 \otimes T} & R_2 \otimes_{R_1} R_1\langle\mathbb{N}\rangle \end{array}$$

For a schematic R_1 module $(\mathbb{M}, \mathbb{N}, T)$, we define $h_{\text{sch}}(\mathbb{M}, \mathbb{N}, T)$ to be the schematic R_2 module $(h_{\text{sch}}\mathbb{M}, h_{\text{sch}}\mathbb{N}, h_{\text{sch}}T)$. The isomorphism C.1 induces a canonical isomorphism

$$(C.2) \quad R_2\langle h_{\text{sch}}(\mathbb{M}, \mathbb{N}, T)\rangle \cong R_2 \otimes_{R_1} R_1\langle(\mathbb{M}, \mathbb{N}, T)\rangle$$

For a map of R_1 modules $U : (\mathbb{M}, \mathbb{N}, T) \rightarrow (\mathbb{M}', \mathbb{N}', T')$, we define $h_{\text{sch}}U$ to be the map defined by the following diagram.

$$\begin{array}{ccc} R_2 \langle h_{\text{sch}}(\mathbb{M}, \mathbb{N}, T) \rangle & \xrightarrow{h_{\text{sch}}U} & R_2 \langle h_{\text{sch}}(\mathbb{M}', \mathbb{N}', T') \rangle \\ \cong \downarrow & & \cong \downarrow \\ R_2 \otimes_{R_1} R_1 \langle (\mathbb{M}, \mathbb{N}, T) \rangle & \xrightarrow{1 \otimes U} & R_2 \otimes_{R_1} R_1 \langle (\mathbb{M}', \mathbb{N}', T') \rangle \end{array}$$

This gives a functor h_{sch} from schematic R_1 modules to schematic R_2 modules. If $h' : R_2 \rightarrow R_3$ is a homomorphism we have $(h' \circ h)_{\text{sch}} = h'_{\text{sch}} \circ h_{\text{sch}}$, and thus the category of schematic R bf modules is a functor of R .

C.6. The ad theory $\text{ad}_{\text{Rel, sch}}^R$. First we translate Definition C.5 into the language of schematic modules:

Definition C.20. (i) A *schematic Relaxed quasi-symmetric complex of dimension n* is a quadruple $(\mathbb{C}, \mathbb{E}, \gamma, \phi)$, where \mathbb{C} is a schematic R chain complex, \mathbb{E} is a schematic $(R^{\text{op}} \otimes R)$ chain complex, $R \langle \mathbb{C} \rangle$ is homotopy finite, $(R^{\text{op}} \otimes R) \langle \mathbb{E} \rangle$ is a homotopy finite chain complex over $R^{\text{op}} \otimes R$ with a $\mathbb{Z}/2$ action for which the generator acts quasi-linearly, γ is a $\mathbb{Z}/2$ equivariant $R^{\text{op}} \otimes R$ -linear quasi-isomorphism $(R \langle \mathbb{C} \rangle)^t \otimes R \langle \mathbb{C} \rangle \rightarrow (R^{\text{op}} \otimes R) \langle \mathbb{E} \rangle$, and ϕ is an n -dimensional element of $(R \otimes_{R^{\text{op}} \otimes R} (R^{\text{op}} \otimes R) \langle \mathbb{E} \rangle)^{\mathbb{Z}/2}$ (where the $\mathbb{Z}/2$ action is given by Lemma C.4).

(ii) We define a category $\mathcal{A}_{\text{Rel, sch}}^R$ as follows. The objects of $\mathcal{A}_{\text{Rel, sch}}^R$ are the schematic Relaxed quasi-symmetric complexes. A morphism $(\mathbb{C}, \mathbb{E}, \gamma, \phi) \rightarrow (\mathbb{C}', \mathbb{E}', \gamma', \phi')$ is a pair $(f : R \langle \mathbb{C} \rangle \rightarrow R \langle \mathbb{C}' \rangle, g : (R^{\text{op}} \otimes R) \langle \mathbb{E} \rangle \rightarrow (R^{\text{op}} \otimes R) \langle \mathbb{E}' \rangle)$, where f is an R -linear chain map and g is a $\mathbb{Z}/2$ equivariant $R^{\text{op}} \otimes R$ -linear chain map, such that $g\gamma = \gamma'(f \otimes f)$, and (if $\dim \phi = \dim \phi'$) $(1 \otimes g)_*(\phi) = \phi'$.

$\mathcal{A}_{\text{Rel, sch}}^R$ is a balanced \mathbb{Z} -graded category, where i takes $(\mathbb{C}, \mathbb{E}, \gamma, \phi)$ to $(\mathbb{C}, \mathbb{E}, \gamma, -\phi)$.

There is a morphism

$$\Lambda : \mathcal{A}_{\text{Rel, sch}}^R \rightarrow \mathcal{A}_{\text{Rel}}^R$$

of \mathbb{Z} -graded categories which takes $(\mathbb{C}, \mathbb{E}, \gamma, \phi)$ to $(R \langle \mathbb{C} \rangle, (R^{\text{op}} \otimes R) \langle \mathbb{E} \rangle, \gamma, \phi)$; this is an equivalence of categories.

Definition C.21. A K -ad with values in $\mathcal{A}_{\text{Rel, sch}}^R$ is a pre K -ad F for which $\Lambda \circ F$ is a K -ad.

We write $\text{ad}_{\text{Rel, sch}}^R(K)$ for the set of K -ads with values in $\mathcal{A}_{\text{Rel, sch}}^R$.

Proposition C.22. (i) $\text{ad}_{\text{Rel, sch}}^R$ is an ad theory.

(ii) Λ induces a morphism of ad theories which is an isomorphism on bordism groups.

This is an easy consequence of Theorem C.12 and the following lemma.

Lemma C.23. *Let K be a ball complex and L a subcomplex. Given a commutative diagram*

$$\begin{array}{ccc} \mathcal{C}ell(L) & \xrightarrow{F} & \mathcal{A} \\ \downarrow & & \downarrow I \\ \mathcal{C}ell(K) & \xrightarrow{G} & \mathcal{A}' \end{array}$$

in which I is an equivalence of categories, there is a functor $H : \mathcal{C}ell(K) \rightarrow \mathcal{A}$ such that $H|_{\mathcal{C}ell(L)} = F$ and $I \circ H$ is naturally isomorphic to G . \square

C.7. $\text{ad}_{\text{Rel,sch}}^R$ is a functor of R . Let $h : R \rightarrow S$ be a homomorphism of rings with involution.

Definition C.24. Define a functor

$$h_{\text{Rel,sch}} : \mathcal{A}_{\text{Rel,sch}}^R \rightarrow \mathcal{A}_{\text{Rel,sch}}^S$$

as follows. For an object $(\mathbb{C}, \mathbb{E}, \gamma, \phi)$ of $\mathcal{A}_{\text{Rel,sch}}^R$, Let

$$h_{\text{Rel}}(\mathbb{C}, \mathbb{E}, \gamma, \phi) = (h_{\text{sch}} \mathbb{C}, (h \otimes h)_{\text{sch}} \mathbb{E}, \delta, \psi),$$

where h_{sch} and $(h \otimes h)_{\text{sch}}$ are given in Subsection C.5, and (letting $C = R\langle \mathbb{C} \rangle$ and $E = (R^{\text{op}} \otimes R)\langle \mathbb{E} \rangle$, and using the notation of Definition C.14 and the isomorphism of Equation (C.2)), δ is defined by the diagram

$$\begin{array}{ccc} S\langle h_{\text{sch}} \mathbb{C} \rangle^t \otimes S\langle h_{\text{sch}} \mathbb{C} \rangle & \xrightarrow{\delta} & (S^{\text{op}} \otimes S)\langle (h \otimes h)_{\text{sch}} \mathbb{E} \rangle \\ \cong \downarrow & & \cong \downarrow \\ (S \otimes_R C)^t \otimes (S \otimes_R C) & \xrightarrow{\gamma'} & (S^{\text{op}} \otimes S) \otimes_{R^{\text{op}} \otimes R} E \end{array}$$

and ψ is the image of ϕ' under the isomorphism

$$(S \otimes_{S^{\text{op}} \otimes S} (S^{\text{op}} \otimes S) \otimes_{R^{\text{op}} \otimes R} E)^{\mathbb{Z}/2} \cong (S \otimes_{S^{\text{op}} \otimes S} (S^{\text{op}} \otimes S) \langle (h \otimes h)_{\text{sch}} \mathbb{E} \rangle)^{\mathbb{Z}/2}.$$

Proposition C.25. $h_{\text{Rel,sch}}$ takes ads to ads.

Proof. Let K be a ball complex and let $F \in \text{ad}_{\text{Rel,sch}}^R(K)$. By Definition C.21, we only need to show that $\Lambda \circ h_{\text{Rel,sch}} \circ F$ is in $\text{ad}_{\text{Rel}}^R(K)$. But $\Lambda \circ h_{\text{Rel,sch}} \circ F$ is naturally isomorphic to $h_{\text{Rel}} \circ \Lambda \circ F$, which is in $\text{ad}_{\text{Rel}}^R(K)$ by Definition C.21 and Proposition C.15. \square

The reader can check that if $k : S \rightarrow T$ is another ring homomorphism then $(kh)_{\text{Rel,sch}}$ is equal to $k_{\text{Rel,sch}}h_{\text{Rel,sch}}$, so $\text{ad}_{\text{Rel,sch}}^R$ is a functor of R as required.

REFERENCES

- [And] Ricardo Andrade (mathoverflow.net/users/21095), *Multisimplicial geometric realization*, MathOverflow, <http://mathoverflow.net/questions/124837> (version: 2013-03-19).
- [BL17] Nils A. Baas and Gerd Laures, *Singularities and Quinn spectra*, Münster J. Math. **10** (2017), no. 1, 1–17. MR 3624098
- [BLM19] Markus Banagl, Gerd Laures, and James E. McClure, *The L-homology fundamental class for IP-spaces and the stratified Novikov conjecture*, Selecta Math. (N.S.) **25** (2019), no. 1, Paper No. 7, 104. MR 3908854
- [EM06] A. D. Elmendorf and M. A. Mandell, *Rings, modules, and algebras in infinite loop space theory*, Adv. Math. **205** (2006), no. 1, 163–228. MR 2254311 (2007g:19001)
- [FM] Greg Friedman and James McClure, *The symmetric signature of a Witt space*, <http://arxiv.org/abs/1106.4798>.
- [FQ90] Michael H. Freedman and Frank Quinn, *Topology of 4-manifolds*, Princeton Mathematical Series, vol. 39, Princeton University Press, Princeton, NJ, 1990. MR 1201584 (94b:57021)
- [GJ99] Paul G. Goerss and John F. Jardine, *Simplicial homotopy theory*, Progress in Mathematics, vol. 174, Birkhäuser Verlag, Basel, 1999. MR 1711612 (2001d:55012)
- [Hir03] Philip S. Hirschhorn, *Model categories and their localizations*, Mathematical Surveys and Monographs, vol. 99, American Mathematical Society, Providence, RI, 2003. MR 1944041 (2003j:18018)
- [HSS00] Mark Hovey, Brooke Shipley, and Jeff Smith, *Symmetric spectra*, J. Amer. Math. Soc. **13** (2000), no. 1, 149–208. MR 1695653 (2000h:55016)
- [KMM] Phillip Kuehl, Tibor Macko, and Adam Mole, *The total surgery obstruction revisited*, <http://arxiv.org/abs/1104.5092>.
- [LM14] Gerd Laures and James E. McClure, *Multiplicative properties of Quinn spectra*, Forum Math. **26** (2014), no. 4, 1117–1185. MR 3228927
- [Lur] Jacob Lurie, *Algebraic L-theory and surgery*, notes for a course, <http://www.math.harvard.edu/~lurie/287x.html>.
- [May72] J. P. May, *The geometry of iterated loop spaces*, Springer-Verlag, Berlin, 1972, Lectures Notes in Mathematics, Vol. 271. MR 0420610 (54 #8623b)
- [May92] J. Peter May, *Simplicial objects in algebraic topology*, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 1992, Reprint of the 1967 original. MR 1206474 (93m:55025)
- [McC] J.E. McClure, *On semisimplicial objects satisfying the Kan condition*, <http://arxiv.org/abs/1210.5650>, to appear in Homology, Homotopy and Applications.
- [ML63] Saunders Mac Lane, *Natural associativity and commutativity*, Rice Univ. Studies **49** (1963), no. 4, 28–46. MR 0170925 (30 #1160)

- [MM79] Ib Madsen and R. James Milgram, *The classifying spaces for surgery and cobordism of manifolds*, Annals of Mathematics Studies, vol. 92, Princeton University Press, Princeton, N.J., 1979. MR 548575 (81b:57014)
- [Qui95] Frank Quinn, *Assembly maps in bordism-type theories*, Novikov conjectures, index theorems and rigidity, Vol. 1 (Oberwolfach, 1993), London Math. Soc. Lecture Note Ser., vol. 226, Cambridge Univ. Press, Cambridge, 1995, pp. 201–271. MR 1388303 (97h:57055)
- [Ran80a] Andrew Ranicki, *The algebraic theory of surgery. I. Foundations*, Proc. London Math. Soc. (3) **40** (1980), no. 1, 87–192. MR 560997 (82f:57024a)
- [Ran80b] ———, *The algebraic theory of surgery. II. Applications to topology*, Proc. London Math. Soc. (3) **40** (1980), no. 2, 193–283. MR 566491 (82f:57024b)
- [Ran92] A. A. Ranicki, *Algebraic L-theory and topological manifolds*, Cambridge Tracts in Mathematics, vol. 102, Cambridge University Press, Cambridge, 1992. MR 1211640 (94i:57051)
- [Ree] C. L. Reedy, *Homotopy theory of model categories*, available at <http://www-math.mit.edu/~psh>.
- [Sch09] Christian Schlichtkrull, *Thom spectra that are symmetric spectra*, Doc. Math. **14** (2009), 699–748. MR 2578805 (2011f:55017)
- [Seg74] Graeme Segal, *Categories and cohomology theories*, Topology **13** (1974), 293–312. MR 0353298 (50 #5782)
- [Sul05] Dennis P. Sullivan, *Geometric topology: localization, periodicity and Galois symmetry*, K-Monographs in Mathematics, vol. 8, Springer, Dordrecht, 2005, The 1970 MIT notes, Edited and with a preface by Andrew Ranicki. MR 2162361 (2006m:55002)
- [Wei94] Charles A. Weibel, *An introduction to homological algebra*, Cambridge studies in advanced mathematics, Cambridge University Press, Cambridge, 1994.

FAKULTÄT FÜR MATHEMATIK, RUHR-UNIVERSITÄT BOCHUM, NA1/66, D-44780 BOCHUM, GERMANY

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, 150 N. UNIVERSITY STREET, WEST LAFAYETTE, IN 47907-2067