# Essential Building Blocks of Dark Energy

Jerome Gleyzes<sup>a</sup>, David Langlois<sup>b</sup>, Federico Piazza<sup>b,c</sup> and Filippo Vernizzi<sup>a</sup>

<sup>a</sup> CEA, IPhT, 91191 Gif-sur-Yvette cédex, France CNRS, URA-2306, 91191 Gif-sur-Yvette cédex, France

<sup>b</sup> APC, (CNRS-Université Paris 7), 10 rue Alice Domon et Léonie Duquet, 75205 Paris, France

<sup>c</sup> PCCP, 10 rue Alice Domon et Léonie Duquet, 75205 Paris, France

#### Abstract

We propose a minimal description of single field dark energy/modified gravity within the effective field theory formalism for cosmological perturbations, which encompasses most existing models. We start from a generic Lagrangian given as an arbitrary function of the lapse and of the extrinsic and intrinsic curvature tensors of the time hypersurfaces in unitary gauge, i.e. choosing as time slicing the uniform scalar field hypersurfaces. Focusing on linear perturbations, we identify seven Lagrangian operators that lead to equations of motion containing at most two (space or time) derivatives, the background evolution being determined by the time dependent coefficients of only three of these operators. We then establish a dictionary that translates any existing or future model whose Lagrangian can be written in the above form into our parametrized framework. As an illustration, we study Horndeski's—or generalized Galileon—theories and show that they can be described, up to linear order, by only six of the seven operators mentioned above. This implies, remarkably, that the dynamics of linear perturbations can be more general than that of Horndeski while remaining second order. Finally, in order to make the link with observations, we provide the entire set of linear perturbation equations in Newtonian gauge, the effective Newton constant in the quasi-static approximation and the ratio of the two gravitational potentials, in terms of the time-dependent coefficients of our Lagrangian.

## Contents

1	Introduction	<b>2</b>					
2	General Lagrangian in unitary gauge         2.1       Background equations         2.2       Perturbations in the ADM formalism         2.3       The EFT language	<b>5</b> 7 7 9					
3	Generalised Galileons3.1Geometric preliminaries3.2 $L_3$ $L_3$ 3.3 $L_4$ $L_5$ 3.5Summary	<ol> <li>11</li> <li>12</li> <li>13</li> <li>13</li> <li>14</li> <li>15</li> </ol>					
4	Observables         4.1       Perturbation equations	<b>15</b> 16 18					
<b>5</b>	Conclusion						
$\mathbf{A}$	Lagrangian dependence on ${}^{(3)}\!R_{\mu u}K^{\mu u}$						
в	B Tensor modes						
$\mathbf{C}$	C EFT parameters for generalized Galileons						
D	D Coefficients of the perturbation equations						

# 1 Introduction

Dark energy has now become a generic name that includes a huge number of models trying to account for the present cosmic acceleration [1,2]. Given their proliferation, the confrontation of such models with present and future cosmological data would be greatly facilitated by an effective approach that can mediate between observational data and theory. Ideally, such a phenomenological approach would provide an effective parameterisation that minimizes the number of free functions and deals directly with the relevant low-energy degrees of freedom, which in our context are the cosmological perturbations (together with the background evolution). A precise dictionary rephrasing the various models into this common language would then simplify the confrontation with the data and point out possible degeneracies between different theories. Within its unifying picture, this effective approach should have the extra virtue of stimulating theorists to study previously unexplored regions of the parameter space which could lead to interesting new models or, conversely, to better understanding why certain regions might be forbidden.

A few steps in this direction have been undertaken recently. The so-called effective field theory (EFT) of cosmological perturbations is a powerful tool that allows to deal directly with the relevant low-energy degrees of freedom of the problem at hand. Such an approach was proposed and intensively used for inflation [3,4], in particular to characterize high-energy corrections to slow-roll models and to predict high-order correlation functions (see e.g. [5,6]). The EFT of inflation has now become a standard way of parametrising primordial non-Gaussianity and was used, for instance, in the

interpretation of the most recent WMAP [7] and Planck [8] data. This approach has also been applied to dark energy, first in the minimally-coupled case [9] where it was proven a useful tool to study the stability in full generality and, for models with vanishing sound speed, the clustering of dark energy down to very nonlinear scales [10].

More recently, the EFT formalism has been extended to dark energy with non-minimal couplings [11, 12], providing a unifying theoretical framework for practically all single-field dark energy and modified gravity models.<sup>1</sup> This approach relies on two basic steps [11]: *a*) assume the weak equivalence principle and therefore the existence of a metric  $g_{\mu\nu}$  universally coupled to all matter fields (it is straightforward to relax this assumption, but at the price of complicating the formalism); *b*) write the *unitary gauge* action, i.e. the most general gravitational action for such a metric compatible with the (unbroken) spatial diffeomorphism invariance on hypersurfaces of constant dark energy field.

In [11] it was argued that the EFT of dark energy has all the virtues advocated at the beginning of this section. The goal of this article is to provide a systematic procedure to translate an arbitrary dark energy model into the EFT language, as well as to establish a firm minimal setting of Lagrangian operators within this framework. In particular, here we focus our attention on the operators of the unitary gauge action that lead to at most two derivatives in the equations of motion for linear perturbations.<sup>2</sup> This minimal set of operators encompasses most of the theoretical models of dark energy and/or modified gravity discussed in the current literature.

The key ingredient of our derivation is a 3 + 1 decomposition à la ADM, where time slicings coincide with the uniform scalar field hypersurfaces. With this time choice, the dynamics of the underlying degree of freedom is embodied in the dynamics of the 3-dimensional metric. In Sec. 2 we consider a generic Lagrangian given as an arbitrary function of the lapse  $N \equiv 1/\sqrt{-g^{00}}$  and of the 3-dimensional metric  $h_{\mu\nu} \equiv g_{\mu\nu} + n_{\mu}n_{\nu}$ , where  $n^{\mu}$  is the unit vector perpendicular to constant time hypersurfaces, more specifically of its extrinsic and intrinsic curvature tensors, respectively  $K_{\nu}^{\ \mu} \equiv h_{\nu}^{\ \rho} \nabla_{\rho} n^{\mu}$  and  ${}^{(3)}R^{\mu}_{\ \nu}$ ,

$$S = \int d^4x \sqrt{-g} L(N, K^{\mu}_{\ \mu}, K_{\mu\nu}K^{\mu\nu}, {}^{(3)}R, {}^{(3)}R_{\mu\nu}{}^{(3)}R^{\mu\nu}, \dots; t) .$$
(1)

In our construction we include combinations of these 3-dimensional objects without taking their derivatives. This automatically prevents the appearance of higher (more than two) time derivatives in the equations of motion. However, it is not enough to also remove higher spatial derivatives. By expanding this Lagrangian up to quadratic order in the cosmological perturbations and making use of an ADM analysis in unitary gauge (see for instance [3, 19, 20]) we obtain specific conditions that ensure the absence of higher spatial derivatives in Sec. 2.2.

Moreover, we also show how the parameters in front of the standard EFT operators of [11] can be expressed in terms of the time-dependent coefficients of the expansion of (1). Since the action of most of the existing theoretical models can be written as eq. (1), this can be used to derive a dictionary between theoretical models and our EFT language. As an illustration, in Sec. 3 we explicitly derive this dictionary for the most general scalar field theory leading to at most second order equations of motion, i.e. the Horndeski theory [21] (see also [22]), recently rediscovered in the context of the so-called Galileon field [23, 24] under the name of "generalized Galileons" [25, 26].

Let us summarize here the main results of Secs. 2 and 3:

<sup>&</sup>lt;sup>1</sup>An alternative formulation of a background independent effective approach to dark energy and modified gravity was given in [13]. A covariant EFT of cosmological acceleration was developed in [14] for inflation and generalized to the case of dark energy in [15, 16]—see discussion in [11] for a comparison between the latter approach and the one advocated here. For a different unifying framework to cosmological perturbations for dark energy and modified gravity see, for instance, [17].

<sup>&</sup>lt;sup>2</sup>This is sufficient to ensure that we only have a single propagating degree of freedom. Note, however, that higher time derivatives do not lead to higher degrees of freedom if they can be treated perturbatively, i.e. evaluating them using the lower order equations of motion [18].

• The most general EFT action, up to quadratic order, for single-field dark energy, in the Jordan frame, leading to at most second-order equations of motion for *linear* perturbations can be written as

$$S = \int d^4x \sqrt{-g} \left[ \frac{M_*^2}{2} f(t)R - \Lambda(t) - c(t)g^{00} + \frac{M_2^4(t)}{2} (\delta g^{00})^2 - \frac{m_3^3(t)}{2} \,\delta K \delta g^{00} - m_4^2(t) \left( \delta K^2 - \delta K^{\mu}_{\ \nu} \,\delta K^{\nu}_{\ \mu} \right) + \frac{\tilde{m}_4^2(t)}{2} \,{}^{(3)}R \,\delta g^{00} \right] \,,$$

$$(2)$$

where  $\delta g^{00} \equiv g^{00} + 1$ ,  $\delta K_{\mu\nu} \equiv K_{\mu\nu} - Hh_{\mu\nu}$ ,  $K \equiv K^{\mu}_{\ \mu}$  and we have assumed a flat Universe so that  ${}^{(3)}R^{\mu}_{\ \nu}$  vanishes on the background<sup>3</sup>. This action describes the propagation of one scalar degree of freedom with dispersion relation  $\omega^2 = c_s^2 k^2$ , where  $c_s$  is the sound speed of fluctuations given by eq. (35) with the relations (47). Stability (absence of ghosts) is ensured by the positivity of the time kinetic term given in (47). The particular combination appearing in the operator proportional to  $m_4^2$  is such that it does not lead to higher-order spatial derivatives. One can check that also the combination

$${}^{(3)}R^{\mu}_{\nu}\,\delta K^{\nu}_{\mu} - \frac{1}{2}{}^{(3)}R\,\delta K \tag{3}$$

does not generate higher derivatives. However, this operator is not explicitly included in eq. (2) because it can be reexpressed in terms of the others (see App. A).

- In the particular case where  $m_4^2 = \tilde{m}_4^2$ , the above action is equivalent to the *linearized* Horndeski's theory/generalized Galileons and the explicit dictionary between generalized Galileons and this action is given in App. C. This implies that the dynamics of linear scalar perturbations of action (2) is more general than that of Horndeski, while remaining second order in time and space derivatives.
- Expanding the Lagrangian (1) up to quadratic order we also find three operators that lead to higher order space—but not time—derivatives. These are

$$S_{\text{h.s.d.}} = \int d^4x \sqrt{-g} \left[ -\bar{m}_4^2(t) \,\delta K^2 \,+\, \frac{\bar{m}_5(t)}{2} \,^{(3)}R \,\delta K \,+\, \frac{\bar{\lambda}(t)}{2} \,^{(3)}R^2 \right] \,. \tag{4}$$

When one of these operators is present in the action the dispersion relation of the propagating mode receives corrections at large momenta,  $\omega^2 = c_s^2 k^2 + k^4/M^2$ , where M is a mass scale. These corrections may become important in the limit of vanishing sound speed, such as in the model of the Ghost Condensate [27] or for deformations of this particular limit [3,9].

Once a Lagrangian describing matter has been included, the action (2) can be used as a benchmark for the study of physical signatures of dark energy/modified gravity in the linear regime. In this context, cosmological perturbations are usually discussed in Newtonian gauge, which is the one that we employ in Sec. 4. In order to do that, in Sec. 4.1 we restore the covariance via the Stueckelberg trick [3, 4, 27] and we vary this action with respect to all the scalar dynamical degrees of freedom. This allows to derive Einstein's equations and the evolution equation for the fluctuations of the scalar field responsible of the acceleration, recovering and generalizing previous results [9,11]. Interestingly, in the Newtonian gauge Einstein's equations and the scalar equation of motion contain higher order derivatives when  $m_4^2 \neq \tilde{m}_4^2$ , even if the dynamical equation for the true degree of freedom is only second order. Finally, in Sec. 4.2 we use these equations to derive the effective Newton constant and the ratio between the two gravitational potentials.

<sup>&</sup>lt;sup>3</sup>The case of non-vanishing spatial curvature is commented on in footnote 7.

The phenomenology of the operators appearing in action (2) was also studied in [12]. In this reference it was indeed mentioned that these operators are sufficient to describe linear perturbations of Horndeski's theories. However, no proof of this statement was given nor the particular combination in which such operators appear shown.

# 2 General Lagrangian in unitary gauge

In the presence of a scalar field  $\phi$  with a non-vanishing timelike gradient, the so-called unitary gauge corresponds to a choice of time slicing where the constant time hypersurfaces are uniform  $\phi$  hypersurfaces. The use of unitary gauge accomplishes two main objectives. First, as explained at length in Refs. [3, 4, 27], it makes it straightforward to write a generic Lagrangian for cosmological perturbations. Since the dynamics of the scalar field has been "eaten" by the metric, the most generic Lagrangian is simply that for the *metric perturbations* around a FLRW solution, compatible with the unbroken symmetry of 3-dimensional diffeomorphisms.

Second, the 3+1 splitting in unitary gauge easily allows to keep the number of time derivatives under control, while considering higher and higher space derivatives. Therefore, the unitary gauge is helpful to systematically explore the space of higher spatial derivative theories by considering geometric invariants on the  $\phi = constant$  hypersurfaces. In practice, we will use the metric in the ADM form

$$ds^{2} = -N^{2}dt^{2} + h_{ij}\left(dx^{i} + N^{i}dt\right)\left(dx^{j} + N^{j}dt\right),$$
(5)

where the 3-dimensional metric  $h_{ij}$  is used to lower and raise latin indices  $i, j, \dots = 1, 2, 3$ . Since 3dimensional diffeomorphism invariance is preserved in unitary gauge, it is natural to write operators (with up to two spatial derivatives per field) in terms of the extrinsic and intrinsic curvatures  $K_{\mu\nu}$ and  ${}^{(3)}R_{\mu\nu}$  and their possible contractions. The Lagrangian is also an explicit function of the lapse function N in general.

Therefore, in the following, we consider a general action of the form

$$S = \int d^4x \sqrt{-g} L(N, K, \mathcal{S}, \mathcal{R}, \mathcal{Z}; t) , \qquad (6)$$

where the Lagrangian L is an arbitrary function of N and of the following four scalar quantities constructed by contracting the extrinsic and intrinsic curvature tensors:

$$K \equiv K^{\mu}_{\ \mu} , \qquad \mathcal{R} \equiv {}^{(3)}R \equiv {}^{(3)}R^{\mu}_{\ \mu} , \qquad \mathcal{S} \equiv K_{\mu\nu}K^{\mu\nu} , \qquad \mathcal{Z} \equiv {}^{(3)}R_{\mu\nu}{}^{(3)}R^{\mu\nu} . \tag{7}$$

Although one should also allow, in principle, for a dependence on  $\mathcal{Y} \equiv {}^{(3)}R_{\mu\nu}K^{\mu\nu}$ , we have preferred not to include it explicitly in the main body, for simplicity. As shown in App. A, this extra dependence leads to a quadratic Lagrangian of the same form as that found later in this section, with slightly modified coefficients. Indeed, since  $\mathcal{Y}$  is equivalent to  $H\mathcal{R}$  at linear order, the quadratic terms in the expansion of the Lagrangian induced by its dependence on  $\mathcal{Y}$  are analogous to those induced by its dependence on  $\mathcal{R}$ . As for the linear term, one can use the equality

$$\lambda(t)^{(3)}R_{\mu\nu}K^{\mu\nu} = \frac{\lambda(t)}{2}{}^{(3)}R K + \frac{\dot{\lambda}(t)}{2N}{}^{(3)}R + \text{boundary terms}, \qquad (8)$$

which is also shown in App. A.

Moreover, one could also consider scalars that are combinations of three or more tensors, like  $K^{\lambda}_{\ \mu}K^{\mu}_{\ \nu}K^{\nu}_{\ \lambda}$ , but it is easy to show that also in this case they can be re-expressed in terms of the above combinations, plus corrections which are at least cubic in the perturbations. We will show this explicitly for the extended Galileon in the next section. Finally, one could take quadratic combinations of the Riemann tensor such as  ${}^{(3)}R_{\mu\nu\rho\sigma}{}^{(3)}R^{\mu\nu\rho\sigma}$ . However, in three dimensions the

Riemann tensor can be expressed in terms of the Ricci scalar and tensor.<sup>4</sup> Thus, at quadratic order in the perturbations, the action above seems to exhaust all the possibilities compatible with our requirements.

In order to explicitly write the expansion of the action (6) up to second order in the perturbations, it is useful to define the tensors

$$\delta K \equiv K - 3H, \qquad \delta K_{\mu\nu} \equiv K_{\mu\nu} - Hh_{\mu\nu} \,, \tag{10}$$

which vanish on the background, and to use the decompositions

$$S = 3H^2 + \delta S, \qquad \delta S \equiv 2H\delta K + \delta K^{\mu}_{\ \nu} \delta K^{\nu}_{\ \mu}.$$
 (11)

The quantities  $\mathcal{R}$  and  $\mathcal{Z}$  vanish on the background and are therefore already perturbative ( $\mathcal{Z}$  is even a second order quantity).

The expansion of the Lagrangian up to second order in the perturbations yields, after discarding irrelevant boundary terms, the expression

$$L(N, K, \mathcal{S}, \mathcal{R}, \mathcal{Z}) = \bar{L} - \dot{\mathcal{F}} - 3H\mathcal{F} + (\dot{\mathcal{F}} + L_N)\,\delta N + L_{\mathcal{R}}\,\delta \mathcal{R} + \frac{\mathcal{A}}{2}\,\delta K^2 + L_{\mathcal{S}}\,\delta K^{\mu}_{\ \nu}\delta K^{\nu}_{\ \mu} + \left(\frac{1}{2}L_{NN} - \dot{\mathcal{F}}\right)\delta N^2 + \frac{1}{2}L_{\mathcal{R}\mathcal{R}}\,\delta \mathcal{R}^2 + \mathcal{B}\,\delta K\delta N + \mathcal{C}\,\delta K\delta \mathcal{R} + L_{N\mathcal{R}}\,\delta N\delta \mathcal{R} + L_{\mathcal{Z}}\delta \mathcal{Z} + \mathcal{O}(3)\,,$$
(12)

where we have introduced the following notations for some combinations of the partial derivatives of the Lagrangian (denoting  $L_N \equiv \partial L/\partial N$ , etc.), to make this expression more compact:

$$\mathcal{F} \equiv 2HL_{\mathcal{S}} + L_{K},$$

$$\mathcal{A} \equiv 4H^{2}L_{\mathcal{S}\mathcal{S}} + 4HL_{\mathcal{S}K} + L_{KK},$$

$$\mathcal{B} \equiv 2HL_{\mathcal{S}N} + L_{KN},$$

$$\mathcal{C} \equiv 2HL_{\mathcal{S}\mathcal{R}} + L_{K\mathcal{R}}.$$
(13)

The first term,  $\bar{L}$ , is the homogeneous Lagrangian and all partial derivatives of L that appear in the above expression are evaluated on the homogeneous background, i.e. for  $\bar{N} = 1$ ,  $\bar{S} = 3H^2$ ,  $\bar{K} = 3H$ ,  $\bar{\mathcal{R}} = 0$  and  $\bar{\mathcal{Z}} = 0$ . Note that, in order to obtain the expression (12), we have rewritten the term linear in  $\delta K$  as

$$\mathcal{F}\delta K = \mathcal{F}(K - 3H) , \qquad (14)$$

and integrated it by parts using  $K = \nabla_{\mu} n^{\mu}$ ,

$$\int d^4x \sqrt{-g} \,\mathcal{F}K = -\int d^4x \sqrt{-g} \,n^\mu \nabla_\mu \mathcal{F} = -\int d^4x \sqrt{-g} \frac{\dot{\mathcal{F}}}{N} \,, \tag{15}$$

where  $n^{\mu}$  is the unit vector orthogonal to constant time hypersurfaces and, in unitary gauge, has time component  $n^0 = 1/N$ .

$${}^{(3)}R_{\mu\nu\rho\sigma} = {}^{(3)}R_{\mu\rho}h_{\nu\sigma} - {}^{(3)}R_{\nu\rho}h_{\mu\sigma} - {}^{(3)}R_{\mu\sigma}g_{\nu\rho} + {}^{(3)}R_{\nu\sigma}h_{\mu\rho} - \frac{1}{2}{}^{(3)}R(h_{\mu\rho}h_{\nu\sigma} - h_{\mu\sigma}h_{\nu\rho}) .$$
(9)

<sup>&</sup>lt;sup>4</sup>This can be done using the relation

#### 2.1 Background equations

For the background we assume a flat homogeneous FLRW metric, written in the form

$$ds^{2} = -N^{2}(t)dt^{2} + a^{2}(t)\delta_{ij}dx^{i}dx^{j}.$$
(16)

In this case K = 3H/N and  $S = 3H^2/N^2$ , where  $H \equiv \dot{a}/a$  is the Hubble rate. Note that it is crucial to explicitly keep the lapse function N, because the first Friedmann equation is the constraint arising from the invariance under time reparametrization. Linear variation of the homogeneous action  $S_0$  with respect to the lapse N and the scale factor a yields

$$\delta S_0 = \int dt d^3x \left[ a^3 \left( \bar{L} + L_N - 3H\mathcal{F} \right) \delta N + 3a^2 \left( \bar{L} - 3H\mathcal{F} - \dot{\mathcal{F}} \right) \delta a \right] , \qquad (17)$$

where we have used  $\sqrt{-g} = a^3 N$ . Then the Friedmann equations are directly given by<sup>5</sup>

$$3H\mathcal{F} - L - L_N = 0,\tag{18}$$

which depends on first time derivatives at most, and

$$\dot{\mathcal{F}} + 3H\mathcal{F} - \bar{L} = 0, \qquad (19)$$

which determines the dynamics of the scale factor.

As expected, by using the above equations one can check that the first order of the total Lagrangian  $\mathcal{L} \equiv \sqrt{-g} L$  vanishes. Indeed, using  $\sqrt{-g} = \sqrt{h} N$ , where h is the determinant of the 3-dimensional metric  $h_{ij}$  in the ADM decomposition, one easily finds

$$\mathcal{L}_1 = \left(\bar{L} - 3H\mathcal{F} - \dot{\mathcal{F}}\right)\delta\sqrt{h} + a^3(L_N + \bar{L} - 3H\mathcal{F})\delta N + a^3L_{\mathcal{R}}\delta\mathcal{R}, \qquad (20)$$

where the last term is a total derivative and can be ignored.

### 2.2 Perturbations in the ADM formalism

In this sub-section we perform the analysis of the perturbations in unitary gauge and by using the ADM form of the metric, eq. (5). For the action at second order, we will only need to take into account the perturbations of  $\sqrt{-g}$  at first order,  $\delta\sqrt{-g} = \delta\sqrt{h} + a^3\delta N$ , because the second order one multiplies the LHS of eq. (19). Thus, the quadratic Lagrangian for perturbations is given by

$$\mathcal{L}_{2} = \delta \sqrt{h} \left[ (\dot{\mathcal{F}} + L_{N}) \delta N + L_{\mathcal{R}} \, \delta \mathcal{R} \right] + a^{3} \left[ \left( L_{N} + \frac{1}{2} L_{NN} \right) \delta N^{2} + L_{\mathcal{R}} \delta_{2} \mathcal{R} + \frac{1}{2} \mathcal{A} \, \delta K^{2} + \mathcal{B} \, \delta K \delta N + \mathcal{C} \, \delta K \delta \mathcal{R} + L_{\mathcal{S}} \, \delta K^{\mu}_{\nu} \, \delta K^{\nu}_{\mu} + L_{\mathcal{Z}} \, \delta \mathcal{R}^{\mu}_{\nu} \, \delta \mathcal{R}^{\nu}_{\mu} + \frac{1}{2} L_{\mathcal{R}\mathcal{R}} \, \delta \mathcal{R}^{2} + (L_{\mathcal{R}} + L_{N\mathcal{R}}) \, \delta N \delta \mathcal{R} \right] ,$$

$$(21)$$

where  $\delta_2 \mathcal{R}$  denotes the expansion of  $\mathcal{R}$  at second order in the perturbations.

In the ADM decomposition (5) the only relevant components of the extrinsic curvature tensor are given by

$$K_{ij} = \frac{1}{2N} (\dot{h}_{ij} - \nabla_i N_j - \nabla_j N_i) , \qquad (22)$$

where  $\nabla_i$  stands for the covariant derivative associated with the 3-dimensional metric  $h_{ij}$ . For explicit calculations in unitary gauge we choose to describe scalar perturbations of the spatial metric in terms of  $\zeta$  [19],

$$h_{ij} = a^2(t)e^{2\zeta}\delta_{ij}.$$
(23)

<sup>&</sup>lt;sup>5</sup>We have not included explicitly the matter in the Friedmann equations, but it is straightforward to do so.

(We consider the tensor modes separately in App. B.) Thus, the perturbations of the quantities used above are given by

$$\delta\sqrt{h} = 3a^{3}\zeta, \qquad \delta K^{i}{}_{j} = \left(\dot{\zeta} - H\delta N\right)\delta^{i}_{j} - \frac{1}{a^{2}}\delta^{ik}\partial_{(k}N_{j)}, \qquad (24)$$

and

$$\delta \mathcal{R}_{ij} = -\delta_{ij}\partial^2 \zeta - \partial_i \partial_j \zeta , \qquad \delta_2 \mathcal{R} = -\frac{2}{a^2} \left[ (\partial \zeta)^2 - 4\zeta \partial^2 \zeta \right] . \tag{25}$$

Note that in this section  $\partial$  stands for a spatial derivative and  $\partial^2 \equiv \partial_i \partial^i$ . By using the above expressions, the variation of  $\mathcal{L}_2$  with respect to  $\delta N$  yields the Hamiltonian constraint, which reads

$$[L_{NN} + 2L_N + 3H (3H\mathcal{A} + 2HL_S - 2\mathcal{B})] \delta N + 3 (\mathcal{B} - 3H\mathcal{A} - 2HL_S) \dot{\zeta} + 3(L_N + \dot{\mathcal{F}})\zeta - (\mathcal{B} - 3H\mathcal{A} - 2HL_S) \frac{\partial^2 \psi}{a^2} - 4 (L_R + L_{NR} - 3H\mathcal{C}) \frac{\partial^2 \zeta}{a^2} = 0.$$
(26)

By varying  $\mathcal{L}_2$  with respect to the shift

$$N_i \equiv \partial_i \psi , \qquad (27)$$

one obtains the momentum constraint, which implies

$$-\left(\mathcal{B} - 3H\mathcal{A} - 2HL_{\mathcal{S}}\right)\delta N + \left(\mathcal{A} + 2L_{\mathcal{S}}\right)\frac{\partial^{2}\psi}{a^{2}} = \left(3\mathcal{A} + 2L_{\mathcal{S}}\right)\dot{\zeta} - 4\mathcal{C}\frac{\partial^{2}\zeta}{a^{2}}.$$
(28)

By combining the two constraints, one can express both  $\delta N$  and  $\partial^2 \psi$  as functions of  $\zeta$  and its derivatives and then substitute in the action to write it only in terms of  $\zeta$  and its derivatives. In general, a term proportional to  $(\partial^2 \zeta)^2$  will remain. Here, in order to single out the lowest derivatives operators first, we want to find conditions under which this term disappears. If one considers the second order action before the substitution of the constraints, one finds the following terms

$$\frac{1}{a} \left[ \frac{1}{2} \left( \mathcal{A} + 2L_{\mathcal{S}} \right) \left( \partial^2 \psi \right)^2 + 4\mathcal{C} \ \partial^2 \psi \ \partial^2 \zeta + 2 \left( 4L_{\mathcal{RR}} + 3L_{\mathcal{Z}} \right) \left( \partial^2 \zeta \right)^2 \right].$$
<sup>(29)</sup>

Taking into account the momentum constraint (28), one immediately sees that imposing the three conditions<sup>6</sup>

$$\mathcal{A} + 2L_{\mathcal{S}} = 0 , \qquad \mathcal{C} = 0 , \qquad 4L_{\mathcal{R}\mathcal{R}} + 3L_{\mathcal{Z}} = 0 , \qquad (30)$$

implies the elimination of the term proportional to  $(\partial^2 \zeta)^2$  in the final action and the absence of higher derivatives in the equation of motion for  $\zeta$ . As we will see in the next section, all generalized Galileon models satisfy the three conditions (30).

When (30) are satisfied, the momentum constraint reduces to

$$\delta N = \mathcal{D}\dot{\zeta}, \qquad \mathcal{D} \equiv \frac{4L_{\mathcal{S}}}{\mathcal{B} + 4HL_{\mathcal{S}}}.$$
 (31)

By direct substitution into  $\mathcal{L}_2$  and after an integration by parts to get rid of the term  $\dot{\zeta}\partial^2\zeta$  (note that the  $\zeta\dot{\zeta}$  term vanishes because of the background equations of motion), we finally get the following Lagrangian for  $\zeta$ :

$$\mathcal{L}_2 = \frac{a^3}{2} \left[ \mathcal{L}_{\dot{\zeta}\dot{\zeta}} \dot{\zeta}^2 + \mathcal{L}_{\partial_i \zeta \partial_i \zeta} \frac{(\partial_i \zeta)^2}{a^2} \right],\tag{32}$$

<sup>&</sup>lt;sup>6</sup>Note that these conditions are only sufficient. A more general analysis can be performed by explicitly requiring that the coefficient of  $(\partial^2 \zeta)^2$  vanishes once the two constraints have been solved. However, this leads to a very complicated equation involving many of the coefficients of the quadratic Lagrangian and it is not clear whether one can find physically relevant solutions that evade (30).

with

$$\mathcal{L}_{\dot{\zeta}\dot{\zeta}} \equiv 2\left(\frac{1}{2}L_{NN} + L_N - 3H\mathcal{B} - 6H^2L_{\mathcal{S}}\right)\mathcal{D}^2 + 12L_{\mathcal{S}},$$

$$\mathcal{L}_{\partial_i\zeta\partial_i\zeta} \equiv 4\left[L_{\mathcal{R}} - \frac{1}{a}\frac{d}{dt}(a\mathcal{M})\right], \qquad \mathcal{M} \equiv \mathcal{D}(L_{\mathcal{R}} + L_{N\mathcal{R}}).$$
(33)

Classical and quantum stability (absence of ghosts) requires that the time kinetic energy is positive (see, e.g. [3,9]),

$$\mathcal{L}_{\dot{c}\dot{c}} > 0 . \tag{34}$$

The sound speed (squared) of fluctuations can be simply computed by taking the ratio

$$c_s^2 = -\frac{\mathcal{L}_{\partial_i \zeta \partial_i \zeta}}{\mathcal{L}_{\dot{\zeta} \dot{\zeta}}} \,. \tag{35}$$

## 2.3 The EFT language

We are now going to express the conditions on the absence of higher derivatives in terms of the coefficients of the action of the EFT formalism of Refs. [11,12]. The action up to quadratic order in the perturbations can be written as

$$S = \int d^4x \sqrt{-g} \left[ \frac{M_*^2}{2} fR - \Lambda - cg^{00} + \frac{M_2^4}{2} (\delta g^{00})^2 - \frac{\bar{m}_1^3}{2} \delta K \delta g^{00} - \frac{\bar{M}_2^2}{2} \delta K^2 - \frac{\bar{M}_3^2}{2} \delta K_{\nu}^{\mu} \delta K_{\mu}^{\nu} + \frac{\mu_1^2}{2} {}^{(3)}R \delta g^{00} + \frac{\bar{m}_5}{2} {}^{(3)}R \delta K + \frac{\lambda_1}{2} {}^{(3)}R^2 + \frac{\lambda_2}{2} {}^{(3)}R_{\nu}^{\mu} {}^{(3)}R_{\mu}^{\nu} \right],$$
(36)

where R in the first term inside the bracket is the four-dimensional Ricci scalar. Note that, in order to make the comparison with the previous subsection simpler, we have found more convenient to use the 3-dimensional Ricci scalar and tensor in the quadratic terms, instead of the four-dimensional ones used in Ref. [11], since the link with the ADM decomposition is then transparent.

Let us first discuss how the background equations (18) and (19) translate in this language. In action (36) we have used the time-time component of the inverse metric  $g^{00}$  and its perturbation in the expansion of quadratic and higher order operators, as it is customary in the EFT formalism. However, in the previous subsections it was more convenient to work directly with the lapse function N, related to  $g^{00}$  by

$$g^{00} = -\frac{1}{N^2} \,. \tag{37}$$

Only the first three terms in brackets in eq. (36) contribute to  $\overline{L}$ ,  $L_N$  and  $\mathcal{F}$ , and thus to the background equations of motion. Using eq. (37) and employing the decomposition of the four-dimensional curvature scalar,

$$R = {}^{(3)}R + K_{\mu\nu}K^{\mu\nu} - K^2 + 2\nabla_{\nu}(n^{\nu}\nabla_{\mu}n^{\mu} - n^{\mu}\nabla_{\mu}n^{\nu}), \qquad (38)$$

after an integration by parts in the action we can rewrite these terms as

$$L_0 = \frac{M_*^2}{2} \left( f\mathcal{R} + f\mathcal{S} - fK^2 - 2\dot{f}\frac{K}{N} \right) - \Lambda + \frac{c}{N^2} , \qquad (39)$$

(we remind the reader that  $\mathcal{R} \equiv {}^{(3)}R$ ). By expanding at linear order in  $\delta N$ , integrating by parts the terms linear in K, we can match the background and linear terms of this action with the first line of eq. (12), which yields

$$\bar{L} - \dot{\mathcal{F}} - 3H\mathcal{F} = M_*^2 (3fH^2 + 2f\dot{H} + 2\dot{f}H + \ddot{f}) + c - \Lambda ,$$
  
$$\dot{\mathcal{F}} + L_N = M_*^2 (\dot{f}H - 2f\dot{H} - \ddot{f}) - 2c .$$
(40)

From these two relations and using the background equations of motion (18) and (19) one finds that c and  $\Lambda$  are given by

$$c + \Lambda = 3M_*^2 \left( fH^2 + \dot{f}H \right) , \qquad (41)$$

$$\Lambda - c = M_*^2 \left( 2f\dot{H} + 3fH^2 + 2\dot{f}H + \ddot{f} \right) .$$
(42)

This coincides with what was found in Ref. [11] in the absence of matter.

To discuss linear perturbations we only need the second-order expansion of the action (36). By rewriting the first three terms as in eq. (39), expressing  $g^{00}$  in terms of N and using the definitions (10) and (11), one immediately sees that the EFT action is of the form (6). One can thus use the second-order expansion of the Lagrangian (21) with the following dictionary:

$$L_{\mathcal{R}} = \frac{1}{2}M_{*}^{2}f ,$$

$$\frac{1}{2}L_{NN} + L_{N} = c + 2M_{2}^{4} ,$$

$$\mathcal{A} = -M_{*}^{2}f - \bar{M}_{2}^{2} ,$$

$$\mathcal{B} = \dot{f}M_{*}^{2} - \bar{m}_{1}^{3} ,$$

$$\mathcal{C} = \frac{\bar{m}_{5}}{2} ,$$

$$L_{\mathcal{S}} = \frac{1}{2} \left(M_{*}^{2}f - \bar{M}_{3}^{2}\right) ,$$

$$L_{\mathcal{Z}} = \frac{\lambda_{2}}{2} ,$$

$$L_{N\mathcal{R}} = \mu_{1}^{2} ,$$

$$L_{\mathcal{R}\mathcal{R}} = \lambda_{1} ,$$
(43)

which is completed with eq. (40).

With these relations, the conditions for the absence of higher derivatives, eq. (30), can be written in the EFT of dark energy language. They read:

$$\bar{M}_2^2 + \bar{M}_3^2 = 0$$
,  $\bar{m}_5 = 0$ ,  $4\lambda_1 + \frac{3}{2}\lambda_2 = 0$ . (44)

These conditions are straightforward to verify. Using eqs. (24) and (27),  $\delta K^2$  contains a higher derivative term,  $(\partial^2 \psi)^2$ , while  $\delta K^{\mu}_{\ \nu} \delta K^{\nu}_{\mu}$  contains  $(\partial_i \partial_j \psi)^2$ . However, when the first condition in eq. (44) is satisfied the combination of higher derivative terms in eq. (36) gives an irrelevant boundary term. The second condition implies that the operator  ${}^{(3)}R \delta K$ , which contains  $\partial^2 \psi \partial^2 \zeta$ , does not appear. Finally,  ${}^{(3)}R^2 = 16(\partial^2 \zeta)^2/a^4$  and  ${}^{(3)}R_{ij}{}^{(3)}R^{ij} = [5(\partial^2 \zeta)^2 + (\partial_i \partial_j \zeta)^2]/a^4$ : one can check that when the third condition is satisfied the sum of the two operators in eq. (36) vanishes up to a total derivative.

In summary, the most general EFT Lagrangian which does not generate higher derivatives in the linear equations for the perturbations is<sup>7</sup>

$$L = \frac{M_*^2}{2} f(t)R - \Lambda(t) - c(t)g^{00} + \frac{M_2^4(t)}{2} (\delta g^{00})^2 - \frac{m_3^3(t)}{2} \delta K \delta g^{00} - m_4^2(t) \left(\delta K^2 - \delta K^{\mu}_{\ \nu} \,\delta K^{\nu}_{\ \mu}\right) + \frac{\tilde{m}_4^2(t)}{2} \,{}^{(3)}R \,\delta g^{00} , \qquad (45)$$

<sup>&</sup>lt;sup>7</sup>Let us comment here on the case of a non-vanishing spatial curvature, to which our formalism can be extended straightforwardly with the following caveats. Obviously,  $\delta^{(3)}R$  should be used instead of  ${}^{(3)}R$  in the quadratic operators, but apart from this the Lagrangian (45) and its properties are unchanged. The background equations change (see e.g. eqs. 16 and 17 or Ref. [11]), as well as the dictionary (43), because some first order quantity will now contribute already at zeroth order. The explicit expressions (25) of  ${}^{(3)}R$  change by a term linear in  $\zeta$  but with no derivatives, which, therefore, will not produce higher derivatives in the ADM analysis of Sec 2.2.

where

$$m_3^3 \equiv \bar{m}_1^3 , \qquad m_4^2 \equiv \frac{1}{4} (\bar{M}_2^2 - \bar{M}_3^2) , \qquad \tilde{m}_4^2 \equiv \mu_1^2 ,$$
(46)

as in eq. (2). Terms containing  ${}^{(3)}R^2$  and  ${}^{(3)}R^{\mu}_{\nu}{}^{(3)}R^{\nu}_{\mu}$  do not appear because they only contain higher spatial derivatives. By employing the dictionary (43) in eq. (33), the quadratic action for  $\zeta$  is given by eq. (32) where

$$\mathcal{L}_{\dot{\zeta}\dot{\zeta}} = 2\left(c + 2M_2^4 - 3H^2M_*^2f - 3HM_*^2\dot{f} + 3Hm_3^3 - 6H^2m_4^2\right)\mathcal{D}^2 + 6(M_*^2f + 2m_4^2) ,$$

$$\mathcal{L}_{\partial_i\zeta\partial_i\zeta} = 2\left[M_*^2f - \frac{2}{a}\frac{d}{dt}(a\mathcal{M})\right] ,$$
(47)

and

$$\mathcal{D} = \frac{2(M_*^2 f + 2m_4^2)}{2H(M_*^2 f + 2m_4^2) + M_*^2 \dot{f} - m_3^3},$$

$$\mathcal{M} = \frac{\mathcal{D}}{2}(M_*^2 f + 2\tilde{m}_4^2).$$
(48)

The stability of a given model is then determined by the condition  $\mathcal{L}_{\dot{\zeta}\dot{\zeta}} > 0$  and the speed of sound can be straightforwardly computed from eq. (35) by using the relations above. One can check that these results agree with those found in [11] in the limit  $m_4^2 = \tilde{m}_4^2 = 0$ .

Finally, we can also write down the independent operators that generate higher spatial derivatives. These are

$$L_{\text{h.s.d.}} = -\bar{m}_4^2(t)\,\delta K^2 \,+\,\frac{\bar{m}_5(t)}{2}\,{}^{(3)}\!R\,\delta K \,+\,\frac{\bar{\lambda}(t)}{2}{}^{(3)}\!R^2\,. \tag{49}$$

We now turn to study a well known example of scalar-tensor theories of gravity which does not generate equations of motion with higher derivatives and, when restricting to linear perturbations, is contained in the Lagrangian (45).

# **3** Generalised Galileons

In four dimensions, the most general scalar tensor theory having field equations of second order in derivatives is a combination of the following generalized Galileon Lagrangians [21, 25, 26],

$$L_2 = G_2(\phi, X) ,$$
 (50)

$$L_3 = G_3(\phi, X) \Box \phi , \qquad (51)$$

$$L_4 = G_4(\phi, X)R - 2G_{4X}(\phi, X)(\Box \phi^2 - \phi^{;\mu\nu}\phi_{;\mu\nu}) , \qquad (52)$$

$$L_5 = G_5(\phi, X)G_{\mu\nu}\phi^{;\mu\nu} + \frac{1}{3}G_{5X}(\phi, X)(\Box\phi^3 - 3\,\Box\phi\,\phi_{;\mu\nu}\phi^{;\mu\nu} + 2\,\phi_{;\mu\nu}\phi^{;\mu\sigma}\phi^{;\nu}_{;\sigma})\,.$$
(53)

For notational convenience, in this section we mostly indicate covariant differentiation with a semicolon symbol, i.e.  $_{;\mu}$ . Moreover, we have defined  $X \equiv \phi^{;\mu}\phi_{;\mu}$  (note that X is sometimes defined differently, i.e., with a factor of -1/2).

In order to translate the above Lagrangians into our EFT language we will proceed in two steps. First, we will rewrite each of these Lagrangians in terms of 3-dimensional geometrical objects  $(K^{\nu}_{\mu}, ^{(3)}R^{\nu}_{\mu}, \text{etc.})$  so that their unitary gauge expression becomes easily readable. The 3+1 decomposition that we are after loses manifest general covariance but shows straightforwardly the lack of higher *time* derivatives already at the level of the action. The second step will be to compute the corresponding coefficients of the operators (45) by simply inverting the dictionary that we derived in the previous section—eqs (40) and (43):

$$c = -\frac{1}{2} \left( \dot{\mathcal{F}} + L_N \right) + H\dot{L}_{\mathcal{R}} - 2L_{\mathcal{R}}\dot{H} - \ddot{L}_{\mathcal{R}} ,$$

$$\Lambda = -\bar{L} - \frac{1}{2}L_N + \frac{1}{2}\dot{\mathcal{F}} + 3H\mathcal{F} + 2\dot{H}L_{\mathcal{R}} + 6H^2L_{\mathcal{R}} + 5H\dot{L}_{\mathcal{R}} + \ddot{L}_{\mathcal{R}} ,$$

$$f = 2L_{\mathcal{R}}M_*^{-2} ,$$

$$M_2^4 = \frac{1}{4}(L_{NN} + 3L_N + \dot{\mathcal{F}}) - \frac{1}{2}(H\dot{L}_{\mathcal{R}} - 2\dot{H}L_{\mathcal{R}} - \ddot{L}_{\mathcal{R}}) ,$$

$$m_3^3 = 2\dot{L}_{\mathcal{R}} - 2HL_{SN} - L_{KN} = 2\dot{L}_{\mathcal{R}} - \mathcal{B} ,$$

$$m_4^2 = \frac{1}{2} \left( L_{\mathcal{S}} - 2L_{\mathcal{R}} - 2H^2L_{\mathcal{SS}} - 2HL_{\mathcal{SK}} - \frac{1}{2}L_{KK} \right) = \frac{1}{2}(L_{\mathcal{S}} - 2L_{\mathcal{R}}) - \frac{1}{4}\mathcal{A} ,$$

$$\tilde{m}_4^2 = L_{N\mathcal{R}} ,$$
(54)

where we have directly adopted the notation (46) which holds in absence of higher derivatives—more general relations are easily found when eq. (44) is not satisfied.

The main result of this section is that the dynamics of linear perturbations for all generalized Galileons is described by (45), with the further restriction  $m_4^2 = \tilde{m}_4^2$ . This is in agreement with the result [25,26] that also higher *space* derivatives are absent from the equations of motion. This section is rather technical; the reader uninterested in the details of the calculations can skip the following subsections and go directly to Sec. 3.5 where we summarize our main results.

#### 3.1 Geometric preliminaries

In order to express in unitary gauge terms of increasing complexity, it is useful to review the geometric formalism adapted to the 3 + 1 decomposition and separate the quantities into "orthogonal" and "parallel" to the hypersurface  $\phi = const$ . First, we define the future directed unitary vector orthogonal to the hypersurface. Up to a factor  $\gamma$ , this is proportional to the gradient of  $\phi$ ,

$$n_{\mu} = -\gamma \phi_{;\mu}, \qquad \gamma = \frac{1}{\sqrt{-X}}.$$
(55)

The metric induced on the  $\phi = const$ . hypersurface is  $h_{\mu\nu} = n_{\mu}n_{\nu} + g_{\mu\nu}$ . Orthogonal to  $n_{\mu}$  are also various quantities that "live" on the hypersurface, in the sense that they vanish when contracted with  $n_{\mu}$ : the extrinsic curvature and the "acceleration" vector

$$K_{\mu\nu} = h^{\sigma}_{\mu} n_{\nu;\sigma}, \qquad \dot{n}_{\mu} = n^{\nu} n_{\mu;\nu} \,. \tag{56}$$

The last two equations can be inverted by decomposing the derivative of  $n_{\mu}$  into parallel and parallel/orthogonal components,

$$n_{\nu;\mu} = K_{\mu\nu} - n_{\mu} \dot{n}_{\nu} \,. \tag{57}$$

By means of the quantities just defined, we can decompose the second derivative of the scalar field as

$$\phi_{;\mu\nu} = -\gamma^{-1}(K_{\mu\nu} - n_{\mu}\dot{n}_{\nu} - n_{\nu}\dot{n}_{\mu}) + \frac{\gamma^{2}}{2}\phi^{;\lambda}X_{;\lambda}n_{\mu}n_{\nu}.$$
(58)

Again, this decomposition into parallel and orthogonal quantities is useful when calculating complicated products such  $\phi_{;\mu\nu} \phi^{;\nu\sigma} \phi^{;\mu}_{;\sigma}$  that appear in  $L_5$ , see eq. (53).

Other relevant equations are the Gauss-Codazzi equations, relating the Ricci tensor and scalar intrinsic to the hypersurface,  ${}^{(3)}R_{\mu\nu}$  and  ${}^{(3)}R$ , to the four-dimensional ones [28, 29],

$${}^{(3)}R_{\mu\nu} = (R_{\mu\nu})_{\parallel} + (n^{\sigma}n^{\rho}R_{\mu\sigma\nu\rho})_{\parallel} - KK_{\mu\nu} + K_{\mu\sigma}K^{\sigma}_{\ \nu}, \tag{59}$$

$$^{(3)}R = R + K^2 - K_{\mu\nu}K^{\mu\nu} - 2(Kn^{\mu} - \dot{n}^{\mu})_{;\mu}, \qquad (60)$$

where the symbol  $\parallel$  means projection on the hypersurface of all tensor indices, e.g.  $(V_{\mu})_{\parallel} \equiv h_{\mu}^{\nu} V_{\nu}$ .

## **3.2** *L*<sub>3</sub>

Since  $L_2$  is trivial, following [11] we start from  $L_3$  and see how to rewrite it in the EFT of dark energy formalism. First, it is convenient to define an auxiliary function  $F_3(\phi, X)$  such that

$$G_3 \equiv F_3 + 2XF_{3X} . (61)$$

Thus,  $L_3$  in eq. (51) can be written as

$$L_3 = F_3 \Box \phi + 2X F_{3X} \Box \phi . \tag{62}$$

We integrate by parts the first term on the right-hand side and we rewrite the second term using  $\Box \phi = -\gamma^{-1}K + \frac{1}{2}\phi^{;\mu}X_{;\mu}/X$ , which is obtained by tracing eq. (58). This yields

$$L_3 = -(F_{3X}X_{;\mu} + F_{3\phi}\phi_{;\mu})\phi^{;\mu} - 2X\gamma^{-1}F_{3X}K + F_{3X}X_{;\mu}\phi^{;\mu}.$$
(63)

After noticing that the first term inside the parenthesis cancels with the last one we finally obtain an expression for  $L_3$  which is of the form of eq. (12),

$$L_3 = 2(-X)^{3/2} F_{3X} K - X F_{3\phi} , \qquad (64)$$

where we have used  $\gamma = 1/\sqrt{-X}$ . In unitary gauge  $\phi(t, \vec{x}) = \phi_0(t)$ , which implies, for instance,

$$F_{3X}(\phi, X) \to F_{3X}(\phi_0(t), -\dot{\phi}_0^2(t)/N^2)$$
 (65)

Using eq. (54), it is now straightforward to derive the corresponding EFT parameters in terms of the Lagrangian parameters evaluated on the background. They are explicitly given in App. C and coincide with those given in [11]. They only depend on four Lagrangian parameters,  $G_{3\phi}$ ,  $G_{3X}$ ,  $G_{3X\phi}$  and  $G_{3XX}$ , so that the dependence on the auxiliary function  $F_3$  disappears. As expected from eq. (64), f,  $m_4^2$  and  $\tilde{m}_4^2$  all vanish: in order to describe  $L_3$  we only need c,  $\Lambda$ ,  $M_2^4$  and  $m_3^3$ .

## **3.3** *L*<sub>4</sub>

We now proceed with  $L_4$ , defined in eq. (52). Using eq. (58) and its trace we can rewrite this as

$$L_{4} = G_{4}R - 2G_{4X} \left[ \left( \gamma^{-1}K + \frac{\gamma^{2}}{2} \phi^{;\mu} X_{;\mu} \right)^{2} - \gamma^{-2} (K_{\mu\nu} K^{\mu\nu} - 2\dot{n}_{\mu} \dot{n}^{\mu}) - \frac{\gamma^{4}}{4} (\phi^{;\mu} X_{;\mu})^{2} \right]$$

$$= G_{4}R + 2XG_{4X} (K^{2} - K_{\mu\nu} K^{\mu\nu}) + 2G_{4X} X_{;\mu} (Kn^{\mu} - \dot{n}^{\mu}) ,$$
(66)

where in the second line we have used that  $\gamma^{-2} = -X$ . Moreover, for the last term we have replaced  $\gamma^{-1}\phi^{;\mu}$  by  $-n^{\mu}$  and used  $\dot{n}_{\mu} = \frac{\gamma^2}{2}h_{\mu}^{\ \nu}X_{;\nu}$ . In this last term we can employ that  $G_{4X}X_{;\mu} = \partial_{\mu}G_4 - G_{4\phi}\phi_{\mu} = \partial_{\mu}G_4 + \gamma^{-1}G_{4\phi}n_{\mu}$ . After an integration by parts this yields, using  $n_{\mu}\dot{n}^{\mu} = 0$ ,

$$2G_{4X}X_{;\mu}(Kn^{\mu} - \dot{n}^{\mu}) = -2G_4(Kn^{\mu} - \dot{n}^{\mu})_{;\mu} - 2\gamma^{-1}G_{4\phi}K.$$
(67)

The first term on the right-hand side of this expression can be rewritten by using the Gauss-Codazzi equation (59). Plugging all this into the second line of eq. (66) we finally obtain  $L_4$  in 3+1 decomposition,

$$L_4 = G_4^{(3)}R + (2XG_{4X} - G_4)(K^2 - K_{\mu\nu}K^{\mu\nu}) - 2\sqrt{-X}G_{4\phi}K.$$
 (68)

It is now lengthy but straightforward to apply our usual dictionary (54) to derive the corresponding EFT parameters. Their explicit expression can be found in App. C. They depend on the six Lagrangian parameters  $G, G_{4X}, G_{4X\phi}, G_{4XX}, G_{4XX\phi}$  and  $G_{4XXX}$ . We need all the seven parameters of the EFT action (45) to describe  $L_4$  but the last two are equal,  $m_4^2 = \tilde{m}_4^2$ .

## **3.4** *L*<sub>5</sub>

This Galileon Lagrangian is more involved than the others. Let us start working on the first term on the right-hand side of eq. (53),  $G_5 G_{\mu\nu} \phi^{;\mu\nu}$ . Integrating it by parts gives

$$G_5 G_{\mu\nu} \phi^{;\mu\nu} = -G_{5X} X^{;\nu} \phi^{;\mu} G_{\mu\nu} - G_{5\phi} \gamma^{-2} G_{\mu\nu} n^{\mu} n^{\nu} .$$
<sup>(69)</sup>

At this stage, as we did for  $L_3$ , it is convenient to define an auxiliary function  $F_5(\phi, X)$ , such that

$$G_{5X} \equiv F_{5X} + \frac{F_5}{2X} , (70)$$

and use this definition to integrate by parts the term proportional to  $G_{5X}$  in (69). In particular, using that

$$G_{5X}X_{;\rho} = \gamma \nabla_{\rho}(\gamma^{-1}F_5) + F_{5\phi}\gamma^{-1}n_{\rho} , \qquad (71)$$

we obtain

$$G_5 G_{\mu\nu}\phi^{;\mu\nu} = F_5\phi^{;\mu\nu}G_{\mu\nu} + \gamma^{-2}(F_{5\phi} - G_{5\phi})G_{\mu\nu}n^{\mu}n^{\nu} - \frac{\gamma}{2}F_5X^{;\mu}n^{\nu}G_{\mu\nu} .$$
(72)

Let us now work on the second term on the right-hand side of eq. (53). Using eq. (58) we can rewrite this as

$$\frac{1}{3}G_{5X}(\Box\phi^3 - 3\,\Box\phi\phi_{;\mu\nu}\phi^{;\mu\nu} + 2\phi_{;\mu\nu}\phi^{;\mu\sigma}\phi^{;\nu}_{;\sigma}) = -G_{5X}\frac{\gamma^{-3}}{3}\mathcal{K} + G_{5X}\mathcal{J} , \qquad (73)$$

where

$$\mathcal{K} \equiv K^{3} - 3KK_{\mu\nu}K^{\mu\nu} + 2K_{\mu\nu}K^{\mu\sigma}K^{\nu}_{\ \sigma}, \qquad (74)$$

$$\mathcal{J} \equiv -\frac{1}{2}\phi^{;\rho}X_{;\rho}(K^2 - K_{\mu\nu}K^{\mu\nu}) - 2\gamma^{-3}(K\dot{n}_{\mu}\dot{n}^{\mu} - K_{\mu\nu}\dot{n}^{\mu}\dot{n}^{\nu}).$$
(75)

The term proportional to  $\mathcal{J}$  on the right-hand side of eq. (73) can be integrated by parts using the same trick as above, which yields

$$G_{5X}\mathcal{J} = -F_5\gamma^{-1}\left(\frac{1}{2}\mathcal{K} + K^{\mu\nu}n^{\sigma}n^{\rho}R_{\mu\sigma\nu\rho} - Kn^{\sigma}n^{\rho}R_{\sigma\rho} + \dot{n}^{\sigma}n^{\rho}R_{\sigma\rho}\right) - \frac{\gamma^{-2}}{2}F_{5\phi}(K^2 - K_{\mu\nu}K^{\mu\nu}).$$
(76)

For the last part of the calculation we need the (one time-)contracted Gauss-Codazzi relation, eq. (59), which gives

$$K^{\mu\nu}G_{\mu\nu} = K^{\mu\nu}{}^{(3)}R_{\mu\nu} - K^{\mu\nu}n^{\sigma}n^{\rho}R_{\mu\sigma\nu\rho} + KK^{2}_{\mu\nu} - K^{3}_{\mu\nu} - \frac{1}{2}RK.$$
(77)

Replacing  $\phi^{;\mu\nu}$  with eq. (58) in eq. (72) and using this relation, the terms proportional to  $F_5$  in eqs. (72) and (76) combine and simplify to

$$-\gamma^{-1}F_5\left({}^{(3)}G_{\mu\nu}K^{\mu\nu} - \frac{1}{6}\mathcal{K}\right) \,. \tag{78}$$

Using this and putting together all the terms of  $L_5$  from eqs. (72), (73) and (76) we finally obtain

$$L_{5} = -\sqrt{-X}F_{5}\left(K^{\mu\nu(3)}R_{\mu\nu} - \frac{1}{2}K^{(3)}R\right) - \frac{1}{3}(-X)^{3/2}G_{5X}\mathcal{K} + \frac{1}{2}X(G_{5\phi} - F_{5\phi})^{(3)}R + \frac{1}{2}XG_{5\phi}(K^{2} - K_{\mu\nu}K^{\mu\nu}), \qquad (79)$$

where in the last line we have used

$$2G_{\mu\nu}n^{\mu}n^{\nu} = {}^{(3)}R + K^2 - K_{\mu\nu}K^{\mu\nu} .$$
(80)

Operator	f	Λ	с	$M_2^4$	$m_{3}^{3}$	$m_4^2 = \tilde{m}_4^2$
$L_2$	0	$\checkmark$	$\checkmark$	$\checkmark$	0	0
$L_3$	0	$\checkmark$	$\checkmark$	$\checkmark$	$\checkmark$	0
$L_4$	$\checkmark$	$\checkmark$	$\checkmark$	$\checkmark$	$\checkmark$	$\checkmark$
$L_5$	$\checkmark$	$\checkmark$	$\checkmark$	$\checkmark$	$\checkmark$	$\checkmark$

Table 1: A list of the different contributions of the generalized Galileon Lagrangians (50)–(53) to the operators of (45).

Note that the last line of (79) has the same form as the first two terms of  $L_4$  given in eq. (68): by using eq. (70) it can be written as

$$G_4^{(3)}R + (2XG_{4X} - G_4)(K^2 - K_{\mu\nu}K^{\mu\nu}), \qquad (81)$$

with  $G_4 \equiv \frac{1}{2}X(G_{5\phi} - F_{5\phi}).$ 

In order to compute the coefficients of the various EFT operators we use the dictionary (54). To treat the term  $K^{\mu\nu(3)}R_{\mu\nu}$  we employ the prescription described by eq. (126) in App. A. Moreover, it is useful to notice that, up to quadratic order, the combination  $\mathcal{K}$  of the extrinsic curvature tensor can be replaced by an expression that depends only on S and K:

$$\mathcal{K} = 6H^3 - 6H^2K + 3HK^2 - 3HS + \mathcal{O}(3).$$
(82)

The EFT operator coefficients are explicitly given in App. C. One finds that they depend on the six Lagrangian parameters  $G_{5\phi}$ ,  $G_{5X}$ ,  $G_{5X\phi}$ ,  $G_{5XX}$ ,  $G_{5XX\phi}$  and  $G_{5XXX}$ —the dependence on  $F_5$  explicitly cancels out—and, as in the case of  $L_4$ ,  $m_4^2 = \tilde{m}_4^2$ . Thus, at linear order in the perturbations quadratic in the action— $L_5$  does not bring any new operator with respect to  $L_4$ . The difference between  $L_4$  and  $L_5$  appears at the cubic order in the action.

### 3.5 Summary

We have established a dictionary between the generalized Galileon theory, eqs. (50)–(53), and the EFT of dark energy parameters entering the action (2). Such a dictionary is explicitly given in App. C. As summarised in Table 1, the EFT operators and their associated time-dependent parameters that are needed to describe the generalized Galileons are only six: c,  $\Lambda$  and f, the three usual parameters already present at the background level, and  $M_2^4$ ,  $m_3^3$ ,  $m_4^2 = \tilde{m}_4^2$ , progressively appearing in  $L_2$ ,  $L_3$ ,  $L_4$  and  $L_5$ , contributing only to the perturbations. As already stressed, at quadratic order in the perturbations,  $L_4$  contains the same number of independent operators as  $L_5$ —in particular, only the combination  $m_4^2 = \tilde{m}_4^2$  appears in the action. The case  $m_4^2 \neq \tilde{m}_4^2$  encompasses the generalized Galileon class.<sup>8</sup> When  $m_4^2 \neq \tilde{m}_4^2$ , higher derivatives are expected to appear beyond linear order. However, the effect of these higher derivatives can be ignored as long as perturbations remain small and linear theory is a good approximation.

## 4 Observables

Observables describing large scale structures are computed in the framework of linear cosmological perturbation theory. In this section we first derive the perturbation equations describing the dynamics of dark energy and modified gravity. We include a matter sector describing cosmological species such

<sup>&</sup>lt;sup>8</sup>Note that our formalism easily applies to nonlinear extensions of Hordenskis theories, such as described in Ref. [30]. In this particular case, one finds that  $m_4^2 \neq \tilde{m}_4^2$  but the quadratic action contains higher order spatial derivatives.

as cold dark matter, baryons, photons and neutrinos—by adding the matter Lagrangian  $\mathcal{L}_m(g_{\mu\nu}, \psi_m)$  to eq. (2), so that the final Jordan frame action in unitary gauge reads

$$S = \int d^4x \sqrt{-g} \left[ \frac{M_*^2}{2} f(t)R - \Lambda(t) - c(t)g^{00} + \frac{M_2^4(t)}{2} (\delta g^{00})^2 - \frac{m_3^3(t)}{2} \delta K \delta g^{00} - m_4^2(t) \left( \delta K^2 - \delta K_{\mu\nu} \delta K^{\mu\nu} \right) + \frac{\tilde{m}_4^2(t)}{2} {}^{(3)}R \, \delta g^{00} - \bar{m}_4^2(t) \delta K^2 + \frac{\bar{m}_5(t)}{2} {}^{(3)}R \, \delta K + \frac{\bar{\lambda}(t)}{2} {}^{(3)}R^2 + \mathcal{L}_m(g_{\mu\nu}, \psi_m) \right].$$
(83)

We then discuss the modifications of gravity expected in linear theory. We will use Newtonian gauge, which is often used in cosmology, especially to describe cosmological perturbations for modified gravity. Extension to other gauges or to so-called "gauge invariant" formalisms is straightforward.

#### 4.1 Perturbation equations

We will first restore the general covariance of the action above and write it in a generic coordinate system. In order to do that we need to reintroduce the scalar fluctuation  $\pi$  via the Stueckelberg trick [3, 4, 27]. Under the time coordinate change  $t \to t + \pi(t, \vec{x})$ , the four-Ricci scalar R remains invariant, while functions of time such as f and the 3-dimensional quantities change as<sup>9</sup><sup>10</sup>

$$f \to f + \dot{f}\pi + \frac{1}{2}\ddot{f}\pi^2 , \qquad (84)$$

$$g^{00} \to g^{00} + 2g^{0\mu}\partial_{\mu}\pi + g^{\mu\nu}\partial_{\mu}\pi\partial_{\nu}\pi , \qquad (85)$$

$$\delta K_{ij} \to \delta K_{ij} - \dot{H}\pi h_{ij} - \partial_i \partial_j \pi , \qquad (86)$$

$$\delta K \to \delta K - 3\dot{H}\pi - \frac{1}{a^2}\partial^2\pi , \qquad (87)$$

$$^{(3)}R_{ij} \to ^{(3)}R_{ij} + H(\partial_i\partial_j\pi + \delta_{ij}\partial^2\pi) , \qquad (88)$$

$${}^{(3)}R \to {}^{(3)}R + \frac{4}{a^2}H\partial^2\pi$$
 (89)

In the new coordinates we consider a linearly perturbed FLRW metric with only scalar fluctuations,

$$ds^{2} = -(1+2\Phi)dt^{2} + 2\partial_{i}\alpha \, dt dx^{i} + a^{2}(t) \left[(1-2\Psi)\delta_{ij} + 2\chi_{ij}\right] dx^{i} dx^{j} , \qquad (90)$$

where  $\chi_{ij}$  is traceless and given in terms of the scalar perturbation  $\beta$ ,  $\chi_{ij} \equiv (\partial_i \partial_j - \frac{1}{3} \delta_{ij} \partial^2)\beta$ . The extrinsic curvature and the 3-dimensional Ricci tensor of the *new* equal-time hypersurfaces thus read

$$K_{ij} = e^{-\Phi} (H - \dot{\Psi}) h_{ij} + \dot{\chi}_{ij} - \partial_i \partial_j \alpha , \qquad (91)$$

$${}^{(3)}R_{ij} = \partial_i \partial_j \Psi + \delta_{ij} \partial^2 \Psi + 2\partial_k \partial_{(i} \chi_{j)}^k - \partial^2 \chi_{ij} .$$

We also decompose the matter stress-energy tensor at linear order as

$$T^0_{\ 0} \equiv -(\rho_m + \delta \rho_m) , \qquad (92)$$

$$T^{0}_{\ i} \equiv (\rho_m + p_m)\partial_i v = -a^2 T^{i}_{\ 0} , \qquad (93)$$

$$T^{i}_{\ j} \equiv (p_m + \delta p_m)\delta^{i}_{\ j} + \left(\partial^i \partial_j - \frac{1}{3}\delta^{i}_{\ j}\partial^2\right)\sigma , \qquad (94)$$

<sup>&</sup>lt;sup>9</sup>With an abuse of notation, here we denote the extrinsic curvature on hypersurfaces of constant time with  $K_{ij}$  even when we are *not* in unitary gauge. The reader must be aware that  $K_{ij}$  is not the same geometrical object *before* and *after* the Stueckelberg trick. The same also holds for <sup>(3)</sup> $R_{ij}$ . In particular, after the Stueckelberg trick  $K_{ij}$  and <sup>(3)</sup> $R_{ij}$ are respectively given by eq. (91).

<sup>&</sup>lt;sup>10</sup>The operator  $\tilde{m}_4^2$  is also considered in Ref. [12]. However, in v1 of this reference, the variation (89) of <sup>(3)</sup>R under the Stueckelberg trick has been overlooked and the error propagates into the Einstein equations and the various observables. With the authors of [12] there is now agreement on this issue [31].

where  $\rho_m$  and  $p_m$  are respectively the background energy density and pressure and  $\delta \rho_m$  and  $\delta p_m$  their perturbations, v is the 3-velocity potential and  $\sigma$  the scalar component of the anisotropic stress. The background equations derived from the action (83) are [11]

$$c + \Lambda = 3M_*^2 (fH^2 + \dot{f}H) - \rho_m , \qquad (95)$$

$$\Lambda - c = M_*^2 (2f\dot{H} + 3fH^2 + 2\dot{f}H + \ddot{f}) + p_m .$$
<sup>(96)</sup>

Using these expressions and the transformations (84)-(89) allows us to rewrite (83) as an action for the scalar fluctuations  $\Phi$ ,  $\alpha$ ,  $\Psi$ ,  $\beta$  and  $\pi$ . We can vary eq. (83) expanded at second order and then fix the Newtonian gauge by setting  $\alpha = 0 = \beta$  in the equations derived. This yields five equations,

$$0 = \frac{1}{\sqrt{-g}} \frac{\delta S}{\delta \Phi} \bigg|_{\alpha=0=\beta} \equiv A_{\Phi} \Phi + A_{\dot{\Psi}} \dot{\Psi} + A_{\pi} \pi + A_{\dot{\pi}} \dot{\pi} + \frac{k^2}{a^2} (A_{\Psi}^{(2)} \Psi + A_{\pi}^{(2)} \pi) + \delta T_0^0 , \qquad (97)$$

$$0 = \frac{1}{\sqrt{-g}} \frac{\delta S}{\delta \alpha} \Big|_{\alpha=0=\beta} \equiv k^2 \left[ B_{\Phi} \Phi + B_{\dot{\Psi}} \dot{\Psi} + B_{\pi} \pi + B_{\dot{\pi}} \dot{\pi} + \frac{k^2}{a^2} (B_{\Psi}^{(2)} \Psi + B_{\pi}^{(2)} \pi) \right] - i k^i \delta T^0_{\ i} , \qquad (98)$$

$$0 = \frac{1}{\sqrt{-g}} \frac{\delta S}{\delta \Psi} \bigg|_{\alpha=0=\beta} \equiv C_{\Phi} \Phi + C_{\dot{\Phi}} \dot{\Phi} + C_{\dot{\Psi}} \dot{\Psi} + C_{\ddot{\Psi}} \ddot{\Psi} + C_{\pi} \pi + C_{\dot{\pi}} \dot{\pi} + C_{\ddot{\pi}} \ddot{\pi} \\ + \frac{k^2}{a^2} (C_{\Phi}^{(2)} \Phi + C_{\Psi}^{(2)} \Psi + C_{\pi}^{(2)} \pi + C_{\dot{\pi}}^{(2)} \dot{\pi}) + \frac{k^4}{a^4} (C_{\Psi}^{(4)} \Psi + C_{\pi}^{(4)} \pi) - \delta T_k^k , \quad (99)$$
$$0 = \frac{1}{\sqrt{-g}} \frac{\delta S}{\delta \varphi} \bigg|_{\omega=0=\beta} \equiv \left( k_i k^j - \frac{1}{2} \delta_j^j k^2 \right) \left[ k^i k_j \left( D_{\Phi}^{(2)} \Phi + D_{\Psi}^{(2)} \Psi + D_{\dot{\pi}}^{(2)} \dot{\Psi} + D_{\pi}^{(2)} \pi + D_{\dot{\pi}}^{(2)} \dot{\pi} \right] \right]$$

$$\frac{1}{\sqrt{-g}} \frac{\delta S}{\delta \beta} \Big|_{\alpha=0=\beta} \equiv \left( k_i k^j - \frac{1}{3} \delta_i^j k^2 \right) \left[ k^i k_j \left( D_{\Phi}^{(2)} \Phi + D_{\Psi}^{(2)} \Psi + D_{\Psi}^{(2)} \dot{\Psi} + D_{\pi}^{(2)} \pi + D_{\pi}^{(2)} \dot{\pi} + \frac{k^2}{a^2} (D_{\Psi}^{(4)} \Psi + D_{\pi}^{(4)} \pi) \right) - \delta T_j^i \right],$$
(100)

$$0 = \frac{1}{\sqrt{-g}} \frac{\delta S}{\delta \pi} \Big|_{\alpha=0=\beta} \equiv E_{\Phi} \Phi + E_{\dot{\Phi}} \dot{\Phi} + E_{\Psi} \Psi + E_{\dot{\Psi}} \dot{\Psi} + E_{\ddot{\Psi}} \ddot{\Psi} + E_{\pi} \pi + E_{\dot{\pi}} \dot{\pi} + E_{\ddot{\pi}} \ddot{\pi} + \frac{k^2}{a^2} (E_{\Phi}^{(2)} \Phi + E_{\Psi}^{(2)} \Psi + E_{\psi}^{(2)} \dot{\Psi} + E_{\pi}^{(2)} \pi) + \frac{k^4}{a^4} (E_{\Psi}^{(4)} \Psi + E_{\pi}^{(4)} \pi) .$$
(101)

The coefficients  $A_a, B_a, C_a, D_a$  and  $E_a$  of these equations are detailed in App. D.

The explicit expressions of the above coefficients given in the appendix contain also higherderivative terms—those proportional to  $\bar{m}_4^2$ ,  $\bar{m}_5$  and  $\bar{\lambda}$ . To compare with the usual Einstein equations, here we rewrite these equations by replacing the components of the stress-energy tensor  $T^{\mu}_{\nu}$ with their expressions given in eqs. (92)–(94). For simplicity, we set  $\bar{m}_4^2 = \bar{m}_5 = \bar{\lambda} = 0$ . We obtain:

• 00-component  $(\delta S/\delta \Phi = 0)$ :

$$M_*^2 \left[ -2f\left(\frac{k^2}{a^2}\Psi + 3H\dot{\Psi} + 3H^2\Phi\right) + \dot{f}\left(\frac{k^2}{a^2}\pi + 3H^2\pi - 3H(\Phi - \dot{\pi}) - 3(\dot{\Psi} + H\Phi)\right) + 3H\ddot{f}\pi \right] - (\dot{c} + \dot{\Lambda})\pi + (2c + 4M_2^4 + 3Hm_3^3)(\Phi - \dot{\pi}) + (m_3^3 - 4Hm_4^2)\left[-\frac{k^2}{a^2}\pi + 3(H\Phi + \pi\dot{H} + \dot{\Psi})\right] - 4\frac{k^2}{a^2}\tilde{m}_4^2(\Psi + H\pi) = \delta\rho_m .$$
(102)

• 0*i*-component 
$$(\delta S/\delta \alpha = 0)$$
:

$$M_*^2 \left[ (H\dot{f} - \ddot{f})\pi + \dot{f} (\Phi - \dot{\pi}) + 2f(H\Phi + \dot{\Psi}) \right] - 2c\pi - m_3^3 (\Phi - \dot{\pi}) + 4m_4^2 (H\Phi + \dot{\Psi} + \dot{H}\pi)$$
  
=  $- (p_m + \rho_m) v$ . (103)

• *ij*-trace component  $(\delta S/\delta \Psi = 0)$ :

$$\begin{split} M_{*}^{2} \bigg\{ 2f \left[ -\frac{1}{3} \frac{k^{2}}{a^{2}} (\Phi - \Psi) + (3H^{2} + 2\dot{H})\Phi + H(\dot{\Phi} + 3\dot{\Psi}) + \ddot{\Psi} \right] \\ + \dot{f} \left[ -\frac{2}{3} \frac{k^{2}}{a^{2}} \pi + 2H\Phi + 2H(\Phi - \dot{\pi}) - (3H^{2} + 2\dot{H})\pi + 2\dot{\Psi} + \dot{\Phi} - \ddot{\pi} \right] \\ + \ddot{f} \left[ -2H\pi + 2(\Phi - \dot{\pi}) \right] - f^{(3)}\pi \bigg\} + (\dot{\Lambda} - \dot{c})\pi + 2c(\Phi - \dot{\pi}) \end{split}$$
(104)  
$$- \frac{4}{3} \frac{k^{2}}{a^{2}} \left[ \tilde{m}_{4}^{2} (\Phi - \dot{\pi}) + \left( Hm_{4}^{2} + (m_{4}^{2})^{\cdot} \right) \pi + m_{4}^{2} \dot{\pi} \right] \\ + 4(\dot{H}m_{4}^{2})^{\cdot} \pi + 4m_{4}^{2}\dot{H}\dot{\pi} - \left[ (m_{3}^{3})^{\cdot} + 3Hm_{3}^{3} \right] (\Phi - \dot{\pi}) - m_{3}^{3} (\dot{\Phi} - \ddot{\pi}) \\ + 4 \left[ H(m_{4}^{2})^{\cdot} + 3H^{2}m_{4}^{2} + \dot{H}m_{4}^{2} \right] \Phi + 4(m_{4}^{2})^{\cdot} \dot{\Psi} + 4m_{4}^{2}H(3\dot{H}\pi + \dot{\Phi} + 3\dot{\Psi}) + 4m_{4}^{2}\ddot{\Psi} = \delta p_{m} \,. \end{split}$$

• *ij*-traceless component  $(\delta S/\delta\beta = 0)$ :

$$M_*^2 \left[ f(\Phi - \Psi) + \dot{f}\pi \right] + 2 \left[ m_4^2 \dot{\pi} + m_4^2 H \pi + (m_4^2) \dot{\pi} \right] + 2 \tilde{m}_4^2 (\Phi - \dot{\pi}) = \sigma .$$
(105)

By combining eqs. (97) and (98) we obtain the relativistic generalization of the Poisson equation,

$$F_{\Phi}\Phi + F_{\dot{\Psi}}\dot{\Psi} + F_{\pi}\pi + F_{\dot{\pi}}\dot{\pi} + \frac{k^2}{a^2}(F_{\Psi}^{(2)}\Psi + F_{\pi}^{(2)}\pi) = \delta\rho_m - 3H(\rho_m + p_m)v \equiv \rho_m\Delta_m , \qquad (106)$$

which can be also written as:

#### • Generalized Poisson equation:

$$-\frac{k^2}{a^2} \left[ (2fM_*^2 + 4\tilde{m}_4^2)\Psi - (\dot{f}M_*^2 - m_3^3 + 4Hm_4^2 - 4H\tilde{m}_4^2)\pi \right] + (6M_*^2H^2\dot{f} - 6Hc - \dot{c} - \dot{\Lambda} + 3m_3^3\dot{H})\pi - (2c + 4M_2^4)\dot{\pi} - (3M_*^2H\dot{f} - 2c - 4M_2^4)\Phi - 3M_*^2\dot{f}\dot{\Psi} + 3m_3^3(\dot{\Psi} + H\Phi) = \rho_m\Delta_m .$$
(107)

Note that when  $m_4^2 \neq \tilde{m}_4^2$  some of the equations contain terms with higher derivates: for instance, the terms with  $k^2 \dot{\pi}$  in eq. (104), fourth line, and those with  $\dot{\pi}$  in eq. (105). However, the scalar propagating degree of freedom satisfies a second order equation. Indeed, one can use eq. (105) to remove the higher derivative terms from eq. (104) and derive a purely second order equation for  $\Psi$ . This is even clearer in unitary gauge, where higher derivative are explicitly absent—see analysis of Sec. (2.2).

### 4.2 Modification of gravity

In order to derive the effective Newton constant,  $G_{\text{eff}}$ , we consider the quasi static approximation, i.e. we neglect the time derivatives in the equations of motion and we neglect the anisotropic stress,  $\sigma = 0$  in eq. (100). This is a good approximation for scales much smaller than the sound horizon scale, i.e. for  $k \gg aH/c_s$ . For models with small or vanishing sound speed (see e.g. [10]) or on scales longer than the sound horizon, a consistent treatment which takes into account the time derivatives should be undertaken.

In the quasi-static limit,  $G_{\text{eff}}$  is defined by

$$-\frac{k^2}{a^2}\Phi \equiv 4\pi G_{\text{eff}}(t,k)\rho_m \Delta_m .$$
(108)

Following [32, 33], in order to write the Poisson equation in this form we can use eqs. (100), (101) and (106). For  $c_s \sim \mathcal{O}(1)$ , we can neglect  $D_{\Phi}$ ,  $D_{\Psi}$ ,  $D_{\pi}$ ,  $E_{\Phi}$ ,  $E_{\Psi}$ ,  $F_{\Phi}$  and  $F_{\pi}$  from these equations and the effective Newton constant is thus given by

$$4\pi G_{\text{eff}} = -[\mathcal{M}^{-1}]_{13} , \qquad \mathcal{M} \equiv \begin{pmatrix} D_{\Phi}^{(2)} & D_{\Psi}^{(2)} + D_{\Psi}^{(4)}(k/a)^2 & D_{\pi}^{(2)} + D_{\pi}^{(4)}(k/a)^2 \\ E_{\Phi}^{(2)} & E_{\Psi}^{(2)} + E_{\Psi}^{(4)}(k/a)^2 & E_{\pi}(k/a)^{-2} + E_{\pi}^{(2)} + E_{\pi}^{(4)}(k/a)^2 \\ 0 & F_{\Psi}^{(2)} & F_{\pi}^{(2)} \end{pmatrix} .$$
(109)

We can write it in a slightly more compact form as

$$4\pi G_{\rm eff}(k) = \frac{a_{-2}(k/a)^{-2} + a_0 + a_2(k/a)^2 + a_4(k/a)^4}{b_{-2}(k/a)^{-2} + b_0 + b_2(k/a)^2} , \qquad (110)$$

where

$$\begin{aligned} a_{-2} &= D_{\Psi}^{(2)} E_{\pi} ,\\ a_{0} &= D_{\Psi}^{(2)} E_{\pi}^{(2)} - D_{\pi}^{(2)} E_{\Psi}^{(2)} + D_{\Psi}^{(4)} E_{\pi} ,\\ a_{2} &= D_{\Psi}^{(2)} E_{\pi}^{(4)} - D_{\pi}^{(4)} E_{\Psi}^{(2)} - D_{\pi}^{(2)} E_{\Psi}^{(4)} + D_{\Psi}^{(4)} E_{\pi}^{(2)} ,\\ a_{4} &= -D_{\pi}^{(4)} E_{\Psi}^{(4)} + D_{\Psi}^{(4)} E_{\pi}^{(4)} ,\\ b_{-2} &= D_{\Phi}^{(2)} E_{\pi} F_{\Psi}^{(2)} ,\\ b_{0} &= D_{\Psi}^{(2)} E_{\Phi}^{(2)} F_{\pi}^{(2)} - D_{\Phi}^{(2)} E_{\Psi}^{(2)} F_{\pi}^{(2)} - D_{\pi}^{(2)} E_{\Phi}^{(2)} F_{\Psi}^{(2)} + D_{\Phi}^{(2)} E_{\pi}^{(2)} F_{\Psi}^{(2)} ,\\ b_{2} &= -D_{\Phi}^{(2)} E_{\Psi}^{(4)} F_{\pi}^{(2)} - D_{\pi}^{(4)} E_{\Phi}^{(2)} F_{\Psi}^{(2)} + D_{\Phi}^{(2)} E_{\pi}^{(4)} F_{\Psi}^{(2)} + D_{\Psi}^{(4)} E_{\Phi}^{(2)} F_{\pi}^{(2)} . \end{aligned}$$

Another quantity often used to parameterize deviations from General Relativity is the ratio between the gravitational potentials  $\gamma \equiv \Psi/\Phi$ , which is given by

$$\gamma = \frac{[\operatorname{com}(\mathcal{M})]_{32}}{[\operatorname{com}(\mathcal{M})]_{31}}, \qquad (112)$$

where  $com(\mathcal{M})$  denotes the comatrix of  $\mathcal{M}$ . This reads

$$\gamma = \frac{c_{-2}(k/a)^{-2} + c_0 + c_2(k/a)^2}{a_{-2}(k/a)^{-2} + a_0 + a_2(k/a)^2 + a_4(k/a)^4},$$
(113)

with

$$c_{-2} = -D_{\Phi}^{(2)} E_{\pi} , \qquad (114)$$

$$c_0 = D_\pi^{(2)} E_\Phi^{(2)} - D_\Phi^{(2)} E_\pi^{(2)} , \qquad (115)$$

$$c_2 = D_{\pi}^{(4)} E_{\Phi}^{(2)} - D_{\Phi}^{(2)} E_{\pi}^{(4)} .$$
(116)

The expressions for  $G_{\text{eff}}$  and  $\gamma$ , eqs. (110) and (113), generalize those given for instance in [32] in absence of higher derivative operators, in which case  $a_2 = a_4 = b_2 = c_2 = 0$ . When also  $a_{-2} = b_{-2} = c_{-2} = 0$  we recover the results of [11]. Finally, we note that the numerator of  $G_{\text{eff}}$  equals the denominator of  $\gamma$ , which confirms the results of Ref. [33]<sup>11</sup>.

<sup>11</sup>It simply follows from  $[\mathcal{M}^{-1}]_{13} = (\det \mathcal{M})^{-1} [\operatorname{com}(\mathcal{M})]_{31}$ .

# 5 Conclusion

In this paper we lay down the basic building blocks for a systematic phenomenological study of dark energy and its cosmological perturbations. Following [11], our basic assumptions are that a) dark energy/modified gravity brings in at most one scalar propagating degree of freedom and that b) the weak equivalence principle is satisfied—there exists a metric tensor universally coupled to matter. We use the effective field theory formalism developed for inflation in [3,4], that is based on an expansion in number of perturbations rather than in number of fields. Indeed, expanding the action in number of fields [15, 16] becomes unpractical every time that the background field configuration undergoes a large excursion. On the opposite, the main advantage of the present (non-covariant) approach is that an expansion in number of perturbations can always be consistently truncated at the desired order of approximation, in virtue of the empirical fact that perturbations are small on the largest scales. Moreover, our formalism is "ready to go", in the sense that there is no need of solving for the background equations first. Apart from the three operators f, c and  $\Lambda$  responsible for the background evolution [11], every new operator is at least quadratic in the perturbations: it does not affect the background and its dynamical effects can be studied independently.

In particular, we consider only operators that are at most quadratic in the number of perturbations—those needed for the linearized equations of motion—and we single out a set of seven operators that bring up to two derivatives in the equations of motion. To achieve this result, in Sec. 2 we provide a systematic treatment of any Lagrangian that can be written in ADM form as a general function of extrinsic and intrinsic 3-dimensional curvature tensors and of the lapse function. This is already enough to avoid higher time derivatives in the equations of motion. Then, in Sec. 2.3 we identify specific combinations of the EFT operators that are required to avoid higher-order spatial derivatives. Some operators can be re-expressed into other ones, thus simplifying the EFT Lagrangians up to quadratic order.

The entire Horndeski, or "generalized Galileon", theory can be written in this formalism (Sec. 3): a relevant amount of work has gone into re-expressing all the generalized Galileon Lagrangians in their ADM form and obtaining their EFT formulation. At linear order, Horndeski theories can be described by a total of six operators: only three quadratic operators in addition to those—f, c and  $\Lambda$ —accounting for the background (see eq. (2) with  $m_4^2 = \tilde{m}_4^2$ ). This seems a substantial simplification if compared to the full covariant treatment and well represents the power of the noncovariant EFT approach. The two Galileon Lagrangians  $L_4$  and  $L_5$ , despite their scaring looks (52)-(53), are affordable at linear order in the perturbations with the addition of just one operator with respect to those needed for  $L_3$ .

At linear order, Horndeski theory is *not* the most general scalar-tensor theory with second-order dynamics. Indeed, for  $m_4^2 \neq \tilde{m}_4^2$  there exists another operator beyond Horndeski that in unitary gauge gives equations of motion limited to second order in time and space derivatives. In some gauges (for instance in Newtonian gauge, see Sec. 4.1), this operator generates higher derivatives in the equations of motion but one can show that the dynamics of the propagating degree of freedom remains second order. [At linear order, there exists another operator beyond the Horndeski theory (for  $m_4^2 \neq \tilde{m}_4^2$ ) that still gives equations of motion limited to second order in time and space derivatives.] Finally, we analyze also some higher *spatial* derivative operators, those in eq. (4).

The time dependent coefficients of our seven plus three operators described by actions (2) and (4) remain to be constrained or measured by observations. Indeed, in Sec. 4.1 we provide the set of linear perturbation equations in Newtonian gauge by varying these actions with respect to scalar metric and field fluctuations in a generic gauge. As an illustration, using these equations we compute the effective Newton constant in the quasi-static approximation and the ratio between the two gravitational potentials (Sec. 4.2). The computation of these "observables" should be considered as a first step towards a more general and systematic study of the impact of dark energy on cosmological perturbations in order to fully exploit future observational data.

# Acknowledgments

Conversations and/or correspondence with Jolyon Bloomfield, Giulia Gubitosi, Ignacy Sawicki, Lorenzo Sorbo, Shinji Tsujikawa and George Zahariade are gratefully acknowledged. D.L. is partly supported by the ANR (Agence Nationale de la Recherche) grant STR-COSMO ANR-09-BLAN-0157-01. F.V. is partially supported by the ANR *Chaire d'excellence* CMBsecond ANR-09-CEXC-004-01.

# A Lagrangian dependence on ${}^{(3)}R_{\mu\nu}K^{\mu\nu}$

In this appendix we show how to treat a dependence on

$$\mathcal{Y} \equiv {}^{(3)}R_{\mu\nu}K^{\mu\nu} \tag{117}$$

in the unitary gauge Lagrangian.

Let us first show the relation

$$\int d^4x \sqrt{-g}\,\lambda(t)^{(3)}R_{\mu\nu}K^{\mu\nu} = \int d^4x \sqrt{-g} \left[\frac{\lambda(t)}{2}{}^{(3)}R K + \frac{\dot{\lambda}(t)}{2N}{}^{(3)}R\right] , \qquad (118)$$

or, equivalently,

$$\int d^4x \sqrt{-g} \left[ \lambda(t)^{(3)} R_{\mu\nu} K^{\mu\nu} - \frac{\lambda(t)}{2}{}^{(3)} R K - \frac{\dot{\lambda}(t)}{2N}{}^{(3)} R \right] = 0 , \qquad (119)$$

up to some irrelevant boundary terms. Since  $K = \nabla_{\mu} n^{\mu}$ , the last two terms in the above integral can be simplified via an integration by parts, so that the expression reduces to

$$\int d^4x \sqrt{-g}\,\lambda(t)\,\left({}^{(3)}R_{\mu\nu}K^{\mu\nu} + \frac{n^{\mu}}{2}\nabla_{\mu}{}^{(3)}R\right)\,.$$
(120)

It remains to show that this can be written as the integral of a total derivative.

Using the explicit expressions for the extrinsic curvature in the ADM decomposition, eq. (22), and  $n^{\mu} = -Ng^{0\mu}$ , the above expression can be rewritten as

$$\int d^4x \sqrt{h}\lambda(t) \left[ \frac{1}{2} \left( h^{ik} h^{jl} \dot{h}_{kl}{}^{(3)}R_{ij} + {}^{(3)}\dot{R} \right) - \nabla^i N^{j}{}^{(3)}R_{ij} - \frac{1}{2} N^i \nabla_i{}^{(3)}R \right] , \qquad (121)$$

where  $\nabla_i$  is the covariant derivative with respect to the three-metric  $h_{ij}$ . The second term can be integrated by parts and then vanishes when combined with the last term, as a consequence of the Bianchi identity  $\nabla^{i(3)}G_{ij} = 0$ . Finally, the term in parenthesis can be rewritten as

$$h^{ik}h^{jl}\dot{h}_{kl} {}^{(3)}R_{ij} + {}^{(3)}\dot{R} = h^{ik}h^{jl}\dot{h}_{kl} {}^{(3)}R_{ij} + \dot{h}^{ij}{}^{(3)}R_{ij} + h^{ij}{}^{(3)}\dot{R}_{ij} = h^{ij}{}^{(3)}\dot{R}_{ij} .$$
(122)

and it is known that the last expression can be reexpressed as the divergence of a three-vector, i.e.  $h^{ij} {}^{(3)}\dot{R}_{ij} = \nabla_i J^i$  (the very same property is used to derive Einstein's equations from the Einstein-Hilbert action<sup>12</sup>). We have thus proved Eq. (118).

Let us now assume that the Lagrangian L introduced in Eq. (6) also contains an explicit dependence on  $\mathcal{Y}$ . By noting that  $\mathcal{Y}$  is already a perturbative quantity, i.e. vanishes in the background, and can be decomposed as

$$\mathcal{Y} = H\mathcal{R} + {}^{(3)}R_{\mu\nu}\delta K^{\mu\nu} \,, \tag{123}$$

where the first term on the right hand side is a first (and higher) order quantity while the second term is only second order, one immediately finds that the expansion of the Lagrangian, up to quadratic

 $<sup>^{12}</sup>$ See for instance Eq. (7.5.14) of Ref. [29].

order, will yield the following extra terms with respect to the expression (12) obtained in the main body:

$$L(N, \mathcal{S}, K, \mathcal{R}, \mathcal{Y}, \mathcal{Z}) \supset L_{\mathcal{Y}}\mathcal{Y} + (L_{N\mathcal{Y}}\delta N + L_{K\mathcal{Y}}\delta K + L_{\mathcal{S}\mathcal{Y}}\delta \mathcal{S} + L_{\mathcal{R}\mathcal{Y}}\delta \mathcal{R}) H\delta \mathcal{R} + \frac{1}{2}L_{\mathcal{Y}\mathcal{Y}}H^2\delta \mathcal{R}^2 .$$
(124)

The first term can be expressed in terms of  $\mathcal{R}$  and K by using eq. (118) with  $\lambda = L_{\mathcal{Y}}$ . Expanding up to second order then yields

$$L_{\mathcal{Y}}\mathcal{Y} = \frac{1}{2} \left( \dot{L}_{\mathcal{Y}} + 3HL_{\mathcal{Y}} \right) \delta \mathcal{R} + \frac{1}{2} \left( L_{\mathcal{Y}} \delta K - \dot{L}_{\mathcal{Y}} \delta N \right) \delta \mathcal{R} + \mathcal{O}(3) + \text{boundary terms}, \quad (125)$$

so that the expansion of the full Lagrangian now reads

$$L(N, \mathcal{S}, K, \mathcal{R}, \mathcal{Y}, \mathcal{Z}) = \bar{L} - \dot{\mathcal{F}} - 3H\mathcal{F} + L_N \,\delta N + \frac{1}{2} \left( 2L_{\mathcal{R}} + \dot{L}_{\mathcal{Y}} + 3HL_{\mathcal{Y}} \right) \delta \mathcal{R} + L_{\mathcal{S}} \,\delta K^{\mu}_{\nu} \delta K^{\nu}_{\mu} + \left( 2H^2 L_{\mathcal{S}\mathcal{S}} + 2HL_{\mathcal{S}K} + \frac{1}{2}L_{KK} \right) \delta K^2 + \frac{1}{2}L_{NN} \delta N^2 + \frac{1}{2} \left( L_{\mathcal{R}\mathcal{R}} + H^2 L_{\mathcal{Y}\mathcal{Y}} + 2HL_{\mathcal{Y}\mathcal{R}} \right) \,\delta \mathcal{R}^2 + \left( 2HL_{\mathcal{S}N} + L_{KN} \right) \delta K \delta N$$
(126)  
+  $\left( 2HL_{\mathcal{S}\mathcal{R}} + L_{K\mathcal{R}} + HL_{K\mathcal{Y}} + 2H^2 L_{\mathcal{S}\mathcal{Y}} + \frac{1}{2}L_{\mathcal{Y}} \right) \delta K \delta \mathcal{R} + \left( L_{N\mathcal{R}} + HL_{N\mathcal{Y}} - \frac{1}{2}\dot{L}_{\mathcal{Y}} \right) \,\delta N \delta \mathcal{R} + \mathcal{O}(3) \,.$ 

In summary, an explicit dependence of the action on  $\mathcal{Y}$  can easily be included in our treatment, via the following substitutions in Eq. (12),

$$L_{\mathcal{R}} \rightarrow L_{\mathcal{R}} + \frac{1}{2}\dot{L}_{\mathcal{Y}} + \frac{3}{2}HL_{\mathcal{Y}} ,$$

$$L_{\mathcal{R}\mathcal{R}} \rightarrow L_{\mathcal{R}\mathcal{R}} + H^{2}L_{\mathcal{Y}\mathcal{Y}} + 2HL_{\mathcal{Y}\mathcal{R}} ,$$

$$L_{N\mathcal{R}} \rightarrow L_{N\mathcal{R}} + HL_{N\mathcal{Y}} - \frac{1}{2}\dot{L}_{\mathcal{Y}} ,$$

$$\mathcal{C} \rightarrow \mathcal{C} + HL_{K\mathcal{Y}} + 2H^{2}L_{\mathcal{S}\mathcal{Y}} + \frac{1}{2}L_{\mathcal{Y}} .$$
(127)

# **B** Tensor modes

In this appendix we study the propagation of tensor modes in the action (2). We consider the spatial metric [19]

$$h_{ij} = a^2(t)e^{2\zeta}\hat{h}_{ij} , \qquad \det \hat{h} = 1 , \qquad \hat{h}_{ij} = \delta_{ij} + \gamma_{ij} + \frac{1}{2}\gamma_{ik}\gamma_{kj} ,$$
 (128)

with  $\gamma_{ij}$  traceless and divergence-free,  $\gamma_{ii} = 0 = \partial_i \gamma_{ij}$ . Since tensor modes decouple from scalars, we can simply replace this metric into the action (2) by setting scalar perturbations to zero, which yields

$$S_{\gamma}^{(2)} = \int d^4x \, a^3 \frac{M_*^2 f}{8} \left[ \left( 1 + \frac{2m_4^2}{M_*^2 f} \right) \dot{\gamma}_{ij}^2 - \frac{1}{a^2} (\partial_k \gamma_{ij})^2 \right] \,, \tag{129}$$

where we used that, up to integration by parts,

$${}^{(3)}R = -\frac{1}{4a^2} (\partial_i \gamma_{kj})^2 , \qquad K = 3H , \qquad (130)$$

$$\delta K_{ij}^2 = \frac{1}{4} \dot{\gamma}_{ij}^2 , \qquad K_{ij} K^{ij} - K^2 = -6H^2 + \frac{1}{4} \dot{\gamma}_{ij}^2 , \qquad (131)$$

and the Gauss-Codazzi relation (38). Thus, for  $m_4^2 \neq 0$  the speed of sound of gravity waves is different from the speed of light,

$$c_T^2 = \left(1 + \frac{2m_4^2}{M_*^2 f}\right)^{-1} , \qquad (132)$$

which confirms [34,35] in the case of generalised Galileon theories.

# C EFT parameters for generalized Galileons

Here we explicitly give the EFT of dark energy parameters in terms of the Lagrangian (12), for the generalized Galileon Lagrangians eqs. (51)–(53). All quantities in the expressions below are calculated on the background.

• *L*<sub>3</sub>:

$$f = 0, \tag{133}$$

$$\Lambda = \dot{\phi}^2 (\ddot{\phi} + 3H\dot{\phi})G_{3X} , \qquad (134)$$

$$c = \dot{\phi}^2 (-\ddot{\phi} + 3H\dot{\phi})G_{3X} + \dot{\phi}^2 G_{3\phi}, \qquad (135)$$

$$M_2^4 = \frac{\dot{\phi}^2}{2} (\ddot{\phi} + 3H\dot{\phi}) G_{3X} - 3H\dot{\phi}^5 G_{3,XX} - \frac{\dot{\phi}^4}{2} G_{3,X\phi} , \qquad (136)$$

$$m_3^3 = 2\dot{\phi}^3 G_{3X}, \quad m_4^2 = \tilde{m}_4^2 = 0.$$
 (137)

• *L*<sub>4</sub>:

$$M_*^2 f = 2G_4 av{(138)}$$

$$\Lambda = \frac{1}{2}\dot{\tilde{\mathcal{F}}} + 3H\dot{X}G_{4X} - 18H^2G_{4X}\dot{\phi}^2 + 6HG_{4X\phi}\dot{\phi}^3 + 12H^2G_{4XX}\dot{\phi}^4 , \qquad (139)$$

$$c = -\frac{1}{2}\dot{\tilde{\mathcal{F}}} + 3H\dot{X}G_{4X} - 6H^2G_{4X}\dot{\phi}^2 + 6HG_{4X\phi}\dot{\phi}^3 + 12H^2G_{4XX}\dot{\phi}^4 , \qquad (140)$$

$$M_2^4 = \frac{1}{4}\dot{\tilde{\mathcal{F}}} - \frac{3}{2}H\dot{X}G_{4X} + 6HG_{4X\phi}\dot{\phi}^3 + 18H^2G_{4XX}\dot{\phi}^4 - 6HG_{4XX\phi}\dot{\phi}^5 - 12H^2G_{4XXX}\dot{\phi}^6 , \quad (141)$$

$$m_3^3 = 2\dot{X}G_{4X} - 8HG_{4X}\dot{\phi}^2 + 4G_{4X\phi}\dot{\phi}^3 + 16HG_{4XX}\dot{\phi}^4 , \qquad (142)$$

$$m_4^2 = \tilde{m}_4^2 = 2G_{4X}\phi^2 \,, \tag{143}$$

with

$$\tilde{\mathcal{F}} \equiv 2M_*^2 H f + M_*^2 \dot{f} + \mathcal{F} = 2\dot{X}G_{4X} - 8HG_{4X}\dot{\phi}^2.$$
(144)

,

• *L*<sub>5</sub>:

$$M_*^2 f = -G_{5\phi} \dot{\phi}^2 + 2G_{5X} \dot{\phi}^2 \ddot{\phi} , \qquad (145)$$

$$c = -\frac{1}{2}\dot{\tilde{\mathcal{F}}} + \frac{3}{2}M_*^2H\dot{f} - 3H^2G_{5\phi}\dot{\phi}^2 - 3H^3G_{5X}\dot{\phi}^3 + 3H^2G_{5X\phi}\dot{\phi}^4 + 2H^3G_{5XX}\dot{\phi}^5 , \qquad (146)$$

$$\Lambda = \frac{1}{2}\dot{\tilde{\mathcal{F}}} + 3M_*^2 H^2 f + \frac{3}{2}M_*^2 H\dot{f} - 6H^2 G_{5\phi}\dot{\phi}^2 - 7H^3 G_{5X}\dot{\phi}^3 + 3H^2 G_{5X\phi}\dot{\phi}^4 + 2H^3 G_{5XX}\dot{\phi}^5 ,$$
(147)

$$M_{2}^{4} = \frac{1}{4}\dot{\tilde{\mathcal{F}}} - \frac{3}{4}M_{*}^{2}H\dot{f} - \frac{3}{2}H^{3}G_{5X}\dot{\phi}^{3} + 6H^{2}G_{5X\phi}\dot{\phi}^{4} + 6H^{3}G_{5XX}\dot{\phi}^{5} - 3H^{2}G_{5XX\phi}\dot{\phi}^{6} - 2H^{3}G_{5XXX}\dot{\phi}^{7}$$
(148)

$$m_3^3 = M_*^2 \dot{f} - 4HG_{5\phi} \dot{\phi}^2 - 6H^2 G_{5X} \dot{\phi}^3 + 4HG_{5X\phi} \dot{\phi}^4 + 4H^2 G_{5XX} \dot{\phi}^5 , \qquad (149)$$
  
$$m_4^2 = \tilde{m}_4^2 = G_{5\phi} \dot{\phi}^2 + HG_{5X} \dot{\phi}^3 - G_{5X} \dot{\phi}^2 \ddot{\phi} , \qquad (150)$$

with

$$\tilde{\mathcal{F}} \equiv 2M_*^2 H f + M_*^2 \dot{f} + \mathcal{F} = 2M_*^2 f H + M_*^2 \dot{f} - 2H G_{5\phi} \dot{\phi}^2 - 2H^2 G_{5X} \dot{\phi}^3 .$$
(151)

# D Coefficients of the perturbation equations

In this appendix we define the coefficients appearing in eqs. (97)–(101). For convenience we have used  $M_4^2$  defined as

$$M_4^2 \equiv 2m_4^2 + 3\bar{m}_4^2 \,. \tag{152}$$

Moreover, from the background equations (95) and (96) and using the background conservation equation for matter,  $\dot{\rho}_m + 3H(\rho_m + p_m) = 0$ , one obtains

$$\dot{c} + \dot{\Lambda} = -6Hc + 6H^2 M_*^2 \dot{f} + 3M_*^2 \dot{f} \dot{H} .$$
(153)

We will make use of this relation to simplify some of the terms and eliminate the dependence with respect to  $\dot{\Lambda}$  and  $\ddot{\Lambda}$ .

• Variation with respect to  $\Phi$ :

$$A_{\Phi} = 2c + 4M_2^4 - 6H \left[ fHM_*^2 + M_*^2 \dot{f} - m_3^3 + HM_4^2 \right] , \qquad (154)$$

$$A_{\dot{\Psi}} = -3 \left[ 2H \left( f M_*^2 + M_4^2 \right) + M_*^2 \dot{f} - m_3^3 \right] , \qquad (155)$$

$$A_{\pi} = 3H^2 M_*^2 f + 3m_3^3 H - \dot{c} - \Lambda - 6M_4^2 H H + 3M_*^2 f$$
  
=  $6Hc - 3(H^2 + \dot{H})M_*^2 \dot{f} + 3M_*^2 \ddot{f} + 3m_3^3 \dot{H} - 6M_4^2 H \dot{H}$ , (156)

$$A_{\pi} = -2c - 4M_2^4 - 3H(m_3^3 - M_*^2 \dot{f}) , \qquad (157)$$

$$A_{\Psi}^{(2)} = -2fM_*^2 + 6H\bar{m}_5 - 4\tilde{m}_4^2 , \qquad (158)$$

$$A_{\pi}^{(2)} = M_*^2 \dot{f} - m_3^3 + 2HM_4^2 - 4H\tilde{m}_4^2 + 6H^2 \bar{m}_5 .$$
(159)

• Variation with respect to  $\alpha$ :

$$B_{\Phi} = -m_3^3 + 2H\left(fM_*^2 + M_4^2\right) + M_*^2\dot{f} , \qquad (160)$$

$$B_{\dot{\Psi}} = 2\left(fM_*^2 + M_4^2\right) , \qquad (161)$$

$$B_{\pi} = -2c + 2M_4^2 \dot{H} + M_*^2 (H\dot{f} - \ddot{f}) , \qquad (162)$$

$$B_{\pi} = m_3^3 - M_*^2 \dot{f} , \qquad (163)$$

$$B_{\Psi}^{(2)} = -2\bar{m}_5 , \qquad (164)$$

$$B_{\pi}^{(2)} = -2(\bar{m}_4^2 + H\bar{m}_5) . \tag{165}$$

# • Variation with respect to $\Psi$ :

$$C_{\Phi} = 3 \left[ 2c + 2(3H^2 + \dot{H})M_4^2 + 2fM_*^2(3H^2 + 2\dot{H}) - (m_3^3)^{\cdot} + H \left( -3m_3^3 + 4M_*^2\dot{f} + 2(M_4^2)^{\cdot} \right) + 2M_*^2\ddot{f} \right] , \qquad (166)$$

$$C_{\dot{\Phi}} = -3m_3^3 + 6H\left(fM_*^2 + M_4^2\right) + 3M_*^2\dot{f} , \qquad (167)$$

$$C_{\dot{\Psi}} = 6 \left( 3fHM_*^2 + 3HM_4^2 + M_*^2\dot{f} + (M_4^2)^{\cdot} \right) , \qquad (168)$$
  

$$C_{\ddot{\mu}} = 6 \left( fM_*^2 + M_4^2 \right) , \qquad (169)$$

$$C_{\Psi} = 6 \left( j \, M_* + M_4 \right) , \qquad (105)$$

$$C_{\pi} = -3 \left[ \dot{c} - \dot{\Lambda} - 2\dot{H} (M_4^2) - 2(\ddot{H} + 3H\dot{H}) M_4^2 + M_*^2 (2\dot{f}\dot{H} + f^{(3)} + 3H^2\dot{f} + 2H\ddot{f}) \right]$$

$$= -3 \left[ 2\dot{c} + 6Hc - 2\dot{H} (M_4^2) - 2(\ddot{H} + 3H\dot{H}) M_4^2 + M_*^2 \left( f^{(3)} - (3H^2 + \dot{H})\dot{f} + 2H\ddot{f} \right) \right] , \qquad (170)$$

$$C_{\dot{\pi}} = 3\left(-2c + 3Hm_3^3 - 2HM_*^2\dot{f} + 2M_4^2\dot{H} + (m_3^3)\dot{-} - 2M_*^2\ddot{f}\right) , \qquad (171)$$

$$C_{\ddot{\pi}} = 3(m_3^3 - M_*^2 \dot{f}) , \qquad (172)$$

$$C_{\Phi}^{(2)} = -\left(2fM_*^2 - 6H\bar{m}_5 + 4\tilde{m}_4^2\right) , \qquad (173)$$

$$C_{\Psi}^{(2)} = 2fM_*^2 - 6H\bar{m}_5 - 6\dot{\bar{m}}_5 , \qquad (174)$$

$$C_{\pi}^{(2)} = -2\left(M_{*}^{2}\dot{f} + (M_{4}^{2})^{\cdot} + HM_{4}^{2} + 3H^{2}\bar{m}_{5} + 3H\dot{\bar{m}}_{5}\right) , \qquad (175)$$

$$C_{\pi}^{(2)} = -2(M_4^2 - 2\tilde{m}_4^2 + 3H\bar{m}_5), \qquad (176)$$

$$C_{\Psi}^{(4)} = 16\bar{\lambda} ,$$
 (177)

$$C_{\pi}^{(4)} = -2\bar{m}_5 + 16H\bar{\lambda} . \tag{178}$$

• Variation with respect to  $\beta$ :

$$D_{\Phi}^{(2)} = M_*^2 f + 2\tilde{m}_4^2 - 3\bar{m}_5 H , \qquad (179)$$

$$D_{\Psi}^{(2)} = -M_*^2 f , \qquad (180)$$

$$D_{\dot{\Psi}}^{(2)} = -3\bar{m}_5 , \qquad (181)$$

$$D_{\pi}^{(2)} = M_*^2 \dot{f} + 2m_4^2 H + 2(m_4^2) \dot{-} 3\dot{H}\bar{m}_5 , \qquad (182)$$

$$D_{\pi}^{(2)} = 2m_4^2 - 2\tilde{m}_4^2 , \qquad (183)$$

$$D_{\Psi}^{(4)} = -8\bar{\lambda} , \qquad (184)$$

$$D_{\pi}^{(4)} = \bar{m}_5 - 8H\bar{\lambda} . \tag{185}$$

#### • Variation with respect to $\pi$ :

$$E_{\Phi} = 6cH + \dot{c} + H^2 (9m_3^3 - 6M_*^2 \dot{f}) + 3(2m_3^3 - M_*^2 \dot{f})\dot{H} - \dot{\Lambda} + 3H \left[4M_2^4 - 2M_4^2 \dot{H} + (m_3^3)^{\cdot}\right] + 4(M_2^4)^{\cdot}$$
  
$$= 12cH + 2\dot{c} + 3m_3^3 (3H^2 + 2\dot{H}) - 6M_*^2 \dot{f}(2\dot{H} + H^2) + 3H \left[4M_2^4 - 2M_4^2 \dot{H} + (m_3^3)^{\cdot}\right] + 4(M_2^4)^{\cdot},$$
  
(186)

$$E_{\dot{\Phi}} = 2c + 4M_2^4 + 3H(m_3^3 - M_*^2\dot{f}) , \qquad (187)$$

$$E_{\Psi} = 3 \left[ 6cH + \dot{c} + \dot{\Lambda} - 3M_*^2 \dot{f} (2H^2 + \dot{H}) \right] = 0 , \qquad (188)$$

$$E_{\dot{\Psi}} = 3 \left[ 2c + 3Hm_3^3 - 4HM_*^2 \dot{f} - 2M_4^2 \dot{H} + (m_3^3)^{\cdot} \right] , \qquad (189)$$

$$E_{\bar{\Psi}} = 3(m_3^3 - M_*^2 f) , \qquad (190)$$

$$E_{\pi} = -\left[ 6M_4^2 \dot{H}^2 - 3(m_3^3) \dot{H} + 6H\dot{c} - 9H\dot{H}m_3^3 + \ddot{c} - 3M_*^2 \dot{H}\dot{f} - 6H^2 M_*^2 \dot{f} - 3m_3^3 \dot{H} + \dot{\Lambda} \right]$$
  
$$= -\left[ 6M_4^2 \dot{H}^2 - 3(m_3^3) \dot{H} + 6\dot{H}c + 3M_*^2 (\ddot{H} + 4H\dot{H})\dot{f} - 9H\dot{H}m_3^3 - 3m_3^3 \ddot{H} \right] , \qquad (191)$$

$$E_{\pi} = -2 \left[ 3H \left( c + 2M_2^4 \right) + \dot{c} + 2(M_2^4)^{\cdot} \right] , \qquad (192)$$

$$E_{\ddot{\pi}} = -2\left(c + 2M_2^4\right) \,, \tag{193}$$

$$E_{\Phi}^{(2)} = -\left[m_3^3 + H\left(-2M_4^2 - 6H\bar{m}_5 + 4\tilde{m}_4^2\right) - M_*^2\dot{f}\right] , \qquad (194)$$

$$E_{\Psi}^{(2)} = -2 \left[ 2H\tilde{m}_4^2 + M_*^2 \dot{f} - 3\bar{m}_5 \dot{H} + 2(\tilde{m}_4^2)^{\cdot} \right] , \qquad (195)$$

$$E_{\dot{\Psi}}^{(2)} = 2M_4^2 + 6H\bar{m}_5 - 4\tilde{m}_4^2 , \qquad (196)$$

$$E_{\pi}^{(2)} = -\left[2c - 4M_4^2\dot{H} + 4\tilde{m}_4^2\dot{H} + (m_3^3) + 4H^2\tilde{m}_4^2 + Hm_3^3 - 12\bar{m}_5H\dot{H} + 4H(\tilde{m}_4^2)\right], \qquad (197)$$

$$E_{\Psi}^{(4)} = -2\bar{m}_5 + 16H\bar{\lambda} , \qquad (198)$$

$$E_{\pi}^{(4)} = -2(\bar{m}_4^2 + 2H\bar{m}_5) + 16H^2\bar{\lambda}.$$
<sup>(199)</sup>

## • Generalized Poisson equation:

$$F_{\Phi} = -3M_*^2 H \dot{f} + 2c + 4M_2^4 + 3Hm_3^3 , \qquad (200)$$

$$F_{\dot{\Psi}} = -3M_*^2 \dot{f} + 3m_3^3 , \qquad (201)$$

$$F_{\pi} = 6M_*^2 H^2 \dot{f} - 6Hc - \dot{c} - \dot{\Lambda} + 3m_3^3 \dot{H} = -3\dot{H}(M_*^2 \dot{f} - m_3^3), \qquad (202)$$

$$F_{\pi} = -(2c + 4M_2^*) , \qquad (203)$$

$$F_{\Psi}^{(2)} = -(2fM_*^2 + 4\tilde{m}_4^2) , \qquad (204)$$

$$F_{\pi}^{(2)} = \dot{f}M_*^2 - m_3^3 + 4Hm_4^2 - 4H\tilde{m}_4^2 \,. \tag{205}$$

# References

- T. Clifton, P. G. Ferreira, A. Padilla and C. Skordis, "Modified Gravity and Cosmology," Phys. Rept. 513, 1 (2012) [arXiv:1106.2476 [astro-ph.CO]].
- [2] L. Amendola and S. Tsujikawa, "Dark Energy: Theory and Observations," Cambridge U. P. (2011) 506 p
- [3] P. Creminelli, M. A. Luty, A. Nicolis and L. Senatore, "Starting the Universe: Stable Violation of the Null Energy Condition and Non-standard Cosmologies," JHEP 0612, 080 (2006) [hep-th/0606090].
- [4] C. Cheung, P. Creminelli, A. L. Fitzpatrick, J. Kaplan and L. Senatore, "The Effective Field Theory of Inflation," JHEP 0803, 014 (2008) [arXiv:0709.0293 [hep-th]].
- [5] L. Senatore, K. M. Smith and M. Zaldarriaga, "Non-Gaussianities in Single Field Inflation and their Optimal Limits from the WMAP 5-year Data," JCAP 1001, 028 (2010) [arXiv:0905.3746 [astro-ph.CO]].

- [6] P. Creminelli, G. D'Amico, M. Musso, J. Norena and E. Trincherini, "Galilean symmetry in the effective theory of inflation: new shapes of non-Gaussianity," JCAP 1102, 006 (2011) [arXiv:1011.3004 [hep-th]].
- [7] C. L. Bennett, D. Larson, J. L. Weiland, N. Jarosik, G. Hinshaw, N. Odegard, K. M. Smith and R. S. Hill et al., "Nine-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Final Maps and Results," arXiv:1212.5225 [astro-ph.CO].
- [8] P. A. R. Ade *et al.* [Planck Collaboration], "Planck 2013 Results. XXIV. Constraints on primordial non-Gaussianity," arXiv:1303.5084 [astro-ph.CO].
- [9] P. Creminelli, G. D'Amico, J. Norena and F. Vernizzi, "The Effective Theory of Quintessence: the wi-1 Side Unveiled," JCAP 0902, 018 (2009) [arXiv:0811.0827 [astro-ph]].
- [10] P. Creminelli, G. D'Amico, J. Norena, L. Senatore and F. Vernizzi, "Spherical collapse in quintessence models with zero speed of sound," JCAP 1003, 027 (2010) [arXiv:0911.2701 [astro-ph.CO]].
- [11] G. Gubitosi, F. Piazza and F. Vernizzi, "The Effective Field Theory of Dark Energy," JCAP 1302, 032 (2013) [arXiv:1210.0201 [hep-th]].
- [12] J. K. Bloomfield, E. E. Flanagan, M. Park and S. Watson, "Dark Energy or Modified Gravity? An Effective Field Theory Approach," arXiv:1211.7054 [astro-ph.CO].
- [13] R. A. Battye and J. A. Pearson, "Effective action approach to cosmological perturbations in dark energy and modified gravity," JCAP 1207, 019 (2012) [arXiv:1203.0398 [hep-th]].
- [14] S. Weinberg, "Effective Field Theory for Inflation," Phys. Rev. D 77, 123541 (2008) [0804.4291 [hep-th]].
- [15] M. Park, K. M. Zurek and S. Watson, "A Unified Approach to Cosmic Acceleration," Phys. Rev. D 81, 124008 (2010) [arXiv:1003.1722 [hep-th]].
- [16] J. K. Bloomfield and E. E. Flanagan, "A Class of Effective Field Theory Models of Cosmic Acceleration," arXiv:1112.0303 [gr-qc].
- [17] T. Baker, P. G. Ferreira and C. Skordis, "The Parameterized Post-Friedmann Framework for Theories of Modified Gravity: Concepts, Formalism and Examples," Phys. Rev. D 87, 024015 (2013) [arXiv:1209.2117 [astro-ph.CO]].
- [18] J. Z. Simon, "Higher Derivative Lagrangians, Nonlocality, Problems And Solutions," Phys. Rev. D 41, 3720 (1990).
- [19] J. M. Maldacena, "Non-Gaussian features of primordial fluctuations in single field inflationary models," JHEP 0305, 013 (2003) [astro-ph/0210603].
- [20] L. Boubekeur, P. Creminelli, J. Norena, F. Vernizzi and , "Action approach to cosmological perturbations: the 2nd order metric in matter dominance," JCAP 0808, 028 (2008) [arXiv:0806.1016 [astro-ph]].
- [21] G. W. Horndeski, Int. J. Theor. Phys. 10, 363 (1974).
- [22] C. Charmousis, E. J. Copeland, A. Padilla and P. M. Saffin, "General second order scalar-tensor theory, self tuning, and the Fab Four," Phys. Rev. Lett. 108, 051101 (2012) [arXiv:1106.2000 [hep-th]].
- [23] A. Nicolis, R. Rattazzi and E. Trincherini, "The Galileon as a local modification of gravity," Phys. Rev. D 79, 064036 (2009) [arXiv:0811.2197 [hep-th]].
- [24] C. Deffayet, G. Esposito-Farese and A. Vikman, "Covariant Galileon," Phys. Rev. D 79, 084003 (2009) [arXiv:0901.1314 [hep-th]].
- [25] C. Deffayet, S. Deser and G. Esposito-Farese, "Generalized Galileons: All scalar models whose curved background extensions maintain second-order field equations and stress-tensors," Phys. Rev. D 80, 064015 (2009) [arXiv:0906.1967 [gr-qc]].
- [26] C. Deffayet, X. Gao, D. A. Steer and G. Zahariade, "From k-essence to generalised Galileons," Phys. Rev. D 84, 064039 (2011) [arXiv:1103.3260 [hep-th]].
- [27] N. Arkani-Hamed, H. -C. Cheng, M. A. Luty, S. Mukohyama, "Ghost condensation and a consistent infrared modification of gravity," JHEP 0405, 074 (2004). [hep-th/0312099].
- [28] E. Poisson, A Relativists Toolkit: The Mathematics of Black-Hole Mechanics, Cambridge University Press, Cambridge, 1973.
- [29] R. M. Wald, "General Relativity," Chicago, Usa: Univ. Pr. (1984) 491p
- [30] S. A. Appleby, A. De Felice and E. V. Linder, "Fab 5: Noncanonical Kinetic Gravity, Self Tuning, and Cosmic Acceleration," JCAP 1210, 060 (2012) [arXiv:1208.4163 [astro-ph.CO]].
- [31] J. K. Bloomfield, E. E. Flanagan, M. Park and S. Watson, private communication.
- [32] A. De Felice, T. Kobayashi, S. Tsujikawa and , "Effective gravitational couplings for cosmological perturbations in the most general scalar-tensor theories with second-order field equations," Phys. Lett. B 706, 123 (2011) [arXiv:1108.4242 [gr-qc]].

- [33] A. Silvestri, L. Pogosian and R. V. Buniy, "A practical approach to cosmological perturbations in modified gravity," arXiv:1302.1193 [astro-ph.CO].
- [34] X. Gao and D. A. Steer, "Inflation and primordial non-Gaussianities of 'generalized Galileons'," JCAP 1112, 019 (2011) [arXiv:1107.2642 [astro-ph.CO]].
- [35] A. De Felice and S. Tsujikawa, "Inflationary non-Gaussianities in the most general second-order scalar-tensor theories," Phys. Rev. D 84, 083504 (2011) [arXiv:1107.3917 [gr-qc]].