

ON THE PUSH-OUT SPACES

M. FATHY AND M. FAGHFOURI

ABSTRACT. Let $f : M^m \rightarrow \mathbb{R}^{m+k}$ be an immersion where M is a smooth connected m -dimensional manifold without boundary. Then we construct a subspace $\Omega(f)$ of \mathbb{R}^k , namely push-out space. which corresponds to a set of embedded manifolds which are either parallel to f , tubes around f or, ingeneral, partial tubes around f . This space is invariant under the action of the normal holonomy group, $\mathcal{H}ol(f)$. Moreover, we construct geometrically some examples for normal holonomy group and push-out space in \mathbb{R}^3 . These examples will show that properties of push-out space that are proved in the case $\mathcal{H}ol(f)$ is trivial, is not true in general.

1. INTRODUCTION

In this paper we introduce push-out space for an immersion $f : M^m \rightarrow \mathbb{R}^{m+k}$, where M is a smooth connected m -dimensional manifold without boundary. To do this, we give some examples in 3-dimensional Euclidean space, \mathbb{R}^3 , infact, in these examples we calculate normal holonomy group and push-out space geometrically. We consider the case when $\mathcal{H}ol(f)$ is non-trivial. This extends the work of Carter and Senturk [2], who obtained results about the case when $\mathcal{H}ol(f)$ is trivial. In these examples we show that some of the properties of push-out space which they obtained is not true for the case when $\mathcal{H}ol(f)$ is non-trivial.

2. BASIC DEFINITIONS

Definition 2.1 ([2]). Let $f : M^m \rightarrow \mathbb{R}^{m+k}$ be a smooth immersion where M is a smooth connected m -dimensional manifold without boundary. The total space of the normal bundle of f is defined by

$$N(f) = \{(p, x) \in M \times \mathbb{R}^{m+k} : \langle x, v \rangle = 0 \quad \forall v \in f_*T_p(M)\}$$

The endpoint map $\eta : N(f) \rightarrow \mathbb{R}^{m+k}$ is defined by $\eta(p, x) = f(p) + x$ and, the set of singular points of η is subset $\Sigma(f) \subset N(f)$ called the set of critical normals of f and the set of focal points of η is a subset $\eta(\Sigma(f)) \subset \mathbb{R}^{m+k}$.

For $p \in M$, we put $N_p(f) = \{x : (p, x) \in N(f)\}$ and $\Sigma_p(f) = \{x : (p, x) \in \Sigma(f)\}$ respectively, normal space at p and the set can be thought of as focal points with base p .

Definition 2.2 ([1]). For $p_0 \in M$ and $p \in M$ and path $\gamma : [0, 1] \rightarrow M$ from p_0 to p define $\varphi_{p,\gamma} : N_{p_0}(f) \rightarrow N_p(f)$ by parallel transport along γ . The $\varphi_{p,\gamma}$'s are isometries. The normal holonomy group on $N_{p_0}(f)$, is

$$\mathcal{H}ol(f) = \{\varphi_{p_0,\gamma} : \gamma : [0, 1] \rightarrow M, \quad \gamma(0) = \gamma(1) = p_0\}$$

1991 *Mathematics Subject Classification.* 53C40, 53C42.

Key words and phrases. Singular points, Normal holonomy group, push-out space.

If the closed path γ at p_0 is homotopically trivial then $\varphi_{p_0, \gamma}$ is an element of the restricted normal holonomy group $\mathcal{H}ol_0(f)$.

Definition 2.3 ([3]). For a fix $p_0 \in M$ the push-out space for an immersion $f : M^m \rightarrow \mathbb{R}^{m+k}$ is defined by

$$\Omega(f) = \{x \in N_{p_0}(f) : \forall p \in M, \forall \gamma \text{ s.t. } \gamma(0) = p_0, \gamma(1) = p \text{ then } \varphi_{p, \gamma} x \notin \Sigma_p(f)\}$$

(i.e. $\forall p \in M, f(p) + \varphi_{p, \gamma}(x)$ is not a focal point with base p when x belongs to $\Omega(f)$). Therefore $\Omega(f)$ is the set of normals at p_0 , where transported parallelly along all curves, do not meet focal points. So $\Omega(f)$ is invariant under the action of $\mathcal{H}ol(f)$.

Definition 2.4 ([4]). Let $B \subset N(f)$ be a smooth subbundle with type fiber S where

- 1) S is a smooth submanifold of \mathbb{R}^k
- 2) $B \cap \Sigma(f) = \emptyset$
- 3) B is invariant under parallel transport (along any curve in M). Then B is a smooth manifold and $g \equiv \eta|_B : B \rightarrow \mathbb{R}^{m+k}$ is a smooth immersion called a partial tube about f .

Theorem 2.5 ([2]). Let $\mathcal{H}ol(f)$ is trivial and M be a compact manifold. Then each path-connected component of $\Omega(f)$ is open in \mathbb{R}^k .

Theorem 2.6 ([2]). Let $\mathcal{H}ol(f)$ is trivial then Each path-connected component of $\Omega(f)$ is convex.


Remark 2.7. In Example 3.2, if $\frac{\alpha}{\pi}$ is irrational then $\Omega(\bar{f})$ is not open in \mathbb{R}^2 but $M = \mathbb{S}^1$ is compact. Also, in Example 3.5, $\Omega(f) = \{O\}$ hence $\Omega(f)$ is closed in \mathbb{R}^2 but $\mathcal{H}ol(f)$ is trivial. This shows that Theorem 2.5 is false when M is not compact or $\mathcal{H}ol(f)$ is non-trivial.

Remark 2.8. In Example 3.6 one of path-connected components of $\Omega(f)$, which is the complement space of cone and two other components in \mathbb{R}^3 , is not convex. This shows that Theorem 2.6 is false when $\mathcal{H}ol(f)$ is non-trivial.

we conclude that the properties of push-out space that are proved in the case $\mathcal{H}ol(f)$ is trivial, is not true in general.

3. EXAMPLES OF NORMAL HOLONOMY GROUPS AND PUSH-OUT SPACES


Example 3.1. We start with a curve as below



suppose this curve is given by $s \mapsto (\xi(s), \eta(s))$ where $s \in [0, 1]$ and at $(1, 0, 0): s = 0$, $\frac{\partial \xi}{\partial s} = 1, (\frac{\partial}{\partial s})^r \eta = 0$ for all $r \geq 0$ and at $(0, 1, 0): s = 1, \frac{\partial \eta}{\partial s} = 1, (\frac{\partial}{\partial s})^r \xi = 0$ for all $r \geq 0$. Now, we take this curve in \mathbb{R}^3 and consider the same curves in yz -plane and


xz -plane and fit together to make a smooth closed curve in \mathbb{R}^3 . Now by identifying \mathbb{S}^1 with $\frac{\mathbb{R}}{3\mathbb{Z}}$, the curve in \mathbb{R}^3 can be redefined as $f : \mathbb{S}^1 \rightarrow \mathbb{R}^3$ where:

$$f(s) = \begin{cases} (\xi(s), \eta(s), 0) & 0 \leq s \leq 1 \\ (0, \xi(s-1), \eta(s-1)) & 1 \leq s \leq 2 \\ (\eta(s-2), 0, \xi(s-2)) & 2 \leq s \leq 3 \end{cases}$$




To find the normal holonomy group of the above curve, we will consider normal vector to the curve under parallel transport. As each part of the curve lies in a 2-plane, the normal plane at a point of the curve is spanned by the perpendicular direction to the 2-planes.

Step1. Start with the normal vector at $(1,0,0)$, in the diagram, it stays in the xy -plane under parallel transport.




The normal vector $(0,1,0)$ at $(1,0,0)$ goes to normal vector $(-1,0,0)$ at $(0,1,0)$.

Step2. At $(0,1,0)$ the normal vector $(-1,0,0)$ is perpendicular to the yz -plane, it stays perpendicular to the yz -plane under parallel transport from $(0,1,0)$ to $(0,0,1)$.




The normal vector $(-1,0,0)$ at $(0,1,0)$ goes to normal vector $(-1,0,0)$ at $(0,0,1)$.


Step3. The the normal vector $(-1,0,0)$ is in the xz -plane at $(0,0,1)$ and stays in the xz -plane from $(0,0,1)$ to $(1,0,0)$.




The normal vector $(-1,0,0)$ at $(1,0,0)$ by going once around the curve the normal vector will turn about $\frac{\pi}{2}$.



Going around of curve again, the normal vector moves through another $\frac{\pi}{2}$ and after four times around the curve back to its original position. this shows that $\mathcal{Hol}(f)$ is generated by a rotation through $\frac{\pi}{2}$.




Now we find the push-out space of f . Except at end-points of three areas, locally the curve lies in a 2-plane so the focal points with base $s, f(s) + \Sigma_s(f)$, consists of a straight line through the center of curvature, $c(s)$, of the curve at s , perpendicular to the line joining $c(s)$ and $f(s)$.




At end-points of three areas, and possibly some other points, the focal set is empty as the center of curvature "at infinity".

so $\Sigma_s(f)$ is a line in $N_s(f)$. The image of $\Sigma_s(f)$ under normal holonomy group is obtained by rotating it through $\frac{\pi}{2}$ until it returns to the original position.



Now, fix the normal plane $N_{s_0}(f)$ at $f(S_0) = (1, 0, 0)$ where $s_0 = 0$ and use parallel transport to identify all the normal planes with the normal plane $N_{s_0}(f)$. The push-out space is complement of all the $\Sigma_s(f)$ and their images under normal holonomy group.



Therefore the push-out space of $f : \mathbb{S}^1 \rightarrow \mathbb{R}^3$ is an open square Q with sides of length 2ρ where ρ is the minimum absolute value of the radius of curvature of the original curve in the xy -plane. (i.e. $\Omega(f)$ is the interior of the smallest square on $N_{s_0}(f)$.)


Example 3.2. We consider the immersion \bar{f} as in Example 3.1 except that the xz -plane is tilted through an angle α .

In other words, $\bar{f} = Lof$ where f is the immersion in example 3.1 and L is the linear transformation given by


$$L = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & \tan \alpha \\ 0 & 0 & 1 \end{pmatrix} \text{ where } 0 < \alpha < \frac{\pi}{2}. \text{ So, we have}$$

$$\bar{f}(s) = \begin{cases} (\xi(s), \eta(s), 0) & 0 \leq s \leq 1 \\ (0, \xi(s-1) + \eta(s-1) \tan \alpha, \eta(s-1)) & 1 \leq s \leq 2 \\ (\eta(s-2), \xi(s-2) \tan \alpha, \xi(s-2)) & 2 \leq s \leq 3 \end{cases}$$

The end-points of three areas of this immersion are $(1,0,0)$, $(0,1,0)$ and $(0, \tan \alpha, 1)$. Note that at these points the tangent to the curve is the radial line form $(0,0,0)$ so unit tangent at $(1,0,0)$ is $(1,0,0)$ and unit tangent at $(0, \tan \alpha, 1)$ is $\frac{(0, \tan \alpha, 1)}{\sqrt{1 + \tan^2 \alpha}}$ etc. As in Example 3.1, under parallel transport, the normal vector $(0,1,0)$ at $(1,0,0)$ goes to the normal vector $(-1,0,0)$ at $(0,1,0)$, which goes to the normal vector $(-1,0,0)$ at $(0, \tan \alpha, 1)$, which goes to the normal vector $\frac{(0, \tan \alpha, 1)}{\sqrt{1 + \tan^2 \alpha}}$ at $(1,0,0)$. So going once around the curve the normal vector has moved through $\frac{\pi}{2} - \alpha$.



This shows that $\mathcal{Hol}(\bar{f})$ is generated by a rotation through $\frac{\pi}{2} - \alpha$. As in Example 3.1, the image of $\Sigma_s(\bar{f})$ under normal holonomy group is obtained by rotating the line $\Sigma_s(\bar{f})$ through $\frac{\pi}{2} - \alpha$. It depends on α and is obtained as:



If R is the rotating through an angle $\frac{\pi}{2} - \alpha$ then $\Omega(\bar{f}) = \bigcap \{(R)^n Q : n \in \mathbb{Z}\}$ where Q is a square as in Example 3.1, thus if $\frac{\alpha}{\pi}$ is rational then $(R)^n(\Sigma_s(\bar{f})) = \Sigma_s(\bar{f})$ for some $n \in \mathbb{Z}$ and so, $\Omega(\bar{f})$ is the interior of the smallest polygon. If $\frac{\alpha}{\pi}$ is irrational then $(R)^n(\Sigma_s(\bar{f})) \neq \Sigma_s(\bar{f})$ for any $n \in \mathbb{Z}$ and so, $\Omega(\bar{f})$ is an open disk of radius ρ together with a dense set of points on the boundary circle where ρ is minimum absolute value of the radius of curvature of the immersed curve by \bar{f} .

Example 3.3. In Example 3.2, we replace the immersion \bar{f} with the immersion $\bar{f}oh$ where $h : \mathbb{R} \rightarrow \mathbb{S}^1 \cong \frac{\mathbb{R}}{2\pi\mathbb{Z}}$ is covering projection. Since \mathbb{R} is simply connected, for any arbitrary point $s \in \mathbb{R}$, any closed path at s is nullhomotopic with constant path at s , hence definition 2.2 shows that, the normal holonomy group of $\bar{f}oh$ is trivial (i.e. $\mathcal{H}ol(\bar{f}oh) = \mathcal{H}ol_0(\bar{f}oh)$). To calculate $\Omega(\bar{f}oh)$, we prove the theorem 3.4, in general. It will show that $\Omega(\bar{f}oh) = \Omega(\bar{f})$.

Theorem 3.4. *Let $f : M^m \rightarrow \mathbb{R}^{m+k}$ be an immersion and \hat{M} be any covering space with covering projection $h : \hat{M} \rightarrow M$. If $\hat{f} = foh$, then $\Omega(\hat{f}) = \Omega(f)$.*

Proof. Let $x \in \Omega(f)$ and fix $p_0 \in M$. Then definition 2.3 implies that, $\forall p \in M, \forall \gamma$ s.t. $\gamma(0) = p_0, \gamma(1) = p; \varphi_{p,\gamma} x \neq \Sigma_p(f)$.

we define the total space of the normal bundle of \hat{f} by

$$N(\hat{f}) = \{(\hat{p}, x) \in \hat{M} \times \mathbb{R}^{m+k} : \langle x, v \rangle = 0 \quad \forall v \in \hat{f}_* T_{\hat{p}}(\hat{M})\}$$

Also, for any $\hat{p} \in h^{-1}(p)$ we have

$$\begin{aligned} \hat{f}_* T_{\hat{p}}(\hat{M}) &= (foh)_* T_{\hat{p}}(\hat{M}) \\ &= (f_* oh_*) T_{\hat{p}}(\hat{M}) \\ &= f_* T_p(M) \end{aligned}$$

this shows that, for any $\hat{p} \in h^{-1}(p)$ we have $N_{\hat{p}}(\hat{f}) = N_p(f)$ and so $\Sigma_{\hat{p}}(\hat{f}) = \Sigma_p(f)$. Further, we fix $\hat{p}_0 \in h^{-1}(p_0)$ then $\hat{\varphi}_{\hat{p},\hat{\gamma}} = \varphi_{p,\gamma}$ where $\hat{\gamma} : [0, 1] \rightarrow \hat{M}$ s.t. $\hat{\gamma}(0) = \hat{p}_0, \hat{\gamma}(1) = \hat{p}$. Therefore, $\forall \hat{p} \in \hat{M}, \forall \hat{\gamma}; \hat{\varphi}_{\hat{p},\hat{\gamma}} x \neq \Sigma_{\hat{p}}(\hat{f})$. Now using definition 2.3 again, follows that, $x \in \Omega(\hat{f})$. By the same way proves that $\Omega(\hat{f}) \subseteq \Omega(f)$. \square


Example 3.5. If $\frac{\pi}{2} - \alpha = \frac{2\pi}{n}$, then Example 3.3 can be modified by replacing h by the n -fold covering $\bar{h} : \mathbb{S}^1 \rightarrow \mathbb{S}^1$.

Going once around the first \mathbb{S}^1 in $\bar{h} : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ corresponds to moving n times around the second \mathbb{S}^1 so parallelly transporting a normal n times around the second \mathbb{S}^1 which gives a rotation of

$$\begin{aligned} n\left(\frac{\pi}{2} - \alpha\right) &= n\left(\frac{2\pi}{n}\right) \\ &= 2\pi \end{aligned}$$

i.e. the identity, so $\mathcal{H}ol(\bar{f}o\bar{h}) = \mathcal{H}ol_0(\bar{f}o\bar{h})$. Since, the immersed curve by $\bar{f}o\bar{h}$ and the immersed curve by \bar{f} have same figure in \mathbb{R}^3 and $\mathcal{H}ol(\bar{f}o\bar{h})$ is trivial so the singular sets of them also the same (i.e. $\Sigma(\bar{f}o\bar{h}) = \Sigma(\bar{f})$). This implies that $\Omega(\bar{f}o\bar{h}) = \Omega(\bar{f})$.

Example 3.6. We consider a sequence of curves f_n in \mathbb{R}^3 defined as in Example 3.1 except that $\|f_n(s)\|$ and the curvature tends to infinity with n when $s = \frac{1}{2}, \frac{3}{2}$ or $\frac{5}{2}$ but is bounded otherwise.



Now, we define the immersion $f : \mathbb{R} \rightarrow \mathbb{R}^3$ by $f(s \pm 3n) = f_n(s)$. When n tends to infinity, the immersion $f : \mathbb{R} \rightarrow \mathbb{R}^3$ has a sequence of points where the curvature tends to infinity and the radius of curvature at these points can be arbitrary small; in other words, $\exists s$ where $\Sigma_s(f)$ is arbitrary close to "O" in $N_s(f)$. So $\{O\}$ is the only point not in the image of $\Sigma_s(f)$ under normal holonomy group for all $s \in \mathbb{R}$. Then $\Omega(f) = \{O\}$. In this case because \mathbb{R} is simply connected then $\mathcal{H}ol(f)$ is trivial.

The following results have been proved in [2], when $\mathcal{H}ol(f)$ is trivial.

Theorem 3.7. *Let M be a compact manifold, then each path-connected component of $\Omega(f)$ is open in \mathbb{R}^k .*

Theorem 3.8. *Each path-connected component of $\Omega(f)$ is convex.*

Remark 3.9. In Example 3.2, if $\frac{\alpha}{\pi}$ is irrational then $\Omega(\bar{f})$ is not open in \mathbb{R}^2 but $M = \mathbb{S}^1$ is compact. Also, in Example 3.5 $\Omega(f) = \{O\}$ so $\Omega(f)$ is closed in \mathbb{R}^2 but $\mathcal{H}ol(f)$ is trivial. This shows that Theorem 3.7 is false when M is not compact or $\mathcal{H}ol(f)$ is non-trivial.

Remark 3.10. In Example 3.6 one of the path-connected components of $\Omega(f)$, which is the complement space of cone and two other components in \mathbb{R}^3 , is not convex. This shows that Theorem 3.8 is false when $\mathcal{Hol}(f)$ is non-trivial.

Thus the properties of push-out space that are proved in the case $\mathcal{Hol}(f)$ is trivial, is not true in general.

REFERENCES

- [1] J. Berndt, S. Console and C. Olmos, Submanifolds and Holonomy, Research Notes in Mathematics 434, CHAPMAN & HALL/CRC, 2003.
- [2] S. Carter and Z. Senturk, The space of immersions parallel to a given immersion, J. London Math. Soc. (2) **50** (1994), 404-416.
- [3] S. Carter, Z. Senturk and A. West, The push-out space of a submanifold, Geometry and Topology of submanifolds VI, (1994), 50-57.
- [4] S. Carter and A. West, partial tubes about immersed manifolds, Geom. Dedicata 54 (1995), 145-169.

UNIVERSITY OF APPLIED SCIENCE, AND TECHNOLOGY OF EAST AZARBAYJAN COOPERATION,,
TABRIZ, IRAN

E-mail address: mortaza.fathy@yahoo.com

UNIVERSITY OF TABRIZ,, TABRIZ, IRAN

E-mail address: faghfouritabrizu.ac.ir