

KÄHLER CURRENTS AND NULL LOCI

TRISTAN C. COLLINS AND VALENTINO TOSATTI*

Dedicated to D.H. Phong with admiration on the occasion of his 60th birthday.

ABSTRACT. We prove that the non-Kähler locus of a nef and big class on a compact complex manifold bimeromorphic to a Kähler manifold equals its null locus. In particular this gives an analytic proof of a theorem of Nakamaye and Ein-Lazarsfeld-Mustață-Nakamaye-Popa. As an application, we show that finite time non-collapsing singularities of the Kähler-Ricci flow on compact Kähler manifolds always form along analytic subvarieties, thus answering a question of Feldman-Ilmanen-Knopf and Campana. We also extend the second author's results about noncollapsing degenerations of Ricci-flat Kähler metrics on Calabi-Yau manifolds to the nonalgebraic case.

1. INTRODUCTION

The general topic of this paper is the study of the boundary of the Kähler cone of a compact Kähler manifold, and our goal is to understand how much cohomology classes on the boundary of the Kähler cone deviate from being Kähler. Let (X, ω) be an n -dimensional compact Kähler manifold. If α is a closed real $(1, 1)$ form on X , then we denote by $[\alpha]$ its cohomology class in $H^{1,1}(X, \mathbb{R})$. The set of cohomology classes of Kähler metrics is the Kähler cone $\mathcal{K} \subset H^{1,1}(X, \mathbb{R})$, which is an open convex cone. Classes in its closure $\overline{\mathcal{K}}$ are called nef (or numerically effective). A class $[\alpha]$ is said to have positive self-intersection if $\int_X \alpha^n > 0$. Cohomology classes which are nef and with positive self-intersection are quite close to being Kähler. In fact, thanks to a fundamental result of Demailly-Păun [20], such classes contain a Kähler current, which roughly speaking is a singular Kähler metric (see Section 2 for definitions). As a result, we will also refer to them as nef and big classes. Furthermore, thanks to Demailly's regularization theorem [17], the Kähler current can be chosen to have analytic singularities, which in particular implies that it is a smooth Kähler metric on a Zariski open subset of X . The largest possible subset of X that we can obtain in this way is called the ample locus of $[\alpha]$, denoted by $\text{Amp}(\alpha)$. Its complement $E_{nK}(\alpha) = X \setminus \text{Amp}(\alpha)$ is called the non-Kähler locus of $[\alpha]$, and it is a proper analytic subvariety of X . In a sense $E_{nK}(\alpha)$ measures the deviation from $[\alpha]$ to being Kähler, since $E_{nK}(\alpha) = \emptyset$ if and only if $[\alpha]$ is Kähler. Ample

*Supported in part by a Sloan Research Fellowship and NSF grants DMS-1236969 and DMS-1308988.

and non-Kähler loci were introduced by Boucksom [6], and they have been much studied recently in connection with the regularity for complex Monge-Ampère equations in big cohomology classes [3, 7, 19].

Given a nef and big class $[\alpha]$, there is another subset of X which measures its non-Kählerity, namely the null locus

$$\text{Null}(\alpha) = \bigcup_{\int_V \alpha^{\dim V} = 0} V,$$

where the union is over all positive dimensional irreducible analytic subvarieties $V \subset X$ where $[\alpha]$ integrates to zero. Note that since $\int_X \alpha^n > 0$, all such subvarieties are proper subsets of X . The null locus was explicitly first considered by Nakamaye [32], although it was probably studied earlier. Our main theorem is the following:

Theorem 1.1. *Let X be a compact complex manifold which is bimeromorphic to a Kähler manifold, let α be a closed smooth real $(1,1)$ -form on X whose class $[\alpha] \in H^{1,1}(X, \mathbb{R})$ is nef with $\int_X \alpha^n > 0$. Then*

$$(1.1) \quad E_{nK}(\alpha) = \text{Null}(\alpha).$$

Recall that compact complex manifolds bimeromorphic to Kähler are said to be in Fujiki's class \mathcal{C} [26, 56]. Equivalently, X is in class \mathcal{C} if there is a modification $\mu : \tilde{X} \rightarrow X$ with \tilde{X} Kähler.

In particular, Theorem 1.1 proves that $\text{Null}(\alpha)$ is a proper analytic subvariety of X , a fact which is far from clear from its definition (see [32] for a proof of this in the algebraic setting). Note that this theorem is trivially true if we drop the hypothesis that $\int_X \alpha^n > 0$, in which case $E_{nK}(\alpha) = \text{Null}(\alpha) = X$.

A simple example where this theorem applies is the following: let (Y^n, ω) be a compact Kähler manifold and $\pi : X \rightarrow Y$ be the blowup of a point $y \in Y$ with exceptional divisor $E = \pi^{-1}(y) \cong \mathbb{C}\mathbb{P}^{n-1}$. Let $[\alpha] = [\pi^*\omega]$, so $[\alpha]$ is nef and has positive self-intersection, with $\text{Null}(\alpha) = E$. It is well-known that there exists a smooth metric h on the line bundle $\mathcal{O}(E)$ with curvature $R(h)$ such that $\pi^*\omega - \varepsilon R(h)$ is a Kähler metric on X for all $\varepsilon > 0$ small. If s is a section of $\mathcal{O}(E)$ that vanishes along E to order 1, then for any ε small $\pi^*\omega + \varepsilon i\partial\bar{\partial} \log |s|_h^2$ is a Kähler current on X in the class α singular precisely along E .

This theorem has several diverse applications, which we now describe. The most direct corollary is an analytic proof of the following theorem in algebraic geometry, due to Nakamaye [32] in the case of \mathbb{Q} -divisors and to Ein-Lazarsfeld-Mustața-Nakamaye-Popa [22, Corollary 5.6] in general:

Corollary 1.2 ([22, 32]). *Let X be an n -dimensional smooth projective variety over \mathbb{C} and D a nef \mathbb{R} -divisor on X with $D^n > 0$. Then the augmented base locus $\mathbb{B}_+(D)$ of D satisfies $\mathbb{B}_+(D) = \text{Null}(D)$.*

Indeed the initial motivation for our work was to extend this result to all compact Kähler manifolds. For different extensions of this result to possibly

singular projective varieties, see [12] in positive characteristic, and [9] in characteristic zero.

From Theorem 1.1, one can also deduce a celebrated result of Demailly-Păun [20]. Our proof is not, by any means, independent of the work of [20], and it relies crucially on the mass concentration technique developed by Demailly-Păun. We are, however, able to avoid the complicated induction on dimension used in [20], which requires working on singular complex analytic spaces and using the results of Păun in [33].

Corollary 1.3 ([20], Theorem 0.1). *Let X be a compact Kähler manifold, and denote by \mathcal{K} the Kähler cone of X . Set*

$$\mathcal{P} := \left\{ [\alpha] \in H^{1,1}(X, \mathbb{R}) \left| \int_Y \alpha^{\dim Y} > 0, \forall Y \subset X \text{ irreducible} \right. \right. \\ \left. \left. \text{analytic subvariety, } \dim Y > 0 \right\}.$$

Then \mathcal{K} is one of the connected components of \mathcal{P} .

Next, as an application of our main Theorem 1.1, we answer affirmatively a question of Feldman-Illmanen-Knopf [25, Question 2, Section 10] and Campana (see [61, Section 4]) about finite time singularities of the Kähler-Ricci flow on compact Kähler manifolds. Let (X, ω_0) be a compact Kähler manifold of dimension n , and consider the Kähler-Ricci flow with initial metric ω_0

$$(1.2) \quad \frac{\partial}{\partial t} \omega = -\text{Ric}(\omega), \quad \omega(0) = \omega_0.$$

The flow always has a solution for short positive time [11], and a result of Tian-Zhang [52] characterizes the maximal existence time T of the flow as

$$T = \sup\{t > 0 \mid [\omega_0] - tc_1(X) \text{ is a Kähler class}\}.$$

Suppose that $T < \infty$, so the flow develops a finite time singularity. The limiting class of the metrics along the flow is $[\alpha] = [\omega_0] - Tc_1(X)$, which is nef but not Kähler. We will say that the Kähler Ricci flow does not develop a singularity at a point $x \in X$ if there is a neighborhood $U \ni x$ and a smooth Kähler metric ω_T on U such that $\omega(t)$ converge to ω_T in the C^∞ topology as $t \rightarrow T^-$. In fact, as we will see in Theorem 4.7, we can equivalently require that the curvature of $\omega(t)$ (or even just the scalar curvature) be bounded on U up to time T , which is a more standard definition of singularity formation in the Ricci flow (see [23]).

The following question was asked by Feldman-Illmanen-Knopf [25, Question 2, Section 10] and Campana (see [61, Section 4]).

Question 1.4. *Do singularities of the Kähler-Ricci flow develop precisely along analytic subvarieties of X ?*

It is not hard to see that if $\int_X \alpha^n = 0$ then the flow develops singularities everywhere on X (see Section 4), so the answer is affirmative in this case. We prove that this is true also when $\int_X \alpha^n > 0$.

Theorem 1.5. *Let (X^n, ω_0) be any compact Kähler manifold. Assume that the solution $\omega(t)$ of the Kähler-Ricci flow (1.2) starting at ω_0 exists on the maximal time interval $[0, T)$ with $T < \infty$, and that the limiting class $[\alpha] = [\omega_0] - Tc_1(X)$ satisfies $\int_X \alpha^n > 0$. Then as $t \rightarrow T^-$ the metrics $\omega(t)$ develop singularities precisely on the Zariski closed set $E_{nK}(\alpha) = \text{Null}(\alpha)$.*

Note that $\text{Null}(\alpha)$ is precisely the union of all irreducible analytic subvarieties of X whose volume with respect to $\omega(t)$ approaches zero as $t \rightarrow T^-$. So it is intuitively clear (and not hard to prove) that singularities should form at least along $\text{Null}(\alpha)$, and our main theorem implies that no singularities develop on its complement. Note also that in this setup, Zhang [62] proved that the Ricci curvature of $\omega(t)$ must be unbounded from below on X as $t \rightarrow T^-$. Our result shows that it remains locally uniformly bounded on compact sets away from $\text{Null}(\alpha)$. We also remark here that Tian has also asked a slightly different question in [50, p.81], [51, Conjecture 2], namely that for any finite-time singularities of the Kähler-Ricci flow, the weak limit as $t \rightarrow T^-$ of $\omega(t)$ as currents should be a current which is smooth at least on a Zariski open set. Theorem 1.5 gives a stronger answer in the case when $\int_X \alpha^n > 0$. Let us also note that in the case when X is projective and $[\omega_0]$ is a rational class, the fact that no singularities develop outside $\text{Null}(\alpha)$ follows from the work of Tian-Zhang [52], together with Nakamaye's result [32].

There are many examples of such finite-time singularities. In the case when the class $[\alpha]$ is the pullback of a Kähler class via the blowdown of an exceptional divisor, a delicate analysis of the singularity formation was done by Song-Weinkove [49]. See also [25, 30, 44, 45, 46, 50, 51, 52, 60] for more results about finite time singularities of the Kähler-Ricci flow. We also remark that if X is compact Kähler with nonnegative Kodaira dimension, and T is any finite time singularity of the Kähler-Ricci flow, then we must necessarily have that $\int_X (\omega_0 - Tc_1(X))^n > 0$, and our Theorem 1.5 applies (see Section 4).

The next application is to the study of degenerations of Ricci-flat Kähler metrics on Calabi-Yau manifolds. Let X be a compact Kähler manifold with $c_1(X) = 0$ in $H^2(X, \mathbb{R})$. We will call such a manifold a Calabi-Yau manifold. Let $\mathcal{K} \subset H^{1,1}(X, \mathbb{R})$ be the Kähler cone of X . Then Yau's theorem [58] says that for every Kähler class $[\alpha] \in \mathcal{K}$ there exists a unique Kähler metric $\omega \in [\alpha]$ with $\text{Ric}(\omega) = 0$. Let now $[\alpha] \in \partial\mathcal{K}$ be a $(1, 1)$ class on the boundary of the Kähler cone (so $[\alpha]$ is nef) with $\int_X \alpha^n > 0$. Let $[\alpha_t] : [0, 1] \rightarrow \overline{\mathcal{K}}$ be a path of $(1, 1)$ classes with $[\alpha_t] \in \mathcal{K}$ for $t > 0$ and with $[\alpha_0] = [\alpha]$. For $0 < t \leq 1$ write ω_t for the unique Ricci-flat Kähler metric in the class $[\alpha_t]$.

Theorem 1.6. *Let X^n be a Calabi-Yau manifold and $[\alpha] \in \partial\mathcal{K}$ be a nef class with $\int_X \alpha^n > 0$. Then there is a smooth incomplete Ricci-flat Kähler metric ω_0 on $X \setminus \text{Null}(\alpha)$ such that given any path $[\alpha_t]$, $0 < t \leq 1$, of Kähler*

classes approaching $[\alpha]$ as $t \rightarrow 0$, the Ricci-flat Kähler metrics ω_t in the class $[\alpha_t]$ converge to ω_0 as $t \rightarrow 0$ in the $C_{\text{loc}}^\infty(X \setminus \text{Null}(\alpha))$ topology. Moreover, (X, ω_t) converges as $t \rightarrow 0$ to the metric completion of $(X \setminus \text{Null}(\alpha), \omega_0)$ in the Gromov-Hausdorff topology.

The same result holds if instead of a path one considers a sequence of Kähler classes $[\alpha_i]$ converging to $[\alpha]$ (the proof is identical). In the special case when X is projective and the class $[\alpha]$ belongs to the real Néron-Severi group (i.e. it is the first Chern class of an \mathbb{R} -divisor), this was proved by the second author in [53], except for the statement about Gromov-Hausdorff convergence which is due to Rong-Zhang [38] (see also [39, 45]). In this case there is a birational contraction map $\pi : X \rightarrow Y$ to a singular Calabi-Yau variety, which is an isomorphism away from $\text{Null}(\alpha)$, and such that ω_0 is the pullback of a singular Kähler-Einstein metric on Y [24]. The contraction map is also used crucially to construct a smooth semipositive representative of the limiting class $[\alpha]$. The interested reader can also consult the surveys [54, 55]. In our more general setup there is no contraction map, but our theorem still shows that set where the metrics ω_t fail to converge smoothly to a limiting Kähler metric is precisely equal to $\text{Null}(\alpha)$, exactly as in the projective case (see Section 5).

The main technical ingredient in the proof of our main theorem is a technical extension-type result for Kähler currents with analytic singularities in a nef and big class which are defined on submanifold of X (Theorem 3.2). The problem of extending positive currents from subvarieties has recently generated a great deal of interest in both analytic and geometric applications [15, 28, 40, 57]. At the moment, the best results available are due to Coman-Guedj-Zeriahi [15], who proved that every positive current in a Hodge class on a (possibly singular) subvariety $V \subset \mathbb{C}\mathbb{P}^N$ is the restriction of a globally defined, positive current. Their proof uses a result of Coltoiu [14] on extending Runge subsets of analytic subsets of \mathbb{C}^N and a delicate, inductive gluing procedure. By contrast, the technique developed here is constructive, and relies only on a gluing property of plurisubharmonic functions with analytic singularities. The main technical tool is the use of resolution of singularities in order to obtain estimates which allow us to glue plurisubharmonic functions near their singularities. The ideas used here are similar, and indeed inspired by, techniques developed by Collins-Greenleaf-Pramanik [13] to obtain sharp estimates for singular integral operators. In turn, these were motivated by the seminal work of Phong, Stein and Sturm [34, 35, 36]. The advantage of these local techniques is that we can work on general Kähler manifolds and with arbitrary Kähler classes. In a forthcoming paper, we will refine these techniques and prove an extension theorem for Kähler currents with analytic singularities in Kähler classes on complex submanifolds.

This paper is organized as follows. In Section 2 we give some background and definitions, and show how Corollaries 1.2 and 1.3 follow from Theorem 1.1. The main Theorem 1.1 is proved in Section 3, where we also establish

an extension-type result for Kähler currents (Theorem 3.2). Section 4 is devoted to the study of finite time singularities of the Kähler-Ricci flow, and contains the proof of Theorem 1.5. Finally, Theorem 1.6 on degenerations of Calabi-Yau metrics is proved in Section 5.

Acknowledgments. We would like to thank D.H. Phong for all his advice and support, and S. Boucksom, S. Dinew, G. Székelyhidi, B. Weinkove, S. Zelditch, A. Zeriahi, Y. Zhang and Z. Zhang for comments and discussions. We also thank the referee for a useful suggestion which simplified our original proof of Theorem 3.2.

2. PRELIMINARIES

In this section we give the necessary definitions and background, and prove Corollaries 1.2 and 1.3 assuming our main Theorem 1.1.

Throughout this section and the next one, X is an n -dimensional compact complex manifold in Fujiki's class \mathcal{C} [26] (i.e. bimeromorphic to a compact Kähler manifold), and ω is a Hermitian metric on X . We will denote by $H^{1,1}(X, \mathbb{R})$ the Bott-Chern cohomology of d -closed real $(1, 1)$ forms (or currents) modulo $\partial\bar{\partial}$ -exact ones.

Let α be a smooth closed real $(1, 1)$ -form on X . We will say that a function φ on X is α -plurisubharmonic (α -PSH) if φ is in $L^1(X)$ and $\alpha + i\partial\bar{\partial}\varphi \geq 0$ as currents. In this case $T = \alpha + i\partial\bar{\partial}\varphi$ is a closed positive $(1, 1)$ current on X , in the cohomology class $[\alpha] \in H^{1,1}(X, \mathbb{R})$, and conversely every closed positive $(1, 1)$ current in the class $[\alpha]$ is of this form. For details about this, and other basic facts on closed positive currents, we refer the reader to the book of Demailly [18].

Closed positive currents are in general singular, and a convenient way to measure their singularities is the Lelong number $\nu(T, x)$. We can also define the Lelong sublevel sets

$$E_c(T) = \{x \in X \mid \nu(T, x) \geq c\},$$

where $c > 0$, and

$$E_+(T) := \bigcup_{c>0} E_c(T).$$

A fundamental result of Siu [42] shows that the Lelong sublevel sets $E_c(T)$ are in fact closed, analytic subvarieties of X . It follows that $E_+(T)$ is a countable union of analytic subvarieties. We also define a Kähler current to be a closed positive $(1, 1)$ current T such that $T \geq \varepsilon\omega$ holds as currents on X for some $\varepsilon > 0$. Thanks to a result of Demailly-Păun [20], a compact complex manifold is in class \mathcal{C} if and only if it admits a Kähler current. If a $(1, 1)$ class $[\alpha]$ contains a Kähler current, it is called big. This terminology comes from algebraic geometry: if X is projective and L is a holomorphic line bundle on X , then L is big (in the sense that $h^0(X, L^k) \sim k^n$ for k large) if and only if $c_1(L)$ is big in the above sense [16].

If X is Kähler, the set of all Kähler classes on X is the Kähler cone $\mathcal{K} \subset H^{1,1}(X, \mathbb{R})$. Classes in its closure $\overline{\mathcal{K}}$ are called nef (or numerically effective). A class $[\alpha]$ is nef if and only if for every $\varepsilon > 0$ there exists a smooth function ρ_ε on X such that $\alpha + i\partial\bar{\partial}\rho_\varepsilon \geq -\varepsilon\omega$. One then takes this as the definition of a nef class on a general compact complex manifold. It is easy to see that the pullback of a nef class by a holomorphic map between compact complex manifolds is still nef. Moreover, it is clear that if the class $c_1(L)$ is nef in the above sense then the line bundle L is nef in the usual algebro-geometric sense (i.e. $L \cdot C \geq 0$ for all curves $C \subset X$), and the converse is true when X is projective [16] and also when X is Moishezon thanks to a result of Păun [33], but false for general compact Kähler manifolds (a generic complex torus \mathbb{C}^n/Λ provides many counterexamples).

A useful notion, introduced by Demailly [17], is that of currents with analytic singularities. These are closed positive $(1, 1)$ currents $T = \alpha + i\partial\bar{\partial}\varphi$ such that given each point $x \in X$ there is an open neighborhood U of x with holomorphic functions f_1, \dots, f_N such that on U we have

$$(2.1) \quad \varphi = \delta \log \left(\sum_j |f_j|^2 \right) + g,$$

where g is a smooth function on U and δ is a nonnegative real number. In this case the current T is smooth away from an analytic subvariety of X , which in this case is exactly $E_+(T)$ (and also equal to $E_c(T)$ for some $c > 0$). Then Demailly's regularization procedure [17] shows that if a class $[\alpha]$ is big, then it also contains a Kähler current with analytic singularities. We will also say that a positive $(1, 1)$ current T has *weakly analytic singularities* if its potentials are locally of the form (2.1) with g continuous.

A fundamental theorem of Demailly-Păun [20, Theorem 2.12] states that if X is a compact Kähler manifold, and $[\alpha]$ is nef with $\int_X \alpha^n > 0$, then $[\alpha]$ contains a Kähler current (i.e. $[\alpha]$ is big).

Finally, we introduce a number of different subsets of X , all of which aim to measure the defect from a class $[\alpha]$ being Kähler. In the rational case, the sets measure the defect from the corresponding \mathbb{Q} -divisor being ample.

Definition 2.1. *Let (X, ω) be an n -dimensional compact complex manifold in class \mathcal{C} equipped with a Hermitian metric, and let $[\alpha] \in H^{1,1}(X, \mathbb{R})$ be a big class.*

- (i) *Following [6], we define the non-Kähler locus, denoted $E_{nK}(\alpha)$, to be the set*

$$E_{nK}(\alpha) := \bigcap_{T \in [\alpha]} E_+(T),$$

where the intersection is taken over all Kähler currents in the class $[\alpha]$.

(ii) We define the null locus, denoted by $\text{Null}(\alpha)$, to be the set

$$\text{Null}(\alpha) := \bigcup_{\int_V \alpha^{\dim V} = 0} V$$

where the union is taken over all positive dimensional irreducible analytic subvarieties of X .

(iii) Following [7], we define the ample locus, denoted by $\text{Amp}(\alpha)$ to be the set of points $x \in X$ such that there exists a Kähler current $T \in [\alpha]$ with analytic singularities, which is smooth in a neighborhood of x .

Finally, suppose X is a smooth projective variety, and

$$[\alpha] \in NS(X, \mathbb{R}) = (H^2(X, \mathbb{Z})_{\text{free}} \cap H^{1,1}(X)) \otimes \mathbb{R},$$

belongs to the real Néron-Severi group, so $[\alpha] = c_1(D)$ for D an \mathbb{R} -divisor:

(iv) We define the augmented base locus, denoted $\mathbb{B}_+(\alpha)$, or $\mathbb{B}_+(D)$ to be the stable base locus of $D - A$ for some small, ample \mathbb{R} -divisor A with $D - A$ a \mathbb{Q} -divisor. That is, $\mathbb{B}_+(\alpha)$ is the base locus of the linear series $|m(D - A)|$ for m sufficiently large and divisible.

The definition of $\mathbb{B}_+(D)$ in (iv) is well-posed, since we get the same stable base locus for all sufficiently small ample \mathbb{R} -divisors A with $D - A$ a \mathbb{Q} -divisor [21, Proposition 1.5], and $\mathbb{B}_+(D)$ depends only on the numerical class of D (so the notation $\mathbb{B}_+(\alpha)$ is justified). Furthermore, $\mathbb{B}_+(D)$ equals the intersection of the supports of E for all “Kodaira’s Lemma”-type decompositions $D = A + E$, where A is an ample \mathbb{R} -divisor and E is an effective \mathbb{R} -divisor [21, Remark 1.3].

In [6, Theorem 3.17 (ii)], Boucksom observed that in a big class we can always find a Kähler current with analytic singularities precisely along $E_{nK}(\alpha)$. Combining this statement with the theorem of Demailly-Păun [20] and Demailly’s regularization [17], we get the following:

Theorem 2.2. *Let X^n be a compact complex manifold in class \mathcal{C} and $[\alpha] \in H^{1,1}(X, \mathbb{R})$ be a nef class with $\int_X \alpha^n > 0$. Then there exists a Kähler current T in the class $[\alpha]$, with analytic singularities such that*

$$E_+(T) = E_{nK}(\alpha).$$

In particular $E_{nK}(\alpha)$ is an analytic subvariety of X .

Indeed, since X is in class \mathcal{C} there is a modification $\mu : \tilde{X} \rightarrow X$ with \tilde{X} a compact Kähler manifold [26, 56]. The class $[\mu^*\alpha]$ is then nef and satisfies $\int_{\tilde{X}} (\mu^*\alpha)^n = \int_X \alpha^n > 0$, and so it contains a Kähler current T by Demailly-Păun [20]. It is then easy to check that the pushforward μ_*T is a Kähler current on X in the class $[\alpha]$. Then Demailly’s regularization [17] and Boucksom’s argument [6, Theorem 3.17 (ii)] (neither of which needs X to be Kähler) complete the proof.

Using this result, we can show how Corollary 1.3 follows from Theorem 1.1.

Proof of Corollary 1.3. The inclusion $\mathcal{K} \subset \mathcal{P}$ is clear. Moreover, \mathcal{K} is clearly connected and open. Suppose $[\alpha] \in \overline{\mathcal{K}} \cap \mathcal{P}$. Then $[\alpha]$ is nef and $\int_X \alpha^n > 0$. Moreover, since $[\alpha] \in \mathcal{P}$, we have $\text{Null}(\alpha) = \emptyset$. By Theorem 1.1 we have that $E_{nK}(\alpha) = \emptyset$. Thanks to Theorem 2.2 there exists $T \in [\alpha]$ a Kähler current with analytic singularities and with $E_+(T) = \emptyset$, which means that T is a smooth Kähler metric. Thus, $[\alpha] \in \mathcal{K}$, and so \mathcal{K} is closed in \mathcal{P} . Since \mathcal{K} is connected, the corollary follows. \square

There are some relations among the subsets of X introduced in Definition 2.1. The following two propositions are available in the literature, and we include proofs for the convenience of the reader.

Proposition 2.3 (Boucksom [6]). *Let X^n be a compact complex manifold in class \mathcal{C} and $[\alpha] \in H^{1,1}(X, \mathbb{R})$ be a nef class with $\int_X \alpha^n > 0$. Then $E_{nK}(\alpha) = \text{Amp}(\alpha)^c$.*

Proof. Apply Theorem 2.2, to get a Kähler current $T \in [\alpha]$ with analytic singularities with $E_+(T) = E_{nK}(\alpha)$. Since T has analytic singularities, it is smooth precisely wherever its Lelong number vanishes, so T is smooth off $E_+(T)$. This shows that $\text{Amp}(\alpha)^c \subset E_{nK}(\alpha)$. The reverse inclusion is trivial. \square

Corollary 1.2 follows immediately from Theorem 1.1 and the following:

Proposition 2.4 (Boucksom [4], Corollary 2.2.8). *If $[\alpha]$ is the class of a big and nef \mathbb{R} -divisor D on a smooth projective variety X over \mathbb{C} , then $E_{nK}(\alpha) = \mathbb{B}_+(\alpha)$.*

Proof. For the inclusion $\mathbb{B}_+(\alpha) \subset E_{nK}(\alpha)$ we essentially follow [6, Proposition 5.2]. If $x \notin E_{nK}(\alpha)$, then there exists a Kähler current T in the class $[\alpha]$ with analytic singularities and smooth in a coordinate neighborhood U of x . Choose A an ample divisor and $\delta > 0$ a small constant such that $D - \delta A$ is a \mathbb{Q} -divisor. Fixing ω a Kähler metric in $c_1(A)$, and taking δ smaller if necessary, we can assume that $T - \delta\omega$ is still a Kähler current. Let θ be a smooth cutoff function supported in U and identically 1 near x , and let

$$\tilde{T} = T - \delta\omega + \varepsilon i \partial \bar{\partial} (\theta \log |z - x|),$$

where $\varepsilon > 0$ is small enough so that \tilde{T} is a Kähler current in $c_1(D - \delta A)$. The Lelong number of \tilde{T} at x is ε , and \tilde{T} is smooth on $U \setminus \{x\}$. Let β be a smooth representative of $-c_1(X)$, and let $\tilde{T}_m = m\tilde{T} - \beta$. For m sufficiently large, \tilde{T}_m is a Kähler current in $c_1(m(D - \delta A) - K_X)$, smooth on $U \setminus \{x\}$ and with Lelong number $m\varepsilon$ at x . If m is sufficiently divisible, then $m(D - \delta A)$ corresponds to a holomorphic line bundle L_m . Thanks to Skoda's lemma [6, Lemma 5.1], if m is large, the subvariety cut out by the multiplier ideal sheaf $\mathcal{I}(\tilde{T}_m)$ of \tilde{T}_m intersects U in the isolated point x . Nadel's vanishing theorem then implies that $H^1(X, L_m \otimes \mathcal{I}(\tilde{T}_m)) = 0$, and therefore the restriction map $H^0(X, L_m) \rightarrow (L_m)_x$ is surjective. Hence there is a global section of L_m that does not vanish at x . This means that $x \in \mathbb{B}_+(\alpha)^c$.

To prove the reverse inclusion, take a point $x \in \mathbb{B}_+(\alpha)^c$, so by definition there is an ample \mathbb{R} -divisor A such that the stable base locus of $D - A$ does not contain x . This means that there is a large integer m such that $mD - mA = c_1(L)$ for some line bundle L and there is an effective \mathbb{Z} -divisor E linearly equivalent to $mD - mA$ which does not pass through x . Take $s \in H^0(X, L)$ a defining section for E , and complete it to a basis $\{s = s_1, s_2, \dots, s_N\}$ of $H^0(X, L)$. Fixing a smooth Hermitian metric h on L with curvature form $R(h)$ we can define a closed positive current $T_L = R(h) + i\partial\bar{\partial} \log \sum_i |s_i|_h^2 \in c_1(L)$ which has analytic singularities and is smooth near x . If ω is a Kähler metric in $c_1(A)$, then $\frac{1}{m}T_L + \omega$ is then a Kähler current on X in the class $[\alpha]$ with analytic singularities and smooth near x , so $x \in \text{Amp}(\alpha)^c$, which equals $E_{nK}(\alpha)^c$ by Proposition 2.3. \square

Let us discuss the structure of $\text{Null}(\alpha)$ when $n = 2$, so X is a complex surface and $[\alpha]$ is a nef class with positive self-intersection which is assumed not to be Kähler. It is well-known that complex surfaces in class \mathcal{C} are Kähler [1]. In this case, $\text{Null}(\alpha)$ is the union of irreducible curves C_i , $1 \leq i \leq N$, with the intersection matrix $(C_i \cdot C_j)$ negative definite. Indeed for any real numbers λ_i we have $\int_X \alpha^2 > 0$ and $\int_{\sum_i \lambda_i C_i} \alpha = 0$, so by the Hodge index theorem [1, Corollary IV.2.16] either $\sum_i \lambda_i C_i = 0$ or $(\sum_i \lambda_i C_i)^2 < 0$. This means that the intersection form is negative definite on the linear span of the curves C_i , which proves our assertion. Grauert's criterion [1, Theorem III.2.1] shows that each connected component of $\text{Null}(\alpha)$ can be contracted, so we get a map $\pi : X \rightarrow Y$ where Y is a normal complex surface. In general though, Y is not a Kähler space.

We conclude this section by proving the easy inclusion in our main Theorem 1.1. An alternative proof of this result (in the case when X is Kähler) can be found in [10, Theorem 2.2].

Theorem 2.5. *Let X be a compact complex manifold in class \mathcal{C} , let α be a closed smooth real $(1, 1)$ -form on X whose class $[\alpha] \in H^{1,1}(X, \mathbb{R})$ is nef with $\int_X \alpha^n > 0$. Then*

$$(2.2) \quad \text{Null}(\alpha) \subset E_{nK}(\alpha).$$

Proof. Fix a point $x \in X$ and an irreducible analytic subvariety $V \subset X$ of dimension k , such that $x \in V$, and $\int_V \alpha^k = 0$. For the sake of obtaining a contradiction, suppose also that $x \in \text{Amp}(\alpha)$ so there exists a Kähler current $T \in [\alpha]$ with analytic singularities, smooth in a neighborhood of x . Since V is a subvariety of X which is in \mathcal{C} , it follows from [26] that V is also in class \mathcal{C} , so there is a modification $\mu : \tilde{V} \rightarrow V$ with \tilde{V} a compact Kähler manifold of dimension k . The class $[\mu^*\alpha]$ is nef and satisfies $\int_{\tilde{V}} (\mu^*\alpha)^k = 0$, and it contains the closed positive $(1, 1)$ current μ^*T (defined as usual by writing $T = \alpha + i\partial\bar{\partial}\varphi$ and letting $\mu^*T = \mu^*\alpha + i\partial\bar{\partial}(\varphi \circ \mu)$). Furthermore μ^*T is a smooth Kähler metric on a nonempty open set $\tilde{U} \subset \tilde{V}$. By [7, Proposition 1.6] the non-pluripolar product $\langle \mu^*T \rangle^k$ is a well-defined closed

positive (k, k) -current on \tilde{V} . By [7, Proposition 1.20] we have that

$$0 \leq \int_{\tilde{V}} \langle \mu^* T \rangle^k \leq \int_{\tilde{V}} (\mu^* \alpha)^k = 0,$$

so $\int_{\tilde{V}} \langle \mu^* T \rangle^k = 0$. But $\mu^* T$ is a smooth Kähler metric on \tilde{U} , and so

$$0 < \int_{\tilde{U}} (\mu^* T)^k = \int_{\tilde{U}} \langle \mu^* T \rangle^k \leq \int_{\tilde{V}} \langle \mu^* T \rangle^k,$$

which is a contradiction. \square

3. PROOF OF THEOREM 1.1

In this section we give the proof of Theorem 1.1. The idea is to observe that if there is a point $x \in \text{Null}(\alpha)^c \cap E_{nK}(\alpha)$, then there is an entire analytic subvariety $V \subset E_{nK}(\alpha)$ of positive dimension, for which the restricted class is again nef and with positive self-intersection. Moreover, there is a Kähler current R in the class $[\alpha]$ with analytic singularities along $E_{nK}(\alpha)$ which by assumption has positive Lelong numbers at every point of V . Assume for the moment that V is smooth. Using the mass concentration technique of Demailly-Păun [20], it follows that V supports a Kähler current T in the class $[\alpha|_V]$ with analytic singularities. If we can use T to produce a Kähler current in the class $[\alpha]$ defined on all of X and $T|_V$ is smooth near the generic point of V , then the theorem follows, since the generic point of V is then contained in $E_{nK}(\alpha)^c$, contradicting the definition of V . The case of V singular is similar after passing to a resolution of singularities.

We start with the following lemma, which is the analytic counterpart of [22, Proposition 1.1] in the algebraic case. Note that by definition $\text{Null}(\alpha)$ does not have any isolated points.

Lemma 3.1. *Let X be a compact complex manifold in class \mathcal{C} and $[\alpha]$ a real $(1, 1)$ class which is nef and with positive self-intersection. Then the analytic set $E_{nK}(\alpha)$ does not have any isolated points.*

Proof. If $x \in E_{nK}(\alpha)$ is an isolated point, then we can find a small open set $U \subset X$ which contains the point x , such that $U \cap E_{nK}(\alpha) = \{x\}$. By shrinking U if necessary, we can assume that U has coordinates $z = (z_1, \dots, z_n)$. If A is a sufficiently large constant, then on U we have

$$\alpha + i\partial\bar{\partial}(A|z|^2 - C) \geq \omega,$$

for any constant C . Let $R = \alpha + i\partial\bar{\partial}F$ be a Kähler current in $[\alpha] \in H^{1,1}(X, \mathbb{R})$ which has analytic singularities, which exists thanks to a theorem of Demailly-Păun [20] and Demailly's regularization [17]. Furthermore, we can choose R with the property that $E_+(R) = E_{nK}(\alpha)$, thanks to Theorem 2.2, and R is smooth outside of $E_+(R)$. In particular F is smooth near ∂U , and so we can choose C sufficiently large such that $\max\{F, A|z|^2 - C\} = F$ near ∂U . Fix a small $\varepsilon > 0$ and let \max_ε be the regularized maximum function (see, e.g. [18, I.5.18]). Then $\tilde{F} = \max_\varepsilon(F, A|z|^2 - C)$ equals F near

∂U and equals $A|z|^2 - C$ near x , is smooth on U , and so $\tilde{R} = \alpha + i\partial\bar{\partial}\tilde{F}$ is a Kähler current on X in the class $[\alpha]$ with analytic singularities and with $x \notin E_+(\tilde{R})$, from which it follows that $x \notin E_{nK}(\alpha)$, a contradiction. \square

Proof of Theorem 1.1. Thanks to Theorem 2.5, we only need to show that $E_{nK}(\alpha) \subset \text{Null}(\alpha)$. For the sake of obtaining a contradiction, assume the result does not hold, and there exists $x \in \text{Null}(\alpha)^c \cap E_{nK}(\alpha)$. Let V be an irreducible component of the analytic set $E_{nK}(\alpha)$ passing through x . Here and in the following whenever we deal with analytic sets, we always consider the underlying reduced variety. From Lemma 3.1, we must have $\dim V > 0$. Thus, we have that $V \subset E_{nK}(\alpha)$ is a positive dimensional irreducible analytic subvariety, with $x \in V$. Moreover, since $x \in \text{Null}(\alpha)^c$, we have that $[\alpha|_V]$ has positive self-intersection, and if V is smooth then $[\alpha|_V]$ is also nef (this is true also for V singular, provided one properly defines nef classes on singular varieties, which we will not need to do). We will need the following theorem:

Theorem 3.2. *Let X be a compact complex manifold in class \mathcal{C} and α a closed smooth real $(1,1)$ form on X with $[\alpha]$ nef and $\int_X \alpha^n > 0$. Let $E = V \cup \bigcup_{i=1}^I Y_i$ be an analytic subvariety of X , with V, Y_i its irreducible components, and V a positive dimensional compact complex submanifold of X . Let $R = \alpha + i\partial\bar{\partial}F$ be a Kähler current in the class $[\alpha]$ on X with analytic singularities precisely along E and let $T = \alpha|_V + i\partial\bar{\partial}\varphi$ be a Kähler current in the class $[\alpha|_V]$ on V with analytic singularities. Then there exists a Kähler current $\tilde{T} = \alpha + i\partial\bar{\partial}\Phi$ in the class $[\alpha]$ on X with $\tilde{T}|_V$ smooth in a neighborhood of the generic point of V .*

Let us complete the proof of Theorem 1.1 assuming this result. Assume first that V is smooth. Then by Demailly-Păun [20] (see Theorem 2.2) there exists a Kähler current $T = \alpha|_V + i\partial\bar{\partial}\varphi \in [\alpha|_V]$. Furthermore thanks to Demailly's regularization [17], we can choose T to have analytic singularities, so in particular T is smooth at the generic point of V . We also write

$$E_{nK}(\alpha) = V \cup \bigcup_{i=1}^I Y_i,$$

for distinct irreducible components Y_i of $E_{nK}(\alpha)$, which are all positive dimensional thanks to Lemma 3.1.

We then apply Theorem 3.2 to $E = E_{nK}(\alpha)$, with the current R constructed as in the proof of Lemma 3.1, and we conclude that there is a Kähler current $\tilde{T} = \alpha + i\partial\bar{\partial}\Phi$ in the class $[\alpha]$ on X with $\tilde{T}|_V$ smooth in a neighborhood of the generic point of V . In particular, for a generic $x \in V$ we have $\nu(\tilde{T}|_V, x) = 0$. But we always have that the Lelong number of \tilde{T} at x calculated in the ambient space X satisfies $\nu(\tilde{T}, x) \leq \nu(\tilde{T}|_V, x)$, hence \tilde{T} has Lelong number zero at a generic point of V , and so $V \not\subset E_+(\tilde{T})$. This

immediately leads to a contradiction in this case, because by definition

$$V \subset E_{nK}(\alpha) \subset E_+(\tilde{T}).$$

If V is not smooth we take $\pi : \tilde{X} \rightarrow X$ be an embedded resolution of singularities of V , obtained by blowing up smooth centers (which exists thanks to work of Hironaka), and let \tilde{V} be the strict transform of V , which is now a smooth submanifold of \tilde{X} . The manifold \tilde{X} is in class \mathcal{C} , and $\pi : \tilde{V} \rightarrow V$ is an isomorphism over V_{reg} . Note that since π is bimeromorphic, we have $\pi_*\pi^*\alpha = \alpha$ as currents. Therefore, if T is a Kähler current in the class $[\pi^*\alpha]$ on \tilde{X} then its pushforward π_*T is a Kähler current in the class $[\alpha]$ on X .

The class $[(\pi^*\alpha)|_{\tilde{V}}]$ is nef and has positive self-intersection on \tilde{V} , and hence by Demailly-Păun [20, Theorem 2.12] and Demailly's regularization there exists a Kähler current $T \in [(\pi^*\alpha)|_{\tilde{V}}]$ with analytic singularities. Furthermore, the class $[\pi^*\alpha]$ is nef and has positive self-intersection on \tilde{X} , so by the same reason there exists a Kähler current $R \in [\pi^*\alpha]$ with analytic singularities along an analytic subset E of \tilde{X} . We must have that $\tilde{V} \subset E$, because if not then R is smooth near the generic point of \tilde{V} , and then the pushforward π_*R would be a Kähler current in the class $[\alpha]$ on X which is smooth at the generic point of V , which contradicts $V \subset E_{nK}(\alpha)$.

We can then apply Theorem 3.2 and get a Kähler current \tilde{T} on \tilde{X} in the class $[\pi^*\alpha]$ with $\tilde{T}|_{\tilde{V}}$ smooth near the generic point of \tilde{V} . Therefore $\pi_*\tilde{T}$ is then a Kähler current on X in the class $[\alpha]$ whose restriction to V is smooth near the generic point of V (note that, in general, it may be the case that $\pi_*\tilde{T}$ has non-vanishing Lelong numbers along V_{sing} , which is however a proper subvariety of V). Again, we have a contradiction, which proves our main theorem. \square

We now turn to the proof of Theorem 3.2. The proof is somewhat technical, so we briefly outline the main steps. First, we perform a resolution of singularities to replace $\cup_i Y_i$ with a divisor with simple normal crossings with the strict transform \tilde{V} of V . We then construct local extensions on a finite open covering of \tilde{V} , which have the same singularities on any overlap. This is a crucial point, which allows us to use a technique of Richberg [37] to glue the local potentials to construct a Kähler current in a neighborhood of \tilde{V} . Lastly, after possibly reducing the Lelong numbers of the global Kähler current R , we can glue the Kähler current on this neighborhood to R , to obtain a global Kähler current.

Proof of Theorem 3.2. Define an analytic set $G \subset X$ by

$$G = \bigcup_i Y_i \cup E_+(T),$$

and let \mathcal{I}_G be its defining ideal sheaf. Let $\pi_1 : \tilde{X}_1 \rightarrow X$ be a principalization of \mathcal{I}_G obtained by blowing up smooth centers, so that $\pi_1^{-1}(G)$ is a union

of smooth divisors with simple normal crossings. Composing π_1 with an embedded resolution of the strict transform of V in \tilde{X}_1 we obtain a birational morphism $\pi : \tilde{X} \rightarrow X$ (with \tilde{X} smooth), which is a composition of blowups with smooth centers, such that

$$\pi^{-1}\mathcal{I}_G = \mathcal{O}_{\tilde{X}} \left(- \sum_{\ell} a_{\ell} D_{\ell} \right), \quad \pi^{-1}(G) = \bigcup_{\ell} D_{\ell} \cup \bigcup_{\ell} E_{\ell},$$

where $a_{\ell} > 0$, $\text{Exc}(\pi) = \sum_{\ell} E_{\ell}$ and $\cup_{\ell} D_{\ell} \cup \cup_{\ell} E_{\ell}$ is a union of smooth divisors with simple normal crossings, which furthermore has normal crossings with \tilde{V} , the strict transform of V . This means that \tilde{V} is a smooth submanifold of \tilde{X} , which can be covered by finitely many charts $\{W_j\}_{1 \leq j \leq N}$ such that on each W_j there are local coordinates (z_1, \dots, z_n) with $\tilde{V} \cap W_j = \{z_1 = \dots = z_{n-k} = 0\}$, where $k = \dim V$, and with $\pi^{-1}(G) \cap W_j = \{z_{i_1} \dots z_{i_p} = 0\}$, $n - k < i_1, \dots, i_p \leq n$ (see e.g. [2, Theorem 2.3]). Write $z = (z_1, \dots, z_{n-k})$ and $z' = (z_{n-k+1}, \dots, z_n)$. Note that in general some divisor in $\sum_{\ell} E_{\ell}$ might also appear in $\sum_{\ell} D_{\ell}$, but this will not be an issue.

Choose $\varepsilon > 0$ small enough so that

$$T = \alpha|_V + i\partial\bar{\partial}\varphi \geq 3\varepsilon\omega|_V,$$

holds as currents on V . Pulling back to \tilde{X} we obtain

$$\pi^*\alpha|_{\tilde{V}} + i\partial\bar{\partial}(\varphi \circ \pi) \geq 3\varepsilon\pi^*\omega|_{\tilde{V}}.$$

Let s_D be a defining section of $\mathcal{O}_{\tilde{X}}(\sum_{\ell} a_{\ell} D_{\ell})$ and s_E a defining section for $\mathcal{O}_{\tilde{X}}(\sum_{\ell} E_{\ell})$. The smooth form $\pi^*\omega$ is not Hermitian on \tilde{X} , but there is a small $\delta > 0$ such that

$$\pi^*\omega + \delta i\partial\bar{\partial} \log |s_E|_{h_E}^2 \geq 2\tilde{\omega},$$

for some Kähler metric $\tilde{\omega}$ on \tilde{X} , where h_E is a suitable smooth metric on $\mathcal{O}_{\tilde{X}}(\sum_{\ell} E_{\ell})$ (see e.g. [27, p.187]). We fix h_D any smooth metric on $\mathcal{O}_{\tilde{X}}(\sum_{\ell} a_{\ell} D_{\ell})$, and we have

$$i\partial\bar{\partial} \log |s_D|_{h_D}^2 \geq -R(h_D),$$

as currents on \tilde{X} , where $R(h_D)$ denotes the curvature form of h_D . Hence we can choose $\delta_1 > 0$ small enough so that

$$\pi^*\omega + \delta i\partial\bar{\partial} \log |s_E|_{h_E}^2 + \delta_1 i\partial\bar{\partial} \log |s_D|_{h_D}^2 \geq \tilde{\omega},$$

on \tilde{X} . Then

$$\pi^*\alpha|_{\tilde{V}} + i\partial\bar{\partial}((\varphi \circ \pi) + 3\varepsilon\delta \log |s_E|_{h_E}^2 + 3\varepsilon\delta_1 \log |s_D|_{h_D}^2) \geq 3\varepsilon\tilde{\omega}|_{\tilde{V}}.$$

For simplicity of notation, define $\hat{\varphi} = (\varphi \circ \pi) + 3\varepsilon\delta \log |s_E|_{h_E}^2 + 3\varepsilon\delta_1 \log |s_D|_{h_D}^2$, which is a function on \tilde{V} . Define a function φ_j on W_j (with analytic singularities) by

$$\varphi_j(z, z') = (\varphi \circ \pi)(z') + A|z|^2 + 3\varepsilon\delta \log |s_E|_{h_E}^2 + 3\varepsilon\delta_1 \log |s_D|_{h_D}^2,$$

where $A > 0$ is a constant. If we shrink the W_j 's slightly, still preserving the property that $\tilde{V} \subset \cup_j W_j$, we can choose A sufficiently large so that

$$\pi^* \alpha + i\partial\bar{\partial}\varphi_j \geq 2\varepsilon\tilde{\omega},$$

holds on W_j for all j . It will also be useful to fix slightly smaller open sets $W'_j \Subset U_j \Subset W_j$ such that $\cup_j W'_j$ still covers \tilde{V} . Note that since φ is smooth at the generic point of V , by construction all functions φ_j are also smooth in a neighborhood of the generic point of $\tilde{V} \cap W_j$.

We wish to glue the functions φ_j together to produce a Kähler current defined in a neighborhood of \tilde{V} in \tilde{X} . This would be straightforward if the functions φ_j were continuous, thanks to a procedure of Richberg [37], but in our case the functions φ_j have poles along $\cup_\ell D_\ell \cup \cup_\ell E_\ell \cap W_j$. However, as we will now show, on any nonempty intersection $W_i \cap W_j$ the differences $\varphi_i - \varphi_j$ are continuous, and this is enough to apply the argument of Richberg.

So take two open sets W_1, W_2 in the covering, with $W_1 \cap W_2 \cap \tilde{V}$ nonempty. Let (z_1^1, \dots, z_n^1) be the coordinates on W_1 as above, and (z_1^2, \dots, z_n^2) the ones on W_2 . If none of the divisors D_ℓ, E_ℓ intersects $W_1 \cap W_2$, then $\varphi_i - \varphi_j$ is clearly smooth there, so we may assume that at least one of them, D_ℓ say, intersects $W_1 \cap W_2$. Up to reordering the coordinates, we can write $D_\ell \cap W_1 = \{z_\ell^1 = 0\}$ and $D_\ell \cap W_2 = \{z_\ell^2 = 0\}$. Therefore on $W_1 \cap W_2$ we must have $z_\ell^2 = u_\ell \cdot z_\ell^1$, where u_ℓ is a never-vanishing holomorphic function.

From the construction of the resolution π , and the fact that φ_j have analytic singularities along $\cup_\ell D_\ell \cup \cup_\ell E_\ell \cap W_j$ (in fact, we will only use the fact that they have weakly analytic singularities), we have on W_1

$$(3.1) \quad \varphi_1 = c_1 \log \left(\prod_k |z_{i_k}^1|^{2\alpha_{i_k}^1} \right) + g_1,$$

where $c_1, \alpha_{i_k}^1$ are constants, g_1 is a continuous function, and $\{\prod_k |z_{i_k}^1|^{2\alpha_{i_k}^1} = 0\}$ equals $\cup_\ell D_\ell \cup \cup_\ell E_\ell \cap W_1$. A similar formula holds for φ_2 on W_2 , and since $\varphi_1|_{\tilde{V}} = \varphi_2|_{\tilde{V}}$ and $z_{i_k}^j|_{\tilde{V}} \neq 0, j = 1, 2$, we must have $c_1 = c_2$ and $\alpha_{i_k}^1 = \alpha_{i_k}^2$. Thanks to the relations $z_{i_k}^2 = u_{i_k} \cdot z_{i_k}^1$ we see that $\varphi_1 - \varphi_2$ is indeed continuous on $W_1 \cap W_2$. Furthermore, on $W_1 \cap W_2$ the function $\max\{\varphi_1, \varphi_2\}$ differs from φ_1 and φ_2 by a continuous function, and hence it has weakly analytic singularities of the same type as φ_1 and φ_2 , and restricts to $\hat{\varphi}$ on $\tilde{V} \cap W_1 \cap W_2$.

Now, for the convenience of the reader, we recall the gluing argument of Richberg [37]. We start by considering two open sets W_1, W_2 in the covering with $W'_1 \cap W'_2 \cap \tilde{V}$ nonempty, and fix a compact set $K \subset \tilde{V}$ with $(W'_1 \cup W'_2) \cap \tilde{V} \subset K \subset (U_1 \cup U_2) \cap \tilde{V}$. Let $M_1 = K \cap \partial U_2, M_2 = K \cap \partial U_1$, so that M_1 and M_2 are disjoint compact subsets of \tilde{V} . This setup is depicted in figure 1.

Pick θ_1 a smooth nonnegative cutoff function which is identically 1 in a neighborhood of M_2 in \tilde{X} and θ_2 a smooth nonnegative cutoff function




FIGURE 1. The setup for the local Richberg argument.

which is identically 1 in a neighborhood of M_1 in \tilde{X} so that the supports of θ_1 and θ_2 are disjoint. Then, if we choose $\eta > 0$ small, the functions

$$(3.2) \quad \tilde{\varphi}_j = \varphi_j - \eta\theta_j,$$

$j = 1, 2$ have analytic singularities and satisfy $\pi^*\alpha + i\partial\bar{\partial}\tilde{\varphi}_j \geq \varepsilon\tilde{\omega}$ on W_j . On $W_1 \cap W_2$ we then define

$$\tilde{\varphi}_0 = \max\{\tilde{\varphi}_1, \tilde{\varphi}_2\},$$

which has weakly analytic singularities, it satisfies $\pi^*\alpha + i\partial\bar{\partial}\tilde{\varphi}_0 \geq \varepsilon\tilde{\omega}$ and it restricts to $\tilde{\varphi}$ on $\tilde{V} \cap W_1 \cap W_2$. Consider now a neighborhood of M_2 in \tilde{X} , small enough so that $\theta_1 = 1$ on M_2 . Since φ_1, φ_2 agree on \tilde{V} and $\varphi_1 - \varphi_2$ is continuous on this neighborhood, we see that there exists a possibly smaller such neighborhood where

$$\tilde{\varphi}_1 = \varphi_1 - \eta < \varphi_2 = \tilde{\varphi}_2,$$

so that $\tilde{\varphi}_0 = \tilde{\varphi}_2$ there. Similarly, on any sufficiently small neighborhood of M_1 we have $\tilde{\varphi}_0 = \tilde{\varphi}_1$. Therefore there is an open neighborhood W_0 of $K \cap \overline{U_1} \cap \overline{U_2}$ in \tilde{X} such that $\tilde{\varphi}_0 = \tilde{\varphi}_1$ on $W_0 \setminus \overline{U_2}$ and $\tilde{\varphi}_0 = \tilde{\varphi}_2$ on $W_0 \setminus \overline{U_1}$. Therefore we can define

$$W' = W_0 \cup (U_1 \setminus \overline{U_2}) \cup (U_2 \setminus \overline{U_1}),$$

which is a neighborhood of K in \tilde{X} , and define a function φ' on W' to be equal to $\tilde{\varphi}_0$ on W_0 , equal to $\tilde{\varphi}_1$ on $U_1 \setminus \overline{U_2}$ and equal to $\tilde{\varphi}_2$ on $U_2 \setminus \overline{U_1}$. Then φ' satisfies $\pi^*\alpha + i\partial\bar{\partial}\varphi' \geq \varepsilon\tilde{\omega}$, it has weakly analytic singularities of the same type as φ_1 and φ_2 , and restricts to $\tilde{\varphi}$ on $W' \cap \tilde{V}$. Clearly, W' contains $(W'_1 \cup W'_2) \cap \tilde{V}$.

We then replace W_1 and W_2 with W' , replace φ_1 and φ_2 with φ' , and repeat the same procedure with two other open sets in this new covering. After at most N such steps, we end up with an open neighborhood W of \tilde{V}

in \tilde{X} with a function φ'' defined on W which satisfies $\pi^*\alpha + i\partial\bar{\partial}\varphi'' \geq \varepsilon'\tilde{\omega}$ for some $\varepsilon' > 0$, it has weakly analytic singularities, and it restricts to $\hat{\varphi}$ on \tilde{V} . This ends the gluing procedure of Richberg.

Now we have a Kähler current $\pi^*\alpha + i\partial\bar{\partial}\varphi''$ defined on W , with $\varphi'' = -\infty$ on all the divisors D_ℓ, E_ℓ which intersect W , and the generic Lelong number of φ'' along each such divisor is strictly positive. On the other hand we also have the function $F \circ \pi$ on \tilde{X} , singular along $\cup_\ell D_\ell \cup \cup_\ell E_\ell \cup \tilde{V}$, which satisfies $\pi^*\alpha + i\partial\bar{\partial}\tilde{F} \geq \gamma\pi^*\omega$ for some small $\gamma > 0$. Hence

$$\pi^*\alpha + i\partial\bar{\partial}(F \circ \pi + \gamma\delta \log |s_E|_{h_E}^2) \geq \gamma\tilde{\omega},$$

is a Kähler current. Let $\tilde{F} = F \circ \pi + \gamma\delta \log |s_E|_{h_E}^2$. Since the class $[\pi^*\alpha]$ is nef, for any small $\nu > 0$ there exists a smooth function ρ_ν on \tilde{X} such that

$$\alpha + i\partial\bar{\partial}\rho_\nu \geq -\nu\gamma\tilde{\omega},$$

holds on \tilde{X} , and we can normalize ρ_ν so that it is strictly positive. Then we have

$$\pi^*\alpha + i\partial\bar{\partial}(\nu\tilde{F} + (1-\nu)\rho_\nu) \geq \nu^2\gamma\omega.$$

Let us shrink W slightly, so that φ'' is defined on an open set containing W in its interior. Pick any point $x \in \pi^{-1}(G) \cap \partial W$, fix a neighborhood U of x with $U \cap \tilde{V} = \emptyset$ and with local coordinates as before, so that on U we can write

$$\begin{aligned} \varphi'' &= c_1 \log \left(\prod_k |z_{i_k}|^{2\alpha_{i_k}} \right) + g_1, \\ \tilde{F} &= c_2 \log \left(\prod_k |z_{i_k}|^{2\beta_{i_k}} \right) + g_2, \end{aligned}$$

where $c_1, c_2, \alpha_{i_k}, \beta_{i_k}$ are constants, g_1, g_2 are continuous functions. Then, since the polar locus of \tilde{F} equals $\pi^{-1}(G) \cup \tilde{V}$ while the polar locus of φ'' equals $\pi^{-1}(G)$, we must have $\alpha_{i_k} > 0$ iff $\beta_{i_k} > 0$. Therefore, if $\nu > 0$ is sufficiently small, we will have that $\nu\tilde{F} + (1-\nu)\rho_\nu > \varphi''$ on a possibly smaller open neighborhood of x .

Repeating this argument on a finite covering of $\pi^{-1}(G) \cap \partial W$, we can find $\nu > 0$ and a neighborhood Z of $\pi^{-1}(G)$ such that $\nu\tilde{F} + (1-\nu)\rho_\nu > \varphi''$ on $Z \cap \partial W$. Since $\tilde{F} > -\infty$ on $(\partial W) \setminus Z$, we can choose $A > 0$ large enough such that on $(\partial W) \setminus Z$ we have $\nu\tilde{F} + (1-\nu)\rho_\nu > \varphi'' - A$. Altogether, this means that $\nu\tilde{F} + (1-\nu)\rho_\nu > \varphi'' - A$ holds in a whole neighborhood of ∂W .

Therefore we can finally define

$$\tilde{\Phi} = \begin{cases} \max\{\varphi'', \nu\tilde{F} + (1-\nu)\rho_\nu + A\} & \text{on } W \\ \nu\tilde{F} + (1-\nu)\rho_\nu + A, & \text{on } \tilde{X} \setminus W, \end{cases}$$

which is defined on the whole of \tilde{X} , it has weakly analytic singularities and satisfies $\tilde{T}_1 = \pi^*\alpha + i\partial\bar{\partial}\tilde{\Phi} \geq \varepsilon''\tilde{\omega}$ for some $\varepsilon'' > 0$. Furthermore, since $\tilde{F} = -\infty$ on \tilde{V} , while φ'' is continuous near the generic point of \tilde{V} , we

see that $\tilde{\Phi} = \varphi''$ in a neighborhood of the generic point of \tilde{V} . Therefore, $\tilde{\Phi}|_{\tilde{V}} = \hat{\varphi}$, which is smooth in a neighborhood (in \tilde{V}) of the generic point of \tilde{V} .

The pushforward $\tilde{T} = \pi_* \tilde{T}_1$ is then a Kähler current on X in the class $[\alpha]$ whose restriction to V is smooth in a neighborhood (in V) of the generic point of V . This completes the proof of Theorem 3.2. \square

4. FINITE TIME SINGULARITIES OF THE KÄHLER-RICCI FLOW

In this section we study finite time singularities of the Kähler-Ricci flow, and give the proof of Theorem 1.5.

Let (X, ω_0) be a compact Kähler manifold of dimension n , and consider Kähler metrics $\omega(t)$ evolving under the Kähler-Ricci flow (1.2) with initial metric ω_0 . Suppose that the maximal existence time of the flow is $T < \infty$. Then the class $[\alpha] = [\omega_0] - Tc_1(X)$ is nef, but not Kähler. Moreover, assume that $\int_X \alpha^n > 0$, so the volume of $(X, \omega(t))$ does not go to zero as $t \rightarrow T^-$. Let α be a smooth representative of this limiting class, and set

$$\begin{aligned}\hat{\omega}_t &= \frac{1}{T} ((T-t)\omega_0 + t\alpha) \in [\omega_0] - tc_1(X) \\ \chi &= \frac{\partial}{\partial t} \hat{\omega}_t = \frac{1}{T} (\alpha - \omega_0) \in -c_1(X).\end{aligned}$$

Let Ω be a volume form so that $i\partial\bar{\partial} \log \Omega = \chi$, and $\int_X \Omega = \int_X \omega_0^n$. Then the Kähler-Ricci flow can be written as the parabolic complex Monge-Ampère equation

$$\frac{\partial}{\partial t} \varphi = \log \frac{(\hat{\omega}_t + i\partial\bar{\partial}\varphi)^n}{\Omega}, \quad \hat{\omega}_t + i\partial\bar{\partial}\varphi > 0, \quad \varphi(0) = 0,$$

so that

$$\omega(t) = \hat{\omega}_t + i\partial\bar{\partial}\varphi,$$

solves (1.2). Since the limiting class is nef and has positive self-intersection, Theorem 2.2 shows that there exists a Kähler current

$$(4.1) \quad R = \alpha + i\partial\bar{\partial}\psi \geq \varepsilon\omega_0,$$

which has analytic singularities, and such that

$$\{\psi = -\infty\} = E_+(R) = E_{nK}(\alpha).$$

By subtracting a constant to ψ , we can assume that $\sup_X \psi \leq 0$.

Theorem 4.1. *There is a closed positive real (1,1) current ω_T on X in the class $[\alpha]$, which is smooth precisely away from $E_{nK}(\alpha)$ and has everywhere vanishing Lelong numbers, such that as $t \rightarrow T^-$ we have that $\omega(t)$ converges to ω_T in $C_{\text{loc}}^\infty(X \setminus E_{nK}(\alpha))$ as well as currents on X .*

Theorem 1.5 follows easily from this result.

Proof of Theorem 1.5. Theorem 4.1 says that no singularities develop on $X \setminus E_{nK}(\alpha)$. Thanks to the main Theorem 1.1, $E_{nK}(\alpha) = \text{Null}(\alpha)$. Finally, we show that the metrics $\omega(t)$ must develop singularities everywhere along $\text{Null}(\alpha)$. Indeed, assume that there is a point $x \in \text{Null}(\alpha)$ with an open neighborhood U where the metrics $\omega(t)$ converge smoothly to a limit Kähler metric ω_U on U . Since $x \in \text{Null}(\alpha)$, there is an irreducible k -dimensional analytic subvariety $V \subset X$ with $x \in V$ and $\int_V \alpha^k = 0$. Then

$$\int_V \omega(t)^k \geq \int_{V \cap U} \omega(t)^k \rightarrow \int_{V \cap U} \omega_U^k > 0,$$

but at the same time $\int_V \omega(t)^k \rightarrow \int_V \alpha^k = 0$, a contradiction. \square

The same proof shows that if $T < \infty$ is a finite time singularity with $\int_X \alpha^n = 0$, then singularities develop on the whole of X .

As a side remark, we have the following:

Proposition 4.2. *Let (X^n, ω_0) be a compact Kähler manifold with nonnegative Kodaira dimension. Then if the Kähler-Ricci flow (1.2) develops a singularity at a finite time T , we must necessarily have that*

$$\int_X (\omega_0 - Tc_1(X))^n > 0,$$

and Theorem 1.5 then applies.

Proof. Indeed, nonnegative Kodaira dimension means that $H^0(X, \ell K_X) \neq 0$ for some $\ell \geq 1$. Since $c_1(K_X) = -c_1(X)$, this implies that the class $-Tc_1(X)$ is pseudoeffective (i.e. it contains a closed positive current S), and therefore the class $[\alpha] = [\omega_0] - Tc_1(X)$ is big since it contains the Kähler current $S + \omega_0$. Then [5, Theorem 4.7] shows that the volume $v(\alpha)$ is strictly positive, and since $[\alpha]$ is also nef we can apply [5, Theorem 4.1] and conclude that $\int_X \alpha^n = v(\alpha) > 0$. \square

Let us also remark that when $n = 2$ if $[\alpha] = \omega_0 - Tc_1(X)$ is the limiting class as above, then $\text{Null}(\alpha)$ is a disjoint union of finitely many (-1) -curves (see [48, Theorem 8.3]) which can be contracted to give a new Kähler surface Y and $[\alpha]$ is the pullback of a Kähler class on Y . The Kähler-Ricci flow can then be restarted on Y , and the whole process is continuous in the Gromov-Hausdorff topology thanks to work of Song-Weinkove [49].

Now we start with the proof of Theorem 4.1. We first need several lemmas.

Lemma 4.3. *There is a constant $C > 0$ such that on $X \times [0, T)$ we have*

- (i) $\dot{\varphi}(t) \leq C$,
- (ii) $\omega^n \leq C\Omega$,
- (iii) $\varphi(t) \leq C$.

The proof is extremely easy (see [48, Lemma 7.1]). Note that (i) is equivalent to (ii), and that integrating (i) from 0 to T gives (iii).

Lemma 4.4. *There is a constant $C > 0$ such that on $X \times [0, T)$ we have*

- (i) $\dot{\varphi}(t) \geq C\psi - C$,
- (ii) $\omega^n \geq C^{-1}e^{C\psi}\Omega$.

Proof. It is enough to prove (i), since (ii) is clearly equivalent to (i).

To prove (i), note that

$$\hat{\omega}_t + i\partial\bar{\partial}\psi = \frac{1}{T} \left((T-t)(\omega_0 + i\partial\bar{\partial}\psi) + t(\alpha + i\partial\bar{\partial}\psi) \right).$$

Now we have

$$\omega_0 + i\partial\bar{\partial}\psi \geq \omega_0 - \alpha \geq -C\omega_0, \quad \alpha + i\partial\bar{\partial}\psi = R \geq \varepsilon\omega_0,$$

where these inequalities are in the sense of currents. Therefore, there is a small $\delta > 0$ such that for $t \in [T - \delta, T)$ we have

$$(4.2) \quad \hat{\omega}_t + i\partial\bar{\partial}\psi \geq \frac{\varepsilon}{2}\omega_0,$$

again as currents. Let

$$Q = \dot{\varphi} + A\varphi - A\psi + Bt,$$

where A and B are large constants to be fixed soon. Our goal is to show that $Q \geq -C$ on $X \times [0, T)$, since then Lemma 4.3 (iii) implies (i).

We fix the constant A large enough so that

$$(A-1)(\hat{\omega}_t + i\partial\bar{\partial}\psi) + \chi \geq \omega_0,$$

holds on $[T - \delta, T)$. To fix the value of B , we first use the arithmetic-geometric mean inequality to get on $[T - \delta, T)$

$$(4.3) \quad \frac{\varepsilon}{2}\text{tr}_\omega\omega_0 \geq C_0^{-1} \left(\frac{\Omega}{\omega^n} \right)^{1/n},$$

where C_0 is a uniform constant. But since the function $y \mapsto A \log y - C_0^{-1}y^{1/n}$ is bounded above for $y > 0$, we have that

$$(4.4) \quad C_0^{-1} \left(\frac{\Omega}{\omega^n} \right)^{1/n} + A \log \frac{\omega^n}{\Omega} \geq -C_1,$$

for another uniform constant C_1 . We then set $B = C_1 + An$.

From the definition of Q we see that $Q \geq -C$ holds on $X \times [0, T - \delta]$, for a uniform constant C . Therefore, to prove that $Q \geq -C$ on $X \times [0, T)$, it suffices to show that given any $T - \delta < T' < T$, the minimum of Q on $X \times [0, T']$ is always achieved on $[0, T - \delta]$. Let then (x, t) be a point in $X \times [T - \delta, T']$ where Q achieves a minimum on $X \times [0, T']$. Since Q approaches $+\infty$ along $E_{nK}(\alpha)$, we conclude that $x \notin E_{nK}(\alpha)$. At (x, t) we

have

$$\begin{aligned}
0 &\geq \left(\frac{\partial}{\partial t} - \Delta \right) Q = \text{tr}_\omega \chi + A\dot{\varphi} - An + A\text{tr}_\omega(\hat{\omega}_t + i\partial\bar{\partial}\psi) + B \\
&= \text{tr}_\omega(\hat{\omega}_t + i\partial\bar{\partial}\psi) + \text{tr}_\omega((A-1)(\hat{\omega}_t + i\partial\bar{\partial}\psi) + \chi) \\
&\quad + A \log \frac{\omega^n}{\Omega} - An + B \\
&\geq \frac{\varepsilon}{2} \text{tr}_\omega \omega_0 + A \log \frac{\omega^n}{\Omega} + \text{tr}_\omega \omega_0 - An + B,
\end{aligned}$$

where $\Delta = \Delta_{\omega(t)}$ is the Laplacian of the moving metric. Thanks to (4.3), (4.4), we conclude that

$$0 \geq \left(\frac{\partial}{\partial t} - \Delta \right) Q \geq \text{tr}_\omega \omega_0 - C_1 - An + B = \text{tr}_\omega \omega_0 > 0,$$

a contradiction. \square

Integrating Lemma 4.4 (i) from 0 to T gives the bound $\varphi(t) \geq C\psi - C$. We now sharpen this:

Lemma 4.5. *There is a constant $C > 0$ such that on $X \times [0, T)$ we have*

$$(4.5) \quad \varphi(t) \geq \psi - C.$$

Proof. Let $Q = \varphi - \psi + At$, for A a large constant to be determined. We need to show that $Q \geq -C$ on $X \times [0, T)$. Clearly, $Q \geq -C$ holds on $X \times [0, T - \delta]$ (where δ is as in Lemma 4.4), so we fix $T - \delta < T' < T$ and suppose that Q achieves its minimum on $X \times [0, T']$ at (x, t) with $t \in [T - \delta, T']$. Since ψ approaches $-\infty$ along $E_{nK}(\alpha)$, we have that $x \in X \setminus E_{nK}(\alpha)$. At (x, t) we then have

$$\begin{aligned}
0 &\geq \frac{\partial Q}{\partial t} = \log \frac{(\hat{\omega}_t + i\partial\bar{\partial}\psi + i\partial\bar{\partial}Q)^n}{\Omega} + A \\
&\geq \log \frac{(\hat{\omega}_t + i\partial\bar{\partial}\psi)^n}{\Omega} + A \\
&\geq \log \frac{(\varepsilon\omega_0/2)^n}{\Omega} + A \geq -C + A,
\end{aligned}$$

using (4.2). If we choose $A > C$, this gives a contradiction. \square

Lemma 4.6. *There is a constant $C > 0$ such that on $X \times [0, T)$ we have*

$$(4.6) \quad \text{tr}_{\omega_0} \omega \leq Ce^{-C\psi}.$$

Proof. Let

$$Q = \log \text{tr}_{\omega_0} \omega - A\varphi + A\psi,$$

where A is a large constant to be determined soon. Thanks to Lemma 4.3 (iii), we will be done if we show that $Q \leq C$ on $X \times [0, T)$. Again, from the definition of Q we see that $Q \leq C$ holds on $X \times [0, T - \delta]$, for a uniform constant C . Fix then any $T - \delta < T' < T$ and suppose that Q achieves its

maximum on $X \times [0, T']$ at (x, t) with $t \in [T - \delta, T']$. Then at (x, t) we can use (4.2) to estimate

$$\begin{aligned} 0 &\leq \left(\frac{\partial}{\partial t} - \Delta \right) Q \leq C \operatorname{tr}_\omega \omega_0 + A \log \frac{\Omega}{\omega^n} + An - A \operatorname{tr}_\omega (\dot{\omega}_t + i\partial\bar{\partial}\psi) \\ &\leq (C - A\varepsilon/2) \operatorname{tr}_\omega \omega_0 + A \log \frac{\Omega}{\omega^n} + An. \end{aligned}$$

Hence if we choose A large so that $C - A\varepsilon/2 \leq -1$, we see that at (x, t) we have

$$\operatorname{tr}_\omega \omega_0 \leq C \log \frac{\Omega}{\omega^n} + C.$$

Hence at the maximum of Q ,

$$\operatorname{tr}_{\omega_0} \omega \leq \frac{1}{(n-1)!} (\operatorname{tr}_\omega \omega_0)^{n-1} \frac{\omega^n}{\omega_0^n} \leq C \frac{\omega^n}{\Omega} \left(\log \frac{\Omega}{\omega^n} \right)^{n-1} + C \leq C,$$

because we know that $\frac{\omega^n}{\Omega} \leq C$ (Lemma 4.3 (ii)) and $x \mapsto x |\log x|^{n-1}$ is bounded above for x close to zero. Thanks to Lemma 4.5, this implies that Q is bounded from above at its maximum, and we are done. \square

Proof of Theorem 4.1. Fix any compact set $K \subset X \setminus E_{nK}(\alpha)$. Thanks to Lemmas 4.4 (ii) and 4.6, there is a constant $C_K > 0$ such that

$$C_K^{-1} \omega_0 \leq \omega(t) \leq C_K \omega_0 \quad \text{on } K \times [0, T].$$

Applying the local higher order estimates of Sherman-Weinkove [41], we get uniform C^∞ estimates for $\omega(t)$ on compact subsets of $X \setminus E_{nK}(\alpha)$. This implies that given every compact set K there exists a constant C_K such that

$$\frac{\partial}{\partial t} \omega = -\operatorname{Ric}(\omega) \leq C_K \omega,$$

which implies that $e^{-C_K t} \omega(t)$ is decreasing in t as well as being bounded from below. This implies that $\omega(t)$ has a limit as $t \rightarrow T$, and since we have uniform estimates away from $E_{nK}(\alpha)$, we see that $\omega(t)$ converges in C^∞ on compact sets to a smooth Kähler metric ω_T on $X \setminus E_{nK}(\alpha)$. Furthermore, by weak compactness of currents, it follows easily that ω_T extends as a closed positive current on X and $\omega(t)$ converges to ω_T as currents on X .

Finally we show that the limiting current ω_T on X has vanishing Lelong numbers everywhere. Indeed, since the class $[\alpha]$ is nef, for any $\nu > 0$ there is a smooth function ρ_ν on X such that $\alpha + i\partial\bar{\partial}\rho_\nu \geq -\nu\varepsilon\omega_0$. Then

$$\alpha + i\partial\bar{\partial}(\nu\psi + (1-\nu)\rho_\nu) \geq \nu^2\varepsilon\omega_0.$$

Then the same argument as Lemma 4.5, replacing ψ with $\nu\psi + (1-\nu)\rho_\nu$, shows that $\varphi(t) \geq \nu\psi - C_\nu$ on $X \times [0, T]$. This implies that the Lelong numbers of ω_T are at most $C\nu$, for any $\nu > 0$, and we are done. \square

Related estimates can be found in [8] under slightly different assumptions.

We now remark that the set of points where the Kähler-Ricci flow develops singularities can also be characterized using the curvature of the evolving metrics. We are grateful to Zhou Zhang for pointing out this result to us.

Following Enders-Müller-Topping [23], we will call Σ the complement of the set of points $x \in X$ such that there exists an open neighborhood U of x and a constant $C > 0$ with $|\text{Rm}(t)|_{g(t)} \leq C$ on $U \times [0, T)$. Also, call Σ' the complement of the set of points $x \in X$ such that there exists an open neighborhood U of x and a constant $C > 0$ with $R(t) \leq C$ on $U \times [0, T)$, where $R(t)$ is the scalar curvature of $\omega(t)$.

Theorem 4.7. *Let the setup be the same as in Theorem 1.5. Then*

$$\text{Null}(\alpha) = \Sigma = \Sigma'.$$

Proof. The inclusion $\Sigma' \subset \Sigma$ is trivial, and thanks to Theorem 4.1 we have the inclusion $\Sigma \subset \text{Null}(\alpha)$. It then suffices to prove that if $R(t) \leq C$ on $U \times [0, T)$, then on a smaller open neighborhood U' of x we have that $\omega(t)$ converge smoothly to a Kähler metric $\omega_{U'}$ as $t \rightarrow T^-$.

The proof is contained in the work of Zhang [61, 60] (see also the exposition [48, Theorem 7.6]). In general, $R \geq -C$ along the flow. From the flow equation

$$\dot{\varphi} = \log \frac{\omega^n}{\Omega}, \quad \frac{\partial}{\partial t} \dot{\varphi} = -R,$$

so $\frac{\partial}{\partial t} \dot{\varphi} \leq C$ on $M \times [0, T)$. Integrating from 0 to T we get $\dot{\varphi} \leq C$ and $\varphi \leq C$ on $M \times [0, T)$. Furthermore, on $U \times [0, T)$ we have $|\frac{\partial}{\partial t} \dot{\varphi}| \leq C$ and hence $|\dot{\varphi}| + |\varphi| \leq C$. The quantity $H = t\dot{\varphi} - \varphi - nt$ is therefore bounded on $U \times [0, T)$, and satisfies

$$\left(\frac{\partial}{\partial t} - \Delta \right) H = -\text{tr}_{\omega} \omega_0.$$

On the other hand a ‘‘Schwarz-Lemma’’-type calculation (see for example [47, 48]) gives

$$\left(\frac{\partial}{\partial t} - \Delta \right) \log \text{tr}_{\omega} \omega_0 \leq C \text{tr}_{\omega} \omega_0.$$

Therefore if we let $Q = \log \text{tr}_{\omega} \omega_0 + AH$, then

$$\left(\frac{\partial}{\partial t} - \Delta \right) Q \leq -\text{tr}_{\omega} \omega_0 \leq 0,$$

provided A is large enough. Therefore given any $T' < T$, the maximum of Q on $M \times [0, T')$ is achieved at $t = 0$, and hence

$$\log \text{tr}_{\omega} \omega_0 + A(t\dot{\varphi} - \varphi - nt) \leq C.$$

This means that on $M \times [0, T)$ we have

$$\text{tr}_{\omega} \omega_0 \leq C e^{-At\dot{\varphi} + A\varphi + Ant} \leq C e^{-At\dot{\varphi}}.$$

Since $\text{tr}_\omega \omega_0 / \Omega = e^{\dot{\phi}}$, and

$$\text{tr}_{\omega_0} \omega \leq C(\text{tr}_\omega \omega_0)^{n-1} \frac{\omega^n}{\Omega} \leq C e^{-((n-1)At-1)\dot{\phi}}.$$

On $U \times [0, T)$ we therefore have $C^{-1}\omega_0 \leq \omega \leq C\omega_0$. The local higher order estimates of Sherman-Weinkove [41], then give us C^∞ estimates for ω on $U' \times [0, T)$ for a slightly smaller open neighborhood U' of x , and from here we conclude as in the proof of Theorem 4.1. \square

5. DEGENERATIONS OF RICCI-FLAT KÄHLER METRICS

In this section we study degenerations of Ricci-flat Kähler metrics on Calabi-Yau manifolds, and prove Theorem 1.6.

Let X be a Calabi-Yau manifold, which is a compact Kähler manifold with $c_1(X) = 0$ in $H^2(X, \mathbb{R})$. Let $\mathcal{K} \subset H^{1,1}(X, \mathbb{R})$ be the Kähler cone of X . Then Yau's theorem [58] says that for every Kähler class $[\alpha] \in \mathcal{K}$ there exists a unique Kähler metric $\omega \in [\alpha]$ with $\text{Ric}(\omega) = 0$. Let α be a smooth closed real $(1, 1)$ form on X with $[\alpha] \in \partial\mathcal{K}$ nef and $\int_X \alpha^n > 0$. Let $[\alpha_t] : [0, 1] \rightarrow \bar{\mathcal{K}}$ be a continuous path of $(1, 1)$ classes with $[\alpha_t] \in \mathcal{K}$ for $t > 0$ and with $[\alpha_0] = [\alpha]$. For $0 < t \leq 1$ write ω_t for the unique Ricci-flat Kähler metric in the class $[\alpha_t]$.

Proof of Theorem 1.6. To start, let us see that we can choose a continuously varying path of reference $(1, 1)$ forms α_t cohomologous to $[\alpha_t]$ with $\alpha_0 = \alpha$. Indeed, it is easy to find $(1, 1)$ forms $\tilde{\alpha}_t$ representing the classes $[\alpha_t]$ and varying continuously in t . Since $\tilde{\alpha}_0$ and α are cohomologous, we may write $\tilde{\alpha}_0 = \alpha + i\partial\bar{\partial}\eta$, and then let $\alpha_t = \tilde{\alpha}_t - i\partial\bar{\partial}\eta$. In general the forms α_t do not satisfy any positivity property. For convenience we will assume that the metric ω is Ricci-flat. Recall the following construction from [7]: for any $x \in X$ let

$$V_\alpha(x) = \sup\{\varphi(x) \mid \varphi \text{ is } \alpha\text{-PSH, } \sup_X \varphi \leq 0\}$$

be the extremal function associated to the form α . Then $\alpha + i\partial\bar{\partial}V_\alpha \geq 0$ and V_α has minimal singularities among all α -PSH functions. Similarly, for $0 \leq t \leq 1$ we let $V_t = V_{\alpha_t}$, so that $V_0 = V_\alpha$. Then the function V_α is continuous on $X \setminus E_{nK}(\alpha)$, while V_t is continuous everywhere on X .

Using the theorem of Demailly-Păun together with Demailly's regularization, we see that there exists an α -PSH function ψ with analytic singularities, with $\sup_X \psi = 0$ and with $\alpha + i\partial\bar{\partial}\psi \geq \delta\omega$ on X , for some $\delta > 0$. Furthermore, thanks to Theorem 2.2, the function ψ can be chosen to be smooth on $X \setminus E_{nK}(\alpha)$ (and singular on $E_{nK}(\alpha)$). In general V_α has strictly weaker singularities than ψ , i.e. it need not be singular on $E_{nK}(\alpha)$. For example if Y is a compact Kähler surface, $\pi : X \rightarrow Y$ is the blowup of a point, $E \subset X$ is the exceptional divisor, and $\alpha = \pi^*\omega$ for a Kähler metric ω on Y , then V_α vanishes identically while ψ is singular along E .

Note that for t sufficiently small, we will have

$$(5.1) \quad \alpha_t + i\partial\bar{\partial}\psi = \alpha + i\partial\bar{\partial}\psi + (\alpha_t - \alpha) \geq \delta\omega/2.$$

Therefore, from the definition of V_t , we have that

$$(5.2) \quad V_t \geq \psi,$$

holds on X for all t sufficiently small.

For $t > 0$ we can write $\omega_t = \alpha_t + i\partial\bar{\partial}\varphi_t$, with $\sup_X \varphi_t = 0$. Then the functions φ_t solve the complex Monge-Ampère equation

$$(5.3) \quad \omega_t^n = (\alpha_t + i\partial\bar{\partial}\varphi_t)^n = c_t\omega^n,$$

where the constant c_t equals $\int_X \alpha_t^n / \int_X \omega^n$ (and is therefore bounded uniformly away from zero and infinity). The crucial result is then the following estimate of Boucksom-Eyssidieux-Guedj-Zeriahi [7, Theorem 4.1, Remark 4.5]

$$(5.4) \quad \varphi_t \geq V_t - C,$$

for a uniform constant C independent of t . This is generalization of a seminal result of Kołodziej [29]. Thanks to (5.2), we conclude that

$$(5.5) \quad \varphi_t \geq \psi - C,$$

for all t sufficiently small. Furthermore [7, Theorem 4.1] also gives us a unique α -PSH function φ_0 with $\sup_X \varphi_0 = 0$ solving

$$\langle \alpha + i\partial\bar{\partial}\varphi_0 \rangle^n = c_0\omega^n,$$

on $X \setminus E_{nK}(\alpha)$, and with $\varphi_0 \geq V_\alpha - C$ (here $\langle \cdot \rangle^n$ is the non-pluripolar product defined in [7]). Now the same simple argument as [7, Lemma 5.3] shows that as $t \rightarrow 0$ the functions φ_t converge to φ_0 in $L^1(X)$. To get higher order estimates of φ_t on $X \setminus E_{nK}(\alpha)$ we proceed as follows. Let Δ_t be the Laplacian of ω_t , A a large constant, and $Q_t = \log \operatorname{tr}_\omega \omega_t - A\varphi_t + A\psi$. A standard calculation (see e.g. [58]) gives

$$\Delta_t Q_t \geq -C \operatorname{tr}_{\omega_t} \omega - C - An + A \operatorname{tr}_{\omega_t} (\alpha_t + i\partial\bar{\partial}\psi).$$

We can assume that t is sufficiently small so that (5.1) holds, and choose A sufficiently large so that

$$\Delta_t Q_t \geq \operatorname{tr}_{\omega_t} \omega - C.$$

Note that for any $t > 0$ the function Q_t approaches $-\infty$ along $E_{nK}(\alpha)$, so its maximum is achieved on its complement. For any sufficiently small t , let $x \in X \setminus E_{nK}(\alpha)$ be a point where the maximum of Q_t is achieved. Then $\operatorname{tr}_{\omega_t} \omega(x) \leq C$, and using (5.3) we see that $\operatorname{tr}_\omega \omega_t(x) \leq C$ and

$$Q_t \leq \log C - A\varphi_t(x) + A\psi(x) \leq C,$$

thanks to (5.5). Therefore $Q_t \leq C$ holds on X for all small $t > 0$. Since $\sup_X \varphi_t = 0$, we have proved that

$$\operatorname{tr}_\omega \omega_t \leq C e^{-A\psi}.$$

This implies that ω_t is uniformly equivalent to ω on any compact set $K \subset X \setminus E_{nK}(\alpha)$, independent of t . The higher order estimates (either the Calabi C^3 estimate [58] or the Evans-Krylov theory [43]) are local, and so we get uniform C_{loc}^∞ estimates for φ_t on $X \setminus E_{nK}(\alpha)$. It follows that as $t \rightarrow 0$ the Ricci-flat metrics ω_t converge to $\omega_0 = \alpha + i\partial\bar{\partial}\varphi_0$ in $C_{\text{loc}}^\infty(X \setminus E_{nK}(\alpha))$. Note that this gives a slightly simpler proof of the fact that $\varphi_0 \in C^\infty(X \setminus E_{nK}(\alpha))$ than the one given in [7, Theorem 4.1]

Next, the metrics ω_t have a uniform upper bound on their diameter thanks to [53, 59]. Once we show the Gromov-Hausdorff convergence statement, it will also follow that $(X \setminus E_{nK}(\alpha), \omega_0)$ has finite diameter, and hence is incomplete. The Gromov-Hausdorff convergence is proved exactly along the lines of [38, Lemma 5.1], who dealt with the case when X is projective and $[\alpha]$ is rational. The key point is that thanks to our main Theorem 1.1, $E_{nK}(\alpha) = \text{Null}(\alpha)$, so the exact same argument as [38, Lemma 5.1] applies, with the subvariety E there replaced by $\text{Null}(\alpha)$. \square

Finally, let us remark that the same proof as in Theorem 2.5 or Theorem 1.5 shows that the metrics ω_t cannot converge smoothly to a Kähler metric near any point of $\text{Null}(\alpha)$.

REFERENCES

- [1] Barth, W.P., Hulek, K., Peters, C.A.M., Van de Ven, A. *Compact complex surfaces. Second edition*, Springer, 2004.
- [2] Benito, A., Encinas, S., Villamayor U., O.E. *Some natural properties of constructive resolution of singularities*, Asian J. Math. **15** (2011), no. 2, 141–191.
- [3] R.J. Berman, J.-P. Demailly *Regularity of plurisubharmonic upper envelopes in big cohomology classes*, in *Perspectives in analysis, geometry, and topology*, 39–66, Progr. Math., 296, Birkhäuser/Springer, New York, 2012.
- [4] S. Boucksom *Cônes positifs des variétés complexes compactes*, Ph.D. Thesis, Institut Fourier Grenoble, 2002.
- [5] S. Boucksom *On the volume of a line bundle*, Internat. J. Math. **13** (2002), no. 10, 1043–1063.
- [6] S. Boucksom *Divisorial Zariski decompositions on compact complex manifolds*, Ann. Sci. École Norm. Sup. (4) **37** (2004), no. 1, 45–76.
- [7] S. Boucksom, P. Eyssidieux, V. Guedj, A. Zeriahi *Monge-Ampère equations in big cohomology classes*, Acta Math. **205** (2010), no. 2, 199–262.
- [8] S. Boucksom, V. Guedj *Kähler-Ricci flows on singular varieties*, in *An introduction to the Kähler-Ricci flow*, Lecture Notes in Math., 2086, Springer, Heidelberg, 2013.
- [9] S. Cacciola, A.F. Lopez *Nakamaye’s theorem on log canonical pairs*, arXiv:1303.7156.
- [10] S. Cantat, A. Zeghib *Holomorphic actions, Kummer examples, and Zimmer program*, Ann. Sci. École Norm. Sup. (4) **45** (2012), no. 3, 447–489.
- [11] H.-D. Cao *Deformation of Kähler metrics to Kähler-Einstein metrics on compact Kähler manifolds*, Invent. Math. **81** (1985), no. 2, 359–372.
- [12] P. Cascini, J. McKernan, M. Mustatǎ *The augmented base locus in positive characteristic*, arXiv:1111.3236.
- [13] T.C. Collins, A. Greenleaf, M. Pramanik *A multi-dimensional resolution of singularities with applications to analysis*, to appear in Amer. J. Math. 2013.
- [14] M. Coltoiu *Traces of Runge domains on analytic subsets*, Math. Ann. **290** (1991), 545–548.

- [15] D. Coman, V. Guedj, A. Zeriahi *Extension of plurisubharmonic functions with growth control*, J. Reine Angew. Math. **676** (2013), 33–49.
- [16] J.-P. Demailly *Singular Hermitian metrics on positive line bundles*, in *Complex algebraic varieties (Bayreuth, 1990)*, 87–104, Lecture Notes in Math., 1507, Springer, Berlin, 1992.
- [17] J.-P. Demailly *Regularization of closed positive currents and intersection theory*, J. Algebraic Geom. **1** (1992), no. 3, 361–409.
- [18] J.-P. Demailly *Complex Analytic and Differential Geometry*, available on the author’s webpage.
- [19] J.-P. Demailly, S. Dinew, V. Guedj, P.H. Hiep, S. Kołodziej, A. Zeriahi *Hölder continuous solutions to Monge-Ampère equations*, arXiv:1112.1388.
- [20] J.-P. Demailly, M. Păun *Numerical characterization of the Kähler cone of a compact Kähler manifold*, Ann. of Math., **159** (2004), no. 3, 1247–1274.
- [21] L. Ein, R. Lazarsfeld, M. Mustață, M. Nakamaye, M. Popa *Asymptotic invariants of base loci*, Ann. Inst. Fourier (Grenoble) **56** (2006), no. 6, 1701–734.
- [22] L. Ein, R. Lazarsfeld, M. Mustață, M. Nakamaye, M. Popa *Restricted volumes and base loci of linear series*, Amer. J. Math. **131** (2009), no. 3, 607–651.
- [23] J. Enders, R., Müller, P.M., Topping *On type-I singularities in Ricci flow*, Comm. Anal. Geom. **19** (2011), no. 5, 905–922.
- [24] P. Eyssidieux, V. Guedj, A. Zeriahi *Singular Kähler-Einstein metrics*, J. Amer. Math. Soc. **22** (2009), 607–639.
- [25] M. Feldman, T. Ilmanen, D. Knopf *Rotationally symmetric shrinking and expanding gradient Kähler-Ricci solitons*, J. Differential Geom. **65** (2003), no. 2, 169–209.
- [26] A. Fujiki *Closedness of the Douady spaces of compact Kähler spaces*, Publ. Res. Inst. Math. Sci. **14** (1978/79), no. 1, 1–52.
- [27] P. Griffiths, J. Harris *Principles of algebraic geometry*, Pure and Applied Mathematics. Wiley-Interscience, New York, 1978.
- [28] T. Hisamoto *Remarks on L^2 -jet extension and extension of singular Hermitian metric with semi positive curvature*, arXiv:1205.1953.
- [29] S. Kołodziej *The complex Monge-Ampère equation*, Acta Math. **180** (1998), 69–117.
- [30] G. La Nave, G. Tian *Soliton-type metrics and Kähler-Ricci flow on symplectic quotients*, arXiv:0903.2413.
- [31] R. Lazarsfeld *Positivity in algebraic geometry I & II*, Springer-Verlag, Berlin, 2004.
- [32] M. Nakamaye *Stable base loci of linear series*, Math. Ann., **318** (2000), no. 4, 837–847.
- [33] M. Păun *Sur l’effectivité numérique des images inverses de fibrés en droites*, Math. Ann. **310** (1998), no. 3, 411–421.
- [34] D.H. Phong, E.M. Stein, J. Sturm *On the growth and stability of real-analytic functions*, Amer. J. Math. **121** (1999), no. 3, 519–554.
- [35] D.H. Phong, J. Sturm *Algebraic estimates, stability of local zeta functions, and uniform estimates for distribution functions*, Ann. of Math. (2) **152** (2000), no. 1, 277–329.
- [36] D.H. Phong, J. Sturm *On the algebraic constructibility of varieties of integrable rational functions on \mathbb{C}^n* , Math. Ann. **323** (2002), no. 3, 453–484.
- [37] R. Richberg *Stetige streng pseudokonvexe Funktionen*, Math. Ann. **175** (1968), 257–286.
- [38] X. Rong, Y. Zhang *Continuity of extremal transitions and flops for Calabi-Yau manifolds*, J. Differential Geom. **89** (2011), no. 2, 233–269.
- [39] X. Rong, Y. Zhang *Degenerations of Ricci-flat Calabi-Yau manifolds*, to appear in Commun. Contemp. Math. 2013.
- [40] G. Schumacher *Asymptotics of Kähler-Einstein metrics on quasi-projective manifolds and an extension theorem on holomorphic maps*, Math. Ann. **311** (1998), no. 4, 631–645.

- [41] M. Sherman, B. Weinkove *Interior derivative estimates for the Kähler-Ricci flow*, Pacific J. Math. **257** (2012), no. 2, 491–501.
- [42] Y.-T. Siu *Analyticity of sets associated to Lelong numbers and the extension of closed positive currents*, Invent. Math. **27** (1974), 53–156.
- [43] Y.-T. Siu *Lectures on Hermitian-Einstein metrics for stable bundles and Kähler-Einstein metrics*, DMV Seminar, 8, Birkhäuser Verlag, Basel, 1987.
- [44] J. Song *Finite time extinction of the Kähler-Ricci flow*, arXiv:0905.0939.
- [45] J. Song *Ricci flow and birational surgery*, arXiv:1304.2607.
- [46] J. Song, G. Székelyhidi, B. Weinkove *The Kähler-Ricci flow on projective bundles*, Int. Math. Res. Not. **2013**, no. 2, 243–257.
- [47] J. Song, G. Tian *The Kähler-Ricci flow on surfaces of positive Kodaira dimension*, Invent. Math. **170** (2007), no. 3, 609–653.
- [48] J. Song, B. Weinkove *Lecture notes on the Kähler-Ricci flow*, arXiv:1212.3653.
- [49] J. Song, B. Weinkove *Contracting exceptional divisors by the Kähler-Ricci flow*, Duke Math. J. **162** (2013), no. 2, 367–415.
- [50] G. Tian *New results and problems on Kähler-Ricci flow*, in *Géométrie différentielle, physique mathématique, mathématiques et société. II*, Astérisque No. **322** (2008), 71–92.
- [51] G. Tian *Finite-time singularity of Kähler-Ricci flow*, Discrete Contin. Dyn. Syst. **28** (2010), no. 3, 1137–1150.
- [52] G. Tian, Z. Zhang *On the Kähler-Ricci flow on projective manifolds of general type*, Chinese Ann. Math. Ser. B **27** (2006), no. 2, 179–192.
- [53] V. Tosatti *Limits of Calabi-Yau metrics when the Kähler class degenerates*, J. Eur. Math. Soc. (JEMS) **11** (2009), no.4, 755–776.
- [54] V. Tosatti *Degenerations of Calabi-Yau metrics*, in *Geometry and Physics in Cracow*, Acta Phys. Polon. B Proc. Suppl. **4** (2011), no.3, 495–505.
- [55] V. Tosatti *Calabi-Yau manifolds and their degenerations*, Ann. N.Y. Acad. Sci. **1260** (2012), 8–13.
- [56] J. Varouchas *Kähler spaces and proper open morphisms*, Math. Ann. **283** (1989), no. 1, 13–52.
- [57] D. Wu *Higher canonical asymptotics of Kähler-Einstein metrics on quasi-projective manifolds*, Comm. Anal. Geom. **14** (2006), no. 4, 795–845.
- [58] S.-T. Yau *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation, I*, Comm. Pure Appl. Math. **31** (1978), no.3, 339–411.
- [59] Y. Zhang *Convergence of Kähler manifolds and calibrated fibrations*, Ph.D. thesis, Nankai Institute of Mathematics, 2006.
- [60] Z. Zhang *Scalar curvature behavior for finite-time singularity of Kähler-Ricci flow*, Michigan Math. J. **59** (2010), no. 2, 419–433.
- [61] Z. Zhang *General weak limit for Kähler-Ricci flow*, arXiv:1104.2961.
- [62] Z. Zhang *Ricci lower bound for Kähler-Ricci flow*, arXiv:1110.5954.

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, 2990 BROADWAY, NEW YORK, NY 10027

E-mail address: `tcollins@math.columbia.edu`

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, 2033 SHERIDAN ROAD, EVANSTON, IL 60201

E-mail address: `tosatti@math.northwestern.edu`