

A Time-Scale Variational Approach to Inflation, Unemployment and Social Loss*

Monika Dryl¹
monikadryl@ua.pt

Agnieszka B. Malinowska²
a.malinowska@pb.edu.pl

Delfim F. M. Torres¹
delfim@ua.pt

¹CIDMA—Center for Research and Development in Mathematics and Applications,
Department of Mathematics, University of Aveiro, 3810-193 Aveiro, Portugal

²Faculty of Computer Science, Białystok University of Technology,
15-351 Białystok, Poland

Abstract

Both inflation and unemployment inflict social losses. When a tradeoff exists between the two, what would be the best combination of inflation and unemployment? A well known approach in economics to address this question consists to write the social loss as a function of the rate of inflation p and the rate of unemployment u , with different weights, and then, using known relations between p , u , and the expected rate of inflation π , to rewrite the social loss function as a function of π . The answer is achieved by applying the theory of the calculus of variations in order to find an optimal path π that minimizes the total social loss over a given time interval. Economists dealing with this question use a continuous or a discrete variational problem. Here we propose to use a time-scale model, unifying available results in the literature. Moreover, the new formalism allow us to obtain new insights to the classical models when applied to real data of inflation and unemployment.

Keywords: calculus on time scales; calculus of variations; delta derivatives; dynamic model; inflation; unemployment.

2010 Mathematics Subject Classification: 34N05; 49K05; 91B50; 91B62.

1 Introduction

Time-scale calculus is a recent and exciting mathematical theory that unifies two existing approaches to dynamic modelling — difference and differential equations — into a general framework called dynamic models on time scales [3, 9, 14]. Since a more general approach to dynamic modelling, it allows to consider more complex time domains, such as $h\mathbb{Z}$, $q^{\mathbb{N}_0}$ or complex hybrid domains [1].

Both inflation and unemployment inflict social losses. When a Phillips tradeoff exists between the two, what would be the best combination of inflation and unemployment? A well-known approach consists to write the social loss function as a function of the rate of inflation p and the rate of unemployment u , with different weights; then, using relations between p , u and the expected rate of inflation π , to rewrite the social loss function as a function of π ; finally, to apply the theory of the calculus of variations in order to find an optimal path π that minimizes the total

*Part of first author's Ph.D., which is carried out at the University of Aveiro under the Doctoral Programme *Mathematics and Applications* of Universities of Aveiro and Minho.

This is a preprint of a paper whose final and definitive form will appear in *Control and Cybernetics*. Paper submitted 31-Oct-2012; revised 02-April-2013; accepted for publication 17-April-2013.

social loss over a certain time interval $[0, T]$ under study. Economists dealing with this question implement the above approach using both continuous and discrete models [5, 16]. Here we propose a new, more general, time-scale model. We claim that such model describes better the reality.

We compare solutions to three models — the continuous, the discrete, and the time-scale model with $\mathbb{T} = h\mathbb{Z}$ — using real data from the USA over a period of 11 years, from 2000 to 2010. Our results show that the solutions to the classical continuous and discrete models do not approximate well the reality. Therefore, while predicting the future, one cannot base predictions on the two classical models only. The time-scale approach proposed here shows, however, that the classical models are adequate if one uses an appropriate data sampling process. Moreover, the proper times for data collection can be computed from the theory of time scales.

The paper is organized as follows. Section 2 provides all the necessary definitions and results of the delta-calculus on time scales, which will be used throughout the text. This section makes the paper accessible to Economists with no previous contact with the time-scale calculus. In Section 3 we present the economical model under our consideration, in continuous, discrete, and time-scale settings. Section 4 contains our results. Firstly, we derive in Section 4.1 necessary (Theorem 27 and Corollary 30) and sufficient (Theorem 33) optimality conditions for the variational problem that models the economical situation. For the time scale $\mathbb{T} = h\mathbb{Z}$ with appropriate values of h , we obtain an explicit solution for the global minimizer of the total social loss problem (Theorem 34). Secondly, we apply those conditions to the model with real data of inflation [10] and unemployment [17] (Section 4.2). We end with Section 5 of conclusions.

2 Preliminaries

In this section we introduce basic definitions and theorems that will be useful in the sequel. For more on the theory of time scales we refer to the gentle books [3, 4]. For general results on the calculus of variations on time scales we refer the reader to [8, 11, 13] and references therein.

A time scale \mathbb{T} is an arbitrary nonempty closed subset of \mathbb{R} . Let $a, b \in \mathbb{T}$ with $a < b$. We define the interval $[a, b]$ in \mathbb{T} by $[a, b]_{\mathbb{T}} := [a, b] \cap \mathbb{T} = \{t \in \mathbb{T} : a \leq t \leq b\}$.

Definition 1 ([3]). The backward jump operator $\rho : \mathbb{T} \rightarrow \mathbb{T}$ is defined by $\rho(t) := \sup\{s \in \mathbb{T} : s < t\}$ for $t \neq \inf \mathbb{T}$ and $\rho(\inf \mathbb{T}) := \inf \mathbb{T}$ if $\inf \mathbb{T} > -\infty$. The forward jump operator $\sigma : \mathbb{T} \rightarrow \mathbb{T}$ is defined by $\sigma(t) := \inf\{s \in \mathbb{T} : s > t\}$ for $t \neq \sup \mathbb{T}$ and $\sigma(\sup \mathbb{T}) := \sup \mathbb{T}$ if $\sup \mathbb{T} < +\infty$. The backward graininess function $\nu : \mathbb{T} \rightarrow [0, \infty)$ is defined by $\nu(t) := t - \rho(t)$, while the forward graininess function $\mu : \mathbb{T} \rightarrow [0, \infty)$ is defined by $\mu(t) := \sigma(t) - t$.

Example 2. The two classical time scales are \mathbb{R} and \mathbb{Z} , representing the continuous and the purely discrete time, respectively. The other example of interest to the present study is the periodic time scale $h\mathbb{Z}$. It follows from Definition 1 that if $\mathbb{T} = \mathbb{R}$, then $\sigma(t) = t$, $\rho(t) = t$, and $\mu(t) = 0$ for all $t \in \mathbb{T}$; if $\mathbb{T} = h\mathbb{Z}$, then $\sigma(t) = t + h$, $\rho(t) = t - h$, and $\mu(t) = h$ for all $t \in \mathbb{T}$.

A point $t \in \mathbb{T}$ is called *right-dense*, *right-scattered*, *left-dense* or *left-scattered* if $\sigma(t) = t$, $\sigma(t) > t$, $\rho(t) = t$, and $\rho(t) < t$, respectively. We say that t is *isolated* if $\rho(t) < t < \sigma(t)$, that t is *dense* if $\rho(t) = t = \sigma(t)$.

2.1 The delta derivative and the delta integral

We collect here the necessary theorems and properties concerning differentiation and integration on a time scale. To simplify the notation, we define $f^{\sigma}(t) := f(\sigma(t))$. The delta derivative is defined for points in the set

$$\mathbb{T}^{\kappa} := \begin{cases} \mathbb{T} \setminus \{\sup \mathbb{T}\} & \text{if } \rho(\sup \mathbb{T}) < \sup \mathbb{T} < \infty, \\ \mathbb{T} & \text{otherwise.} \end{cases}$$

Definition 3 (Section 1.1 of [3]). We say that a function $f : \mathbb{T} \rightarrow \mathbb{R}$ is Δ -differentiable at $t \in \mathbb{T}^\kappa$ if there is a number $f^\Delta(t)$ such that for all $\varepsilon > 0$ there exists a neighborhood O of t such that

$$|f^\sigma(t) - f(s) - f^\Delta(t)(\sigma(t) - s)| \leq \varepsilon|\sigma(t) - s| \quad \text{for all } s \in O.$$

We call to $f^\Delta(t)$ the Δ -derivative of f at t .

Theorem 4 (Theorem 1.16 of [3]). *Let $f : \mathbb{T} \rightarrow \mathbb{R}$ and $t \in \mathbb{T}^\kappa$. The following holds:*

1. *If f is differentiable at t , then f is continuous at t .*
2. *If f is continuous at t and t is right-scattered, then f is differentiable at t with*

$$f^\Delta(t) = \frac{f^\sigma(t) - f(t)}{\mu(t)}.$$

3. *If t is right-dense, then f is differentiable at t if, and only if, the limit*

$$\lim_{s \rightarrow t} \frac{f(t) - f(s)}{t - s}$$

exists as a finite number. In this case,

$$f^\Delta(t) = \lim_{s \rightarrow t} \frac{f(t) - f(s)}{t - s}.$$

4. *If f is differentiable at t , then $f^\sigma(t) = f(t) + \mu(t)f^\Delta(t)$.*

Example 5. If $\mathbb{T} = \mathbb{R}$, then item 3 of Theorem 4 yields that $f : \mathbb{R} \rightarrow \mathbb{R}$ is delta differentiable at $t \in \mathbb{R}$ if, and only if,

$$f^\Delta(t) = \lim_{s \rightarrow t} \frac{f(t) - f(s)}{t - s}$$

exists, i.e., if, and only if, f is differentiable (in the ordinary sense) at t : $f^\Delta(t) = f'(t)$. If $\mathbb{T} = h\mathbb{Z}$, then point 2 of Theorem 4 yields that $f : \mathbb{Z} \rightarrow \mathbb{R}$ is delta differentiable at $t \in h\mathbb{Z}$ if, and only if,

$$f^\Delta(t) = \frac{f(\sigma(t)) - f(t)}{\mu(t)} = \frac{f(t+h) - f(t)}{h}. \quad (1)$$

In the particular case $h = 1$, $f^\Delta(t) = \Delta f(t)$, where Δ is the usual forward difference operator.

Theorem 6 (Theorem 1.20 of [3]). *Assume $f, g : \mathbb{T} \rightarrow \mathbb{R}$ are Δ -differentiable at $t \in \mathbb{T}^\kappa$. Then,*

1. *The sum $f + g : \mathbb{T} \rightarrow \mathbb{R}$ is Δ -differentiable at t with $(f + g)^\Delta(t) = f^\Delta(t) + g^\Delta(t)$.*
2. *For any constant α , $\alpha f : \mathbb{T} \rightarrow \mathbb{R}$ is Δ -differentiable at t with $(\alpha f)^\Delta(t) = \alpha f^\Delta(t)$.*
3. *The product $fg : \mathbb{T} \rightarrow \mathbb{R}$ is Δ -differentiable at t with*

$$(fg)^\Delta(t) = f^\Delta(t)g(t) + f^\sigma(t)g^\Delta(t) = f(t)g^\Delta(t) + f^\Delta(t)g^\sigma(t).$$

4. *If $g(t)g^\sigma(t) \neq 0$, then f/g is Δ -differentiable at t with*

$$\left(\frac{f}{g}\right)^\Delta(t) = \frac{f^\Delta(t)g(t) - f(t)g^\Delta(t)}{g(t)g^\sigma(t)}.$$

Definition 7 (Definition 1.71 of [3]). A function $F : \mathbb{T} \rightarrow \mathbb{R}$ is called an antiderivative of $f : \mathbb{T} \rightarrow \mathbb{R}$ provided $F^\Delta(t) = f(t)$ for all $t \in \mathbb{T}^\kappa$.

Definition 8 ([3]). A function $f : \mathbb{T} \rightarrow \mathbb{R}$ is called rd-continuous provided it is continuous at right-dense points in \mathbb{T} and its left-sided limits exists (finite) at all left-dense points in \mathbb{T} .

The set of all rd-continuous functions $f : \mathbb{T} \rightarrow \mathbb{R}$ is denoted by $C_{rd} = C_{rd}(\mathbb{T}) = C_{rd}(\mathbb{T}, \mathbb{R})$. The set of functions $f : \mathbb{T} \rightarrow \mathbb{R}$ that are Δ -differentiable and whose derivative is rd-continuous is denoted by $C_{rd}^1 = C_{rd}^1(\mathbb{T}) = C_{rd}^1(\mathbb{T}, \mathbb{R})$.

Theorem 9 (Theorem 1.74 of [3]). *Every rd-continuous function f has an antiderivative F . In particular, if $t_0 \in \mathbb{T}$, then F defined by*

$$F(t) := \int_{t_0}^t f(\tau) \Delta\tau, \quad t \in \mathbb{T},$$

is an antiderivative of f .

Definition 10. Let \mathbb{T} be a time scale and $a, b \in \mathbb{T}$. If $f : \mathbb{T}^\kappa \rightarrow \mathbb{R}$ is a rd-continuous function and $F : \mathbb{T} \rightarrow \mathbb{R}$ is an antiderivative of f , then the Δ -integral is defined by

$$\int_a^b f(t) \Delta t := F(b) - F(a).$$

Example 11. Let $a, b \in \mathbb{T}$ and $f : \mathbb{T} \rightarrow \mathbb{R}$ be rd-continuous. If $\mathbb{T} = \mathbb{R}$, then

$$\int_a^b f(t) \Delta t = \int_a^b f(t) dt,$$

where the integral on the right side is the usual Riemann integral. If $\mathbb{T} = h\mathbb{Z}$, $h > 0$, then

$$\int_a^b f(t) \Delta t = \begin{cases} \sum_{k=\frac{a}{h}}^{\frac{b}{h}-1} f(kh)h, & \text{if } a < b, \\ 0, & \text{if } a = b, \\ -\sum_{k=\frac{b}{h}}^{\frac{a}{h}-1} f(kh)h, & \text{if } a > b. \end{cases}$$

Theorem 12 (Theorem 1.75 of [3]). *If $f \in C_{rd}$ and $t \in \mathbb{T}^\kappa$, then*

$$\int_t^{\sigma(t)} f(\tau) \Delta\tau = \mu(t)f(t).$$

Theorem 13 (Theorem 1.77 of [3]). *If $a, b \in \mathbb{T}$, $a \leq c \leq b$, $\alpha \in \mathbb{R}$, and $f, g \in C_{rd}(\mathbb{T}, \mathbb{R})$, then:*

1. $\int_a^b (f(t) + g(t)) \Delta t = \int_a^b f(t) \Delta t + \int_a^b g(t) \Delta t,$
2. $\int_a^b \alpha f(t) \Delta t = \alpha \int_a^b f(t) \Delta t,$
3. $\int_a^b f(t) \Delta t = -\int_b^a f(t) \Delta t,$
4. $\int_a^b f(t) \Delta t = \int_a^c f(t) \Delta t + \int_c^b f(t) \Delta t,$

5. $\int_a^a f(t)\Delta t = 0,$
6. $\int_a^b f(t)g^\Delta(t)\Delta t = f(t)g(t)|_{t=a}^{t=b} - \int_a^b f^\Delta(t)g^\sigma(t)\Delta t,$
7. $\int_a^b f^\sigma(t)g^\Delta(t)\Delta t = f(t)g(t)|_{t=a}^{t=b} - \int_a^b f^\Delta(t)g(t)\Delta t,$
8. *if $f(t) \geq 0$ for all $a \leq t < b$, then $\int_a^b f(t)\Delta t \geq 0.$*

2.2 Delta dynamic equations

We now recall the definition and main properties of the delta exponential function. The general solution to a linear and homogenous second-order delta differential equation with constant coefficients is given.

Definition 14 (Definition 2.25 of [3]). We say that a function $p : \mathbb{T} \rightarrow \mathbb{R}$ is regressive if

$$1 + \mu(t)p(t) \neq 0$$

for all $t \in \mathbb{T}^\kappa$. The set of all regressive and rd-continuous functions $f : \mathbb{T} \rightarrow \mathbb{R}$ is denoted by $\mathcal{R} = \mathcal{R}(\mathbb{T}) = \mathcal{R}(\mathbb{T}, \mathbb{R})$.

Definition 15 (Definition 2.30 of [3]). If $p \in \mathcal{R}$, then we define the exponential function by

$$e_p(t, s) := \exp \left(\int_s^t \xi_{\mu(\tau)}(p(\tau))\Delta\tau \right), \quad s, t \in \mathbb{T},$$

where ξ_μ is the cylinder transformation (see [3, Definition 2.21]).

Example 16. Let \mathbb{T} be a time scale, $t_0 \in \mathbb{T}$, and $\alpha \in \mathcal{R}(\mathbb{T}, \mathbb{R})$. If $\mathbb{T} = \mathbb{R}$, then $e_\alpha(t, t_0) = e^{\alpha(t-t_0)}$ for all $t \in \mathbb{T}$. If $\mathbb{T} = h\mathbb{Z}$, $h > 0$, and $\alpha \in \mathbb{C} \setminus \{-\frac{1}{h}\}$ is a constant, then

$$e_\alpha(t, t_0) = (1 + \alpha h)^{\frac{t-t_0}{h}} \quad \text{for all } t \in \mathbb{T}. \quad (2)$$

Theorem 17 (Theorem 2.36 of [3]). *Let $p, q \in \mathcal{R}$ and $\ominus p(t) := \frac{-p(t)}{1+\mu(t)p(t)}$. The following holds:*

1. $e_0(t, s) \equiv 1$ and $e_p(t, t) \equiv 1;$
2. $e_p(\sigma(t), s) = (1 + \mu(t)p(t))e_p(t, s);$
3. $\frac{1}{e_p(t, s)} = e_{\ominus p}(t, s);$
4. $e_p(t, s) = \frac{1}{e_p(s, t)} = e_{\ominus p}(s, t);$
5. $\left(\frac{1}{e_p(t, s)} \right)^\Delta = \frac{-p(t)}{e_p^\sigma(t, s)}.$

Theorem 18 (Theorem 2.62 of [3]). *Suppose $y^\Delta = p(t)y$ is regressive, that is, $p \in \mathcal{R}$. Let $t_0 \in \mathbb{T}$ and $y_0 \in \mathbb{R}$. The unique solution to the initial value problem*

$$y^\Delta(t) = p(t)y(t), \quad y(t_0) = y_0,$$

is given by $y(t) = e_p(t, t_0)y_0$.

Let us consider the following linear second-order dynamic homogeneous equation with constant coefficients:

$$y^{\Delta\Delta} + \alpha y^\Delta + \beta y = 0, \quad \alpha, \beta \in \mathbb{R}. \quad (3)$$

We say that the dynamic equation (3) is regressive if $1 - \alpha\mu(t) + \beta\mu^2(t) \neq 0$ for $t \in \mathbb{T}^\kappa$, i.e., $\beta\mu - \alpha \in \mathcal{R}$.

Definition 19 (Definition 3.5 of [3]). Given two delta differentiable functions y_1 and y_2 , we define the Wronskian $W(y_1, y_2)(t)$ by

$$W(y_1, y_2)(t) := \det \begin{bmatrix} y_1(t) & y_2(t) \\ y_1^\Delta(t) & y_2^\Delta(t) \end{bmatrix}.$$

We say that two solutions y_1 and y_2 of (3) form a fundamental set of solutions (or a fundamental system) for (3), provided $W(y_1, y_2)(t) \neq 0$ for all $t \in \mathbb{T}^\kappa$.

Theorem 20 (Theorem 3.16 of [3]). *If (3) is regressive and $\alpha^2 - 4\beta \neq 0$, then a fundamental system for (3) is given by $e_{\lambda_1}(\cdot, t_0)$ and $e_{\lambda_2}(\cdot, t_0)$, where $t_0 \in \mathbb{T}^\kappa$ and λ_1 and λ_2 are given by*

$$\lambda_1 := \frac{-\alpha - \sqrt{\alpha^2 - 4\beta}}{2}, \quad \lambda_2 := \frac{-\alpha + \sqrt{\alpha^2 - 4\beta}}{2}.$$

Theorem 21 (Theorem 3.32 of [3]). *Suppose that $\alpha^2 - 4\beta < 0$. Define $p = \frac{-\alpha}{2}$ and $q = \frac{\sqrt{4\beta - \alpha^2}}{2}$. If p and $\mu\beta - \alpha$ are regressive, then a fundamental system of (3) is given by $\cos_{\frac{q}{(1+\mu p)}}(\cdot, t_0)e_p(\cdot, t_0)$ and $\sin_{\frac{q}{(1+\mu p)}}(\cdot, t_0)e_p(\cdot, t_0)$, where $t_0 \in \mathbb{T}^\kappa$.*

Theorem 22 (Theorem 3.34 of [3]). *Suppose $\alpha^2 - 4\beta = 0$. Define $p = \frac{-\alpha}{2}$. If $p \in \mathcal{R}$, then a fundamental system of (3) is given by*

$$e_p(t, t_0) \quad \text{and} \quad e_p(t, t_0) \int_{t_0}^t \frac{1}{1 + p\mu(\tau)} \Delta\tau,$$

where $t_0 \in \mathbb{T}^\kappa$.

Theorem 23 (Theorem 3.7 of [3]). *If functions y_1 and y_2 form a fundamental system of solutions for (3), then $y(t) = \alpha y_1(t) + \beta y_2(t)$, where α, β are constants, is a general solution to (3), i.e., every function of this form is a solution to (3) and every solution of (3) is of this form.*

2.3 Calculus of variations on time scales

Consider the following problem of the calculus of variations on time scales:

$$\mathcal{L}(y) = \int_a^b L(t, y(t), y^\Delta(t)) \Delta t \longrightarrow \min \quad (4)$$

subject to the boundary conditions

$$y(a) = y_a, \quad y(b) = y_b, \quad (5)$$

where $L : [a, b]_{\mathbb{T}}^\kappa \times \mathbb{R}^2 \rightarrow \mathbb{R}$, $(t, y, v) \mapsto L(t, y, v)$, is a given function, and $y_a, y_b \in \mathbb{R}$.

Definition 24. A function $y \in C_{rd}^1([a, b]_{\mathbb{T}}, \mathbb{R})$ is said to be an admissible path to problem (4) if it satisfies the given boundary conditions (5).

We assume that $L(t, \cdot, \cdot)$ is differentiable in (y, v) ; $L(t, \cdot, \cdot)$, $L_y(t, \cdot, \cdot)$ and $L_v(t, \cdot, \cdot)$ are continuous at (y, y^Δ) uniformly at t and rd-continuously at t for any admissible path; the functions $L(\cdot, y(\cdot), y^\Delta(\cdot))$, $L_y(\cdot, y(\cdot), y^\Delta(\cdot))$ and $L_v(\cdot, y(\cdot), y^\Delta(\cdot))$ are Δ -integrable on $[a, b]$ for any admissible path y .

Definition 25. We say that an admissible function \hat{y} is a local minimizer to problem (4)–(5) if there exists $\delta > 0$ such that $\mathcal{L}(\hat{y}) \leq \mathcal{L}(y)$ for all admissible functions $y \in C_{rd}^1$ satisfying the inequality $\|y - \hat{y}\| < \delta$. The following norm in C_{rd}^1 is considered:

$$\|y\| := \sup_{t \in [a, b]_{\mathbb{T}}^{\kappa}} |y(t)| + \sup_{t \in [a, b]_{\mathbb{T}}^{\kappa}} |y^{\Delta}(t)|.$$

Theorem 26 (Corollary 1 of [6]). *If y is a local minimizer to problem (4)–(5), then y satisfies the Euler–Lagrange equation*

$$L_v(t, y(t), y^{\Delta}(t)) = \int_a^{\sigma(t)} L_y(\tau, y(\tau), y^{\Delta}(\tau)) \Delta\tau + c \quad (6)$$

for some constant $c \in \mathbb{R}$ and all $t \in [a, b]_{\mathbb{T}}^{\kappa}$.

3 Economical model

The inflation rate, p , affects decisions of the society regarding consumption and saving, and therefore aggregated demand for domestic production, which in turn affects the rate of unemployment, u . A relationship between the inflation rate and the rate of unemployment is described by the Phillips curve, the most commonly used term in the analysis of inflation and unemployment [15]. Having a Phillips tradeoff between u and p , what is then the best combination of inflation and unemployment over time? To answer this question, we follow here the formulations presented in [5, 16]. The Phillips tradeoff between u and p is defined as

$$p := -\beta u + \pi, \quad \beta > 0, \quad (7)$$

where π is the expected rate of inflation that is captured by the equation

$$\pi' = j(p - \pi), \quad 0 < j \leq 1. \quad (8)$$

The government loss function, λ , is specified in the following quadratic form:

$$\lambda = u^2 + \alpha p^2, \quad (9)$$

where $\alpha > 0$ is the weight attached to government's distaste for inflation relative to the loss from income deviating from its equilibrium level. Combining (7) and (8), and substituting the result into (9), we obtain that

$$\lambda(\pi(t), \pi'(t)) = \left(\frac{\pi'(t)}{\beta j} \right)^2 + \alpha \left(\frac{\pi'(t)}{j} + \pi(t) \right)^2,$$

where α , β , and j are real positive parameters that describe the relations between all variables that occur in the model [16]. The problem consists to find the optimal path π that minimizes the total social loss over the time interval $[0, T]$. The initial and the terminal values of π , π_0 and π_T , respectively, are given with $\pi_0, \pi_T > 0$. To recognize the importance of the present over the future, all social losses are discounted to their present values via a positive discount rate δ . Two models are available in the literature: the *continuous model*

$$\Lambda_C(\pi) = \int_0^T \lambda(\pi(t), \pi'(t)) e^{-\delta t} dt \longrightarrow \min, \quad (10)$$

subject to given boundary conditions

$$\pi(0) = \pi_0, \quad \pi(T) = \pi_T, \quad (11)$$

and the *discrete model*

$$\Lambda_D(\pi) = \sum_{t=0}^{T-1} \lambda(\pi(t), \Delta\pi(t))(1 + \delta)^{-t} \longrightarrow \min, \quad (12)$$

also subject to given boundary conditions (11). In both cases (10) and (12),

$$\lambda(t, \pi, v) := \left(\frac{v}{\beta j}\right)^2 + \alpha \left(\frac{v}{j} + \pi\right)^2. \quad (13)$$

Here we propose the more general *time-scale model*

$$\Lambda_{\mathbb{T}}(\pi) = \int_0^T \lambda(t, \pi(t), \pi^\Delta(t)) e_{\ominus\delta}(t, 0) \Delta t \longrightarrow \min \quad (14)$$

subject to boundary conditions (11) and with λ defined by (13). Clearly, the time-scale model includes both the discrete and continuous models as special cases: our time-scale functional (14) reduces to (10) when $\mathbb{T} = \mathbb{R}$ and to (12) when $\mathbb{T} = \mathbb{Z}$.

4 Main results

Standard dynamic economic models are set up in either continuous or discrete time. Since time scale calculus can be used to model dynamic processes whose time domains are more complex than the set of integers or real numbers, the use of time scales in economy is a flexible and capable modelling technique. In this section we show the advantage of using (14) with the periodic time scale. We begin by obtaining in Section 4.1 a necessary and also a sufficient optimality condition for our economic model (14): Theorem 27 and Theorem 33, respectively. For $\mathbb{T} = h\mathbb{Z}$, $h > 0$, the explicit solution $\hat{\pi}$ to the problem (14) subject to (11) is given (Theorem 34). Afterwards, we use such results with empirical data (Section 4.2).

4.1 Theoretical results

Let us consider the problem

$$\mathcal{L}(\pi) = \int_0^T L(t, \pi(t), \pi^\Delta(t)) \Delta t \longrightarrow \min \quad (15)$$

subject to boundary conditions

$$\pi(0) = \pi_0, \quad \pi(T) = \pi_T. \quad (16)$$

As explained in Section 3, we are particularly interested in the situation where

$$L(t, \pi(t), \pi^\Delta(t)) = \left[\left(\frac{\pi^\Delta(t)}{\beta j}\right)^2 + \alpha \left(\frac{\pi^\Delta(t)}{j} + \pi(t)\right)^2 \right] e_{\ominus\delta}(t, 0). \quad (17)$$

For simplicity, in the sequel we use the notation $[\pi](t) := (t, \pi(t), \pi^\Delta(t))$.

Theorem 27. *If $\hat{\pi}$ is a local minimizer to problem (15)–(16) and the graininess function μ is a Δ -differentiable function on $[0, T]_{\mathbb{T}}^{\kappa}$, then $\hat{\pi}$ satisfies the Euler–Lagrange equation*

$$(L_v[\pi](t))^\Delta = (1 + \mu^\Delta(t)) L_y[\pi](t) + \mu^\sigma(t) (L_y[\pi](t))^\Delta \quad (18)$$

for all $t \in [0, T]_{\mathbb{T}}^{\kappa^2}$.

Proof. If $\hat{\pi}$ is a local minimizer to (15)–(16), then, by Theorem 26, $\hat{\pi}$ satisfies the following equation:

$$L_v[\pi](t) = \int_0^{\sigma(t)} L_y[\pi](\tau) \Delta\tau + c.$$

Using the properties of the Δ -integral (see Theorem 12), we can write that $\hat{\pi}$ satisfies

$$L_v[\pi](t) = \int_0^t L_y[\pi](\tau) \Delta\tau + \mu(t) L_y[\pi](t) + c. \quad (19)$$

Taking the Δ -derivative to both sides of (19), we obtain equation (18). \square

Using Theorem 27, we can immediately write the classical Euler–Lagrange equations for the continuous (10) and the discrete (12) models.

Example 28. Let $\mathbb{T} = \mathbb{R}$. Then, $\mu \equiv 0$ and (18) with the Lagrangian (17) reduces to

$$(1 + \alpha\beta^2) \pi''(t) - \delta(1 + \alpha\beta^2) \pi'(t) - \alpha j \beta^2 (\delta + j) = 0. \quad (20)$$

This is the Euler–Lagrange equation for the continuous model (10).

Example 29. Let $\mathbb{T} = \mathbb{Z}$. Then, $\mu \equiv 1$ and (18) with the Lagrangian (17) reduces to

$$(\alpha j \beta^2 - \alpha \beta^2 - 1) \Delta^2 \pi(t) + (\alpha j^2 \beta^2 + \delta \alpha \beta + \delta) \Delta \pi(t) + \alpha j \beta^2 (\delta + j) \pi(t) = 0. \quad (21)$$

This is the Euler–Lagrange equation for the discrete model (12).

Corollary 30. Let $\mathbb{T} = h\mathbb{Z}$, $h > 0$, $\pi_0, \pi_T \in \mathbb{R}$, and $T = Nh$ for a certain integer $N > 2h$. If $\hat{\pi}$ is a solution to the problem

$$\begin{aligned} \Lambda_h(\pi) &= \sum_{t=0}^{T-h} L(t, \pi(t), \pi^\Delta(t)) h \longrightarrow \min, \\ \pi(0) &= \pi_0, \quad \pi(T) = \pi_T, \end{aligned}$$

then $\hat{\pi}$ satisfies the Euler–Lagrange equation

$$(L_v[\pi](t))^\Delta = L_y[\pi](t) + h(L_y[\pi](t))^\Delta \quad (22)$$

for all $t \in \{0, \dots, T - 2h\}$.

Proof. Follows from Theorem 27 by choosing \mathbb{T} to be the periodic time scale $h\mathbb{Z}$. \square

Example 31. The Euler–Lagrange equation for problem (14) on $\mathbb{T} = h\mathbb{Z}$ is given by (22):

$$(1 + \alpha\beta^2 - \alpha\beta^2 j h) \pi^{\Delta\Delta} + (-\delta - \alpha\beta^2 \delta - \alpha\beta^2 j^2 h) \pi^\Delta + (-\alpha\beta^2 \delta j - \alpha\beta^2 j^2) \pi = 0. \quad (23)$$

Assume that $1 + \alpha\beta^2 - \alpha\beta^2 j h \neq 0$. Then equation (23) is regressive and we can use the well known theorems in the theory of dynamic equations on time scales (see Section 2.2), in order to find its general solution. Introducing the quantities

$$\Omega := 1 + \alpha\beta^2 - \alpha\beta^2 j h, \quad A := -(\delta + \alpha\beta^2 \delta + \alpha\beta^2 j^2 h), \quad B := \alpha\beta^2 j (\delta + j), \quad (24)$$

we rewrite equation (23) as

$$\pi^{\Delta\Delta} + \frac{A}{\Omega} \pi^\Delta - \frac{B}{\Omega} \pi = 0. \quad (25)$$

The characteristic equation for (25) is

$$\varphi(\lambda) = \lambda^2 + \frac{A}{\Omega}\lambda - \frac{B}{\Omega} = 0$$

with determinant

$$\zeta = \frac{A^2 + 4B\Omega}{\Omega^2}. \quad (26)$$

In general we have three different cases depending on the sign of the determinant ζ : $\zeta > 0$, $\zeta = 0$ and $\zeta < 0$. However, with our assumptions on the parameters, simple computations show that the last case cannot occur. Therefore, we consider the two possible cases:

1. If $\zeta > 0$, then we have two different characteristic roots:

$$\lambda_1 = \frac{-A + \sqrt{A^2 + 4B\Omega}}{2\Omega} > 0 \text{ and } \lambda_2 = \frac{-A - \sqrt{A^2 + 4B\Omega}}{2\Omega} < 0,$$

and by Theorem 20 and Theorem 23 we get that

$$\pi(t) = C_1 e_{\lambda_1}(t, 0) + C_2 e_{\lambda_2}(t, 0) \quad (27)$$

is the general solution to (25), where C_1 and C_2 are constants determined using the given boundary conditions (11). Using (2), we rewrite (27) as

$$\pi(t) = C_1 (1 + \lambda_1 h)^{\frac{t}{h}} + C_2 (1 + \lambda_2 h)^{\frac{t}{h}}.$$

2. If $\zeta = 0$, then by Theorems 22 and 23 we get that

$$\pi(t) = K_1 e_p(t, 0) + K_2 e_p(t, 0) \int_0^t \frac{\Delta\tau}{1 + p\mu(\tau)} \quad (28)$$

is the general solution to (25), where K_1 and K_2 are constants, determined using the given boundary conditions (11), and $p = -\frac{A}{2\Omega} \in \mathcal{R}$. Using Example 11 and (2), we rewrite (28) as

$$\pi(t) = K_1 \left(1 - \frac{A}{2\Omega}h\right)^{\frac{t}{h}} + K_2 \left(1 - \frac{A}{2\Omega}h\right)^{\frac{t}{h}} \frac{2\Omega t}{2\Omega - Ah}.$$

In certain cases one can show that the Euler–Lagrange extremals are indeed minimizers. In particular, this is true for the Lagrangian (17) under study. We recall the notion of jointly convex function (cf., e.g., [12, Definition 1.6]).

Definition 32. Function $(t, u, v) \mapsto L(t, u, v) \in C^1([a, b]_{\mathbb{T}} \times \mathbb{R}^2; \mathbb{R})$ is jointly convex in (u, v) if

$$L(t, u + u_0, v + v_0) - L(t, u, v) \geq \partial_2 L(t, u, v)u_0 + \partial_3 L(t, u, v)v_0$$

for all $(t, u, v), (t, u + u^0, v + v^0) \in [a, b]_{\mathbb{T}} \times \mathbb{R}^2$.

Theorem 33. Let $(t, u, v) \mapsto L(t, u, v)$ be jointly convex with respect to (u, v) for all $t \in [a, b]_{\mathbb{T}}$. If \hat{y} is a solution to the Euler–Lagrange equation (6), then \hat{y} is a global minimizer to (4)–(5).

Proof. Since L is jointly convex with respect to (u, v) for all $t \in [a, b]_{\mathbb{T}}$,

$$\begin{aligned} \mathcal{L}(y) - \mathcal{L}(\hat{y}) &= \int_a^b [L(t, y(t), y^\Delta(t)) - L(t, \hat{y}(t), \hat{y}^\Delta(t))] \Delta t \\ &\geq \int_a^b [\partial_2 L(t, \hat{y}(t), \hat{y}^\Delta(t)) \cdot (y(t) - \hat{y}(t)) + \partial_3 L(t, \hat{y}(t), \hat{y}^\Delta(t)) \cdot (y^\Delta(t) - \hat{y}^\Delta(t))] \Delta t \end{aligned}$$

for any admissible path y . Let $h(t) := y(t) - \hat{y}(t)$. Using boundary conditions (5), we obtain that

$$\begin{aligned} \mathcal{L}(y) - \mathcal{L}(\hat{y}) &\geq \int_a^b h^\Delta(t) \left[- \int_a^{\sigma(t)} \partial_2 L(\tau, \hat{y}(\tau), \hat{y}^\Delta(\tau)) \Delta\tau + \partial_3 L(t, \hat{y}(t), \hat{y}^\Delta(t)) \right] \Delta t \\ &\quad + h(t) \int_a^b \partial_2 L(t, \hat{y}(t), \hat{y}^\Delta(t)) \Delta t \Big|_a^b \\ &= \int_a^b h^\Delta(t) \left[- \int_a^{\sigma(t)} \partial_2 L(\tau, \hat{y}(\tau), \hat{y}^\Delta(\tau)) \Delta\tau + \partial_3 L(t, \hat{y}(t), \hat{y}^\Delta(t)) \right] \Delta t. \end{aligned}$$

From (6) we get

$$\mathcal{L}(y) - \mathcal{L}(\hat{y}) \geq \int_a^b h^\Delta(t) c \Delta t = 0$$

for some $c \in \mathbb{R}$. Hence, $\mathcal{L}(y) - \mathcal{L}(\hat{y}) \geq 0$. \square

Combining Examples 16 and 31 and Theorem 33, we obtain the central result to be applied in Section 4.2.

Theorem 34 (Solution to the total social loss problem of the calculus of variations in the time scale $\mathbb{T} = h\mathbb{Z}$, $h > 0$). *Let us consider the economic problem*

$$\begin{aligned} \Lambda_h(\pi) &= \sum_{t=0}^{T-h} \left[\left(\frac{\pi^\Delta(t)}{\beta j} \right)^2 + \alpha \left(\frac{\pi^\Delta(t)}{j} + \pi(t) \right)^2 \right] \left(1 - \frac{h\delta}{1+h\delta} \right)^{\frac{t}{h}} h \rightarrow \min, \\ \pi(0) &= \pi_0, \quad \pi(T) = \pi_T, \end{aligned} \quad (29)$$

discussed in Section 3 with $\mathbb{T} = h\mathbb{Z}$, $h > 0$, and the delta derivative given by (1). More precisely, let $T = Nh$ for a certain integer $N > 2h$, $\alpha, \beta, \delta, \pi_0, \pi_T \in \mathbb{R}^+$, and $0 < j \leq 1$ be such that $h > 0$ and $1 + \alpha\beta^2 - \alpha\beta^2 jh \neq 0$. Let Ω , A and B be given as in (24).

1. If $A^2 + 4B\Omega > 0$, then the solution $\hat{\pi}$ to problem (29) is given by

$$\hat{\pi}(t) = C \left(1 - \frac{A - \sqrt{A^2 + 4B\Omega}}{2\Omega} h \right)^{\frac{t}{h}} + (\pi_0 - C) \left(1 - \frac{A + \sqrt{A^2 + 4B\Omega}}{2\Omega} h \right)^{\frac{t}{h}}, \quad (30)$$

$t \in \{0, \dots, T - 2h\}$, where

$$C := \frac{\pi_T - \pi_0 \left(\frac{2\Omega - hA - h\sqrt{A^2 + 4B\Omega}}{2\Omega} \right)^{\frac{T}{h}}}{\left(\frac{2\Omega - hA + h\sqrt{A^2 + 4B\Omega}}{2\Omega} \right)^{\frac{T}{h}} - \left(\frac{2\Omega - hA - h\sqrt{A^2 + 4B\Omega}}{2\Omega} \right)^{\frac{T}{h}}}.$$

2. If $A^2 + 4B\Omega = 0$, then the solution $\hat{\pi}$ to problem (29) is given by

$$\hat{\pi}(t) = \left(1 - \frac{A}{2\Omega} h \right)^{\frac{t}{h}} \pi_0 + \left(1 - \frac{A}{2\Omega} h \right)^{\frac{t}{h}} \left[\pi_T \left(\frac{2\Omega}{2\Omega - Ah} \right)^{\frac{T}{h}} - \pi_0 \right] \frac{t}{T}, \quad (31)$$

$t \in \{0, \dots, T - 2h\}$.

Proof. From Example 31, $\hat{\pi}$ satisfies the Euler–Lagrange equation for problem (29). Moreover, the Lagrangian of functional Λ_h of (29) is a convex function because it is the sum of convex functions. Hence, by Theorem 33, $\hat{\pi}$ is a global minimizer. \square




Figure 1: The expected rate of inflation $\hat{\pi}(t)$ during the year of 2000 in USA, obtained from the classical discrete model (12) (upper function) and the classical continuous model (10) (lower function), with boundary conditions (11) from January ($t = 0$) and December ($t = 11$), together with the empirical rate of inflation with real data from 2000 [10, 17] (function in the middle).

4.2 Empirical results

We have three forms for the total social loss: continuous (10), discrete (12), and on a time scale \mathbb{T} (14). Our idea is to compare the implications of one model with those of another using empirical data: the rate of inflation p from [10] and the rate of unemployment u from [17], which were being collected each month in the USA over 11 years, from 2000 to 2010. We consider the coefficients

$$\beta := 3, \quad j := \frac{3}{4}, \quad \alpha := \frac{1}{2}, \quad \delta := \frac{1}{4},$$

borrowed from [5]. Therefore, the time-scale total social loss functional for one year is

$$\Lambda_{\mathbb{T}}(\pi) = \int_0^{11} \left[\frac{16}{9} (\pi^{\Delta}(t))^2 + \frac{1}{2} \left(\frac{4}{3} \pi^{\Delta}(t) + \pi(t) \right)^2 \right] e_{\ominus \frac{1}{4}}(t, 0) \Delta t. \quad (32)$$

Empirical values π_E of the expected rate of inflation, π , for all months in each year, are calculated using (7) and appropriate values of p and u [10, 17]. In the sequel, the boundary conditions $\pi(0)$ and $\pi(11)$ will be selected from empirical data in January and December, respectively. We shall compare the minimum values of the total social loss functional (32) obtained from continuous and discrete models and the value for empirical data, i.e., the value of the discrete functional $\Lambda_D(\pi_E) =: \Lambda_E$ computed with empirical data π_E .

In the continuous case we use the Euler–Lagrange equation (20) with appropriate boundary conditions in order to find the optimal path that minimizes Λ_C over each year. Then, we calculate the optimal values of Λ_C for each year (see 2nd column of Table 1). In the 3rd column of Table 1 we collect empirical values of total social loss Λ_E for each year, which are obtained by (12) from empirical data. We find the optimal path that minimizes Λ_D over each year using the Euler–Lagrange equation (21) with appropriate boundary conditions. The optimal values of Λ_D for each year are given in the 5th column of Table 1. The paths obtained from the three approaches, using empirical data from 2000, are presented in Figure 1. The implications obtained from the three methods in a fixed year are very different, independently of the year we chose. Table 2 shows the relative errors between Λ_C and Λ_E (the 3rd column), Λ_D and Λ_E (the 4th column). Our research




Figure 2: The three functions of Figure 1 together with the one obtained from our time-scale model (14) and Theorem 34, illustrating the fact that the expected rate of inflation given by (30) with $h = 0,22$ approximates well the empirical rate of inflation.

was motivated by these discrepancies. Why are the results so different? Is it caused by poor design of the model or maybe by something else?

We focus on the data collection time sampling and consider it as a cause of those differences in the results. There may exist other reasons, but we examine here the data gathering. Let us consider our time-scale model in which we consider functional (32) over a periodic time scale $\mathbb{T} = h\mathbb{Z}$. In each year we change the time scale by changing h , in such a way that the sum in the functional makes sense, and we are seeking such value of h for which the absolute error between the minimal values of the functional (32) and Λ_E is minimal. In Table 1, the 6th column presents the values of the most appropriate h and the 4th column the minimal values of the total social loss that correspond to them. Figure 2 presents the optimal paths for the continuous, discrete and time-scale models together with the empirical path, obtained using real data from 2000 [10, 17]. In the 2nd column of Table 2 we collect the relative errors between the minimal values of functional Λ_E and Λ_h .

year	The value of the functional in different approaches				the best h
	continuous	empirical	time scales	discrete	
	Λ_C	Λ_E	Λ_h	Λ_D	
2000	37,08888039	457,1493181	487,1508715	2470	0,22
2001	52,78839446	522,8060796	536,0298868	3040	0,11
2002	63,88123645	673,399954	663,2573844	3820	0,11
2003	62,01139398	811,1909476	853,5383036	4520	0,2
2004	61,72908568	703,7663513	699,714732	4130	0,11
2005	56,01553586	672,0977499	665,8735854	4060	0,1
2006	45,73885179	592,0374216	594,1793342	3700	0,1
2007	53,65457721	505,8743517	511,5351347	2910	0,1
2008	73,4472459	785,9852316	746,8126214	4260	0,11
2009	144,2965207	1352,738181	1357,167459	6330	0,22
2010	153,4630805	1819,572063	1865,77131	11400	0,1
11years	12,89356177	480,5729081	446,1625854	2E+91	0,11

Table 1: Comparison of the values of the total social loss functionals in different approaches.

year	Relative error between the empirical value Λ_E and the result in		
	time scale $h\mathbb{Z}$ with the best h	continuous approach	discrete classic approach
	$\frac{\Lambda_h - \Lambda_E}{\Lambda_E}$	$\frac{\Lambda_C - \Lambda_E}{\Lambda_E}$	$\frac{\Lambda_D - \Lambda_E}{\Lambda_E}$
2000	6,562747053	91,88692208	440,3048637
2001	2,529390479	89,90287288	481,4775533
2002	1,506173195	90,51362625	467,270606
2003	5,220393068	92,35551208	457,2054291
2004	0,575705174	91,2287529	486,8424929
2005	0,926080247	91,66556712	504,0787967
2006	0,361786602	92,27433096	524,9604949
2007	1,119009687	89,39369489	475,2416564
2008	4,98388629	90,6553911	441,9949165
2009	0,327430545	89,33300451	367,939775
2010	2,539017164	91,56597952	526,5209404
11 years	7,160271027	97,31704356	4,1617E+90

Table 2: Relative errors.

5 Conclusions

We introduced a time-scale model to the total social loss over a certain time interval under study. During examination of the proposed time-scale model for $\mathbb{T} = h\mathbb{Z}$, $h > 0$, we changed the graininess parameter. Our goal was to obtain the most similar value of the total social loss functional Λ_h to its real value, i.e., the value from empirical data. We analyzed 11 years with real data from [10, 17]. With a well-chosen time scale, we found a small relative error between the real value of the total social loss and the value obtained from our time-scale model (see the 2nd column of Table 2). We conclude that the lack of accurate results by the classical models arise due to an inappropriate frequency data collection. Indeed, if one measures the level of inflation and unemployment about once a week, which is suggested by the values of h obtained from the time-scale model, e.g., $h = 0,11$ or $h = 0,2$ (here $h = 1$ corresponds to one month), the credibility of the results obtained from the classical methods will be much higher. In other words, similar results to the ones obtained by our time-scale model can be obtained with the classical models, if a higher frequency of data collection could be used. In practical terms, however, to collect the levels of inflation and unemployment on a weekly basis is not realizable, and the calculus of variations on time scales [2, 7] assumes an important role.

Acknowledgements

This work was supported by *FEDER* funds through *COMPETE* — Operational Programme Factors of Competitiveness (“Programa Operacional Factores de Competitividade”) and by Portuguese funds through the *Center for Research and Development in Mathematics and Applications* (University of Aveiro) and the Portuguese Foundation for Science and Technology (“FCT — Fundação para a Ciência e a Tecnologia”), within project PEst-C/MAT/UI4106/2011 with *COMPETE* number FCOMP-01-0124-FEDER-022690. Dryl was also supported by FCT through the Ph.D. fellowship SFRH/BD/51163/2010; Malinowska by Bialystok University of Technology grant S/WI/02/2011; and Torres by EU funding under the 7th Framework Programme FP7-PEOPLE-2010-ITN, grant agreement number 264735-SADCO.

The authors are grateful to two anonymous referees for valuable suggestions and comments, which improved the quality of the paper.

References

- [1] R. Almeida and D. F. M. Torres, Isoperimetric problems on time scales with nabla derivatives, *J. Vib. Control* **15** (2009), no. 6, 951–958. [arXiv:0811.3650](#)
- [2] Z. Bartosiewicz and D. F. M. Torres, Noether’s theorem on time scales, *J. Math. Anal. Appl.* **342** (2008), no. 2, 1220–1226. [arXiv:0709.0400](#)
- [3] M. Bohner and A. Peterson, *Dynamic equations on time scales*, Birkhäuser Boston, Boston, MA, 2001.
- [4] M. Bohner and A. Peterson, *Advances in dynamic equations on time scales*, Birkhäuser Boston, Boston, MA, 2003.
- [5] A. C. Chiang, *Elements of dynamic optimization*, McGraw-Hill, Inc., Singapore, 1992.
- [6] R. A. C. Ferreira, A. B. Malinowska and D. F. M. Torres, Optimality conditions for the calculus of variations with higher-order delta derivatives, *Appl. Math. Lett.* **24** (2011), no. 1, 87–92. [arXiv:1008.1504](#)
- [7] E. Girejko, A. B. Malinowska and D. F. M. Torres, A unified approach to the calculus of variations on time scales, Proceedings of 2010 CCDC, Xuzhou, China, May 26-28, 2010. In: IEEE Catalog Number CFP1051D-CDR, 2010, 595–600. DOI:10.1109/CCDC.2010.5498972 [arXiv:1005.4581](#)
- [8] E. Girejko, A. B. Malinowska and D. F. M. Torres, The contingent epiderivative and the calculus of variations on time scales, *Optimization* **61** (2012), no. 3, 251–264. [arXiv:1007.0509](#)
- [9] S. Hilger, Differential and difference calculus—unified!, *Nonlinear Anal.* **30** (1997), no. 5, 2683–2694.
- [10] InflationData.Com, Current inflation, http://inflationdata.com/Inflation/Inflation_Rate/CurrentInflation
- [11] A. B. Malinowska and D. F. M. Torres, Euler-Lagrange equations for composition functionals in calculus of variations on time scales, *Discrete Contin. Dyn. Syst.* **29** (2011), no. 2, 577–593. [arXiv:1007.0584](#)
- [12] A. B. Malinowska and D. F. M. Torres, *Introduction to the fractional calculus of variations*, Imp. Coll. Press, London, 2012.
- [13] N. Martins and D. F. M. Torres, Generalizing the variational theory on time scales to include the delta indefinite integral, *Comput. Math. Appl.* **61** (2011), no. 9, 2424–2435. [arXiv:1102.3727](#)
- [14] D. Mozyrska and D. F. M. Torres, A study of diamond-alpha dynamic equations on regular time scales, *Afr. Diaspora J. Math. (N.S.)* **8** (2009), no. 1, 35–47. [arXiv:0902.1380](#)
- [15] P. A. Samuelson and W. D. Nordhaus, *Economics*, 18th ed., McGraw-Hill/Irwin, Boston, MA, 2004.
- [16] D. Taylor, Stopping inflation in the Dornbusch model: Optimal monetary policies with alternate price-adjustment equations, *Journal of Macroeconomics* **11** (1989), no. 2, 199–216.
- [17] UnemploymentData.com, Unemployment rate, <http://unemploymentdata.com/unemployment-rate>

Corrigendum to “A time-scale variational approach to inflation, unemployment and social loss”*

Monika Dryl¹
monikadryl@ua.pt

Agnieszka B. Malinowska²
a.malinowska@pb.edu.pl

Delfim F. M. Torres¹
delfim@ua.pt

¹Center for Research and Development in Mathematics and Applications,
Department of Mathematics, University of Aveiro, 3810-193 Aveiro, Portugal

²Faculty of Computer Science, Białystok University of Technology,
15-351 Białystok, Poland

Abstract

The aim of this note is to point out an error in the empirical results of Section 4.2 of [Control Cybernet. 42 (2013), no. 2, 399–418].

We regret to announce that there was an error in the **Maple** computer file that was used to compute the values for the Example 7 of [1], which adversely affects the results of the paper. Our aim is to clarify here this matter.

Example 7 of [1] is based on real data: the rate of inflation p and the rate of unemployment u , taken respectively from *InflationData.com* and *UnemoloymentData.com*, and which were being collected each month in the USA over 11 years, from 2000 to 2010. The time-scale economic model given in Section 4.1 of [1] is

$$\mathcal{L}(\pi) = \int_0^T L(t, \pi(t), \pi^\Delta(t)) \Delta t \longrightarrow \min$$

subject to boundary conditions

$$\pi(0) = \pi_0, \quad \pi(T) = \pi_T,$$

where the Lagrangian L is given by

$$L(t, \pi(t), \pi^\Delta(t)) = \left[\left(\frac{\pi^\Delta(t)}{\beta j} \right)^2 + \alpha \left(\frac{\pi^\Delta(t)}{j} + \pi(t) \right)^2 \right] e_{\ominus\delta}(t, 0).$$

All parameters of this model, that is, β , j , α and δ , are specified in Section 4.2 of [1]. The characteristic equation of the Euler–Lagrange equation associated to this problem is given in Example 7 of [1]:

$$\lambda^2 + \frac{A}{\Omega}\lambda - \frac{B}{\Omega} = 0. \tag{1}$$

The characteristic roots λ_1 and λ_2 of (1) are of different signs. They are presented correctly on page 410 of [1] but unfortunately they were coded incorrectly in the Maple computer file. It turns

*Scheduled to appear in *Control and Cybernetics* (ISSN: 0324-8569) 43 (2014), no. 4.

out that the wrong values for the characteristic roots lead to very interesting results while this is not the case with the correct ones. The wrong values used in the Maple computer file were

$$\lambda_1(h) = \frac{\frac{11}{8} + \frac{81h}{32} + \frac{1}{2} \sqrt{4 \frac{(-\frac{11}{8} - \frac{81h}{32})^2}{(\frac{11}{2} - \frac{27h}{8})^2} + \frac{54}{\frac{11}{2} - \frac{27h}{8}}}}{11 - \frac{27h}{4}}$$

and

$$\lambda_2(h) = \frac{\frac{11}{8} + \frac{81h}{32} - \frac{1}{2} \sqrt{4 \frac{(-\frac{11}{8} - \frac{81h}{32})^2}{(\frac{11}{2} - \frac{27h}{8})^2} + \frac{54}{\frac{11}{2} - \frac{27h}{8}}}}{11 - \frac{27h}{4}}.$$

References

- [1] M. Dryl, A. B. Malinowska, D. F. M. Torres, *A time-scale variational approach to inflation, unemployment and social loss*, Control Cybernet. **42** (2013), no. 2, 399–418. [arXiv:1304.5269](https://arxiv.org/abs/1304.5269)