

A Stochastic Gronwall Lemma

Michael Scheutzow*

April 22, 2013

Abstract

We prove a stochastic Gronwall lemma of the following type: if Z is an adapted nonnegative continuous process which satisfies a linear integral inequality with an added continuous local martingale M and a process H on the right hand side, then for any $p \in (0, 1)$ the p -th moment of the supremum of Z is bounded by a constant κ_p (which does not depend on M) times the p -th moment of the supremum of H . Our main tool is a martingale inequality which is due to D. Burkholder. We provide an alternative simple proof of the martingale inequality which provides an explicit numerical value for the constant c_p appearing in the inequality which is at most four times as large as the optimal constant.

2010 *Mathematics Subject Classification* Primary 60G44 Secondary 60H10, 60E15, 26D15

Keywords. stochastic Gronwall lemma, martingale inequality.

In this note we first state a martingale inequality which is due to D. Burkholder [1] and which estimates the p -th moment of the supremum of a continuous local martingale by a constant c_p times the p -th moment of its negative infimum for $0 < p < 1$. Then we apply the martingale inequality and prove a stochastic Gronwall lemma for a nonnegative process Z . The stochastic Gronwall lemma is useful when proving existence and uniqueness of solutions to stochastic differential equations satisfying only a one-sided Lipschitz condition (where the usual proof using the Burkholder-Davis-Gundy inequality does not apply). The point of the stochastic Gronwall lemma is that it provides an upper bound for the p -th moment of Z which does not depend on the local martingale M on the right-hand side of the inequality. The price one has to pay for this uniformity with respect to M is that one has to assume $p < 1$.

We start by formulating the martingale inequality.

Proposition 1. *For each $p \in (0, 1)$ and each continuous local martingale $M(t)$, $t \geq 0$ starting at $M(0) = 0$, we have*

$$\mathbf{E}\left(\sup_{t \geq 0} M^p(t)\right) \leq c_p \mathbf{E}\left(\left(-\inf_{t \geq 0} M(t)\right)^p\right), \quad (1)$$

where $c_p := \left(4 \wedge \frac{1}{p}\right) \frac{\pi p}{\sin(\pi p)}$.

*Institut für Mathematik, MA 7-5, Fakultät II, Technische Universität Berlin, Straße des 17. Juni 136, 10623 Berlin, FRG; ms@math.tu-berlin.de

Remark 2. The inequality was first proved by D. Burkholder ([1], Theorem 1.4) even a bit more generally (for a larger class of functions than the p -th power) but without an explicit estimate of the numerical value of c_p . We provide a short and elementary alternative proof below.

Remark 3. It is clear that the previous proposition does not extend to $p \geq 1$: consider the continuous martingale $M(t) := W(\tau_{-1} \wedge t)$ where W is standard Brownian motion and $\tau_x := \inf\{s \geq 0 : W(s) = x\}$. Then the left hand side of (1) is infinite for each $p \geq 1$ while the right hand side is finite. This example also shows that even though the constant c_p is certainly not optimal, it is at most off from the optimal constant by the factor $4 \wedge (1/p)$ (which converges to one as p approaches one). It is also clear that the proposition does not extend to right-continuous martingales: consider a martingale which is constant except for a single jump at time 1 of height 1 with probability δ and height $-\frac{\delta}{1-\delta}$ with probability $1 - \delta$ where $\delta \in (0, 1)$. It is straightforward to check that for an inequality of type $(\mathbf{E} \sup_{t \geq 0} M^p(t))^{1/p} \leq c_{p,q} (\mathbf{E} (-\inf_{t \geq 0} M(t))^q)^{1/q}$ to hold for this class of examples for some finite $c_{p,q}$, we require that $q \geq 1$ irrespective of the value of $p \in (0, 1)$.

Proof of Proposition 1. Since M is a continuous local martingale starting at 0 it can be represented as a time-changed Brownian motion W (on a suitable probability space). We can and will assume that M converges almost surely (otherwise there is nothing to prove), so there exists an almost surely finite stopping time T for W such that $A := \sup_{0 \leq t \leq T} W(t) = \sup_{0 \leq t} M(t)$ and $B := -\inf_{0 \leq t \leq T} W(t) = -\inf_{0 \leq t} M(t)$. Let $0 = a_0 < a_1 < \dots$ be a sequence which converges to ∞ and define

$$\tau_i := \inf\{t \geq 0 : W(t) = -a_i\}, \quad Y_i := \sup_{\tau_{i-1} \leq t \leq \tau_i} W(t), \quad i \in \mathbf{N}, \quad N := \inf\{i \in \mathbf{N} : \tau_i \geq T\}.$$

The Y_i are independent by the strong Markov property of W and for $p \in (0, 1)$ and $i \in \mathbf{N}$ we have

$$\Gamma_i := \mathbf{E}(Y_i \vee 0)^p = \frac{a_i - a_{i-1}}{a_i^{1-p}} \int_0^\infty \frac{1}{1 + y^{1/p}} dy = \frac{a_i - a_{i-1}}{a_i^{1-p}} \frac{\pi p}{\sin(\pi p)}.$$

Therefore,

$$\begin{aligned} \mathbf{E}A^p &\leq \sum_{n=1}^{\infty} \mathbf{E}\left(\sup\{Y_1, \dots, Y_n\}^p \mathbf{1}_{N=n}\right) \leq \sum_{n=1}^{\infty} \sum_{i=1}^n \mathbf{E}\left((Y_i \vee 0)^p \mathbf{1}_{N=n}\right) \\ &= \sum_{i=1}^{\infty} \mathbf{E}\left((Y_i \vee 0)^p \mathbf{1}_{N \geq i}\right) = \sum_{i=1}^{\infty} \Gamma_i \mathbf{P}\{N \geq i\}, \end{aligned}$$

where the last equality again follows from the strong Markov property. Inserting the formula for Γ_i , choosing the particular values $a_i = c\gamma^i$ for some $c > 0$ and $\gamma > 1$, and observing that

$\mathbf{P}\{N \geq i\} \leq \mathbf{P}\{B \geq a_{i-1}\}$, we get

$$\begin{aligned} \mathbf{E}A^p &\leq \frac{\pi p}{\sin(\pi p)} c^p \left(\gamma^p + \left(1 - \frac{1}{\gamma}\right) \sum_{i=2}^{\infty} \gamma^{ip} \mathbf{P}\{B \geq c\gamma^{i-1}\} \right) \\ &= \frac{\pi p}{\sin(\pi p)} c^p \left(\gamma^p + \left(1 - \frac{1}{\gamma}\right) \sum_{j=2}^{\infty} \mathbf{P}\{B \in [c\gamma^{j-1}, c\gamma^j]\} \left(\frac{\gamma^{p(j+1)} - 1}{\gamma^p - 1} - 1 - \gamma^p \right) \right) \\ &\leq \frac{\pi p}{\sin(\pi p)} \left(c^p \gamma^p + \left(1 - \frac{1}{\gamma}\right) \frac{\gamma^{2p}}{\gamma^p - 1} \mathbf{E}B^p - c^p \left(1 - \frac{1}{\gamma}\right) \left(\frac{1}{\gamma^p - 1} + 1 + \gamma^p \right) \mathbf{P}\{B \geq c\gamma\} \right). \end{aligned}$$

Dropping the last (negative) term, letting $c \rightarrow 0$ and observing that the function of γ in front of $\mathbf{E}B^p$ converges to $1/p$ as $\gamma \rightarrow 1$ and that $\inf_{\gamma>1} \gamma^{2p}/(\gamma^p - 1) = 4$ we obtain the assertion. \square

Next, we apply the martingale inequality to prove a stochastic Gronwall lemma. A similar stochastic Gronwall lemma was proved and used in [2] in order to prove existence and uniqueness of a solution to a stochastic functional differential equation satisfying a one-sided Lipschitz condition only. That result was slightly more general in the sense that on the right hand side of equation (2) Z was replaced by its running supremum, but it was less general concerning the function ψ and it required higher moments of H^* . The proof did not explicitly use a martingale inequality.

For a real-valued process denote $Y^*(t) := \sup_{0 \leq s \leq t} Y(s)$.

Theorem 4. *Let c_p be as in Theorem 1. Let Z and H be nonnegative, adapted processes with continuous paths and assume that ψ is nonnegative and progressively measurable. Let M be a continuous local martingale starting at 0. If*

$$Z(t) \leq \int_0^t \psi(s) Z(s) \, ds + M(t) + H(t) \quad (2)$$

holds for all $t \geq 0$, then for $p \in (0, 1)$, and $\mu, \nu > 1$ such that $\frac{1}{\mu} + \frac{1}{\nu} = 1$ and $p\nu < 1$, we have

$$\mathbf{E} \sup_{0 \leq s \leq t} Z^p(s) \leq (c_{p\nu} + 1)^{1/\nu} \left(\mathbf{E} \exp\{p\mu \int_0^t \psi(s) \, ds\} \right)^{1/\mu} \left(\mathbf{E}(H^*(t))^{p\nu} \right)^{1/\nu}. \quad (3)$$

If ψ is deterministic, then

$$\mathbf{E} \sup_{0 \leq s \leq t} Z^p(s) \leq (c_p + 1) \exp\{p \int_0^t \psi(s) \, ds\} \left(\mathbf{E}(H^*(t))^p \right), \quad (4)$$

and

$$\mathbf{E}Z(t) \leq \exp\left\{ \int_0^t \psi(s) \, ds \right\} \mathbf{E}H^*(t). \quad (5)$$

Proof. Let $L(t) := \int_0^t \exp\{-\int_0^s \psi(u) \, du\} \, dM(s)$. Applying the usual Gronwall Lemma (for each fixed $\omega \in \Omega$) to Z and integrating by parts, we obtain

$$Z(t) \leq \exp\left\{ \int_0^t \psi(s) \, ds \right\} (L(t) + H^*(t)). \quad (6)$$

Since Z is nonnegative, we have $-L(t) \leq H^*(t)$ for all $t \geq 0$. Therefore, using Proposition 1 and Hölder's inequality, we get

$$\begin{aligned} \mathbf{E}(Z^*)^p(t) &\leq \left(\mathbf{E} \exp\left\{p\mu \int_0^t \psi(s) \, ds\right\} \right)^{1/\mu} \left(\mathbf{E}(L^*(t))^{p\nu} + \mathbf{E}(H^*(t))^{p\nu} \right)^{1/\nu} \\ &\leq \left(\mathbf{E} \exp\left\{p\mu \int_0^t \psi(s) \, ds\right\} \right)^{1/\mu} (c_{p\nu} + 1)^{1/\nu} \left(\mathbf{E}(H^*(t))^{p\nu} \right)^{1/\nu}, \end{aligned}$$

which is (3). Inequality (4) follows similarly. The final statement follows by applying (6) to $\tau_n \wedge t$ for a sequence of localizing stopping times τ_n for L and applying Fatou's Lemma. \square

Acknowledgement. It is a pleasure to thank R. Bañuelos for drawing my attention to the cited paper of Donald Burkholder.

References

- [1] Burkholder, D. L. (1975). One-sided maximal functions and H^p , *J. Func. Anal.* **18**, 429–454.
- [2] v. Renesse, M., Scheutzow, M. (2010). Existence and uniqueness of solutions of stochastic functional differential equations, *Random Oper. Stoch. Equ.* **18**, 267–284.