Direct and inverse spectral problems for a class of non-selfadjoint band matrices

Natalia Bebiano and Mikhail Tyaglov

June 18, 2018

Abstract

The spectral properties of a class of band matrices are investigated. The reconstruction of matrices of this special class from given spectral data is also studied. Necessary and sufficient conditions for that reconstruction are found. The obtained results extend some results on the direct and inverse spectral problems for periodic Jacobi matrices and for some non-self-adjoint tridiagonal matrices.

AMS: 65 F15, 15A18

Keywords: Inverse eigenvalue problem, stable polynomials, Hermite-Biehler theorem, Jacobi matrix, periodic Jacobi matrix

1 Introduction

Inverse eigenvalue problems arise in mathematics as well as in many areas of engineering and science such as chemistry, geology, physics etc. Often the mathematical model describing a certain physical system involves matrices whose spectral data allow the prediction of the behavior of the system. Determining the spectra of those matrices is the so-called *direct problem*, while the *inverse problem* consists in the reconstruction of the matrices from the knowledge of the behavior of the system, frequently expressed by spectral data.

Inverse eigenvalue problems, in general, and for structured matrices, in particular, have attracted attention of many researchers, some of them motivated by the numerous applications of this scientific area (see e.g. [1], [16]). The mathematical background employed in those investigations may involve rather sophisticated techniques such as algebraic curves, functional analysis, matrix theory, etc. (see [23], [10], [11], [2], [25] and the references therein).

Inverse eigenvalue problems for band matrices have been actively investigated, e.g. see [10] and their references. The inverse spectral problem for a *periodic Jacobi matrix*, that is, a real symmetric matrix of the form

$$L_{n} = \begin{pmatrix} a_{1} & b_{1} & 0 & \dots & 0 & b_{n} \\ b_{1} & a_{2} & b_{2} & \dots & 0 & 0 \\ 0 & b_{2} & a_{3} & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & a_{n-1} & b_{n-1} \\ b_{n} & 0 & 0 & \dots & b_{n-1} & a_{n} \end{pmatrix}, \qquad b_{i} > 0,$$
(1.1)

deserved the attention of researchers, see [11, 13, 2, 26] and their references. These matrices appear in studies of the periodic Toda lattice, inverse eigenvalue problems for Sturm-Liouville equations and Hill's equation [10, 13]. If $b_n = 0$, the the matrices L_n of the form (1.1) reduce to tridiagonal symmetric matrices called the *Jacobi matrices*. The Jacobi matrices motivated intensive study as an useful tool in the investigation of orthogonal polynomials, in the theory of continued fractions, and in numerical analysis [27, 9, 8]. Namely, the inverse problems for Jacobi matrices have been an intensive topic of research since the seminal papers by Hochstadt and Hald [20, 19] in the seventies of the last century. In the present work, we study spectral properties of complex matrices of the form

$$J_{n} = \begin{pmatrix} c_{1} & b_{1} & 0 & \dots & 0 & b_{n} \\ \overline{b}_{1} & c_{2} & b_{2} & \dots & 0 & 0 \\ 0 & \overline{b}_{2} & c_{3} & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & c_{n-1} & b_{n-1} \\ b_{n} & 0 & 0 & \dots & \overline{b}_{n-1} & a_{n} \end{pmatrix},$$
(1.2)

where $b_1, \ldots, b_{n-1}, b_n \in \mathbb{C} \setminus \mathbb{R}, c_1, \ldots, c_{n-1} \in \mathbb{R}$, and $a_n \in \mathbb{C}$, and solve the direct problem for such matrices. (Here \overline{z} means the complex conjugate of z.) The matrices of the form (1.2) constitute the class \mathcal{J}_n .

We also solve the inverse spectral problem for matrices from the subclass $\widehat{\mathcal{J}}_n$ of the class \mathcal{J}_n . This class consists of the matrices of the form

$$\widehat{J}_{n} = \begin{pmatrix} \widehat{c}_{1} & \widehat{b}_{1} & 0 & \dots & 0 & \widehat{b}_{n} \\ \widehat{b}_{1} & \widehat{c}_{2} & \widehat{b}_{2} & \dots & 0 & 0 \\ 0 & \widehat{b}_{2} & \widehat{c}_{3} & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & \widehat{c}_{n-1} & \widehat{b}_{n-1} \\ \widehat{b}_{n} & 0 & 0 & \dots & \widehat{b}_{n-1} & \widehat{a}_{n} \end{pmatrix},$$

$$(1.3)$$

where $\hat{b}_1, \ldots, \hat{b}_{n-1}, \hat{c}_1, \ldots, \hat{c}_{n-1} \in \mathbb{R}$, and $\hat{a}_n, \hat{b}_n \in \mathbb{C}, \hat{b}_n \neq 0$.

Note that in [3] (see also [4]), Arlinskii and Tsekhanovskii considered the matrices of the form (1.3) with $b_n = 0$, $a_n \in \mathbb{C} \setminus \mathbb{R}$, and solved the direct and inverse eigenvalue problems for those matrices. In [26], the direct and inverse spectral problems for the matrices the form (1.3) with $b_n \in \mathbb{R} \setminus \{0\}$ and $a_n \in \mathbb{R}$ (that is, for the matrices of the form (1.1)) were solved, and necessary and sufficient conditions for solvability of the inverse problem were found.

Recall that in [26], it was established that the necessary and sufficient conditions for the inverse spectral problem for the matrices of the form (1.1) to be solvable are

$$\prod_{j=1}^{n} |\mu_k - \lambda_j| \ge (-1)^{n-k-1}\beta, \qquad k = 1, \dots, n-1.$$
(1.4)

Here $\beta = b_1 \cdots b_n$ and the sets $\{\lambda_j\}_{j=1}^n$ and $\{\mu_k\}_{k=1}^{n-1}$ are the spectra of the matrix (1.1) and its $(n-1) \times (n-1)$ leading principal submatrix, respectively.

In [3] (see also [4]), it was established that any matrix of the form (1.3) with $\hat{b}_n = 0$ and $\text{Im } a_n > 0$ has its eigenvalues in the open half-plane of the complex plane. The inverse spectral problem of such matrices was also solved in [3].

In the present work, we extend the results of the work [26] and find necessary and sufficient conditions for solvability of the inverse spectral problem for the matrices from the class $\hat{\mathcal{J}}_n$. Those conditions are also necessary for the matrices from the class \mathcal{J}_n and generalize the inequality (1.4). Also we extended the results of the paper [3] to the matrices from the class \mathcal{J}_n .

The paper is organized as follows. In Section 2, we survey some general properties of the real symmetric tridiagonal matrices. Section 3 is devoted to the study of spectral properties of matrices from the class \mathcal{J}_n . We study the eigenvalue location for real and nonreal numbers $\beta = b_1 \cdots b_n$ and a_n , and find a necessary condition for the spectra of those matrices. In Section 4, we show that for any matrix from the class \mathcal{J}_n , there exists a matrix from the class $\hat{\mathcal{J}}_n$ with the same spectral data. In this section, we solve the inverse spectral problem for the matrices from the class $\hat{\mathcal{J}}_n$, and establish that the necessary condition obtained in Section 3 is also sufficient for the inverse spectral problem to be solvable.

In Sections 3 and 4, we follow the approach developed in [26]. However, we simplify the substantiation of their technique and extend it to a more wide class of matrices.

$\mathbf{2}$ **Preliminaries**

Let us denote by J_k , k = 1, ..., n-1, the kth principal submatrix of J_n , and consider the matrix J_{n-1}

$$J_{n-1} = \begin{pmatrix} c_1 & b_1 & 0 & \dots & 0 & 0 \\ \overline{b}_1 & c_2 & b_2 & \dots & 0 & 0 \\ 0 & \overline{b}_2 & c_3 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & \underline{c}_{n-2} & b_{n-2} \\ 0 & 0 & 0 & \dots & \overline{b}_{n-2} & c_{n-1} \end{pmatrix}.$$
 (2.1)

This matrix is self-adjoint, so its characteristic polynomial

$$\chi_{n-1}(\lambda) = \det(\lambda I_{n-1} - J_{n-1})$$

has only real zeroes, the eigenvalues of the matrix J_{n-1} . Here I_{n-1} is the identity matrix of size n-1. Note that the spectrum of J_{n-1} coincides with the spectrum of the matrix (cf.[14])

$$\widetilde{J}_{n-1} = \begin{pmatrix} c_1 & |b_1| & 0 & \dots & 0 & 0 \\ |b_1| & c_2 & |b_2| & \dots & 0 & 0 \\ 0 & |b_2| & c_3 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & c_{n-2} & |b_{n-2}| \\ 0 & 0 & 0 & \dots & |b_{n-2}| & c_{n-1} \end{pmatrix}.$$
(2.2)

But this matrix is a real Jacobi matrix so its eigenvalues are simple. Thus we obtain that the spectrum of the matrix J_{n-1} , $\sigma(J_{n-1}) = \{\mu_1, \ldots, \mu_{n-1}\}$, is real and simple (e.g., see [14]):

$$\mu_1 < \mu_2 < \cdots < \mu_{n-1}$$

 \mathbf{SO}

$$\operatorname{sign}\left(\chi_{n-1}'(\mu_k)\right) = (-1)^{n-k-1},$$
(2.3)

where $\chi'_{n-1}(\lambda)$ is the derivative of the polynomial $\chi_{n-1}(\lambda)$ By $\mathbf{u}_k = (u_{k1}, \dots, u_{k,n-1})^T \in \mathbb{C}^{n-1}$ we denote the eigenvector of J_{n-1} corresponding to the eigenvalue $\mu_k, k = 1, \ldots, n-1$ such that

$$\overline{\mathbf{u}}_{j}^{T}\mathbf{u}_{k} = \delta_{jk}, \qquad (2.4)$$

where δ_{jk} is the Kronecker symbol. Then the *resolvent* of the matrix J_{n-1} has the form

$$\left(\lambda I_{n-1} - J_{n-1}\right)^{-1} = \sum_{k=1}^{n-1} \frac{\mathbf{u}_k \overline{\mathbf{u}}_k^T}{\lambda - \mu_k}.$$
(2.5)

It is easy to see that

$$\mathbf{e}_{1}^{T}(\lambda I_{n-1}-J_{n-1})^{-1}\mathbf{e}_{n-1}=\frac{b_{1}b_{2}\cdots b_{n-2}}{\chi_{n-1}(\lambda)},$$

where $\mathbf{e}_1 = (1, 0, \dots, 0, 0)^T \in \mathbb{C}^{n-1}$ and $\mathbf{e}_{n-1} = (0, 0, \dots, 0, 1)^T \in \mathbb{C}^{n-1}$. The formula (2.5) gives us

$$\frac{b_1\cdots b_{n-2}}{\chi_{n-1}(\lambda)} = \sum_{k=1}^{n-1} \frac{u_{k1}\overline{u}_{k,n-1}}{\lambda - \mu_k},$$

so we obtain

$$b_n u_{k1} b_{n-1} \overline{u}_{k,n-1} = \frac{\beta}{\chi'_{n-1}(\mu_k)}, \quad k = 1, \dots, n-1,$$
 (2.6)

where

$$\beta = b_1 b_2 \cdots b_n \neq 0. \tag{2.7}$$

In particular, $u_{k1} \neq 0$ and $u_{k,n-1} \neq 0$.

The formula
$$(2.6)$$
 implies that

$$|b_n u_{k1}|^2 |b_{n-1}\overline{u}_{k,n-1}|^2 = \frac{|\beta|^2}{[\chi'_{n-1}(\mu_k)]^2}, \quad k = 1, \dots, n-1.$$
(2.8)

3 Spectral properties of the matrices in \mathcal{J}_n

In this section, we characterize the spectra of the matrices in \mathcal{J}_n . Note first that any matrix J_n of the form (1.2) can be represented as follows

$$J_n = \begin{pmatrix} J_{n-1} & \mathbf{y} \\ \overline{\mathbf{y}}^T & a_n \end{pmatrix},$$

where $\mathbf{y} = (\overline{b}_n, 0, 0, \dots, 0, b_{n-1})^T \in \mathbb{C}^{n-1}$. By the Schur determinant formula [29], we obtain (cf. [26])

$$\frac{\chi_n(\lambda)}{\chi_{n-1}(\lambda)} = \lambda - a_n - \overline{\mathbf{y}}^T (\lambda I_{n-1} - J_{n-1})^{-1} \mathbf{y}, \qquad (3.1)$$

where $\chi_n(\lambda)$ and $\chi_{n-1}(\lambda)$ are the characteristic polynomials of the matrices J_n and J_{n-1} , respectively:

$$\chi_n(\lambda) = \det(\lambda I_n - J_n), \quad \chi_{n-1}(\lambda) = \det(\lambda I_{n-1} - J_{n-1}).$$

Furthermore, the formula (2.5) implies

$$\overline{\mathbf{y}}^{T} (\lambda I_{n-1} - J_{n-1})^{-1} \mathbf{y} = \sum_{k=1}^{n-1} \frac{|\overline{\mathbf{y}}^{T} \mathbf{u}_{k}|^{2}}{\lambda - \mu_{k}} = \sum_{k=1}^{n-1} \frac{|b_{n} u_{k1} + \overline{b}_{n-1} u_{k,n-1}|^{2}}{\lambda - \mu_{k}}.$$
 (3.2)

From (3.1) and (3.2) we get

$$\frac{\chi_n(\lambda)}{\chi_{n-1}(\lambda)} = \lambda - a_n - \sum_{k=1}^{n-1} \frac{\alpha_k}{\lambda - \mu_k},$$
(3.3)

where

$$\alpha_k = -\frac{\chi_n(\mu_k)}{\chi'_{n-1}(\mu_k)} = |b_n u_{k1} + \overline{b}_{n-1} u_{k,n-1}|^2 \ge 0$$
(3.4)

or

$$\alpha_k = |b_n u_{k1}|^2 + |b_{n-1} u_{k,n-1}|^2 + 2\operatorname{Re}(b_n u_{k1} b_{n-1} \overline{u}_{k,n-1}) \ge 0.$$
(3.5)

Note that (3.4)–(3.5) and (2.3) imply that

$$(-1)^{n-k}\chi_n(\mu_k) \ge 0. \tag{3.6}$$

Thus, the function

$$R_{n-1}(\lambda) = \lambda - \sum_{k=1}^{n-1} \frac{\alpha_k}{\lambda - \mu_k}$$

maps the upper half-plane of the complex plane to itself, since

$$\operatorname{sign} \operatorname{Im} \frac{-\alpha_k}{\lambda - \mu_k} = \operatorname{sign} \operatorname{Im} \frac{-\alpha_k (\overline{\lambda} - \mu_k)}{|\lambda - \mu_k|^2} = \operatorname{sign} \operatorname{Im} \lambda.$$

Therefore, the zeroes and poles of $R_{n-1}(\lambda)$ are real, simple and interlacing (see e.g. [21] and references therein), so (3.3) can be represented as follows

$$\frac{\chi_n(\lambda)}{\chi_{n-1}(\lambda)} = R_{n-1}(\lambda) - a_n.$$
(3.7)

By (3.4), $\chi_n(\mu_j) = 0$ for some j, j = 1, ..., n-1, if and only if $\alpha_j = 0$, or equivalently, if and only if

$$b_n u_{k1} + \overline{b}_{n-1} u_{k,n-1} = 0.$$

Let $N = \{j_1, \ldots, j_m\} \subset \{1, 2, \ldots, n-1\}$ be the set of indices such that $\alpha_j = 0$ for $j \in N$. Then the function $R_{n-1}(\lambda)$ has the form

$$R_{n-1}(\lambda) = \lambda - \sum_{\substack{k=1\\k\notin N}}^{n-1} \frac{\alpha_k}{\lambda - \mu_k},$$
(3.8)

and the polynomial $\chi_n(\lambda)$ has *m* zeroes in common with $\chi_{n-1}(\lambda)$, $\mu_{j_1}, \mu_{j_2}, \ldots, \mu_{j_m}$ while the its other zeroes are the solutions of the equation

$$R_{n-1}(\lambda) = a_n. \tag{3.9}$$

If $a_n \in \mathbb{R}$, then the function $R_{n-1}(\lambda) - a_n$ maps the upper half-plane to itself, so the solutions of the equation (3.9) are real and simple and interlace the numbers μ_k , $k \in \{1, 2, ..., n-1\}\setminus N$. However, they may coincide with some numbers μ_j , $j \in N$. So for $a_n \in \mathbb{R}$, the eigenvalues of J_n are real and of multiplicity at most two. Multiple eigenvalues are always eigenvalues of J_{n-1} . This property of the spectrum of J_n with $b_k \in \mathbb{C}$, k = 1, 2..., n is the same as for real b_k , e.g. [26].

Suppose now that a_n is nonreal, and recall that $R_{n-1}(\lambda)$ maps the upper (lower) half-plane to itself. Therefore, all solutions of the equation (3.9) lie in the upper (lower) half-plane of the complex plane whenever Im $a_n > 0$ (Im $a_n < 0$), and cannot be real since the functions mapping the upper half-plane to itself are always real on the real line (see e.g. [21] and references there). Note that the famous Hermite-Biehler theorem can be proved by the same technique, see [24].

Finally, we note that the formulæ (3.4) and (2.6) imply that

$$\alpha_{k} = \frac{[|b_{1}u_{k1}|^{2} + \operatorname{Re}(b_{n}u_{k1}b_{n-1}\overline{u}_{k,n-1})]^{2} + [\operatorname{Im}(b_{n}u_{k1}b_{n-1}\overline{u}_{k,n-1})]^{2}}{|b_{1}u_{k1}|^{2}} = \frac{[\chi_{n-1}'(\mu_{k})|b_{1}u_{k1}|^{2} + \operatorname{Re}\beta]^{2} + (\operatorname{Im}\beta)^{2}}{|b_{1}u_{k1}|^{2} [\chi_{n-1}'(\mu_{k})]^{2}}.$$
(3.10)

This formula shows that if $\text{Im } \beta \neq 0$, then $\alpha_k \neq 0$ for any $k = 1, \ldots, n-1$. At the same time, if $\beta \in \mathbb{R} \setminus \{0\}$, then

$$\alpha_k = \frac{[\chi'_{n-1}(\mu_k)|b_1u_{k1}|^2 + \beta]^2}{|b_1u_{k1}|^2 [\chi'_{n-1}(\mu_k)]^2}.$$

Therefore, for real nonzero β , the number α_k equals zero if and only if $\chi'_{n-1}(\mu_k)|b_1u_{k1}|^2 + \beta = 0$, that is equivalent to the following equality

$$|\chi_{n-1}'(\mu_k)||b_1u_{k1}|^2 = (-1)^{n-k}\beta,$$

by (2.3). This equality and (3.6) give us: for $\beta > 0$

$$(-1)^{n-k}\chi_n(\mu_k) > 0, \qquad k = n - 1, n - 3, n - 5, \dots, (-1)^{n-k}\chi_n(\mu_k) \ge 0, \qquad k = n - 2, n - 4, n - 6, \dots,$$
 (3.11)

$$(-1)^{n-k}\chi_n(\mu_k) \ge 0, \qquad k = n - 1, n - 3, n - 5, \dots, (-1)^{n-k}\chi_n(\mu_k) > 0, \qquad k = n - 2, n - 4, n - 6, \dots.$$
(3.12)

If additionally $a_n \in \mathbb{R}$, then the eigenvalues of the matrices J_n and J_{n-1} are distributed as follows:

for $\beta < 0$

 $\dots \leqslant \lambda_{n-5} < \mu_{n-4} < \lambda_{n-4} \leqslant \mu_{n-3} \leqslant \lambda_{n-2} < \mu_{n-2} < \lambda_{n-1} \leqslant \mu_{n-1} \leqslant \lambda_n, \tag{3.13}$

and for $\beta > 0$

$$\dots < \lambda_{n-5} \leqslant \mu_{n-4} \leqslant \lambda_{n-4} < \mu_{n-3} < \lambda_{n-2} \leqslant \mu_{n-2} \leqslant \lambda_{n-1} < \mu_{n-1} < \lambda_n.$$
(3.14)

Thus, we come to the following statements.

Theorem 3.1. Let β be nonreal. If $a_n \in \mathbb{R}$, then all the eigenvalues of the matrix J_n defined in (1.2) are real and simple and interlace the eigenvalues of J_{n-1} , which are real and simple as well.

If $a_n \in \mathbb{C}\setminus\mathbb{R}$, then all the eigenvalues of the matrix J_n lie in the **open** upper (lower) half-plane of the complex plane whenever $\operatorname{Im} a_n > 0$ ($\operatorname{Im} a_n < 0$).

Moreover, for complex a_n (real or nonreal) the characteristic polynomial of the matrix J_n satisfies the inequalities

$$(-1)^{n-k}\chi_n(\mu_k) > 0, \qquad k = 1, \dots, n-1.$$

Theorem 3.2. Let $\beta \in \mathbb{R}$. If $a_n \in \mathbb{R}$, then all the eigenvalues of the matrix J_n given in (1.2) are real and of multiplicity at most 2. Any multiple eigenvalue of J_n is an eigenvalue of J_{n-1} .

If $a_n \in \mathbb{C}\setminus\mathbb{R}$, then all the eigenvalues of the matrix J_n lie in the **closed** upper (lower) half-plane of the complex plane whenever $\operatorname{Im} a_n > 0$ ($\operatorname{Im} a_n < 0$). An eigenvalue of J_n is real if and only if it is an eigenvalue of J_{n-1} .

Moreover, the eigenvalues of J_n and J_{n-1} are distributed as in (3.13)–(3.14) for real a_n , or satisfy the inequalities (3.11)–(3.12) for any complex a_n .

Remark 3.3. From (3.3) it follows that if a_n is nonreal, then

$$\chi_n(\lambda) = \chi_{n-1}(\lambda)(\lambda - \operatorname{Re} a_n) - \sum_{k=1}^n \alpha_k \prod_{\substack{j=1\\j \neq k}}^{n-1} (\lambda - \mu_j) - i \operatorname{Im} a_n \cdot \chi_{n-1}(\lambda), \qquad (3.15)$$

that is, $\chi_{n-1}(\lambda)$ is the imaginary part (up to the constant factor $\operatorname{Im} a_n$) of the polynomial $\chi_n(\lambda)$, so $\chi_n(\mu_k) \in \mathbb{R}, k = 1, ..., n-1$.

We now study a necessary condition which the spectra of the matrices in the class \mathcal{J}_n must satisfy.

Theorem 3.4. Let $J_n \in \mathcal{J}_n$, $\chi_n(\lambda) = \det(\lambda I_n - J_n)$, and $\sigma(J_{n-1}) = \{\mu_1, \ldots, \mu_{n-1}\}$. Then

• for nonreal β

 $(-1)^{n-k}\chi_n(\mu_k) > 0 \quad and \quad |\chi_n(\mu_k) + 2\operatorname{Re}\beta| \ge 2|\beta|, \quad k = 1, \dots, n-1,$ (3.16)

• for real nonzero β

$$(-1)^{n-k}\chi_n(\mu_k) \ge 0 \quad and \quad |\chi_n(\mu_k)| \ge 4(-1)^{n-k-1}\beta, \quad k = 1, \dots, n-1,$$
(3.17)

Here β is defined in (2.7):

$$\beta = b_1 b_2 \cdots b_n \neq 0.$$

Proof. In fact, taking into account that $b_n u_{k1} \neq 0$ by (2.6) and $\chi'_{n-1}(\mu_k) \neq 0$ by (2.3), from (3.4) and (3.10) we obtain that $X_k := |b_n u_{k1}|^2$ satisfies the following equation

$$[\chi'_{n-1}(\mu_k)X_k + \operatorname{Re}\beta]^2 + \chi_n(\mu_k)\chi'_{n-1}(\mu_k)X_k + (\operatorname{Im}\beta)^2 = 0, \qquad (3.18)$$

or

$$[\chi'_{n-1}(\mu_k)]^2 X_k^2 + (\chi_n(\mu_k) + 2\operatorname{Re}\beta)\chi'_{n-1}(\mu_k)X_k + (\operatorname{Im}\beta)^2 = 0.$$
(3.19)

The solutions of this equations have the form

$$X_k^{(1,2)} = \frac{(-1)^{n-k} (\chi_n(\mu_k) + 2\operatorname{Re}\beta) \pm \sqrt{[\chi_n(\mu_k) + 2\operatorname{Re}\beta]^2 - 4|\beta|^2}}{2|\chi_{n-1}'(\mu_k)|}.$$
(3.20)

Since $b_n u_{k1} \neq 0$, X_k is positive, so we have for any (real or nonreal) β and a_n that (see (3.6))

$$\left[\chi_n(\mu_k) + 2\operatorname{Re}\beta\right]^2 - 4|\beta|^2 \ge 0, \qquad (3.21)$$

$$(-1)^{n-k}(\chi_n(\mu_k) + 2\operatorname{Re}\beta) \ge 0, \qquad (3.22)$$

$$(-1)^{n-k}\chi_n(\mu_k) \ge 0, \tag{3.23}$$

and $[\chi_n(\mu_k) + 2\operatorname{Re}\beta]^2 - 4|\beta|^2 \ge 0$ and $(-1)^{n-k}(\chi_n(\mu_k) + 2\operatorname{Re}\beta) \ge 0$ do not equal zero simultaneously.

We now show that the inequality in (3.22) follows from (3.21) and (3.23). Indeed, the inequality (3.21) implies

$$(\chi_n(\mu_k) + 4\operatorname{Re}\beta)\chi_n(\mu_k) \ge 4(\operatorname{Im}\beta)^2 \ge 0,$$

 \mathbf{SO}

$$(-1)^{n-k}(\chi_n(\mu_k) + 4\operatorname{Re}\beta) \ge 0.$$

Therefore, if $(-1)^{n-k} \operatorname{Re} \beta < 0$, then

$$(-1)^{n-k}(\chi_n(\mu_k) + 2\operatorname{Re}\beta) > (-1)^{n-k}(\chi_n(\mu_k) + 4\operatorname{Re}\beta) \ge 0,$$

and if $(-1)^{n-k} \operatorname{Re} \beta \ge 0$, then

$$(-1)^{n-k}(\chi_n(\mu_k) + 2\operatorname{Re}\beta) \ge (-1)^{n-k}\chi_n(\mu_k) \ge 0.$$

Thus, we obtain that for any complex (real or nonreal) β and a_n , the eigenvalues of the matrices J_n and J_{n-1} satisfy the inequalities (3.21) and (3.23) that is equivalent to (3.16).

Let now $\beta \in \mathbb{R} \setminus \{0\}$, so $\operatorname{Re} \beta = \beta$ and $\operatorname{Im} \beta = 0$. In this case, the inequality (3.21) has the form

$$\chi_n(\mu_k)[\chi_n(\mu_k) + 4\beta] \ge 0,$$

$$\chi_n(\mu_k) = 0, \tag{3.24}$$

or

 \mathbf{SO}

$$|\chi_n(\mu_k)| \ge 4\beta(-1)^{n-k-1}.$$
 (3.25)

Moreover, for real β the eigenvalues of J_n and J_{n-1} satisfy the inequalities (3.11)–(3.12), which include (3.24).

It is easy to see that (3.11)–(3.12) and (3.25) imply (3.17), as required.

Remark 3.5. By the Hermite–Biehler theorem (see e.g. [24]), the condition $(-1)^{n-k}\chi_n(\mu_k) > 0$ in (3.16) can be changed to $\chi_n(\mu_k) \in \mathbb{R}$, since for $\operatorname{Im} a_n > 0$ ($\operatorname{Im} a_n < 0$) the polynomial $\chi_n(\lambda)$ has all roots in the open upper (lower) half-plane of the complex plane, and since $\chi_{n-1}(\lambda)$ is its imaginary part (see Remark 3.3).

Remark 3.6. The inequalities (3.17) were established in [26] under the assumption of reality of all the entries of the matrix J_n and $b_n > 0$.

Remark 3.7. If β is a pure imaginary number, that is, $\beta = i\gamma$, $\gamma \in \mathbb{R} \setminus \{0\}$, then the conditions (3.16) have the form

$$|\chi_n(\mu_k)| \ge 2|\gamma|, \quad k = 1, \dots, n-1.$$

Remark 3.8. The formulæ (3.4) and (2.6) also imply that

$$\alpha_{k} = \frac{\left[|\overline{b}_{n-1}u_{k,n-1}|^{2} + \operatorname{Re}(b_{n}u_{k1}b_{n-1}\overline{u}_{k,n-1})]^{2} + \left[\operatorname{Im}(b_{n}u_{k1}b_{n-1}\overline{u}_{k,n-1})\right]^{2}}{|\overline{b}_{n-1}u_{k,n-1}|^{2}} = \frac{\left[\chi_{n-1}'(\mu_{k})|\overline{b}_{n-1}u_{k,n-1}|^{2} + \operatorname{Re}\beta\right]^{2} + (\operatorname{Im}\beta)^{2}}{|\overline{b}_{n-1}u_{k,n-1}|^{2}\left[\chi_{n-1}'(\mu_{k})\right]^{2}}.$$
(3.26)

Hence from (3.4) and (3.26) we get that $X_k := |b_{n-1}u_{k,n-1}|^2$ satisfies the equation (3.18). Therefore, if $X_k^{(1,2)}$ are solutions of (3.18), then $X_k^{(1)} = |b_1u_{k1}|^2$, $X_k^{(2)} = |\overline{b}_{n-1}u_{k,n-1}|^2$, or $X_k^{(1)} = |\overline{b}_{n-1}u_{k,n-1}|^2$, $X_k^{(2)} = |b_1u_{k1}|^2$.

Finally, recall that the system of vectors $\mathbf{u}_1, \ldots, \mathbf{u}_{n-1}$ is orthonormal, so the matrix $U = [|u_{kj}|]_{k,j=1}^{n-1}$ is unitary. Therefore,

$$\sum_{k=1}^{n-1} |u_{k1}|^2 = 1$$

Since $X_k = |b_n u_{k1}|^2$, we have

$$\sum_{k=1}^{n-1} X_k^{(1,2)} = \sum_{k=1}^{n-1} |b_n u_{k1}|^2 = |b_n|^2 \sum_{k=1}^{n-1} |u_{k1}|^2 = |b_n|^2.$$
(3.27)

Thus, (3.20) and (3.27) imply

$$|b_n| = \left(\sum_{k=1}^{n-1} \frac{(-1)^{n-k} (\chi_n(\mu_k) + 2\operatorname{Re}\beta) \pm \sqrt{[\chi_n(\mu_k) + 2\operatorname{Re}\beta]^2 - 4|\beta|^2}}{2|\chi'_{n-1}(\mu_k)|}\right)^{\frac{1}{2}}, \qquad (3.28)$$

and

$$|u_{k1}|^{2} = \frac{(-1)^{n-k}(\chi_{n}(\mu_{k}) + 2\operatorname{Re}\beta) \pm \sqrt{[\chi_{n}(\mu_{k}) + 2\operatorname{Re}\beta]^{2} - 4|\beta|^{2}}}{2|b_{n}|^{2}|\chi_{n-1}'(\mu_{k})|} > 0, \qquad (3.29)$$

for k = 1, ..., n - 1.

By the same reasoning as above and by Remark 3.8 we obtain that

$$|b_{n-1}| = \left(\sum_{k=1}^{n-1} \frac{(-1)^{n-k} (\chi_n(\mu_k) + 2\operatorname{Re}\beta) \mp \sqrt{[\chi_n(\mu_k) + 2\operatorname{Re}\beta]^2 - 4|\beta|^2}}{2|\chi'_{n-1}(\mu_k)|}\right)^{\frac{1}{2}},$$
(3.30)

and

$$|u_{k,n-1}|^{2} = \frac{(-1)^{n-k}(\chi_{n}(\mu_{k}) + 2\operatorname{Re}\beta) \mp \sqrt{[\chi_{n}(\mu_{k}) + 2\operatorname{Re}\beta]^{2} - 4|\beta|^{2}}}{2|b_{n-1}|^{2}|\chi_{n-1}'(\mu_{k})|} > 0, \qquad (3.31)$$

for k = 1, ..., n - 1.

If $\beta \in \mathbb{R} \setminus \{0\}$, then the formulæ (3.28)–(3.29) can be represented in the following form:

$$|b_n| = \left(\sum_{k=1}^{n-1} \frac{(-1)^{n-k} (\chi_n(\mu_k) + 2\beta) \pm \sqrt{\chi_n(\mu_k)(\chi_n(\mu_k) + 4\beta)}}{2|\chi'_{n-1}(\mu_k)|}\right)^{\frac{1}{2}},$$
(3.32)

and

$$|u_{k1}|^2 = \frac{(-1)^{n-k}(\chi_n(\mu_k) + 2\beta) \pm \sqrt{\chi_n(\mu_k)(\chi_n(\mu_k) + 4\beta)}}{2|b_n|^2|\chi'_{n-1}(\mu_k)|} > 0,$$
(3.33)

for k = 1, ..., n - 1.

$$|b_{n-1}| = \left(\sum_{k=1}^{n-1} \frac{(-1)^{n-k} (\chi_n(\mu_k) + 2\beta) \mp \sqrt{\chi_n(\mu_k) (\chi_n(\mu_k) + 4\beta)}}{2|\chi'_{n-1}(\mu_k)|}\right)^{\frac{1}{2}},$$
(3.34)

and

$$|u_{k,n-1}|^2 = \frac{(-1)^{n-k}(\chi_n(\mu_k) + 2\beta) \mp \sqrt{\chi_n(\mu_k)(\chi_n(\mu_k) + 4\beta)}}{2|b_{n-1}|^2|\chi'_{n-1}(\mu_k)|} > 0,$$
(3.35)

for k = 1, ..., n - 1.

4 Inverse problems for matrices in \mathcal{J}_n

In this section, we show that the necessary conditions (3.16)–(3.17) on the spectra of the matrices in the class \mathcal{J}_n are also sufficient. This means that given 2n numbers $\lambda_1, \ldots, \lambda_n, \mu_1, \ldots, \mu_{n-1}, \beta$ satisfying (3.16) or (3.17) we can reconstruct a matrix $J_n \in \mathcal{J}_n$ whose eigenvalues are $\lambda_1, \ldots, \lambda_n$, the eigenvalues of its leading principal submatrix J_{n-1} are μ_1, \ldots, μ_{n-1} , and $\beta = b_1 \cdots b_n$.

But any matrix in the class \mathcal{J}_n has 3n + 1 parameters, namely, c_k , $k = 1, \ldots, n - 1$, Re a_n , Im a_n , Re b_j , Im b_j , $j = 1, \ldots, n$. At the same time, the spectral data $\lambda_1, \ldots, \lambda_n, \mu_1, \ldots, \mu_{n-1}$, β consist of only 2n or 2n + 1 parameters. This is obvious if all the λ_j are real. But if the λ_j are nonreal, they depend on the numbers μ_k , since the numbers $\prod_{j=1}^n (\mu_k - \lambda_j)$ are real by (3.16). Thus, if we want to reconstruct finitely many matrices from the spectral data mentioned above, we should restrict ourselves to a subclass of the class \mathcal{J}_n .

Thus, in this section, we consider the subclass $\widehat{\mathcal{J}}_n$ of the class \mathcal{J}_n consisting of matrices of the form

$$\hat{J}_{n} = \begin{pmatrix} \hat{c}_{1} & \hat{b}_{1} & 0 & \dots & 0 & \hat{b}_{n} \\ \hat{b}_{1} & \hat{c}_{2} & \hat{b}_{2} & \dots & 0 & 0 \\ 0 & \hat{b}_{2} & \hat{c}_{3} & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & \hat{c}_{n-1} & \hat{b}_{n-1} \\ \hat{b}_{n} & 0 & 0 & \dots & \hat{b}_{n-1} & \hat{a}_{n} \end{pmatrix},$$

$$(4.1)$$

with real $\hat{c}_k, \hat{b}_k, k = 1, ..., n-1$ and complex (real or nonreal) \hat{a}_n and \hat{b}_n . This subclass has the following important property.

Lemma 4.1. For any matrix in the class \mathcal{J}_n , there exists a matrix in the class $\widehat{\mathcal{J}}_n$ with the same spectral data, that is, with the same spectra of the matrix itself and of its leading $(n-1) \times (n-1)$ submatrix, and the same number β (see (2.7)).

Proof. Indeed, by (3.15) the spectrum of a matrix J_n in the class \mathcal{J}_n depends on a_n , μ_k , and α_k , $k = 1, \ldots, n-1$, where μ_k are the eigenvalues of the leading principal submatrix J_{n-1} . At the same time, by (3.10) the numbers α_k depend on μ_k , β , and $|b_1 u_{k1}|^2$, where β is defined in (2.7).

Thus, given a matrix J_n in the class \mathcal{J}_n , we must find a matrix \widehat{J}_n in the class $\widehat{\mathcal{J}}_n$ with the same a_n , β , the spectrum of the submatrix \widehat{J}_{n-1} , and the same numbers $|b_1u_{k1}|^2$.

So we consider a matrix

$$J_n = \begin{pmatrix} c_1 & b_1 & 0 & \dots & 0 & \overline{b}_n \\ \overline{b}_1 & c_2 & b_2 & \dots & 0 & 0 \\ 0 & \overline{b}_2 & c_3 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & c_{n-1} & b_{n-1} \\ b_n & 0 & 0 & \dots & \overline{b}_{n-1} & a_n \end{pmatrix}$$

in the class \mathcal{J}_n and construct the matrix

$$\widehat{J}_{n} = \begin{pmatrix} \widehat{c}_{1} & \widehat{b}_{1} & 0 & \dots & 0 & \overline{b}_{n} \\ \widehat{b}_{1} & \widehat{c}_{2} & \widehat{b}_{2} & \dots & 0 & 0 \\ 0 & \widehat{b}_{2} & \widehat{c}_{3} & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & \widehat{c}_{n-1} & \widehat{b}_{n-1} \\ \widehat{b}_{n} & 0 & 0 & \dots & \widehat{b}_{n-1} & \widehat{a}_{n} \end{pmatrix},$$

such that

$$\hat{c}_j = c_j, \qquad j = 1, \dots, n-1,$$

$$\hat{b}_k = |b_k|, \qquad k = 1, \dots, n-1,$$

$$\hat{a}_n = a_n,$$
(4.2)

and

$$\widehat{b}_n = \frac{b_1 b_2 \cdots b_n}{|b_1 b_2 \cdots b_{n-1}|} = \frac{\beta}{\widehat{b}_1 \widehat{b}_2 \cdots \widehat{b}_{n-1}}.$$
(4.3)

Obviously, $\widehat{J}_n \in \widehat{\mathcal{J}}_n$.

The formulæ (4.2) show that the spectra of the submatrices J_{n-1} and \hat{J}_{n-1} coincide (see [14]):

$$\sigma(J_n) = \sigma(J_n) = \{\mu_1, \dots, \mu_{n-1}\}$$

Moreover, from (4.3) we have

$$\beta = b_1 b_2 \cdots b_n = \widehat{b}_1 \widehat{b}_2 \cdots \widehat{b}_n = \widehat{\beta}.$$

Therefore, to establish that $\sigma(\widehat{J}_n) = \sigma(J_n)$ it suffices to prove that

$$\left(\widehat{b}_1\widehat{u}_{k1}\right)^2 = |b_1u_{k1}|^2, \qquad k = 1, \dots, n-1.$$
 (4.4)

The formula (2.5) and the equalities (4.2) imply that

$$\sum_{k=1}^{n-1} \frac{|b_1 u_{k1}|^2}{\lambda - \mu_k} = |b_1|^2 \mathbf{e}_1^T (\lambda I_{n-1} - J_{n-1})^{-1} \mathbf{e}_1 =$$

$$= \frac{|b_1|^2 \det(\lambda I_{n-2} - J_{n-2})}{\det(\lambda I_{n-1} - J_{n-1})} = \frac{\widehat{b}_1^2 \det(\lambda I_{n-2} - \widehat{J}_{n-2})}{\det(\lambda I_{n-1} - \widehat{J}_{n-1})} =$$

$$= \widehat{b}_1^2 \mathbf{e}_1^T (\lambda I_{n-1} - \widehat{J}_{n-1})^{-1} \mathbf{e}_1 = \sum_{k=1}^{n-1} \frac{\left(\widehat{b}_1 \widehat{u}_{k1}\right)^2}{\lambda - \mu_k},$$

so (4.4) holds, as required.

Now we establish that the conditions (3.16)–(3.17) on the spectral data are also sufficient to reconstruct finitely many matrices in the class $\widehat{\mathcal{J}}_n$ with this spectral data. We consider 4 cases.

Theorem 4.2. Let β be a nonreal number, and let $\{\mu_k\}_{k=1}^{n-1}$ be a set of real distinct numbers. Suppose that $\{\lambda_j\}_{j=1}^n$ is a set of complex numbers such that $\operatorname{Im} \lambda_j > 0$, $j = 1, \ldots, n-1$. Then there exists a matrix \widehat{J}_n in the class $\widehat{\mathcal{J}}_n$ with $\operatorname{Im} \widehat{a}_n > 0$ and $\beta = \widehat{b}_1 \cdots \widehat{b}_n$ such that $\sigma(\widehat{J}_n) = \{\lambda_1, \ldots, \lambda_n\}$ and $\sigma(\widehat{J}_{n-1}) = \{\mu_1, \ldots, \mu_{n-1}\}$ if and only if

$$(-1)^{n-k}\chi_n(\mu_k) > 0 \quad and \quad |\chi_n(\mu_k) + 2\operatorname{Re}\beta| \ge 2|\beta|, \quad k = 1, \dots, n-1,$$
 (4.5)

where $\chi_n(\lambda) = \prod_{j=1}^n (\lambda - \lambda_j).$

Moreover, if there are $m \ (0 \leq m \leq n-1)$ equalities in (4.5), then there exist exactly 2^{n-m-1} matrices in the class $\widehat{\mathcal{J}}_n$ with this spectral data.

Proof. The necessity of the conditions (4.5) is provided by Theorem 3.4.

Suppose that we have a nonreal number $\hat{\beta}$, distinct real numbers $\{\mu_k\}_{k=1}^{n-1}$, and the numbers $\{\lambda_j\}_{j=1}^n$ in the open upper half-plane, and suppose that they satisfy (4.5).

We also assume that there are exactly $m, 0 \leq m \leq n-1$, equalities in (4.5). Then one has exactly 2^{n-m-1} ways to construct the number $|\hat{b}_n|$ and the corresponding values $|u_{k1}|^2$, $k = 1, \ldots, n-1$, by the formulæ (3.28) and (3.29). The conditions (4.5) ensure positivity of the obtained numbers $|\hat{b}_n|$ and $|u_{k1}|^2$, $k = 1, \ldots, n-1$. Indeed, (4.5) imply positivity of the numbers $(-1)^{n-k}[\chi_{n-1}(\mu_k) + 2\operatorname{Re}\beta]$, $k = 1, \ldots, n-1$, as we established in the proof of Theorem 3.4. At the same time, positivity of these numbers and the conditions (4.5) obviously imply positivity of the numbers $X_k^{(1,2)}$ defined in (3.20). Thus, the conditions (4.5) guaranty the existence of positive numbers $|\hat{b}_n|$ and $|u_{k1}|^2$, $k = 1, \ldots, n-1$.

With the values $|u_{k1}|^2$, k = 1, ..., n - 1, we construct the rational function

$$\sum_{k=1}^{n-1} \frac{|u_{k1}|^2}{\lambda - \mu_k} := \frac{\psi(\lambda)}{\chi_{n-1}(\lambda)},\tag{4.6}$$

where the polynomial $\psi(\lambda)$ of degree n-2 is uniquely determined by the numbers $|u_{k1}|^2$ and μ_k , $k = 1, \ldots, n-1$. Moreover, since $|u_{k1}|^2 > 0$ by construction, the function $\psi(\lambda)/\chi_{n-1}(\lambda)$ maps the upper half-plane to the lower half-plane of the complex plane, so the zeroes of ψ are real and simple and interlace the zeroes of χ_{n-1} .

If we know the function (4.6), we can always reconstruct a unique Jacobi matrix of the form

$$\begin{pmatrix} \widehat{c}_{1} & \widehat{b}_{1} & 0 & \dots & 0 & 0\\ \widehat{b}_{1} & \widehat{c}_{2} & \widehat{b}_{2} & \dots & 0 & 0\\ 0 & \widehat{b}_{2} & \widehat{c}_{3} & \dots & 0 & 0\\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots\\ 0 & 0 & 0 & \dots & \widehat{c}_{n-2} & \widehat{b}_{n-2}\\ 0 & 0 & 0 & \dots & \widehat{b}_{n-2} & \widehat{c}_{n-1} \end{pmatrix} = \widehat{J}_{n-1},$$

$$(4.7)$$

where $\hat{b}_1, \ldots, \hat{b}_{n-1} > 0, \hat{c}_1, \ldots, \hat{c}_{n-1} \in \mathbb{R}$. There exist a few algorithms to make such a reconstruction [20, 19, 16, 15].

Thus, given the numbers μ_k and $|u_{k1}|^2$ the rational function (4.6) uniquely determines the matrix J_{n-1} whose eigenvalues are μ_k , while $\psi(\lambda) = \chi_{n-2}(\lambda)$. Furthermore, we put

$$\widehat{a}_n := \left(\sum_{j=1}^n \lambda_j - \sum_{k=1}^{n-1} \mu_k\right)$$
$$\widehat{b}_n := |\widehat{b}_n| e^{i \arg \beta}$$
$$\widehat{b}_n \downarrow := \frac{\beta}{\beta}$$

and

$$\widehat{b}_{n-1} := \frac{\beta}{\widehat{b}_1 \cdots \widehat{b}_{n-2} \widehat{b}_n}.$$

So we constructed a matrix of the form (4.1) such that $\sigma(\hat{J}_{n-1}) = \{\mu_1, \ldots, \mu_{n-1}\}$ and $\hat{b}_1 \cdots \hat{b}_n = \beta$. To finish the proof it suffices to establish that $\sigma(\hat{J}_n) = \{\lambda_1, \ldots, \lambda_n\}$. To do this we consider the following formula obtained from (3.1)–(3.2)

$$\frac{\det(\lambda I_n - \hat{J}_n)}{\det(\lambda I_{n-1} - \hat{J}_{n-1})} = \lambda + \hat{a}_n - \sum_{k=1}^{n-1} \frac{|\hat{b}_n u_{k1} + \overline{\hat{b}}_{n-1} u_{k,n-1}|^2}{\lambda - \mu_k},$$
(4.8)

where u_{k1} and $u_{k,n-1}$ are the first and the last entries of the orthonormal eigenvector \mathbf{u}_k of the submatrix \widehat{J}_{n-1} corresponding to the eigenvalue μ_k , $k = 1, \ldots, n-1$. These entries are related as in (2.6), and u_{k1} satisfies (3.29) by construction. Then the numbers \widehat{b}_{n-1} and $|u_{k,n-1}|^2$, $k = 1, \ldots, n-1$, must satisfy the equalities (3.30)–(3.31). Thus $|\widehat{b}_n u_{k1}|^2$ and $|\overline{b}_{n-1} u_{k,n-1}|^2$ are the solutions of the equation (3.19) so we have

$$|\widehat{b}_n u_{k1}|^2 + |\overline{\widehat{b}}_{n-1} u_{k,n-1}|^2 = -\frac{\chi_n(\mu_k) + 2\operatorname{Re}\beta}{\chi'_{n-1}(\mu_k)}.$$

This formula together with (2.6) gives us

$$\widehat{b}_{n}u_{k1} + \overline{\widehat{b}}_{n-1}u_{k,n-1}|^{2} = |\widehat{b}_{n}u_{k1}|^{2} + |\overline{\widehat{b}}_{n-1}u_{k,n-1}|^{2} + 2\operatorname{Re}(\widehat{b}_{n}u_{k1}\widehat{b}_{n-1}\overline{u}_{k,n-1}) = -\frac{\chi_{n}(\mu_{k})}{\chi_{n-1}'(\mu_{k})} + \frac{2\operatorname{Re}\beta}{\chi_{n-1}'(\mu_{k})} = -\frac{\prod_{j=1}^{n}(\mu_{k} - \lambda_{j})}{\prod_{\substack{r=1\\r \neq k}}^{n-1}(\mu_{k} - \mu_{r})}$$

so det $(\lambda I_n - \widehat{J}_n) = \prod_{j=1}^n (\lambda - \lambda_j)$, as required.

Theorem 4.3. Let β be a nonreal number, and let $\{\mu_k\}_{k=1}^{n-1}$ and $\{\lambda_j\}_{j=1}^n$ be sets of distinct real numbers with no common elements.

Then there exists a matrix \widehat{J}_n in the class $\widehat{\mathcal{J}}_n$ with $\widehat{a}_n \in \mathbb{R}$ and $\beta = \widehat{b}_1 \cdots \widehat{b}_n$ such that $\sigma(\widehat{J}_n) = \{\lambda_1, \ldots, \lambda_n\}$ and $\sigma(\widehat{J}_{n-1}) = \{\mu_1, \ldots, \mu_{n-1}\}$ if and only if the conditions (4.5) hold.

Moreover, if there are $m \ (0 \leq m \leq n-1)$ equalities in (4.5), then there exist exactly 2^{n-m-1} matrices in the class $\widehat{\mathcal{J}}_n$ with these spectral data.

For real nonzero β we have the following results.

Theorem 4.4. Let β be a real nonzero number, and let $\{\mu_k\}_{k=1}^{n-1}$ be a set of real distinct numbers. Suppose that $\{\lambda_j\}_{j=1}^n$ is a set of complex numbers in the **closed** upper half-plane of the complex plane, $\operatorname{Im} \lambda_j \ge 0, j = 1, \ldots, n-1$.

Then there exists a matrix \widehat{J}_n in the class $\widehat{\mathcal{J}}_n$ with $\operatorname{Im} \widehat{a}_n > 0$ and $\beta = \widehat{b}_1 \cdots \widehat{b}_n$ such that $\sigma(\widehat{J}_n) = \{\lambda_1, \dots, \lambda_n\}$ and $\sigma(\widehat{J}_{n-1}) = \{\mu_1, \dots, \mu_{n-1}\}$ if and only if

 $(-1)^{n-k}\chi_n(\mu_k) \ge 0, \quad k = 1, \dots, n-1,$ (4.9)

and

$$|\chi_n(\mu_k)| \ge 4(-1)^{n-k-1}\beta, \quad k = 1, \dots, n-1,$$
(4.10)

where $\chi_n(\lambda) = \prod_{j=1}^n (\lambda - \lambda_j).$

Moreover, if there are exactly m_1 equalities in (4.9) and exactly m_2 equalities in (4.10) ($0 \leq m_1 + m_2 \leq n-1$), then there exist exactly $2^{n-m_1-m_2-1}$ matrices in the class $\hat{\mathcal{J}}_n$ with these spectral data.

Theorem 4.5. Let β be a real number, and let $\{\mu_k\}_{k=1}^{n-1}$ and $\{\lambda_j\}_{j=1}^n$ be sets of distinct real numbers that may have common elements.

Then there exists a matrix \widehat{J}_n in the class $\widehat{\mathcal{J}}_n$ with $\widehat{a}_n \in \mathbb{R}$ and $\beta = \widehat{b}_1 \cdots \widehat{b}_n$ such that $\sigma(\widehat{J}_n) = \{\lambda_1, \ldots, \lambda_n\}$ and $\sigma(\widehat{J}_{n-1}) = \{\mu_1, \ldots, \mu_{n-1}\}$ if and only if the numbers $\{\mu_k\}_{k=1}^{n-1}$ and $\{\lambda_j\}_{j=1}^n$ satisfy the inequalities (4.9)–(4.10).

Moreover, if there are exactly m_1 equalities in (4.9) and exactly m_2 equalities in (4.10) ($0 \leq m_1 + m_2 \leq n-1$), then there exist exactly $2^{n-m_1-m_2-1}$ matrices in the class $\hat{\mathcal{J}}_n$ with these spectral data.

Theorems 4.3–4.5 can be proved in the same way as we proved Theorem 4.2. Note that the numbers m_1 and m_2 do not exceed a half of n (approximately). The exact upper bounds on m_1 and m_2 depend on the parity of n and on the sign of β , and can be obtained from the inequalities (4.9)–(4.10).

Theorem 4.5 was established in [26, Theorems 6–7]. As well as in [26, Corollary 8] we note that in Theorems 4.2–4.5 the constructed matrix \hat{J}_n is unique if and only if m = n-1 or $m_1+m_2 = n-1$. This is possible for specific distributions of the λ_j and the μ_k .

5 Conclusions

In this paper, we showed that the technique developed by Y.-H. Xu and E.-X. Jiang [26] for periodic Jacobi matrices can be extended to a class of complex band matrices. We also gave some new explanations of the technique using the theory of the location of roots of polynomials. Our results extend the results of Arlinskii and Tsekhanovskii [3] (see also [4]) who studied a one-dimensional imaginary perturbation of symmetric real Jacobi matrices.

We believe that the combination of the methods of the works [26] and [4] can be helpful to study direct and inverse spectral problems of some other classes of band matrices.

References

- [1] N.L. Akhiezer, The Classical Momentum Problem, Hafner, New York, 1965 (Engl. Transl.).
- [2] S.A. Andrea and T.G. Berry, Continued fractions and periodic Jacobi matrices, *Linear Algebra Appl.* 161, 1992, 117–134.
- [3] Y. Arlinskii and E. Tsekhanovskii, Non-self-adjoint Jacobi matrices with a rank-one imaginary part, Journal of Func. Anal. 241, 2006, 383–438.
- [4] Y. Arlinskii, S. Belyi, and E. Tsekhanovskii, Conservative realizations of Herglotz-Nevanlinna functions, Operator Theory: Advances and Appl. 217, 2011.
- [5] Yu. S. Barkovsky, Lectures on the Routh-Hurwitz problem. arXiv.org: math/0802.1805, 2008.
- [6] N. Bebiano and J. da Providência, Inverse eigenvalue problems for pseudo-Jacobi matrices: existence and uniqueness, *Inverse Problems* 27, no. 2, 2011, 025005, 12pp.
- [7] N. Bebiano, C.M. da Fonseca, J. da Providência, An inverse eigenvalue problem for periodic Jacobi matrices in Minkowski spaces, *Linear Algebra Appl.* 435, 2011, 2033–2045.
- [8] B. Beckerman, Complex Jacobi matrices, J. Comp. Appl. Math. 127, no. 1-2, 2001, 17–65.
- B. Beckerman, On the classification of the spectrum of second order difference operator, Math. Nachr. 916, 2000, 45–59.
- [10] D. Boley and G.H. Golub, Inverse eigenvalue problems for band matrices, *Lecture Notes in Math.* 630, Springer, Berlin, 1978, 23–31.
- [11] D. Boley and G.H. Golub, A modified method for reconstructing periodic Jacobi matrices, Math. Comp. 42, no. 165, 1984, 143–150.
- [12] C. de Boor and G.H. Golub, The numerically stable reconstruction of a Jacobi matrix from the spectral data, *Linear Algebra Appl.* 21, 1978, 245–260.
- [13] W.E. Ferguson, The construction of Jacobi and periodic Jacobi matrices with prescribed spectra, Math. Comp. 35, 1980, 1203–1220.
- [14] Ф. Р. Гантмахер, М. Г. Крейн. Осцилляционные матрицы и ядра и малые колебания механических систем. (Russian), Москва-Ленинград, ГТТИ, 1941.

English transl.: F. Gantmacher, M. Krein., Oscillation matrices and kernels and small vibrations of mechanical systems. AMS Chelsea Publishing, Providence, RI, revised edition, 2002. Translation based on the 1941 Russian original. Edited and with a preface by Alex Eremenko.

- [15] F. Gesztesy, B. Simon, m-functions and inverse spectral analysis for finite and semi-infinite Jacobi matrices, *Journal d'Analyse Mathématique*, **73** (1997), 267-297.
- [16] G.M.L. Gladwell, *Inverse Problems in Vibrations*, Monographs and Textbooks on Mechanics of Solids and Fluids: Mechanics. Dynamical Systems, 9, Martinus Nijhoff Publishers, Dordrecht, 1986.
- [17] L.J. Gray and D.G. Wilson, Construction of a Jacobi matrix from spectral data, *Linear Algebra Appl.* 14, 1976, 131–134.
- [18] G. Sh. Guseinov, An inverse spectral problem for complex Jacobi matrices, Commun. Nonlinear Sci. Numer. Simulat. 15, 2010, 840–851.
- [19] O. Hald, Inverse eigenvalue problems for Jacobi matrices, Linear Algebra Appl. 14, 1976, 63–85.
- [20] H. Hochstadt, On the construction of a Jacobi matrix from spectral data, *Linear Algebra Appl.* 8, 1974, 435-446.
- [21] O. Holtz and M. Tyaglov, Structured matrices, continued fractions, and root localization of polynomials, SIAM Rev. 54, no. 3, 2012, 421–509.
- [22] M. Marden, Geometry of polynomials, Math. Servey Monographs 3, Amer. Math. Soc., Providence, RI, 1966.
- [23] P. van Moerbeke, The spectrum of Jacobi matrices, Invent. Math. 37, no. 1, 1976, 45–81.
- [24] Obreschkoff N., Verteilung und Berechnung der Nullstellen reeller Polynome, VEB Deutscher Verlag der Wissenschaften, Berlin, 1963.
- [25] G. Teschl, Jacobi Operators and Completely Integrable Nonlinear Lattices, Math. Surv. and Mon. 72, Amer. Math. Soc., Rhode Island, 2000.

- [26] Y.-H. Xu and E.-X. Jiang, An inverse eigenvalue problem for periodic Jacobi matrices, *Inverse Problems* 23, no. 1, 2007, 165–181.
- [27] H.S.Wall, Analytic Theory of Continued Fractions, Chelsea, Bronx NY, 1973.
- [28] Y.-H Xu, E.-X. Jiang, An algorithm for an inverse eigenvalue problem on Jacobi matrices, J. Shangai Univ. 12, no. 4, 2008, 289–293.
- [29] F. Zhang (Ed.), The Schur complement and its applications, Numerical Methods and Algorithms 4, Springer-Verlag, New York, 2005.