

A REMARK ON THE REEB FLOW FOR SPHERES

ROGER CASALS AND FRANCISCO PRESAS

ABSTRACT. We produce homotopically non-trivial 2-spheres in the space of contact structures of a hypercontact manifold. In particular, this proves the non-triviality of the Reeb flow for the standard contact spheres \mathbb{S}^{2n+1} , $n \neq 3$, inside the fundamental group of their contactomorphism group.

Let (M, ξ) be a contact manifold. Consider the space $\mathcal{C}(M, \xi)$ of contact structures isotopic to ξ . This space has been studied in special cases. See [El] for the 3-sphere and [Bo], [G2] for torus bundles. In the present note we prove the non-triviality of its second homotopy group for the class of manifolds introduced in [G1]:

Theorem 1. *Let (M, ξ) be a hypercontact manifold, then $\text{rk}(\pi_2(\mathcal{C}(M, \xi))) \geq 1$.*

Let $(\mathbb{S}^{4n+3}, \xi_0)$ be the standard contact sphere. Such non-trivial spheres in $\mathcal{C}(\mathbb{S}^{4n+3}, \xi_0)$ allow us to answer a question posed in [Gi]:

Remarque 2.10: On peut se demander s'il n'y a pas, dans $\text{Cont}(\mathbb{S}^{2n+1}, \xi_0)$, un lacet positif contractile plus simple que dans $\mathbb{P}U(n, 1)$ et par exemple si le lacet ρ_t , $t \in \mathbb{S}^1$, n'est pas contractile. C'est peu probable mais je n'en ai pas la preuve.

The answer we provide is the following

Corollary 2. *The class in $\pi_1(\text{Cont}(\mathbb{S}^{2n+1}, \xi_0))$ generated by the Reeb flow is a non-trivial element of infinite order for $n \neq 3$.*

In Section 1 we introduce the objects of interest and necessary notation. The geometric construction underlying the results is explained in Section 2. Theorem 1 is concluded. Section 3 contains the argument deducing Corollary 2. Section 4 extends the results to higher homotopy groups.

1. PRELIMINARIES

1.1. Contact structures.

Definition 3. *Let M^{2n+1} be a smooth manifold. A codimension-1 regular distribution ξ is a contact distribution if there exists a 1-form $\alpha \in \Omega^1(M)$ such that $\ker \alpha = \xi$ and $\alpha \wedge d\alpha^n$ is a volume form.*

The structure described above is known as a cooriented contact structure. Since the non-coorientable case is not considered in this article, we refer to a cooriented contact structure simply as a contact structure. The smooth manifold M will be assumed to be oriented. The contact structures to be considered will be positively cooriented, i.e. the induced orientation coincides with that prescribed on M .

The definition is independent of the choice of 1-form $\alpha' = e^f \alpha$, for any $f \in C^\infty(M, \mathbb{R})$. Let $\text{Cont}(M, \xi) = \{s \in \text{Diff}(M) : ds_* \xi = \xi\}$ be the space of diffeomorphisms that preserve the contact structure. These diffeomorphisms are called contactomorphisms. The connected component of the identity of $\text{Cont}(M, \xi)$ will be denoted by $\text{Cont}_0(M, \xi)$. $\mathcal{C}(M, \xi)$ will stand for the space of positive contact structures in M isotopic to ξ . The unique vector field R such that

$$i_R \alpha = 1, \quad i_R d\alpha = 0,$$

is called the Reeb vector field associated to α .

2010 *Mathematics Subject Classification.* Primary: 53D10.
Key words and phrases. contact structures, Reeb flow.

A vector field $X \in \Gamma(TM)$ preserves the contact structure if it satisfies the following pair of equations

$$\begin{aligned} i_X \alpha &= H, \\ i_X d\alpha &= -dH + (i_R dH)\alpha, \end{aligned}$$

for a choice of α and a function $H \in C^\infty(M, \mathbb{R})$. Such a function is called the Hamiltonian associated to the vector field. This correspondance defines a linear isomorphism between the space of vector fields $\Gamma_\xi(TM)$ preserving the contact structure ξ and the vector space of smooth functions $C^\infty(M, \mathbb{R})$. By definition, a contactomorphism $\phi \in \text{Cont}_0(M, \xi)$ admits an expression as $\phi = \phi_1$ for a time dependent flow $\{\phi_t\}_{t \in [0,1]}$ generated by a time dependent family $X_t \in \Gamma_\xi(TM)$. Therefore, its flow $\{\phi_t\}$ can be generated by a time dependent family of smooth functions $\{H_t\}$.

1.2. Contact fibrations. A smooth fibration $\pi : X \rightarrow B$ is said to be contact for a codimension-1 distribution $\xi \subset TX$ if for any fiber $F_p = \pi^{-1}(p) \xrightarrow{e} X$, the restriction of the distribution $e^*\xi$ is a contact structure on the fibre.

Let $\pi : X \rightarrow B$ be a smooth fibration. The vertical subbundle $V \subset TX$ is defined fiberwise by $V_x = \ker d\pi(x), \forall x \in X$. An *Ehresmann connection* is a smooth choice of a fiberwise complementary linear space H_x for V_x inside $T_x X$. Therefore, the map $d\pi_x : H_x \rightarrow TB_{\pi(x)}$ is a linear isomorphism and there is a well-defined notion of parallel transport.

There is a canonical connection once a contact fibration $(\pi, \xi = \ker \alpha)$ is fixed. The connection H is defined at a point $x \in X$ to be the annihilator of the vector subspace $V_x \cap \xi_x$ with respect to the quadratic form $(\xi, d\alpha)$. It is complementary to V_x since $V_x \cap \xi_x$ is a symplectic space for the 2-form $d\alpha$. The connection is independent of the choice of form α . See [Pr] for details on the following facts.

Lemma 4. *The parallel transport of the canonical connection associated to a contact fibration is by contactomorphisms.*

Lemma 5. *Let $(F, \ker \alpha_0)$ be a closed contact manifold. Let $\pi : F \times \mathbb{D}^2 \rightarrow \mathbb{D}^2$ be a contact fibration with fibration distribution defined by the kernel of $\alpha = \alpha_0 + Hd\theta$, for some function $H : F \times \mathbb{D}^2 \rightarrow \mathbb{R}$ satisfying $|H| \in O(r^2)$. Fix a loop $\gamma : \mathbb{S}^1 \rightarrow \mathbb{D}^2$, defined as $\gamma(\theta) = \gamma(r_0, \theta)$ in polar coordinates. Then, the contactomorphism of the fiber $F \times (r_0, 0)$ defined by the parallel transport along γ is generated by the family of Hamiltonian functions $\{G_\theta(p) = -H(p, r_0, \theta)\}_{\theta \in [0, 2\pi]}$.*

Let us study general contact fibrations over a 2-disk \mathbb{D}^2 . Fix a contact fibration $\pi : X \rightarrow \mathbb{D}^2$ with distribution $\xi = \ker \alpha$. Consider the radial vector field $Y = \partial_r$, defined on $\mathbb{D}^2 \setminus \{0\}$. It can be lifted to X by using the canonical contact connection. This produces a vector field $\tilde{Y} : Y \setminus F_0 \rightarrow TY$. Once an angle θ_0 is fixed it can be uniquely extended to $0 \in \mathbb{D}^2$. In such a case, denote by $\phi_{r, \theta_0} : F_0 \rightarrow F_{(r, \theta_0)}$ the associated flow at time r . It identifies via contactomorphisms the fibers over $0 \in \mathbb{D}^2$ and over $(r, \theta_0) \in \mathbb{D}^2$. Define the diffeomorphism:

$$\begin{aligned} \Phi : F_0 \times D^2 &\rightarrow X \\ (p, r, \theta) &\mapsto \phi_{r, \theta}(p). \end{aligned}$$

Then the definition of the contact connection implies $\Phi^* \alpha = e^g(\alpha_0 + Hd\theta)$, where $g : M \times D^2 \rightarrow \mathbb{R}$ and $H : M \times D^2 \rightarrow \mathbb{R}$ are arbitrary smooth functions. We can choose as fibration form $\alpha' = e^{-g}\alpha$ and trivialize the fibration using Φ . Then we obtain the expression

$$(1) \quad \Phi^* \alpha' = (\alpha_0 + Hd\theta).$$

Given a contact fibration over the disk, the trivialization constructed above is called *radial*. It is convenient to observe that the radial trivialization construction can be made parametric for families of contact fibrations over the disk.

1.3. Loops at infinity. Fix a contact fibration $\pi : X \rightarrow \mathbb{S}^2$ with distribution ξ , fibre F and a point $N \in \mathbb{S}^2$. This point will be referred as North pole or infinity. Define the restriction fibration $\pi_N : X \setminus \pi^{-1}(N) \rightarrow \mathbb{S}^2 \setminus N \simeq \mathbb{D}^2$. Trivialize the contact fibration π_N radially from $S = \{0\} \in \mathbb{D}^2$ to obtain a new contact fibration $\hat{\pi} : F \times \mathbb{D}^2 \rightarrow \mathbb{D}^2$ with fibration form $\alpha_0 + Hd\theta$. Denoting by

$\Phi : F \times \mathbb{D}^2 \rightarrow X \setminus \pi^{-1}(N)$ the trivialization map, we obtain $\Phi^*\xi = \ker \alpha_0 + Hd\theta$. Therefore, the map is connection-preserving. Consider the family of loops

$$\begin{aligned} \gamma_r : \mathbb{S}^1 &\longrightarrow D^2 \\ \theta &\longmapsto (r, \theta). \end{aligned}$$

Composing with the embedding $\mathbb{D}^2 \hookrightarrow \mathbb{S}^2$, for $r \rightarrow 1$, they are smaller and smaller loops around the North pole $N \in \mathbb{S}^2$. By Lemma 5, the parallel transport associated to the loop γ_r is generated by a family of Hamiltonians $\{G_\theta^r\}_{\theta \in \mathbb{S}^1}$, defined by $G_\theta^r(p) = -H(p, r, \theta)$. The limit function

$$G_\theta = \lim_{r \rightarrow 1} G_\theta^r$$

exists because the connection associated to ξ is a smooth connection over \mathbb{S}^2 . It is clear that $\{G_\theta\}$ defines a loop in $\text{Cont}(M, \xi_0 = \ker \alpha_0)$. This will be called the *loop at infinity* associated to (π, ξ) . Continuous families of contact fibrations with marked fibre produce continuous families of loops at infinity.

Definition 6. A contact sphere is a smooth map $e : \mathbb{S}^2 \rightarrow \mathcal{C}(M, \xi)$.

There is a canonical contact fibration over \mathbb{S}^2 associated to any contact sphere e . It is defined as

$$X = M \times \mathbb{S}^2 \rightarrow \mathbb{S}^2,$$

with the distribution at $(p, z) \in M \times \mathbb{S}^2$ being $\xi^e(p, z) = e(z)_p \oplus T_z \mathbb{S}^2 \subset T_p M \oplus T_z \mathbb{S}^2$.

Denote by $C^\infty(\mathbb{S}^2, \mathcal{C}(M, \xi))$ the space of smooth maps from \mathbb{S}^2 to $\mathcal{C}(M, \xi)$. The smooth loop space of $\text{Cont}_0(M, \xi)$ is denoted as $\Omega(\text{Cont}_0(M, \xi), id)$.

Lemma 7. The previous construction induces a continuous map

$$C^\infty(\mathbb{S}^2, \mathcal{C}(M, \xi)) \rightarrow \Omega(\text{Cont}_0(M, \xi), id).$$

Therefore, it provides a morphism

$$\pi_2(\mathcal{C}(M, \xi)) \rightarrow \pi_1(\text{Cont}_0(M, \xi)).$$

1.4. Homotopy sequence. The group $\text{Diff}_0(M)$ acts transitively on $\mathcal{C}(M, \xi)$ because of the Gray's Stability Theorem. It is a Serre fibration with homotopy fibre $\text{Cont}(M, \xi) \cap \text{Diff}_0(M)$. This homotopy fibre might be disconnected. Its identity component is $\text{Cont}_0(M, \xi)$. Hence the fibration induces a long exact sequence

$$(2) \quad \dots \rightarrow \pi_2(\text{Diff}_0(M)) \rightarrow \pi_2(\mathcal{C}(M, \xi)) \xrightarrow{\partial_2} \pi_1(\text{Cont}_0(M, \xi)) \rightarrow \pi_1(\text{Diff}_0(M)) \rightarrow \dots$$

The map ∂_2 is the one provided by Lemma 7. The study of this sequence will provide Corollary 2.

Note that the geometric lifting map

$$(3) \quad \pi_j(\mathcal{C}(M, \xi)) \xrightarrow{\partial_j} \pi_{j-1}(\text{Cont}_0(M, \xi))$$

can be analogously defined. It provides a geometric representative of the connecting morphism. This generalizes the previous constructions. It will be used in Section 4.

2. SPHERES IN $\mathcal{C}(M, \xi)$

2.1. Almost contact structures. Let M be an oriented $(2n + 1)$ -dimensional manifold. Denote by $\text{Dist}(M)$ the space of smooth codimension-1 regular cooriented distributions on M . Concerning orientations, an almost complex structure on a cooriented distribution will be positive if the induced orientation coincides with the prescribed one. Define the space of almost contact structures as

$$\mathcal{A}(M) = \{(\xi, j) : \xi \in \text{Dist}(M), j \in \text{End}(\xi), j^2 = -id, j \text{ positive}\}.$$

It has a subset defined by

$$\mathcal{AC}(M, \xi) = \{(\eta, j) : \eta \in \mathcal{C}(M, \xi), j \in \text{End}(\eta), j^2 = -id, j \text{ compatible with } \eta\}.$$

The space of almost complex structures compatible with a fixed symplectic form is contractible. Thus, the forgetful map $\mathcal{AC}(M, \xi) \rightarrow \mathcal{C}(M, \xi)$ has a contractible homotopy fibre. Hence there exists a homotopy inverse $\iota : \mathcal{C}(M, \xi) \rightarrow \mathcal{AC}(M, \xi)$ given by a choice of a compatible almost complex structure

on the contact distribution.

Fix a point $p \in M$ and a framing $\tau : T_p M \xrightarrow{\cong} \mathbb{R}^{2n+1}$. Define the evaluation map

$$e_{(p,\tau)} : \mathcal{A}(M) \longrightarrow \mathcal{A}(\mathbb{R}^{2n+1}), \quad e_{(p,\tau)}(\xi, \mathbf{j}) = (\tau_* \xi_p, \tau_* \mathbf{j}_p).$$

This is a continuous map and thus induces $\tilde{e}_{(p,\tau)} : \pi_2(\mathcal{A}(M)) \longrightarrow \pi_2(\mathcal{A}(\mathbb{R}^{2n+1}))$. Therefore, we obtain

$$\varepsilon_{(p,\tau)} = \tilde{e}_{(p,\tau)} \circ 1_* : \pi_2(\mathcal{C}(M, \xi)) \longrightarrow \pi_2(\mathcal{A}(\mathbb{R}^{2n+1}))$$

Lemma 8. $\pi_2(\mathcal{A}(\mathbb{R}^{2n+1})) \cong \mathbb{Z}$.

Proof. The space $\mathcal{A}(\mathbb{R}^{2n+1})$ is isomorphic to the homogeneous space $SO(2n+1)/U(n)$. The standard inclusion $SO(2n) \longrightarrow SO(2n+1)$ descends to a map

$$SO(2n)/U(n) \longrightarrow SO(2n+1)/U(n)$$

with homotopy fibre \mathbb{S}^{2n} . The long exact sequence for homotopy fibration implies that $\pi_2(SO(2n)/U(n))$ and $\pi_2(SO(2n+1)/U(n))$ are isomorphic for $n \geq 2$. It is simple to show that $SO(2n+1)/U(n)$ is also isomorphic to $SO(2n+2)/U(n+1)$. Since $SO(4)/U(2)$ is a 2–sphere, the statement follows. \square

Thus the evaluation map can be seen as an integer–valued map for $\pi_2(\mathcal{C}(M, \xi))$.

Lemma 9. *The map $\varepsilon_{(p,\tau)} : \pi_2(\mathcal{C}(M, \xi)) \longrightarrow \pi_2(\mathcal{A}(\mathbb{R}^{2n+1}))$ is independent of the choice of p and τ .*

Proof. Let $p, q \in M$ and τ_p, τ_q be framings of $T_p M, T_q M$ respectively. Consider a continuous path of pairs $\{(p_t, \tau_t)\}$ connecting (p, τ_p) and (q, τ_q) . The continuous family of maps

$$e_{(p_t, \tau_t)} : \mathcal{A}(M) \longrightarrow \mathcal{A}(\mathbb{R}^{2n+1}), \quad e_{(p_t, \tau_t)}(\xi, \mathbf{j}) = (\tau_{t*} \xi_p, \tau_{t*} \mathbf{j}_p)$$

provides a homotopy between $e_{(p, \tau_p)}$ and $e_{(q, \tau_q)}$. \square

2.2. Linear Contact Spheres.

Definition 10. *A linear contact sphere is a contact sphere $\iota : \mathbb{S}^2 \longrightarrow \mathcal{C}(M, \xi)$ such that there exist three contact forms $(\alpha_0, \alpha_1, \alpha_2)$ satisfying*

$$\iota(p) = \ker(e_0 \alpha_0 + e_1 \alpha_1 + e_2 \alpha_2)$$

for the standard embedding $(e_0, e_1, e_2) : \mathbb{S}^2 \longrightarrow \mathbb{R}^3$.

Remark 11. *Such spheres can only exist in a $(4n+3)$ –dimensional manifold. The fact that α and $-\alpha$ do not induce the same volume form in dimensions congruent to 1 modulo 4 yields an obstruction for their existence.*

Note that for a 3–fold the triple $(\alpha_0, \alpha_1, \alpha_2)$ constitutes a framing of the cotangent bundle.

Lemma 12. *Let M be a 3–fold and S a linear contact sphere. The class $[S] \in \pi_2(\mathcal{C}(M, \xi))$ is non–trivial and has infinite order.*

Proof. Let $p \in M$ be a point and consider the framing $\tau = (\alpha_0, \alpha_1, \alpha_2)_p$. In the three–dimensional case $\mathcal{A}(\mathbb{R}^3)$ is homotopic to a 2–sphere. This homotopy can be realized by projection π onto the space of cooriented 2–plane distributions. The degree of the evaluation map is computed via

$$\begin{aligned} \mathbb{S}^2 &\xrightarrow{\varepsilon_{(p,\tau)}} \mathcal{A}(T_p M) \xrightarrow{\pi} \text{Dist}(\mathbb{R}^3) \cong \mathbb{S}^2 \\ z &\longmapsto e_0(z) \alpha_0(p) + e_1(z) \alpha_1(p) + e_2(z) \alpha_2(p) \longmapsto (e_0(z), e_1(z), e_2(z)). \end{aligned}$$

Being the identity, this map has degree 1. \square

2.3. Hypercontact manifolds. Let us define a class of contact manifolds with natural linear contact spheres. These objects have been studied in the literature, see [G1].

Definition 13. *Let M be a smooth manifold and consider the embedding $M \xrightarrow{e} M \times \mathbb{R}$. Let t be the coordinate on the \mathbb{R} factor. The manifold M is hypercontact if there exists a hyperkähler cone metric g on $M \times \mathbb{R}$ such that the compatible complex structures (I, J, K) satisfy:*

- 1) *For any $\mathbf{j} \in \{I, J, K\}$, the complex tangencies $e^*(e_* TM \cap \mathbf{j} e_* TM) = \ker(\alpha_{\mathbf{j}})$ are contact structures, where $\alpha_{\mathbf{j}}(v) = g(e_* v, \mathbf{j} \partial_t)$.*
- 2) *The three symplectic structures $d(e^t \alpha_{\mathbf{j}})$ on $M \times \mathbb{R}$ define the same hyperkähler structure.*

Remark 14. *In a hypercontact manifold, the associated linear sphere defined by $(\alpha_I, \alpha_J, \alpha_K)$ is a contact sphere.*

Theorem 15. *Let M^{4n+3} be a hypercontact manifold. The class of a linear contact sphere associated to the hypercontact structure is an element of infinite order in $\pi_2(\mathcal{C}(M, \ker(\alpha_I)))$.*

Proof. Let $p \in M$ and note that the $4n$ -distribution $\eta = \ker \alpha_I \cap \ker \alpha_J \cap \ker \alpha_K$ is (I, J, K) -invariant. Thus, it can be identified with the quaternionic vector space \mathbb{H}^n by fixing a quaternionic framing $v = \{v_1, \dots, v_n\}$. This induces a real framing $\tau = \{v, Iv, Jv, Kv\}$ for η , identifying it with \mathbb{R}^{4n} endowed with the standard quaternionic structure.

Consider the Reeb vector fields R_I, R_J, R_K associated to α_I, α_J , and α_K . Extend the framing τ to $\tilde{\tau} = \{\tau, R_I, R_J, R_K\}$. Interpret the space $\mathcal{A}(\mathbb{R}^{4n+3})$ as pairs of (v, \mathfrak{j}) , where $v \in \mathbb{S}^{4n+2} \subset \mathbb{R}^{4n+3}$ is a unit vector and \mathfrak{j} an almost complex structure in its orthogonal space. Define

$$(4) \quad h : \mathcal{A}(\mathbb{R}^{4n+3}) \longrightarrow \mathcal{J}(\mathbb{R}^{4n+3} \oplus \mathbb{R}), \quad (v, \mathfrak{j}) \longmapsto \tilde{\mathfrak{j}} : \langle v \rangle^\perp \oplus \langle v \rangle \oplus \langle \partial_t \rangle \longrightarrow \langle v \rangle^\perp \oplus \langle v \rangle \oplus \langle \partial_t \rangle$$

where the almost complex structure is $\tilde{\mathfrak{j}} = \begin{pmatrix} \mathfrak{j} & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}$. This induces a morphism of second ho-

motopy groups. Through the above identification the linear contact sphere generated by $(\alpha_I, \alpha_J, \alpha_K)$ evaluates in a sphere $\langle (\xi_I, I), (\xi_J, J), (\xi_K, K) \rangle \in \mathcal{A}(\mathbb{R}^{4n+3})$. This sphere maps via (4) to the sphere of complex structures generated by the triple (I, J, K) in $\mathcal{J}(\mathbb{R}^{4n+4})$.

It is left to prove that the class of that sphere is an infinite order element of $\pi_2(SO(4n+4)/U(2n+2))$. Let us write $m = n + 1$ to ease notation. The homotopy fibration

$$U(2m) \longrightarrow SO(4m) \longrightarrow SO(4m)/U(2m)$$

induces an injection $\pi_2(SO(4m)/U(2m)) \longrightarrow \pi_1(U(2m)) \cong \mathbb{Z}$.

Let $(\theta, \phi) \in [0, 2\pi] \times [0, \pi]$ be spherical angles. Define

$$J_\theta = \cos \theta J + \sin \theta K, \quad \tilde{I} = \cos \phi I + \sin \phi J_\theta, \quad P_{\theta, \phi} = \cos(\phi/2)I + \sin(\phi/2)J_\theta.$$

The sphere is represented by \tilde{I} , we shall compute its image under the boundary morphism. Note that $P_{\theta, \phi} \in SO(4m)$ and $\tilde{I} = P_{\theta, \phi}^\dagger I P_{\theta, \phi}$. Further $P_{\theta, \pi} = J_\theta = (\cos \theta \cdot id + \sin \theta I)J$, with $\cos \theta \cdot id + \sin \theta I \in U(2m)$ and $J \in SO(4m)$. This decomposition provides a representative in $\pi_2(SO(4m)/U(2m))$. Thus the loop in $\pi_1(U(2m))$ is provided by $\cos \theta \cdot id + \sin \theta I$ with $\theta \in [0, 2\pi]$. Since the identification $\pi_1(U(2m)) \cong \mathbb{Z}$ is given by the complex determinant, the degree of the sphere is $2m$. \square

The argument above applies to a broader class of manifolds:

Definition 16. *A contact manifold (M, ξ_0) is said to possess an almost-quaternionic sphere if it admits a sphere $\mathbb{S}^2 \xrightarrow{\xi} \mathcal{C}(M, \xi_0)$ such that:*

- 1) *There exists a family $\{\mathfrak{j}_p\}_{p \in \mathbb{S}^2}$ compatible with the contact distributions $\xi_p = \xi(p)$,*
- 2) *There exists a point $q \in M$ and a framing τ for $T_q M$ such that $e_{q, \tau}(\xi(\mathbb{S}^2))$ becomes the linear sphere associated to $\langle (\xi_I, I), (\xi_J, J), (\xi_K, K) \rangle \in \mathcal{A}(\mathbb{R}^{4n+3})$.*

Corollary 17. *An almost-quaternionic sphere inside a contact manifold (M, ξ) generates a class of infinite order in $\pi_2(\mathcal{C}(M, \xi))$.*

3. REEB FLOW FOR SPHERES

Let us prove Corollary 2. The standard contact sphere will be denoted (\mathbb{S}^{2n+1}, ξ) . The relevant case is that of the spheres \mathbb{S}^{2k+1} with k odd. Indeed, for the spheres \mathbb{S}^{2k+1} with $k = 2n$ the Reeb flow is non-trivial in $\pi_1(SO(4n+1)) \subset \pi_1(\text{Diff}(\mathbb{S}^{4n+1}))$. Thus it cannot be contractible in $\text{Cont}_0(M, \xi) \subset \text{Diff}_0(\mathbb{S}^{4n+1})$. In order to conclude the case \mathbb{S}^{4n+3} we detail the construction in Sections 1 and 2.

Consider the endomorphisms I, J, K of $\mathbb{R}^{4(n+1)}$ obtained by direct sum of the corresponding endomorphisms i, j, k of \mathbb{R}^4 , satisfying the quaternionic relations

$$i^2 = j^2 = k^2 = ijk = -1.$$

The endomorphisms I, J, K anti-commute and hence any of their linear combinations is a complex structure. Let $e = (e_0, e_1, e_2) : \mathbb{S}^2 \rightarrow \mathbb{R}^3$ be the standard embedding of the 2-sphere in Euclidean 3-space with azimuthal angle θ and polar angle ϕ :

$$e_0 = \cos \theta \sin \phi, \quad e_1 = \sin \theta \sin \phi, \quad e_2 = \cos \phi, \quad (\theta, \phi) \in [0, 2\pi] \times [0, \pi].$$

A complex structure $j \in \text{End}(\mathbb{R}^{4n+4})$ induces the real $(4n+2)$ -distribution

$$\xi_j = T\mathbb{S}^{4n+3} \cap jT\mathbb{S}^{4n+3}$$

of j -complex tangencies on the sphere \mathbb{S}^{4n+3} . There exists a unique $U(j, n)$ -invariant 1-form α_j such that $\ker \alpha_j = \xi_j$. It is given by $\alpha(z) = z^t j dz$. We use the following three 1-forms

$$\alpha_0 = \alpha_I, \quad \alpha_1 = \alpha_J, \quad \alpha_2 = \alpha_K.$$

Their respective Reeb vector fields R_0, R_1 and R_2 are linearly independent and their flows are given by the family of rotations generated by I, J and K . Consider the 1-form $\alpha = e_0 \alpha_0 + e_1 \alpha_1 + e_2 \alpha_2$. The form α is a contact form on \mathbb{S}^{4n+3} for each value of e . Although not used in the rest of the article, it is simple to prove the following

Lemma 18. $(\mathbb{S}^2 \times \mathbb{S}^{4n+3}, \ker \alpha)$ is a contact manifold.

Let us compute the loop at infinity for the trivial contact fibration

$$\mathbb{S}^2 \times \mathbb{S}^{4n+3} \rightarrow \mathbb{S}^2, (e, p) \mapsto e.$$

In the spherical coordinates above, we will obtain the loop at infinity corresponding to $\phi = \pi$. The contact connection allows us to lift a vector field X in the base \mathbb{S}^2 . The lift \tilde{X} is the unique vector field on $\mathbb{S}^2 \times \mathbb{S}^{4n+3}$ conforming the two conditions

$$\alpha(\tilde{X}) = 0, \quad d\alpha(\tilde{X}, V) = 0, \quad \text{with } V \text{ an arbitrary vertical vector field.}$$

Since uniqueness is provided once a solution is found, the following assertion can be readily verified

Lemma 19. The lift of the polar vector field ∂_ϕ to the contact connection given by α is

$$\tilde{X}_\phi = \partial_\phi + \frac{1}{2}(-\sin \theta R_0 + \cos \theta R_1).$$

The Hamiltonian will appear once we pull-back the contact form α with the π -time flow of the lift \tilde{X}_ϕ . Consider the linear endomorphism $F_\theta = \frac{1}{2}(-\sin \theta I + \cos \theta J)$. The flow associated to \tilde{X}_ϕ induces a diffeomorphism between the central fibre $\{\phi = 0\}$ and the fibre at an arbitrary ϕ . This diffeomorphism can be expressed as

$$\varphi_\phi : \mathbb{S}^{4n+3} \rightarrow \mathbb{S}^{4n+3}, \quad \varphi(p) = e^{F_\theta \phi} p.$$

This is understood as a map in complex space \mathbb{C}^{2n+2} restricted to the sphere. The theory explained in Section 1, in particular formula (1), implies that the pull-back will be of the form $\alpha_2 + H(p, \phi)d\theta$. A computation yields

Lemma 20. $\varphi_\phi^*(\alpha) = \alpha_2 + \sin^2(\phi/2)d\theta$

The loops correspond to the flow of the vector field associated to $G = -\sin^2(\phi/2)$. The loop at infinity has Hamiltonian $G|_{\phi=\pi} \equiv -1$. Thus it is the Reeb flow.

We have geometrically realized the boundary map of the long exact homotopy sequence (2). The non-contractibility of the Reeb flow will follow from an understanding of the contact sphere above and the group $\pi_2(\text{Diff}_0(\mathbb{S}^{4n+3}))$. Regarding the former we have the following

Lemma 21. Let S be the sphere of complex structures

$$S = \{e_0 I + e_1 J + e_2 K : e \in \mathbb{S}^2\} \subset SO(4n+4)/U(2n+2).$$

- 1) S represents a non-trivial element of $\pi_2(SO(4n+4)/U(2n+2)) \cong \mathbb{Z}$.
- 2) $S \subset \mathcal{C}(\mathbb{S}^{4n+3}, \xi)$ and generates an infinite cyclic subgroup in $\pi_2(\mathcal{C}(\mathbb{S}^{4n+3}, \xi))$.

Proof. Both statements follow from the argument provided in the proof of Theorem 15. □

Concerning the group $\text{Diff}_0(\mathbb{S}^{4n+3})$, the following lemma will suffice.

Lemma 22. $\pi_2(\text{Diff}(\mathbb{S}^{4n+3})) \otimes \mathbb{Q} = 0$ for $n \geq 2$.

Proof. This is a result in algebraic topology. Let $\text{Diff}_0(\mathbb{D}^l, \partial)$ be the group of diffeomorphisms of the l -disk restricting to the identity at the boundary. Note the homotopy equivalence

$$\text{Diff}_0(\mathbb{S}^l) \simeq SO(l+1) \times \text{Diff}_0(\mathbb{D}^l, \partial)$$

and that $\pi_2(SO(l+1)) = 0$ since $SO(l+1)$ is a Lie group. Let $\phi(l) = \min\{(l-4)/3, (l-7)/2\}$. In the stable concordance range $0 \leq j < \phi(l)$ we have

$$(5) \quad \pi_j(\text{Diff}_0(\mathbb{D}^l, \partial)) \otimes \mathbb{Q} = 0, \quad \text{if } l \text{ even or } 4 \nmid j + 1.$$

See [We] for details. In particular $\pi_2(\text{Diff}_0(\mathbb{D}^l, \partial)) \otimes \mathbb{Q} = 0$ for $l > 11$. We are thus able to conclude

$$\pi_2(\text{Diff}_0(\mathbb{S}^{4n+3})) \otimes \mathbb{Q} \cong \pi_2(\text{Diff}_0(\mathbb{D}^{4n+3}, \partial)) \otimes \mathbb{Q} = 0, \quad n > 2.$$

For the case $n = 2$ we provide a more *ad hoc* argument. Let $C(\mathbb{D}^{11})$ be the space of pseudo-isotopies for the disk \mathbb{D}^{11} . There exists a homotopy fibration

$$\text{Diff}_0(\mathbb{D}^{12}, \partial) \longrightarrow C(\mathbb{D}^{11}) \longrightarrow \text{Diff}_0(\mathbb{D}^{11}, \partial)$$

Algebraic K -theory implies $\pi_1 C(\mathbb{D}^{11}) \otimes \mathbb{Q} = \pi_2 C(\mathbb{D}^{11}) \otimes \mathbb{Q} = 0$. Observe that (5) implies that $\pi_1(\text{Diff}(\mathbb{D}^{12}, \partial))$ is a torsion group. The long exact homotopy sequence of the above fibration gives

$$\dots \longrightarrow \pi_2(C(\mathbb{D}^{11})) \xrightarrow{\rho_2} \pi_2(\text{Diff}_0(\mathbb{D}^{11}, \partial)) \xrightarrow{\partial} \pi_1(\text{Diff}_0(\mathbb{D}^{12}, \partial)) \xrightarrow{i_1} \pi_1(C(\mathbb{D}^{11})) \longrightarrow \dots$$

This implies the short exact sequence of Abelian groups

$$0 \longrightarrow A \longrightarrow \pi_2(\text{Diff}_0(\mathbb{D}^{11}, \partial)) \longrightarrow B \longrightarrow 0,$$

where $A = \ker \partial = \text{im } \rho_2$ and $B = \text{im } i_1 = \text{coker } \rho_2$. Thus $\pi_2(\text{Diff}_0(\mathbb{D}^{11}, \partial))$ is a torsion group. \square

Remark 23. *The Smale conjecture* $\text{Diff}_0(\mathbb{S}^3) \simeq SO(3)$ holds for \mathbb{S}^3 , see [Ha].

In order to conclude Corollary 2 for \mathbb{S}^{4n+3} consider the class of the Reeb loop in $\pi_1(\text{Cont}_0(\mathbb{S}^{4n+3}, \xi))$. The construction explained above shows that it lies in the image of the boundary morphism

$$\partial : \mathcal{C}(\mathbb{S}^{4n+3}, \xi) \longrightarrow \pi_1(\text{Cont}_0(\mathbb{S}^{4n+3}, \xi)).$$

If the Reeb class were to be zero the sphere S would lie in the image of $\pi_2(\text{Diff}_0(\mathbb{S}^{4n+3}))$ in (2). Lemma 22 implies that such a sphere needs to be a torsion class if $n \geq 2$. Lemma 21 contradicts this statement. Thus proving Corollary 2.

4. HIGHER HOMOTOPY GROUPS

The previous arguments can be modified for n -dimensional homotopy spheres. This allows us to conclude properties of the higher homotopy type of the contactomorphism group. Consider the evaluation map

$$e_{p,\tau} : \mathcal{A}(M) \longrightarrow \mathcal{A}(\mathbb{R}^{2n+1}).$$

Composition with the homotopy inverse $\iota : \mathcal{C}(M) \rightarrow \mathcal{AC}(M)$ defines higher homotopy maps

$$\pi_k(e_{p,\tau} \circ \iota) : \pi_k(\mathcal{C}(M)) \longrightarrow \pi_k(\mathcal{A}(\mathbb{R}^{2n+1})), \quad k \geq 1.$$

Let us provide a simple application. Define the natural inclusion

$$i_{\mathcal{J}} : \mathcal{J}(\mathbb{R}^{2n+2}) \longrightarrow \mathcal{C}(\mathbb{S}^{2n+1}, \xi), \quad i_{\mathcal{J}}(j) = T\mathbb{S}^{2n+1} \cap jT\mathbb{S}^{2n+1}.$$

Lemma 24. *The map $i_{\mathcal{J}}$ is a homotopy inclusion.*

Proof. Consider the following chain of maps

$$c : \mathcal{J}(\mathbb{R}^{2n+2}) \xrightarrow{i_{\mathcal{J}}} \mathcal{C}(\mathbb{S}^{2n+1}, \xi) \xrightarrow{e_{p,\tau} \circ \iota} \mathcal{A}(\mathbb{R}^{2n+1}) \xrightarrow{h} \mathcal{J}(\mathbb{R}^{2n+2}).$$

The definition of each map implies $c = \text{id}$. Therefore, it induces the identity in homotopy:

$$\pi_k(c) = \text{id} : \pi_k(\mathcal{J}(\mathbb{R}^{2n+2})) \xrightarrow{\pi_k(i_{\mathcal{J}})} \pi_k(\mathcal{C}(\mathbb{S}^{2n+1}, \xi)) \xrightarrow{\pi_k(e_{p,\tau} \circ \iota)} \pi_k(\mathcal{A}(\mathbb{R}^{2n+1})) \xrightarrow{\pi_k(h)} \pi_k(\mathcal{J}(\mathbb{R}^{2n+2})).$$

Thus the map $i_{\mathcal{J}}$ induces an injection $\pi_k(i_{\mathcal{J}})$. \square

This lemma can be combined with results on the homotopy type of the group $\text{Diff}(\mathbb{S}^{2n+1})$. We can then conclude the existence of infinite order elements in certain homotopy groups of $\text{Cont}(\mathbb{S}^{2n+1}, \xi)$. Among many others, a simple instance is the following

Lemma 25. *The group $\pi_5(\text{Cont}(\mathbb{S}^{8k-1}), \xi)$ has an element of infinite order, for $k \geq 3$.*

Proof. Using the connecting map ∂_6 , as described in equation (3), the statement is reduced to the following two assertions:

- $\pi_6(\mathcal{J}(\mathbb{R}^{8k})) = \pi_6(SO(8k)/U(4k)) = \mathbb{Z}$ and therefore, by Lemma 24, $\text{rk}(\pi_6(\mathcal{C}(\mathbb{S}^{8k-1}, \xi))) \geq 1$.
- $\pi_6(\text{Diff}(\mathbb{S}^{8k-1})) \otimes \mathbb{Q} = 0$, for $k \geq 3$. This is again a consequence of the results in [We].

□

These techniques can be adapted for general contact manifolds as long as there is a partial understanding of the homotopy type of their group of diffeomorphisms.

REFERENCES

- [Bo] F. Bourgeois, Contact homology and homotopy groups of the space of contact structures, *Math. Res. Lett.* 13 (2006), 71–85.
- [El] Y. Eliashberg, Contact 3-manifolds twenty years since J. Martinet’s work, *Ann. Inst. Fourier* 46 (1992), 165–192.
- [G1] H. Geiges, C.B. Thomas, Hypercontact manifolds, *J. London Math. Soc.* 2 (2) 51 (1995), 342–352.
- [G2] H. Geiges, J. Gonzalo, On the topology of the space of contact structures on torus bundles, *Bull. London Math. Soc.* 36 (2004), 640–646.
- [Gi] E. Giroux, Sur la géométrie et la dynamique des transformations de contact, *Séminaire Bourbaki* (2009), n.1004.
- [Ha] A. E. Hatcher, A proof of the Smale conjecture, *Ann. of Math.* 117 (1983), 553–607.
- [Pr] F. Presas, A class of non-fillable contact structures, *Geometry&Topology* 11 (2007), 2203–2225.
- [We] M. Weiss, B. Williams, Automorphisms of manifolds, *Surveys on Surgery Theory. Vol. II Papers Dedicated to C.T.C Wall*, *Annals of Mathematics Studies* 149 (2001).

INSTITUTO DE CIENCIAS MATEMÁTICAS – CSIC. C. NICOLÁS CABRERA, 13–15, 28049, MADRID, SPAIN.
E-mail address: `casals.roger@icmat.es`

INSTITUTO DE CIENCIAS MATEMÁTICAS – CSIC. C. NICOLÁS CABRERA, 13–15, 28049, MADRID, SPAIN.
E-mail address: `fpresas@icmat.es`