

# Bayesian abstract fuzzy economies, random quasi-variational inequalities with random fuzzy mappings and random fixed point theorems

Monica Patriche

*University of Bucharest Faculty of Mathematics and Computer Science  
14 Academiei Street  
010014 Bucharest, Romania  
monica.patriche@yahoo.com*

---

## Abstract

In this paper, we introduce an abstract fuzzy economy model with a measure space of agents which generalizes Patriche model (2009), we obtain a theorem of fuzzy equilibrium existence and we prove the existence of the solutions for two types of random quasi-variational inequalities with random fuzzy mappings. As a consequence, we obtain random fixed point theorems.

*Keywords:* Bayesian abstract fuzzy economy, Bayesian fuzzy equilibrium, incomplet information, random fixed point, random quasi-variational inequalities, random fuzzy mapping.

---

AMS Subject Classification: 58E35, 47H10, 91B50, 91A44.

## 1. INTRODUCTION

The concept of a fuzzy game (or a fuzzy abstract economy) has been introduced in [11] and the existence of the equilibrium for 1-person fuzzy game has been proven. The theory of fuzzy sets, initiated by Zadeh [29], has become a good framework for obtaining results concerning fuzzy equilibrium existence for abstract fuzzy economies. The existence of equilibrium points of fuzzy games has been also studied in [2], [9]-[13],[17]-[19], [23].

Firstly, we introduce a fuzzy extension of Patriche's model of the Bayesian abstract economy [16] and we prove the existence of Bayesian fuzzy equilibrium. All economic activity in a society is made under conditions of uncertainty (incomplet information) and the model we propose captures this

meaning. Actually, we define an abstract fuzzy economy with asymmetric information and a measure set of agents, each of which is characterized by a private information set, a fuzzy action (strategy) mapping, a random fuzzy constraint one and a random fuzzy preference mapping. The fuzzy equilibrium concept is an extension of the deterministic equilibrium. Our model generalizes the former deterministic ones introduced by Debreu [3], Shafer and Sonnenschein [21] or Yannelis and Prabhakar [25] or Patriche [16]. For an overview of the results concerning the equilibrium of abstract economies, the reader is referred to [20].

As application, we study random quasi-variational inequalities with random fuzzy mappings. The variational inequalities were introduced in 1960s by Fichera and Stampacchia, who studied equilibrium problems arising from mechanics. Since then, this domain has been extensively studied and has been found very useful in many diverse fields of pure and applied sciences, such as mechanics, physics, optimization and control theory, operations research and several branches of engineering sciences. Recently, the random variational inequality and random quasi-variational inequality problems have been introduced and studied in [7], [8], [15], [22], [28].

In this paper, we introduce an abstract fuzzy economy model with a measure space of agents which generalizes Patriche model (2009), we prove a theorem of fuzzy equilibrium existence and prove the existence of the solution for two types of random quasi-variational inequalities with random fuzzy mappings. As a consequence, we obtain random fixed point theorems.

The paper is organized as follows. In the next section, some notational and terminological conventions are given. We also present, for the reader's convenience, some results on Bochner integration. In Section 3, the model of differential information abstract fuzzy economy and the main result are presented. Section 4 contains existence theorems for solutions of random quasi-variational inequalities with random fuzzy mappings.

## 2. NOTATION AND DEFINITION

Throughout this paper, we shall use the following notation:

1.  $\mathbb{R}_{++}$  denotes the set of strictly positive reals.  $\text{co}D$  denotes the convex hull of the set  $D$ .  $\overline{\text{co}}D$  denotes the closed convex hull of the set  $D$ .  $2^D$  denotes the set of all non-empty subsets of the set  $D$ . If  $D \subset Y$ , where  $Y$  is a topological space,  $\text{cl}D$  denotes the closure of  $D$ .

For the reader's convenience, we review a few basic definitions and results from continuity and measurability of correspondences, Bochner integrable functions and the integral of a correspondence.

Let  $Z$  and  $Y$  be sets.

**Definition 1.** *The graph of the correspondence  $P : Z \rightarrow 2^Y$  is the set  $G_P = \{(z, y) \in Z \times Y : y \in P(z)\}$ .*

Let  $Z, Y$  be topological spaces and  $P : Z \rightarrow 2^Y$  be a correspondence.

1.  $P$  is said to be *upper semicontinuous* if for each  $z \in Z$  and each open set  $V$  in  $Y$  with  $P(z) \subset V$ , there exists an open neighborhood  $U$  of  $z$  in  $Z$  such that  $P(y) \subset V$  for each  $y \in U$ .

2.  $P$  is said to be *lower semicontinuous* if for each  $z \in Z$  and each open set  $V$  in  $Y$  with  $P(z) \cap V \neq \emptyset$ , there exists an open neighborhood  $U$  of  $z$  in  $Z$  such that  $P(y) \cap V \neq \emptyset$  for each  $y \in U$ .

**Lemma 1.** *(see [27]). Let  $Z$  and  $Y$  be two topological spaces and let  $D$  be an open subset of  $Z$ . Suppose  $P_1 : Z \rightarrow 2^Y$ ,  $P_2 : Z \rightarrow 2^Y$  are upper semicontinuous such that  $P_2(z) \subset P_1(z)$  for all  $z \in D$ . Then the correspondence  $P : Z \rightarrow 2^Y$  defined by*

$$P(z) = \begin{cases} P_1(z), & \text{if } z \notin D, \\ P_2(z), & \text{if } z \in D \end{cases}$$

*is also upper semicontinuous.*

**Definition 2.** *Let  $Y$  be a metric space and  $Y'$  be its dual.  $P : Y \rightarrow 2^{Y'}$  is said to be *monotone* if  $\operatorname{Re}\langle u - v, y - x \rangle \geq 0$  for all  $u \in P(y)$  and  $v \in P(x)$  and  $x, y \in Y$ .*

Let now  $(\Omega, \mathcal{F}, \mu)$  be a complete, finite measure space, and  $Y$  be a topological space.

**Definition 3.** *The correspondence  $P : \Omega \rightarrow 2^Y$  is said to have a *measurable graph* if  $G_P \in \mathcal{F} \otimes \beta(Y)$ , where  $\beta(Y)$  denotes the Borel  $\sigma$ -algebra on  $Y$  and  $\otimes$  denotes the product  $\sigma$ -algebra.*

**Lemma 2.** (see [14]). Let  $P_n : \Omega \rightarrow 2^Y$ ,  $n = 1, 2, \dots$  be a sequence of correspondences with measurable graphs. Then the correspondences  $\cup_n P_n$ ,  $\cap_n P_n$  and  $Y \setminus P_n$  have measurable graphs.

Let  $(\Omega, F, \mu)$  be a measure space and  $Y$  be a Banach space.

It is known (see [14], Theorem 2, p.45) that, if  $x : \Omega \rightarrow Y$  is a  $\mu$ -measurable function, then  $x$  is Bochner integrable if and only if  $\int_{\Omega} \|x(\omega)\| d\mu(\omega) < \infty$ .

It is denoted by  $L_1(\mu, Y)$  the space of equivalence classes of  $Y$ -valued Bochner integrable functions  $x : \Omega \rightarrow Y$  normed by  $\|x\| = \int_{\Omega} \|x(\omega)\| d\mu(\omega)$ .

Also it is known (see [4], p.50) that  $L_1(\mu, Y)$  is a Banach space.

**Definition 4.** The correspondence  $P : \Omega \rightarrow 2^Y$  is said to be integrably bounded if there exists a map  $h \in L_1(\mu, \mathbb{R})$  such that  $\sup\{\|x\| : x \in P(\omega)\} \leq h(\omega)$   $\mu$ -a.e.

We denote by  $S_P^1$  the set of all selections of the correspondence  $P : \Omega \rightarrow 2^Y$  that belong to the space  $L_1(\mu, Y)$ , i.e.

$$S_P^1 = \{x \in L_1(\mu, Y) : x(\omega) \in P(\omega) \text{ } \mu\text{-a.e.}\}.$$

Further, we will see the conditions under which  $S_P^1$  is nonempty and weakly compact in  $L_1(\mu, Y)$ . Aumann measurable selection theorem (see Appendix) and Diestel's Theorem (see Appendix) are necessary.

Let  $F(Y)$  be a collection of all fuzzy sets over  $Y$ .

**Definition 5.** A mapping  $P : \Omega \rightarrow F(Y)$  is called a fuzzy mapping. If  $P$  is a fuzzy mapping from  $\Omega$ ,  $P(\omega)$  is a fuzzy set on  $Y$  and  $P(\omega)(y)$  is the membership function of  $y$  in  $P(\omega)$ .

Let  $A \in F(Y)$ ,  $\alpha \in [0, 1]$ , then the set  $(A)_{\alpha} = \{y \in Y : A(y) \geq \alpha\}$  is called an  $\alpha$ -cut set of fuzzy set  $A$ .

**Definition 6.** A fuzzy mapping  $P : \Omega \rightarrow F(Y)$  is said to be measurable if for any given  $\alpha \in [0, 1]$ ,  $(P(\cdot))_{\alpha} : \Omega \rightarrow 2^Y$  is a measurable set-valued mapping.

**Definition 7.** We say that a fuzzy mapping  $P : \Omega \rightarrow F(Y)$  is said to have a measurable graph if for any given  $\alpha \in [0, 1]$ , the set-valued mapping  $(P(\cdot))_{\alpha} : \Omega \rightarrow 2^Y$  has a measurable graph.

**Definition 8.** A fuzzy mapping  $P : \Omega \times X \rightarrow F(Y)$  is called a random fuzzy mapping if for any given  $x \in X$ ,  $P(\cdot, x) : \Omega \rightarrow F(Y)$  is a measurable fuzzy mapping.

### 3. FUZZY EQUILIBRIUM EXISTENCE FOR BAYESIAN ABSTRACT FUZZY ECONOMIES

#### 3.1. THE MODEL OF AN ABSTRACT FUZZY ECONOMY WITH A MEASURABLE SPACE OF AGENTS

We introduce the next model of the abstract fuzzy economy which generalizes the classical deterministic models introduced by Debreu [3], Shafer and Sonnenschein [21], Yannelis and Prabhakar [25] and Patriche [16].

Let  $(\Omega, F, \mu)$  be a complete finite separable measure space, where  $\Omega$  denotes the set of states of nature of the world and the  $\sigma$ -algebra  $F$  denotes the set of events. Let  $Y$  be a separable Banach space, denoting the commodity or strategy space.

Let us consider  $X : T \times \Omega \rightarrow \mathcal{F}(Y)$  and  $z \in (0, 1]$ .

Let  $S_{X_t}^1 = \{y(t) \in L_1(\mu, Y) : y(t, \cdot) : \Omega \rightarrow Y \text{ is } F_t \text{-measurable and } y(t, \omega) \in (X(t, \omega))_z \mu - a.e.\}$ . Notice that  $S_{X_t}^1$  is the set of all Bochner integrable and  $F_t$ -measurable selections from the random strategy of agent  $t$ . In essence this is the set, out of which agent  $t$  will pick his/her optimal choices. In particular, an element  $x_t$  in  $S_{X_t}^1$  is called a *strategy* for agent  $t$ . The typical element of  $S_{X_t}^1$  is denoted by  $\tilde{x}_t$  and that of  $X(t, \omega)$  by  $x_t(\omega)$ . Let  $S_X^1 = \{\tilde{x} \in L_1(v, L_1(\mu, Y)) : \tilde{x}(t, \cdot) \in S_{X_t}^1 \text{ } v - a.e.\}$ . An element of  $S_X^1$  will be a *joint strategy profile*.

We introduce the following model.

**Definition 9.** *A Bayesian abstract fuzzy economy (or social system) with differential information and a measure space of agents  $(T, \tau, v)$  is a set  $G = \{(X, F_t, A, P, a, b, z), t \in T\}$ , where*

1)  $X : T \times \Omega \rightarrow \mathcal{F}(Y)$  is the random fuzzy action (strategy) mapping, where,  $X(t, \omega) \subset \mathcal{F}(Y)$  is interpreted as the strategy set of agent  $t$  of the state of nature  $\omega$ ;

2)  $F_t$  is a sub  $\sigma$ -algebra of  $F$  which denotes the private information of agent  $t$ ;

3) for each  $t \in T$ ,  $A(t, \cdot, \cdot) : \Omega \times S_X^1 \rightarrow \mathcal{F}(Y)$  is the random fuzzy constraint mapping of agent  $t$ ;

4) for each  $t \in T$ ,  $P(t, \cdot, \cdot) : \Omega \times S_X^1 \rightarrow \mathcal{F}(Y)$  is the random fuzzy preference mapping of agent  $t$ ;

5)  $z \in (0, 1]$  such that for all  $(t, \omega, x) \in T \times \Omega \times S_X^1$ ,  $(A(t, \omega, \tilde{x}))_{a(\tilde{x})} \subset (X(t, \omega))_z$  and  $(P(t, \omega, \tilde{x}))_{p(\tilde{x})} \subset (X(t, \omega))_z$ ;

5)  $a : S_X^1 \rightarrow (0, 1]$  is a random fuzzy constraint function and  $p : S_X^1 \rightarrow (0, 1]$  is a random fuzzy preference function.

We propose the next concept of fuzzy equilibrium for an abstract fuzzy economy with a measure space of agents.

**Definition 10.** *A Bayesian fuzzy equilibrium for  $G$  is a strategy profile  $\tilde{x}^* \in S_X^1$  such that for  $v - a.e.$ ,*

- i)  $\tilde{x}^*(t, \omega) \in (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)} \mu - a.e.$ ;
- ii)  $(A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)} \cap (P(t, \omega, \tilde{x}^*))_{p(\tilde{x}^*)} = \emptyset \mu - a.e.$

### 3.2. EXISTENCE OF THE BAYESIAN FUZZY EQUILIBRIUM

Now, we are establishing the next equilibrium existence theorem for Bayesian abstract fuzzy economies with a measure space of agents and with upper semi-continuous fuzzy mappings. Our result generalizes that one we proved in [16].

Let  $(\Omega, F, \mu)$  be a complete finite measure separable space, where  $\Omega$  denotes the set of states of nature of the world and the  $\sigma$ -algebra  $F$  denotes the set of events. Let  $Y$  denote the strategy or commodity space, where  $Y$  is a separable Banach space.

**Theorem 3.** *Let  $(T, \tau, \nu)$  be a measure space of agents and be the Bayesian fuzzy abstract economy  $G = \{(X, F_t, A, P, a, p, z), t \in T\}$  satisfying (A.1)-(A.5). Then, there exists a Bayesian equilibrium for  $G$ .*

A.1)

(a)  $X : T \times \Omega \rightarrow F(Y)$  is such that  $(t, \omega) \rightarrow (X(t, \omega))_z : T \times \Omega \rightarrow 2^Y$  is a non-empty convex weakly compact-valued and integrably bounded fuzzy mapping;

(b) For each fixed  $t \in T$ ,  $(X(t, \cdot))_z$  has an  $F_t$ -measurable graph;

A.2)

(a) the correspondence  $(t, \omega, \tilde{x}) \rightarrow (A(t, \omega, \tilde{x}))_{a(\tilde{x})} : T \times \Omega \times S_X^1 \rightarrow 2^Y$  has a measurable graph;

(b) For each  $(t, \omega) \in T \times \Omega$ ,  $\tilde{x} \rightarrow (A(t, \omega, \tilde{x}))_{a(\tilde{x})} : S_X^1 \rightarrow 2^Y$  is an upper semicontinuous correspondence with closed, convex and non-empty values.

A.3)

(a) the correspondence  $(t, \omega, \tilde{x}) \rightarrow (P(t, \omega, \tilde{x}))_{p(\tilde{x})} : T \times \Omega \times S_X^1 \rightarrow 2^Y$  has a measurable graph;

(b) For each  $(t, \omega) \in T \times \Omega$ ,  $\tilde{x} \rightarrow (P(t, \omega, \tilde{x}))_{p(\tilde{x})} : S_X^1 \rightarrow 2^Y$  is an upper semicontinuous correspondence with closed, convex and non-empty values.

A.4) The correspondence  $t \rightarrow S_{X_t}^1$  has a measurable graph.

A.5)

(a) For each  $(t, \omega) \in T \times \Omega$ , for each  $\tilde{x} \in S_X^1$ ,  $\tilde{x}(t, \omega) \notin (A(t, \omega, \tilde{x}))_{a(\tilde{x})} \cap (P(t, \omega, \tilde{x}))_{p(\tilde{x})}$

(b) the set  $U^{(t, \omega)} = \{\tilde{x} \in S_X^1 : (A(t, \omega, \tilde{x}))_{a(\tilde{x})} \cap (P(t, \omega, \tilde{x}))_{p(\tilde{x})} \neq \emptyset\}$  is weakly open in  $S_X^1$ .

*Proof.* Define  $\Phi : T \times \Omega \times S_X^1 \rightarrow 2^Y$  by  $\Phi(t, \omega, \tilde{x}) = (A(t, \omega, \tilde{x}))_{a(\tilde{x})} \cap (P(t, \omega, \tilde{x}))_{p(\tilde{x})}$ . We shall prove first that  $S_X^1$  is non-empty, convex, weakly compact.

Since  $(\Omega, F, \mu)$  is a complete finite measure space,  $Y$  is a separable Banach space and  $X(t, \cdot) : \Omega \rightarrow 2^Y$  has measurable graph, by Aumann's selection theorem (see Appendix) it follows that there exists a function  $f(t, \cdot) : \Omega \rightarrow Y$  such that  $f(t, \omega) \in X(t, \omega)$   $\mu - a.e.$  Since  $X(t, \cdot)$  is integrably bounded, we have that  $f(t, \cdot) \in L_1(\mu, Y)$ , hence  $S_{X_t}$  is non-empty and  $S_X = \prod_{t \in T} S_{X_t}$  is non-empty.  $S_{X_t}^1$  is convex and  $S_X^1$  is also convex. Since  $X(t, \cdot) : \Omega \rightarrow 2^Y$  is integrably bounded and has convex weakly compact values, by Diestel's Theorem (see Appendix) it follows that  $S_{X_t}^1$  is a weakly compact subset of  $L_1(\mu, Y)$  and so is  $S_X^1$ . We have that  $S_X^1$  is a metrizable set as being a weakly compact subset of the separable Banach space  $L_1(\nu, L_1(\mu, Y))$  (Dunford-Schwartz [1958, p 434]).

Let denote  $A^* : T \times \Omega \times S_X^1 \rightarrow 2^Y$ ,  $A^*((t, \omega, \tilde{x})) = (A(t, \omega, \tilde{x}))_{a(\tilde{x})}$ .

Since all the values of the correspondence  $A^*$  are contained in the compact set  $X(\cdot, \cdot)$  and  $A^*$  is closed and convex valued (hence weakly closed), it follows that  $A^*$  is weakly compact valued.

Then  $\Phi$  is convex valued and for each  $(t, \omega) \in T \times \Omega$ ,  $\Phi(t, \omega, \cdot)$  is upper semicontinuous. We have that  $\Phi(\cdot, \cdot, \tilde{x})$  has a measurable graph for each  $\tilde{x} \in S_X^1$ . Let  $U = \{(t, \omega, \tilde{x}) \in \Omega \times S_X^1 : \Phi(t, \omega, \tilde{x}) \neq \emptyset\}$ . For each  $\tilde{x} \in S_X^1$ , let  $U^x = \{(t, \omega) \in T \times \Omega : \Phi(t, \omega, \tilde{x}) \neq \emptyset\}$  and for each  $\omega \in \Omega$ , let  $U^{(t, \omega)} = \{\tilde{x} \in S_X^1 : \Phi(t, \omega, \tilde{x}) \neq \emptyset\}$ . Define  $G : T \times \Omega \times S_X^1 \rightarrow 2^Y$  by 
$$G(t, \omega, \tilde{x}) = \begin{cases} \Phi(t, \omega, \tilde{x}) & \text{if } (t, \omega, \tilde{x}) \in U \\ (A(t, \omega, \tilde{x}))_{a(\tilde{x})} & \text{if } (t, \omega, \tilde{x}) \notin U. \end{cases}$$

For each  $\tilde{x} \in S_X^1$ , the correspondence  $G(\cdot, \cdot, \tilde{x})$  has a measurable graph.

By assumption A5) (b), the set  $U^{(t, \omega)} = \{\tilde{x} \in S_X^1 : (A(t, \omega, \tilde{x}))_{a(\tilde{x})} \cap (P(t, \omega, \tilde{x}))_{p(\tilde{x})} \neq \emptyset\}$  is weakly open in  $S_X^1$ . For each  $(t, \omega) \in T \times \Omega$ ,  $G(t, \omega, \cdot) : S_X^1 \rightarrow 2^Y$  is upper semi-continuous.

Let  $V$  be weakly open in  $S_X^1$  and  $\tilde{x} \in S_X^1$ .

$$\begin{aligned} W &= \{\tilde{x} \in S_X^1 : G(t, \omega, \tilde{x}) \subset V\} = \\ &= \{\tilde{x} \in U^{(t, \omega)} : G(t, \omega, \tilde{x}) \subset V\} \cup \{\tilde{x} \in S_X^1 \setminus U^{(t, \omega)} : G(t, \omega, \tilde{x}) \subset V\} \\ &= \{\tilde{x} \in U^{(t, \omega)} : \Phi(t, \omega, \tilde{x}) \subset V\} \cup \{\tilde{x} \in S_X^1 : G(t, \omega, \tilde{x}) \subset V\}. \end{aligned}$$

$W$  is an open set, because  $U^{(t, \omega)}$  is open,  $\Phi(t, \omega, \cdot)$  is a upper semicontinuous map on  $U^{(t, \omega)}$  and the set  $\{\tilde{x} \in S_X^1 : G(t, \omega, \tilde{x}) \subset V\}$  is open since  $G(t, \omega, \cdot)$  is u.s.c. Moreover,  $G$  is convex and non-empty valued.

Define the correspondence  $\varphi : T \times S_X^1 \rightarrow 2^{L_1(\mu, Y)}$  by  $\varphi(t, \tilde{x}) = \{\tilde{y}(t) \in L_1(\mu, Y) : \tilde{y}(t, \omega) \in G(t, \omega, \tilde{x}) \mu - a.e.\} \cap S_{X_t}^1$ .

By the measurability lifting theorem (Theorem 16, see Appendix), the correspondence  $t \rightarrow \{\tilde{y}(t) \in L_1(\mu, Y) : \tilde{y}(t, \omega) \in G(t, \omega, \tilde{x}) \mu - a.e.\}$  has a measurable graph and so does  $t \rightarrow S_{X_t}^1$  by (A.4). Thus, for each fixed  $\tilde{x} \in S_X^1$ ,  $\varphi(\cdot, \tilde{x})$  has a measurable graph.

Since for each fixed  $\tilde{x} \in S_X^1$ ,  $G(\cdot, \cdot, \tilde{x})$  has a measurable graph and it is nonempty valued, then by the Aumann measurable selection theorem (see Appendix), it admits a measurable selection and we can conclude that  $\varphi$  is nonempty valued. It follows by the u.s.c. lifting theorem that for each fixed  $t$ ,  $\varphi(t, \cdot)$  is weakly u.s.c.

Define  $G' : S_X^1 \rightarrow 2^{S_X^1}$ , by  $G'(\tilde{x}) = \{\tilde{y} \in S_X^1 : \tilde{y}(t) \in \varphi(t, \tilde{x}) \nu - a.e.\}$ .

Another application of the u.s.c. lifting theorem enables us to conclude that  $G'$  is a weakly u.s.c. correspondence which is obviously convex valued (since  $\varphi$  is convex valued) and also nonempty valued (recall once more the Aumann measurable selection theorem and notice that the set  $S_X^1$  is metrizable).

$G'$  is an upper semicontinuous correspondence and has also non-empty convex closed values.

By Fan-Glicksberg's fixed-point theorem in [6], there exists  $\tilde{x}^* \in S_X^1$  such that  $\tilde{x}^* \in G'(\tilde{x}^*)$ . It follows that  $\tilde{x}^* \in S_X^1$  and  $\tilde{x}^*(t) \in \varphi(t, \tilde{x}^*) \nu - a.e.$  Thus, we have that  $\tilde{x}^* \in S_X^1$  and  $\tilde{x}^*(t, \omega) \in G(t, \omega, \tilde{x}^*) \mu - a.e., \nu - a.e.$

By assumption A.4.a, it follows that  $\tilde{x}^*(t, \omega) \notin (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)} \cap (P(t, \omega, \tilde{x}^*))_{p(\tilde{x}^*)}$ , then we have that  $\tilde{x}^* \notin U$  and  $\tilde{x}^*(t, \omega) \in (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)}$ .

Therefore, for  $\nu - a.e.$  :

- i)  $\tilde{x}^*(t, \omega) \in (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)} \mu - a.e.$ ;
- ii)  $(A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)} \cap (P(t, \omega, \tilde{x}^*))_{p(\tilde{x}^*)} = \emptyset \mu - a.e. \square$



#### 4. RANDOM QUASI-VARIATIONAL INEQUALITIES

In this section, we are establishing new random quasi-variational inequalities with random fuzzy mappings and random fixed point theorems. The proofs rely on the theorem of Bayesian fuzzy equilibrium existence for the abstract fuzzy economy with a measure space of agents.

This is our first theorem.

**Theorem 4.** *Let  $(T, \tau, \nu)$  be a measure space,  $(\Omega, F, \mu)$  be a complete finite separable measure space, and let  $Y$  be a separable Banach space. Suppose that the following conditions are satisfied:*

A.1)

(a)  $X : T \times \Omega \rightarrow F(Y)$  and  $z \in (0, 1]$  are such that  $(t, \omega) \rightarrow (X(t, \omega))_z : T \times \Omega \rightarrow 2^Y$  is a non-empty convex weakly compact-valued and integrably bounded correspondence;

(b) For each fixed  $t \in T$ ,  $F_t$  is a sub  $\sigma$ -algebra of  $F$  and  $(X(t, \cdot))_z$  has an  $F_t$ -measurable graph;

A.2)

$A : T \times \Omega \times S_X^1 \rightarrow \mathcal{F}(Y)$  and  $a : S_X^1 \rightarrow (0, 1]$  are such that

(a) the correspondence  $(t, \omega, \tilde{x}) \rightarrow (A(t, \omega, \tilde{x}))_{a(\tilde{x})} : T \times \Omega \times S_X^1 \rightarrow 2^Y$  has a measurable graph;

(b) For each  $(t, \omega) \in T \times \Omega$ ,  $\tilde{x} \rightarrow (A(t, \omega, \tilde{x}))_{a(\tilde{x})} : S_X^1 \rightarrow 2^Y$  is an upper semicontinuous correspondence with closed, convex and non-empty values.

A.3) The correspondence  $t \rightarrow S_{X_t}^1$  has a measurable graph;

A.4)

$\psi : T \times \Omega \times S_X^1 \times Y \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$  and  $p : S_X^1 \rightarrow (0, 1]$  are such that:

(a)  $(t, \tilde{x}) \rightarrow \{y \in Y : \psi(t, \omega, \tilde{x}, y) > 0\}$  is upper semicontinuous with compact values on  $T \times S_X^1$  for each fixed  $\omega \in \Omega$ ;

(b)  $\tilde{x}(t, \omega) \notin \text{cl}\{y \in Y : \psi(t, \omega, \tilde{x}, y) > 0\}$  for each fixed  $(t, \omega, \tilde{x}) \in T \times \Omega \times S_X^1$ ;

(c) for each  $(t, \omega, \tilde{x}) \in T \times \Omega \times S_X^1$ ,  $\psi(t, \omega, \tilde{x}, \cdot)$  is concave;

(d) for each  $(t, \omega) \in T \times \Omega$ ,  $\{\tilde{x} \in S_X^1 : \alpha(t, \omega, \tilde{x}) > 0\}$  is weakly open in  $S_X^1$ , where  $\alpha : T \times \Omega \times S_X^1 \rightarrow \mathbb{R}$  is defined by  $\alpha(t, \omega, \tilde{x}) = \sup_{y \in (A(t, \omega, \tilde{x}))_{a(\tilde{x})}} \psi(t, \omega, \tilde{x}, y)$  for each  $(t, \omega, \tilde{x}) \in T \times \Omega \times S_X^1$ ;

(d)  $\{(t, \omega, \tilde{x}) : \psi(t, \omega, \tilde{x}, y) > 0\} \in \tau \otimes \mathcal{F}_t \otimes B(S_X^1)$ .

Then, there exists  $\tilde{x}^* \in S_X^1$  such that  $\nu - a.e.$ ,

- i)  $\tilde{x}^*(t, \omega) \in (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)} \mu - a.e;$
- ii)  $\sup_{y \in (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)}} \psi(t, \omega, \tilde{x}^*, y) \leq 0 \mu - a.e.$

*Proof.* Let  $P : T \times \Omega \times S_X^1 \rightarrow \mathcal{F}(Y)$  and  $p : S_X^1 \rightarrow (0, 1]$  such that  $(P(t, \omega, \tilde{x}))_{p(\tilde{x})} = \{y \in Y : \psi(t, \omega, \tilde{x}, y) > 0\}$  for each  $(t, \omega, \tilde{x}) \in T \times \Omega \times S_X^1$ .

We shall show that the abstract fuzzy economy  $G = \{(X, F_t, A, P, a, p, z), t \in T\}$  satisfies all hypotheses of Theorem 4.

Suppose  $(t, \omega) \in T \times \Omega$ .

According to A4 a), we have that for each  $(t, \omega) \in T \times \Omega$ ,  $\tilde{x} \rightarrow (P(t, \omega, \tilde{x}))_{p(\tilde{x})} : S_X^1 \rightarrow 2^Y$  is an upper semicontinuous correspondence with closed, convex and non-empty values and according to A4 b),  $\tilde{x}(t, \omega) \notin (P(t, \omega, \tilde{x}))_{p(\tilde{x})}$  for each  $\tilde{x} \in S_X^1$ . Assumption A4 c) implies that  $\tilde{x} \rightarrow (P(t, \omega, \tilde{x}))_{p(\tilde{x})} : S_X^1 \rightarrow 2^Y$  has convex values.

According to the definition of  $\alpha$ , we note that  $\{\tilde{x} \in S_X^1 : (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)} \cap (P(t, \omega, \tilde{x}^*))_{p(\tilde{x}^*)} \neq \emptyset\} = \{\tilde{x} \in S_X^1 : \alpha(t, \omega, \tilde{x}) > 0\}$  so that  $\{\tilde{x} \in S_X^1 : (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)} \cap (P(t, \omega, \tilde{x}^*))_{p(\tilde{x}^*)} \neq \emptyset\}$  is weakly open in  $S_X^1$  by A4d).

According to A2 a) and A4 e), it follows that the correspondences  $(t, \omega, \tilde{x}) \rightarrow (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)}$  and  $(t, \omega, \tilde{x}) \rightarrow (P(t, \omega, \tilde{x}^*))_{p(\tilde{x}^*)}$  have measurable graphs.

Thus the abstract fuzzy economy  $G = \{(X, F_t, A, P, a, p, z), t \in T\}$  satisfies all hypotheses of Theorem 3. Therefore, there exists  $\tilde{x}^* \in S_X^1$  such that  $v - a.e. :$

- $x^*(t, \omega) \in (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)} \mu - a.e$  and
- $(A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)} \cap (P(t, \omega, \tilde{x}^*))_{p(\tilde{x}^*)} = \emptyset \mu - a.e;$
- that is, there exists  $\tilde{x}^* \in S_X^1$  such that  $v - a.e. :$
- i)  $x^*(t, \omega) \in (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)} \mu - a.e;$
- ii)  $\sup_{y \in (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)}} \psi(t, \omega, \tilde{x}^*, y) \leq 0 \mu - a.e.$

If  $|\Gamma|=1$ , we obtain the following corollary.

**Corollary 5.** *Let  $(\Omega, F, \mu)$  be a complete finite separable measure space, and let  $Y$  be a separable Banach space. Suppose that the following conditions are satisfied:*

A.1)

(a)  $X : \Omega \rightarrow F(Y)$  and  $z \in (0, 1]$  are such that  $\omega \rightarrow (X(\omega))_z : T \times \Omega \rightarrow 2^Y$  is a non-empty convex weakly compact-valued and integrably bounded correspondence;

(b)  $(X(\cdot))_z$  has an  $F$ -measurable graph;

A.2)

$A : \Omega \times S_X^1 \rightarrow \mathcal{F}(Y)$  and  $a : S_X^1 \rightarrow (0, 1]$  are such that

(a) the correspondence  $(\omega, \tilde{x}) \rightarrow (A(\omega, \tilde{x}))_{a(\tilde{x})} : \Omega \times S_X^1 \rightarrow 2^Y$  has a measurable graph;

(b) For each  $\omega \in \Omega$ ,  $\tilde{x} \rightarrow (A(\omega, \tilde{x}))_{a(\tilde{x})} : S_X^1 \rightarrow 2^Y$  is an upper semicontinuous correspondence with closed, convex and non-empty values.

A.3)

$\psi : \Omega \times S_X^1 \times Y \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$  and  $p : S_X^1 \rightarrow (0, 1]$  are such that:

(a)  $\tilde{x} \rightarrow \{y \in Y : \psi(\omega, \tilde{x}, y) > 0\}$  is upper semicontinuous with compact values on  $S_X^1$  for each fixed  $\omega \in \Omega$ ;

(b)  $\tilde{x}(\omega) \notin \text{cl}\{y \in Y : \psi(\omega, \tilde{x}, y) > 0\}$  for each fixed  $(\omega, \tilde{x}) \in \Omega \times S_X^1$ ;

(c) for each  $(\omega, \tilde{x}) \in \Omega \times S_X^1$ ,  $\psi(\omega, \tilde{x}, \cdot)$  is concave;

(d) for each  $\omega \in \Omega$ ,  $\{\tilde{x} \in S_X^1 : \alpha(\omega, \tilde{x}) > 0\}$  is weakly open in  $S_X^1$ , where  $\alpha : \Omega \times S_X^1 \rightarrow \mathbb{R}$  is defined by  $\alpha(\omega, \tilde{x}) = \sup_{y \in (A(\omega, \tilde{x}))_{a(\tilde{x})}} \psi(\omega, \tilde{x}, y)$  for each  $(\omega, \tilde{x}) \in \Omega \times S_X^1$ ;

(d)  $\{(\omega, \tilde{x}) : \psi(\omega, \tilde{x}, y) > 0\} \in \mathcal{F}_t \otimes B(S_X^1)$ .

Then, there exists  $\tilde{x}^* \in S_X^1$  such that:

i)  $\tilde{x}^*(t, \omega) \in (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)}$   $\mu - a.e.$ ;

ii)  $\sup_{y \in (A(\omega, \tilde{x}^*))_{a(\tilde{x}^*)}} \psi(\omega, \tilde{x}^*, y) \leq 0$   $\mu - a.e.$

As a consequence of Theorem 4, we prove the following random generalized quasi-variational inequality with random fuzzy mappings. Our result generalizes Theorem 5.1 in [22].

**Theorem 6.** Let  $(T, \tau, \nu)$  be a measure space,  $(\Omega, F, \mu)$  be a complete finite separable measure space, and let  $Y$  be a separable Banach space. Suppose that the following conditions are satisfied:

A.1)

(a)  $X : T \times \Omega \rightarrow \mathcal{F}(Y)$  and  $z \in (0, 1]$  are such that  $(t, \omega) \rightarrow (X(t, \omega))_z : T \times \Omega \rightarrow 2^Y$  is a non-empty convex weakly compact-valued and integrably bounded correspondence;

(b) For each fixed  $t \in T$ ,  $F_t$  is a sub  $\sigma$ -algebra of  $F$  and  $(X(t, \cdot))_z$  has an  $F_t$ -measurable graph;

A.2)

$A : T \times \Omega \times S_X^1 \rightarrow \mathcal{F}(Y)$  and  $a : S_X^1 \rightarrow (0, 1]$  are such that

(a) the correspondence  $(t, \omega, \tilde{x}) \rightarrow (A(t, \omega, \tilde{x}))_{a(\tilde{x})} : T \times \Omega \times S_X^1 \rightarrow 2^Y$  has a measurable graph;

(b) For each  $(t, \omega) \in T \times \Omega$ ,  $\tilde{x} \rightarrow (A(t, \omega, \tilde{x}))_{a(\tilde{x})} : S_X^1 \rightarrow 2^Y$  is an upper semicontinuous correspondence with closed, convex and non-empty values.

A.3) The correspondence  $t \rightarrow S_{X_t}^1$  has a measurable graph;

A.4)

$G : T \times \Omega \times Y \rightarrow \mathcal{F}(Y')$  and  $g : Y \rightarrow (0, 1]$  are such that:

(a) for each  $(t, \omega) \in T \times \Omega$ ,  $y \rightarrow (G(t, \omega, y))_{g(y)} : Y \rightarrow 2^{Y'}$  is monotone with non-empty values;

(b) for each  $(t, \omega) \in T \times \Omega$ ,  $y \rightarrow (G(t, \omega, y))_{g(y)} : L \cap Y \rightarrow 2^{Y'}$  is lower semicontinuous from the relative topology of  $Y$  into the weak\*-topology  $\sigma(Y', Y)$  of  $Y'$  for each one-dimensional flat  $L \subset Y$ ;

A.5)

(a)  $f : T \times \Omega \times S_X^1 \times Y \rightarrow \mathbb{R} \cup \{\infty, -\infty\}$  is such that  $\tilde{x} \rightarrow f(t, \omega, \tilde{x}, y)$  is lower semicontinuous on  $S_X^1$  for each fixed  $(t, \omega, y) \in T \times \Omega \times Y$ ,  $f(t, \omega, \tilde{x}, \tilde{x}(t, \omega)) = 0$  for each  $(t, \omega, \tilde{x}) \in T \times \Omega \times S_X^1$  and  $y \rightarrow f(t, \omega, \tilde{x}, y)$  is concave on  $Y$  for each fixed  $(t, \omega, x) \in T \times \Omega \times S_X^1$ ;

(b) for each fixed  $(t, \omega) \in T \times \Omega$ , the set

$\{\tilde{x} \in S_X^1 : \sup_{y \in (A(t, \omega, \tilde{x}))_{a(\tilde{x})}} [\sup_{u \in (G(t, \omega, y))_{g(y)}} \text{Re}\langle u, \tilde{x} - y \rangle + f(t, \omega, \tilde{x}, y)] > 0\}$  is weakly open in  $S_X^1$

(c)  $\{(t, \omega, \tilde{x}) : \sup_{u \in (G(t, \omega, y))_{g(y)}} \text{Re}\langle u, \tilde{x} - y \rangle + f(t, \omega, \tilde{x}, y) > 0\} \in \tau \otimes \mathcal{F}_t \otimes B(S_X^1)$ .

Then, there exists  $\tilde{x}^* \in S_X^1$  such that  $\nu$ -a.e.:

i)  $\tilde{x}^*(t, \omega) \in (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)}$   $\mu$ -a.e.;

ii)  $\sup_{u \in (G(t, \omega, y))_{g(y)}} \text{Re}\langle u, \tilde{x}^*(\omega) - y \rangle + f(t, \omega, \tilde{x}^*, y) \leq 0$  for all  $y \in (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)}$   $\mu$ -a.e.

*Proof.* Let us define  $\psi : T \times \Omega \times S_X^1 \times Y \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$  by

$\psi(t, \omega, \tilde{x}, y) = \sup_{u \in (G(t, \omega, y))_{g(y)}} \text{Re}\langle u, \tilde{x} - y \rangle + f(t, \omega, \tilde{x}, y)$  for each  $(t, \omega, \tilde{x}, y) \in T \times \Omega \times S_X^1 \times Y$ .

According to assumption A5 a),  $\tilde{x} \rightarrow f(t, \omega, \tilde{x}, y)$  is lower semicontinuous on  $S_X^1$  for each fixed  $(t, \omega, y) \in T \times \Omega \times Y$  and  $f(t, \omega, \tilde{x}, \tilde{x}(t, \omega)) = 0$  for each  $(t, \omega, \tilde{x}) \in T \times \Omega \times S_X^1$  implies that  $\tilde{x}(t, \omega) \notin \{y \in Y : \psi(t, \omega, \tilde{x}, y) > 0\}$  for each fixed  $(t, \omega, \tilde{x}) \in T \times \Omega \times S_X^1$ .

We also have that for each  $(t, \omega, \tilde{x}) \in T \times \Omega \times S_X^1$ ,  $\psi(t, \omega, \tilde{x}, \cdot)$  is concave. This fact is a consequence of assumption A4 a).

All the hypotheses of Theorem 4 are satisfied. According to Theorem 4, there exists  $\tilde{x}^* \in S_X^1$  such that  $\nu$ -a.e.

$\tilde{x}^*(t, \omega) \in (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)}$   $\mu$ -a.e.,

and

(1)  $\sup_{y \in (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)}} \sup_{(G(t, \omega, y))_{g(y)}} [\text{Re}\langle u, \tilde{x}^*(t, \omega) - y \rangle + f(t, \omega, \tilde{x}^*, y)] \leq 0$   $\mu$ -a.e.

Finally, we will prove that  $v - a.e.$ ,

$$\sup_{y \in A(t, \omega, \tilde{x}^*)_{a(\tilde{x}^*)}} \sup_{u \in (G(t, \omega, \tilde{x}^*))_{g(\tilde{x}^*)}} [Re \langle u, \tilde{x}^*(\omega) - y \rangle + f(t, \omega, \tilde{x}^*, y)] \leq 0$$

$\mu - a.e.$

In order to do that, let us consider the fixed points  $t \in T$  and  $\omega \in \Omega$ .

Let  $y \in v$ ,  $\lambda \in [0, 1]$  and  $z_\lambda(t, \omega) := \lambda y + (1 - \lambda)\tilde{x}^*(t, \omega)$ . According to assumption A2 b),  $z_\lambda(t, \omega) \in (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)}$ .

According to (1), we have  $\sup_{u \in (G(t, \omega, z_\lambda(t, \omega)))_{g(z_\lambda(t, \omega))}} [Re \langle u, \tilde{x}^*(t, \omega) - z_\lambda(t, \omega) \rangle + f(t, \omega, \tilde{x}^*, z_\lambda(t, \omega))] \leq 0$  for each  $\lambda \in [0, 1]$ .

According to assumption A5 a),  $f(t, \omega, \tilde{x}^*, \tilde{x}^*(t, \omega)) = 0$ . For each  $y_1, y_2 \in Y$  and for each  $\lambda \in [0, 1]$ , we also have that  $f(t, \omega, \tilde{x}^*, \lambda y_1 + (1 - \lambda)y_2) \geq \lambda f(t, \omega, \tilde{x}^*, y_1) + (1 - \lambda)f(t, \omega, \tilde{x}^*, y_2)$ .

Therefore, for each  $\lambda \in [0, 1]$ , we have that

$$\begin{aligned} & t \{ \sup_{u \in (G(t, \omega, z_\lambda(t, \omega)))_{g(z_\lambda(t, \omega))}} [Re \langle u, \tilde{x}^*(t, \omega) - y \rangle + f(t, \omega, \tilde{x}^*, y)] \} \leq \\ & \sup_{u \in (G(t, \omega, z_\lambda(t, \omega)))_{g(z_\lambda(t, \omega))}} t [Re \langle u, \tilde{x}^*(t, \omega) - y \rangle + f(t, \omega, \tilde{x}^*, z_\lambda(t, \omega))] = \\ & \sup_{u \in (G(t, \omega, z_\lambda(t, \omega)))_{g(z_\lambda(t, \omega))}} [Re \langle u, \tilde{x}^*(t, \omega) - z_\lambda(t, \omega) \rangle + f(t, \omega, \tilde{x}^*, z_\lambda(t, \omega))] \leq \\ & 0. \end{aligned}$$

It follows that for each  $\lambda \in [0, 1]$ ,

$$(2) \sup_{u \in (G(t, \omega, z_\lambda(t, \omega)))_{g(z_\lambda(t, \omega))}} [Re \langle u, \tilde{x}^*(t, \omega) - y \rangle + f(t, \omega, \tilde{x}^*, y)] \leq 0.$$

Now, we are using the lower semicontinuity of  $F(t, \omega, \cdot)$  in order to show the conclusion. For each  $z_0 \in (G(t, \omega, \tilde{x}^*(t, \omega)))_{g(\tilde{x}^*(t, \omega))}$  and  $e > 0$  let us consider  $U_{z_0}^t$ , the neighborhood of  $z_0$  in the topology  $\sigma(Y', Y)$ , defined by  $U_{z_0}^t := \{z \in Y' : |Re \langle z_0 - z, \tilde{x}^*(t, \omega) - y \rangle| < e\}$ . As  $y \rightarrow (G(t, \omega, y))_{g(y)} : L \cap Y \rightarrow 2^{Y'}$  is lower semicontinuous, where  $L = \{z_\lambda(t, \omega) : \lambda \in [0, 1]\}$  and  $U_{z_0}^t \cap (G(t, \omega, \tilde{x}^*(t, \omega)))_{g(\tilde{x}^*(t, \omega))} \neq \emptyset$ , there exists a non-empty neighborhood  $N(\tilde{x}^*(t, \omega))$  of  $\tilde{x}^*(t, \omega)$  in  $L$  such that for each  $z \in N(\tilde{x}^*(t, \omega))$ , we have that  $U_{z_0}^t \cap F(t, \omega, z) \neq \emptyset$ . Then there exists  $\delta \in (0, 1]$ ,  $t \in (0, \delta)$  and  $u \in (G(t, \omega, z_\lambda(t, \omega)))_{g(z_\lambda(t, \omega))} \cap U_{z_0}^t \neq \emptyset$  such that  $Re \langle z_0 - u, \tilde{x}^*(t, \omega) - y \rangle < e$ . Therefore,  $Re \langle z_0, \tilde{x}^*(t, \omega) - y \rangle < Re \langle u, \tilde{x}^*(t, \omega) - y \rangle + e$ .

It follows that

$$Re \langle z_0, \tilde{x}^*(t, \omega) - y \rangle + f(t, \omega, \tilde{x}^*, y) < Re \langle u, \tilde{x}^*(t, \omega) - y \rangle + f(t, \omega, \tilde{x}^*, y) + e < e.$$

The last inequality comes from (2). Since  $e > 0$  and  $z_0 \in (G(t, \omega, \tilde{x}^*(t, \omega)))_{g(\tilde{x}^*(t, \omega))}$  have been chosen arbitrarily, the next relation holds:

$$Re \langle z_0, \tilde{x}^*(t, \omega) - y \rangle + f(t, \omega, \tilde{x}^*, y) < 0.$$

Hence,  $\mu - a.e.$ , we have that  $\sup_{u \in (G(t, \omega, \tilde{x}^*(t, \omega)))_{g(\tilde{x}^*(t, \omega))}} [Re \langle z_0, \tilde{x}^*(t, \omega) - y \rangle + f(t, \omega, \tilde{x}^*, y)] \leq 0$  for every  $y \in (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)}$ .

If  $|\mathbb{T}|=1$ , we can make abstraction of  $\mathbb{T}$  and we obtain the following corollary.

**Theorem 7.** *Let  $(\Omega, F, \mu)$  be a complete finite separable measure space, and let  $Y$  be a separable Banach space. Suppose that the following conditions are satisfied:*

A.1)

(a)  $X : \Omega \rightarrow F(Y)$  and  $z \in (0, 1]$  are such that  $\omega \rightarrow (X(\omega))_z : \Omega \rightarrow 2^Y$  is a non-empty convex weakly compact-valued and integrably bounded correspondence;

(b)  $(X(\cdot, \cdot))_z$  has an  $F$ -measurable graph;

A.2)

$A : \Omega \times S_X^1 \rightarrow \mathcal{F}(Y)$  and  $a : S_X^1 \rightarrow (0, 1]$  are such that

(a) the correspondence  $(\omega, \tilde{x}) \rightarrow (A(\omega, \tilde{x}))_{a(\tilde{x})} : T \times \Omega \times S_X^1 \rightarrow 2^Y$  has a measurable graph;

(b) For each  $\omega \in \Omega$ ,  $\tilde{x} \rightarrow (A(\omega, \tilde{x}))_{a(\tilde{x})} : S_X^1 \rightarrow 2^Y$  is an upper semicontinuous correspondence with closed, convex and non-empty values;

A.3)

$G : \Omega \times Y \rightarrow \mathcal{F}(Y')$  and  $g : Y \rightarrow (0, 1]$  are such that:

(a) for each  $\omega \in \Omega$ ,  $y \rightarrow (G(\omega, y))_{g(y)} : Y \rightarrow 2^{Y'}$  is monotone with non-empty values;

(b) for each  $\omega \in \Omega$ ,  $y \rightarrow (G(\omega, y))_{g(y)} : L \cap Y \rightarrow 2^{Y'}$  is lower semicontinuous from the relative topology of  $Y$  into the weak\*-topology  $\sigma(Y', Y)$  of  $Y'$  for each one-dimensional flat  $L \subset Y$ ;

A.5)

(a)  $f : \Omega \times S_X^1 \times Y \rightarrow \mathbb{R} \cup \{\infty, -\infty\}$  is such that  $\tilde{x} \rightarrow f(\omega, \tilde{x}, y)$  is lower semicontinuous on  $S_X^1$  for each fixed  $(\omega, y) \in \Omega \times Y$ ,  $f(\omega, \tilde{x}, \tilde{x}(t, \omega)) = 0$  for each  $(\omega, \tilde{x}) \in \Omega \times S_X^1$  and  $y \rightarrow f(\omega, \tilde{x}, y)$  is concave on  $Y$  for each fixed  $(\omega, \tilde{x}) \in \Omega \times S_X^1$ ;

(b) for each fixed  $\omega \in \Omega$ , the set

$$\{\tilde{x} \in S_X^1 : \sup_{y \in (A(\omega, \tilde{x}))_{a(\tilde{x})}} [\sup_{u \in (G(\omega, y))_{g(y)}} \operatorname{Re} \langle u, \tilde{x} - y \rangle + f(t, \omega, \tilde{x}, y)] > 0\}$$

is weakly open in  $S_X^1$

$$(c) \{(\omega, \tilde{x}) : \sup_{u \in (G(\omega, y))_{g(y)}} \operatorname{Re} \langle u, \tilde{x} - y \rangle + f(t, \omega, \tilde{x}, y) > 0\} \in \mathcal{F}_t \otimes B(S_X^1).$$

Then, there exists  $\tilde{x}^* \in S_X^1$  such that:

i)  $\tilde{x}^*(\omega) \in (A(\omega, \tilde{x}^*))_{a(\tilde{x}^*)}$   $\mu - a.e.$ ;

ii)  $\sup_{u \in (G(\omega, y))_{g(y)}} \operatorname{Re} \langle u, \tilde{x}^*(\omega) - y \rangle + f(t, \omega, \tilde{x}^*(\omega), y) \leq 0$  for all  $y \in (A(\omega, \tilde{x}^*))_{a(\tilde{x}^*)}$   $\mu - a.e.$

We obtain the following random fixed point theorem with fuzzy mapping as a particular case of Theorem 6.

**Theorem 8.** *Let  $(T, \tau, \nu)$  be a measure space,  $(\Omega, F, \mu)$  be a complete finite separable measure space, and let  $Y$  be a separable Banach space. Suppose that the following conditions are satisfied:*

A.1)

(a)  $X : T \times \Omega \rightarrow F(Y)$  and  $z \in (0, 1]$  are such that  $(t, \omega) \rightarrow (X(t, \omega))_z : T \times \Omega \rightarrow 2^Y$  is a non-empty convex weakly compact-valued and integrably bounded correspondence;

(b) For each fixed  $t \in T$ ,  $F_t$  is a sub  $\sigma$ -algebra of  $F$  and  $(X(t, \cdot))_z$  has an  $F_t$ -measurable graph;

A.2)

$A : T \times \Omega \times S_X^1 \rightarrow \mathcal{F}(Y)$  and  $a : S_X^1 \rightarrow (0, 1]$  are such that

(a) the correspondence  $(t, \omega, \tilde{x}) \rightarrow (A(t, \omega, \tilde{x}))_{a(\tilde{x})} : T \times \Omega \times S_X^1 \rightarrow 2^Y$  has a measurable graph;

(b) For each  $(t, \omega) \in T \times \Omega$ ,  $\tilde{x} \rightarrow (A(t, \omega, \tilde{x}))_{a(\tilde{x})} : S_X^1 \rightarrow 2^Y$  is an upper semicontinuous correspondence with closed, convex and non-empty values.

A.3) The correspondence  $t \rightarrow S_{X_t}^1$  has a measurable graph;

Then, there exists  $\tilde{x}^* \in S_X^1$  such that  $\nu - a.e.$ ,  $\tilde{x}^*(t, \omega) \in (A(t, \omega, \tilde{x}^*))_{a(\tilde{x}^*)}$   $\mu - a.e.$

If  $|T|=1$ , we obtain the following result.

**Theorem 9.** *Let  $(\Omega, F, \mu)$  be a complete finite separable measure space, and let  $Y$  be a separable Banach space. Suppose that the following conditions are satisfied:*

A.1)

(a)  $X : \Omega \rightarrow F(Y)$  and  $z \in (0, 1]$  are such that  $\omega \rightarrow (X(\omega))_z : T \times \Omega \rightarrow 2^Y$  is a non-empty convex weakly compact-valued and integrably bounded correspondence;

(b)  $(X(\cdot))_z$  has an  $F$ -measurable graph;

A.2)

$A : \Omega \times S_X^1 \rightarrow \mathcal{F}(Y)$  and  $a : S_X^1 \rightarrow (0, 1]$  are such that

(a) the correspondence  $(\omega, \tilde{x}) \rightarrow (A(\omega, \tilde{x}))_{a(\tilde{x})} : \Omega \times S_X^1 \rightarrow 2^Y$  has a measurable graph;

(b) For each  $\omega \in \Omega$ ,  $\tilde{x} \rightarrow (A(\omega, \tilde{x}))_{a(\tilde{x})} : S_X^1 \rightarrow 2^Y$  is an upper semicontinuous correspondence with closed, convex and non-empty values.  
Then, there exists  $\tilde{x}^* \in S_X^1$  such that  $\tilde{x}^*(\omega) \in (A(\omega, \tilde{x}))_{a(\tilde{x})}$   $\mu - a.e.$

## 5. APPENDIX

The results below have been used in the proof of our theorems. For more details and further references see the paper quoted.

**Theorem 10.** (*Projection theorem*). Let  $(\Omega, F, \mu)$  be a complete, finite measure space, and  $Y$  be a complete separable metric space. If  $H$  belongs to  $F \otimes \beta(Y)$ , its projection  $\text{Proj}_\Omega(H)$  belongs to  $F$ .

**Theorem 11.** (*Aumann measurable selection theorem [24]*). Let  $(\Omega, F, \mu)$  be a complete finite measure space,  $Y$  be a complete, separable metric space and  $T : \Omega \rightarrow 2^Y$  be a nonempty valued correspondence with a measurable graph, i.e.,  $G_T \in F \otimes \beta(Y)$ . Then there is a measurable function  $f : \Omega \rightarrow Y$  such that  $f(\omega) \in T(\omega)$   $\mu - a.e.$

**Theorem 12.** (*Diestel's Theorem [24, Theorem 3.1]*). Let  $(\Omega, F, \mu)$  be a complete finite measure space,  $X$  be a separable Banach space and  $T : \Omega \rightarrow 2^Y$  be an integrably bounded, convex, weakly compact and nonempty valued correspondence. Then  $S_T = \{x \in L_1(\mu, Y) : x(\omega) \in T(\omega) \mu\text{-a.e.}\}$  is weakly compact in  $L_1(\mu, Y)$ .

**Theorem 13.** (*Carathéodory-type selection theorem [14]*). Let  $(\Omega, F, \mu)$  be a complete measure space,  $Z$  be a complete separable metric space and  $Y$  a separable Banach space. Let  $X : \Omega \rightarrow 2^Y$  be a correspondence with a measurable graph, i.e.,  $G_X \in F \otimes \beta(Y)$  and let  $T : \Omega \times Z \rightarrow 2^Y$  be a convex valued correspondence (possibly empty) with a measurable graph, i.e.,  $G_T \in F \otimes \beta(Z) \otimes \beta(Y)$  where  $\beta(Y)$  and  $\beta(Z)$  are the Borel  $\sigma$ -algebras of  $Y$  and  $Z$ , respectively.

Suppose that:

- (1) for each  $\omega \in \Omega$ ,  $T(\omega, x) \subset X(\omega)$  for all  $x \in Z$ .
- (2) for each  $\omega \in \Omega$ ,  $T(\omega, \cdot)$  has open lower sections in  $Z$ , i.e., for each  $\omega \in \Omega$  and  $y \in Y$ ,  $T^{-1}(\omega, y) = \{x \in Z : y \in T(\omega, x)\}$  is open in  $Z$ .



(3) for each  $(\omega, x) \in \Omega \times Z$ , if  $T(\omega, x) \neq \emptyset$ , then  $T(\omega, x)$  has a non-empty interior in  $X(\omega)$ .

Let  $U = \{(\omega, x) \in \Omega \times Z : T(\omega, x) \neq \emptyset\}$  and for each  $x \in Z$ ,  $U^x = \{\omega \in \Omega : (\omega, x) \in U\}$  and for each  $\omega \in \Omega$ ,  $U^\omega = \{x \in Z : (\omega, x) \in U\}$ . Then for each  $x \in Z$ ,  $U^x$  is a measurable set in  $\Omega$  and there exists a Caratheodory-type selection from  $T|_U$ , i.e., there exists a function  $f : U \rightarrow Y$  such that  $f(\omega, x) \in T(\omega, x)$  for all  $(\omega, x) \in U$ , for each  $x \in Z$ ,  $f(\cdot, x)$  is measurable on  $U^x$  and for each  $\omega \in \Omega$ ,  $f(\omega, \cdot)$  is continuous on  $U^\omega$ . Moreover,  $f(\cdot, \cdot)$  is jointly measurable.

**Theorem 14.** (*U. s. c. Lifting Theorem. [24]*). Let  $Y$  be a separable space,  $(\Omega, F, \mu)$  be a complete finite measure space and  $X : \Omega \rightarrow 2^Y$  be an integrably bounded, nonempty, convex valued correspondence such that for all  $\omega \in \Omega$ ,  $X(\omega)$  is a weakly compact, convex subset of  $Y$ . Denote by  $S_X$  the set  $\{x \in L_1(\mu, Y) : x(\omega) \in X(\omega) \mu - a.e.\}$ . Let  $T : \Omega \times S_X \rightarrow 2^Y$  be a nonempty, closed, convex valued correspondence such that  $T(\omega, x) \subset X(\omega)$  for all  $(\omega, x) \in \Omega \times S_X$ . Assume that for each fixed  $x \in S_X$ ,  $T(\cdot, x)$  has a measurable graph and that for each fixed  $\omega \in \Omega$ ,  $T(\omega, \cdot) : S_X \rightarrow 2^Y$  is u.s.c. in the sense that the set  $\{x \in S_X : T(\omega, x) \subset V\}$  is weakly open in  $S_X$  for every norm open subset  $V$  of  $Y$ . Define the correspondence  $\Phi : S_X \rightarrow 2^{S_X}$  by

$$\Phi(x) = \{y \in S_X : y(\omega) \in T(\omega, x) \mu - a.e.\}.$$

Then  $\Phi$  is weakly u.s.c., i.e., the set  $\{x \in S_X : \Phi(x) \subset V\}$  is weakly open in  $S_X$  for every weakly open subset  $V$  of  $S_X$ .

**Theorem 15.** (*Measurability Lifting Theorem*) [1]. Let  $Y$  and  $E$  be separable Banach spaces, and  $(T, \tau, \nu)$  and  $(\Omega, F, \mu)$  be finite complete separable measure spaces. Let  $\gamma : T \times \Omega \times E \rightarrow 2^Y$  be a nonempty valued correspondence. Suppose that for each  $y \in E$ ,  $\gamma(\cdot, \cdot, y)$  has a measurable graph. Define the correspondence  $\psi : \Omega \times E \rightarrow 2^{L_1(\mu, Y)}$  by

$$\psi(t, y) = \{x(t) \in L_1(\mu, Y) : x(t, \omega) \in \gamma(t, \omega, y) \mu - a.e.\}.$$

Then for each  $y \in E$ ,  $\psi(\cdot, y)$  has a measurable graph.

## References

- [1] E.J.Balder, N.C. Yannelis, Equilibrium in random and Bayesian games with a continuum of players, M.A.Khan and N.C. Yannelis (eds.), Equilibrium Theory in Infinite Dimensional Spaces, Springer 1991

- [2] S.-S. Chang and K. K. Tan, *Equilibria and maximal elements of abstract fuzzy economies and qualitative fuzzy games*, Fuzzy Sets and Systems **125** (2002), 389-399.
- [3] G. Debreu, A social equilibrium existence theorem, Proc. Natl. Acad. Sci. USA **38** (1952), 886-903.
- [4] J. Diestel, J. Uhl, Vector Measures, Mathematical Surveys, Vol. 15. Amer. Math. Soc., Providence 1977
- [5] N. Dunford, J.T. Schwartz, Linear Operators, Vol I Interscience, New York 1958
- [6] K. Fan, Fixed-point and minimax theorems in locally convex topological linear spaces, Proc. Nat. Acad. Sci. U.S.A. **38** (1952), 121-126.
- [7] J. Gwinner and F. Raciti, On a class of random variational inequalities on random sets, Numerical Functional Analysis and Optimization, 27(5-6):619-636, 2006
- [8] J. Gwinner and F. Raciti, Some equilibrium problems under uncertainty and random variational inequalities, Ann. Oper. Res. (2012), 200: 299-319.
- [9] N.-J. Huang, *A new equilibrium existence theorem for abstract fuzzy economies*, Appl. Math. Letters **12** (1999), 1-5.
- [10] N.-J. Huang, *Existence of equilibrium for generalized abstract fuzzy economies*, Fuzzy Sets and Systems **117** (2001), 151-156.
- [11] W. K. Kim and K. H. Lee, *Fuzzy fixed point and existence of equilibria of fuzzy games*, J. Fuzzy Math. **6** (1998), 193-202.
- [12] W. K. Kim and K. H. Lee, *Existence of equilibria in generalized fuzzy games*, J. Chungcheong Math. Soc. **12** (1999), 53-61.
- [13] W. K. Kim and K. H. Lee, *Generalized fuzzy games and fuzzy equilibria*, Fuzzy Sets and Systems **122** (2001), 293-301.
- [14] T. Kim, K. Prikry and N.C. Yannelis, On a Carathéodory-type selection theorem, J. Math. Anal. Appl. 135 (1988), 664-670.

- [15] Noor, M.A. and Elsanousi, S.A., Iterative algorithms for random variational inequalities, Pan Amer. Math. Journal 3 (1993), 39-50.
- [16] M. Patriche, Bayesian abstract economy with a measure space of agents, Abstract and Applied Analysis, Volume 2009 (2009), Article ID 523619, doi:10.1155/2009/52361
- [17] M. Patriche, *Equilibria for free abstract fuzzy economies*, An. St. Ovidius University of Constanta, Volume 17(2), 2009, 143-154.
- [18] M. Patriche, *Abstract fuzzy economies and fuzzy equilibrium pairs*, Mathematical Reports, 12(62), 4 (2010), 373 – 382
- [19] M. Patriche, *Existence of fuzzy equilibria for fuzzy abstract economies with  $Q^1$ -majorized correspondences*, Recent Researches in Neural Networks, Fuzzy Systems, Evolutionary Computing and Automation, International Conference on Fuzzy Systems, 11 – 13 April 2011, Brasov 2011.
- [20] M. Patriche, Equilibrium in games and competitive economies, The Publishing House of the Romanian Academy, 2011;
- [21] W. Shafer and H. Sonnenschein, Equilibrium in abstract economies without ordered preferences. J. Math. Econom. **2** (1975), 345-348.
- [22] K.-K. Tan and G. X.-Z. Yuan, Random equilibria of random generalized games with applications to non-compact random quasi-variational inequalities, Topological Methods in Nonlinear Analysis, Journal of the Juliusz Schauder Center, Volume 5, 1995, 59–82
- [23] L Wang, Y Je Cho, N Huang, *The robustness of generalized abstract fuzzy economies in generalized convex spaces*, Fuzzy Sets and Systems, 176 (1) (2011), 56-63.
- [24] N.C. Yannelis, On the Existence of Bayesian Nash Equilibrium. In: Agramovich, Y., Avgerinos, E., Yannelis, N.C. eds., Functional Analysis and Economic Theory, Springer-Verlag 1998
- [25] N. C. Yannelis and N. D. Prabhakar, Existence of maximal elements and equilibrium in linear topological spaces, J. Math. Econom. **12** (1983), 233-245.

- [26] N.C. Yannelis, Integration of Banach-valued correspondences. In:Khan, M.A., Yannelis, N.C. eds, Equilibrium Theory in Infinite Dimensional Spaces (Studies in Economic Theory). Springer-Verlag 1991b
- [27] X. Z. Yuan, The Study of Minimax inequalities and Applications to Economies and Variational inequalities, Memoirs of the American Society, vol **132**, (1988), Nr 625.
- [28] X.-Z. Yuan and J.-M. Roy, Some new deterministic and random variational inequalities and their applications, Journal of Applied Mathematics and Stochastic Analysis **8** (4) (1995), 381-391.
- [29] L. A. Zadeh, *Fuzzy sets*, Inform and Control **8** (1965), 338-353.