

Dimension Four Wins the Same Game as the Standard Model Group

H.B. Nielsen *

The Niels Bohr Institute, Copenhagen, Denmark

Abstract

In a previous article Don Bennett and I looked for, found and proposed a game in which the Standard Model Gauge Group $S(U(2) \times U(3))$ gets singled out as the “winner”. This “game” means that the by Nature chosen gauge group should be just the one that has the maximal value for a quantity which is a modification of the ratio of the quadratic Casimir for the adjoint representation divided by that for a “smallest” faithful representation. Here I propose to extend this “game” to construct a corresponding game between different potential dimensions for space-time. The idea is to formulate, how the same competition as the one between the potential gauge groups would run out, if restricted to the potential Lorentz or Poincare groups achievable for different dimensions of space-time d . The remarkable point is that it is the experimental space-time *dimension 4 which wins*. So the same function defined over Lie groups seems to single out *both* the gauge group *and* the dimension of space time in nature. This seems a rather strange coincidence, unless there really is some similar physical reason behind causing our game-variable (or goal variable) to be selected to be maximal. It has crudely to do with that the groups preferred are easily represented on very “small” but faithful representations.

1 Introduction

The main idea of the present series of articles is to seek some game, that at the same time can select out the gauge group observed in nature - let us suppose we should have the Standard Model Group $S(U(2) \times U(3))$ in nature, say - and *also* the gauge group (whatever that means) of the (gravitational) general relativity by saying that nature has chosen the “winner” in this game. That is to say we look for a group-characteristic quantity (“goal quantity”) which happens to be largest possible for both the gauge group of the Standard Model and a group associate with the Lorentz transformations, the Lorentz group say. We could then claim that such a goal quantity specifies both the gauge group for the Standard Model say and the Lorentz group meaning the dimension of space time. If the quantity is reasonably simple this could be an explanation for both the gauge group and the dimension of space time. In the previous article[1] we sought in this way to invent a game or rather a “goal quantity” - which were at first the ratio C_A/C_F of the quadratic Casimirs for the group in question for the adjoint representation to the quadratic Casimir of some “small” but still faithful representation- in such a way that this ratio would take its largest value for the by nature chosen (gauge) group. Actually N. Brene and I had already earlier proposed another game that essentially pointed also to the Standard Model Gauge group being the winner [6], but it is the more recent proposal with the quadratic

* hbech@nbi.dk, hbechnbi@gmail.com

Cassimirs or rather their ratio C_A/C_F which we seek to generalize to determine the dimension of space time in this article. This concept of the gauge group for general relativity may be a bit imprecise, and so I want at first to simplify a little bit by making a few ad hoc decisions to extract a group that essentially is the gauge group of general relativity, even if we should not have chosen the definition of this concept completely clearly yet.

A first candidate, which is for me rather attractive for the purpose, is simply the Lorentz group meaning the group of Lorentz boosts and rotations.

You could consider the attitude of the present article and the foregoing in the series [1] as attempts to extract the information[7] discussed in the parameters or details of the Standard Model in order to use it for finding the hypothesized more fundamental theory beyond the Standard Model.

1.1 Why the Standard Model Group

The first question, which the present series of articles, namely this and the previous work[1], attacks is “Why did Nature choose just the Standard Model group $S(U(2) \times U(3))$, it could seemingly have chosen among a lot of Lie groups?” Historically this Standard Model group has appeared as pieces arising from different types of interactions, one $U(1)$ -subgroup came from electromagnetic interactions, built in a bit complicated way into what as *group* stands as $U(2)$ (namely $SU(2) \times U(1)$ Lie-algebraic) contains also the weak nuclear forces, and finally the QCD describing strong interaction connected to the $SU(3)$ part of the Standard Gauge group. The Standard model is so to speak phenomenologically found piecewise subalgebra for subalgebra. It is first afterwards or at least by inclusion of further possible pieces of the gauge group that speculations of grand unification models of various types have appeared.

The specification of some appropriate GUT model, say as the simplest and most promising $SU(5)$, together with some breaking scheme, makes up an explanation for the Standard Model gauge Lie algebra and also easily for the gauge *group* - which we take to be implemented as a restriction on which representations are allowed, so that we can indeed claim a phenomenologically accessible element of knowledge - being the one realized in nature. Truly a major part of the success of the GUT $SU(5)$ model is that the representations of the $SU(5)$ gauge group are automatically representations of the subgroup of the $SU(5)$ with the Lie algebra of the Standard Model, and this *subgroup* is just the $S(U(2) \times U(3))$, so that the GUT $SU(5)$ precisely can explain the same restrictions on the allowed representations as can the Liegroup $S(U(2) \times U(3))$. If we can explain this *group* structure and not only the Lie algebra structure of the Standard Model group, then we would have less need for grand unification, because it would mean obtaining similar predictions for the representations of say quarks and leptons under the gauge group. It is simply so that $S(U(2) \times U(3)) \subset SU(5)$ while e.g. the group $U(1) \times SU(2) \times SU(3)$ is not a *subgroup* of $SU(5)$. Therefore the restrictions on the representation from $SU(5)$ interpreted as containing the Standard Model after some breaking leads to the restrictions of the subgroup $S(U(2) \times U(3))$. Other grand unified groups that have success typically contains $SU(5)$ as a subgroup and thus can reproduce the same representation restrictions corresponding to the Standard Model group $S(U(2) \times U(3))$. Especially we can mention the $SO(10)$ -group, which comes out of [5]. In principle a different model behind the Standard Model is the flipped $SU(5)$.

Often super string theories would go through some of these grand unification type models, but one can also construct models going to the Standard Model without going via the unifying groups.

I think one can say that the possibilities are so many that one should admit that it is not unless you somehow have already got to know the grand unified group, then the resulting group

could be so many different possible ones that there is not much predictive power in these models.

So only if one has a prediction of the unified group in a model should we consider the standard model group explained by grand unification.

In this respect the unification of spin and charge model of Norma Mankoc Borstnik (et al)[5] is better by leading to $SO(N)$ groups just from the spirit of it. To get just $SO(10)$ as is needed for getting the Standard Model the dimension $13 + 1$ should be put in, and so after all the prediction of the Standard Model does not come quite without seemingly ad hoc numbers being put in. We think of the number 14 for the space time dimension.

The type of explanation for the Standard Model group, which we are after in the present series of works is rather to seek to characterize the group by properties defined for abstract Lie-group, and then postulate some number-valued group-characteristica - which we should guess so cleverly that it will specify just the wanted Standard Model group - to by Nature arranged to be maximal.

One attempt of this type were our work, Brene et al. [6] in which we define a concept of “skewness” in a quantitative way, so that one obtains a number for each Lie group (we never got the idea completely developed to specify how many $U(1)$ invariant subgroups there should be, but I think we essentially get to there being just one $U(1)$ factor). If we to help the project a bit assumed that we should only consider the possible gauge groups with just one $U(1)$ -factor and construct a measure for the degree of symmetry as the logarithm of the number of outer automorphisms o divided by the rank of the Lie algebra r

$$“symmetry” = \frac{\ln(o)}{r}, \tag{1}$$

we get the Standard Model group to have the minimal value of this quantity “symmetry”. In this sense the Standard Model group is among the most “skew” in the sense of being the least symmetric.

The subject of the present series of papers [1] and the present article is a different approach of the same character as the just mentioned “skewness” characterization of the Standard Model group. However, our present attempt to characterize the gauge group is by a different quantity from the “symmetry” quantity; it is a new attempt. It must though be admitted that there appears one overlapping element in the two different quantities to be extremized: some of the potential outer automorphisms that are to be counted in o can be gotten away with by dividing out of the group a subgroup of the center. Actually such a division out of a subgroup of the center seems to be a rather characteristic property of the Standard Model group so that it is quite helpful if a quantity to determine the Standard Model group has a strong dependence on the appearance of an effect of such a division out of a subgroup of the center, so as to favour a gauge group with a lot of division out of the center. This division out is not relevant in the very first version of our present proposal, namely the ratio C_A/C_F of the quadratic Casimir of the adjoint representation A , denoted C_A to the quadratic Casimir of that faithful representation F having the smallest quadratic Casimir C_F . However, when we begin to help the construction of our quantity by specifying the details concerning the Abelian $U(1)$ components in the Lie group for which our quadratic Casimir ratio is not a priori welldefined, we manage to one could almost say “sneak in” the division out so that we by the details of defining what to do when we have $U(1)$ factors get the groups with a complicated division out of the center get favoured to win the game and become the gauge group chosen by nature.

In this way our present proposal for what nature has chosen to maximize and our older proposal of maximizing “skewness” is not completely different because they have an overlap by both favouring the complicate division out of the center. But apart from that they look superficially seen very different. Thus our present proposal is quite new after all.

I shall review the successive small improvements in finding partly phenomenologically our final suggestion for the quantity or the game that should specify the gauge group to be chosen in section 2 below.

1.2 History of Explaining 3+1=4 Dimensions

As said the main purpose of the present article is to use the idea from our earlier article [1] to explain why nature should have chosen just 4 (meaning 3+1) space time dimensions. This way to be explained below is new relativ to earlier attempts to explain, why we shall just have four dimensions:

One of the earlier attempts is my own [9] starting the idea of “Random Dynamics” by pointing out that in a *non-Lorentz invariant* theory - being a quantum field theory in which neither rotational nor boost invariance is present, but only translational invariance - one finds generically that assuming an appropriate Fermisurface an *effective Lorentz invariance with 3+1 dimensions* appears automatically! In this sense I claimed to derive under very general assumptions - as almost unavoidable - the appearance of both Lorentz and thereby rotational invariance and of just the right number of dimensions, $4=3+1$. The success of this dimensionality post-diction [9] were for me the introduction to a long series of works seeking to derive from almost nothing or a random theory - not obeying many of the usual principles - which I gave the name “Random Dynamics” [9, 10, 11, 12, ?], many of the known physical laws. Really we may consider the present work as an alternative attempt to derive the dimension of space time much in Random Dynamics way, in as far as we end up suggesting the philosophy that it would be most easy/likely to get just the experimental dimensionality by accident. (If successful then of course the present work would be a second derivation of 3+1 dimensions in Random Dynamics).

Also Max Tegmark has derived 3+1 dimensions from a similar Random Dynamics like philosophy of “all mathematics being realized”[2]. Max Tegmark considers the differential equations for the time development of fields so as to guarantee equations with predictivity, as well as the stability of atoms. For the anyway from his arguments needed case of just one time his figure shows that the field equation would be elliptic and thus unpredictable for $d=1$, too simple for $d=2$ and $d=3$, and unstable meaning unstable atoms say for $d=5$ or more. The latter point goes back to Ehrenfest in 1917 [3], who argued that neither atoms nor planetary systems could be stable in more than four space time dimensions.

Also known is the story that in say two spatial dimensions (corresponding to $3=2+1$ space time dimensions) an animal - as ourselves - having an intestine channel would fall apart into two pieces. Thus by an antropic principle $3=2+1$ should not be possible to have us.

According to a review of antropic questions by Gordon Kane [4]:

“One aspect of our universe we want to understand is the fact that we live in three space dimensions. There is an anthropic explanation. It was realized about a century ago[3] that planetary orbits are not stable in four or more space dimensions, so planets would not orbit a sun long enough for life to originate. For the same reason atoms are not stable in four or more space dimensions. And in two or one space dimensions there can be neither blood flow nor large numbers of neuron connections. Thus interesting life can only exist in three dimensions. Alternatively, it may be that we can derive the fact that we live in three dimensions, because the unique ground state of the relevant string theory turns out to have three large dimensions (plus perhaps some small ones we are not normally aware of). Or string theory may have many states with three space dimensions, and all of them may give universes that contain life”.

Further one has considered the renormalizability of quantum field theories not being possible for higher than 4 dimensions, except for the scalar ϕ^3 coupling theory, which is anyway not good

[14].

In theories which like string theories or Norma Mankoc Borstnik's model [5] are Kaluza-Klein-like the question of understanding the effective dimension for long distances being 3 space plus 1 time dimension would a priori mean an understanding of why precisely there is that number of extra dimensions being somehow compactified that just three space dimensions survive as essentially flat and extended. In superstring theory the consistency requires fundamentally 9 +1 dimensions of space time. If one takes it that it is needed that the compact space described by the as extra dimensions appearing dimensions must be a Callaby-Yau space, then since the latter has 6 dimensions the observed or flat dimensions must make up $(9+1) - (6+0) = 3+1$. So the combination of the superstring with the requirement of using Callaby-Yau compactification do indeed explain why we have just the experimental number of dimensions[15, 16].

In [17] you find:

“Now to make contact with our 4-dimensional world we need to compactify the 10-dimensional superstring theory on a 6-dimensional compact manifold. Needless to say, the Kaluza Klein picture described above becomes a bit more complicated. One way could simply be to put the extra 6 dimensions on 6 circles, which is just a 6-dimensional Torus. As it turns out this would preserve too much supersymmetry. It is believed that some supersymmetry exists in our 4-dimensional world at an energy scale above 1 TeV (this is the focus of much of the current and future research at the highest energy accelerators around the world!). To preserve the minimal amount of supersymmetry, $N=1$ in 4 dimensions, we need to compactify on a special kind of 6-manifold called a Calabi-Yau manifold. ”

1.3 Development of Goal Quantities

In the present article we shall ignore these antropic principle arguments and seek to get a statement that the experimental number of dimensions just maximizes some quantity, that is a relatively simple function of the group structure of say the Lorentz group, and which we then call a “goal quantity”.

Let me therefore list some of the first approximation simplified proposals which we suggest for this *goal quantity*. But this is for the dimension a two step procedure: 1) we first use the proposals in our previous article [1] to give a number - a goal quantity - for any Lie group. 2) we have to specify on which group we shall take and use the procedure of previous work; shall it be the Lorentz group, its covering group or somehow an attempt with the Poincare group ? :

- a. Just take the Lorentz group and calculate for that the Dynkin index[8] or rather the quantity which we already used as goal quantity in the previous article [1] C_A/C_F . This gets especially simple for the except for dimension $d = 2$ or smaller semi-simple Lorentz groups (simple in the mathematical sense of not having any invariant nontrivial subgroup; semi-simple: no abelian invariant subgroup); since though the global structure of the Lorentz group is not fixed until we assign it a meaning we are really here having in mind: The Lorentz group shall “have simple Lie algebra” . For simple groups we can namely ignore the minor corrections invented for the improvement in the case of an Abelian component present in the potential gauge group.
- b. We supplement in a somewhat ad hoc way the above *a.*, i.e. C_A/C_F by taking its $\frac{d+1}{d-1}$ th power. The idea behind this proposal is that we think of the Poincare group instead of as under *a.* only on the Lorentz group part, though still in a crude way. This means we think of a group, which is the Poincare group, except that we for simplicity ignore that the translation generators do not commute with the Lorentz group part.

Then we assign in accordance with the ad hoc rule used in [1] the abelian sub-Lie algebra a formal replacement 1 for the ratio of the quadratic Cassimirs C_A/C_f :I.e. we put “ $C_A/C_F|_{Abelian\ formal} = 1$ ”. Next we construct an “average” averaged *in a logarithmic way* (meaning that we average the logathms and then exponentiate again) weighted with the dimension of the Lie groups over all the dimensions of the Poincare Lie group. Since the dimension of the Lorentz group for d dimensional space-time is $\frac{d(d-1)}{2}$ while the Poincare group has dimension $\frac{d(d-1)}{2} + d = \frac{d(d+1)}{2}$ the logarithmic averageing means that we get

$$\exp\left(\frac{\frac{d(d-1)}{2} \ln(C_A/C_F)|_{Lorentz} + \ln(1) * d}{d(d+1)/2}\right) = (C_A/C_F)|_{Lorentz}^{\frac{d(d-1)/2 + d}{d+1}} = (C_A/C_F)|_{Lorentz}^{\frac{d-1}{d+1}} \quad (2)$$

That is to say we shall make a certain ad hoc partial inclusion of the abelian dimensions in the Poincare groups.

To be concrete we here propose to say crudely: let the poincare group have of course d “abelian” generators or dimensions. Let the dimension of the Lorentz group be $d_{Lor} = d(d-1)/2$; then the total dimension of the Poincare group is $d_{Poi} = d + d_{Lor} = d(d+1)/2$. If we crudely followed the idea of weighting proposed in the previous article [1] as if the d “abelian” generators were just simple cross product factors - and not as they really are: not quite usual by not commuting with the Lorentz generators - then since we formally are from this previous article suggested to use the *as if number 1 for the abelian groups*, we should use the quantity

$$(C_A/C_F)|_{Lor}^{\frac{d_{Lor}}{d_{Poi}}} = (C_A/C_F)|_{Lor}^{\frac{d-1}{d+1}} \quad (3)$$

as goal quantity.

- c. We could improve the above proposals for goal quantities *a.* or *b.* by including into the quadratic Cassimir C_A for the adjoint representation also contributions from the translation generating generators, so as to define a quadratic Cassimir for the whole Poincare group. This would mean that we for calculating our goal quantity would do as above but

$$\mathbf{Replace} : C_A \rightarrow C_A + C_V, \quad (4)$$

where C_V is the vector representation quadratic Casimir, meaning the representation under which the translation generators transform under the Lorentz group. Since in the below table we in the lines denoted “no fermions” have taken the “small representation” F to be this vector representation V , this replacement means that we replace the goal quantity ratio C_A/C_F like this:

$$\mathbf{(S)O(d), \quad “no spinors”}: \quad (5)$$

$$C_A/C_F = C_A/C_V \quad \rightarrow \quad (C_A + C_V)/C_F = C_A/C_F + 1 \quad (6)$$

$$\mathbf{Spin(d), \quad “with spinors”}: \quad (7)$$

$$C_A/C_F \quad \rightarrow \quad (C_A + C_V)/C_F \quad (8)$$

$$= C_A/C_F + (C_A/C_V)^{-1}(C_A/C_F) \quad (9)$$

$$= (1 + (C_A/C_F)|_{\text{no spinors}}^{-1})C_A/C_F. \quad (10)$$

Let me stress though that this proposal c. is not quite “fair” in as far as it is based on the Poincare group, while the representations considered are not faithfull w.r.t. to the whole Poincare group, but only w.r.t. the Lorentz group

- d. To make the proposal *c.* a bit more “fair” we should at least say: Since we in *c.* considered a representation which were only faithful w.r.t. the Lorentz subgroup of the Poincare group we should at least correct the quadratic Casimir - expected crudely to be “proportional” to the number of dimensions of the (Lie)group - by a factor $\frac{d+1}{d-1}$ being the ratio of the dimension of the Poincare (Lie)group, $d + d(d-1)/2$ to that of actually faithfully represented Lorentz group $d(d-1)/2$. That is to say we should before forming the ratio of the improved C_A meaning $C_A + C_V$ (as calculated under *c.*) to C_F replace this C_F by $\frac{d+1}{d-1} * C_F$, i.e. we perform the replacement:

$$C_F \rightarrow C_F * \frac{d(d-1)/2 + d}{d(d-2)/2} = C_F * \frac{d+1}{d-1}. \quad (11)$$

Inserted into $(C_A + C_V)/C_F$ from *c.* we obtain for the in this way made more “fair” approximate “goal quantity”

$$\text{“goal quantity”}|_{\text{no spinor}} = (C_A/C_F + 1) * \frac{d-1}{d+1} \quad (12)$$

$$\text{“goal quantity”}|_{\text{w. spinor}} = (1 + (C_A/C_F)|_{\text{no spinor}}^{-1}) * C_A/C_F * \frac{d-1}{d+1} \quad (13)$$

This proposal d. should then at least be crudely balanced with respect to how many dimensions that are represented faithfully.

The reader should consider these different proposals for a quantity to maximize (= use as goal quantity) as rather closely related versions of a quantity suggested by a perhaps a bit vague idea being improved successively by treating the from our point of view a bit more difficult to treat Abelian part (the translation part of the Poincare group) at least in an approximate way. One should have in mind, that this somewhat vague basic idea behind is: The group selected by nature is the one that counted in a “normalization determined from the Lie algebra of the group” can be said to have a faithful representation (F) the matrices of which move as little as possible when the group element being represented move around in the group.

Let me at least clarify a bit, what is meant by this statement:

We think by representations as usual on linear representations, and thus it really means representation of the group by means of a homomorphism of the group into a group of matrices. The requirement of the representation being faithful then means, that this group of matrices shall actually be an isomorphic image of the original group. Now on a system of matrices we have a natural metric, namely the metric in which the distance between two matrices \mathbf{A} and \mathbf{B} is given by the square root of the trace of the numerical square of the difference

$$dist = \sqrt{tr((\mathbf{A} - \mathbf{B})(\mathbf{A} - \mathbf{B})^+)}. \quad (14)$$

To make a comparizon of one group and some representation of it with another group and its representation w.r.t. to how fast the representation matrices move for a given motion of the group elements we need a normalization giving us a weldefined metric on the groups w.r.t. which we can ask for the rate of variation of the representations. In my short statement I suggested that this “normalization should be determined from the Lie algebra of the group”. This is to be taken to mean more precisely, that one shall consider the *adjoint* representation, which is in fact completely given by the Lie algebra, and then use the same distance concept as we just proposed for the matrix representation $\sqrt{tr((\mathbf{A} - \mathbf{B})(\mathbf{A} - \mathbf{B})^+)}$. In this way the quantity to minimize would be the ratio of the motion-distance in the representation - F say - and in the Lie algebra

representation - i.e. the adjoint representation. But that ratio is just for infinitesimal motions $\sqrt{C_F/C_A}$. So if we instead of talking about what to minimize, inverted it and claimed we should maximize we would get $\sqrt{C_A/C_F}$ to be *maximized*. Of course the square root does not matter, and we thus obtain in this way a means to look at the ratio C_A/C_F as a measure for the motion of an element in the group compared to the same element motion on the representation.

It might not really be so wild to think that a group which can be represented in a way so that the representation varies little when the group element moves around would be easier to get realized in nature than one that varies more. If one imagine that the potential groups become good symmetries by accident, then at least it would be less of an accident required the less the degrees of freedom moves around under the group to have the symmetry (approximately). It is really such a philosophy of it being easier to get some groups approximately being good symmetries than other, and those with biggest C_A/C_F should be the easiest to become good symmetries by accident, I argue for. That is indeed the speculation behind the present article as well as the previous one [1] that symmetries may appear by accident (then perhaps being strengthened to be exact by some means [10, 18]).

But let us stress that you can also look at the present work and the previous one in the following phenomenological philosophy:

We wonder, why Nature has chosen just 4 (=3+1) dimensions and why Nature - at the present experimentally accessible scale at least - has chosen just the Standard Model group $S(U(2) \times U(3))$? Then we speculate that there might be some quantity characterizing groups, which measures how well they “are suited” to be the groups for Nature. And then we begin to *seek* that quantity as being some function defined on the class of abstract groups - i.e. giving a number for each abstract (Lie?) group - of course by proposing for ourselves at least various versions or ideas for what such a *relatively simple* function defined on the abstract Lie groups could be. Then the present works - this paper and the previous one - represents the present status of the search: We found that with small variations the types of such functions representing the spirit of the *little motion of the “best” faithful representation*, i.e. essentially the largest C_A/C_F , turned out truly to bring Nature's choices to be (essentially) the winners.

In this sense we may then claim that we have found phenomenologically that at least the “direction” of a quantity like C_A/C_F or light modifications of it is a very good quantity to make up a “theory” for why we have got the groups we got!

1.4 Outline

In the following section 2 we review the main results of the C_A/C_F quantity, which we in the previous article studied for the various Lie groups in order to discuss, that the Standard Model group could be made to be favoured. In section 3 we then extract and concentrate on those groups which can be Lorentz groups. The main content of both these sections are actually the tables listing the results of the proposed to be maximized quantities for the relevant groups. In section 4 we resume and conclude, that actually we may be on the track to have found a *common* reason or explanation for, that we have 3 +1 dimensions, and for the gauge group of the Standard Model!

2 Our Previous Numbers

In the previous work by D. Bennett and myself [1] we essentially collected the ratios (related to Dynkin index [8]) C_A/C_F , where we for the representation of the group in question G selected that representation F , which would give the largest value for this ratio C_A/C_F . (In the table we

give in a few cases two proposals for F , but really it is what one would loosely call the smallest faithful representation). But we shall have in mind that this ratio is only welldefined for the simple Lie groups; and it is thus only for the simple groups we could make a clean table as the one just below. For semi-simple Lie groups it is strictly speaking needed to specify a replacement quantity, that can be the needed generalization to semisimple Lie groups. One shall naturally construct a logarithmic average weighted with the dimensions of the various simple group factors contained in the semisimple Lie group written as a product of its simple invariant subgroups. Extending the generalization even to inclusion of $U(1)$ -factors in the Lie group gets even a bit more arbitrary, but we did choose the rule of counting the $U(1)$ factors as, if they had C_A/C_F equal to unity. The problem with the $U(1)$'s, the Abelian groups, is that the adjoint quadratic Cassimir C_A is just zero and does not provide a good normalization. But although we thus have to declare formally C_A/C_F to be unity at first for $U(1)$ we take the opportunity to - and we think it is very natural - to include a correction depending not only on the Lie-*algebra* but also on the *group* structure in a way roughly describing that a $U(1)$ representation with a small "charge" is "smaller" than one with a larger charge in a very similar way to the way in which a small quadratic Cassimir signals a "small" representation.

Our Ratio of Adjoint to "Simplest" (or smallest) Quadratic Casimirs C_A/C_F

$$\frac{C_A}{C_F}|_{A_n} = \frac{2(n+1)^2}{n(n+2)} = \frac{2(n+1)^2}{(n+1)^2-1} = \frac{2}{1-\frac{1}{(n+1)^2}} \quad (15)$$

$$\frac{C_A}{C_F \text{ vector}}|_{B_n} = \frac{2n-1}{n} = 2 - \frac{1}{n} \quad (16)$$

$$\frac{C_A}{C_F \text{ spinor}}|_{B_n} = \frac{2n-1}{\frac{2n^2+n}{8}} = \frac{16n-8}{n(2n+1)} \quad (17)$$

$$\frac{C_A}{C_F}|_{C_n} = \frac{n+1}{n/2+1/4} = \frac{4(n+1)}{2n+1} \quad (18)$$

$$\frac{C_A}{C_F \text{ vector}}|_{D_n} = \frac{2(n-1)}{n-1/2} = \frac{4(n-1)}{2n-1} \quad (19)$$

$$\frac{C_A}{C_F \text{ spinor}}|_{D_n} = \frac{2(n-1)}{\frac{2n^2-n}{8}} = \frac{16(n-1)}{n(2n-1)} \quad (20)$$

$$\frac{C_A}{C_F}|_{G_2} = \frac{4}{2} = 2 \quad (21)$$

$$\frac{C_A}{C_F}|_{F_4} = \frac{9}{6} = \frac{3}{2} \quad (22)$$

$$\frac{C_A}{C_F}|_{E_6} = \frac{12}{\frac{26}{3}} = \frac{18}{13} \quad (23)$$

$$\frac{C_A}{C_F}|_{E_7} = \frac{18}{\frac{57}{4}} = \frac{72}{57} = \frac{24}{19} \quad (24)$$

$$\frac{C_A}{C_F}|_{E_8} = \frac{30}{30} = 1 \quad (25)$$

For calculation of this table seek help in[20, 19]

In the just above table we have of course used the conventional notation for the classification of Lie algebras, wherein the index n on the capital letter denotes the rank of the Lie algebra, and:

- A_N is $SU(n+1)$,

- B_n are the symplectic Lie algebras.
- C_n is the odd orthogonal Lie algebra for $SO(2n+1)$ or for its covering group $\text{Spin}(2n+1)$,
- D_n is the even dimension orthogonal Lie algebra for $SO(2n)$ or its covering group $\text{Spin}(2n)$,
- while F_4 , G_2 , and E_n for $n = 6, 7, 8$ are the exceptional Lie algebras.

The words *spinor* or *vector* following in the index the letter F which itself denotes the “small” representation - i.e. most promising for giving a small quadratic Cassimir C_F - means that we have used F respectively the spinor and vector representation.

It may be reassuring to check that our goal quantity for the simple groups C_A/C_F becomes the same for the cases of isomorphic Lie algebras such as $B_2 \cong C_2$ and $C_3 \cong A_3$.

2.1 Development of the Gauge Group Determination Proposal

It may be best to describe the proposal for the quantity to be maximized for the gauge group by describing how a phenomenological discussion adjusting small problems can be guided towards of final rule. Let me review the work [1] as a successive discussion towards finding a goal quantity that would make the Standard Model group win the game of making it maximal:

The following successive proposals are then the following:

- I The ground idea for the goal quantity is the ratio C_A/C_F in which the symbols C_A and C_F are the quadratic Casimirs for the group in question for respectively the adjoint representation A and another representation F which then in the search for a maximal ratio C_A/C_F will lead to choosing F with minimal C_F . To avoid that F be the trivial representation we shall what is very reasonable we would say require F to be faithful.

Now the reader shall have in mind that

- I.1 The ratio C_A/C_F does not suffer from the normalization problem of the generators representing the Lie algebra, because we take the ratio so that scaling the convention for the Lie algebra basis if changed will change the numerator C_A and the denominator C_F by the same factor.
- I.2 But at first even this *ratio* C_A/C_F is only welldefined for a simple Lie group. In the case of even a still semisimple Lie group there is namely an ambiguity in the in the normalization of the basis vectors for the Lie algebra.
- I.3 If we do not specify the normalizations of the basis vectors and thereby their representations, then we get a notation dependent quantity.
- I.4 If we have $U(1)$ as factors in the \times -product, then we have the obvious trouble with our first proposal C_A/C_F that the adjoint representation of $U(1)$ is trivial, or rather the Abelian Lie group $U(1)$ has no (meaningful) adjoint representation and thus C_A becomes meaningless for $U(1)$.
- II Next let us improve the first proposal C_A/C_F for the goal quantity, by generalizing it in a good way to the *semisimple Lie groups*, meaning in practice for us a Lie group of the form of a cross product of some $U(1)$ groups and some simple Lie groups finally modified by dividing out some discrete subgroup of the center. That is to say we have in mind groups of the form

$$G = U(1) \times U(1) \times \cdots \times U(1) \times SU(2) \times \cdots \dots \times G_{max}/D \quad (26)$$

where the \times -product runs over a numbe of occurences of all the possible simple Lie groups as classified say by their Dynkin diagrams up to the last one for the group G in question here denoted G_{max} . Finally then the *group* rather than Lie algebra structure is achieved by dividing some discrete subgroup D out of the center of the group achieved by the cross product without modification. The reader should have in mind that the typical covering groups as e.g. $SU(N) = A_{N-1}$ has often a nontrivial center with a finite number of elements. E.g. $SU(N)$ has a center isomorphic to the group Z_N of the integer numbers counted modulo N . The whole center in the cross product thus becomes the cross product of the typically discrete centers for the simple Lie groups crossed further with the $U(1)$'s which each of them are all center (since they are Abelian). For example the center of the cross product $U(1) \times SU(2) \times SU(3)$ that shall be used to produce the Standard Model group (and which has the Standard Model Lie algebra) is $U(1) \times Z_2 \times Z_3$. By dividing out of this a cleverly chosen with Z_6 isomorphic (discrete) subgroup D generated by the element $(2\pi, 1(mod2), 1(mod3))$ (where "1(mod3)" e.g. the element in Z_3 being the class of integers with rest 1 modulo 3) we obtain a *group* which does not have as representations of the group all the representations of the cross product $U(1) \times SU(2) \times SU(3)$ or better its covering group, but nevertheless those representations occuring for particles in the Standard Model.

The problem which we first have to solve in extending in a meaningfull way the proposed quantity is to even ignoring at first the $U(1)$ groups and restricting ourselves to semisimple groups find some way of defining a quantity like the for the simple Lie group or algebra welldefined quantity C_A/C_F .

Since for a simple group the ratio C_A/C_F is welldefined the obvious idea to make analogous expression for a semisimple, which is just a \times -product of several simple Lie groups $S_1 \times S_2 \times \dots \times S_n$ by taking some sort of average over the sepearate simple groups S_i of the quantities $C_A/C_F|_{S_i}$ for the various simple groups. The proposal which we thought were reasonable was to average logarithmically and weighting with the dimension of the Lie groups. That means we proposed to use as the average that should replace the C_A/C_F for the simple group in the semisimple case

$$\begin{aligned} & \text{"}C_A/C_F\text{replacement for } S_1 \times S_2 \times \dots \times S_n'' \quad (27) \\ & = (C_A/C_F|_{S_1})^{\frac{dim(S_1)}{\sum_i dim(S_i)}} * (C_A/C_F|_{S_2})^{\frac{dim(S_2)}{\sum_i dim(S_i)}} * \dots * (C_A/C_F|_{S_n})^{\frac{dim(S_n)}{\sum_i dim(S_i)}} \quad . \quad (28) \end{aligned}$$

It is of course a priori an ad hoc choice to weight just with the dimensions $dim(S_i)$ of the simple groups S_i , but is an extremely natural choice, but of course the type of choice we could revise if we should look for some little adjustment of our proposal to make it be more successful.

You may consider the quadratic Casimir as representing a metric tensor describing distances in the representation space for coordinates originating from the Lie algebra or Lie group. If one would think of the volume (of dimension $dim(G)$ of course) for a faithful representation F say relative to the adjoint representation volume, it would be $(C_F/C_A)^{dim(G)/2}$. So you could see our quantity "C_A/C_Freplacement for $S_1 \times S_2 \times \dots \times S_n$ " as being the $dim(S_1 \times S_2 \times \dots \times S_n)/2$ 'th root of the volume ratio of the adjoining representation A , i.e. $Vol(A)$ relative to that of the faithful representation F , i.e. $Vol(F)$,

$$\text{"}C_A/C_F\text{replacement for } S_1 \times S_2 \times \dots \times S_n'' = \left(\frac{Vol(A)}{Vol(F)} \right)^{\frac{1}{dim(G)}}. \quad (29)$$

This simple and nice interpretation supports estetically the use of the dimensionality of the various simple Lie groups being used to weight the logarithmic average. We can namely instead of talking at first about the quadratic Casimirs say that we talk about the volume ratio of the adjoint representation and the representation F from the start. The ratio of the volumes of two representations in the natural metric defined above in (14) is a very simple and beautyfull quantity. We then take the root of (half) the dimensionality of the group to make it depend on the structure of the various simple subgroups rather than on the total number of them or hteir dimension in a too strong way. By using this root choice (29) we obtain the good feature that you cannot obtain the large quantity just by taking a group with a high dimension by taking for instance a cross product of a lot of groups. For single simple groups we also achieve that our root quantity becomes just the C_A/C_F from which we started. And you can even see from the table above that in the limit of the rank going to infinity the various series of infinitely many simple Lie groups have our quanti y go nicely to 2. In this way we get a very *ballanced* quantity, not favouring at first neither large nor small dimensions dramatically.

Hereby we think we have proposed a very nice and beautyfull quantity for the semisimple groups.

- III Next we have the problem with groups having $U(1)$ -factors:

For the $U(1)$ our C_A/C_F hardly makes any sense, and so we have to invent a replacement essentially arbitrarily. In order to do that let us have in mind that whenever our quantity C_A/C_F makes sense it is for trivial reasons allways bigger than unity. We could namely always as a special possibilty for the faithful representation F use the adjoint representation itself, in which case of course the ratio $C_A/C_F = C_A/C_A = 1$. So since we shall choose the representation F so as to maximize the ratio C_A/C_F , it must allways be larger than or equal to this possibility value 1.

When we so to speak shall invent a value for the replacement of the C_A/C_F for the Abelian group $U(1)$, we must at least not to violate the trivial lower bound at least choose the value larger than or equal to 1. In the light of that at first we have apart from there being a “charge” value, which is hard to specify, to determine a representation of $U(1)$, suggest that there is in some way only one representation of it and thus we have from the trivial consideration just the lower bound value - which occurs when you can not find an F different from the A - to be the replacement value to formally be used in specifying our game. In other words we propose to choose:

$$“C_A/C_F\text{replacement for }U(1)” = 1. \quad (30)$$

It is of course then the obvious generalization to the groups being cross products of $U(1)$'s with a semisimple group that we shall average this 1 logarithmically with the C_A/C_F 's for the simple groups weighting as above with the dimension of the Lie groups.

This means that we have now come to the proposal:

$$“C_A/C_F\text{replacement for }U(1) \times \dots \times U(1) \times S_1 \times \dots \times S_n” \quad (31)$$

$$= 1^{\frac{1}{\#U(1)'s + \sum_i \dim(S_i)}} * \dots * 1^{\frac{1}{\#U(1)'s + \sum_i \dim(S_i)}} \quad (32)$$

$$*(C_A/C_F|_{S_1})^{\frac{\dim(S_1)}{\#U(1)'s + \sum_i \dim(S_i)}} * \dots * (C_A/C_F|_{S_n})^{\frac{\dim(S_n)}{\#U(1)'s + \sum_i \dim(S_i)}} \quad (33)$$

$$= (C_A/C_F|_{S_1})^{\frac{\dim(S_1)}{\#U(1)'s + \sum_i \dim(S_i)}} * \dots * (C_A/C_F|_{S_n})^{\frac{\dim(S_n)}{\#U(1)'s + \sum_i \dim(S_i)}}. \quad (34)$$

Here $\#U(1)'s$ means the number of $U(1)$ -factors in the \times -product forming the group G under study/evaluation w.r.t. the goal quantity. Of course the full dimension of the Lie group G is just

$$\dim(G) = \#U(1)'s + \sum_i \dim(S_i). \quad (35)$$

At this stage it will of course *not* w.r.t. maximizing the goal quantity “ C_A/C_F replacement for $U(1) \times \dots \times U(1) \times S_1 \times \dots \times S_n''$ pay to have any $U(1)$'s at all. So we have at this stage in de-velopping the goal quantity no chance to make group that like the Standard Model has an invariant Abelian subgroup $U(1)$ have any chance of winning the game of maximization.

- IV Improvement for Abelian and the *group* structure:

In order to make it at all pay for the maximization to have at least some factor $U(1)$ like the Standard Model happens to have we must open for the possibility of letting a $U(1)$ invariant subgroup contribute more than just absolute minimum 1 as replacement for its meaningless C_A/C_F . But the irreducible representations of the $U(1)$ are all just one-dimensional representations with the single element in the unitary matrix being just a phase factor $\exp(ie\delta)$, where δ is the phase describing the element in $U(1)$ and e is a “charge” for the representation in question. It is wellknown that the various representations of $U(1)$ are characterized by such “charges” e . The quadratic Casimirs are given as the square of the “charges” $C_R \rightarrow e_R^2$, where e_R is the “charge” for the representation R .

Now let us have in mind that for our purpose studying gauge groups we mainly have in mind the charges for various particles, and that when we give a physical meaning to the gauge *group* rather than just to the gauge Lie algebra we do that on the basis of phenomenologically found restrictions on the representations/particles occurring in the model. We can then ask the questions: What is the lowest non-zero charge e_A say on a particle in accordance with the restrictions from the *group*-structure, when this particle has no non-abelian transformations(i.e. when it transforms trivially/i.e. not at all under the other factors in the group? And we can also ask what is the absolutely (numerically) smallest “charge” e_F say on any particle allowed under the group rule(whatever its couplings to the non-abelian Lie groups might be)? The indices suggested here were chosen to suggest to use the here defined minimal charges e_A and e_F should be used to form a “replacement” for the C_A/C_F for a $U(1)$ being instead of the at first proposed 1 now improved to e_A^2/e_F^2 . We can only obtain this ratio e_A^2/e_F^2 to be different from unity by having a gauge group obtained by dividing out a discrete subgroup of the center of the starting pure cross product. Actually this choice is very reasonable in as far as the e_F charge is the smallest possible non-trivial charge quite analogous to our in the non-abelian case choice of the representation F being the smallest faithful one. So the only a little bit ad hoc choice is to replace the adjoint representation for the non-abelian case by the smallest representation of the $U(1)$ that does *not* mix up with non-abelian groups. We think this is pretty much the simplest replacement.

Choosing this procedure we get to the final proposal:

$$\text{“}C_A/C_F\text{ replacement for }U(1)^{(1)} \times \dots \times U(1)^{(m)} \times S_1 \times \dots \times S_n'' = \quad (36)$$

$$= \left(\frac{e_A^{(1)2}}{e_F^{(1)2}} * \dots * \frac{e_A^{(m)2}}{e_F^{(m)2}} * \frac{C_A}{C_F} \Big|_{S_1}^{dim(S_1)} * \dots * \frac{C_A}{C_F} \Big|_{S_n}^{dim(S_n)} \right)^{\frac{1}{\#U(1)'s + \sum_i \dim(S_i)}}. \quad (37)$$

Here of course $m = \#U(1)'s$ and the index in the round brackets on the charges enumerates the various $U(1)$ factors in the cross product. It should be had in mind that the non-trivial (i.e. not just 1) ratios $\frac{e_A^{(i)2}}{e_F^{(i)2}}$ only come in play when a discrete subgroup of the center of the cross product have been divided out. Remember that I above told that this division out of a discrete subgroup of the center were in the “skewness” estimation [6] an important ingredient in reducing the symmetry and thus got favoured by asking for “skewness”. So letting this division out be favoured in the game gives a favouring of the Standard Model *group* which has relatively much such “division out”. This is how we got in this last step introduced a favouring of the “division out” although we started out by the C_A/C_F ratio which were purely dependent on the Lie algebra alone.

2.2 Standard Model Group Wins

Using the table (25) inserting it into (37) we may now contemplate which group should win the “game” of obtaining the largest value for the goal quantity (37).

First we see from the table (25) that the groups favoured even in the system that were made ballanced in such a way that the ratio C_A/C_F goes to a constant -actually 2 - for very large ranks $r \rightarrow \infty$ are the small rank ones: $SU(2) = A_1$ is the winner among the simple group with its $C_A/C_F = \frac{2}{1 - \frac{1}{(n+1)^2}}$ for A_n which for $n = 1$ gives 8/3.

The next among the simplest Lie groups is $SO(5) = B_2 = C_2$ with $C_A/C_F = 12/5$, which is obtained by using the spinor representation for B_2 giving $\frac{16n-8}{n(2n+1)}|_{n=2} = \frac{12}{5}$, while the vector representation gives the less competitive $C_A/C_F|_{B_2 \text{ vector}} = 3/5$, and checking that the isomorphic C_2 also gives as it should 12/5. First at the third place we find the Lie group $SU(3) = A_2$ with its $C_A/C_F|_{A_2} = 9/4$ the group we would hope to win over the $SO(5)$, because it is $SU(3)$ and not $SO(5)$ which occurs in the Standard Model.

But now in the competition between the $SO(5)$ and the $SU(3)$ comes in some help for $SU(3)$ in our final proposal:

- Dimension of $SU(3)$ is lower than that of $SO(5)$ Comparing the semisimple groups formed by crossing $SU(2)$ with respectively $SO(5)$ and $SU(3)$ we obtain the C_A/C_F ratios when weighted according to (28) to be

$$\frac{C_A}{C_F} \text{“replacement”}|_{SU(3) \times SO(5)} = \left(\frac{8}{3}\right)^{3/13} * \left(\frac{12}{5}\right)^{10/13} = 2.459068704 \quad (38)$$

$$\frac{C_A}{C_F} \text{“replacement”}|_{SU(3) \times SU(3)} = \left(\frac{8}{3}\right)^{3/11} * \left(\frac{9}{4}\right)^{8/11} = 2.356709384 \quad (39)$$

$$(40)$$

The $(\frac{12}{5}/\frac{9}{4} - 1) * 100\% = 6.6666667\%$ higher value for $SO(5)$ over $SU(3)$ is by the logarithmic dimensional weighting reduced to $(\frac{2.459068704}{2.356709384} - 1) * 100\% = 4.343315332\%$.

- Involving a $U(1)$ and the division out of a central subgroup

According to the details of the definition of our “goal quantity” when involving $U(1)$ cross product factors we have the possibility of obtaining the dimensional root of factors e_A^2/e_F^2 from (37). As by our definitonal choice above the difference between the charge e_A and e_F is that e_A should be for representation *with only the $U(1)$ charge but trivial under the non-Abelian groups* while e_F can be chosen for any faithfull representation, the ratio e_A/e_F can only be bigger than unity by involving a rule for allowed representations of the *group*

(rather than just the Lie algebra). That is to say we need to involve the center so to speak of one of the covering groups of the non-Abelian Lie groups. We can then e.g. for $SU(3)$ which has a center isomorphic to the group of integers modulo 3, i.e. Z_3 , obtain a division out of a Z_3 and get a ratio of 3 for e_A over e_f if we wish. The reader should check that using our formula (37) for the goal quantity and imagining various *groups* obtained by various division-outs of the center we can with only *one* $U(1)$ cross product factor only divide out say Z_2 once if we want to get the effect of this division out getting felt for the ratios e_A/e_F obtainable, whereas we can even with only one $U(1)$ manage to get both Z_2 and Z_3 give rise to effective factors of the type $(e_A/e_F)^2$. With just one $U(1)$ factor we can thus gain a factor $(e_A^2/e_F^2)^{1/dim(G)}$ which gives $(1/3^2)^{1/dim(G)}$ for $SU(3)$ and $(1/2^2)^{1/dim(G)}$ for $SU(2)$ in the goal quantity (where $dim(G)$ is the dimension of the full group). However, once we have already gotten such a gain from $SU(2)$ we cannot gain one more from $SO(5)$ unless we incorporate yet another $U(1)$. In this way $SU(3)$ gets favoured not only by having a Z_3 isomorphic center compared to the only Z_2 isomorphic center of the $SO(5)$ covering group, meaning a $3^2 = 9$ factor compared to the $2^2 = 4$ only for $SO(5)$, but the $SO(5)$ cannot get its Z_2 in play without one more $U(1)$. So in reality now $SU(3)$ gets in front by a factor 9 (before one takes the $dim(G)$'s root.).

So now let us compare the two groups obtained from the semisimple ones in (39) by cross multiplying them with a $U(1)$ and successively dividing appropriately a discrete group out of the center:

We calculate the following goal quantities

$$\text{“}C_A/C_F\text{replacement for }SU(2) \times SU(3) \times U(1)/Z_6'' = \left(6^2 * \left(\frac{8}{3}\right)^3 * \left(\frac{9}{4}\right)^8\right)^{\frac{1}{12}} \quad (41)$$

$$= 2.957824511 \quad (42)$$

$$\text{“}C_A/C_F\text{replacement for }SU(2) \times Spin(5) \times U(1)/Z_2'' = \left(2^2 * \left(\frac{8}{3}\right)^3 * \left(\frac{12}{5}\right)^{10}\right)^{\frac{1}{14}} \quad (43)$$

$$= 2.54602555 \quad (44)$$

Here $Spin(5)$ just stands for the covering group of $SO(5)$, the numbers for the three involved simple Lie groups are the C_A/C_F ratios respectively $8/3$, $9/4$, and $12/5$ for $SU(2)$, $SU(3)$ and $Spin(5)$ (or just think $SO(5)$). The numbers 36 and 4 come from the “charge” ratio and are essentially the squares of the number of elements in the divided out subgroup in the cases here. The final root taking is of course because of the averaging with the dimension of the full group- respectively 12 and 14 - to finally be divided out of the logarithm.

It might be nice to have in mind what the significance of e.g. the factor 3 in the “charge ratio” e_A/e_F due to the $SU(3)$ contributes, namely a factor 9 before the 12th root is taken. Indeed $9^{\frac{1}{12}} = 1.200936955$, which means that obtaining the charge ratio due to the $SU(3)$ rather than there being no factor with $SO(5)$ ($Spin(5)$) not even a new Z_2 to divide out when we already have done so using $SU(2)$, because we only allow the charge for the $U(1)$ to be reduced by the factor 2 once.

The only 4.343315332% advantage of the semisimple $SO(5)$ over the $SU(3)$ when combined to $SU(2) \times Spin(5)$ and $SU(2) \times SU(3)$ respectively is thus rather easily overshadowed by the effect of the e_A/e_F from the $SU(3)$, which is of the order of 20 % in the goal quantity.

After the inclusion of the Abelian charge type of ratio we found that the final advantage of the Standard Model *group* $S(U(2) \times U(3)) = U(1) \times SU(2) \times SU(3)/Z_6$ compared to the

group with the $SU(3)$ replaced by the competing $Spin(5) SO(5)$ namely $U(1) \times SU(2) \times Spin(5)/Z_2$ is $(\frac{2.957824511}{2.54602555} - 1) * 100\% = 16.174188\%$

2.3 Some Property of Our Goal Quantiy

We must have in mind the property of our “goal quantity” due to its logarithmic averaging has the property that taking a repeated cross product of whatever group with itself necessarily leads to groups with the same goal quantiy as the one mutiplied up. Thus if say the Standard Model group wins, then at the same time any number of crossings of the Standard Model with itself will stand even and share the first plac with the Standard Model alone.

We above essentially had the discussion that lead to the Standard Model winning except that we did not sufficiently carefully compare groups with different numbers of simple group factors. For instance the obvious and very serious competitor to the Standard Model is simply $U(2) = U(1) \times SU(2)/Z_2$, which obtains the goal quantiy

$$“C_A/C_F\text{replacement for }U(2)” = \left(2^2 * \left(\frac{8}{3}\right)^3\right)^{\frac{1}{4}} = 2.951151786. \quad (45)$$

This is truly exceedingly close run to the Standard Model, but the Standard Model indeed wins over even $U(2)$ on the fourth ciffer. Indeed the advange of the Standard Model group over the so closely competing $U(2)$ (which would physically be that there were no strong interactions, but only the Weinberg Salam Glashow model say) is by $(\frac{2.957824511}{2.951151786} - 1) * 100\% = .2261058\%$ Since the contribution from the Abelian invariant subgroup $U(1)$ namely the “charge ratio” is so important we might look for the winning group by first noticing that for the same center of a Lie group with a simple Lie algebra different such simple Lie algebras will play the same role w.r.t. the division out of the center and the “charges for the Abelian group(s). So we can concentrate on choosing from each class of simple Lie algebras giving a given center group for the covering group. This would mean that we among simple Lie algebras having center Z_2 in their covering group can concentration on $SU(2)$; for those with center Z_3 in their covering group we can concentrate on $SU(3)$ etc.

One may rather easily see that involving the more complicated center groups in the simple Lie algebras hardly shall pay.

If you seek as would be best a Z_k center with k being prime w.r.t. the other k values say 2 and 3 we get up to $k=5$ and $SU(5)$ already has the high dimension 24 and would largely weight out the effect of even a factor 5^2 ; in fact $(5^2)^{1/24} = 1.143529836$. We may look at the series:

$$“C_A/C_F\text{replacement for }U(1)” = (1)^1 = 1 \quad (46)$$

$$“C_A/C_F\text{replacement for }U(1) \times SU(2))/Z_2'' = (2^2 * (8/3)^3)^{\frac{1}{4}} \quad (47)$$

$$= 2.951151786 \quad (48)$$

$$“C_A/C_F\text{replacement for }U(1) \times SU(2) \times SU(3))/Z_6'' = (6^2 * (8/3)^3 * (9/4)^8)^{\frac{1}{24}} \quad (49)$$

$$= 2.957824511 \quad (50)$$

$$“C_A/C_F\text{replacement for }U(1) \times SU(2) \times SU(3) \times SU(5))/Z_{30}'' = \quad (51)$$

$$(6^2 * (8/3)^3 * (9/4)^8 * (25/12)^{24})^{\frac{1}{36}} = 2.341513375 \quad (52)$$

$$(53)$$

We see that in this series of the most promissing candidates with given centers of the covering groups for the simple Lie algebras the Standard Model lies at the (flat) maximum.

The reader can himself check in detail and get help by studying our earlier work[1], and see that indeed the Standard Model wins our game with its value 2.957824511, sharply followed by the group $U(2)$, which achieves 2.951151786 (for its silver medal).

We think it is remarkable that such a relatively simple proposal for a goal quantity as our slightly ad hoc extended essentially Dynkin index C_A/C_F , the precise definition of which is largely determined from requirements of not depending too much on the notation choice, leads to just the gauge group that Nature has chosen! One should think that there is truly something in it. By this that it is largely fixed by independence of notation, we mean that it had to be a *ratio* of quadratic Casimirs, if it shall be given by quadratic Casimirs at all; otherwise it would depend on the normalization of the quadratic Casimir, which would make it much more complicated to define. Our just called “ad hoc” extension to the inclusion of $U(1)$ cross product factors is really very analogous to the C_A/C_F , so indeed it is not such a series arbitrary choice.

We even have a speculative physical mechanism behind which though might be thought to be more replacable by some other version.

3 Competition Among Lorentz Groups on C_A/C_F and the Like

Caption: In the below tables we evaluate for different dimensions d of the Minkowski space-time - for simplicity here replaced by the Euklideanized d dimensional space-time, but that makes no difference for our calculation here - the first two goal-quantities proposed, $a.$ and $b.$ in subsection 1.3 written in respectively 5th and 7th columns. Because of the ambiguity of the global structure of the Lorentz-**group** the group in d dimension may be either $O(d)$ (essentially $SO(d)$ if we do not include parity) or Spinor(d) if we use the covering group. We have therefore for each value of the dimension d two items corresponding to these two global extensions of the Lie algebra of the Lorentz group, and they are denoted by “no spinors” and “with spinors” respectively.

Dimension	Lorentz group	Spinor or not	Rank	Ratio C_A/C_F max a)	$\frac{d_{Lor}}{d_{Poi}}$	$(C_A/C_F)^{\frac{d_{Lor}}{d_{Poi}}}$ max b)
d=1	discrete		0	-	0	indefinite 0 ⁰
d=2 ¹	(S)O(2) = U(1) U(1)	no spinor with spinor	1 1	- (formally 1) -(formally 2)	1/3 1/3	- (formally 1) -(formally 2 ^{1/3} = 1.26)
d=3	(S)O(3) Spin(3)=SU(2)	no spinor with spinor	1 1	1 8/3= 2.6667	1/2 1/2	1 $\sqrt{8/3}$ =1.632993162
d=4	(S)O(4) Spin(4) = SU(2) × SU(2)	no spinor with spin	2 2	4/3=1.3333 8/3=2.6667	3/5 3/5	$(4/3)^{3/5}$ =1.188401639 $(8/3)^{3/5}$ =1.801280051
d=5	(S)O(5) Spin(5)	no spinor with spinor	2 2	3/2=1.5 12/5=2.4	2/3 2/3	$(3/2)^{2/3}$ =1.310370697 $(12/5)^{2/3}$ =1.792561899
d=6	(S)O(6) Spin(6) = SU(4)	no spinor with spinor	3 3	8/5=1.6 32/15=2.1333	5/7 5/7	$(8/5)^{5/7}$ = 1.398942897 $(32/15)^{5/7}$ =1.718074304
d=7	(S)O(7) Spin(7)	no spinor with spinor	3 3	13/7=1.8571 40/21=1.9048	3/4 3/4	$(13/7)^{3/4}$ =1.590867407 $(40/21)^{3/4}$ =1.621363987
d=8	(S)O(8) Spin(8)	no spinor with spinor	4 4	12/7=1.7143 12/7=1.7143	7/9 7/9	$(12/7)^{7/9}$ =1.520774129 $(12/7)^{7/9}$ =1.520774129
d=9	(S)O(9) Spin(9)	no spinor with spinor	4 4	7/4=1.75 14/9=1.5556	4/5 4/5	$(7/4)^{4/5}$ =1.564697681 $(14/9)^{4/5}$ = 1.423994858
d=10	(S)O(10) Spin(10)	no spinor with spinor	5 5	16/9=1.7778 64/45=1.4222	9/11 9/11	$(16/9)^{9/11}$ =1.601198613 $(64/45)^{9/11}$ =1.33399805
d=11	(S)O(11) Spin(11)	no spinor with spinor	5 5	9/5=1.8 72/55= 1.3091	5/6 5/6	$(9/5)^{5/6}$ =1.632026054 $(72/55)^{5/6}$ =1.251626758
d=12	(S)O(12) Spin(12)	no spinor with spinor	6 6	44/23= 1.9130 40/33= 1.2121	11/13 11/13	$(44/23)^{11/13}$ =1.731340775 $(40/33)^{11/13}$ =1.176773318
d=13	(S)O(13) Spin(13)	no spinor with spinor	6 6	25/13 =1.9231 44/39 =1.1282	6/7 6/7	$(25/13)^{6/7}$ 1.75156277 $(44/39)^{6/7}$ =1.108929813
d=14	(S)O(14) Spin(14)	no spinor 18 with spinor	7 7	24/13 =1.8461 104/105 =0.9905	13/15 13/15	$(24/13)^{13/15}$ = 1.701239682 $(104/105)^{13/15}$

Dimension	Lorentz group	Spinor or not	Rank	Ratio C_A/C_F max a)	$\frac{d_{Lor}}{d_{Poi}}$	$(C_A/C_F)^{\frac{d_{Lor}}{d_{Poi}}}$ max b)
d odd	(S)O(d)	no spinor	$n=(d-1)/2$	$2-1/n$ $=2-2/(d-1)$	$\frac{d-1}{d+1}$	$(2-\frac{1}{d-1})^{\frac{d-1}{d+1}}$
	Spin(d)	with spinor	$n=(d-1)/2$	$\frac{8(2n-1)}{n(2n+1)}$ $=\frac{16(d-2)}{d(d-1)}$	$\frac{d-1}{d+1}$	$(\frac{16(d-2)}{d(d-1)})^{\frac{d-1}{d+1}}$
d even	(S)O(d)	no spin	$n=d/2$	$\frac{4(n-1)}{2n-1}$ $=\frac{2(d-2)}{d-1}$	$\frac{d-1}{d+1}$	$(\frac{2(d-1)}{d-1})^{\frac{d-1}{d+1}}$
	Spin(d)	with spinor	$n=d/2$	$\frac{16(d-2)}{d(d-1)}$	$\frac{d-1}{d+1}$	$(\frac{16(d-2)}{d(d-1)})^{\frac{d-1}{d+1}}$
$d \rightarrow \infty$		no spinor	$c \rightarrow \infty$	$\rightarrow 2$	$\rightarrow 1$	$\rightarrow 2$
		with spinor	$\rightarrow \infty$	$\rightarrow 0$	$\rightarrow 1$	$\rightarrow 0$

¹ The case $d=2$ is special because the Lorentz group is Abelian $U(1)$ for $d=2$, and we must apply the formal extension definition $C_A/C_F = 1$ for $U(1)$ from our previous work [1], and even include an extra factor connected with dividing out a subgroup of the center, or better say that the formal quadratic Cassimir shall behave like the charge squared for the $U(1)$.

3.1 Discussion of Table

Motivated either by the fact that we have spinor transforming particles in nature a- namely the Fermions - or because the goal-numbers for the spinor groups are anyway the biggest (most competitive) we should think of the Lorentz *group* as the spinor group and therefore in the above table read the Spin(d) entrances rather than the (S)O(d)-entrances:

Concentrating on the *Spin(d)*-entrances we then find that with the proposal *a.* of subsection 1.3 the dimensions $d = 3$ and $d= 4$ stand even with the same goal-number $8/3 = 2.6667$. But note that at least the experimental dimension 4 already is in the sample of the “winners” with the simple choice of *a.* meaning, that we only consider the genuine Lorentz group - but totally ignore the abelian part of the Poincare group.

Next when we go to the slightly more complicated version of a goal-quantity namely *b.* we get the separation between also $d=3$ and $d=4$, and it is the $d=4$ dimension, that “wins”, because we get for $d=3$ only 1.6330, while we for $d=4$ obtain 1.8013. Thus in this approximate treatment of the Abelian part also being included the (after all rather) “little” difference between the two schemes *a.* and *b.* leads to giving the $d=4$ case - the experimental case the little push forward making the experimental dimension $d=4$ be the only winner!

Caption: We have put the goal-numbers for the third proposal *c* in which I (a bit more in detail) seek to make an analogon to the number used in the reference [1] in which we studied the gauge group of the Standard Model. The purpose of *c.* is to approximate using the *Poincare group* a bit more detailed, but still not by making a true representation of the Poincare group. I.e. it is still not truly the Poincare group we represent faithfully, but only the Lorentz group, or here in the table only the covering group *Spin(d)* of the Lorentz group. However, I include in the column marked “*c.*, max c)” in the quadratic Cassimir C_A of the Lorentz group an extra term coming from the structure constants describing the non-commutativity of the Lorentz group generators with the translation generators C_V so as to replace C_A in the starting expression of ours C_A/C_F by $C_A + C_V$. In the column marked “*d.*, max d)” we correct the ratio to be more “fair” by counting at least that because of truly faithfully represented part of the Poincare group

in the representations, I use, has only dimension $d(d-1)/2$ (it is namely only the Lorentz group) while the full Poincare group - which were already in c . but also in d . used in the improved C_A being $C_A + C_V$ - is $d(d-1)/2 + d = d(d+1)/2$. The correction is crudely made by the dimension ratio $\dim(\text{Lorentz})/\dim(\text{Poincare}) = (d-1)/(d+1)$ given in the next to last column.

Dimension	Lorentz group (covering group)	Ratio C_A/C_F for spinor	Ratio C_A/C_V as “no spinor”	c -quantity max c)	$\frac{d-1}{d+1}$	d -quantity max d)
2^2	U(1)	-(formally 2)	-(formally 1)	4	1/3	4/3=1.33
3	spin(3)	$\frac{8}{3} = 2.6667$	1	$\frac{16}{3} = 5.3333$	$\frac{2}{4} = .5$	$\frac{8}{3} = 2.6667$
4	$Spin(4)$ $= SU(2) \times SU(2)$	$\frac{8}{3} = 2.6667$	$\frac{4}{3}$	$\frac{14}{3} = 4.6667$	$\frac{3}{5} = .6$	$\frac{14}{5} = 2.8$
5	Spin(5)	$\frac{12}{5} = 2.4$	$\frac{3}{2} = 1.5$	4	$\frac{4}{6} = .667$	$\frac{8}{3} = 2.6667$
6	$Spin(6)$	$\frac{32}{15}$	$\frac{8}{5} = 1.6$	$\frac{52}{15} = 3.4667$	$\frac{5}{7} = .714$	$\frac{52}{21} = 2.4762$
d odd	Spin(d)	$\frac{8(2n-1)}{n(2n+1)}$ $= \frac{16(d-2)}{d(d-1)}$	$2 - 1/n$ $= 2 - 2/(d-1)$	$\frac{8(3d-5)}{d(d-1)}$	$\frac{d-1}{d+1}$	$\frac{8(3d-5)}{d(d+1)}$
d even	$Spin(d)$	$\frac{16(d-2)}{d(d-1)}$	$\frac{4(n-1)}{2n-1} = \frac{2d-4}{d-1}$	$\frac{8(3d-5)}{d(d-1)}$	$\frac{d-1}{d+1}$	$\frac{8(3d-5)}{d(d+1)}$
d odd $\rightarrow \infty$	Spin(d)	$\approx 16/d$	$\rightarrow 2$	$\approx 24/d$	$\rightarrow 1$	$\approx 24/d \rightarrow 0$
d even $\rightarrow \infty$	Spin(d)	$\approx 16/d$	$\rightarrow 2$	$\approx 24/d$	$\rightarrow 1$	$\approx 24/d \rightarrow 0$

² The case $d=2$ is only getting its C_A/C_F rather formally by seeking to use the rules roughly of our previous article[1], because the Lorentz group is Abelian.

We see from the table here for simplicity made only for the most competitive case of “with spinor” in the terminology the foregoing table that with column c -goal numbers actually it is $d=3$ rather than the experimental dimension $d=4$ that “wins” (i.e. is largest), but this series of proposed numbers c . is not truly “fair” in as far as one has effectively used only the Lorentz group in the denominator C_F but at least crudely the full Poincare group in the numerator $C_A + C_V$. Thus in order to avoid a simple wrong expected variation of a quadratic Cassimir with the dimensionality of the Lie group, we should at least correct the denominator C_F by multiplying it by the ratio of the dimension of the Poincare Lie group over that of the Lorentz Lie group, $(d+1)/(d-1)$. When we make this “fairness correction” at least crudely getting no obvious wrong Lie-group-dimension-dependent factor in, then the dimension $d=4$ becomes (again) the winner! In fact we get for $d=4$ (the experimental dimension) the goal quantity in column d . equal to $14/5=2.8$ while accidentally the two neighboring dimensions $d=3$ and $d=5$ both gets in stead $8/3 = 2.66667$, which is less.

But notice that it is a rather smooth peaked curve with the peak near the experimental dimension 4, so that the latter becomes the winner among integers, but it is only by a tiny bit it wins. That is to be expected from the smoothness of the variation of the goal number with the dimension d . This smallness of the excess making the $d=4$ be the winner of course makes the uncertainty bigger and my “crude” corrections rather than exactly calculating some welldefined quantity is thus not so convincing. Anyway I think, that the accuracy may be good enough, and the simplicity of the proposed goal quantities sufficient to make it at least highly suggestive, that the coincidence of the winning dimension and the experimental one means that we are on the right track!

4 Conclusion

We have found that a couple of very reasonable specifications of what the extension of our previous [1] quantity to be maximized to obtain the Standard Model gauge group leads to that the maximization of the generalized quantity gives as the “winner” the dimension $d=4$ as is empirically the dimension! That is to say we have found a possible *explanation* for, why we have just 4 (meaning 3+1) dimensions of space-time!

In fact we have extended the main idea of claiming the maximization of essentially the to the Dynkin index related group dependent quantity C_A/C_F (with C_A and C_F being the quadratic Casimirs for respectively the adjoint representation C_A and for that (essentially) faithful representation F chosen so as to maximize the ratio C_A/C_F .) to lead to the experimentally realized group (the Standard Model group). For honesty it should be admitted that the victory of exactly the Standard Model Group were dependent on our slightly ad hoc treatment of the Abelian invariant subgroup - i.e. the $U(1)$ - needed because of the ratio C_A/C_F being rather meaningless a priori for an abelian group. For honesty we must also admit that in reference [1] we sneaked in a dependence on the *group* rather than only *Lie algebra* by considering the volume of the *group* which depends on the identification of center elements, properties being revealed phenomenologically via the representations of the gauge group (except for the case of $d=2$ this detail is though not relevant for the Lorentz group and thus the dimension).

Our extension consists in that we in stead of the gauge group rather consider the *Lorentz group*, or we even seek to use the *Poincare group*. Then of course our quantity C_A/C_F or slight modifications/“improvements” of it - enumerated *a.*, *b.*, *c.*, *d.* - will depend on the dimension d of space time. The dimension d gives of course a different Lorentz group for each value of d . We then inserted this d -dependent Lorentz group in stead of the gauge group which were studied in last paper[1]. The various modifications, *a.*, *b.*, *c.*, *d.*, shall be considered attempts to use the *Poincare group* instead of the Lorentz group, but rather than truly doing that, make some approximate treatment as if crudely using the Poincare group.

The results of the search for the dimension having the largest “goal-quantity”, using various proposals for the exact form such as *a.* meaning C_A/C_F simply, are the following:

- *a.* The simple quantity C_A/C_F for the *Lorentz group* with the same formal assignment for abelian group as used in [1], here making $d=2$ noncompetitive (but at least having a score formally), leads to $d=3$ and $d=4$ standing even, both scoring the same number $C_A/C_F = 8/3$.
- *b.* Making a crude correction to consider instead the quantity $(C_A/C_F)^{(d-1)/(d+1)}$ leads to that the experimental dimension of space time $d=4$ gets the largest score. The meaning of this slight modification of *a.* is that we make an attempt to take the group to replace the gauge group in our previous paper[1] to be the Poincare group rather than the Lorentz group. We, however, only make a crude attempt in that direction. Since the Poincare has the translation subgroups which are by themselves Abelian we naturally tend to use formally - just like in reference [1] - to assign a factor 1 to the abelian groups. Then we average in the logarithm our goal numbers for the various factors into which the group falls weighting with the dimension in the Lie algebra.

The inclusion of the Poincare group is not done in a fully correct way though in as far as we only consider the faithful representations of the Lorentz group and only extend a bit speculatively to weight as if we had the Poincare group.

- *c.* Still thinking of crudely using the Poincare group rather than the Lorentz group, we

proposed to still take a representation F only of the Lorentz group, but evaluating the quadratic Cassimir for the Poincare group, although that sounds not quite “fair”. The quadratic Cassimir we used here under $c.$ for “the Poincare group” were taken to $C_A + C_V$, where V denotes the vector representation and thus C_V its quadratic Cassimir. In this “unfair” game the dimension for space time $d=3=2+1$ got the highest score. So our hoped for victory of the experimental dimension failed in this “unfair” proposal. *But since I stress the “unfairness” of this proposal, we should not take this proposal seriously.*

- $d.$ This last proposal in the present article is a crude attempt at least to correct for, that the ratio of the dimensions of the Poincare and the Lorentz Lie groups is space-time dimension d dependent. That is to say, we argue that the quadratic Cassimir C_F for the representation F of the Lorentz group should at least be scaled so as to correspond to a representation rather of the Poincare group by being multiplied by the ratio of the Lie group dimensions of the Poincare group relative to that of the Lorentz group, $\frac{d(d-1)/2+d}{d(d-1)/2} = \frac{d+1}{d-1}$. That is to say we perform the crude correction of replacing

$$C_F \rightarrow C_F * \frac{d+1}{d-1}. \quad (54)$$

Since the quantity C_F occurs in the denominator of the quantity $(C_F + C_V)/C_F$ maximized under $c.$ of course this quantity is scaled the opposite way, and the goal-quantity in this proposal $d.$ is taken as

$$\text{“goal quantity } d.” = \frac{(C_A + C_V)(d-1)}{C_F(d+1)}. \quad (55)$$

Now the result becomes that *the experimental dimension $d=4$ has the largest value for the goal quantity $d.$* , in as far as it gets

$$\frac{(C_A + C_V)(d-1)}{C_F(d+1)} \Big|_{d=4} = \frac{14}{5} = 2.8, \quad (56)$$

while by accident the two neighboring space time dimensions 3 and 5 score only $\frac{8}{3} = 2.6667$. So indeed the *experimental space time dimension 4 won the most developed suggestion $d.$*

This means that apart from the “unfair” proposal $c.$, all the four proposals here have the space time dimension $d=4$ realized in nature obtain a largest “goal quantity” among the winners! In $a.$ $d=3$ and $d=4$ share the winner place, but in the two other “fair” proposals $b.$ and $d.$ it is indeed space time dimension $d=4$, the experimental one, that gives the highest “goal quantity”.

Taking this result serious, and not as being just accidental coincidence or a result of inventive construction, it must tell us about the reason for that the dimensionality became just $d=4$. Taking our result serious we must look for what is the spirit behind the proposals above, so as to obtain an answer to “Why did we get just $d=4$ space time dimensions?”. This “spirit” behind the proposals here set up to select the experimental dimension $d=4$ seems to be that the group - the Poincare or Lorentz group or the gauge group say - should be *representable in a way where the matrices or other objects on which the group is represented are relatively slowly varying under the group.*

We may, if taking this “slowness” of the motion of the representative in the representation with the group element serious, seek to invent a model behind the 4-dimensional space time and the Standard Model gauge group that could explain that slowness. One possible such explanation could be:

Truly the fundamental physics model or theory is “random” and that without the symmetries we seek to explain. Then “by accident” there appears approximately some symmetries - and we here hope for say some Lorentz invariance symmetry -. Now we dream that there may be some way in which such an approximate symmetry can be automatically become exact in practice. We, Førster, Ninomiya and me [10](see also Damgaard et al. [18]), have actually argued that gauge symmetry with electrodynamics (and Yang Mills theory) as example can occur in a whole phase in practice giving precisely the *massless* photon in that phase. Thus we can for symmetries that can somehow be considered gauge symmetry - as can the Lorentz symmetry in general relativity - speculate that such symmetries appear in practice as exact provided though, that they are there approximately at first[22, 23].. But now the crux of the matter is that *if a symmetry is represented by slowly moving matrices or whatever, then one must expect that statistically it would be easier to get the symmetry approximately by accident.* If it were such that the fundamental theory could be considered random and only obtaining some symmetries by accident - at first approximately, but perhaps made exact by some mechanism[10, 22, 23, 18] - then we could consider the practically random Lagrange or action as taking random values for regions of some (small) size in the value space for the *representaion* of the group which gives the transformation properties of the fields or degrees of freedom under the group in question. Now when a group is represented by a representation, which in some sense is the represented matrix or field, and these fields or matrices move slowly for an appropriately normalized motion of the group element represented, then one can vary the group element much before one varies the representation field much. But this means that one needs less good luck to get a symmetry accidentally the slower the representation moves, because the displacement inside the group (itself) corresponding to one of the (small) size regions (over which we assume essential constancy of the action) becomes bigger the slower the representaion motion rate.

The crucial point should be that one would with the in some sense random action have a better chance to obtain by accident a certain symmetry, when this symmetry is represented on the fields or degrees of freedom by a “slowly moving representaion”, so such a symmetry would more often occur by accident, if one thinks this random action way.

So when our various “goal quantities” favour the experimentally found gauge group and the dimension of space time, it means that the groups realized in nature are the ones that have the optimal chance to come out of a random action model. This is so because these goal quantities being large means that the representation motion is slow.

So the message from the gauge group and the dimension is that such a random action philosophy is one possible mechanism behind the choice by nature of the gauge groups and dimension.

The idea of there being made in some sense a lot of attempts randomly of groups to be tested off could be said to have remiscense of the idea of a gaugeglass [21], which though rather means the action is random quenched randomly locally, but that the gauge group is given from the start; but the spirit is similar.

A priori one should speculate about possible other machineries that could explain that precisely our type of “goal quantities”; but at first it seems that the random action type of model allowing symmetries of the type with highest goal quantities is a good idea and very likely something like that could be the reason behind the choice by nature of the gauge group (of the Standard Model) and of the dimension.

In any case we have found a surprisingly simple principle - the maximization of our to each other rather closely related “goal quantities” - leading to both the gauge group of the Standard Model *and* the dimension of space time being 4.

Let me stress that a work of the present type and of [1] - finding a goal quantity leading to

the realized groups - is an attempt to ask phenomenologically whether there is some signal in the details of the phenomenologically present theory that successively can give us hint(s) about the more fundamental theory behind the presently working Standard Model with its gauge group $S(U(2) \times U(3))$ and the seeming dimension 4.

5 Acknowledgement

I would like to thank the Niels Bohr Institute for allowing me status as professor emeritus and for economical support and Matjaz Breskvar support for visiting the Bled Conference "Beyond the Standard Models" wherein the first paper[1] in the present series of two articles were presented together with Don Bennett - although a year earlier though -, who is also thanked together with colleagues discussing there the previous work.

References

- [1] Don Bennett and H. B. Nielsen, "Seeking...", Contribution to the workshop "Beyond the Standard Models", Bled 2011. arXiv:1306.2963v1.
- [2] Max Tegmark, "On the dimensionality of space time", *Class. Quantum Grav.* 14, 1.69 - 1.75 (1997).
- [3] Ehrenfest,P., 1917 *Proc. Amsterdam Acad.* **20** 200
- [4] Gordon Kane : <http://particle-theory.physics.lsa.umich.edu/kane/modern.html>
- [5] Norma Mankoc et al. see many contributions to the present and previous Bled Proceedings.
- [6] H. B. Nielsen and N. Brene, "Spontaneous Emergence Of Gauge Symmetry," IN *KRAKOW 1987, PROCEEDINGS, SKYRMIONS AND ANOMALIES*, 493-498 AND COPENHAGEN UNIV. - NBI-HE-87-28 (87,REC.JUN.) 6p H. B. Nielsen and N. Brene, "Skewness Of The Standard Model: Possible Implications," *Physicalia Magazine, The Gardener of Eden*, 12 (1990) 157; NBI-HE-89-38; H. B. Nielsen and N. Brene, "What Is Special About The Group Of The Standard Model?," *Phys. Lett. B* **223** (1989) 399.
- [7] H.B. Nielsen, S.E. Rugh and C. Surlykke, Seeking Inspiration from the Standard Model in Order to Go Beyond It, *Proc. of Conference held on Korfu* (1992)
- [8] Philippe Di Francesco, Pierre Mathieu, David Snychal, *Conformal Field Theory*, 1997 Springer-Verlag New York, ISBN 0-387-94785-X
- [9] RANDOM DYNAMICS: H. B. Nielsen, "Dual Strings," "Fundamentals of quark models", In: *Proc. of the Seventeenth Scott. Univ. Summer School in Physics*, St. Andrews, august 1976. I.M. Barbour and A.T. Davies(eds.), Univ. of Glasgow , 465-547 (publ. by the Scott.Univ. Summer School in Physics, 1977) (CITATION = NBI-HE-74-15;)
H.B. Nielsen, Har vi brug for fundamentale naturlove(in Danish) (meaning:"Do we need laws of Nature?") *Gamma* 36 page 3-16, 1978(1. part) and *Gamma* 37 page 35-46, 1978 (2. part)
H.B. Nielsen and C. D. Froggatt, "Statistical Analysis of quark and lepton masses", *Nucl. Phys.* **B164**(1979) 114 - 140.

- [10] D. Førstner, H.B. Nielsen, and M. Ninomiya, “Dynamical stability of local gauge symmetry. Creation of light from chaos.” *Phys. Lett.* **B94**(1980) 135 -140
- [11] H.B. Nielsen, *Lecture notes in Physics* **181**, “Gauge Theories of the Eighties” In: *Proc. of the Arctic School of Physics 1982, Akaeslompolo, Finland, Aug. 1982.* R. Raitio and J. Lindfors(eds.). Springer, Berlin, 1983,p. 288-354.
H.B. Nielsen, D.L. Bennett and N. Brene: “The random dynamics project from fundamental to human physics”. *In: Recent developments in quantum field theory.* J. Ambjoern, B.J. Durhuus and J.L. Petersen(eds.), Elsevier Sci.Publ. B.V., 1985, pp. 263-351
- [12] See the “home page of Random Dynamics”: <http://www.nbi.dk/~kleppe/random/qa/qa.html>
- [13] H.B. Nielsen and D. L. Bennett, “The Gauge Glass: A short review”, Elaborated version of talk at the Conf. on Disordered Systems, Copenhagen, September 1984. Nordita preprint 85/23.
- [14] see e.g. John Baez: <http://math.ucr.edu/home/baez/renormalizability.html>, November 2006
- [15] P. Candelas, Gary T. Horowitch, Andrew Strominger, and Edward Witten, “Vacuum configurations for superstrings”,NFS-ITP-84-170.
- [16] Candelas...Witten, *Nuclear Physics B* 258: 46 % G % 74, Bibcode:1985NuPhB.258...46C, doi:10.1016/0550-3213(85)90602-9
- [17] See e.g. John M. Pierre, “Superstrings Extra dimensions” <http://www.sukidog.com/jpierre/strings/extradim.htm>
- [18] P.H. Damgaard, N. Kawamoto, K. Shigemoto: *Phys. Rev. Lett.*53, 2211 (1984)
- [19] A J Macfarlane and Hendryk Pfeiffer, *J. Phys. A: Math. Gen.* 36 (2003) 2305 200 2232317 PII: S0305-4470(03)56335-1 Representations of the exceptional and other Lie algebras with integral eigenvalues of the Casimir operator
- [20] T. van Ritbergen, A. N. Shellekens, J. A. M. Vermaseren, UM-TH-98-01 NIKHEF-98-004 Group theory factors for Feynman ... [www.nikhef.nl/ form/maindir/oldversions/.../packages/.../color.ps](http://www.nikhef.nl/form/maindir/oldversions/.../packages/.../color.ps)
- [21] *Physics Letters B* Volume 208, Issue 2, 14 July 1988, Pages 275-280, <http://arxiv.org/abs/hep-ph/9311321>
<http://arxiv.org/abs/hep-ph/9607341>
Physics Letters B Volume 178, Issues 2-3, 2 October 1986, Pages 179-186
H.B. Nielsen and N. Brene, Gauge Glass, *Proc. of the XVIII International Symposium on the Theory of Elementary Particles, Ahrenshoop, 1985* (Institut fur Hochenergiphysik, Akad. der Wissenschaften der DDR, Berlin-Zeuthen, 1985);
- [22] Lehto, M., Nielsen H. B., and Ninomiya, M. (1986). Pregeometric quantum lattice: A general discussion. *Nuclear Physics B*, 272, 213-227. Lehto, M., Nielsen H. B., and Ninomiya, M. (1986). Diffeomorphism symmetry in simplicial quantum gravity. *Nuclear Physics B*, 272, 228-252. Lehto, M., Nielsen H. B., and Ninomiya, M. (1989). Time translational symmetry. *Physics Letters B*, 219, 87-91.

- [23] M. Lehto, H. B. Nielsen, and Masao Ninomiya “A correlation decay theorem at high temperature” *Comm. Math. Phys.* Volume 93, Number 4 (1984), 483-493.