

Convolution properties of some harmonic mappings in the right-half plane

Raj Kumar^a*, Michael Dorff^{b†}, Sushma Gupta^a, and Sukhjit Singh^a

^a Sant Longowal Institute of Engineering and Technology, Longowal-148106 (Punjab), India.

^b Department of Mathematics, Brigham Young University, Provo, Utah, 84602, USA.

Abstract

Dorff, proved in [2] that the convolution of two harmonic right-half plane mappings is convex in the direction of real axis provided that the convolution is locally univalent and sense preserving. Later, it was shown in [3] that the condition of locally univalent and sense preserving can be dropped in some special cases. In this paper, we generalize the main result from [3].

Key Words : Univalent harmonic mappings, mappings in right half-plane, convolution.

AMS Subject Classification: 30C45.

1 Introduction

Let $f = u + iv$ be a continuous complex-valued harmonic mapping in the open unit disk $E = \{z : |z| < 1\}$, where both u and v are real-valued harmonic functions in E . Such a mapping can be decomposed into two parts and can be expressed as $f = h + \bar{g}$. Here h is known as the analytic part and g the co-analytic part of f . Lewy's Theorem implies that a harmonic mapping $f = h + \bar{g}$ defined in E , is locally univalent and sense preserving if and only if the Jacobian of the mapping, defined by $J_f = |h'|^2 - |g'|^2$, is positive or equivalently, if and only if $h'(z) \neq 0$ and the dilatation function ω of f , defined by $\omega(z) = \frac{g'(z)}{h'(z)}$, satisfies $|\omega(z)| < 1$ in E . We denote by S_H the class of all harmonic, sense-preserving and univalent mappings $f = h + \bar{g}$, defined in E which are normalized by the conditions $h(0) = 0$ and $h_z(0) = 1$. Therefore, a function $f = h + \bar{g}$ in class S_H has the representation,

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n + \sum_{n=1}^{\infty} \overline{b_n z^n}, \quad (1)$$

*rajgarg2012@yahoo.co.in

†mdorff@math.byu.edu

for all z in E . The class of functions of the type (1) with $\bar{b}_1 = 0$ is denoted by S_H^0 which is a subset of S_H . Further let K_H (respectively K_H^0) be the subclass of S_H (respectively S_H^0) consisting of functions which map the unit disk E onto convex domains. A domain Ω is said to be convex in the direction $\phi, 0 \leq \phi < \pi$, if every line parallel to the line joining 0 and $e^{i\phi}$ has a connected intersection with Ω . In particular, a domain convex in horizontal direction is denoted by CHD.

Definition 1.1. Convolution or Hadamard product of two harmonic mappings $F(z) = H + \bar{G} = z + \sum_{n=2}^{\infty} A_n z^n + \sum_{n=1}^{\infty} \bar{B}_n \bar{z}^n$ and $f(z) = h + \bar{g} = z + \sum_{n=2}^{\infty} a_n z^n + \sum_{n=1}^{\infty} \bar{b}_n \bar{z}^n$ in S_H is defined as

$$\begin{aligned} (F * f)(z) &= (H * h)(z) + \overline{(G * g)(z)} \\ &= z + \sum_{n=2}^{\infty} a_n A_n z^n + \sum_{n=1}^{\infty} \overline{b_n B_n} \bar{z}^n. \end{aligned}$$

Let $f_a = h_a + \bar{g}_a \in K_H$ be the mapping in the right half-plane given by $h_a + g_a = \frac{z}{1-z}$ with dilatation function $\omega_a(z) = \frac{a-z}{1-az}$ ($|a| < 1, a \in \mathbb{R}$). Then, by using the shearing technique (see [1]), we get

$$h_a(z) = \frac{\frac{1}{1+a}z - \frac{1}{2}z^2}{(1-z)^2} \quad \text{and} \quad g_a(z) = \frac{\frac{a}{1+a}z - \frac{1}{2}z^2}{(1-z)^2}. \quad (2)$$

By setting $a = 0$, we get $f_0 = h_0 + \bar{g}_0 \in K_H^0$, the standard right half-plane mapping, where

$$h_0(z) = \frac{z - \frac{1}{2}z^2}{(1-z)^2} \quad \text{and} \quad g_0(z) = \frac{-\frac{1}{2}z^2}{(1-z)^2}. \quad (3)$$

Unlike the case of analytic functions, the convolution of two univalent convex harmonic functions is not necessarily convex harmonic. It may not even be univalent. So, it is interesting to explore the convolution properties of mappings in the class K_H . In [2] and [3], the authors obtained several results in this direction. In particular they proved the following:

Theorem A. (See [2]) Let $f_1 = h_1 + \bar{g}_1, f_2 = h_2 + \bar{g}_2 \in S_H^0$ with $h_i + g_i = \frac{z}{1-z}$ for $i = 1, 2$. If $f_1 * f_2$ is locally univalent and sense preserving, then $f_1 * f_2 \in S_H^0$ and is CHD.

Theorem B. (See [3]) Let $f = h + \bar{g} \in K_H^0$ with $h + g = \frac{z}{1-z}$ and $\omega(z) = e^{i\theta} z^n$ ($n \in \mathbb{N}$ and $\theta \in \mathbb{R}$). If $n = 1, 2$, then $f_0 * f \in S_H^0$ and is CHD, where f_0 is given by (3).

In Theorem B, the authors established that the requirement that the convolution should be locally univalent and sense preserving in Theorem A can be dropped. Recently, Li and Ponnusamy [5,6] also obtained some results involving convolutions of right half-plane and slanted right half-plane harmonic mappings. In [6], they proved:

Theorem C. Let f_0 be given by (3). If $f = h + \bar{g}$, is a slanted right half-plane mapping, given by $h + e^{-2i\alpha}g = \frac{z}{1 - e^{i\alpha}z}$ ($0 \leq \alpha < 2\pi$) with $\omega(z) = e^{i\theta}z^n$ ($\theta \in \mathbb{R}$), then for $n = 1, 2$, $f * f_0 \in S_H^0$ and is convex in the direction of $-\alpha$.

In [3] and [6], authors showed that Theorem B and Theorem C do not hold for $n \geq 3$. The aim of the present paper is to investigate the convolution properties of harmonic mappings f_a ($a \in \mathbb{R}, |a| < 1$) defined by (2) with right half-plane mappings $f_n = h + \bar{g}$ where $h + g = \frac{z}{1 - z}$ and dilatation $\omega(z) = e^{i\theta}z^n$ ($\theta \in \mathbb{R}, n \in \mathbb{N}$). We establish that $f_a * f_n$ are in S_H and are CHD for all $a \in \left[\frac{n-2}{n+2}, 1\right)$ and for all $n \in \mathbb{N}$. A condition is also determined under which $f_a * f_b$ is CHD and belongs to S_H .

2 Main Results

We begin by proving the following lemma.

Lemma 2.1. Let $f_a = h_a + \bar{g}_a$ be defined by (2) and $f = h + \bar{g} \in S_H$ be the right half-plane mapping, where $h + g = \frac{z}{1 - z}$ with dilatation $\omega(z) = \frac{g'(z)}{h'(z)}$ ($h'(z) \neq 0, z \in E$). Then $\tilde{\omega}_1$, the dilatation of $f_a * f$, is given by

$$\tilde{\omega}_1 = \left[\frac{2\omega(a - z)(1 + \omega) + z\omega'(a - 1)(1 - z)}{2(1 - az)(1 + \omega) + z\omega'(a - 1)(1 - z)} \right]. \quad (4)$$

Proof. From $h + g = \frac{z}{1 - z}$ and $g' = \omega h'$, we immediately get

$$h'(z) = \frac{1}{(1 + \omega(z))(1 - z)^2} \quad \text{and} \quad h''(z) = \frac{2(1 + \omega(z)) - \omega'(z)(1 - z)}{(1 + \omega(z))^2(1 - z)^3}.$$

Let

$$f_a * f = h_a * h + \overline{g_a * g} = h_1 + \bar{g}_1, \text{ where}$$

$$\begin{aligned} h_1(z) &= \frac{1}{2} \left[\frac{z}{1 - z} + \frac{(1 - a)z}{(1 + a)(1 - z)^2} \right] * h \\ &= \frac{1}{2} \left[h + \frac{(1 - a)}{(1 + a)} zh' \right] \end{aligned}$$

and

$$\begin{aligned} g_1(z) &= \frac{1}{2} \left[\frac{z}{1 - z} - \frac{(1 - a)z}{(1 + a)(1 - z)^2} \right] * g \\ &= \frac{1}{2} \left[g - \frac{(1 - a)}{(1 + a)} zg' \right]. \end{aligned}$$

Now the dilatation $\tilde{\omega}_1$ of $f_a * f$ is given by

$$\begin{aligned}\tilde{\omega}_1(z) &= \frac{g'_1(z)}{h'_1(z)} = \left[\frac{2ag' - (1-a)zg''}{2h' + (1-a)zh''} \right] \\ &= \left[\frac{2a\omega h' - z(1-a)(\omega h'' + \omega' h')}{2h' + (1-a)zh''} \right] \\ &= \left[\frac{2\omega(a-z)(1+\omega) + z\omega'(a-1)(1-z)}{2(1-az)(1+\omega) + z\omega'(a-1)(1-z)} \right].\end{aligned}$$

□

We shall also need the following forms of Cohn's rule and Schur-Cohn's algorithm .

Lemma A. (Cohn's rule [7, p.375]) Given a polynomial

$$t(z) = a_0 + a_1z + a_2z^2 + \dots + a_nz^n$$

of degree n , let

$$t^*(z) = z^n \overline{\left(\frac{1}{z} t \right)} = \bar{a}_n + \bar{a}_{n-1}z + \bar{a}_{n-2}z^2 + \dots + \bar{a}_0z^n.$$

Denote by r and s the number of zeros of $t(z)$ inside and on the unit circle $|z| = 1$, respectively.

If $|a_0| < |a_n|$, then

$$t_1(z) = \frac{\bar{a}_n t(z) - a_0 t^*(z)}{z}$$

is of degree $n-1$ and has $r_1 = r-1$ and $s_1 = s$ number of zeros inside the unit circle and on it, respectively.

Lemma B. (Schur-Cohn's algorithm [7, p.383]) Given a polynomial

$$r(z) = a_0 + a_1z + \dots + a_nz^n + a_{n+1}z^{n+1}$$

of degree $n+1$, let

$$M_k = \det \begin{pmatrix} \bar{B}_k^T & A_k \\ \bar{A}_k^T & B_k \end{pmatrix} \quad (k = 1, 2, \dots, n+1),$$

where A_k and B_k are the triangular matrices

$$A_k = \begin{pmatrix} a_0 & a_1 & \dots & a_{k-1} \\ & a_0 & \dots & a_{k-2} \\ & & \ddots & \vdots \\ & & & a_0 \end{pmatrix}, \quad B_k = \begin{pmatrix} \bar{a}_{n+1} & \bar{a}_n & \dots & \bar{a}_{(n+1)-k+1} \\ & \bar{a}_{n+1} & \dots & \bar{a}_{(n+1)-k+2} \\ & & \ddots & \vdots \\ & & & \bar{a}_{n+1} \end{pmatrix}.$$

Then $r(z)$ has all its zeros inside the unit circle $|z| = 1$ if and only if the determinants M_1, M_2, \dots, M_{n+1} are all positive.

We now proceed to state and prove our main result.

Theorem 2.2. *Let $f_a = h_a + \overline{g_a}$ be given by (2). If $f_n = h + \overline{g}$ is the right half-plane mapping given by $h + g = \frac{z}{1-z}$ with $\omega(z) = e^{i\theta} z^n$ ($\theta \in \mathbb{R}, n \in \mathbb{N}$), then $f_a * f_n \in S_H$ and is CHD for $a \in [\frac{n-2}{n+2}, 1)$.*

Proof. In view of Theorem A, it suffices to show that the dilatation of $f_a * f_n = \tilde{\omega}_1$ satisfies $|\tilde{\omega}_1(z)| < 1$, for all $z \in E$. Setting $\omega(z) = e^{i\theta} z^n$ in (4), we get

$$\begin{aligned} \tilde{\omega}_1(z) &= -z^n e^{2i\theta} \left[\frac{z^{n+1} - az^n + \frac{1}{2}(2 + an - n)e^{-i\theta}z + \frac{1}{2}(n - 2a - an)e^{-i\theta}}{\frac{1}{2}(n - 2a - an)e^{i\theta}z^{n+1} + \frac{1}{2}(2 + an - n)e^{i\theta}z^n - az + 1} \right] \\ &= -z^n e^{2i\theta} \frac{p(z)}{p^*(z)}, \text{ where} \end{aligned} \quad (5)$$

$$p(z) = z^{n+1} - az^n + \frac{1}{2}(2 + an - n)e^{-i\theta}z + \frac{1}{2}(n - 2a - an)e^{-i\theta} \quad (6)$$

and $p^*(z) = z^{n+1} \overline{p\left(\frac{1}{\overline{z}}\right)}$.

Obviously, if z_0 is a zero of p then $\frac{1}{\overline{z_0}}$ is a zero of p^* . Hence, if A_1, A_2, \dots, A_{n+1} are the zeros of p (not necessarily distinct), then we can write

$$\tilde{\omega}_1(z) = -z^n e^{2i\theta} \frac{(z - A_1)}{(1 - \overline{A_1}z)} \frac{(z - A_2)}{(1 - \overline{A_2}z)} \dots \frac{(z - A_{n+1})}{(1 - \overline{A_{n+1}}z)}.$$

Now for $|A_i| \leq 1$, $\frac{(z - A_i)}{(1 - \overline{A_i}z)}$ maps $\overline{E} = \{z : |z| \leq 1\}$ onto \overline{E} . So in order to prove our theorem, we will show that A_1, A_2, \dots, A_{n+1} lie inside or on the unit circle $|z| = 1$ for $a \in \left(\frac{n-2}{n+2}, 1\right)$ (in the case $a = \frac{n-2}{n+2}$, from (5) we see that $|\tilde{\omega}_1(z)| = |-z^n e^{i\theta}| < 1$). To do this we will use Lemma B by considering the following two cases.

Case 1. *When $n = 1$.* In this case $p(z) = z^2 + \left[-a + \frac{1}{2}(1+a)e^{-i\theta}\right]z + \frac{1}{2}(1-3a)e^{-i\theta}$ and $a \in \left(-\frac{1}{3}, 1\right)$. Thus, by comparing $p(z)$ and $r(z)$ of Lemma B, we have

$$\begin{aligned} M_1 &= \det \begin{pmatrix} a_2 & a_0 \\ \overline{a_0} & \overline{a_2} \end{pmatrix} = \det \begin{pmatrix} 1 & \frac{1}{2}(1-3a)e^{-i\theta} \\ \frac{1}{2}(1-3a)e^{i\theta} & 1 \end{pmatrix} \\ &= \frac{3}{4}(1-a)(1+3a) > 0, \quad \text{and} \end{aligned}$$

$$\begin{aligned}
M_2 &= \det \begin{pmatrix} a_2 & 0 & a_0 & a_1 \\ a_1 & a_2 & 0 & a_0 \\ \bar{a}_0 & 0 & \bar{a}_2 & \bar{a}_1 \\ \bar{a}_1 & \bar{a}_0 & 0 & \bar{a}_2 \end{pmatrix} \\
&= \det \begin{pmatrix} 1 & 0 & \frac{1}{2}(1-3a)e^{-i\theta} & -a + \frac{1}{2}(1+a)e^{-i\theta} \\ -a + \frac{1}{2}(1+a)e^{-i\theta} & 0 & 1 & \frac{1}{2}(1-3a)e^{i\theta} \\ \frac{1}{2}(1-3a)e^{i\theta} & 0 & 1 & -a + \frac{1}{2}(1+a)e^{i\theta} \\ -a + \frac{1}{2}(1+a)e^{i\theta} & \frac{1}{2}(1-3a)e^{i\theta} & 0 & 1 \end{pmatrix} \\
&= \frac{1}{2}(1-a)^2(1+3a)^2 \cos^2 \frac{\theta}{2} > 0, \text{ for } \theta \neq (2m+1)\pi, m \in \mathbb{N}.
\end{aligned}$$

If $\theta = (2m+1)\pi$, $m \in \mathbb{N}$, then $p(z) = z^2 - \frac{1}{2}(1+3a)z - \frac{1}{2}(1-3a)$. Obviously, $z = 1$ and $z = -\frac{1}{2}(1-3a)$ are zeros of $p(z)$ and lie on and inside the unit circle $|z| = 1$, respectively, for $-\frac{1}{3} < a < 1$.

Case 2. When $n \geq 2$. In this case $p(z) = z^{n+1} - az^n + \frac{1}{2}(2+an-n)e^{-i\theta}z + \frac{1}{2}(n-2a-an)e^{-i\theta}$. Again comparing $p(z)$ and $r(z)$, let

$$M_k = \det \begin{pmatrix} \bar{B}_k^T & A_k \\ \bar{A}_k^T & B_k \end{pmatrix} \quad (k = 1, 2, 3, \dots, n+1),$$

where A_k and B_k are as defined in Lemma B with $a_{n+1} = 1, a_n = -a, a_{n-1} = 0, \dots, a_2 = 0, a_1 = \frac{1}{2}(2+an-n)e^{-i\theta}$ and $a_0 = \frac{1}{2}(n-2a-an)e^{-i\theta}$. Since $a_{n+1} = 1$, therefore $\det(B_k) = 1$ and so,

$$\begin{pmatrix} \bar{B}_k^T & A_k \\ \bar{A}_k^T & B_k \end{pmatrix} \begin{pmatrix} I & 0 \\ -B_k^{-1}\bar{A}_k^T & I \end{pmatrix} = \begin{pmatrix} \bar{B}_k^T - A_k B_k^{-1} \bar{A}_k^T & A_k \\ 0 & B_k \end{pmatrix}$$

which gives

$$M_k = \det \begin{pmatrix} \bar{B}_k^T & A_k \\ \bar{A}_k^T & B_k \end{pmatrix} = \det [\bar{B}_k^T - A_k B_k^{-1} \bar{A}_k^T].$$

Now we consider the following two subcases.

Subcase 1. When $k = 1, 2, 3, \dots, n$. We will show that in this case, $M_k = \left(\frac{1}{4}\right)^k n^{k-1}(n+2k)(2-n+2a+an)^k(1-a)^k$, which is positive for $a \in \left(\frac{n-2}{n+2}, 1\right)$.

In this case A_k and B_k are the following $k \times k$ matrices;

$$A_k = \begin{pmatrix} a_0 & a_1 & a_2 & \cdots & a_{n-1} \\ 0 & a_0 & a_1 & \cdots & a_{n-2} \\ 0 & 0 & a_0 & \cdots & a_{n-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & a_0 \end{pmatrix} = \begin{pmatrix} a_0 & a_1 & 0 & \cdots & 0 \\ 0 & a_0 & a_1 & \cdots & 0 \\ 0 & 0 & a_0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & a_0 \end{pmatrix}, \quad B_k = \begin{pmatrix} \bar{a}_{n+1} & \bar{a}_n & \bar{a}_{n-1} & \cdots & \bar{a}_2 \\ 0 & \bar{a}_{n+1} & \bar{a}_n & \cdots & \bar{a}_3 \\ 0 & 0 & \bar{a}_{n+1} & \cdots & \bar{a}_4 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \bar{a}_{n+1} \end{pmatrix} = \begin{pmatrix} 1 & -a & 0 & \cdots & 0 \\ 0 & 1 & -a & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}.$$

We can compute

$$\bar{B}_k^T - A_k B_k^{-1} \bar{A}_k^T = \begin{pmatrix} 1 - \bar{a}_0 a_0 - \bar{a}_1 (a a_0 + a_1) & -\bar{a}_0 (a a_0 + a_1) - \bar{a}_1 a (a a_0 + a_1) & a[-\bar{a}_0 (a a_0 + a_1) - \bar{a}_1 a (a a_0 + a_1)] & \cdots & -\bar{a}_0 a^{k-2} (a a_0 + a_1) \\ -a - \bar{a}_1 a_0 & 1 - \bar{a}_0 a_0 - \bar{a}_1 (a a_0 + a_1) & -\bar{a}_0 (a a_0 + a_1) - \bar{a}_1 a (a a_0 + a_1) & \cdots & -\bar{a}_0 a^{k-3} (a a_0 + a_1) \\ 0 & -a - \bar{a}_1 a_0 & 1 - \bar{a}_0 a_0 - \bar{a}_1 (a a_0 + a_1) & \cdots & -\bar{a}_0 a^{k-4} (a a_0 + a_1) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 - \bar{a}_0 a_0 \end{pmatrix}.$$

Now

- (a) $1 - \bar{a}_0 a_0 - \bar{a}_1 (a a_0 + a_1) = \frac{1}{4} n (2 - n + 2a + an) (1 - a) (2 - a)$.
- (b) $-\bar{a}_0 (a a_0 + a_1) - \bar{a}_1 a (a a_0 + a_1) = -\frac{1}{4} n (2 - n + 2a + an) (1 - a)^3$.
- (c) $\bar{a}_0 a^{k-m} (a a_0 + a_1) = \frac{1}{4} a^{k-m} (n - 2a - an) (2 - n + 2a + an) (1 - a)$, $m = 1, 2, 3, \dots, k$.
- (d) $-a - \bar{a}_1 a_0 = -\frac{1}{4} n (2 - n + 2a + an) (1 - a)$.
- (e) $1 - \bar{a}_0 a_0 = \frac{1}{4} (n + 2) (2 - n + 2a + an) (1 - a)$.

Therefore,

$$\begin{aligned} M_k &= \left[\left(\frac{1}{4} \right)^k n^{k-1} (2 - n + 2a + an)^k (1 - a)^k \right] \det \begin{pmatrix} (2-a) & -(1-a)^2 & -a(1-a)^2 & \cdots & -a^{k-2}(n-2a-an) \\ -1 & (2-a) & -(1-a)^2 & \cdots & -a^{k-3}(n-2a-an) \\ 0 & -1 & (2-a) & \cdots & -a^{k-4}(n-2a-an) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & (n+2) \end{pmatrix} \\ &= \left[\left(\frac{1}{4} \right)^k n^{k-1} (2 - n + 2a + an)^k (1 - a)^k \right] \det \begin{pmatrix} (2-a) & -(1-a)^2 & -a(1-a)^2 & \cdots & -a^{k-2}(n-2a-an) \\ 0 & \frac{3-2a}{2-a} & -\frac{2(1-a)^2}{2-a} & \cdots & -\frac{2a^{k-3}(n-2a-an)}{2-a} \\ 0 & 0 & \frac{4-3a}{3-2a} & \cdots & -\frac{3a^{k-4}(n-2a-an)}{3-2a} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \frac{n+2k}{k-(k-1)a} \end{pmatrix} \\ &= \left(\frac{1}{4} \right)^k n^{k-1} (n + 2k) (2 - n + 2a + an)^k (1 - a)^k. \end{aligned}$$

Subcase 2. When $k = n + 1$. In this case,

$$A_{n+1} = \begin{pmatrix} a_0 & a_1 & a_2 & \cdots & a_n \\ 0 & a_0 & a_1 & \cdots & a_{n-1} \\ 0 & 0 & a_0 & \cdots & a_{n-2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & a_0 \end{pmatrix} = \begin{pmatrix} a_0 & a_1 & 0 & \cdots & -a \\ 0 & a_0 & a_1 & \cdots & 0 \\ 0 & 0 & a_0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & a_0 \end{pmatrix} \quad \text{and} \quad B_{n+1} = \begin{pmatrix} \bar{a}_{n+1} & \bar{a}_n & \bar{a}_{n-1} & \cdots & \bar{a}_1 \\ 0 & \bar{a}_{n+1} & \bar{a}_n & \cdots & \bar{a}_2 \\ 0 & 0 & \bar{a}_{n+1} & \cdots & \bar{a}_3 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \bar{a}_{n+1} \end{pmatrix} = \begin{pmatrix} 1 & -a & 0 & \cdots & \bar{a}_1 \\ 0 & 1 & -a & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}.$$

We compute that

$$\overline{B}_{n+1}^T - A_{n+1}B_{n+1}^{-1}\overline{A}_{n+1}^T = \begin{pmatrix} 1 - a_0\overline{a}_0 - \overline{a}_1(aa_0 + a_1) + a^n(aa_0 + a_1) - a(a_0\overline{a}_1 + a) & -\overline{a}_0(aa_0 + a_1) - a\overline{a}_1(aa_0 + a_1) & \cdots \\ -a - \overline{a}_1a_0 + a^{n-1}(aa_0 + a_1) & 1 - \overline{a}_0a_0 - \overline{a}_1(aa_0 + a_1) & \cdots \\ a^{n-2}(aa_0 + a_1) & -a - \overline{a}_1a_0 & \cdots \\ \vdots & \vdots & \ddots \\ aa_0 + a_1 & 0 & \cdots \end{pmatrix}$$

$$\left. \begin{array}{l} a^{n-2}[-\overline{a}_0(aa_0 + a_1) - a\overline{a}_1(aa_0 + a_1)] + \overline{a}_1(\overline{a}_1a_0 + a) \\ a^{n-3}[-\overline{a}_0(aa_0 + a_1) - a\overline{a}_1(aa_0 + a_1)] \\ a^{n-4}[-\overline{a}_0(aa_0 + a_1) - a\overline{a}_1(aa_0 + a_1)] \\ \vdots \\ -a - \overline{a}_1a_0 \end{array} \right\} \begin{array}{l} -\overline{a}_0a^{n-1}(aa_0 + a_1) + \overline{a}_0(a_0\overline{a}_1 + a) \\ -\overline{a}_0a^{n-2}(aa_0 + a_1) \\ -\overline{a}_0a^{n-3}(aa_0 + a_1) \\ \vdots \\ 1 - a_0\overline{a}_0 \end{array}$$

Let E_j be the j^{th} column of $\overline{B}_{n+1}^T - A_{n+1}B_{n+1}^{-1}\overline{A}_{n+1}^T$, where $j = 1, 2, \dots, n+1$. Note that for E_m ($m = 2, 3, \dots, n-1$), the column entries are identical to those of $\overline{B}_k^T - A_kB_k^{-1}\overline{A}_k^T$ in Subcase 1. However, the entries for E_1 , E_n , and E_{n+1} are different. We split E_1 , E_n and E_{n+1} in the following way:

$$E_1 = F_1 + G_1 + H_1, E_n = F_n + G_n, E_{n+1} = F_{n+1} + G_{n+1}, \text{ where}$$

$$F_1^T = [1 - a_0\overline{a}_0 - \overline{a}_1(aa_0 + a_1), -a - \overline{a}_1a_0, 0, \dots, 0]$$

$$G_1^T = [a^n(aa_0 + a_1), a^{n-1}(aa_0 + a_1), a^{n-2}(aa_0 + a_1), \dots, (aa_0 + a_1)]$$

$$H_1^T = [-a(a_0\overline{a}_1 + a), 0, 0, \dots, 0]$$

$$F_n^T = [-a^{n-2}\overline{a}_0(aa_0 + a_1) - a^{n-1}\overline{a}_1(aa_0 + a_1), -a^{n-3}\overline{a}_0(aa_0 + a_1) - a^{n-2}\overline{a}_1(aa_0 + a_1), \dots, -(a + a_0\overline{a}_1)]$$

$$G_n^T = [-\overline{a}_1(\overline{a}_1a_0 + a), 0, 0, \dots, 0]$$

$$F_{n+1}^T = [-\overline{a}_0a^{n-1}(aa_0 + a_1), -\overline{a}_0a^{n-2}(aa_0 + a_1), \dots, 1 - \overline{a}_0a_0]$$

$$G_{n+1}^T = [\overline{a}_0(a_0\overline{a}_1 + a), 0, 0, \dots, 0].$$

$$\text{Now } \det[\overline{B}_{n+1}^T - A_{n+1}B_{n+1}^{-1}\overline{A}_{n+1}^T] = \det[E_1E_2 \cdots E_nE_{n+1}]$$

$$\begin{aligned} &= \det[F_1E_2 \cdots F_nF_{n+1}] + \det[G_1E_2 \cdots F_nF_{n+1}] + \det[H_1E_2 \cdots F_nF_{n+1}] \\ &+ \det[F_1E_2 \cdots F_nG_{n+1}] + \det[G_1E_2 \cdots F_nG_{n+1}] + \det[H_1E_2 \cdots F_nG_{n+1}] \\ &+ \det[F_1E_2 \cdots G_nF_{n+1}] + \det[G_1E_2 \cdots G_nF_{n+1}] + \det[H_1E_2 \cdots G_nF_{n+1}] \\ &+ \det[F_1E_2 \cdots G_nG_{n+1}] + \det[G_1E_2 \cdots G_nG_{n+1}] + \det[H_1E_2 \cdots G_nG_{n+1}]. \end{aligned}$$

We will compute each of these determinants. From Subcase 1,

$$\det[F_1E_2 \cdots F_nF_{n+1}] = \left(\frac{1}{4}\right)^{n+1} n^n(3n+2)(2-n+2a+an)^{n+1}(1-a)^{n+1}. \quad (7)$$

$$\text{Also, } \det[F_1E_2 \cdots F_nG_{n+1}] = (-1)^n[\overline{a}_0(a_0\overline{a}_1 + a)][(-1)^n((a + \overline{a}_1a_0)^n)]$$

$$= \left(\frac{1}{4}\right)^{n+1} n^n(2-n+2a+an)^{n+1}(1-a)^{n+1} \left[\frac{1}{2}e^{i\theta}n(n-2a-an)\right], \quad (8)$$

$$\begin{aligned}
\text{and } \det[H_1 E_2 \cdots F_n F_{n+1}] &= -a(a_0 \bar{a}_1 + a) \det[\bar{B}_n^T - A_n B_n^{-1} \bar{A}_n^T] \\
&= -\left(\frac{1}{4}\right)^{n+1} a n^n (2-n+2a+an)^{n+1} (1-a)^{n+1} (3n). \tag{9}
\end{aligned}$$

Next $\det[G_1 E_2 \cdots F_n F_{n+1}]$

$$\begin{aligned}
&= \left(\frac{1}{4}\right)^{n+1} n^{n-1} (2-n+2a+an)^{n+1} (1-a)^{n+1} 2e^{-i\theta} \det \begin{pmatrix} a^n & -(1-a)^2 & -a(1-a)^2 & \cdots & -a^{n-1}(n-2a-an) \\ a^{n-1} & (2-a) & -(1-a)^2 & \cdots & -a^{n-2}(n-2a-an) \\ a^{n-2} & -1 & (2-a) & \cdots & a^{n-3}(n-2a-an) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & 0 & \cdots & (n+2) \end{pmatrix} \\
&= \left(\frac{1}{4}\right)^{n+1} n^{n-1} (2-n+2a+an)^{n+1} (1-a)^{n+1} 2e^{-i\theta} \det \begin{pmatrix} a^n & -(1-a)^2 & -a(1-a)^2 & \cdots & -a^{n-1}(n-2a-an) \\ 0 & \frac{1}{a} & 0 & \cdots & 0 \\ 0 & \frac{1-2a}{a^2} & \frac{1}{a} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \frac{(1-a)^2}{a^n} & \frac{(1-a)^2}{a^{n-1}} & \cdots & \frac{n}{a} \end{pmatrix} \\
&= \left(\frac{1}{4}\right)^{n+1} n^n (2-n+2a+an)^{n+1} (1-a)^{n+1} 2e^{-i\theta}. \tag{10}
\end{aligned}$$

Also, $\det[G_1 E_2 \cdots F_n G_{n+1}]$

$$= (-1)^n \left(\frac{1}{4}\right)^{n+1} n^n (2-n+2a+an)^{n+1} (1-a)^{n+1} (n-2a-an) \det \begin{pmatrix} a^{n-1} & (2-a) & -(1-a)^2 & \cdots & -a^{n-3}(1-a)^2 \\ a^{n-2} & -1 & (2-a) & \cdots & -a^{n-4}(1-a)^2 \\ a^{n-3} & 0 & -1 & \cdots & -a^{n-5}(1-a)^2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & 0 & \cdots & -1 \end{pmatrix}.$$

Now

$$\begin{aligned}
&\det \begin{pmatrix} a^{n-1} & (2-a) & -(1-a)^2 & \cdots & -a^{n-3}(1-a)^2 \\ a^{n-2} & -1 & (2-a) & \cdots & -a^{n-4}(1-a)^2 \\ a^{n-3} & 0 & -1 & \cdots & -a^{n-5}(1-a)^2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & 0 & \cdots & -1 \end{pmatrix} = \\
&a^{n-1} \det \begin{pmatrix} -1 & (2-a) & -(1-a)^2 & \cdots & -a^{n-4}(1-a)^2 \\ 0 & -1 & (2-a) & \cdots & -a^{n-5}(1-a)^2 \\ 0 & 0 & -1 & \cdots & -a^{n-6}(1-a)^2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & -1 \end{pmatrix} - a^{n-2} \det \begin{pmatrix} (2-a) & -(1-a)^2 & -a(1-a)^2 & \cdots & -a^{n-3}(1-a)^2 \\ 0 & -1 & (2-a) & \cdots & -a^{n-5}(1-a)^2 \\ 0 & 0 & -1 & \cdots & -a^{n-6}(1-a)^2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & -1 \end{pmatrix} + \\
&a^{n-3} \det \begin{pmatrix} (2-a) & -(1-a)^2 & -a(1-a)^2 & \cdots & -a^{n-3}(1-a)^2 \\ -1 & (2-a) & -(1-a)^2 & \cdots & -a^{n-4}(1-a)^2 \\ 0 & -1 & (2-a) & \cdots & -a^{n-6}(1-a)^2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & -1 \end{pmatrix} - a^{n-4} \det \begin{pmatrix} (2-a) & -(1-a)^2 & -a(1-a)^2 & \cdots & -a^{n-3}(1-a)^2 \\ -1 & (2-a) & -(1-a)^2 & \cdots & -a^{n-4}(1-a)^2 \\ 0 & -1 & (2-a) & \cdots & -a^{n-5}(1-a)^2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & -1 \end{pmatrix} + \dots
\end{aligned}$$

$$\begin{aligned}
&= a^{n-1}(-1)^{n-1} + a^{n-2}(-1)^{n-1}(2-a) + a^{n-3}(-1)^{n-1}(3-2a) + a^{n-4}(-1)^{n-1}(4-3a) \cdots + \\
&\quad a(-1)^{n-1}[(n-1) - (n-2)a] + (-1)^{n-1}[n - (n-1)a] \\
&= (-1)^{n-1}n. \text{ Therefore,}
\end{aligned}$$

$$\det[G_1 E_2 \cdots F_n G_{n+1}] = -\left(\frac{1}{4}\right)^{n+1} n^n (2-n+2a+an)^{n+1} (1-a)^{n+1} (n-2a-an)n. \quad (11)$$

In addition

$$\det[F_1 E_2 \cdots G_n F_{n+1}]$$

$$= \det \begin{pmatrix} 1 - \bar{a}_0 a_0 - \bar{a}_1 (aa_0 + a_1) & -\bar{a}_0 (aa_0 + a_1) - \bar{a}_1 a (aa_0 + a_1) & \cdots & \bar{a}_1 (\bar{a}_1 a_0 + a) & -\bar{a}_0 a^{n-1} (aa_0 + a_1) \\ -a - \bar{a}_1 a_0 & 1 - \bar{a}_0 a_0 - \bar{a}_1 (aa_0 + a_1) & \cdots & 0 & -\bar{a}_0 a^{n-2} (aa_0 + a_1) \\ 0 & -a - \bar{a}_1 a_0 & \cdots & 0 & -\bar{a}_0 a^{n-3} (aa_0 + a_1) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 - \bar{a}_0 a_0 \end{pmatrix}$$

$$= \left(\frac{1}{4}\right)^{n+1} n^n (2-n+2a+an)^{n+1} (1-a)^{n+1} \left[\frac{e^{i\theta}}{2} (n+2)(2-n+an) \right]. \quad (12)$$

Also, $\det[G_1 E_2 \cdots G_n F_{n+1}]$

$$\begin{aligned}
&= \det \begin{pmatrix} a^n (aa_0 + a_1) & -\bar{a}_0 (aa_0 + a_1) - \bar{a}_1 a (aa_0 + a_1) & \cdots & \bar{a}_1 (\bar{a}_1 a_0 + a) & -\bar{a}_0 a^{n-1} (aa_0 + a_1) \\ a^{n-1} (aa_0 + a_1) & 1 - \bar{a}_0 a_0 - \bar{a}_1 (aa_0 + a_1) & \cdots & 0 & -\bar{a}_0 a^{n-2} (aa_0 + a_1) \\ a^{n-2} (aa_0 + a_1) & -a - \bar{a}_1 a_0 & \cdots & 0 & -\bar{a}_0 a^{n-3} (aa_0 + a_1) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ (aa_0 + a_1) & 0 & 0 & \cdots & 1 - \bar{a}_0 a_0 \end{pmatrix} \\
&= \left(\frac{1}{4}\right)^{n+1} n^{n-1} (2-n+2a+an)^{n+1} (1-a)^{n+1} 2e^{-i\theta} (-1)^{n+1} \bar{a}_1 \det \begin{pmatrix} a^{n-1} & (2-a) & -(1-a)^2 & \cdots & -a^{n-2}(n-2a-an) \\ a^{n-2} & -1 & (2-a) & \cdots & -a^{n-3}(n-2a-an) \\ a^{n-3} & 0 & -1 & \cdots & -a^{n-4}(n-2a-an) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & 0 & \cdots & n+2 \end{pmatrix}.
\end{aligned}$$

Now

$$\begin{aligned}
&\det \begin{pmatrix} a^{n-1} & (2-a) & -(1-a)^2 & \cdots & -a^{n-2}(n-2a-an) \\ a^{n-2} & -1 & (2-a) & \cdots & -a^{n-3}(n-2a-an) \\ a^{n-3} & 0 & -1 & \cdots & -a^{n-4}(n-2a-an) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & 0 & \cdots & n+2 \end{pmatrix} = \\
&a^{n-1} \det \begin{pmatrix} -1 & (2-a) & -(1-a)^2 & \cdots & -a^{n-3}(n-2a-an) \\ 0 & -1 & (2-a) & \cdots & -a^{n-4}(n-2a-an) \\ 0 & 0 & -1 & \cdots & -a^{n-5}(n-2a-an) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & n+2 \end{pmatrix} - a^{n-2} \det \begin{pmatrix} (2-a) & -(1-a)^2 & -a(1-a)^2 & \cdots & -a^{n-2}(n-2a-an) \\ 0 & -1 & (2-a) & \cdots & -a^{n-4}(n-2a-an) \\ 0 & 0 & -1 & \cdots & -a^{n-5}(n-2a-an) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & n+2 \end{pmatrix} + \\
&a^{n-3} \det \begin{pmatrix} (2-a) & -(1-a)^2 & -a(1-a)^2 & \cdots & -a^{n-2}(n-2a-an) \\ -1 & (2-a) & -(1-a)^2 & \cdots & -a^{n-3}(n-2a-an) \\ 0 & 0 & -1 & \cdots & -a^{n-5}(n-2a-an) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & n+2 \end{pmatrix} + \cdots (-1)^{n+1} \det \begin{pmatrix} (2-a) & -(1-a)^2 & -a(1-a)^2 & \cdots & -a^{n-2}(n-2a-an) \\ -1 & (2-a) & -(1-a)^2 & \cdots & -a^{n-3}(n-2a-an) \\ 0 & -1 & (2-a) & \cdots & -a^{n-4}(n-2a-an) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & -(n-2a-an) \end{pmatrix}
\end{aligned}$$

$$\begin{aligned}
&= a^{n-1}(-1)^{n-2}(n+2) + a^{n-2}(-1)^{n-2}(n+2)(2-a) + a^{n-3}(-1)^{n-2}(n+2)(3-2a) + \cdots + (-1)^{n-2}(n-1)[n - (n+2)a] \\
&= (-1)^{n-2}n(n-1). \text{ Hence,}
\end{aligned}$$

$$\det[G_1 E_2 \cdots G_n F_{n+1}] = - \left(\frac{1}{4}\right)^{n+1} n^n (2-n+2a+an)^{n+1} (1-a)^{n+1} (2+an-n)(n-1). \quad (13)$$

Finally,

$$\begin{aligned}
\det[F_1 E_2 \cdots G_n G_{n+1}] &= \det[H_1 E_2 \cdots G_n F_{n+1}] = \det[H_1 E_2 \cdots G_n G_{n+1}] = \\
& \det[G_1 E_2 \cdots G_n G_{n+1}] = \det[H_1 E_2 \cdots F_n G_{n+1}] = 0 \quad (14)
\end{aligned}$$

Using equations (7)-(14), we get

$$\begin{aligned}
M_{n+1} &= \det[\overline{B}_{n+1}^T - A_{n+1} B_{n+1}^{-1} \overline{A}_{n+1}^T] = \left(\frac{1}{4}\right)^{n+1} n^n (1-a)^{n+1} (2-n+2a+an)^{n+1} [(3n+2) + \\
& \frac{e^{i\theta}}{2}n(n-2a-an) - 3an + 2e^{-i\theta} - n(n-2a-an) + \frac{e^{i\theta}}{2}(n+2)(2+an-n) - (n-1)(2+an-n)] \\
&= \left(\frac{1}{4}\right)^{n+1} n^n (1-a)^{n+1} (2-n+2a+an)^{n+1} (4+4\cos\theta).
\end{aligned}$$

Therefore $M_{n+1} > 0$, if $\theta \neq (2m+1)\pi, m \in \mathbb{N}$.

When $\theta = (2m+1)\pi, m \in \mathbb{N}$, then $p(z) = z^{n+1} - az^n + \frac{1}{2}(n-2-an)z - \frac{1}{2}(n-2a-an)$. As $z=1$ is a zero of $p(z)$, therefore we can write

$$\begin{aligned}
p(z) &= (z-1)[z^n + (1-a)z^{n-1} + (1-a)z^{n-2} + \cdots + (1-a)z + \frac{1}{2}(n-2a-an)] \\
&= (z-1)q(z).
\end{aligned}$$

It suffices to show that zeros of $q(z)$ lie inside $|z|=1$. Since $|\frac{1}{2}(n-2a-an)| < 1$ whenever $a \in (\frac{n-2}{n+2}, 1)$, by applying Lemma A on $q(z)$ (by comparing it with $t(z)$), we get

$$\begin{aligned}
q_1(z) &= \frac{\overline{a}_n q(z) - a_0 q^*(z)}{z} \\
&= (1-a)[1 - (\frac{1}{2}(n-2a-an))] \left\{ (1 + \frac{n}{2})z^{n-1} + z^{n-2} + \cdots + z + 1 \right\}.
\end{aligned}$$

By Lemma A, the number of zeros of $q_1(z)$ inside the unit circle is one less than the number of zeros of $q(z)$ inside the unit circle. Let $p_1(z) = (1 + \frac{n}{2})z^{n-1} + z^{n-2} + \cdots + z + 1$. Again $1 < |1 + \frac{n}{2}|$, therefore

$$\begin{aligned}
q_2(z) &= \frac{\overline{a}_{n-1} p_1(z) - a_0 p_1^*(z)}{z} \\
&= \frac{n}{2} [(2 + \frac{n}{2})z^{n-2} + z^{n-3} + \cdots + z + 1].
\end{aligned}$$

Again the number of zeros of $q_2(z)$ inside the unit circle is two less than the number of zeros of $q(z)$ inside the unit circle. Continuing in this manner we derive that

$$q_k(z) = \left[(k-2) + \frac{n}{2} \right] \left\{ \left(k + \frac{n}{2} \right) z^{n-k} + z^{n-(k+1)} + \dots + z^{n-(n-1)} + 1 \right\}, \quad k = 2, 3, \dots, n-1.$$

In particular for $k = n-1$

$$q_{n-1}(z) = \left((n-3) + \frac{n}{2} \right) \left\{ \left((n-1) + \frac{n}{2} \right) z + 1 \right\},$$

which has $(n-1)$ less number of zeros inside the unit circle than the number of zeros of $q(z)$ inside the unit circle. But the zero of q_{n-1} is $-\frac{2}{3n-2}$ which lies inside the unit circle $|z| = 1$ for $n \geq 2$. Consequently all zeros of $q(z)$ lie inside $|z| = 1$ and the proof of our theorem is now complete. \square

Remark 2.3. If we set $a = 0$, then n is restricted to $n = 1$ or $n = 2$ and we get the same result as Theorem B stated in Section 1.

Next we give an example showing that for a given value of n , if we go beyond the range of real number a as specified in Theorem 2.2, then convolution no longer remains locally univalent and sense preserving. If we take $n = 1$, then the range of real constant a comes out to be $[-\frac{1}{3}, 1)$.

Example 2.4. If we take $n = 1$, $a = -0.34 < -\frac{1}{3}$ and $\theta = \pi$, in (5) we get

$$\begin{aligned} \tilde{\omega}_1(z) &= -z \left[\frac{z^2 + 0.01z - 1.01}{1 + 0.01z - 1.01z^2} \right] \\ &= -zR(z). \end{aligned}$$

We prove that there exists some point z_0 in E such that $|\tilde{\omega}_1(z_0)| > 1$. Assume that this is not true. It is easy to see that for each α , $|R(e^{i\alpha})| = 1$ and $R(z)\overline{R(\frac{1}{\bar{z}})} = 1$. So the function $R(z)$ preserves the symmetry about the unit circle and maps the closed disk $|z| \leq 1$ onto itself. Therefore, $R(z)$ can be written as a Blaschke product of order two. However, the product of the moduli of zeros of R in the unit disk is 1.01, which is a contradiction.

The image of E under $f_a * f_1$ for $a = -0.34$ is shown (using the applet *Complex Tool* (see [4])) in Figure 1. Figure 2 is a zoomed version of Figure 1 showing that the images of two outer most concentric circles in E are intersecting and so $f_a * f_1$ is not univalent.

In the next result we find the condition under which the convolution, $f_a * f_b \in S_H$ and is CHD.

Theorem 2.5. *If $f_b = h + \bar{g} \in K_H$ is given by $h + g = \frac{z}{1-z}$ with dilatation $\omega(z) = \frac{b-z}{1-bz}$ ($|b| < 1, b \in \mathbb{R}$), then $f_a * f_b \in S_H$ and is CHD for $b \geq -\frac{1+3a}{3+a}$, where f_a is as in Theorem 2.2.*




Figure 1




Figure 2

Proof. If $\tilde{\omega}_1(z)$ is the dilatation of $f_a * f_b$, then in view of Theorem A it is sufficient to prove that $|\tilde{\omega}_1(z)| < 1$, for all $z \in E$. Substituting $\omega(z) = \frac{b-z}{1-bz}$ in (4), we get

$$\begin{aligned} \tilde{\omega}_1(z) &= \left[\frac{2 \left(\frac{b-z}{1-bz} \right) (a-z) \left(1 + \frac{b-z}{1-bz} \right) + \frac{b^2 z - z^2}{(1-bz)^2} (a-1)(1-z)}{2(1-az) \left(1 + \frac{b-z}{1-bz} \right) + \frac{b^2 z - z^2}{(1-bz)^2} (a-1)(1-z)} \right] \\ &= \frac{z^2 + \frac{1}{2}(ab - 3a - 3b + 1)z + ab}{abz^2 + \frac{1}{2}(ab - 3a - 3b + 1)z + 1} = \frac{m(z)}{m^*(z)} \end{aligned}$$

$$\begin{aligned} \text{where } m(z) &= z^2 + \frac{1}{2}(ab - 3a - 3b + 1)z + ab \\ &= a_2 z^2 + a_1 z + a_0 \quad \text{and} \\ m^*(z) &= z^2 m\left(\frac{1}{z}\right). \end{aligned}$$

As before, if z_0 is a zero of m then $\frac{1}{z_0}$ is a zero of m^* , and we can write

$$\tilde{\omega}_1(z) = \frac{(z+A)(z+B)}{(1+\bar{A}z)(1+\bar{B}z)}.$$

It suffices to show that either both the zeros $-A, -B$ lie inside unit circle $|z| = 1$ or one of the zeros lies inside $|z| = 1$ and other lies on it. As $|a_0| = |ab| < 1 = |a_2|$, using Lemma A on m , we have

$$\begin{aligned} m_1(z) &= \frac{\bar{a}_2 m(z) - a_0 m^*(z)}{z} \\ &= (1-ab) \left[(1+ab)z + \frac{1}{2}(-3a-3b+1+ab) \right]. \end{aligned}$$

But $z_0 = \frac{3}{2} \left(\frac{a+b}{1+ab} \right) - \frac{1}{2}$ is the zero of m_1 which lies in or on the unit circle $|z| = 1$ if

$b \geq -\frac{(1+3a)}{(3+a)}$. So, either both zeros of m lie inside $|z| = 1$ or at least one zero lies in and other lies on $|z| = 1$. Hence $|\tilde{\omega}_1(z)| < 1$. □

Corollary 2.6. *The convolution $f_a * f_a \in S_H$ and CHD for $a \in [-3 + 2\sqrt{2}, 1)$.*

Acknowledgement: The first author is thankful to Council of Scientific and Industrial Research, New Delhi, for financial support vide (grant no. 09/797/0006/2010 EMR-1).

References

- [1] J. Clunie and T. Shiel-Small, *Harmonic univalent functions*, Ann. Acad. Sci. Fenn. Ser. A. I Math., **9** (1984), 3-25.
- [2] M. Dorff, *Convolutions of planar harmonic convex mappings*, Comp. Var. Theory Appl. **45** (2001), no. 3, 263-271.
- [3] M. Dorff, M. Nowak and M. Woloszkiwicz, *Convolutions of harmonic convex mappings*, Comp. Var. Elliptic Eqn., **57**(2012), no. 5, 489-503.
- [4] M. Dorff and J. Rolf, *Anamorphosis, mapping problems, and harmonic univalent functions. Explorations in complex analysis*, 197-269, Math. Assoc. of America, Washington, DC, 2012 (or see www.maa.org/ebooks/EXCA/applets.html).
- [5] L. Li and S. Ponnusamy, *Solution to an open problem on convolutions of harmonic mappings*, Comp. Var. Elliptic Eqn. (2012), accepted.
- [6] L. Li and S. Ponnusamy, *Convolution of slanted right half-plane harmonic mappings*, Analysis (2012) accepted (or see arXiv:1206.4364).
- [7] Q.I. Rahman and G. Schmeisser, *Analytic Theory of Polynomials*, London Mathematical Society Monographs New Series, Vol. 26, Oxford University Press, Oxford, 2002.