

L_∞ -algebras of local observables from higher prequantum bundles

Domenico Fiorenza
 Department of Mathematics
 Sapienza Università di Roma
 Rome, Italy
 fiorenza@mat.uniroma1.it

Christopher L. Rogers
 Mathematics Institute
 Georg-August Universität Göttingen
 Göttingen, Germany
 crogers@uni-math.gwdg.de

Urs Schreiber
 Mathematics Institute
 Radboud Universiteit Nijmegen
 Nijmegen, The Netherlands
 urs.schreiber@gmail.com

March 11, 2019

Abstract

The second author has defined a class of L_∞ -algebras that are naturally associated to manifolds equipped with a closed higher-degree differential form, which reduce to the underlying Lie algebra of the usual Poisson algebra in the case of symplectic manifolds. Here we show how these L_∞ -algebras of local observables can be interpreted as the infinitesimal autoequivalences of higher prequantum bundles covering higher Hamiltonian symplectomorphisms. More precisely, we define a dg Lie algebra of such infinitesimal autoequivalences and exhibit an explicit homotopy equivalence between it and the L_∞ -algebra of local observables. By truncation of the connection data for the prequantum bundle, this produces higher analogues of the Lie algebra of sections of the Atiyah Lie algebroid and the Lie 2-algebra of sections of the Courant Lie 2-algebroid. Finally we exhibit the L_∞ -cocycle that realizes the L_∞ -algebras of local observables as a Kostant-Souriau-type L_∞ -extension of the Hamiltonian vector fields. When restricted along a Lie algebra action, this yields L_∞ -analogs of the Heisenberg Lie algebras and their classifying cocycles. In particular, this recovers the string Lie 2-algebra of a semisimple Lie algebra as well as the 3-cocycle that classifies it.

Contents

1	Introduction	2
2	Higher prequantum geometry over smooth manifolds	4
2.1	n -Plectic manifolds and their Hamiltonian vector fields	4
2.2	Prequantization of (pre-) n -plectic manifolds	5
3	The L_∞-algebra of local observables and its Kostant-Souriau L_∞-cocycle	7
3.1	L_∞ -algebras, extensions and cocycles	7
3.2	The L_∞ -algebra of local observables	12
3.3	The Kostant-Souriau L_∞ -algebra cocycle	13
3.4	The Kostant-Souriau L_∞ -extension	15
3.5	The Heisenberg L_∞ -extension	16

4	The dg-Lie algebra of infinitesimal quantomorphisms	17
4.1	Quantomorphism n -groups	17
4.2	Infinitesimal quantomorphisms as a strict model for the L_∞ -algebra of observables	19
5	Inclusion into Atiyah and Courant L_∞-algebras	20
5.1	The $n = 1$ case	22
5.2	The $n = 2$ case	23
A	An explicit weak equivalence between $L_\infty(X, \omega)$ and $\mathrm{dgLie}_{\mathrm{Qu}}(X, \bar{A})$	27

1 Introduction

Geometric objects, such as manifolds, orbifolds, or stacks, equipped with a closed differential form play important roles in many areas of current mathematical interest. The archetypical example are closed 2-forms in (pre-)symplectic geometry. Higher degree closed forms play crucial roles, for example, in covariant quantum field theory, in Hitchin’s generalized complex/Riemannian geometry, and in differential cohomology. It is becoming clear that it is advantageous to consider these forms, in one way or another, as higher degree generalizations of symplectic structures.

In all of these applications, there is a particular focus on integral closed forms. This is because such forms correspond to the curvatures of higher geometric structures known as $U(1)$ - n -bundles with connection (or $U(1)$ - $(n - 1)$ -bundle gerbes with connection). Here we refer to these as *higher prequantum bundles*, in analogy with the role that $U(1)$ -principal connections play in the geometric prequantization of symplectic manifolds [16, 28]. (A modern review can be in [6].) In the companion article [10] we present general aspects of such higher geometric prequantum structures; here we work out details of the general theory specialized to the higher differential geometry over smooth manifolds. In particular, we use homotopy Lie theory to study the infinitesimal autoequivalences of higher prequantum bundles covering infinitesimal diffeomorphisms of the base manifold i.e. the infinitesimal quantomorphisms.

It is well-known that every pre-symplectic manifold induces a Lie algebra of Hamiltonian functions whose bracket is the Poisson bracket given by the closed 2-form. When the manifold is equipped with a prequantum bundle, this Lie algebra is isomorphic to the Lie algebra of infinitesimal autoequivalences of that structure i.e. those vector fields on the bundle whose flow preserves the underlying bundle and its connection under pullback. These are also called prequantum operators. More generally, manifolds equipped with higher degree forms also have Hamiltonian vector fields, which form a Lie algebra just like in symplectic geometry. The differential form induces a bilinear skew-symmetric bracket not on functions, but on higher degree differential forms. However, this bracket fails to satisfy the Jacobi identity. The observation made in [22] was that, for the case of non-degenerate forms, this failure is controlled by coherent homotopy. Hence, instead of being a problem, the lack of a genuine Lie bracket indicates the presence of a natural, but higher (homotopy-theoretic) structure. More precisely, the higher Poisson bracket gives rise to a strong-homotopy Lie algebra or L_∞ -algebra. The construction in [22] extends trivially to the case of degenerate forms, and we call the resulting algebra the ‘ L_∞ -algebra of local observables’. In this paper, we illuminate its conceptual role further.

Summary of results. We identify the higher Kostant-Souriau L_∞ -algebra cocycle that classifies the L_∞ -algebra of local observables as an extension of the Hamiltonian vector fields (theorem 3.4.1) and show how this result immediately gives a construction of ‘higher Heisenberg L_∞ -algebras’ (section 3.5). As an example, we obtain a direct re-derivation (example 3.5.4) of the $\mathbf{string}_{\mathfrak{g}}$ -Lie 2-algebra as the Heisenberg Lie 2-algebra of a compact simple Lie group G [3].

We briefly recall the construction of the higher prequantum automorphism group of a higher prequantum bundle, which is described with more detail in [10]. We construct a dg Lie algebra (def. 4.2.1) which can be thought of as modeling the “infinitesimal elements” of this higher automorphism group in terms of the Čech-Deligne cocycle for the prequantum bundle. (Similar dg Lie models for the “infinitesimal symmetries” of a $U(1)$ -bundle gerbe were constructed by Collier [9].)

We prove explicitly that our dg Lie algebra of infinitesimal quantomorphisms is equivalent, as an L_∞ -algebra, to the L_∞ -algebra of local observables of the corresponding pre- n -plectic form (theorem 4.2.2).

Finally, we show that this construction induces an inclusion of the L_∞ -algebra of local observables into higher Courant and higher Atiyah Lie algebras (section 5).

Remark 1.0.1. All of the constructions and results that we discuss here apply to the general context of pre- n -plectic manifolds, i.e. manifolds equipped with a closed $(n + 1)$ -form. Non-degeneracy conditions on the differential form do not play a role. Nevertheless, our formalism allows us to restrict to the case of non-degenerate forms, and it may be interesting to do so in specific applications. This is analogous to the well-known fact that non-degeneracy is not needed to prequantize a symplectic manifold. Indeed, one can proceed even further in this case; the full geometric quantization of pre-symplectic manifolds is a well-defined and interesting endeavour in its own right (e.g. [7]).

Acknowledgements. We thank Yael Frégier and Marco Zambon for sharing a preliminary draft [13] of their work (joint with C.L.R.) which inspired some of the ideas developed here, and Ruggero Bandiera, Christian Blohmann and Marco Manetti for the invaluable inspiring conversations they had with D.F. on the homotopy fibers of L_∞ -morphisms and on higher symplectic geometry.

C.L.R. acknowledges support from the German Research Foundation (Deutsche Forschungsgemeinschaft (DFG)) through the Institutional Strategy of the University of Göttingen

Notations and conventions.

Notations for Cartan calculus

The Schouten bracket of two decomposable multivector fields $u_1 \wedge \cdots \wedge u_m, v_1 \wedge \cdots \wedge v_n \in \bigwedge^\bullet \mathfrak{X}(X)$ is

$$[u_1 \wedge \cdots \wedge u_m, v_1 \wedge \cdots \wedge v_n] = \sum_{i=1}^m \sum_{j=1}^n (-1)^{i+j} [u_i, v_j] \wedge u_1 \wedge \cdots \wedge \hat{u}_i \wedge \cdots \wedge u_m \wedge v_1 \wedge \cdots \wedge \hat{v}_j \wedge \cdots \wedge v_n, \quad (1.0.1)$$

where $[u_i, v_j]$ is the usual Lie bracket of vector fields.

Given a form $\alpha \in \Omega^\bullet(X)$, the **interior product** of a decomposable multivector field $v_1 \wedge \cdots \wedge v_n$ with α is

$$\iota_{v_1 \wedge \cdots \wedge v_n} \alpha = \iota_{v_n} \cdots \iota_{v_1} \alpha,$$

where $\iota_{v_i} \alpha$ is the usual interior product of vector fields and differential forms. The interior product of an arbitrary multivector field is obtained by extending the above formula by $C^\infty(X; \mathbb{R})$ -linearity.

The **Lie derivative** \mathcal{L}_v of a differential form along a multivector field $v \in \bigwedge^\bullet \mathfrak{X}(X)$ is defined via the graded commutator of d and $\iota(v)$:

$$\mathcal{L}_v \alpha = d \iota_v \alpha - (-1)^{|v|} \iota_v d \alpha, \quad (1.0.2)$$

where $\iota(v)$ is considered as a degree $-|v|$ operator.

The last identity we will need involving multivector fields is for the graded commutator of the Lie derivative and the interior product. Given $u, v \in \bigwedge^\bullet \mathfrak{X}(X)$, it follows from [12, Proposition A3] that

$$\iota_{[u,v]} \alpha = (-1)^{(|u|-1)|v|} \mathcal{L}_u \iota_v \alpha - \iota_v \mathcal{L}_u \alpha. \quad (1.0.3)$$

Conventions on chain and cochain complexes

We will mostly work with chain complexes and homological degree conventions. The differential of a chain complex (A_\bullet, d) will have degree -1:

$$\cdots \rightarrow A_{n+1} \xrightarrow{d} A_n \xrightarrow{d} A_{n-1} \rightarrow \cdots$$

The shift functor $A_\bullet \mapsto A[1]_\bullet$ will act by shifting on the left, i.e., $A[1]_k = A_{k-1}$. In particular, if V is a vector space, seen as a chain complex concentrated in degree zero, $V[n]$ will be the chain complex consisting of V concentrated in degree n .

A cochain complex (A^\bullet, d) will have a differential of degree $+1$,

$$\dots \rightarrow A^{n-1} \xrightarrow{d} A^n \xrightarrow{d} A^{n+1} \rightarrow \dots,$$

and will be identified with a chain complex (with the same differential) by the rule $A_k = A^{-k}$. In particular chain complexes concentrated in non-negative degree will correspond to cochain complexes concentrated in nonpositive degree, and vice versa. Also on cochain complexes the shift functor $A^\bullet \mapsto A[1]^\bullet$ will act by shifting on the left, i.e., we have $A[1]^k = A^{k+1}$.

Conventions on stacks and higher stacks

While this article focuses on homotopy Lie theory, we do mention at some points the corresponding constructions in higher smooth stacks, according to [10]. A detailed overview of this formalism is given in Sec. 3.1 in [11].

Smooth stacks are taken to be stacks over the category of all smooth manifolds equipped with its standard Grothendieck topology of good open covers. Equivalently but more conveniently this are stacks over just the subcategory CartSp of Cartesian spaces $\{\mathbb{R}^n\}_{n \in \mathbb{N}}$ (or equivalently of open n -balls), regarded as smooth manifolds. A higher smooth stack may always be presented as a Kan-complex valued functor on $\text{CartSp}^{\text{op}}$ and the homotopy theory \mathbf{H} of smooth stacks is given by the category of such functors with stalkwise homotopy equivalences of Kan complexes universally turned into actual homotopy equivalences:

$$\mathbf{H} := L_{\text{the}} \text{Func}(\text{CartSp}^{\text{op}}, \text{KanCplx}).$$

In the applications of the present article all examples of such objects are either given by sheaves of chain complexes A_\bullet of abelian groups in non-negative degrees under the Dold-Kan correspondence

$$\text{DK} : \text{Ch}_{\geq \bullet}(\text{Ab}) \xrightarrow{\simeq} \text{AbGrp}^{\Delta^{\text{op}}} \xrightarrow{\text{forget}} \text{KanCplx}$$

or are the Čech nerve $\check{C}(\mathcal{U})$ of an open cover $\mathcal{U} = \{U_i \rightarrow X\}_i$ of a smooth manifold X . If \mathcal{U} is a good cover and if A_\bullet is CartSp -acyclic (which it is in all the examples we consider), then the function complex $\mathbf{H}(X, A) \simeq \text{Func}(\check{C}(\mathcal{U}), \text{DK}(A_\bullet))$ is the traditional cocycle complex of Čech hypercohomology of X with coefficients in A_\bullet .

2 Higher prequantum geometry over smooth manifolds

We briefly review here the basic notions of higher prequantum geometry over smooth manifolds which will be used throughout the article. First in 2.1 we recall the notion of pre- n -plectic manifolds and their Hamiltonian vector fields and then in 2.2 their pre-quantization by Čech-Deligne cocycles.

2.1 n -Plectic manifolds and their Hamiltonian vector fields

In [2] the following terminology has been introduced.

Definition 2.1.1. A *pre- n -plectic manifold* (X, ω) is a smooth manifold X equipped with a closed $(n+1)$ -form $\omega \in \Omega_{\text{cl}}^{n+1}(X)$. If the contraction map $\hat{\omega} : TX \rightarrow \Lambda^n T^*X$ is injective, then ω is called *non-degenerate* or *n -plectic* and (X, ω) is called an *n -plectic manifold*.

Example 2.1.2. For $n = 1$ an n -plectic manifold is equivalently an ordinary symplectic manifold.

Example 2.1.3. Let G be a compact connected simple Lie group. Equipped with its canonical left invariant differential 3-form $\omega := \langle -, [-, -] \rangle$ this is a 2-plectic manifold.

Definition 2.1.4. Let (X, ω) be a pre- n -plectic manifold. If a vector field v and an $(n-1)$ -form H are related by

$$\iota_v \omega + dH = 0$$

then we say that v is a Hamiltonian field for H and that H is a Hamiltonian form for v .

Definition 2.1.5. We denote by

$$\text{Ham}^{n-1}(X) \subseteq \mathfrak{X}(X) \oplus \Omega^{n-1}(X)$$

the subspace of pairs (v, H) such that $\iota_v \omega + dH = 0$. We call this the space of *Hamiltonian pairs*. The image $\mathfrak{X}_{\text{Ham}}(X) \subseteq \mathfrak{X}(X)$ of the projection $\text{Ham}^{n-1}(X) \rightarrow \mathfrak{X}(X)$ is called the space of *Hamiltonian vector fields* of (X, ω) .

Remark 2.1.6. Given a pre- n -plectic manifold (X, ω) We have a short exact sequence of vector spaces

$$0 \rightarrow \Omega_{\text{cl}}^{n-1}(X) \rightarrow \text{Ham}^{n-1}(X) \rightarrow \mathfrak{X}_{\text{Ham}}(X) \rightarrow 0,$$

i.e., closed $(n-1)$ -forms are Hamiltonian, with zero Hamiltonian vector field.

Remark 2.1.7. It is immediate from the definition that Hamilton vector fields preserve the pre- n -plectic form ω , i.e., $\mathcal{L}_v \omega = 0$. Indeed, since ω is closed, we have $\mathcal{L}_v \omega = d\iota_v \omega = -d^2 H_v = 0$. Therefore the integration of a Hamiltonian vector field gives a diffeomorphism of X preserving the pre- n -plectic form: a *Hamiltonian n -plectomorphism*.

Lemma 2.1.8. *The subspace $\mathfrak{X}_{\text{Ham}}(X)$ is a Lie subalgebra of $\mathfrak{X}(X)$.*

Proof. Let v_1 and v_2 be Hamiltonian vector fields, and H_1, H_2 their respective Hamiltonian forms. By $\mathcal{L}_{v_1} \omega = 0$ and by the Cartan formulas, we get

$$\iota_{[v_1, v_2]} \omega = [\mathcal{L}_{v_1}, \iota_{v_2}] \omega = -\mathcal{L}_{v_1} dH_2 = -d\mathcal{L}_{v_1} H_2 = d\iota_{v_1} \iota_{v_2} \omega,$$

i.e., $[v_1, v_2]$ is Hamiltonian, with Hamiltonian $\iota_{v_1} \wedge \iota_{v_2} \omega$. □

Remark 2.1.9. Hamiltonian vector fields on a pre- n -plectic manifold (X, ω) are by definition those vector fields v such that $\iota_v \omega$ is exact. One may relax this condition and consider *symplectic vector fields* instead, i.e., those vector fields v such that $\iota_v \omega$ is closed. Then the arguments in Remark 2.1.7 and in Lemma 2.1.8 show that symplectic vector fields form a Lie subalgebra $\mathfrak{X}_{\text{symp}}(X)$ of $\mathfrak{X}(X)$ and that $\mathfrak{X}_{\text{symp}}(X) \subseteq \mathfrak{X}_{\text{Ham}}(X)$ is a Lie ideal.

2.2 Prequantization of (pre-) n -plectic manifolds

The traditional notion of prequantization of a presymplectic manifold (X, ω) is equivalently a lift of the presymplectic form, regarded as a de Rham 2-cocycle, to a degree 2 cocycle in *ordinary differential cohomology* (for instance [6]). Equivalently, this is a lift of ω to a $U(1)$ -principal connection ∇ on X with curvature $F_\nabla = \omega$. Accordingly, the prequantization of a pre- n -plectic manifold is naturally defined to be a lift of ω regarded as a degree $(n+1)$ cocycle in de Rham cohomology to a cocycle of degree $(n+1)$ in ordinary differential cohomology.

Definition 2.2.1. For X a smooth manifold and $\mathcal{U} = \{U_i \rightarrow X\}$ an open cover, we write $(\text{Tot}(\mathcal{U}, \Omega), d_{\text{Tot}})$ for the corresponding Čech-de Rham total complex, i.e. the cochain complex whose underlying graded vector space is

$$\text{Tot}^n(\mathcal{U}, \Omega) = \bigoplus_{i+j=n} \check{C}^i(\mathcal{U}, \Omega^j)$$

and whose differential is given on elements $\bar{\theta} = \sum_{i=0}^n \theta^{n-i} \in \text{Tot}^n(\mathcal{U}, \Omega)$ with $\theta^{n-i} \in \check{C}^i(\mathcal{U}, \Omega^{n-i})$ by

$$d_{\text{Tot}} \theta^{n-i} = \delta \theta^{n-i} + (-1)^i d \theta^{n-i}.$$

Definition 2.2.2. The cochain complex of sheaves

$$C^\infty(-; U(1)) \xrightarrow{d\log} \Omega^1(-) \xrightarrow{d} \Omega^2(-) \xrightarrow{d} \dots \xrightarrow{d} \Omega^n(-) \xrightarrow{d} \Omega^{n+1}(-) \rightarrow \dots,$$

with $C^\infty(-; U(1))$ in degree zero, will be called the *Deligne complex* and will be denoted by the symbol $\underline{U}(1)_{\text{Del}}$. Its truncation

$$C^\infty(-; U(1)) \xrightarrow{d\log} \Omega^1(-) \xrightarrow{d} \Omega^2(-) \xrightarrow{d} \dots \xrightarrow{d} \Omega^n(-) \rightarrow 0 \rightarrow 0 \rightarrow \dots$$

will be denoted by $\underline{U}(1)_{\text{Del}}^{\leq n}$.

It follows from the above definition that a degree n Čech-Deligne cocycle in $\underline{U}(1)_{\text{Del}}^{\leq n}$ is

$$\bar{A} = \sum_{i=0}^n A^{n-i}, \quad A^{n-i} \in \check{C}^i(\mathcal{U}, \Omega^{n-i}), \quad A^0 \in \check{C}^n(\mathcal{U}, \underline{U}(1))$$

satisfying

$$\begin{aligned} \delta A^{n-i} &= (-1)^i dA^{n-i-1}, \quad i = 0, \dots, n-2 \\ \delta A^1 &= (-1)^{n-1} d\log A^0 \\ \delta A^0 &= 1 \end{aligned} \tag{2.2.1}$$

Definition 2.2.3. We denote by $\mathbf{B}^n U(1)_{\text{conn}}$ the n -stack of principal $U(1)$ - n -bundles (or $(n-1)$ -bundle gerbes) with connection. This is the n -stack presented via applying the Dold-Kan construction to the presheaf $\underline{U}(1)_{\text{Del}}^{\leq n}[n]$, regarded as a presheaf of chain complexes concentrated in non-negative degree. The commutative diagram

$$\begin{array}{ccccccc} C^\infty(-; U(1)) & \xrightarrow{d\log} & \Omega^1(-) & \xrightarrow{d} & \dots & \xrightarrow{d} & \Omega^{n-1}(-) & \xrightarrow{d} & \Omega^n(-) \\ \downarrow & & \downarrow & & & & \downarrow & & \downarrow d \\ 0 & \longrightarrow & 0 & \longrightarrow & \dots & \longrightarrow & 0 & \longrightarrow & \Omega^{n+1}(-)_{\text{cl}} \end{array}$$

presents the morphism of stacks

$$F : \mathbf{B}^n U(1)_{\text{conn}} \rightarrow \Omega^{n+1}(-)_{\text{cl}}$$

which maps a principal $U(1)$ - n -bundle with connection to its *curvature* $(n+1)$ -form.

Definition 2.2.4. Let (X, ω) be a pre- n -plectic manifold. A *prequantization* of (X, ω) is a lift

$$\begin{array}{ccc} & \mathbf{B}^n U(1)_{\text{conn}} & \\ \nabla \nearrow & & \downarrow F \\ X & \xrightarrow{\omega} & \Omega^{n+1}(-)_{\text{cl}}. \end{array}$$

We call the triple (X, ω, ∇) a prequantized pre- n -plectic manifold.

Local data for a prequantization (X, ω, ∇) are conveniently expressed in terms of the Čech-Deligne total complex. The following result is a simple exercise in the definitions.

Proposition 2.2.5. *Let X be a smooth manifold and let \mathcal{U} be a good cover of X . Then a pre- n -plectic structure on X is the datum of a closed element ω in $\check{C}^0(\mathcal{U}, \underline{U}(1)_{\text{Del}}^{n+1})$ and a prequantization of ω is an element in $\text{Tot}^n(\mathcal{U}, \underline{U}(1)_{\text{Del}})$ such that $d_{\text{Tot}} A = \omega$.*

Remark 2.2.6. It is a well know fact that (X, ω) admits a prequantization if and only if the closed form ω represents an integral class in de Rham cohomology; see, e.g., [6]. Indeed, since the shifted Deligne complex $\underline{U}(1)_{\text{Del}}[n]$ is an acyclic resolution of the cochain complex of sheaves $\mathfrak{b}\mathbf{B}^n U(1)$ consisting of locally constant $U(1)$ -valued functions placed in degree $-n$, we see that a pre- n -plectic structure ω is prequantizable if and only if ω defines the trivial class in the degree n Čech cohomology of X with coefficients in the discrete abelian group $U(1)$. By the short exact sequence of groups

$$0 \rightarrow \mathbb{Z} \rightarrow \mathbb{R} \rightarrow U(1) \rightarrow 1$$

and by the Čech-de Rham isomorphism $H_{\text{dR}}^n(X, \mathbb{R}) \cong \check{H}^n(X, \mathbb{R})$, this is equivalent to requiring that the de Rham class of ω is an integral class.

3 The L_∞ -algebra of local observables and its Kostant-Souriau L_∞ -cocycle

To any pre- n -plectic manifold (X, ω) one can associate an L_∞ -algebra $L_\infty(X, \omega)$, as defined in [13, 22], which we may think of as the higher local observables on (X, ω) . This is an L_∞ -extension of the Lie algebra of Hamiltonian vector fields on (X, ω) by the $(n-1)$ -shifted truncated de Rham complex of X . After a general discussion of some aspects of the homotopy theory of L_∞ -algebra in 3.1, we briefly recall this construction in 3.2 below.

For (V, ω) an ordinary symplectic vector space, we may regard it as a symplectic manifold that is canonically equipped with a V -action by Hamiltonian vector fields, with V regarded as the abelian Lie algebra of constant (left invariant) vector fields on itself. The evaluation map at zero $\iota_{-\wedge-\omega}|_0 : V \times V \rightarrow \mathbb{R}$ of the symplectic form is then a Lie algebra 2-cocycle on V and hence defines an extension of Lie algebras. This is famous as the *Heisenberg Lie algebra* extension and $\iota_{-\wedge-\omega}|_0$ is the *Kostant-Souriau cocycle* that classifies it (see example 3.3.5 below). More generally, for any symplectic manifold, the Kostant-Souriau 2-cocycle classifies the underlying Lie algebra of the Poisson algebra as a central extension of the Hamiltonian vector fields [16, 28]. For symplectic vector spaces, the restriction of the Kostant-Souriau 2-cocycle to the constant Hamiltonian vector fields is precisely the above cocycle. We describe in 3.3 below a further generalization of this to a class of L_∞ -algebra $(n+1)$ -cocycles on Hamiltonian vector fields over pre- n -plectic manifolds. We call these the *higher Kostant-Souriau L_∞ -cocycles*.

In 3.4 we prove that the L_∞ -algebra extension which is classified by the Kostant-Souriau $(n+1)$ -cocycle is indeed again the Poisson-bracket L_∞ -algebra of local observables.

3.1 L_∞ -algebras, extensions and cocycles

Here we first recall some details concerning the homotopy theory of L_∞ -algebras which we shall use throughout the text. We then give an explicit characterization of certain homotopy fibers for L_∞ -morphisms which we use to construct L_∞ -extensions classified by L_∞ -cocycles. Below we follow homological degree conventions, as in [17].

Definition 3.1.1. An L_∞ -algebra \mathfrak{g} is an unbounded chain complex \mathfrak{g}_\bullet (of real vector spaces) equipped with a sequence $\{l_k : L^{\otimes k} \rightarrow L\}_{k=2}^\infty$ of k -ary graded skew symmetric brackets which constitute the components of a degree (-1) codifferential on the co-free graded co-commutative coalgebra on $\mathfrak{g}_\bullet[-1]$. A *morphism* of L_∞ -algebras is a morphism of the corresponding dg-coalgebras (sometimes called an “sh-map” of L_∞ -algebras). In other words, if dgCoCAlg denotes the category of connected dg co-commutative coalgebras, then the category of L_∞ -algebras is the full subcategory

$$\text{CE}: L_\infty\text{Alg} \hookrightarrow \text{dgCoCAlg}$$

consisting of those objects whose underlying graded co-commutative coalgebras (hence forgetting the differential) are co-free on a chain complex. Restricting a morphism $\mathfrak{g} \rightarrow \mathfrak{h}$ of L_∞ -algebras to the subspace of

primitive elements induces a chain map $\mathfrak{g}_\bullet \rightarrow \mathfrak{h}_\bullet$ on the underlying complexes, and hence a forgetful functor from $L_\infty\text{Alg}$ to the category of unbounded chain complexes.

Example 3.1.2 (Lie n -algebras). All examples of L_∞ -algebras \mathfrak{g} given here have their underlying chain complex \mathfrak{g}_\bullet concentrated in non-negative degree. It is common to call an L_∞ -algebra concentrated in degrees 0 through $(n-1)$ a *Lie n -algebra*. A *Lie 1-algebra* is equivalently an ordinary Lie algebra.

Example 3.1.3 (dg Lie algebras). An L_∞ -algebra whose k -ary brackets for $k > 2$ are trivial is equivalently a dg-Lie algebra.

Example 3.1.4 (Abelian L_∞ -algebras). An L_∞ -algebra whose k -ary brackets for $k \geq 2$ are trivial, i.e. a plain chain complex, is called an *abelian L_∞ -algebra*. If \mathfrak{h} is an abelian L_∞ -algebra with underlying chain complex \mathfrak{h}_\bullet , then we also write $\mathbf{B}\mathfrak{h}$ for the abelian L_∞ -algebra with underlying chain complex $\mathfrak{h}_\bullet[1]$.

For $n \in \mathbb{N}$ we write $\mathbb{R}[n]$ for the abelian L_∞ -algebra whose underlying chain complex is \mathbb{R} concentrated in degree n , hence $\mathbb{R}[n] = \mathbf{B}^n\mathbb{R}$.

Example 3.1.5 (L_∞ -morphisms). A morphism of L_∞ -algebras $f : \mathfrak{g} \rightarrow A$ from a Lie n -algebra to a dg-Lie algebra $(A, d_A, [\cdot, \cdot]_A)$ is equivalently given by a sequence of n skew-symmetric linear maps

$$f_k : \mathfrak{g}_\bullet^{\otimes k} \rightarrow A, \quad k = 1, \dots, n \quad |f_k| = k - 1.$$

which satisfy the following identity for all $m \geq 0$:

$$\begin{aligned} d_A f_m(x_1, \dots, x_m) + \sum_{j+k=m+1} \sum_{\sigma \in \text{Sh}(k, m-k)} \chi(\sigma) (-1)^{k(j-1)+1} f_j(l_k(x_{\sigma(1)}, \dots, x_{\sigma(k)}), x_{\sigma(k+1)}, \dots, x_{\sigma(m)}) \\ + \sum_{s+t=m} \sum_{\substack{\tau \in \text{Sh}(s, m-s) \\ \tau(1) < \tau(s+1)}} \chi(\tau) (-1)^{s-1} (-1)^{(t-1)\sum_{p=1}^s |x_{\tau(p)}|} [f_s(x_{\tau(1)}, \dots, x_{\tau(s)}), f_t(x_{\tau(s+1)}, \dots, x_{\tau(m)})]_A = 0. \end{aligned} \quad (3.1.1)$$

See for instance [17, Def. 5.2].

As a special case of the above, an L_∞ -algebra morphism $f : \mathfrak{g} \rightarrow \mathfrak{h}$ from a Lie 1-algebra to an abelian Lie $(n+1)$ -algebra \mathfrak{h} is equivalently a collection of linear maps

$$\{f_k : \wedge^k \mathfrak{g}_\bullet \rightarrow \mathfrak{h}_\bullet\}_{k=1}^{n+1}$$

such that following holds for all $k \geq 1$

$$d_{\mathfrak{h}} f_k(v_1 \wedge \dots \wedge v_k) = \sum_{i < j} (-1)^{i+j+1} f_{k-1}([v_i, v_j]_{\mathfrak{g}} \wedge v_1 \wedge \dots \wedge \widehat{v}_i \wedge \dots \wedge \widehat{v}_j \wedge \dots \wedge v_k). \quad (3.1.2)$$

Example 3.1.6 (Cocycles as morphisms). L_∞ -morphisms of the form $\mathfrak{g} \rightarrow \mathbb{R}[n] = \mathbf{B}^n\mathbb{R}$ are in natural bijection with closed elements of degree $(n+1)$ in the Chevalley-Eilenberg cochain complex of \mathfrak{g} . If \mathfrak{g} is an ordinary Lie algebra, then these are just the traditional cocycles used in Lie algebra cohomology.

Quillen's adjunction and rectification

As mentioned previously, an L_∞ -algebra whose k -ary brackets for $k > 2$ are trivial is equivalently a dg-Lie algebra. Moreover, every morphism of dg-Lie algebras is a morphism of L_∞ -algebras under this identification. This non-full inclusion is part of an adjunction

$$(\mathcal{R} \dashv i) : L_\infty\text{Alg} \begin{array}{c} \xleftarrow{i} \\ \xrightarrow{\mathcal{R}} \end{array} \text{dgLie} . \quad (3.1.3)$$

We will call $i \circ \mathcal{R}$ the *rectification functor* for L_∞ -algebras. See for instance [18, Proposition 11.4.5]. We will often leave the (non-full) embedding i notationally implicit.

The general properties of the adjunction (3.1.3) give rise to the following structure, which we use below in the computation of homotopy pullbacks.

Let \mathfrak{g} be an L_∞ -algebra. There is a canonical L_∞ -algebra homomorphism $\mathfrak{g} \xrightarrow{v_{\mathfrak{g}}} \mathcal{R}(\mathfrak{g})$, namely, the unit of the adjunction, such that every L_∞ morphism $f_\infty : \mathfrak{g} \rightarrow A$ to a dg-Lie algebra A uniquely factors as

$$\mathfrak{g} \xrightarrow{v_{\mathfrak{g}}} \mathcal{R}(\mathfrak{g}) \xrightarrow{\xi_A \circ \mathcal{R}(f_\infty)} A,$$

where $\xi_A : \mathcal{R}(A) \rightarrow A$ is the dg-Lie algebra morphism in the factorization of the identity of A as

$$A \xrightarrow{v_A} \mathcal{R}(A) \xrightarrow{\xi_A} A.$$

Here $\xi_A \circ \mathcal{R}(f_\infty)$ is the *adjunct* of f_∞ . That the adjunction unit is natural means that for any morphism of L_∞ -algebras $\varphi : \mathfrak{g} \rightarrow \mathfrak{h}$ we have a commutative diagram in $L_\infty\text{Alg}$ of the form

$$\begin{array}{ccc} \mathfrak{g} & \xrightarrow{v_{\mathfrak{g}}} & \mathcal{R}(\mathfrak{g}) \\ \varphi \downarrow & & \downarrow \mathcal{R}(\varphi) \\ \mathfrak{h} & \xrightarrow{v_{\mathfrak{h}}} & \mathcal{R}(\mathfrak{h}) \end{array} .$$

Analogously, that the counit of the adjunction is natural means for the any morphism of dg-Lie algebras $\varphi : A \rightarrow B$ we have a commutative diagram of the form

$$\begin{array}{ccc} \mathcal{R}(A) & \xrightarrow{\xi_A} & A \\ \mathcal{R}(\varphi) \downarrow & & \downarrow \varphi \\ \mathcal{R}(B) & \xrightarrow{\xi_B} & B \end{array} .$$

Homotopy theory

There is a wealth of presentations for the homotopy theory of L_∞ -algebras in which homotopy fiber sequences may be interpreted, given by a web of model category structures with Quillen equivalences between them [20]. Here we make use of the model structures due to [14, 15], from which one can distill the following statement.

Theorem 3.1.7. *The category dgLie of dg-Lie algebras (over the real numbers) carries a model category structure in which the weak equivalences are the quasi-isomorphisms on the underlying chain complexes, and the fibrations are the degreewise surjections on the underlying chain complexes.*

Moreover, if we define a morphism $\mathfrak{g} \rightarrow \mathfrak{h}$ in $L_\infty\text{Alg}$ to be a weak equivalence iff the underlying morphism of complexes $\mathfrak{g}_\bullet \rightarrow \mathfrak{h}_\bullet$ is a quasi-isomorphism, then the adjunction $(\mathcal{R} \dashv i)$ induces an equivalence between the homotopy theories of dg-Lie algebras and L_∞ -algebras. In particular, the components of the unit $\mathfrak{g} \xrightarrow{v_{\mathfrak{g}}} \mathcal{R}(\mathfrak{g})$ and counit $\mathcal{R}(A) \xrightarrow{\xi_A} A$ of this adjunction are weak equivalences.

Remark 3.1.8. In [15], weak equivalences in the model category dgCoCAlg are defined to be those coalgebra morphisms $F : C \rightarrow C'$ such that $\mathcal{R}(F)$ is a weak equivalence in dgLie . It then follows from [18, Proposition 11.4.5] that a L_∞ -morphism $\mathfrak{g} \rightarrow \mathfrak{h}$ is a weak equivalence if and only if it is an L_∞ quasi-isomorphism, i.e., if $\mathfrak{g}_\bullet \rightarrow \mathfrak{h}_\bullet$ is a quasi-isomorphism of chain complexes.

Remark 3.1.9. Throughout, we consider the category of unbounded chain complexes equipped with its projective model structure, whose weak equivalences are quasi-isomorphisms, and whose fibrations are surjections.

Corollary 3.1.10. *Let $f_\infty : \mathfrak{g} \rightarrow \mathfrak{h}$ be an L_∞ -morphism between two L_∞ -algebras. Then an L_∞ -algebra \mathfrak{k} presents the homotopy fiber of f_∞ if \mathfrak{k} is L_∞ -quasi-isomorphic to the homotopy fiber of $\mathcal{R}(f_\infty) : \mathcal{R}(\mathfrak{g}) \rightarrow \mathcal{R}(\mathfrak{h})$ in the category of dglas.*

Definition 3.1.11. An L_∞ -algebra cocycle on an L_∞ -algebra \mathfrak{g} with coefficients in the abelian L_∞ -algebra A is a morphism

$$c : \mathfrak{g} \longrightarrow \mathbf{B}A .$$

The L_∞ -extension $\widehat{\mathfrak{g}}$ of \mathfrak{g} classified by c is the homotopy fiber of c which sits in the long homotopy fiber sequence of L_∞ -algebras:

$$\begin{array}{ccc} A & \longrightarrow & \widehat{\mathfrak{g}} \\ & & \downarrow \text{hfib}(c) \\ & & \mathfrak{g} \xrightarrow{c} \mathbf{B}A \end{array} .$$

In the broader context of principal ∞ -bundles, L_∞ -extensions induced by L_∞ -cocycles are discussed in [19]. (That generalized geometric cohomology is modeled by homotopy classes of maps in a suitable geometric homotopical context goes back to [5].)

We now establish a recognition principle for homotopy fibers of L_∞ -algebra morphisms (Thm. 3.1.13). We begin with a preliminary fact:

Lemma 3.1.12. *Let \mathfrak{g} be an L_∞ -algebra, A be a dgla and $f_\infty : \mathfrak{g} \rightarrow A$ an L_∞ morphism. Let $p_A : B \rightarrow A$ be a fibrant replacement of the zero morphism $0 \rightarrow A$ in the category of dglas. The fiber product*

$$\begin{array}{ccc} \mathcal{R}(\mathfrak{g}) \times_A B & \xrightarrow{\pi_B} & B \\ \pi_{\mathcal{R}(\mathfrak{g})} \downarrow & & \downarrow p_A \\ \mathcal{R}(\mathfrak{g}) & \xrightarrow{\xi_A \circ \mathcal{R}(f_\infty)} & A \end{array}$$

is a dgla model for the homotopy fiber of f_∞ .

Proof. Consider the commutative diagram of dglas

$$\begin{array}{ccccc} \mathcal{R}(\mathfrak{g}) \times_A B & \xrightarrow{(\mathcal{R}(f_\infty), \text{id}_B)} & \mathcal{R}(A) \times_A B & \xrightarrow{\pi_B} & B \\ \pi_{\mathcal{R}(\mathfrak{g})} \downarrow & & \downarrow \pi_{\mathcal{R}(A)} & & \downarrow p_A \\ \mathcal{R}(\mathfrak{g}) & \xrightarrow{\mathcal{R}(f_\infty)} & \mathcal{R}(A) & \xrightarrow{\xi_A} & A \end{array}$$

where the rightmost diagram and the outer diagram are pullbacks. By the pasting law, also the leftmost diagram is a pullback. Since p_A is a fibration and ξ_A is a weak equivalence, the map $\pi_{\mathcal{R}(A)}$ is a fibration and the map π_B is a weak equivalence. It follows that $\pi_{\mathcal{R}(A)}$ is a fibrant replacement of $0 \rightarrow \mathcal{R}(A)$. Hence $\mathcal{R}(\mathfrak{g}) \times_A B$ is a model for the homotopy fiber of $\mathcal{R}(f_\infty)$ in the category of dglas. \square

Theorem 3.1.13. *Let \mathfrak{g} be an L_∞ -algebra, A be a dgla and $f_\infty : \mathfrak{g} \rightarrow A$ an L_∞ morphism. Let $p_A : B \rightarrow A$ be a fibrant replacement of the zero morphism $0 \rightarrow A$ in the category of dglas. Consider the fiber product of chain complexes*

$$\begin{array}{ccc} \mathfrak{g} \times_A B & \xrightarrow{\pi_B} & B \\ \pi_{\mathfrak{g}} \downarrow & & \downarrow p_A \\ \mathfrak{g} & \xrightarrow{f} & A, \end{array}$$

where $f = U(f_\infty)$ is the linear part of f_∞ . Assume there exists an L_∞ -structure Q on $\mathfrak{g} \times_A B$ and L_∞ -morphisms $\pi_{B,\infty} : (\mathfrak{g} \times_A B, Q) \rightarrow B$ and $\pi_{\mathfrak{g},\infty} : (\mathfrak{g} \times_A B, Q) \rightarrow \mathfrak{g}$ such that

- $U(\pi_{B,\infty}) = \pi_B$;
- $U(\pi_{\mathfrak{g},\infty}) = \pi_{\mathfrak{g}}$;
- the diagram

$$\begin{array}{ccc} (\mathfrak{g} \times_A B, Q) & \xrightarrow{\pi_{B,\infty}} & B \\ \pi_{\mathfrak{g}} \downarrow & & \downarrow p_A \\ \mathfrak{g} & \xrightarrow{f_\infty} & A \end{array}$$

is a commutative diagram of L_∞ -algebras.

Then $(\mathfrak{g} \times_A B, Q)$ is a model for the homotopy fiber of f_∞ .

Proof. Applying the rectification functor to the diagram of L_∞ -morphisms above we get a commutative diagram of dglas

$$\begin{array}{ccc} \mathcal{R}(\mathfrak{g} \times_A B, Q) & \xrightarrow{\mathcal{R}(\pi_{B,\infty})} & \mathcal{R}(B) \\ \mathcal{R}(\pi_{\mathfrak{g},\infty}) \downarrow & & \downarrow \mathcal{R}(p_A) \\ \mathcal{R}(\mathfrak{g}) & \xrightarrow{\mathcal{R}(f_\infty)} & \mathcal{R}(A) \end{array}$$

Using the counit of the adjunction (3.1.3), we can extend this to a commutative diagram of dglas

$$\begin{array}{ccccc} \mathcal{R}(\mathfrak{g} \times_A B, Q) & \xrightarrow{\mathcal{R}(\pi_{B,\infty})} & \mathcal{R}(B) & \xrightarrow{\xi_B} & B \\ \mathcal{R}(\pi_{\mathfrak{g},\infty}) \downarrow & & \downarrow \mathcal{R}(p_A) & & \downarrow p_A \\ \mathcal{R}(\mathfrak{g}) & \xrightarrow{\mathcal{R}(f_\infty)} & \mathcal{R}(A) & \xrightarrow{\xi_A} & A \end{array}$$

By the universal property of the pullback of dglas, the outer rectangle commutative diagram

$$\begin{array}{ccc} \mathcal{R}(\mathfrak{g} \times_A B, Q) & \xrightarrow{\xi_B \circ \mathcal{R}(\pi_{B,\infty})} & B \\ \mathcal{R}(\pi_{\mathfrak{g},\infty}) \downarrow & & \downarrow p_A \\ \mathcal{R}(\mathfrak{g}) & \xrightarrow{\xi_A \circ \mathcal{R}(f_\infty)} & A \end{array}$$

is equivalent to the datum of a morphism of dglas

$$\psi : \mathcal{R}(\mathfrak{g} \times_A B, Q) \rightarrow \mathcal{R}(\mathfrak{g}) \times_A B,$$

where $\mathcal{R}(\mathfrak{g}) \times_A B$ is the pullback of dglas

$$\begin{array}{ccc} \mathcal{R}(\mathfrak{g}) \times_A B & \xrightarrow{\pi_B} & B \\ \pi_{\mathcal{R}(\mathfrak{g})} \downarrow & & \downarrow p_A \\ \mathcal{R}(\mathfrak{g}) & \xrightarrow{\xi_A \circ \mathcal{R}(f_\infty)} & A \end{array}$$

The morphism ψ will satisfy

$$\pi_B \circ \psi = \xi_B \circ \mathcal{R}(\pi_{B,\infty}); \quad \pi_{\mathcal{R}(\mathfrak{g})} \circ \psi = \mathcal{R}(\pi_{\mathfrak{g},\infty}).$$

By Lemma 3.1.12, the dgla $\mathcal{R}(\mathfrak{g}) \times_A B$ is a dgla model for the homotopy fiber of f_∞ . Then to conclude we only need to show that ψ is a quasi-isomorphism. This is equivalent to proving that the L_∞ -morphism η given by the composition

$$\eta : (\mathfrak{g} \times_A B, Q) \xrightarrow{v_{(\mathfrak{g} \times_A B, Q)}} \mathcal{R}(\mathfrak{g} \times_A B, Q) \xrightarrow{\psi} \mathcal{R}(\mathfrak{g}) \times_A B$$

is a quasi-isomorphism. The linear part $U(\eta)$ of η is determined by its compositions with the linear projections to $\mathcal{R}(\mathfrak{g})$ and to B . We have

$$\pi_B \circ U(\eta) = U(\pi_B \circ \eta) = U(\pi_B \circ \psi \circ v_{(\mathfrak{g} \times_A B, Q)}) = U(\xi_B \circ \mathcal{R}(\pi_{B, \infty}) \circ v_{(\mathfrak{g} \times_A B, Q)}) = U(\pi_{B, \infty}) = \pi_B$$

and

$$\pi_{\mathcal{R}(\mathfrak{g})} \circ U(\eta) = U(\pi_{\mathcal{R}(\mathfrak{g})} \circ \eta) = U(\pi_{\mathcal{R}(\mathfrak{g})} \circ \psi \circ v_{(\mathfrak{g} \times_A B, Q)}) = U(\mathcal{R}(\pi_{\mathfrak{g}, \infty}) \circ v_{(\mathfrak{g} \times_A B, Q)}) = U(v_{\mathfrak{g}} \circ \pi_{\mathfrak{g}, \infty}) = U(v_{\mathfrak{g}}) \circ \pi_{\mathfrak{g}}$$

This means that the map of chain complexes

$$U(\eta) : \mathfrak{g} \times_A B \rightarrow \mathcal{R}(\mathfrak{g}) \times_A B$$

is

$$U(\eta) = (U(v_{\mathfrak{g}}), \text{id}_B).$$

Now consider the commutative diagram

$$\begin{array}{ccccc} \mathfrak{g} \times_A B & \xrightarrow{(U(v_{\mathfrak{g}}), \text{id}_B)} & \mathcal{R}(\mathfrak{g}) \times_A B & \xrightarrow{\pi_B} & B \\ \pi_{\mathfrak{g}} \downarrow & & \downarrow \pi_{\mathcal{R}(\mathfrak{g})} & & \downarrow p_A \\ \mathfrak{g} & \xrightarrow{U(v_{\mathfrak{g}})} & \mathcal{R}(\mathfrak{g}) & \xrightarrow{\xi_A \circ \mathcal{R}(f_\infty)} & A \end{array}$$

The rightmost subdiagram is a pullback by definition, while the total diagram is

$$\begin{array}{ccc} \mathfrak{g} \times_A B & \xrightarrow{\pi_B} & B \\ \pi_{\mathfrak{g}} \downarrow & & \downarrow p_A \\ \mathfrak{g} & \xrightarrow{f} & A \end{array} ,$$

since

$$\xi_A \circ \mathcal{R}(f_\infty) \circ U(v_{\mathfrak{g}}) = U(\xi_A \circ \mathcal{R}(f_\infty) \circ v_{\mathfrak{g}}) = U(f_\infty) = f,$$

and so it is a pullback by hypothesis. Then, by the pasting law, also the leftmost subdiagram is a pullback. The map $\pi_{\mathcal{R}(\mathfrak{g})}$ is a fibration, since p_A is fibration, and all chain complexes are fibrant. Hence, since $U(v_{\mathfrak{g}})$ is a quasi-isomorphism, its pullback $U(\eta) = (U(v_{\mathfrak{g}}), \text{id}_B)$ is also a quasi-isomorphism. \square

3.2 The L_∞ -algebra of local observables

We recall the construction of the L_∞ -algebra of local observables associated to a pre- n -plectic manifold. It is best seen in the light of the following immediate consequence of Cartan's "magic formula" $\mathcal{L}_v = dt_v + \iota_v d$.

Lemma 3.2.1. *Let X be a smooth manifold and let β be an n -form (not necessarily closed) on X . For all $k \geq 1$ and all vector fields v_1, \dots, v_k we have:*

$$\begin{aligned} (-1)^k dt_{v_1 \wedge \dots \wedge v_k} \beta &= \sum_{1 \leq i < j \leq k} (-1)^{i+j} \iota_{[v_i, v_j]} \wedge v_1 \wedge \dots \wedge \hat{v}_i \wedge \dots \wedge \hat{v}_j \wedge \dots \wedge v_k \beta \\ &+ \sum_{i=1}^k (-1)^i \iota_{v_1 \wedge \dots \wedge \hat{v}_i \wedge \dots \wedge v_k} \mathcal{L}_{v_i} \beta \\ &+ \iota_{v_1 \wedge \dots \wedge v_k} d\beta. \end{aligned}$$

A special case of the above appeared as Lemma 3.7 in [22]. We thank M. Zambon for pointing out to us this generalization.

Proposition 3.2.2 (Thm. 5.2 [22]; Thm. 4.7 [13]). *Let (X, ω) be a pre- n -plectic manifold. There exists a Lie n -algebra $L_\infty(X, \omega)$ whose underlying chain complex is*

$$\Omega^0(X) \xrightarrow{d} \Omega^1(X) \xrightarrow{d} \dots \xrightarrow{d} \Omega^1(X) \xrightarrow{(0,d)} \text{Ham}^{n-1}(X),$$

with $\text{Ham}^{n-1}(X)$ in degree zero, and whose multilinear brackets l_i are

$$l_1(x) = \begin{cases} 0 \oplus dx & \text{if } |x| = 1, \\ dx & \text{if } |x| > 1, \end{cases}$$

$$l_2(x_1, x_2) = \begin{cases} [v_1, v_2] + \iota_{v_1 \wedge v_2} \omega & \text{if } |x_1 \otimes x_2| = 0, \\ 0 & \text{otherwise,} \end{cases}$$

and, for $k > 2$:

$$l_k(x_1, \dots, x_k) = \begin{cases} -(-1)^{\binom{k+1}{2}} \iota_{v_1 \wedge \dots \wedge v_k} \omega & \text{if } |x_1 \otimes \dots \otimes x_k| = 0, \\ 0 & \text{otherwise,} \end{cases}$$

where $x = v + \eta^\bullet$ denotes a generic element $(\eta^0, \eta^1, \dots, v + \eta^{n-1})$ in the chain complex and $|x_1 \otimes \dots \otimes x_k|$ is the total degree of the element $x_1 \otimes \dots \otimes x_k$.

Definition 3.2.3. We call the Lie n -algebra $L_\infty(X, \omega)$ defined in the statement of Proposition 3.2.2 the L_∞ -algebra of local observables on (X, ω) .

Remark 3.2.4. The projection map of def. 2.1.5 uniquely extends to a morphism of L_∞ -algebras of the form

$$\begin{array}{c} L_\infty(X, \omega) \\ \downarrow \pi_L \\ \mathfrak{X}_{\text{Ham}}(X) \end{array},$$

i.e., local observables of (X, ω) cover Hamiltonian vector fields. Below in 3.3 we turn to the classification of this map by an L_∞ -algebra cocycle.

Example 3.2.5. If $n = 1$ then (X, ω) is a pre-symplectic manifold and the chain complex underlying $L_\infty(X, \omega)$ is

$$\text{Ham}^0(X) = \{v + H \in \mathfrak{X}(X) \oplus C^\infty(X; \mathbb{R}) \mid \iota_v \omega + dH = 0\},$$

and the Lie bracket is

$$[v_1 + H_1, v_2 + H_2] = [v_1, v_2] + \iota_{v_1 \wedge v_2} \omega.$$

If moreover ω is non-degenerate so that (X, ω) is symplectic, then the projection $v + H \mapsto H$ is a linear isomorphism $\text{Ham}^0(X) \xrightarrow{\cong} C^\infty(X; \mathbb{R})$. It is easy to see that under this isomorphism $L_\infty(X, \omega)$ is the underlying Lie algebra of the usual Poisson algebra of functions. See also Prop. 2.3.9 in [6].

3.3 The Kostant-Souriau L_∞ -algebra cocycle

Here we present an L_∞ -algebra cocycle (def. 3.1.11) on the Lie algebra of Hamiltonian vector fields on a pre- n -plectic manifold, which generalizes the traditional Kostant-Souriau cocycle and the Heisenberg cocycle to higher geometry.

Definition 3.3.1. For X a smooth manifold, denote by $\mathbf{BH}(X, \mathfrak{b}\mathbf{B}^{n-1}\mathbb{R})$ the abelian Lie $(n+1)$ -algebra given by the chain complex

$$\Omega^0(X) \xrightarrow{d} \Omega^1(X) \xrightarrow{d} \dots \xrightarrow{d} \Omega^{n-1}(X) \xrightarrow{d} d\Omega^{n-1}(X),$$

with $d\Omega^{n-1}(X)$ in degree zero.

Remark 3.3.2. The complex of def. 3.3.1 serves as a resolution of the cocycle complex

$$\Omega^0(X) \xrightarrow{d} \Omega^1(X) \xrightarrow{d} \dots \xrightarrow{d} \Omega_{\text{cl}}^{n-1}(X) \rightarrow 0,$$

for the de Rham cohomology of X up to degree $n-1$ once delooped (i.e., shifted).

Proposition 3.3.3. *Let (X, ω) be a pre- n -plectic manifold. The multilinear maps*

$$\omega_{[1]} : v \mapsto -\iota_v \omega; \quad \omega_{[2]} : v_1 \wedge v_2 \mapsto \iota_{v_1} \wedge v_2 \omega; \quad \dots \quad \omega_{[n+1]} : v_1 \wedge v_2 \wedge \dots \wedge v_{n+1} \mapsto -(-1)^{\binom{n+1}{2}} \iota_{v_1 \wedge v_2 \wedge \dots \wedge v_{n+1}} \omega$$

define an L_∞ -morphism

$$\omega_{[\bullet]} : \mathfrak{X}_{\text{Ham}}(X) \rightarrow \mathbf{BH}(X, \mathfrak{b}\mathbf{B}^{n-1}\mathbb{R}),$$

and hence an L_∞ -algebra $(n+1)$ -cocycle on the Lie algebra of Hamiltonian vector fields, def. 2.1.5, with values in the abelian $(n+1)$ -algebra of def. 3.3.1.

Proof. First notice that the underlying map on chain complexes is indeed well defined: by definition of Hamiltonian vector fields, if v is Hamiltonian, then there exists an $(n-1)$ -form H such that $\iota_v \omega + dH = 0$ and so $\omega_{[\bullet]}$ takes values in $\mathbf{BH}(X, \mathfrak{b}\mathbf{B}^{n-1}\mathbb{R})$. To check that the morphism is indeed a L_∞ -morphism, notice that in the present situation the equations (3.1.2) of example 3.1.5 become

$$d\iota_{v_1 \wedge \dots \wedge v_k} \omega = \sum_{i < j} (-1)^{i+j+k} \iota_{[v_i, v_j]} \wedge v_1 \wedge \dots \wedge \widehat{v}_i \wedge \dots \wedge \widehat{v}_j \wedge \dots \wedge v_{k+1} \omega.$$

These are satisfied – since the ω is closed and the v_i are Hamiltonian – by Lemma 3.2.1. □

Definition 3.3.4. The degree $(n+1)$ higher Kostant-Souriau L_∞ -cocycle associated to the pre- n -plectic manifold (X, ω) is the L_∞ -morphism

$$\omega_{[\bullet]} : \mathfrak{X}_{\text{Ham}}(X) \longrightarrow \mathbf{BH}(X, \mathfrak{b}\mathbf{B}^{n-1}\mathbb{R})$$

given in Prop. 3.3.3.

If $\rho : \mathfrak{g} \rightarrow \mathfrak{X}_{\text{Ham}}(X)$ is an L_∞ -morphism encoding an action of an L_∞ -algebra \mathfrak{g} on (X, ω) by Hamiltonian vector fields, then we call the composite $\rho^* \omega_{[\bullet]}$ the corresponding *Heisenberg L_∞ -algebra cocycle*. This terminology is motivated by the following example 3.3.5. Further discussion of this aspect is below in Section 3.5.

Example 3.3.5. Let V be a vector space equipped with a skew-symmetric multilinear form $\omega : \Lambda^{n+1}V \rightarrow \mathbb{R}$. Since V is an abelian Lie group, we obtain via left-translation of ω a unique closed invariant form, which we also denote as ω . By identifying V with left-invariant vector fields on V , the Poincaré lemma implies that we have a canonical inclusion

$$j_V : V \hookrightarrow \mathfrak{X}_{\text{Ham}}(V)$$

of V regarded as an abelian Lie algebra into the Hamiltonian vector fields on (V, ω) regarded as a pre- n -plectic manifold. Since V is contractible as a topological manifold, we have, by remark 3.3.2, a quasi-isomorphism

$$\mathbf{BH}(V; \mathfrak{b}\mathbf{B}^{n-1}\mathbb{R}) \xrightarrow{\cong} \mathbb{R}[n]$$

By inspection and comparison with prop. 3.2.2 it is easy to see that the fiber product of chain complexes of K and $\mathfrak{X}_{\text{Ham}}(X)$ over $\mathbf{BH}(X, \mathfrak{b}\mathbf{B}^{n-1}\mathbb{R})$ is the chain complex $L_\infty(X, \omega)_\bullet$ that underlies the L_∞ -algebra of local observables:

$$\begin{array}{ccc} L_\infty(X, \omega)_\bullet & \xrightarrow{f_1} & K \\ \pi_L \downarrow & & \downarrow \pi_R \\ \mathfrak{X}_{\text{Ham}}(X) & \xrightarrow{\omega^{[1]}} & \mathbf{BH}(X, \mathfrak{b}\mathbf{B}^{n-1}\mathbb{R}), \end{array},$$

where f_1 is the morphism

$$f_1 : v + \eta^\bullet \mapsto \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 & 0 & \\ & \eta^0 & \eta^1 & \cdots & \eta^{n-3} & \eta^{n-2} & \eta^{n-1} \end{pmatrix}.$$

As we already observed in remark 3.2.4, the chain map underlying π_L uniquely extends to an L_∞ -algebra morphism. Therefore to complete the proof, it is sufficient to show that we can lift the horizontal chain map f_1 above to a morphism of L_∞ -algebras which makes the diagram

$$\begin{array}{ccc} L_\infty(X, \omega) & \xrightarrow{f} & K \\ \pi_L \downarrow & & \downarrow \pi_R \\ \mathfrak{X}_{\text{Ham}}(X) & \xrightarrow{\omega^{[\bullet]}} & \mathbf{BH}(X, \mathfrak{b}\mathbf{B}^{n-1}\mathbb{R}), \end{array}$$

commute. This is easily realized by defining the ‘‘Taylor coefficients’’ of f for $k \geq 2$ to be the degree $(1-k)$ maps $f_k : \wedge^k L_\infty(X, \omega) \rightarrow K$ given by

$$f_k : (v_1 + \eta_1^\bullet) \wedge \cdots \wedge (v_k + \eta_k^\bullet) \mapsto \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 & 0 \\ & 0 & 0 & \cdots & 0 & -(-1)^{\binom{k+1}{2}} \iota_{v_1 \wedge \cdots \wedge v_k} \omega & 0 & \cdots & 0 \end{pmatrix}.$$

□

3.5 The Heisenberg L_∞ -extension

If a Lie algebra \mathfrak{g} acts on an n -plectic manifold by Hamiltonian vector fields, then the Kostant-Souriau L_∞ -extension of $\mathfrak{X}_{\text{Ham}}(X)$, discussed above in 3.4, restricts to an L_∞ -extension of \mathfrak{g} . This is a generalization of Kostant’s construction [16] of central extensions of Lie algebras to the context of L_∞ -algebras. Perhaps the most famous of these central extensions is the Heisenberg Lie algebra, which is the inspiration behind the following terminology:

Definition 3.5.1. Let (X, ω) be a pre- n -plectic manifold and let $\rho : \mathfrak{g} \rightarrow \mathfrak{X}_{\text{Ham}}(X)$ be a Lie algebra homomorphism encoding an action of \mathfrak{g} on X by Hamiltonian vector fields. The corresponding *Heisenberg L_∞ -algebra extension* $\mathfrak{heis}_\rho(\mathfrak{g})$ of \mathfrak{g} is the extension classified by the composite L_∞ -morphism $\omega^{[\bullet]} \circ \rho$, i.e. the homotopy pullback on the left of

$$\begin{array}{ccccc} \mathfrak{heis}_\rho(\mathfrak{g}) & \longrightarrow & L_\infty(X, \omega) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow \\ \mathfrak{g} & \xrightarrow{\rho} & \mathfrak{X}_{\text{Ham}}(X) & \xrightarrow{\omega^{[\bullet]}} & \mathbf{BH}(X, \mathfrak{b}\mathbf{B}^{n-1}\mathbb{R}) \end{array}.$$

Remark 3.5.2. It is natural to call an L_∞ -morphism with values in the L_∞ -algebra of observables of a pre- n -plectic manifold (X, ω) an ‘ L_∞ co-moment map’, which generalizes the familiar notion in symplectic geometry. Hence, one could say that an action ρ of a Lie algebra \mathfrak{g} on a pre- n -plectic manifold (X, ω) via Hamiltonian vector fields naturally induces such a co-moment map from the Heisenberg L_∞ -algebra $\mathfrak{heis}_\rho(\mathfrak{g})$.

Example 3.5.3. For (V, ω) a symplectic vector space regarded as a symplectic manifold, the translation action of V on itself is via Hamiltonian vector fields (see example 3.3.5). If one denotes by

$$\rho : V \rightarrow \mathfrak{X}_{\text{Ham}}(X)$$

this action, then the induced Heisenberg L_∞ -extension is the traditional Heisenberg Lie algebra.

Example 3.5.4. Let G be a (connected) compact simple Lie group, regarded as a 2-plectic manifold with its canonical 3-form $\omega := \langle -, [-, -] \rangle$ as in example 2.1.3. The infinitesimal generators of the action of G on itself by right translation are the left invariant vector fields \mathfrak{g} , which are Hamiltonian. We have $H_{\text{dR}}^1(G) \cong H_{\text{CE}}^1(\mathfrak{g}, \mathbb{R}) = 0$, and therefore a weak equivalence:

$$\mathbf{BH}(G, \mathfrak{b}\mathbf{B}\mathbb{R}) \xrightarrow{\simeq} \mathbb{R}[2]$$

given by the evaluation at the identity element of G . The resulting composite cocycle

$$\langle -, [-, -] \rangle : \mathfrak{g} \xrightarrow{\rho} \mathfrak{X}_{\text{Ham}}(X) \xrightarrow{\omega[\bullet]} \mathbb{R}[2]$$

is exactly the 3-cocycle which classifies the String Lie-2-algebra, namely just $\langle -, [-, -] \rangle$ regarded as a Lie algebra 3-cocycle. By theorem 3.1.13 the String Lie 2-algebra is the homotopy fiber of this cocycle, in that we have a homotopy pullback square of L_∞ -algebras

$$\begin{array}{ccc} \mathfrak{string}_{\mathfrak{g}} & \longrightarrow & 0 \\ \downarrow & & \downarrow \\ \mathfrak{g} & \xrightarrow{\langle -, [-, -] \rangle} & \mathbb{R}[2] \end{array} .$$

Hence the String Lie 2-algebra is the Heisenberg Lie 2-algebra of the 2-plectic manifold $(G, \langle -, [-, -] \rangle)$ with its canonical \mathfrak{g} -action ρ :

$$\mathfrak{heis}_{\rho}(\mathfrak{g}) \simeq \mathfrak{string}_{\mathfrak{g}} .$$

The relationship between $\mathfrak{string}_{\mathfrak{g}}$ and $L_\infty(G, \omega)$ was first explored in [3].

4 The dg-Lie algebra of infinitesimal quantomorphisms

The L_∞ -algebra $L_\infty(X, \omega)$ discussed above in Section 3 has the nice property that the definition of its brackets generalizes the definition of the traditional Poisson bracket in an elegant way. We now present another L_∞ -algebra which looks a little less elegant in components, but has a more manifest conceptual interpretation, namely as the dg-Lie algebra of infinitesimal automorphisms of a $U(1)$ - n -bundle with connection which cover the diffeomorphisms of the base.

A main result of this section is Thm. 4.2.2, which establishes a weak equivalence between the aforementioned dg Lie algebra and $L_\infty(M, \omega)$.

4.1 Quantomorphism n -groups

Since, by definition, a prequantization of a pre- n -plectic manifold (X, ω) is a morphism of higher stacks $X \rightarrow \mathbf{B}^n U(1)_{\text{conn}}$, a prequantized pre- n -plectic manifold is naturally an object in the overcategory (or “slice topos”) $\mathbf{H}/_{\mathbf{B}^n U(1)_{\text{conn}}}$. This leads to the following definition.

Definition 4.1.1. Let $\nabla_0, \nabla_1 : X \rightarrow \mathbf{B}^n U(1)_{\text{conn}}$ be two morphisms representing (or “modulating”) principal $U(1)$ - n -bundles with connection on X . A 1-morphism $(\phi, \eta) : \nabla_0 \rightarrow \nabla_1$ in $\mathbf{H}/_{\mathbf{B}^n U(1)_{\text{conn}}}$ is a homotopy

commutative diagram of the form

$$\begin{array}{ccc}
X & \xrightarrow{\phi} & X \\
\searrow \nabla_0 & \swarrow \eta & \swarrow \nabla_1 \\
& \mathbf{B}^n U(1)_{\text{conn}} &
\end{array}$$

A 2-morphism $(k, h): (\phi_1, \eta_1) \rightarrow (\phi_2, \eta_2)$ is only between 1-morphisms such that $\phi_1 = \phi_2$ and is given by a homotopy commutative diagram of the form

$$\begin{array}{ccc}
X & \xrightarrow{\phi_1 = \phi_2} & X \\
\searrow \nabla_1 & \swarrow \eta_1 & \swarrow \nabla_0 \\
& \mathbf{B}^n U(1)_{\text{conn}} & \\
\swarrow k \downarrow & & \\
\nabla_1 & & \nabla_0
\end{array}
,$$

where one has the (undisplayed) 2-arrow η_2 on the back face of the diagrams and an undisplayed 3-arrow $h: k \circ \eta_1 \rightarrow \eta_2$ decorating the bulk of the 3-simplex. Higher morphisms are defined similarly.

Remark 4.1.2. Since we are dealing with a commutative diagram of morphisms between (higher) stacks, we have the homotopy η appearing here as part of the data of the commutative diagram defining a 1-morphism. In particular, isomorphisms (or better, equivalences) between ∇_0 and ∇_1 will be pairs (ϕ, η) consisting of a diffeomorphism $\phi: X \rightarrow X$ and a gauge transformation of higher connections

$$\eta: \phi^* \nabla_1 \xrightarrow{\cong} \nabla_0.$$

In particular, for the 1-plectic (i.e. symplectic) case, ∇_0 and ∇_1 correspond to principal $U(1)$ -bundles with connection. If X is compact, then the 1-morphisms between them correspond to “strict contactomorphisms” $(P_0, A_0) \rightarrow (P_1, A_1)$ between the total spaces of the bundles with their connection 1-forms $A_i \in \Omega^1(P_i; \mathbb{R})$ regarded as “regular” contact forms. If $\nabla = \nabla_0 = \nabla_1$ and ∇ is regarded as the prequantization of its curvature i.e. the symplectic 2-form ω , then such a contactomorphism is often called a *quantomorphism* in the geometric quantization literature.

The automorphisms $\mathbf{Aut}_{/\mathbf{B}^n U(1)_{\text{conn}}}(\nabla)$ of any object $\nabla \in \mathbf{H}_{/\mathbf{B}^n U(1)_{\text{conn}}}$ form an “ n -group” (see, for example, Sec. 2.3 of [19]). And so, motivated by the terminology used in the above remark, we introduce the following definition.

Definition 4.1.3. Let $\nabla: X \rightarrow \mathbf{B}^n U(1)_{\text{conn}}$ be a morphism modulating a $U(1)$ - n -bundle with connection. The *quantomorphism n -group* of ∇ :

$$\mathbf{QuantMorph}(\nabla)$$

is the automorphism n -group $\mathbf{Aut}_{/\mathbf{B}^n U(1)_{\text{conn}}}(\nabla)$ equipped with its natural smooth structure.

Remark 4.1.4. In the above definition we described $\mathbf{QuantMorph}(\nabla)$ as a “smooth n -group”. In order to make this precise, we need to say what a smooth family of automorphisms is. This is systematically done by working with smooth families from the very beginning, i.e., by replacing the hom-spaces $\mathbf{H}(X, \mathbf{B}^n U(1)_{\text{conn}})$ by what we call the “concretification” of the internal homs (the higher mapping stacks) $[X, \mathbf{B}^n U(1)_{\text{conn}}]$. See Sec. 2.3.2 of [10] for precise discussion of this aspect. The intuition behind this smooth structure - which is all that we need for our purposes here - is that all local bundle data depends smoothly on a parameter varying in the base. For instance, a smooth path in the concretification of $[X, \mathbf{B}U(1)_{\text{conn}}]$ is given by

1. smooth 1-forms $A_\alpha \in \Omega^1(U_\alpha \times [0, 1], \mathbb{R})$ such that $\iota_{\partial/\partial t} A_\alpha = 0$;
2. smooth functions $g_{\alpha\beta}: U_{\alpha\beta} \times [0, 1] \rightarrow U(1)$;

where $\{U_\alpha \rightarrow X\}_\alpha$ is a good open cover of X , such that

1. $A_\beta = A_\alpha + d_X \log g_{\alpha\beta}$ on $U_{\alpha\beta}$;
2. $g_{\alpha\beta} g_{\beta\gamma} g_{\gamma\alpha} = 1$ on $U_{\alpha\beta\gamma}$.

In the above formulae, t is a coordinate on $[0, 1]$ and d_X is the de Rham differential in the X -direction. In even more explicit terms, if $\{u^1, \dots, u^n\}$ are coordinates on U_α , we are requiring that the 1-form A_α has a coordinate expression of the form $A_{\alpha,i}(t, u^1, \dots, u^n) du^i$ and that

$$A_{\beta,i}(t, u^1, \dots, u^n) = A_{\alpha,i}(t, u^1, \dots, u^n) + \frac{\partial \log g_{\alpha\beta}(t, u^1, \dots, u^n)}{\partial u^i}$$

for any i .

4.2 Infinitesimal quantomorphisms as a strict model for the L_∞ -algebra of observables

Since the quantomorphism n -group $\mathbf{QuantMorph}(\nabla)$ is equipped with a smooth structure, it has a notion of “tangent vectors”. Roughly speaking, these correspond to maps out of the formal interval: $\mathrm{Spec}(\mathbb{R}[\epsilon]/(\epsilon^2)) \rightarrow \mathbf{QuantMorph}(\nabla)$. So it is not surprising that there is also an abstract notion of “Lie differentiation” in this context which, when applied to the smooth n -group $\mathbf{QuantMorph}(\nabla)$, produces not a Lie algebra, but rather a Lie n -algebra. (See Sec. 3.10.9 and Sec. 4.5.1.2 in [26] for more details). This Lie n -algebra is technically only defined up to weak equivalence, and although it will not make an explicit appearance here, for the purposes of discussion we denote it as $\mathrm{Lie}(\mathbf{QuantMorph}(\nabla))$.

The defining equations of $\mathrm{Lie}(\mathbf{QuantMorph}(\nabla))$ can be conceptually described as the infinitesimal versions of the defining equations for the quantomorphism n -group. In particular, a degree zero element in $\mathrm{Lie}(\mathbf{QuantMorph}(\nabla))$ will be an infinitesimal version of a pair $(\phi : X \xrightarrow{\sim} X, h : \phi^* \nabla \xrightarrow{\sim} \nabla)$, i.e. a pair (v, b) consisting of a vector field v on X and an “infinitesimal homotopy” b such that

$$b : \mathcal{L}_v \nabla \rightarrow 0,$$

where \mathcal{L}_v is the Lie derivative along v . Degree 1 elements in $\mathrm{Lie}(\mathbf{QuantMorph}(\nabla))$ will be homotopies between the b 's, and so on.

The notion of taking the Lie derivative of a morphism of higher stacks may give pause, but it has an obvious interpretation if we represent the map $\nabla : X \rightarrow \mathbf{B}^n U(1)_{\mathrm{conn}}$ as a Čech-Deligne cocycle \bar{A} on X (def. 2.2). In this context, $\mathcal{L}_v \nabla$ corresponds to the usual Lie derivative $\mathcal{L}_v \bar{A}$ for vector fields acting on local differential forms. Moreover, in this context the Dold-Kan correspondence tells us that, for example, an infinitesimal homotopy $b : \mathcal{L}_v \nabla \rightarrow 0$ is simply an element $\bar{\theta}$ of the total Čech-de Rham complex $(\mathrm{Tot}^\bullet(\mathcal{U}, \Omega), d_{\mathrm{Tot}})$ satisfying

$$\mathcal{L}_v \bar{A} = d_{\mathrm{Tot}} \bar{\theta}.$$

The above discussion is the intuition behind the following:

Definition/Proposition 4.2.1. Let X be a smooth manifold and $n \in \mathbb{N}$. If \bar{A} is a Čech-Deligne $(n+1)$ -cocycle on X relative to some cover \mathcal{U} , then the *dg Lie algebra of infinitesimal quantomorphisms*

$$\mathrm{dgLie}_{\mathrm{Qu}}(X, \bar{A})$$

is the strict Lie n -algebra whose underlying complex is

$$\begin{aligned} \mathrm{dgLie}_{\mathrm{Qu}}(X, \bar{A})^0 &= \{v + \bar{\theta} \in \mathfrak{X}(M) \oplus \mathrm{Tot}^{n-1}(\mathcal{U}, \Omega) \mid \mathcal{L}_v \bar{A} = d_{\mathrm{Tot}} \bar{\theta}\}, \\ \mathrm{dgLie}_{\mathrm{Qu}}(X, \bar{A})^i &= \mathrm{Tot}^{n-1-i}(\mathcal{U}, \Omega) \quad \text{for } 1 \leq i \leq n-1, \end{aligned}$$

with differential

$$\mathrm{dgLie}_{\mathrm{Qu}}(X, \bar{A})^{n-1} \xrightarrow{d_{\mathrm{Tot}}} \mathrm{dgLie}_{\mathrm{Qu}}(X, \bar{A})^{n-2} \xrightarrow{d_{\mathrm{Tot}}} \dots \xrightarrow{d_{\mathrm{Tot}}} \mathrm{dgLie}_{\mathrm{Qu}}(X, \bar{A})^1 \xrightarrow{0 \oplus d_{\mathrm{Tot}}} \mathrm{dgLie}_{\mathrm{Qu}}(X, \bar{A})^0,$$

and whose graded Lie bracket is the semidirect product bracket for the Lie algebra of vector fields acting on differential forms by Lie derivative:

$$\begin{aligned} \llbracket v_1 + \bar{\theta}_1, v_2 + \bar{\theta}_2 \rrbracket &= [v_1, v_2] + \mathcal{L}_{v_1} \bar{\theta}_2 - \mathcal{L}_{v_2} \bar{\theta}_1 \\ \llbracket v + \bar{\theta}, \bar{\eta} \rrbracket &= - \llbracket \bar{\eta}, v + \bar{\theta} \rrbracket = \mathcal{L}_v \bar{\eta} \\ \llbracket \bar{\eta}, \bar{\eta} \rrbracket &= 0. \end{aligned} \tag{4.2.1}$$

The next theorem reveals the relationship between the above dgla of infinitesimal quantomorphisms and the L_∞ -algebra of local observables. It is the higher analog of the well-known fact in traditional prequantization that the underlying Lie algebra of the Poisson algebra on a prequantized symplectic manifold is isomorphic to the Lie algebra of $U(1)$ -invariant connection-preserving vector fields on the total space of the prequantum bundle.

Theorem 4.2.2. *If (X, ω) is a pre- n -plectic manifold (def. 2.1.1) and ∇ is a prequantization of (X, ω) (def. 2.2.4) presented by a Čech-Deligne cocycle $\bar{A} \in \check{C}^n(\mathcal{U}, \underline{U}(1))$, then there is a natural weak equivalence (in the sense of Thm. 3.1.7)*

$$L_\infty(X, \omega) \xrightarrow{\simeq} \mathrm{dgLie}_{\mathrm{Qu}}(X, \bar{A}) ,$$

in $L_\infty\mathrm{Alg}$ between the L_∞ -algebra of local observables (def. 3.2.3) and the dgla of infinitesimal quantomorphisms. (def. 4.2.1).

Proof. We obtain the weak equivalence by first constructing a chain map using the familiar quasi-isomorphism between the de Rham complex and the total Čech-de Rham complex. We then show explicitly that this map lifts to a L_∞ -morphism. The proof itself is a lengthy but straightforward computation. We report it in Appendix A. \square

Corollary 4.2.3. *The image of the natural projection $\mathrm{dgLie}_{\mathrm{Qu}}(X, \bar{A}) \rightarrow \mathfrak{X}(X)$ is the subspace $\mathfrak{X}_{\mathrm{Ham}}(X)$ of Hamiltonian vector fields. That is, the infinitesimal quantomorphisms cover infinitesimal Hamiltonian n -plectomorphisms.*

Remark 4.2.4. Theorem 4.2.2 implies that $\mathrm{dgLie}_{\mathrm{Qu}}(X, \bar{A})$ is independent, up to equivalence, of the choice of prequantization \bar{A} of ω . The theorem also says that $\mathrm{dgLie}_{\mathrm{Qu}}(X, \bar{A})$ is a ‘rectification’ or ‘semi-strictification’ of the L_∞ -algebra $L_\infty(X, \omega)$.

5 Inclusion into Atiyah and Courant L_∞ -algebras

If (X, ω) is a prequantized symplectic manifold, and (P, A) is the corresponding prequantum bundle, then there is an embedding, induced by the morphism given in Thm. 4.2.2, of the Lie algebra of observables on X into the Lie algebra of $U(1)$ -invariant vector fields on P . The latter is the Lie algebra of global sections of the Atiyah algebroid of P (see, for example, Sec. 2 of [23] and Def. 5.1.1 below). The integrated analog of this embedding is a canonical map from the group of quantomorphisms to the group of bisections [8, Chap. 15] of the Lie groupoid which integrates the Atiyah algebroid. This groupoid is usually called the gauge groupoid of P , but we prefer to call it the ‘Atiyah groupoid’. Likewise, we call its group of bisections the ‘Atiyah group’. Such a bisection is just an equivariant diffeomorphism of P covering a diffeomorphism of the base X , and hence it “forgets” the connection 1-form A .

In analogy with the above, we now explain how similar embeddings of quantomorphisms naturally arise in the higher case. This provides the motivation for the Lie-theoretic results presented in this section.

Higher Atiyah groups and the Courant n -group

Recall from Section 2.2 that the n -stack $\mathbf{B}^n U(1)_{\text{conn}}$ is presented via the Dold-Kan correspondence by the presheaf of chain complexes

$$C^\infty(-; U(1)) \xrightarrow{d\log} \Omega^1(-) \xrightarrow{d} \Omega^2(-) \xrightarrow{d} \dots \xrightarrow{d} \Omega^n(-)$$

with $\Omega^n(-)$ in degree zero. We can also consider, for example, the n -stack $\mathbf{B}(\mathbf{B}^{n-1}U(1)_{\text{conn}})$, which is the delooping of the $(n-1)$ stack $\mathbf{B}^{n-1}U(1)_{\text{conn}}$. It is presented by the presheaf

$$C^\infty(-; U(1)) \xrightarrow{d\log} \Omega^1(-) \xrightarrow{d} \Omega^2(-) \xrightarrow{d} \dots \xrightarrow{d} \Omega^{n-1}(-) \rightarrow 0$$

with $\Omega^{n-1}(-)$ in degree 1. Hence there is a commutative diagram

$$\begin{array}{ccccccc} C^\infty(-; U(1)) & \xrightarrow{d\log} & \Omega^1(-) & \xrightarrow{d} & \dots & \xrightarrow{d} & \Omega^{n-1}(-) & \xrightarrow{d} & \Omega^n(-) \\ \downarrow & & \downarrow & & & & \downarrow & & \downarrow \\ C^\infty(-; U(1)) & \xrightarrow{d\log} & \Omega^1(-) & \xrightarrow{d} & \dots & \xrightarrow{d} & \Omega^{n-1}(-) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & & & \downarrow & & \downarrow \\ \dots & & \dots & & \dots & & \dots & & \dots \\ \downarrow & & \downarrow & & & & \downarrow & & \downarrow \\ C^\infty(-; U(1)) & \longrightarrow & 0 & \xrightarrow{d} & \dots & \longrightarrow & 0 & \longrightarrow & 0 \end{array}$$

corresponding to a sequence of natural forgetful morphisms of stacks

$$\mathbf{B}^n U(1)_{\text{conn}} \rightarrow \mathbf{B}(\mathbf{B}^{n-1}U(1)_{\text{conn}}) \rightarrow \dots \rightarrow \mathbf{B}^n U(1),$$

where at each step the top differential form data for the connection are forgotten.

If $\nabla : X \rightarrow \mathbf{B}^n U(1)_{\text{conn}}$ is the morphism representing a $U(1)$ - n -bundle with connection on a smooth manifold X , then the forgetful morphisms realize X both as an object over $\mathbf{B}(\mathbf{B}^{n-1}U(1)_{\text{conn}})$ and as an object over $\mathbf{B}^n U(1)$. Therefore we have a sequence of automorphism n -groups¹ of ∇ :

$$\mathbf{Aut}_{/\mathbf{B}^n U(1)_{\text{conn}}}(\nabla) \rightarrow \mathbf{Aut}_{/\mathbf{B}(\mathbf{B}^{n-1}U(1)_{\text{conn}})}(\nabla) \rightarrow \mathbf{Aut}_{/\mathbf{B}^n U(1)}(\nabla).$$

We call the n -group $\mathbf{Aut}_{/\mathbf{B}^n U(1)}(\nabla)$ the ‘Atiyah n -group’ of ∇ , since for the case $n = 1$, it is the previously mentioned Atiyah group. We call $\mathbf{Aut}_{/\mathbf{B}(\mathbf{B}^{n-1}U(1)_{\text{conn}})}(\nabla)$ the ‘Courant n -group’ of ∇ , since for the $n = 2$ case it can be thought of as the object that integrates the Courant Lie 2-algebra. (see Def. 5.2.2 below). A more detailed discussion of these objects as the bisections of smooth ∞ -groupoids appears in [10].

Conceptually speaking, the infinitesimal analog of the above sequence of n -groups is a sequence of Lie n -algebras

$$\mathbf{LieQuantMorph}(\nabla) \rightarrow \mathbf{LieCourant}(\nabla) \rightarrow \mathbf{LieAtiyah}(\nabla), \quad (5.0.2)$$

where $\mathbf{LieQuantMorph}(\nabla)$ is the Lie n -algebra of infinitesimal quantomorphisms described in the beginning of Sec. 4.2. The elements of $\mathbf{LieAtiyah}(\nabla)$ are to be thought of as those infinitesimal autoequivalences which preserve only the underlying $U(1)$ - n -bundle of ∇ , while $\mathbf{LieCourant}(\nabla)$ consists of those infinitesimal autoequivalences which preserve all of the connection data on the n -bundle except the highest degree part.

Recall that we modeled the Lie n -algebra $\mathbf{LieQuantMorph}(\nabla)$ by using the dg Lie algebra $\text{dgLie}_{\text{Qu}}(X, \bar{A})$ given in Def/Prop. 4.2.1. Similarly, we define below dg Lie algebras which can be thought of as models for $\mathbf{LieAtiyah}(\nabla)$ and $\mathbf{LieCourant}(\nabla)$ for the $n = 1$ and $n = 2$ cases. We consider these particular cases in order to relate our results to the traditional theory of prequantum $U(1)$ -bundles and also more recent work on Courant algebroids and $U(1)$ -bundle gerbes.

¹All of these automorphism n -groups have a smooth structure and are ‘concretified’ in the sense of Remark 4.1.4.

5.1 The $n = 1$ case

Here (X, ω) is an ordinary pre-symplectic manifold, and the algebra of local observables $L_\infty(X, \omega)$ (Def. 3.2.3) is the underlying Lie algebra of the Poisson algebra of Hamiltonian functions. A prequantization ∇ is an ordinary $U(1)$ -principal bundle with connection over X .

From any closed 2-form ω , one can construct a Lie algebroid over X whose global sections form the following Lie algebra (see for example, Sec. 2 of [23]):

Definition 5.1.1. Let (X, ω) be a presymplectic manifold. The *Atiyah Lie algebra* $\mathfrak{atiyah}(X, \omega)$ is the vector space $\mathfrak{X}(X) \oplus C^\infty(X; \mathbb{R})$ endowed with the Lie bracket

$$\llbracket v_1 + c_1, v_2 + c_2 \rrbracket_{\text{at}} = [v_1, v_2] + \mathcal{L}_{v_1} c_2 - \mathcal{L}_{v_2} c_1 - \omega(v_1, v_2).$$

Obviously the underlying vector space of the Lie algebra $L_\infty(X, \omega)$ is a subspace of $\mathfrak{atiyah}(X, \omega)$. A straightforward calculation shows that the inclusion

$$L_\infty(X, \omega) \hookrightarrow \mathfrak{atiyah}(X, \omega) \tag{5.1.1}$$

is also a Lie algebra morphism.

Just like in our construction of $\text{dgLie}_{\text{Qu}}(X, \bar{A})$ (Def/Prop. 4.2.1), we now represent the prequantization ∇ by a Čech-Deligne 1-cocycle (2.2), and obtain a model for $\text{LieAtiyah}(\nabla)$.

Definition 5.1.2. If $\bar{A} = A^1 + A^0$ is a Čech-Deligne 1-cocycle on X relative to some cover \mathcal{U} , then $\text{Lie}_{\text{At}}(X, \bar{A})$ is the Lie algebra whose underlying vector space is

$$\text{LieAlg}_{\text{At}}(X, \bar{A}) = \{v + \bar{\theta} \in \mathfrak{X}(X) \oplus \check{C}^0(\mathcal{U}, \Omega^0) \mid \mathcal{L}_v A^0 = \delta \bar{\theta}\}$$

with Lie bracket

$$\llbracket v_1 + \bar{\theta}_1, v_2 + \bar{\theta}_2 \rrbracket_{\text{At}} = [v_1, v_2] + \mathcal{L}_{v_1} \bar{\theta}_2 - \mathcal{L}_{v_2} \bar{\theta}_1 \tag{5.1.2}$$

Since $\mathcal{L}_v A^0 = \iota_v d \log A^0$, it is easy to see that an element of $\text{LieAlg}_{\text{At}}(X, \bar{A})$ is the local data corresponding to a $U(1)$ -invariant vector field on the total space P of the prequantum bundle, i.e. a global section of the Atiyah algebroid $TP/U(1) \rightarrow X$. Moreover, by construction, there is an inclusion of Lie algebras

$$\text{dgLie}_{\text{Qu}}(X, \bar{A}) \hookrightarrow \text{LieAlg}_{\text{At}}(X, \bar{A}) \tag{5.1.3}$$

exhibiting the infinitesimal quantomorphisms as the Lie subalgebra of vector fields on P which preserve the connection.

The following proposition describes the relationship between $\text{LieAlg}_{\text{At}}(X, \bar{A})$ and $\mathfrak{atiyah}(X, \omega)$, which one can think of as an extension of Thm. 4.2.2 for the $n = 1$ case.

Proposition 5.1.3. *There exists a natural Lie algebra isomorphism*

$$\psi: \mathfrak{atiyah}(X, \omega) \xrightarrow{\cong} \text{LieAlg}_{\text{At}}(X, \bar{A})$$

such that the following diagram commutes

$$\begin{array}{ccc} \mathfrak{atiyah}(X, \omega) & \xrightarrow{\psi} & \text{LieAlg}_{\text{At}}(X, \bar{A}) \\ \uparrow & & \uparrow \\ L_\infty(X, \omega) & \xrightarrow{f} & \text{dgLie}_{\text{Qu}}(X, \bar{A}) \end{array}$$

where $f: L_\infty(M, \omega) \xrightarrow{\cong} \text{dgLie}_{\text{Qu}}(X, \bar{A})$ is the isomorphism of Lie algebras given in Thm. 4.2.2, and the vertical morphisms are the inclusions (5.1.1) and (5.1.3).

Proof. It follows from Prop. A.1 that the isomorphism $f: L_\infty(M, \omega) \xrightarrow{\cong} \text{dgLie}_{\text{Qu}}(X, \bar{A})$ is $f(v + c) = v - c|_{U_\alpha} + \iota_v A^1$. Hence, we define $\psi: \text{atiyah}(X, \omega) \rightarrow \text{LieAlg}_{\text{At}}(X, \bar{A})$ to be

$$\psi(v + c) = v - c|_{U_\alpha} + \iota_v A^1. \quad (5.1.4)$$

Note that if $\mathcal{L}_v A^0 = \delta \bar{\theta}$, then $\delta(\bar{\theta} + \iota_v A^1) = 0$. Hence ψ is an isomorphism of vector spaces. The fact that ψ preserves the Lie brackets follows from the equalities

$$\mathcal{L}_{v_1} \iota_{v_2} A^1 - \mathcal{L}_{v_2} \iota_{v_1} A^1 = \iota_{[v_1, v_2]} A^1 + \iota_{v_2} \iota_{v_1} dA^1 = \iota_{[v_1, v_2]} A^1 + \iota_{v_1 \wedge v_2} \omega.$$

□

Remark 5.1.4. Note that the isomorphism ψ (5.1.4) uses the connection A to horizontally lift a vector field on M to a vector field on P in the standard way.

5.2 The $n = 2$ case

Here (X, ω) is a pre-2-plectic manifold. A prequantization ∇ of (X, ω) is a $U(1)$ -bundle gerbe (or principal $U(1)$ 2-bundle) over X equipped with a 2-connection.

In addition to the Lie 2-algebra of local observables $L_\infty(X, \omega)$, there are two other Lie 2-algebras one can build directly from any closed 3-form ω . It seems that the first of these has not appeared previously in the literature, while the second one originates in Roytenberg and Weinstein's work on Courant algebroids [25]. (The 2-term truncation we use here is due to subsequent work by Roytenberg [24].)

Definition/Proposition 5.2.1. Let ω be a 2-plectic structure on X . The *Atiyah Lie 2-algebra* $\text{atiyah}(\omega)$ is the graded vector space

$$\begin{aligned} \text{atiyah}(X, \omega)^0 &= \mathfrak{X}(X) \\ \text{atiyah}(X, \omega)^1 &= \Omega^0(X); \end{aligned}$$

endowed with the brackets

$$\begin{aligned} \llbracket \eta \rrbracket_1^{\text{a}} &= 0 \\ \llbracket v_1, v_2 \rrbracket_2^{\text{a}} &= [v_1, v_2] \\ \llbracket v, \eta \rrbracket_2^{\text{a}} &= \mathcal{L}_v \eta \\ \llbracket v_1, v_2, v_3 \rrbracket_3^{\text{a}} &= -\iota_{v_1 \wedge v_2 \wedge v_3} \omega \end{aligned}$$

(with all other brackets zero by degree reasons).

Definition/Proposition 5.2.2. Let ω be a 2-plectic structure on X . The *Courant Lie 2-algebra* $\text{courant}(\omega)$ is the graded vector space

$$\begin{aligned} \text{courant}(X, \omega)^0 &= \mathfrak{X}(X) \oplus \Omega^1(X) \\ \text{courant}(X, \omega)^1 &= \Omega^0(X); \end{aligned}$$

endowed with the brackets

$$\begin{aligned} \llbracket \eta \rrbracket_1^{\text{c}} &= d\eta \\ \llbracket v_1 + \theta_1, v_2 + \theta_2 \rrbracket_2^{\text{c}} &= [v_1, v_2] + \mathcal{L}_{v_1} \theta_2 - \mathcal{L}_{v_2} \theta_1 \\ &\quad - \frac{1}{2} d(\iota_{v_1} \theta_2 - \iota_{v_2} \theta_1) - \iota_{v_1 \wedge v_2} \omega \\ \llbracket v + \theta, \eta \rrbracket_2^{\text{c}} &= \frac{1}{2} \iota_v d\eta \\ \llbracket v_1 + \theta_1, v_2 + \theta_2, v_3 + \theta_3 \rrbracket_3^{\text{c}} &= -\frac{1}{6} \left(\langle \llbracket v_1 + \theta_1, v_2 + \theta_2 \rrbracket_2^{\text{c}}, v_3 + \theta_3 \rangle + \text{cyc. perm.} \right) \end{aligned}$$

where \langle , \rangle is the natural symmetric pairing between sections of $T^*X \oplus TX$, i.e.,

$$\langle v_1 + \theta_1, v_2 + \theta_2 \rangle := \iota_{v_1}\theta_2 + \iota_{v_2}\theta_1$$

(and with all other brackets zero by degree reasons).

The relationship between these Lie 2-algebras is given by the next proposition.

Proposition 5.2.3. *There exists a natural sequence of L_∞ morphisms*

$$L_\infty(X, \omega) \xrightarrow{\phi} \mathbf{courant}(X, \omega) \xrightarrow{\psi} \mathbf{atiyah}(X, \omega),$$

where the nontrivial components of the morphism ϕ are

$$\begin{aligned} \phi_1(v + \theta) &= v + \theta \\ \phi_1(\eta) &= \eta \\ \phi_2(v_1 + \theta_1, v_2 + \theta_2) &= -\frac{1}{2}(\iota_{v_1}\theta_2 - \iota_{v_2}\theta_1) \end{aligned}$$

and the nontrivial components of the morphism ψ are

$$\begin{aligned} \psi_1(v + \theta) &= v \\ \psi_1(\eta) &= \eta \\ \psi_2(v_1 + \theta_1, v_2 + \theta_2) &= -\frac{1}{2}(\iota_{v_1}\theta_2 - \iota_{v_2}\theta_1) \end{aligned}$$

Proof. The fact that ϕ is an L_∞ -morphism is the content of Thm. 7.1 in [23]. To show ψ is a L_∞ morphism, we first perform several straightforward computations using the Cartan calculus in order to obtain the following equalities:

$$\begin{aligned} \psi_2(d\eta, v + \theta) &= \psi_1(\llbracket \eta, v + \theta \rrbracket^c) - \llbracket \psi_1(\eta), \psi_1(v + \theta) \rrbracket_2^a, \\ \llbracket v_1 + \theta_1, v_2 + \theta_2, v_3 + \theta_3 \rrbracket_3^c &= -\frac{1}{4}(\iota_{v_3}\mathcal{L}_{v_1}\theta_2 - \iota_{v_3}\mathcal{L}_{v_2}\theta_1 + \text{cyc. perm.}) + \frac{1}{2}\iota_{v_1 \wedge v_2 \wedge v_3}\omega, \\ \psi_2(\llbracket v_1 + \theta_1, v_2 + \theta_2 \rrbracket_2^c, v_3 + \theta_3) + \text{cyc. perm.} &= -\frac{1}{4}(\iota_{v_3}\mathcal{L}_{v_1}\theta_2 - \iota_{v_3}\mathcal{L}_{v_2}\theta_1 + \text{cyc. perm.}) \\ &\quad - (\iota_{v_1 \wedge v_2}d\theta_3 + \text{cyc. perm.}) - \frac{3}{2}\iota_{v_1 \wedge v_2 \wedge v_3}\omega, \\ \llbracket \psi_1(v_1 + \theta_1), \psi_2(v_2 + \theta_2, v_3 + \theta_3) \rrbracket_2^a + \text{cyc. perm.} &= -\frac{1}{2}(\iota_{v_3}\mathcal{L}_{v_1}\theta_2 - \iota_{v_3}\mathcal{L}_{v_2}\theta_1 + \text{cyc. perm.}) \\ &\quad - (\iota_{v_1 \wedge v_2}d\theta_3 + \text{cyc. perm.}). \end{aligned}$$

We then use the above to verify that the equalities given in [1, Def. 34] are satisfied. \square

If (X, ω) is prequantized, then we represent the prequantum 2-bundle $\nabla : X \rightarrow \mathbf{B}^2U(1)_{\text{conn}}$ with a Čech-Deligne 2-cocycle, and obtain dg Lie algebras which we think of as modeling the previously discussed L_∞ -algebras $\mathbf{LieAtiyah}(\nabla)$ and $\mathbf{LieCourant}(\nabla)$. In what follows, $\Omega^{\leq 0}$ and $\Omega^{\leq 1}$ denote the cochain complexes of sheaves

$$\Omega^0(-) \rightarrow 0 \rightarrow 0 \rightarrow 0 \rightarrow \cdots, \quad \Omega^0(-) \xrightarrow{d} \Omega^1(-) \rightarrow 0 \rightarrow 0 \rightarrow \cdots,$$

respectively, with both having $\Omega^0(-)$ in degree zero.

Definition/Proposition 5.2.4. If $\bar{A} = A^2 + A^1 + A^0$ is a Čech-Deligne 2-cocycle on X relative to some cover \mathcal{U} , then we denote by $\mathbf{dgLie}_{\text{At}}(X, \bar{A})$ and $\mathbf{dgLie}_{\text{Cou}}(X, \bar{A})$ the dg-Lie algebras whose underlying complexes are

$$\begin{aligned} \mathbf{dgLie}_{\text{At}}(X, \bar{A})^0 &= \{v + \bar{\theta} \in \mathfrak{X}(X) \oplus \text{Tot}^1(\mathcal{U}, \Omega^{\leq 0}) \mid \mathcal{L}_v A^0 = d_{\text{Tot}}\bar{\theta}\} \\ \mathbf{dgLie}_{\text{At}}(X, \bar{A})^1 &= \text{Tot}^0(\mathcal{U}, \Omega^{\leq 0}) \end{aligned}$$

and

$$\begin{aligned} \mathrm{dgLie}_{\mathrm{Cou}}(X, \bar{A})^0 &= \{v + \bar{\theta} \in \mathfrak{X}(X) \oplus \mathrm{Tot}^1(\mathcal{U}, \Omega^{\leq 1}) \mid \mathcal{L}_v(A^1 + A^0) = d_{\mathrm{Tot}}\bar{\theta}\} \\ \mathrm{dgLie}_{\mathrm{Cou}}(X, \bar{A})^1 &= \mathrm{Tot}^0(\mathcal{U}, \Omega^{\leq 1}); \end{aligned}$$

both equipped with the differential $0 \oplus d_{\mathrm{Tot}}$, and whose graded Lie brackets are (for both cases)

$$\begin{aligned} \llbracket v_1 + \bar{\theta}_1, v_2 + \bar{\theta}_2 \rrbracket &= [v_1, v_2] + \mathcal{L}_{v_1}\bar{\theta}_2 - \mathcal{L}_{v_2}\bar{\theta}_1 \\ \llbracket v + \bar{\theta}, \bar{\eta} \rrbracket &= -\llbracket \bar{\eta}, v + \bar{\theta} \rrbracket = \mathcal{L}_v\bar{\eta} \\ \llbracket \bar{\eta}, \bar{\eta} \rrbracket &= 0. \end{aligned} \tag{5.2.1}$$

The dg Lie algebra $\mathrm{dgLie}_{\mathrm{At}}(X, \bar{A})$ was constructed by Collier [9, Def. 6.11, Thm. 8.18], and he rigorously proved that its degree zero elements correspond to infinitesimal autoequivalences of the $U(1)$ 2-bundle represented by the Čech 2-cocycle A^0 . He also constructed $\mathrm{dgLie}_{\mathrm{Cou}}(X, \bar{A})$ and proved that its degree zero elements are the infinitesimal autoequivalences the $U(1)$ 2-bundle equipped with a connective structure represented by the truncated Čech-Deligne 2-cocycle $A^1 + A^0$ [9, Def. 10.38, Prop. 10.48].

There is an obvious map of dg Lie algebras $\mathrm{dgLie}_{\mathrm{Cou}}(X, \bar{A}) \xrightarrow{p} \mathrm{dgLie}_{\mathrm{At}}(X, \bar{A})$, which in degree zero forgets the $\check{C}^0(\mathcal{U}, \Omega^1)$ component. It is also clear that the dg Lie algebra $\mathrm{dgLie}(X, \bar{A})$ of infinitesimal quantomorphisms (Def/Prop. 4.2.1) embeds into $\mathrm{dgLie}_{\mathrm{Cou}}(X, \bar{A})$. Hence the next result follows automatically by construction.

Proposition 5.2.5. *There is a natural sequence of dg Lie algebras*

$$\mathrm{dgLie}_{\mathrm{Qu}}(X, \bar{A}) \xrightarrow{i} \mathrm{dgLie}_{\mathrm{Cou}}(X, \bar{A}) \xrightarrow{p} \mathrm{dgLie}_{\mathrm{At}}(X, \bar{A})$$

which we interpret as modeling the sequence (5.0.2).

In [9, Theorem 12.50], Collier constructed a weak equivalence of Lie 2-algebras between a local Čech description of the Courant Lie 2-algebra (5.2.2) and the dg Lie algebra $\mathrm{dgLie}_{\mathrm{Cou}}(X, \bar{A})$. We conclude with the following proposition which strengthens this result by incorporating our Thm. 4.2.2 and Prop. 5.2.3. It can also be viewed as the higher analog of Prop. 5.1.3.

Proposition 5.2.6. *If (X, ω) is a prequantized 2-plectic manifold, then there exist natural weak equivalences of Lie 2-algebras*

$$f^a: \mathrm{atiyah}(X, \omega) \xrightarrow{\sim} \mathrm{dgLie}_{\mathrm{At}}(X, \bar{A})$$

and

$$f^c: \mathrm{courant}(X, \omega) \xrightarrow{\sim} \mathrm{dgLie}_{\mathrm{Cou}}(X, \bar{A})$$

such that the following diagram in $L_\infty\mathrm{Alg}$ (strictly) commutes:

$$\begin{array}{ccc} \mathrm{atiyah}(X, \omega) & \xrightarrow{f^a} & \mathrm{dgLie}_{\mathrm{At}}(X, \bar{A}) \\ \psi \uparrow & & \uparrow p \\ \mathrm{courant}(X, \omega) & \xrightarrow{f^c} & \mathrm{dgLie}_{\mathrm{Cou}}(X, \bar{A}) \\ \phi \uparrow & & \uparrow i \\ L_\infty(X, \omega) & \xrightarrow{f} & \mathrm{dgLie}_{\mathrm{Qu}}(X, \bar{A}) \end{array}$$

where $f: L_\infty(M, \omega) \xrightarrow{\sim} \mathrm{dgLie}_{\mathrm{Qu}}(X, \bar{A})$ is the weak equivalence given in Thm. 4.2.2, and the vertical morphisms are those given in Prop. 5.2.3 and Prop. 5.2.5.

Proof. In terms of the notation above, Prop. A.1 and Eqs. A.9 imply that the weak equivalence $f: L_\infty(M, \omega) \xrightarrow{\sim} \text{dgLie}_{\text{Qu}}(X, \bar{A})$ has non-trivial components

$$\begin{aligned} f_1(v + \theta) &= v - \theta + \iota_v(A^2 - A^1) \\ f_1(\eta) &= -\eta \\ f_2(v_1 + \theta_1, v_2 + \theta_2) &= \iota_{v_1}\theta_2 - \iota_{v_2}\theta_1 + \iota_{v_1 \wedge v_2}A^2. \end{aligned}$$

(Above we have suppressed the restriction of global forms on X to open sets $U_\alpha \in \mathcal{U}$.) Hence, we define the non-trivial components of f^c to be

$$\begin{aligned} f_1^c(v + \theta) &= v - \theta + \iota_v(A^2 - A^1) \\ f_1^c(\eta) &= -\eta \\ f_2^c(v_1 + \theta_1, v_2 + \theta_2) &= \frac{1}{2}(\iota_{v_1}\theta_2 - \iota_{v_2}\theta_1) + \iota_{v_1 \wedge v_2}A^2. \end{aligned}$$

Similarly, we define f^a by

$$\begin{aligned} f_1^a(v) &= v - \iota_v A^1 \\ f_1^a(\eta) &= -\eta \\ f_2^a(v_1, v_2) &= \iota_{v_1 \wedge v_2}A^2. \end{aligned}$$

Note that if $v + \bar{\theta}$ is a degree 0 element of $\text{dgLie}_{\text{Cou}}(X, \bar{A})$, then $d_{\text{Tot}}(\bar{\theta} - \iota_v(A^2 - A^1)) = 0$. Similarly, if $v + \bar{\theta}$ is a degree 0 element of $\text{dgLie}_{\text{At}}(X, \bar{A})$, then $d_{\text{Tot}}(\bar{\theta} + \iota_v A^1) = 0$. It then follows from the Poincaré Lemma that both f_1^c and f_1^a are quasi-isomorphisms of chain complexes.

It follows immediately from the definitions (Prop. 5.2.3 and Prop. 5.2.5) that $f_1^c \circ \phi_1 = i \circ f_1$ and $f_1^a \circ \psi_1 = p \circ f_1^c$. Simple calculations show that the following equations hold:

$$\begin{aligned} (f^c \circ \phi)_2(v_1 + \theta_1, v_2 + \theta_2) &:= f_1^c \phi_2(v_1 + \theta_1, v_2 + \theta_2) + f_2^c(\phi_1(v_1 + \theta_1), \phi_1(v_2 + \theta_2)) \\ &= i \circ f_2(v_1 + \theta_1, v_2 + \theta_2), \\ (f^a \circ \psi)_2(v_1 + \theta_1, v_2 + \theta_2) &:= f_1^a \psi_2(v_1 + \theta_1, v_2 + \theta_2) + f_2^a(\psi_1(v_1 + \theta_1), \psi_1(v_2 + \theta_2)) \\ &= p \circ f_2^c(v_1 + \theta_1, v_2 + \theta_2). \end{aligned}$$

Hence the above diagram commutes.

Finally, to show f^c and f^a are indeed L_∞ morphisms, we must check that (3.1.1) holds for the $n = 2$ case (or equivalently the equalities given in [1, Def. 34]). Using the identities from Sec. 1 and the cocycle equation for \bar{A} , we obtain the following equalities:

$$\begin{aligned} \llbracket f_1^c(v_1 + \theta_1), f_1^c(v_2 + \theta_2) \rrbracket^{\text{Cou}} &= [v_1, v_2] - \mathcal{L}_{v_1}\theta_2 + \mathcal{L}_{v_2}\theta_1 + \iota_{[v_1, v_2]}(A^2 - A^1) \\ &\quad + \iota_{v_1 \wedge v_2}\omega - d_{\text{Tot}}(\iota_{v_1 \wedge v_2}A^2) \\ f_2^c(\llbracket v_1 + \theta_1, v_2 + \theta_2 \rrbracket_2^c, v_3 + \theta_3) + \text{cyc. perm.} &= \frac{1}{4}(\iota_{v_3}\mathcal{L}_{v_1}\theta_2 - \iota_{v_3}\mathcal{L}_{v_2}\theta_1 + \text{cyc. perm.}) \\ &\quad + (\iota_{v_1 \wedge v_2}d\theta_3 + \iota_{[v_1, v_2] \wedge v_3}A^2 + \text{cyc. perm.}) + \frac{3}{2}\iota_{v_1 \wedge v_2 \wedge v_3}\omega \\ \llbracket f_1^c(v_1 + \theta_1), f_2^c(v_2 + \theta_2, v_3 + \theta_3) \rrbracket_2^{\text{Cou}} + \text{cyc. perm.} &= \frac{1}{2}(\iota_{v_3}\mathcal{L}_{v_1}\theta_2 - \iota_{v_3}\mathcal{L}_{v_2}\theta_1 + \text{cyc. perm.}) \\ &\quad + (\iota_{v_1 \wedge v_2}d\theta_3 + 2\iota_{[v_1, v_2] \wedge v_3}A^2 + \iota_{v_2 \wedge v_3}\mathcal{L}_{v_1}A^2 + \text{cyc. perm.}). \end{aligned}$$

And similarly for f^a :

$$\begin{aligned} \llbracket f_1^a(v_1), f_1^a(v_2) \rrbracket^{\text{At}} &= [v_1, v_2] - \iota_{[v_1, v_2]}A^1 - d_{\text{Tot}}(\iota_{v_1 \wedge v_2}A^2) \\ f_2^a(\llbracket v_1, v_2 \rrbracket_2^a, v_3) + \text{cyc. perm.} &= \iota_{[v_1, v_2] \wedge v_3}A^2 + \text{cyc. perm.} \\ \llbracket f_1^a(v_1), f_2^a(v_2, v_3) \rrbracket^{\text{At}} + \text{cyc. perm.} &= (2\iota_{[v_1, v_2] \wedge v_3} + \iota_{v_2 \wedge v_3}\mathcal{L}_{v_1})A^2 + \text{cyc. perm.} \end{aligned}$$

Using these in conjunction with Lemma 3.2.1, it is easy to verify that f^c and f^a are L_∞ morphisms. \square

A An explicit weak equivalence between $L_\infty(X, \omega)$ and $\text{dgLie}_{\text{Qu}}(X, \bar{A})$

In this section, we prove Thm. 4.2.2. Namely, given a pre n -plectic manifold (X, ω) and a prequantization presented by a Čech-Deligne n -cocycle \bar{A} with respect to a cover $\mathcal{U} = \{U_\alpha\}$ of X , we shall construct a weak equivalence (L_∞ -quasi-isomorphism)

$$L_\infty(X, \omega) \xrightarrow{\sim} \text{dgLie}_{\text{Qu}}(X, \bar{A}).$$

The proof is a long, but straightforward, calculation which we present below in several steps.

A quasi-isomorphism of chain complexes

Recall that the underlying complex (L, l_1) of the Lie n -algebra $L_\infty(X, \omega)$ from definition 3.2.3 is:

$$\begin{aligned} L_0 &= \{v + H \in \mathfrak{X}_{\text{Ham}}(X) \oplus \Omega^{n-1}(X) \mid dH = -\iota_v \omega\} \\ L_i &= \Omega^{n-1-i}(X) \quad i \geq 1 \end{aligned}$$

with differential

$$L_{n-1} \xrightarrow{d} L_{n-2} \xrightarrow{d} \cdots \xrightarrow{d} L_1 \xrightarrow{0 \oplus d} L_0$$

while the underlying complex (L', l'_1) of the dg Lie algebra $\text{dgLie}_{\text{Qu}}(X, \bar{A})$ from definition 4.2.1 is

$$\begin{aligned} L'_0 &= \{v + \bar{\theta} \in \mathfrak{X}(X) \oplus \text{Tot}^{n-1}(\mathcal{U}, \Omega) \mid \mathcal{L}_v \bar{A} = d_{\text{Tot}} \bar{\theta}\} \\ L'_i &= \text{Tot}^{n-1-i}(\mathcal{U}, \Omega) \quad i \geq 1, \end{aligned}$$

with differential

$$L'_{n-1} \xrightarrow{d_{\text{Tot}}} L'_{n-2} \xrightarrow{d_{\text{Tot}}} \cdots \xrightarrow{d_{\text{Tot}}} L'_1 \xrightarrow{0 \oplus d_{\text{Tot}}} L'_0.$$

We denote by

$$\text{res}: \Omega^\bullet(X) \rightarrow \check{C}^0(\mathcal{U}, \Omega^\bullet)$$

the restriction map

$$\text{res}(\theta)_\alpha = \theta|_{U_\alpha} \in \Omega^\bullet(U_\alpha).$$

For $\bar{A} = \sum_{i=0}^n A^{n-i}$ a Čech-Deligne cocycle, we define for all $m \geq 1$:

$$\bar{A}(m) := \sum_{i=0}^m (-1)^{mi} A^{n-i}. \tag{A.2}$$

Proposition A.1. *The linear map*

$$f_1: L \rightarrow L'$$

defined by

$$f_1(x) = \begin{cases} v - \text{res}(\theta) + \iota_v \bar{A}(1) & \forall x = v + \theta \in L_0 \\ -\text{res}(x) & \forall x \in L_{i>0} \end{cases}$$

is a quasi-isomorphism of chain complexes.

Proof. It is clear from the definition that f_1 is a chain map. Since \bar{A} is a Čech-Deligne cocycle, and since the interior product ι_v commutes with the Čech differential, it is not hard to see that

$$d_{\text{Tot}} \iota_v \bar{A}(1) = d \iota_v A^n + \sum_{i=1}^n \mathcal{L}_v A^{n-i}.$$

This implies that $v + \bar{\theta} \in L'_0$ if and only if

$$d_{\text{Tot}}(\bar{\theta} - \iota_v \bar{A}(1)) = \text{res}(\iota_v \omega).$$

Let \tilde{L} be the complex whose underlying graded vector space is

$$\begin{aligned}\tilde{L}_0 &= \{v + \bar{\eta} \in \mathfrak{X}(X) \oplus \text{Tot}^{n-1}(\mathcal{U}, \Omega) \mid d_{\text{Tot}}\bar{\eta} = \text{res}(\iota_v\omega)\} \\ \tilde{L}_i &= L'_i \quad i > 0,\end{aligned}$$

and whose differential is $\tilde{l}_1 = l'_1$, the same differential as on L' . The chain map f_1 then is equal to the composition:

$$L \xrightarrow{r} \tilde{L} \xrightarrow{\phi} L',$$

where

$$r(x) = \begin{cases} v - \text{res}(\theta) & \forall x = v + \theta \in L_0 \\ -\text{res}(x) & \forall x \in L_{i>0} \end{cases}$$

and

$$\phi(x) = \begin{cases} v + \bar{\eta} + \iota_v \bar{A}(1) & \forall x = v + \bar{\eta} \in \tilde{L}_0 \\ x & \forall x \in \tilde{L}_{i>0}. \end{cases}$$

Note that ϕ is a isomorphism of complexes.

Next, let $\{\rho_\alpha\}$ be a partition of unity subordinate to the cover $\mathcal{U} = \{U_\alpha\}$. Define a map $K: \check{C}^i(\mathcal{U}, \Omega^j) \rightarrow \check{C}^{i-1}(\mathcal{U}, \Omega^j)$ to be

$$(K\theta)_{\alpha_0, \dots, \alpha_{i-1}} = \sum_{\alpha} \rho_\alpha \theta_{\alpha, \alpha_0, \dots, \alpha_{i-1}},$$

and let $D'': \check{C}^i(\mathcal{U}, \Omega^j) \rightarrow \check{C}^i(\mathcal{U}, \Omega^{j+1})$ be the “signed” de Rham differential $D''\theta = (-1)^i d\theta$. Then there exists a chain map [4, Prop. 9.5]

$$j: \text{Tot}^\bullet(\mathcal{U}, \Omega) \rightarrow \Omega^\bullet(X)$$

such that

$$\begin{aligned}j \circ \text{res} &= \text{id}_{\Omega^\bullet(X)} \\ \text{id}_{\text{Tot}^\bullet(\mathcal{U}, \Omega)} - \text{res} \circ j &= d_{\text{Tot}}H - Hd_{\text{Tot}},\end{aligned}\tag{A.3}$$

where $H: \text{Tot}^\bullet(\mathcal{U}, \Omega) \rightarrow \text{Tot}^\bullet(\mathcal{U}, \Omega)$ is the following chain homotopy: If $\bar{\theta} = \sum_{i=0}^m \theta^{m-i} \in \text{Tot}^m(\mathcal{U}, \Omega)$ with $\theta^{m-i} \in \check{C}^i(\mathcal{U}, \Omega^{m-i})$ then

$$H(\bar{\theta}) = \sum_{i=0}^{m-1} (H\bar{\theta})_i$$

where

$$(H\bar{\theta})_i = \sum_{j=i+1}^m K \circ \underbrace{(-D''K) \circ (-D''K) \circ \dots \circ (-D''K)}_{j-(i+1)} \theta^{m-j} \in \check{C}^i(\mathcal{U}, \Omega^{m-1-i}).\tag{A.4}$$

Hence, the restriction map res is a quasi-isomorphism between the de Rham and Čech-de Rham complexes, and j is its homotopy inverse.

Let $\tilde{j}: \tilde{L} \rightarrow L$ to be the chain map

$$\tilde{j}(x) = \begin{cases} v - j(\bar{\eta}) & \forall x = v + \bar{\eta} \in \tilde{L}_0 \\ -j(x) & \forall x \in \tilde{L}_{i>0}, \end{cases}$$

Note that \tilde{j} is well-defined on degree 0 elements since $d_{\text{Tot}}\bar{\eta} = \text{res}(\iota_v\omega)$. Let $\tilde{H}: \tilde{L} \rightarrow \tilde{L}$ be the degree 1 map

$$\tilde{H}(x) = \begin{cases} H(\bar{\eta}) & \forall x = v + \bar{\eta} \in \tilde{L}_0 \\ H(x) & \forall x \in \tilde{L}_{i>0}. \end{cases}$$

We now show that \tilde{H} is a chain homotopy i.e.

$$\text{id}_{\tilde{L}} - r \circ \tilde{j} = \tilde{l}_1 \tilde{H} - \tilde{H} \tilde{l}_1.$$

Since (A.3) holds, it follows from the definition of the maps and differentials that we just need to check Eq. A on degree 0 elements:

$$v + \bar{\eta} - r \circ \tilde{j}(v + \bar{\eta}) = 0 + d_{\text{Tot}} H(\bar{\eta}) \quad (\text{A.5})$$

for all $v + \bar{\eta} \in \tilde{L}_0$. Since we have the equality

$$d_{\text{Tot}} \bar{\eta} = \text{res}(\iota_v \omega) \in \check{C}^0(\mathcal{U}, \Omega^n),$$

it follows from the definition of H (A.4) that

$$H(d_{\text{Tot}} \bar{\eta}) = 0.$$

So (A.3) implies that (A.5) is satisfied as well. Therefore r is a quasi-isomorphism, and hence f_1 is a quasi-isomorphism. \square

The goal of the remainder of the appendix is to show the map $f_1: L \rightarrow L'$ lifts to an L_∞ -quasi-isomorphism between $L_\infty(X, \omega)$ and $\text{dgLie}_{\text{Qu}}(X, \bar{A})$.

Notation

We use the following conventions to help simplify calculations.

- As above, (L, l_1) denotes the underlying complex of $L_\infty(X, \omega)$, and l_2, \dots, l_{n+1} denotes the k -ary brackets of the L_∞ structure given in Def. 3.2.3. (L', l'_1) denotes, as above, the underlying complex of the dgla $\text{dgLie}_{\text{Qu}}(X, \bar{A})$, and $l'_2 = \llbracket \cdot, \cdot \rrbracket$ denotes its Lie bracket given in Def. 4.2.1.

Elements of arbitrary degree in L (resp. L') will be denoted as x_1, x_2, \dots (resp. $\bar{x}_1, \bar{x}_2, \dots$) where

$$x_i := v_i + \theta_i \quad (\text{resp. } \bar{x}_i := v_i + \bar{\theta}_i). \quad (\text{A.6})$$

It is understood for the vector field v_i that we set $v_i = 0$ if $|x_i| > 0$ (resp. $|\bar{x}_i| > 0$).

So for example, for any $x_1, \dots, x_k \in L$ and any $\bar{x}_1, \bar{x}_2 \in L'$ the following equalities hold:

$$\begin{aligned} l_2(x_1, x_2) &= [v_1, v_2] + \iota_{v_1 \wedge v_2} \omega, \\ l_{k \geq 3}(x_1, \dots, x_k) &= -(-1)^{\binom{k+1}{2}} \iota_{v_1 \wedge \dots \wedge v_k} \omega, \\ \llbracket \bar{x}_1, \bar{x}_2 \rrbracket &= [v_1, v_2] + \mathcal{L}_{v_1} \bar{\theta}_2 - \mathcal{L}_{v_2} \bar{\theta}_1 \end{aligned} \quad (\text{A.7})$$

and, also for the map f_1 in Prop. A.1 we have:

$$f_1(x_1) = v_1 - \text{res}(\theta_1) + \iota(v_1) \bar{A}(1).$$

- For all $m \geq 2$ we define a map

$$S_m: L^{\otimes m} \rightarrow L$$

where

$$S_m(x_1, \dots, x_m) = \sum_{i=1}^m (-1)^i \iota_{v_1 \wedge \dots \wedge \bar{v}_i \wedge \dots \wedge v_m} \theta_i. \quad (\text{A.8})$$

It is clear from our above notation that $S_m(x_1, \dots, x_m) = 0$ if two or more arguments have degree > 0 . If all arguments have degree 0, then each θ_i is an $(n-1)$ -form, hence

$$S_m(x_1, \dots, x_m) \in \Omega^{n-1-(m-1)}(X) = L_{m-1}.$$

If $x_i \in L_{k>0} = \Omega^{n-1-k}(X)$ and $|x_j| = 0$ for all $j \neq i$ then

$$S_m(x_1, \dots, x_m) = (-1)^i \iota_{v_1 \wedge \dots \wedge \widehat{v}_i \wedge \dots \wedge v_m} \theta_i \in \Omega^{n-1-(k+m-1)}(X) = L_{k+m-1}.$$

Hence, the above implies that S_m is a graded skew-symmetric map of degree $m-1$ and $S_{m>n} = 0$.

- For all $m \geq 1$ we define the linear maps

$$f_m : L^{\otimes m} \rightarrow L',$$

where $f_1 : L \rightarrow L'$ is the chain map given in Prop. A.1, and

$$f_{2 \leq m \leq n}(x_1, \dots, x_m) = -(-1)^{\binom{m+1}{2}} (\text{res} \circ S_m(x_1, \dots, x_m) + \iota_{v_1 \wedge \dots \wedge v_m} \bar{A}(m)), \quad (\text{A.9})$$

and

$$f_{m>n} = 0.$$

Note that the second term on the right-hand side of (A.9) vanishes if any x_i has degree > 0 . It follows from (A.8) that each f_m is graded skew-symmetric with $|f_m| = m-1$. Below, we will suppress the restriction map in the definition.

The above are the structure maps we will use to lift f_1 to an L_∞ -morphism.

- Finally, we use the structure maps (A.9) and brackets (A.7) to define the following linear maps: For all $m \geq 1$

$$I_{(1)}^m, I_{(2)}^m, I_{(3)}^m : L^{\otimes m} \rightarrow L'$$

where $I_2^1 = I_3^{m<3} = 0$ and

$$\begin{aligned} I_{(1)}^m(x_1, \dots, x_m) &= \sum_{\sigma \in \text{Sh}(1, m-1)} \chi(\sigma) (-1)^m f_m(l_1(x_{\sigma(1)}), \dots, x_{\sigma(m)}), \\ I_{(2)}^{m \geq 2}(x_1, \dots, x_m) &= - \sum_{\sigma \in \text{Sh}(2, m-2)} \chi(\sigma) f_{m-1}(l_2(x_{\sigma(1)}, x_{\sigma(2)}), \dots, x_{\sigma(m)}), \\ I_{(3)}^{m \geq 3}(x_1, \dots, x_m) &= \sum_{k=3}^m \sum_{\sigma \in \text{Sh}(k, m-k)} \chi(\sigma) (-1)^{k(m-k)+1} f_{m+1-k}(l_k(x_{\sigma(1)}, \dots, x_{\sigma(k)}), \dots, x_{\sigma(m)}). \end{aligned} \quad (\text{A.10})$$

Above, as always, $\chi(\sigma) = (-1)^\sigma \epsilon(\sigma)$, where $\epsilon(\sigma)$ is the Koszul sign of the permutation.

We also define for all $m \geq 1$ the maps:

$$J^m : L^{\otimes m} \rightarrow L',$$

where $J^1 = 0$ and

$$\begin{aligned} J^{m \geq 2}(x_1, \dots, x_m) &= \\ & \sum_{s+t=m} \sum_{\substack{\tau \in \text{Sh}(s, m-s) \\ \tau(1) < \tau(s+1)}} \chi(\tau) (-1)^{s-1} (-1)^{(t-1) \sum_{p=1}^s |x_{\tau(p)}|} \llbracket f_s(x_{\tau(1)}, \dots, x_{\tau(s)}), f_t(x_{\tau(s+1)}, \dots, x_{\tau(m)}) \rrbracket. \end{aligned} \quad (\text{A.11})$$

A few technical lemmas

Next we present several small computational results necessary for the proof.

Lemma A.2. *For all $m \geq 2$ and $x_1, \dots, x_m \in L$, we have*

$$I_{(1)}^{m \geq 2}(x_1, \dots, x_m) = -(-1)^{\binom{m+1}{2}} (-1)^m \sum_{i=1}^m (-1)^i \iota_{v_1 \wedge \dots \wedge \widehat{v}_i \wedge \dots \wedge v_m} (l_1 \theta_i). \quad (\text{A.12})$$

Proof. Recalling the definition of $I_{(1)}^m$ (A.10), we rewrite the summation over unshuffles, and use the definition of $f_{m \geq 2}$ (A.9) to give us

$$I_{(1)}^{m \geq 2}(x_1, \dots, x_m) = -(-1)^{\binom{m+1}{2}} (-1)^m \sum_{i=1}^m (-1)^{i-1} \epsilon(\sigma(i)) S_m(l_1 x_i, x_1, \dots, \widehat{x}_i, \dots, x_m). \quad (\text{A.13})$$

In terms of our notation (A.6), the vector field associated to $l_1 x_i = l_1(v_i + \theta_i)$ is zero, hence it follows from the definition of S_m (A.8) that

$$S_m(l_1 x_i, x_1, \dots, \widehat{x}_i, \dots, x_m) = -\iota_{v_1 \wedge \dots \wedge \widehat{v}_i \wedge \dots \wedge v_m} (l_1 \theta_i).$$

Furthermore, the right-hand side above vanishes if any other x_j have degree > 0 , so any terms contributing to the sum (A.13) have $\epsilon(\sigma(i)) = 1$. Hence, the equality A.12 holds. \square

Lemma A.3. *If $x_1, x_2 \in L$, then*

$$I_{(2)}^2(x_1, x_2) = -[v_1, v_2] + \iota_{v_1 \wedge v_2} \omega - \iota_{[v_1, v_2]} \bar{A}(1),$$

and for all $m > 2$ and $x_1, \dots, x_m \in L$, the following equality holds:

$$\begin{aligned} I_{(2)}^{m > 2}(x_1, \dots, x_m) &= -(-1)^{\binom{m}{2}} \left(\binom{m}{2} \iota_{v_1 \wedge \dots \wedge v_m} \omega + \sum_{i < k} (-1)^{i+k} \iota_{[v_i, v_k] \wedge v_1 \wedge \dots \wedge \widehat{v}_i \dots \widehat{v}_k \dots \wedge v_m} \bar{A}(m-1) \right) \\ &+ (-1)^{\binom{m}{2}} \left(\sum_{i < k < j} - \sum_{i < j < k} + \sum_{j < i < k} \right) (-1)^{i+k+j} \iota_{[v_i, v_k] \wedge v_1 \wedge \dots \wedge \widehat{v}_i \dots \widehat{v}_j \dots \widehat{v}_k \dots \wedge v_m} \theta_j \end{aligned} \quad (\text{A.14})$$

Proof. The $m = 2$ case follows immediately from the definitions.

For $m > 2$, let us first replace the summation

$$- \sum_{\sigma \in \text{Sh}(2, m-2)} \chi(\sigma)$$

in the definition of $I_{(2)}^m$ (A.10) with the equivalent

$$\sum_{1 \leq i < k \leq m} (-1)^{i+k} \epsilon(i, k). \quad (\text{A.15})$$

It follows from the definition of f_{m-1} (A.9) that a summand contributing to $I_{(2)}^m$ is of the form

$$\begin{aligned} f_{m-1}(l_2(x_i, x_k), x_1, \dots, \widehat{x}_i, \dots, \widehat{x}_k, \dots, x_m) &= -(-1)^{\binom{m}{2}} \left(S_{m-1}(l_2(x_i, x_k), x_1, \dots, \widehat{x}_i, \dots, \widehat{x}_k, \dots, x_m) \right. \\ &\left. + \iota_{[v_i, v_k] \wedge v_1 \wedge \dots \wedge \widehat{v}_i \dots \widehat{v}_k \dots \wedge v_m} \bar{A}(m-1) \right). \end{aligned} \quad (\text{A.16})$$

The second term on the right-hand side above vanishes if $|x_i| > 0$ for any i , hence taking the summation (A.15) of all such terms gives

$$\sum_{1 \leq i < k \leq m} (-1)^{i+k} \iota_{[v_i, v_k] \wedge v_1 \wedge \dots \wedge \widehat{v}_i \dots \widehat{v}_k \dots \wedge v_m} \bar{A}(m-1). \quad (\text{A.17})$$

Using (A.8), we rewrite the first term on the right-hand side of (A.16) as

$$\begin{aligned}
S_{m-1}(l_2(x_i, x_k), x_1, \dots, \widehat{x}_i, \dots, \widehat{x}_k, \dots, x_m) &= -\iota_{v_1 \wedge \dots \wedge \widehat{v}_i \dots \wedge \widehat{v}_k \dots \wedge v_m} \iota_{v_i \wedge v_k} \omega \\
&+ \left(-\sum_{j=1}^i + \sum_{j=i+1}^{k-1} - \sum_{j=k+1}^m \right) (-1)^j \iota_{[v_i, v_k] \wedge v_1 \dots \wedge \widehat{v}_i \dots \wedge \widehat{v}_j \dots \wedge \widehat{v}_k \dots \wedge v_m} \theta_j.
\end{aligned} \tag{A.18}$$

Our convention for interior product of multivector fields implies that

$$-\iota_{v_1 \wedge \dots \wedge \widehat{v}_i \dots \wedge \widehat{v}_k \dots \wedge v_m} \iota_{v_i \wedge v_k} \omega = (-1)^{i+k} \iota_{v_1 \wedge \dots \wedge v_m} \omega.$$

Since this term will vanish if $|x_i| > 0$ for any i , the summation (A.15) of all such terms is

$$\binom{m}{2} \iota_{v_1 \wedge \dots \wedge v_m} \omega. \tag{A.19}$$

Finally, we take the summation (A.15) over i and k of in second term on the right-hand side of (A.18). This vanishes if two or more of the x_i have degree > 0 . Hence $\epsilon(i, k) = 1$, and the summation is

$$\left(-\sum_{i < k < j} + \sum_{i < j < k} - \sum_{j < i < k} \right) (-1)^{i+k+j} \iota_{[v_i, v_k] \wedge v_1 \dots \wedge \widehat{v}_i \dots \wedge \widehat{v}_j \dots \wedge \widehat{v}_k \dots \wedge v_m} \theta_j$$

Combining the above with (A.17) and (A.19) completes the proof. \square

Lemma A.4. *For all $m \geq 3$ and $x_1, \dots, x_m \in L$, the following equality holds:*

$$I_{(3)}^{m \geq 3}(x_1, \dots, x_m) = (-1)^{\binom{m+1}{2}} (-1)^m \left(\binom{m}{2} - m + 1 \right) \iota_{v_1 \wedge \dots \wedge v_m} \omega. \tag{A.20}$$

Proof. Recalling the definition of $I_{(3)}^m$ (A.10), we consider terms

$$f_{m+1-k}(l_k(x_{\sigma(1)}, \dots, x_{\sigma(k)}), \dots, x_{\sigma(m)})$$

for $3 \leq k \leq m$ and $\sigma \in \text{Sh}(k, m-k)$. If $k = m$, then it follows directly from the definitions of f_1 and l_k (A.7) that

$$f_1(l_m(x_{\sigma(1)}, \dots, x_{\sigma(m)})) = (-1)^{\binom{m+1}{2}} \iota_{v_{\sigma(1)} \wedge \dots \wedge v_{\sigma(m)}} \omega. \tag{A.21}$$

For $k < m$, we have

$$\begin{aligned}
f_{m+1-k}(l_k(x_{\sigma(1)}, \dots, x_{\sigma(k)}), \dots, x_{\sigma(m)}) &= -(-1)^{\binom{m-k+2}{2}} S_{m+1-k}(l_k(x_{\sigma(1)}, \dots, x_{\sigma(k)}), \dots, x_{\sigma(m)}) \\
&= -(-1)^{\binom{m-k+2}{2}} (-1)^1 \iota_{v_{\sigma(k+1)} \wedge \dots \wedge v_{\sigma(m)}} l_k(x_{\sigma(1)}, \dots, x_{\sigma(k)}) \\
&= -(-1)^{\binom{k+1}{2}} (-1)^{\binom{m-k+2}{2}} \iota_{v_{\sigma(1)} \wedge \dots \wedge v_{\sigma(m)}} \omega.
\end{aligned} \tag{A.22}$$

The second-to-last equality above follows from the definition of S_m (A.8), and the fact that $|l_k| > 0$ for $k \geq 3$; the last equality above follows from the definition of l_k and our convention for the interior product of multi-vector fields.

Combining (A.21), and (A.22), with definition of $I_{(3)}^m$ gives

$$I_{(3)}^{m \geq 3}(x_1, \dots, x_m) = \sum_{k=3}^m \sum_{\sigma \in \text{Sh}(k, m-k)} \chi(\sigma) (-1)^{k(m-k)} (-1)^{\binom{k+1}{2}} (-1)^{\binom{m-k+2}{2}} \iota_{v_{\sigma(1)} \wedge \dots \wedge v_{\sigma(m)}} \omega. \tag{A.23}$$

Observe that the sum on the right-hand side above vanishes if, for any i , $|x_i| > 0$ (since, by our notation in (A.6), $v_i = 0$.) Hence, non-zero summands above have $\chi(\sigma) = (-1)^\sigma$. Since ω is skew-symmetric, reordering the vector fields will cancel this sign. The number of unshuffles appearing in the summation is

$$\#\text{Sh}(k, m-k) = \binom{m}{k},$$

therefore, summing over σ gives

$$I_{(3)}^{m \geq 3}(x_1, \dots, x_m) = \sum_{k=3}^m (-1)^{k(m-k)} (-1)^{\binom{k+1}{2}} (-1)^{\binom{m-k+2}{2}} \binom{m}{k} \iota_{v_1 \wedge \dots \wedge v_m} \omega. \quad (\text{A.24})$$

It's easy to see that

$$(-1)^{k(m-k)} (-1)^{\binom{k+1}{2}} (-1)^{\binom{m-k+2}{2}} = -(-1)^{\binom{m+1}{2}} (-1)^m (-1)^k.$$

By substituting the above sign into (A.24) and using the fact that

$$\sum_{k=0}^m \binom{m}{k} (-1)^k = 0,$$

we obtain the equality (A.20). □

Lemma A.5. *For all $m \geq 2$ and $x_1, \dots, x_m \in L$ the following equality holds:*

$$J^{m \geq 2}(x_1, \dots, x_m) = \sum_{i=1}^m (-1)^{i-1} \llbracket f_1(x_i), f_{m-1}(x_1, \dots, \widehat{x}_i, \dots, x_m) \rrbracket. \quad (\text{A.25})$$

Proof. Recalling the definition of J^m (A.11), we observe that for $m = 2$, the only unshuffle $\tau \in \text{Sh}(1, 1)$ satisfying $\tau(1) < \tau(2)$ is the identity. Hence, $J^2(x_1, x_2) = \llbracket f_1(x_1), f_1(x_2) \rrbracket$.

For the $m > 2$ case, it follows from the definition of the bracket (A.7) that the summands on the right-hand side of (A.11) vanish if both $s > 1$ and $t > 1$. Hence, we have

$$\begin{aligned} J^{m \geq 3}(x_1, \dots, x_m) &= \sum_{\substack{\tau \in \text{Sh}(1, m-1) \\ \tau(1) < \tau(2)}} \chi(\tau) (-1)^{(m-2)|x_{\tau(1)}|} \llbracket f_1(x_{\tau(1)}), f_{m-1}(x_{\tau(2)}, \dots, x_{\tau(m)}) \rrbracket \\ &+ \sum_{\substack{\tau \in \text{Sh}(m-1, 1) \\ \tau(1) < \tau(m)}} \chi(\tau) (-1)^m \llbracket f_{m-1}(x_{\tau(1)}, \dots, x_{\tau(m-1)}), f_1(x_{\tau(m)}) \rrbracket. \end{aligned} \quad (\text{A.26})$$

The only unshuffle $\tau \in \text{Sh}(1, m-1)$ satisfying $\tau(1) < \tau(2)$ is the identity. Hence, the first summation on the right-hand side is

$$\begin{aligned} \text{1st sum} &= (-1)^{(m-2)|x_1|} \llbracket f_1(x_1), f_{m-1}(x_2, \dots, x_m) \rrbracket \\ &= \llbracket f_1(x_1), f_{m-1}(x_2, \dots, x_m) \rrbracket. \end{aligned} \quad (\text{A.27})$$

The last equality above follows from the fact that the sum vanishes unless $|x_1| = 0$.

For the second summation, note that the summation over unshuffles $\tau \in \text{Sh}(m-1, 1)$ satisfying $\tau(1) < \tau(m)$ can be rewritten as

$$\text{2nd sum} = \sum_{i \geq 2} \chi(\tau(i)) (-1)^m \llbracket f_{m-1}(x_1, x_2, \dots, \widehat{x}_i, \dots, x_m), f_1(x_i) \rrbracket. \quad (\text{A.28})$$

Above $x_i = x_{\tau(m)}$, so the sign is

$$\chi(\tau(i)) = (-1)^{m-i} \epsilon(\tau(i)) = (-1)^{m-i} (-1)^{|x_i| \sum_{j>i} |x_j|}.$$

Since $m \geq 3$, it follows from the definition of the bracket and structure maps that the i th summand does not contribute to Eq. A.28 if $|x_i| > 0$. So by the antisymmetry of the bracket and the fact that $|f_1| = 0$, we have

$$\text{2nd sum} = \sum_{i \geq 2} (-1)^{i-1} \llbracket f_1(x_i), f_{m-1}(x_1, x_2, \dots, \hat{x}_i, \dots, x_m) \rrbracket. \quad (\text{A.29})$$

Combining the above (A.29) with (A.27) verifies that the equalities (A.25) hold. \square

Lemma A.6. *For all $m \geq 3$ and $x_1, \dots, x_m \in L$ the following equality holds:*

$$\begin{aligned} -(-1)^{\binom{m}{2}} J^{m>2}(x_1, \dots, x_m) &= 2 \left(\sum_{i < k < j} - \sum_{i < j < k} + \sum_{j < i < k} \right) (-1)^{i+j+k} \iota_{[v_i, v_k] \wedge v_1 \wedge \dots \wedge \hat{v}_i \dots \wedge \hat{v}_j \dots \wedge \hat{v}_k \dots \wedge v_m} \theta_j \\ &\quad + \left(\sum_{i < j} - \sum_{j < i} \right) (-1)^{i+j} \iota_{v_1 \wedge \dots \wedge \hat{v}_i \dots \wedge \hat{v}_j \dots \wedge v_m} \mathcal{L}_{v_i} \theta_j \\ &\quad - 2 \sum_{i < j} (-1)^{i+j} \iota_{[v_i, v_j] \wedge v_1 \wedge \dots \wedge \hat{v}_i \dots \wedge \hat{v}_j \dots \wedge v_m} \bar{A}(m-1) \\ &\quad - \sum_i (-1)^i \iota_{v_1 \wedge \dots \wedge \hat{v}_i \dots \wedge v_m} \mathcal{L}_{v_m} \bar{A}(m-1). \end{aligned} \quad (\text{A.30})$$

Proof. Lemma A.5 and the definitions of the bracket $\llbracket \cdot, \cdot \rrbracket$ and f_{m-1} imply that

$$-(-1)^{\binom{m}{2}} J^{m>2}(x_1, \dots, x_m) = \sum_i (-1)^{i-1} (\mathcal{L}_{v_i} S_{m-1}(x_1, \dots, \hat{x}_i, \dots, x_m) + \mathcal{L}_{v_i} \iota_{v_1 \wedge \dots \wedge \hat{v}_i \dots \wedge v_m} \bar{A}(m-1)). \quad (\text{A.31})$$

Consider the first term on the right-hand side above. The definition of S_{m-1} (A.8) implies, after rearranging indices, that

$$\begin{aligned} \sum_i (-1)^{i-1} \mathcal{L}_{v_i} S_{m-1}(x_1, \dots, \hat{x}_i, \dots, x_m) &= \sum_{i < j} (-1)^{i+j+1} \left(\mathcal{L}_{v_j} \iota_{v_1 \wedge \dots \wedge \hat{v}_i \dots \wedge \hat{v}_j \dots \wedge v_m} \theta_i \right. \\ &\quad \left. - \mathcal{L}_{v_i} \iota_{v_1 \wedge \dots \wedge \hat{v}_i \dots \wedge \hat{v}_j \dots \wedge v_m} \theta_j \right) \end{aligned} \quad (\text{A.32})$$

The commutator (1.0.3)

$$\iota_{[v_j, v_i] \wedge v_1 \wedge \dots \wedge \hat{v}_i \dots \wedge \hat{v}_j \dots \wedge v_m} = \mathcal{L}_{v_j} \iota_{v_1 \wedge \dots \wedge \hat{v}_i \dots \wedge \hat{v}_j \dots \wedge v_m} - \mathcal{L}_{v_i} \iota_{v_1 \wedge \dots \wedge \hat{v}_i \dots \wedge \hat{v}_j \dots \wedge v_m} \mathcal{L}_{v_j},$$

and the definition of the Schouten bracket (1.0.1) imply that

$$\begin{aligned} \sum_{i < j} (-1)^{i+j+1} \mathcal{L}_{v_j} \iota_{v_1 \wedge \dots \wedge \hat{v}_i \dots \wedge \hat{v}_j \dots \wedge v_m} \theta_i &= \left(\sum_{k < i < j} - \sum_{i < k < j} + \sum_{i < j < k} \right) (-1)^{i+j+k} \\ &\quad \iota_{[v_j, v_k] \wedge v_1 \wedge \dots \wedge \hat{v}_i \dots \wedge \hat{v}_j \dots \wedge \hat{v}_k \dots \wedge v_m} \theta_i + \sum_{i < j} (-1)^{i+j+1} \iota_{v_1 \wedge \dots \wedge \hat{v}_i \dots \wedge \hat{v}_j \dots \wedge v_m} \mathcal{L}_{v_j} \theta_i \\ &= \left(- \sum_{i < j < k} + 2 \sum_{j < i < k} \right) (-1)^{i+j+k} \iota_{[v_i, v_k] \wedge v_1 \wedge \dots \wedge \hat{v}_i \dots \wedge \hat{v}_j \dots \wedge \hat{v}_k \dots \wedge v_m} \theta_j \\ &\quad + \sum_{j < i} (-1)^{i+j+1} \iota_{v_1 \wedge \dots \wedge \hat{v}_i \dots \wedge \hat{v}_j \dots \wedge v_m} \mathcal{L}_{v_i} \theta_j. \end{aligned} \quad (\text{A.33})$$

The last equality above follows from the skew-symmetry of the bracket $[v_j, v_k] = -[v_k, v_j]$ and index re-labeling. A similar calculation performed on the second term on the right-hand side of (A.32) yields

$$\begin{aligned} - \sum_{i < j} (-1)^{i+j+1} \mathcal{L}_{v_i} \iota_{v_1 \wedge \dots \widehat{v}_i \dots \widehat{v}_j \dots \wedge v_m} \theta_j &= - \left(-2 \sum_{i < k < j} + \sum_{i < j < k} \right) (-1)^{i+j+k} \\ &\quad \iota_{[v_i, v_k] \wedge v_1 \wedge \dots \widehat{v}_i \dots \widehat{v}_j \dots \widehat{v}_k \dots \wedge v_m} \theta_j - \sum_{i < j} (-1)^{i+j+1} \iota_{v_1 \wedge \dots \widehat{v}_i \dots \widehat{v}_j \dots \wedge v_m} \mathcal{L}_{v_i} \theta_j. \end{aligned} \quad (\text{A.34})$$

Hence, combining the above with (A.33) results in the following equality:

$$\begin{aligned} \sum_i (-1)^{i-1} \mathcal{L}_{v_i} S_{m-1}(x_1, \dots, \widehat{x}_i, \dots, x_m) &= 2 \left(\sum_{i < k < j} - \sum_{i < j < k} + \sum_{j < i < k} \right) (-1)^{i+j+k} \iota_{[v_i, v_k] \wedge v_1 \wedge \dots \widehat{v}_i \dots \widehat{v}_j \dots \widehat{v}_k \dots \wedge v_m} \theta_j \\ &\quad + \left(\sum_{i < j} - \sum_{j < i} \right) (-1)^{i+j} \iota_{v_1 \wedge \dots \widehat{v}_i \dots \widehat{v}_j \dots \wedge v_m} \mathcal{L}_{v_i} \theta_j. \end{aligned} \quad (\text{A.35})$$

Finally, we deal with the second term on the right-hand side of (A.31). The commutation relation $[\iota, \mathcal{L}]$ (1.0.3) implies that

$$\begin{aligned} \sum_i (-1)^{i-1} \mathcal{L}_{v_i} \iota_{v_1 \wedge \dots \widehat{v}_i \dots \wedge v_m} \bar{A}(m-1) &= \sum_i (-1)^{i-1} \left(\iota_{[v_i, v_1 \wedge \dots \widehat{v}_i \dots \wedge v_m]} \right. \\ &\quad \left. + \iota_{v_1 \wedge \dots \widehat{v}_i \dots \wedge v_m} \mathcal{L}_{v_i} \right) \bar{A}(m-1) \end{aligned} \quad (\text{A.36})$$

For the first term on the right above, we use the definition of the Schouten bracket:

$$\begin{aligned} &\sum_i (-1)^{i-1} \iota_{[v_i, v_1 \wedge \dots \widehat{v}_i \dots \wedge v_m]} \bar{A}(m-1) \\ &= \left(\sum_{j < i} - \sum_{i < j} \right) (-1)^{i+j} \iota_{[v_i, v_j] \wedge v_1 \wedge \dots \widehat{v}_i \dots \widehat{v}_j \dots \wedge v_m} \bar{A}(m-1) \\ &= -2 \sum_{i < j} (-1)^{i+j} \iota_{[v_i, v_j] \wedge v_1 \wedge \dots \widehat{v}_i \dots \widehat{v}_j \dots \wedge v_m} \bar{A}(m-1), \end{aligned} \quad (\text{A.37})$$

where the last equality follows from skew-symmetry of the Lie bracket of vector fields. Combining this with the remain term in (A.36) and with the equality (A.35) gives the desired expression for $J^{m>2}$ and completes the proof. \square

Lemma A.7. *For all $m \geq 2$ and $x_1, \dots, x_m \in L$ the following equality holds:*

$$l'_1 f_m(x_1, \dots, x_m) = -(-1)^{\binom{m+1}{2}} (dS_m(x_1, \dots, x_m) + (-1)^m \iota_{v_1 \wedge \dots \wedge v_m} \omega + \mathcal{L}_{v_1 \wedge \dots \wedge v_m} \bar{A}(m-1)).$$

Proof. Recalling the definition of f_m for $m > 1$, we see that $l'_1 f_m = d_{\text{Tot}} f_m$ and

$$d_{\text{Tot}} f_m(x_1, \dots, x_m) = -(-1)^{\binom{m+1}{2}} (dS_m(x_1, \dots, x_m) + d_{\text{Tot}} \iota_{v_1 \wedge \dots \wedge v_m} \bar{A}(m)).$$

The Čech differential commutes with interior product. Hence,

$$d_{\text{Tot}} \iota_{v_1 \wedge \dots \wedge v_m} \bar{A}(m) = \iota_{v_1 \wedge \dots \wedge v_m} \delta \bar{A}(m) + d \iota_{v_1 \wedge \dots \wedge v_m} A^n + \sum_{i=1}^n (-1)^{mi+i} d \iota_{v_1 \wedge \dots \wedge v_m} A^{n-i}$$

Since \bar{A} is a Čech-Deligne n -cocycle, $\delta A^{n-j} = (-1)^j dA^{n-j-1}$ for $j = 0, \dots, n-1$. Therefore:

$$\begin{aligned} \iota_{v_1 \wedge \dots \wedge v_m} \delta \bar{A}(m) &= \iota_{v_1 \wedge \dots \wedge v_m} \sum_{j=0}^{n-1} (-1)^{mj} \delta A^{n-j} \\ &= -(-1)^m \iota_{v_1 \wedge \dots \wedge v_m} \sum_{i=1}^n (-1)^{(m-1)i} dA^{n-i}. \end{aligned}$$

Hence, the above plus Cartan's formula (1.0.2):

$$\mathcal{L}_{v_1 \wedge \dots \wedge v_m} = d\iota_{v_1 \wedge \dots \wedge v_m} - (-1)^m \iota_{v_1 \wedge \dots \wedge v_m} d$$

implies that

$$d_{\text{Tot}} \iota_{v_1 \wedge \dots \wedge v_m} \bar{A}(m) = d\iota_{v_1 \wedge \dots \wedge v_m} A^n + \sum_{i=1}^n (-1)^{(m-1)i} \mathcal{L}_{v_1 \wedge \dots \wedge v_m} A^{n-i}.$$

Since ω is the curvature of \bar{A} , we have

$$d\iota_{v_1 \wedge \dots \wedge v_m} A^n = (-1)^m \iota_{v_1 \wedge \dots \wedge v_m} \omega + \mathcal{L}_{v_1 \wedge \dots \wedge v_m} A^n,$$

and the desired result then follows from the definition of $\bar{A}(m-1)$ (A.2). \square

Proof of Theorem 4.2.2

Here we prove that the maps (A.9) $f_k: L^{\otimes k} \rightarrow L'$ assemble to give an L_∞ -morphism, and hence by Prop. A.1 an L_∞ -quasi-isomorphism.

We must verify the follow equality holds $\forall m \geq 1$:

$$\begin{aligned} & l'_1 f_m(x_1, \dots, x_m) + \sum_{j+k=m+1} \sum_{\sigma \in \text{Sh}(k, m-k)} \chi(\sigma) (-1)^{k(j-1)+1} f_j(l_k(x_{\sigma(1)}, \dots, x_{\sigma(k)}), x_{\sigma(k+1)}, \dots, x_{\sigma(m)}) \\ & + \sum_{s+t=m} \sum_{\substack{\tau \in \text{Sh}(s, m-s) \\ \tau(1) < \tau(s+1)}} \chi(\tau) (-1)^{s-1} (-1)^{(t-1) \sum_{p=1}^s |x_{\tau(p)}|} \llbracket f_s(x_{\tau(1)}, \dots, x_{\tau(s)}), f_t(x_{\tau(s+1)}, \dots, x_{\tau(m)}) \rrbracket = 0. \end{aligned}$$

Hence, in terms of the notation established above, we must show:

$$(l'_1 f_m + I_{(1)}^m + I_{(2)}^m + I_{(3)}^m + J^m)(x_1, \dots, x_m) = 0 \quad \forall m \geq 1. \quad (\text{A.38})$$

For $m = 1$, by definition $I_{(2)}^1 = I_{(3)}^1 = J^1 = 0$, and $I_{(1)}^1(x) = f_1(x)$. Hence, Eq. A.38 holds, since f_1 is a chain map.

For $m = 2$, we have $I_{(3)}^2 = 0$ by definition, and it follows from Lemmas A.2 and A.3 that

$$I_{(1)}^2(x_1, x_2) + I_{(2)}^2(x_1, x_2) = -[v_1, v_2] - \iota_{v_2} l_1 \theta_1 + \iota_{v_1} l_1 \theta_2 + \iota_{v_1 \wedge v_2} \omega - \iota_{[v_1, v_2]} \bar{A}(1)$$

By Lemma A.5, $J^2(x_1, x_2) = \llbracket f_1(x_1), f_1(x_2) \rrbracket$. Using the definition of f_1 and the Lie bracket, a simple calculation involving the Cartan calculus yields

$$J^2(x_1, x_2) = [v_1, v_2] - \iota_{v_1} d\theta_2 + \iota_{v_2} d\theta_1 - dS_2(x_1, x_2) + \iota_{[v_1, v_2]} \bar{A}(1) - \mathcal{L}_{v_1 \wedge v_2} \bar{A}(1),$$

Combining all of the above with Lemma A.7 gives the following for the left-hand side of Eq. A.38:

$$(l'_1 f_2 + I_{(1)}^2 + I_{(2)}^2 + I_{(3)}^2 + J^2)(x_1, x_2) = \iota_{v_1} (l_1 - d)\theta_2 + \iota_{v_2} (d - l_1)\theta_1 + 2\iota_{v_1 \wedge v_2} \omega.$$

If $|x_1| = |x_2| = 0$, then $l_1 = 0$ and the θ_i are Hamiltonian, i.e., $-\iota_{v_1} d\theta_2 = \iota_{v_2} d\theta_1 = -\iota_{v_1 \wedge v_2} \omega$. Hence, Eq. A.38 is satisfied. If $|x_i| > 0$, then $v_i = 0$ and $l_1 \theta_i = d\theta_i$, and hence the above calculation again shows that Eq. A.38 holds.

For the $m \geq 3$ case, we first observe that Lemma 3.2.1 combined with Cartan's formula (1.0.2) implies that for any $x_1, \dots, x_m \in L$:

$$\begin{aligned} (-1)^m \mathcal{L}_{v_1 \wedge \dots \wedge v_m} \bar{A}(m-1) &= \sum_{i < j} (-1)^{i+j} \iota_{[v_i, v_j] \wedge v_1 \wedge \dots \wedge \widehat{v}_i \wedge \dots \wedge \widehat{v}_j \wedge \dots \wedge v_m} \bar{A}(m-1) \\ &+ \sum_i (-1)^i \iota_{v_1 \wedge \dots \wedge \widehat{v}_i \wedge \dots \wedge v_m} \mathcal{L}_{v_m} \bar{A}(m-1), \end{aligned}$$

and similarly

$$\begin{aligned}
(-1)^{m-1} \sum_{j=1}^m (-1)^j \mathcal{L}_{v_1 \wedge \dots \wedge \widehat{v}_j \dots \wedge v_m} \theta_j &= \left(\sum_{i < k < j} - \sum_{i < j < k} + \sum_{j < i < k} \right) (-1)^{i+j+k} \iota_{[v_i, v_k] \wedge v_1 \wedge \dots \wedge \widehat{v}_i \dots \wedge \widehat{v}_j \dots \wedge v_k \dots \wedge v_m} \theta_j \\
&\quad + \left(\sum_{i < j} - \sum_{j < i} \right) (-1)^{i+j} \iota_{v_1 \wedge \dots \wedge \widehat{v}_i \dots \wedge \widehat{v}_j \dots \wedge v_m} \mathcal{L}_{v_i} \theta_j \\
&= (-1)^{m-1} dS_m(x_1, \dots, x_m) - \sum_{j=1}^m (-1)^j \iota_{v_1 \wedge \dots \wedge \widehat{v}_j \dots \wedge v_m} d\theta_j,
\end{aligned} \tag{A.39}$$

where the last equality follows from (A.8). Note that the above equations hold regardless of the degree of the x_i 's. Combining the above with Lemmas A.3 and A.6 gives the following equality:

$$\begin{aligned}
(I_{(2)}^m + J^m)(x_1, \dots, x_m) &= -(-1)^{\binom{m}{2}} \left((-1)^{m-1} \sum_{j=1}^m (-1)^j \mathcal{L}_{v_1 \wedge \dots \wedge \widehat{v}_j \dots \wedge v_m} \theta_j \right. \\
&\quad \left. - (-1)^m \mathcal{L}_{v_1 \wedge \dots \wedge v_m} \bar{A}(m-1) + \binom{m}{2} \iota_{v_1 \wedge \dots \wedge v_m} \omega \right).
\end{aligned} \tag{A.40}$$

Using the identity

$$(-1)^{\binom{m}{2}} (-1)^{\binom{m+1}{2}} = (-1)^m, \tag{A.41}$$

we combine Eq. A.40 with the result of Lemma A.4 to obtain:

$$\begin{aligned}
(I_{(2)}^m + I_{(3)}^m + J^m)(x_1, \dots, x_m) &= -(-1)^{\binom{m}{2}} \left((-1)^{m-1} \sum_{j=1}^m (-1)^j \mathcal{L}_{v_1 \wedge \dots \wedge \widehat{v}_j \dots \wedge v_m} \theta_j \right. \\
&\quad \left. - (-1)^m \mathcal{L}_{v_1 \wedge \dots \wedge v_m} \bar{A}(m-1) + (m-1) \iota_{v_1 \wedge \dots \wedge v_m} \omega \right).
\end{aligned} \tag{A.42}$$

Now we include the results of Lemmas A.2 and A.7 to the above and, again use the identity (A.41) to cancel terms:

$$\begin{aligned}
(l'_1 f_m + I_{(1)}^m + I_{(2)}^m + I_{(3)}^m + J^m)(x_1, \dots, x_m) &= \\
&\quad - (-1)^{\binom{m}{2}} \sum_{i=1}^m (-1)^i \iota_{v_1 \wedge \dots \wedge \widehat{v}_i \wedge \dots \wedge v_m} (l_1 \theta_i) \\
&\quad - (-1)^{\binom{m}{2}} \left(- \sum_{j=1}^m (-1)^j \iota_{v_1 \wedge \dots \wedge \widehat{v}_j \dots \wedge v_m} d\theta_j + m \iota_{v_1 \wedge \dots \wedge v_m} \omega \right).
\end{aligned} \tag{A.43}$$

(Note that above we used the last equality in (A.39) to cancel the dS_m term from $l'_1 f_m$.)

If we have $|x_k| > 0$ for some x_k then $v_k = 0$ and $l_1 \theta_k = d\theta_k$. Therefore the right-hand side of the above equality (A.43) vanishes and hence Eq. A.38 is satisfied. If all x_i are degree 0, then $l_1 = 0$ and all θ_i are Hamiltonian. Hence,

$$\begin{aligned}
\sum_{j=1}^m (-1)^j \iota_{v_1 \wedge \dots \wedge \widehat{v}_j \dots \wedge v_m} d\theta_j &= \sum_{j=1}^m (-1)^{j-1} \iota_{v_1 \wedge \dots \wedge \widehat{v}_j \dots \wedge v_m} \iota(v_j) \omega \\
&= \sum_{j=1}^m (-1)^{j-1} \iota_{v_j \wedge v_1 \wedge \dots \wedge \widehat{v}_j \dots \wedge v_m} \omega \\
&= m \iota_{v_1 \wedge \dots \wedge v_m} \omega.
\end{aligned} \tag{A.44}$$

Therefore, the right-hand side of (A.43) also vanishes in this case, and hence Eq. A.38 is satisfied. This completes the proof of the theorem. \square

References

- [1] J. Baez and A. Crans, Higher-dimensional algebra VI: Lie 2-algebras, arXiv:math/0307263v6.
- [2] J. Baez, A. Hoffnung, and C. Rogers, Categorical symplectic geometry and the classical string, *Comm. Math. Phys.* **293** (2010), 701–715. Also available as arXiv:0808.0246.
- [3] J. Baez, C. Rogers, *Categorified symplectic geometry and the string Lie 2-algebra*, Homology Homotopy Appl. 12 (2010), 221-236 Also available as arXiv:0901.4721
- [4] R. Bott and L. W. Tu, *Differential forms in algebraic topology*, Graduate Texts in Mathematics, 82, Springer, New York, 1982.
- [5] K. Brown, *Abstract homotopy theory and generalized sheaf cohomology*, Trans. Amer. Math. Soc. 186 (1973), 419-458.
- [6] J.-L. Brylinski, *Loop Spaces, Characteristic Classes and Geometric Quantization*, Birkhauser, Boston, 1993.
- [7] A. Canas da Silva, Y. Karshon, S Tolman, *Quantization of Presymplectic Manifolds and Circle Actions*, Trans. Amer. Math. Soc. 352 (2000), 525-552. Also available as arXiv:dg-ga/9705008.
- [8] A. Cannas da Silva and A. Weinstein, *Geometric Models for Noncommutative Algebras*, Berkeley Mathematics Lecture Notes **10**, Amer. Math. Soc., Providence, 1999.
- [9] B. Collier, Infinitesimal symmetries of Dixmier-Douady gerbes. Available as arXiv:1108.1525.
- [10] D. Fiorenza, C. Rogers, U. Schreiber, Higher geometric prequantum theory, Available as arXiv:1304.0236
- [11] D. Fiorenza, U. Schreiber, J. Stasheff Čech cocycles for differential characteristic classes, *Adv. Theor. Math. Phys.* 16 (2012) 149-250 Also available as arXiv:1011.4735
- [12] M. Forger, C. Paufler, and H. Römer, The Poisson bracket for Poisson forms in multisymplectic field theory, *Rev. Math. Phys.* **15** (2003), 705–744. Also available as arXiv:math-ph/0202043.
- [13] Y. Frégier, C. Rogers, and M. Zambon, Homotopy moment maps. Available as arXiv:1304.2051
- [14] V. Hinich, Homological algebra of homotopical algebras, *Comm. in Algebra* 25 (1997), 3291-3323.
- [15] V. Hinich, DG coalgebras as formal stacks, *J. Pure Appl. Algebra* **162** (2001), 209–250. Also available as arXiv:math/9812034 [math.AG].
- [16] B. Kostant, Quantization and unitary representations, *Lecture Notes in Math.* **170** (1970), 87–208.
- [17] T. Lada and M. Markl, Strongly homotopy Lie algebras, *Comm. Algebra.* **23** (1995), 2147–2161. Also available as arXiv:hep-th/9406095.
- [18] J.-L. Loday and B. Vallette, *Algebraic operads*, Grundlehren Math. Wiss. **346**, Springer, Heidelberg, 2012.
- [19] T. Nikolaus, U. Schreiber, D. Stevenson, *Principal ∞ -bundles, I: General theory*, Available as arXiv:1207.0248
T. Nikolaus, U. Schreiber, D. Stevenson, *Principal ∞ -bundles, II: Presentations*, Available as arXiv:1207.0249
- [20] J. Pridham, *Unifying derived deformation theories*, Adv. Math. 224 (2010), no.3, 772-826. Also available as arXiv:0705.0344

- [21] D. Quillen, Rational homotopy theory, *Ann. of Math. (2)* **90** (1969), 205–295.
- [22] C. Rogers, L_∞ -algebras from multisymplectic geometry, *Lett. Math. Phys.* **100** (2012), 29–50. Also available as arXiv:1005.2230.
- [23] C. Rogers, 2-plectic geometry, Courant algebroids, and categorified prequantization. *J. Symplectic Geom.* **11** (2013), 53–91. Also available as arXiv:1009.2975.
- [24] D. Roytenberg, On weak Lie 2-algebras, in: P. Kielanowski *et al* (eds.) XXVI Workshop on Geometrical Methods in Physics. AIP Conference Proceedings **956**, pp. 180-198. American Institute of Physics, Melville (2007). Also available as arXiv:0712.3461.
- [25] D. Roytenberg and A. Weinstein, Courant algebroids and strongly homotopy Lie algebras, *Lett. Math. Phys.* **46** (1998), 81–93. Also available as arXiv:math/9802118.
- [26] U. Schreiber, *Differential cohomology in a cohesive ∞ -topos*, <http://ncatlab.org/schreiber/files/cohesivedocumentv032.pdf>
- [27] J.-M. Souriau, *Structure of dynamical systems - A symplectic view of physics*, Birkhäuser (1997)
- [28] J.-M. Souriau, Quantification géométrique: Applications, *Ann. Inst. H. Poincaré Sect. A (N.S.)* **6** (1967), 311–341.