# **IIB** horizons

U. Gran<sup>1</sup>, J. Gutowski<sup>2</sup> and G. Papadopoulos<sup>3</sup>

<sup>1</sup> Fundamental Physics Chalmers University of Technology SE-412 96 Göteborg, Sweden

<sup>2</sup> Department of Mathematics University of Surrey Guildford, GU2 7XH, UK

<sup>3</sup> Department of Mathematics King's College London Strand London WC2R 2LS, UK

#### Abstract

We solve the Killing spinor equations for all near-horizon IIB geometries which preserve at least one supersymmetry. We show that generic horizon sections are 8-dimensional almost Hermitian spin<sub>c</sub> manifolds. Special cases include horizon sections with a Spin(7) structure and those for which the Killing spinor is pure. We also explain how the common sector horizons and the horizons with only 5-form flux are included in our general analysis. We investigate several special cases mainly focusing on the horizons with constant scalars admitting a pure Killing spinor and find that some of these exhibit a generalization of the 2-SCYT condition that arises in the horizons with 5-form fluxes only. We use this to construct new examples of near-horizon geometries with both 3-form and 5-form fluxes.

# 1 Introduction

In understanding the physics of black holes and branes, and in AdS/CFT, horizons have played a central role. In most applications, the horizons are Killing<sup>1</sup>, and in the extreme case can be investigated in a unified way because there is a local model for the near-horizon fields [1, 2]. Typically, some additional assumptions are made depending on the problem. The most common assumption is that the metric and all the other form field strengths<sup>2</sup> are taken to be smooth. This assumption includes many black hole horizons, as well as the horizons of some branes, like the D3-, M2- and M5-branes. Therefore it is expected that a systematic understanding of all near-horizon geometries will have applications in AdS/CFT [3] as well as in the construction of IIB back hole solutions which exhibit horizons with exotic topology and geometry, see e.g. [4]-[16] for earlier related works.

In this paper, we solve the Killing spinor equations (KSEs) of IIB supergravity for all near-horizon geometries preserving at least one supersymmetry and determine the geometry of the horizon sections. This extends the results on M-horizons [17] and heterotic horizons [18] to IIB supergravity. In the investigation of the KSEs for IIB horizons three cases<sup>3</sup> arise depending on the choice of Killing spinor (i) the generic horizons, (ii) the Spin(7) horizons and (iii) the pure SU(4) horizons. We find that the sections  $\mathcal{S}$  of generic near-horizon geometries are 8-dimensional almost Hermitian spin<sub>c</sub> manifolds, and the sections of the Spin(7) horizons admit a Spin(7) structure. The sections of the pure SU(4) horizons, which admit a pure Killing spinor, are again 8-dimensional almost Hermitian spin<sub>c</sub> manifolds. In all the above cases, we also examine the implications on the topology and geometry of horizon sections depending on whether the IIB scalars take values in the hyperbolic upper half-plane or the hyperbolic upper half-plane after an  $SL(2, \mathbb{Z})$  U-duality identification.

We further investigate near-horizon geometries with constant scalars. The structure group of horizon sections both for generic and pure SU(4) horizons reduces further to a subgroup of SU(4). Therefore in these two cases, the near-horizon sections are almost Hermitian spin<sub>c</sub> manifolds with an SU(4) structure. In all IIB horizons without further restrictions on the fields, the KSEs do not impose additional restrictions<sup>4</sup> on the geometry of the horizon sections. We also explore several examples mostly focusing on the pure SU(4) horizons with constant scalars. In the complex case, we find a deformation of the 2-SKT condition,  $\partial \bar{\partial} \omega^2 = 0$ , of horizons with only 5-form fluxes [20] with the addition of a source term which depends on the 3-form fluxes. We use this to give new examples of horizons with 3- and 5-form fluxes. We also describe how the horizons with only 5-form flux and the horizons of the common sector arise as special cases of our IIB horizons.

To prove the above results, we first demonstrate that the KSEs are integrable along the

<sup>&</sup>lt;sup>1</sup>This means that they admit a time Killing vector field which becomes null at a spacetime hypersurface.

<sup>&</sup>lt;sup>2</sup>Non-smooth near-horizon geometries can also be investigated in the special case for which there exists a frame in which the metric is smooth. This class includes all branes. In this case, one works in a frame with respect to which the metric is smooth, and allow singularities in the other fields, such as the scalars of the theory. If all fields are smooth, the choice of frame is not essential.

<sup>&</sup>lt;sup>3</sup>This is similar to the three cases that arise in the solution of the KSEs for IIB backgrounds preserving one supersymmetry in [19].

<sup>&</sup>lt;sup>4</sup>Of course to find solutions, one has to impose in addition the Bianchi identities and field equations.

lightcone directions giving rise to a system of differential and algebraic equations on the horizon section S. We find using the field equations, Bianchi identities and the bilinear matching condition<sup>5</sup> that this system can be considerably simplified giving rise to a parallel transport equation and an algebraic equation on the horizon sections associated with the gravitino and dilatino KSEs, respectively. These two equations are solved using spinorial geometry [21], and as we have already mentioned, there are three cases to consider: the generic horizons, the Spin(7) horizons, and the horizons with a pure Killing spinor. The geometry of the horizon sections is next investigated by computing the Killing spinor bilinears, and in particular those which are associated with nowhere vanishing forms on the horizon sections. In the analysis above, the compactness properties of section S have not been used. Because of this, the results apply to both brane and black hole horizons. However in the construction of examples, like those associated with the deformation of the 2-SKT condition, we take S to be compact without boundary as expected for black hole horizon sections.

This paper has been organized as follows. In section 2, we state the near-horizon metric and the form fluxes of IIB supergravity and decompose the field equations and Bianchi identities along the lightcone and horizon section S directions. Moreover we derive which equations are independent. In section 3, we integrate the KSEs along the lightcone directions and use the field equations, Bianchi identities and the bilinear matching condition to derive the independent KSEs. In section 4, we solve the independent KSEs using spinorial geometry and express the fluxes in terms of the geometry. Moreover, we derive the geometry of the horizon section and describe some aspects of their topology. In section 5, we explain how the horizons with only 5-form flux and those of the common sector are included as special cases in our analysis. In section 6, we explore the geometry of horizon sections admitting a pure Killing spinor and demonstrate that in the complex case the 2-SKT conditions of 5-form horizons is deformed with the 3-form fluxes. In section 7, we state our conclusions, and in appendix A we give the solution of the linear system associated with IIB horizons.

# 2 Fields and dynamics near a horizon

# 2.1 Fields and KSEs

To describe the form field strengths of IIB supergravity, consider the complex line bundle  $\lambda$  over the spacetime. This is the pull-back of the canonical bundle of either the coset space  $\mathbb{H} = SU(1,1)/U(1)$  or after a U-duality identification of  $SL(2,\mathbb{R})\setminus\mathbb{H}$  with respect to the dilaton and axion, the two scalar fields of the theory.  $\mathbb{H}$  is identified with the hyperbolic upper half-plane which is a manifold and  $SL(2,\mathbb{R})\setminus\mathbb{H}$  is identified with the fundamental domain of the moduli space of complex structures of a 2-torus which is an orbifold.

The field strengths of the bosonic fields of IIB supergravity are a  $\lambda^2$ -valued complex 1-form P, a  $\lambda$ -valued complex 3-form G and a self-dual 5-form F. P, G and F are the

<sup>&</sup>lt;sup>5</sup>The bilinear matching condition is the identification of the stationary Killing vector field of a black hole with the Killing spinor vector bilinear.

field strengths of two scalars, two 2-form gauge potentials and a 4-form gauge potential, respectively. Assuming that the horizons under investigation are killing horizons, and under some regularity assumptions, the IIB fields in the extreme near-horizon limit can be expressed as

$$ds^{2} = 2\mathbf{e}^{+}\mathbf{e}^{-} + \delta_{ij}\mathbf{e}^{i}\mathbf{e}^{j} = 2du(dr + rh - \frac{1}{2}r^{2}\Delta du) + \gamma_{IJ}dy^{I}dy^{J} , \qquad (2.1)$$

$$F = r\mathbf{e}^+ \wedge X + \mathbf{e}^+ \wedge \mathbf{e}^- \wedge Y + \star_8 Y , \qquad (2.2)$$

$$G = r\mathbf{e}^+ \wedge L + \mathbf{e}^+ \wedge \mathbf{e}^- \wedge \Phi + H , \qquad (2.3)$$

$$P = \xi , \qquad (2.4)$$

where we have introduced the frame

$$\mathbf{e}^{+} = du, \qquad \mathbf{e}^{-} = dr + rh - \frac{1}{2}r^{2}\Delta du, \qquad \mathbf{e}^{i} = e_{I}^{i}dy^{I}, \qquad (2.5)$$

and used the self-duality<sup>6</sup> of F,

$$F_{M_1M_2M_3M_4M_5} = -\frac{1}{5!} \epsilon_{M_1M_2M_3M_4M_5} {}^{N_1N_2N_3N_4N_5} F_{N_1N_2N_3N_4N_5} , \qquad (2.6)$$

which also requires that

$$X = -\star_8 X . \tag{2.7}$$

IIB supergravity also admits a U(1) connection of  $\lambda$ , denoted by Q. Upon restriction to the near-horizon geometry,

$$Q = \Lambda (2.8)$$

where  $\Lambda$  is r, u-independent, i.e.  $Q_+ = Q_- = 0$ , and so  $\Lambda$  is a connection of  $\lambda$  restricted to S.

The dependence on the coordinates u and r is explicitly given, and so h,  $\Delta$ , X, Y, L,  $\Phi$ , H,  $\Lambda$  and  $\xi$  depend only on the coordinates  $y^I$  of the horizon section S which is the co-dimension 2 subspace given by r = u = 0. In particular, the scalars depend only on y, and  $\xi$  is a section of  $\lambda^2 \otimes \Lambda^1(S)$ , where we have restricted<sup>7</sup>  $\lambda$  to S. Similarly, L,  $\Phi$  and H are sections of  $\lambda \otimes \Lambda^2(S)$ ,  $\lambda \otimes \Lambda^1(S)$  and  $\lambda \otimes \Lambda^3(S)$ , respectively, and X and Y are sections of  $\Lambda^4(S)$  and  $\Lambda^3(S)$ , respectively. Observe that  $V = \frac{\partial}{\partial u}$  is the horizon timelike Killing vector which becomes null at r = 0. The derivation of (2.1)-(2.4) is similar to that of other near-horizon geometries with form fluxes and we shall not elaborate here.

<sup>&</sup>lt;sup>6</sup>We choose  $\epsilon_{0123456789} = 1$  and the spacetime volume form is related to that of the horizon section as  $dvol(M) = \mathbf{e}^+ \wedge \mathbf{e}^- \wedge dvol(\mathcal{S})$ . In particular,  $(\star_8 Y)_{n_1n_2n_3n_4n_5} = \frac{1}{3!} \epsilon_{n_1n_2n_3n_4n_5} {}^{m_1m_2m_3} Y_{m_1m_2m_3}$ . <sup>7</sup>Note that  $\lambda$  is topologically trivial if it is the pull-back of a line bundle over the hyperbolic upper half-

<sup>&</sup>lt;sup>7</sup>Note that  $\lambda$  is topologically trivial if it is the pull-back of a line bundle over the hyperbolic upper halfplane which is a contractible space. This will also be the case after an  $SL(2,\mathbb{Z})$  U-duality identification of the IIB scalars as the j-function maps the fundamental domain homeomorphically to  $\mathbb{C}$ , unless two copies are glued together to a sphere as in the case of 24 cosmic strings [25]. In both cases,  $\lambda$  may be equipped with a connection with non-vanishing curvature.

# 2.2 Bianchi identities and field equations

Before proceeding to an analysis of the KSEs, it is useful to evaluate the Bianchi identities and the field equations of IIB supergravity [22, 23] on the near-horizon fields (2.1)-(2.4). First consider the Bianchi identity

$$dF - \frac{i}{8}G \wedge \bar{G} = 0 . \qquad (2.9)$$

This implies that

$$X = d_h Y - \frac{i}{8} (\Phi \wedge \bar{H} - \bar{\Phi} \wedge H) , \quad d_h Y = dY - h \wedge Y , \qquad (2.10)$$

$$d \star_8 Y = \frac{i}{8} H \wedge \bar{H} , \qquad (2.11)$$

or equivalently

$$\tilde{\nabla}^{\ell} Y_{\ell i j} = -\frac{i}{288} \epsilon_{i j}^{\ell_1 \ell_2 \ell_3 \ell_4 \ell_5 \ell_6} H_{\ell_1 \ell_2 \ell_3} \bar{H}_{\ell_4 \ell_5 \ell_6} , \qquad (2.12)$$

where  $\tilde{\nabla}$  is the Levi-Civita connection on S. Therefore X is not independent. However, notice that the anti-self duality condition (2.7) of X imposes an additional restriction on the components of the fluxes that appear in (2.10).

Next, the Bianchi identity

$$dG - iQ \wedge G + P \wedge \bar{G} = 0 , \qquad (2.13)$$

implies that

$$L = d_h \Phi - i\Lambda \wedge \Phi + \xi \wedge \bar{\Phi} , \qquad (2.14)$$

and

$$dH = i\Lambda \wedge H - \xi \wedge \bar{H} . \tag{2.15}$$

So again L is not independent and can be expressed in terms of the other fluxes.

The Bianchi identity

$$dP - 2iQ \wedge P = 0 , \qquad (2.16)$$

implies

$$d\xi = 2i\Lambda \wedge \xi , \qquad (2.17)$$

and the Bianchi identity

$$dQ = -iP \wedge \bar{P} , \qquad (2.18)$$

implies

$$d\Lambda = -i\xi \wedge \bar{\xi} \ . \tag{2.19}$$

We remark that (2.9) and (2.13) imply two additional identities involving dX and dL, however, these are in fact implied by (2.10), (2.11), (2.14), (2.15), (2.17), (2.19). As a consequence, the latter are necessary and sufficient conditions imposed by the Bianchi identities.

Next we turn to the bosonic field equations. Substituting the near-horizon fields into the 2-form gauge potentials field equations

$$\nabla^C G_{ABC} - iQ^C G_{ABC} - P^C \bar{G}_{ABC} + \frac{2i}{3} F_{ABN_1N_2N_3} G^{N_1N_2N_3} = 0 , \qquad (2.20)$$

one finds that

$$\tilde{\nabla}^{i} \Phi_{i} - i\Lambda^{i} \Phi_{i} - \xi^{i} \bar{\Phi}_{i} + \frac{2i}{3} Y_{\ell_{1}\ell_{2}\ell_{3}} H^{\ell_{1}\ell_{2}\ell_{3}} = 0 , \qquad (2.21)$$

$$-\tilde{\nabla}^{i}L_{im} + i\Lambda^{i}L_{im} + h^{i}L_{im} - \frac{1}{2}dh^{ij}H_{ijm} + \xi^{i}\bar{L}_{im} + \frac{2i}{3}(X_{m\ell_{1}\ell_{2}\ell_{3}}H^{\ell_{1}\ell_{3}\ell_{3}} - 3Y_{m\ell_{1}\ell_{2}}L^{\ell_{1}\ell_{2}}) = 0 ,$$
(2.22)

and

$$\tilde{\nabla}^{\ell} H_{\ell i j} - i\Lambda^{\ell} H_{\ell i j} - h^{\ell} H_{\ell i j} + L_{i j} - \xi^{\ell} \bar{H}_{\ell i j} + \frac{2i}{3} (\star_8 Y_{i j \ell_1 \ell_2 \ell_3} H^{\ell_1 \ell_2 \ell_3} - 6Y_{i j \ell} \Phi^{\ell}) = 0 . \quad (2.23)$$

Similarly, the field equation of the scalars

$$\nabla^A P_A - 2iQ^A P_A + \frac{1}{24}G_{N_1N_2N_3}G^{N_1N_2N_3} = 0 , \qquad (2.24)$$

implies

$$\tilde{\nabla}^{i}\xi_{i} - 2i\Lambda^{i}\xi_{i} - h^{i}\xi_{i} + \frac{1}{24}(-6\Phi^{i}\Phi_{i} + H_{\ell_{1}\ell_{2}\ell_{3}}H^{\ell_{1}\ell_{2}\ell_{3}}) = 0.$$
(2.25)

Note that there is no independent field equation for F because F is self-dual.

It remains to investigate the Einstein equation

$$R_{AB} - \frac{1}{6} F_{AL_1L_2L_3L_4} F_B^{L_1L_2L_3L_4} - \frac{1}{4} G_{(A}^{N_1N_2} \bar{G}_{B)N_1N_2} + \frac{1}{48} g_{AB} G_{N_1N_2N_3} \bar{G}^{N_1N_2N_3} - 2P_{(A} \bar{P}_{B)} = 0.$$
(2.26)

Substituting the near-horizon fields, one finds that +- component gives

$$\frac{1}{2}\tilde{\nabla}^{i}h_{i} - \Delta - \frac{1}{2}h^{2} + \frac{2}{3}Y_{\ell_{1}\ell_{2}\ell_{3}}Y^{\ell_{1}\ell_{2}\ell_{3}} + \frac{3}{8}\Phi^{i}\bar{\Phi}_{i} + \frac{1}{48}H_{\ell_{1}\ell_{2}\ell_{3}}\bar{H}^{\ell_{1}\ell_{2}\ell_{3}} = 0.$$
(2.27)

Similarly, the +i, ij and ++ components imply that

$$-\frac{1}{2}\tilde{\nabla}^{j}dh_{ji} - dh_{ij}h^{j} - \tilde{\nabla}_{i}\Delta + \Delta h_{i} + \frac{4}{3}X_{i\ell_{1}\ell_{2}\ell_{3}}Y^{\ell_{1}\ell_{2}\ell_{3}} \\ -\frac{1}{8}\left(L_{\ell_{1}\ell_{2}}\bar{H}_{i}^{\ell_{1}\ell_{2}} - 2\Phi^{\ell}\bar{L}_{i\ell} + \bar{L}_{\ell_{1}\ell_{2}}H_{i}^{\ell_{1}\ell_{2}} - 2\bar{\Phi}^{\ell}L_{i\ell}\right) = 0 , \qquad (2.28)$$

$$\tilde{R}_{ij} + \tilde{\nabla}_{(i}h_{j)} - \frac{1}{2}h_{i}h_{j} + 4Y_{i\ell_{1}\ell_{2}}Y_{j}^{\ell_{1}\ell_{2}} + \frac{1}{2}\Phi_{(i}\bar{\Phi}_{j)} - 2\xi_{(i}\bar{\xi}_{j)} - \frac{1}{4}H_{\ell_{1}\ell_{2}(i}\bar{H}_{j)}^{\ell_{1}\ell_{2}} + \delta_{ij}\left(-\frac{1}{8}\Phi_{\ell}\bar{\Phi}^{\ell} - \frac{2}{3}Y_{\ell_{1}\ell_{2}\ell_{3}}Y^{\ell_{1}\ell_{2}\ell_{3}} + \frac{1}{48}H_{\ell_{1}\ell_{2}\ell_{3}}\bar{H}^{\ell_{1}\ell_{2}\ell_{3}}\right) = 0 , \qquad (2.29)$$

and

$$\frac{1}{2}\tilde{\nabla}^{2}\Delta - \frac{3}{2}h^{i}\tilde{\nabla}_{i}\Delta - \frac{1}{2}\Delta\hat{\nabla}^{i}h_{i} + \Delta h^{2} + \frac{1}{4}dh_{ij}dh^{ij} - \frac{1}{6}X_{\ell_{1}\ell_{2}\ell_{3}\ell_{4}}X^{\ell_{1}\ell_{2}\ell_{3}\ell_{4}} - \frac{1}{4}L_{ij}\bar{L}^{ij} = 0 ,$$
(2.30)

respectively, where  $\tilde{R}$  is the Ricci tensor of S.

Not all of the above field equations are independent. In particular, on taking the divergence of (2.23), and making use of (2.10), (2.11), (2.14), (2.15), (2.17), (2.19), (2.23), (2.7), one obtains (2.22). Also, on computing the Einstein tensor of  $\mathcal{S}$  using (2.29) and (2.27), and evaluating the Einstein tensor Bianchi identity, one obtains (2.28) upon making use of (2.10), (2.11), (2.14), (2.15), (2.17), (2.21), (2.23), (2.25). Furthermore, on taking the divergence of (2.28), one also obtains (2.30), after making use of (2.7) and (2.10) to compute  $d \star_8 X$ , (2.14) to compute dL, and also (2.28) and (2.23). Hence it follows that (2.22), (2.28) and (2.30) are implied by the other field equations and Bianchi identities.

# 3 KSEs on IIB horizons

## 3.1 Lightcone integrability of KSEs

The gravitino and dilatino KSEs of IIB supergravity [22, 23] are

$$\left(\nabla_{M} - \frac{i}{2}Q_{M} + \frac{i}{48}F_{MN_{1}N_{2}N_{3}N_{4}}\Gamma^{N_{1}N_{2}N_{3}N_{4}}\right)\epsilon - \frac{1}{96}\left(\Gamma_{M}{}^{N_{1}N_{2}N_{3}}G_{N_{1}N_{2}N_{3}} - 9G_{MN_{1}N_{2}}\Gamma^{N_{1}N_{2}}\right)C * \epsilon = 0 , \qquad (3.1)$$

$$P_M \Gamma^M C * \epsilon + \frac{1}{24} G_{N_1 N_2 N_3} \Gamma^{N_1 N_2 N_3} \epsilon = 0 .$$
(3.2)

For our spinor conventions see [19].

As in the analysis of KSEs for near-horizon geometries in other supergravity theories, the IIB KSEs can be integrated along the lightcone directions. In particular solving the - component of the gravitino KSE, one finds

$$\begin{aligned}
\epsilon_{+} &= \phi_{+} \\
\epsilon_{-} &= \phi_{-} + r\Gamma_{-} \left( \frac{1}{4} h_{i} \Gamma^{i} + \frac{i}{12} Y_{n_{1} n_{2} n_{3}} \Gamma^{n_{1} n_{2} n_{3}} \right) \phi_{+} \\
&+ r\Gamma_{-} \left( \frac{1}{96} H_{\ell_{1} \ell_{2} \ell_{3}} \Gamma^{\ell_{1} \ell_{2} \ell_{3}} + \frac{3}{16} \Phi_{i} \Gamma^{i} \right) C * \phi_{+} , 
\end{aligned} \tag{3.3}$$

where  $\phi_{\pm}$  do not depend on r. Similarly, the solution of the + component of the gravitino KSE gives

$$\phi_{+} = \eta_{+} + u\Gamma_{+} \left( \frac{1}{4} h_{i} \Gamma^{i} - \frac{i}{12} Y_{n_{1}n_{2}n_{3}} \Gamma^{n_{1}n_{2}n_{3}} \right) \eta_{-} ,$$
  
+  $u\Gamma_{+} \left( \frac{1}{96} H_{\ell_{1}\ell_{2}\ell_{3}} \Gamma^{\ell_{1}\ell_{2}\ell_{3}} - \frac{3}{16} \Phi_{i} \Gamma^{i} \right) C * \eta_{-}$   
$$\phi_{-} = \eta_{-} , \qquad (3.4)$$

where  $\eta_{\pm}$  do not depend on the *u* and *r* coordinates. Furthermore,  $\eta_{+}$  and  $\eta_{-}$  must also satisfy a number of algebraic conditions. To simplify these, we shall first identify the 1-form associated with the stationary Killing vector field  $V = \partial_u$  of horizons with the 1-form *Z* constructed as a Killing spinor bilinear.

#### 3.1.1 Vector bilinear matching condition

The 1-form Killing spinor bilinear<sup>8</sup> is

$$Z = \langle B(C\epsilon^*)^*, \Gamma_A \epsilon \rangle \ e^A = \langle \Gamma_0 \epsilon, \Gamma_A \epsilon \rangle \ e^A , \qquad (3.5)$$

where  $\epsilon$  is a Killing spinor and  $\langle \cdot, \cdot \rangle$  is the standard Hermitian inner product, see [19]. We require that Z should be proportional to V, where

$$V = -\frac{1}{2}r^2\Delta \mathbf{e}^+ + \mathbf{e}^- . \tag{3.6}$$

First, evaluate Z at r = u = 0, for which  $\epsilon = \eta_+ + \eta_-$ . Requiring that  $Z_+ = 0$  at r = u = 0 forces

$$\eta_{-} = 0$$
 . (3.7)

Then, using r, u independent Spin(8) gauge transformations of the type considered in [19], one can, without loss of generality, take

$$\eta_+ = p + q e_{1234} , \qquad (3.8)$$

<sup>&</sup>lt;sup>8</sup>There are other  $\lambda$ -valued 1-form bilinears which can be constructed from the Killing spinor  $\epsilon$ . However, these cannot be identified with V as they are twisted 1-forms and generically are not associated with a Killing vector.

for p, q complex valued functions. Furthermore, on computing the component  $Z_-$ , one finds that  $|p|^2 + |q|^2$  must be a (non-zero) constant, or equivalently,

$$\langle \eta_+, \eta_+ \rangle = \text{const.}$$
 (3.9)

Next, evaluate  $Z_i$  at  $r \neq 0$ . As this component must vanish, one finds the condition

$$(|p|^{2} + |q|^{2})h_{\alpha} + \frac{3}{4}(pq\bar{\Phi}_{\alpha} + \bar{p}\bar{q}\Phi_{\alpha}) + 2i(|p|^{2} - |q|^{2})Y_{\alpha\beta}^{\beta} - \frac{1}{4}(pq\bar{H}_{\alpha\beta}^{\beta} + \bar{p}\bar{q}H_{\alpha\beta}^{\beta}) + \frac{1}{12}\epsilon_{\alpha\beta_{1}\beta_{2}\beta_{3}}(p^{2}\bar{H}^{\beta_{1}\beta_{2}\beta_{3}} + \bar{q}^{2}H^{\beta_{2}\beta_{2}\beta_{3}}) = 0 , \qquad (3.10)$$

where the 8-dimensional indices are split as  $i = (\alpha, \bar{\alpha})$ .

It will be convenient to define

$$\tau_{+} = \left(\frac{1}{4}h_{i}\Gamma^{i} + \frac{i}{12}Y_{\ell_{1}\ell_{2}\ell_{3}}\Gamma^{\ell_{1}\ell_{2}\ell_{3}}\right)\eta_{+} + \left(\frac{3}{16}\Phi_{i}\Gamma^{i} + \frac{1}{96}H_{\ell_{1}\ell_{2}\ell_{3}}\Gamma^{\ell_{1}\ell_{2}\ell_{3}}\right)C * \eta_{+} .$$
(3.11)

Then the condition (3.10) implies that one can write

$$\tau_{+} = q\mu^{\alpha}e_{\alpha} - \frac{1}{6}p \ \overline{(\mu^{\beta})} \ \epsilon_{\overline{\beta}}{}^{\alpha_{1}\alpha_{2}\alpha_{3}}e_{\alpha_{1}\alpha_{2}\alpha_{3}} \ , \tag{3.12}$$

with

$$(|p|^{2} + |q|^{2})\mu^{\alpha} = \frac{3}{8\sqrt{2}} ((\bar{q})^{2} \Phi^{\alpha} - p^{2} \bar{\Phi}^{\alpha}) + \sqrt{2} i p \bar{q} Y^{\alpha}{}_{\mu}{}^{\mu} - \frac{i}{3\sqrt{2}} (|p|^{2} + |q|^{2}) Y_{\mu_{1}\mu_{2}\mu_{3}} \epsilon^{\mu_{1}\mu_{2}\mu_{3}\alpha} + \frac{1}{8\sqrt{2}} ((\bar{q})^{2} H^{\alpha}{}_{\mu}{}^{\mu} - p^{2} \bar{H}^{\alpha}{}_{\mu}{}^{\mu}) + \frac{\sqrt{2}}{48} (-\bar{p} \bar{q} H_{\mu_{1}\mu_{2}\mu_{3}} + p q \bar{H}_{\mu_{1}\mu_{2}\mu_{3}}) \epsilon^{\mu_{1}\mu_{2}\mu_{3}\alpha} .$$
(3.13)

Also, noting that

$$\Delta = -2r^{-2}\frac{Z_+}{Z_-} , \qquad (3.14)$$

one finds

$$\Delta = 4\delta_{\alpha\bar{\beta}}\mu^{\alpha} \overline{(\mu^{\beta})} , \qquad (3.15)$$

so  $\Delta \ge 0$ , as expected.

### 3.1.2 Algebraic conditions

Returning to the solution of the + and - components of the gravitino KSE, the Killing spinor  $\epsilon$  can be written in terms of  $\eta_+$ ,  $\tau_+$  as

$$\epsilon = \eta_+ + r\Gamma_- \tau_+ . \tag{3.16}$$

Moreover, one also finds the algebraic conditions

$$\left(\frac{1}{2}\Delta - \frac{1}{8}dh_{ij}\Gamma^{ij}\right)\eta_{+} + \frac{1}{16}L_{ij}\Gamma^{ij}C * \eta_{+} + \left(\frac{1}{2}h_{i}\Gamma^{i} - \frac{i}{6}Y_{\ell_{1}\ell_{2}\ell_{3}}\Gamma^{\ell_{1}\ell_{2}\ell_{3}}\right)\tau_{+} + \left(-\frac{3}{8}\Phi_{i}\Gamma^{i} + \frac{1}{48}H_{\ell_{1}\ell_{2}\ell_{3}}\Gamma^{\ell_{1}\ell_{2}\ell_{3}}\right)C * \tau_{+} = 0$$
(3.17)

and

$$(\Delta h_i - \partial_i \Delta)\Gamma^i \eta_+ + \left( -\frac{1}{2} dh_{ij} \Gamma^{ij} + \frac{i}{12} X_{\ell_1 \ell_2 \ell_3 \ell_4} \Gamma^{\ell_1 \ell_2 \ell_3 \ell_4} \right) \tau_+ + \frac{1}{2} L_{ij} \Gamma^{ij} C * \tau_+ = 0 .$$
(3.18)

This completes the integration of the gravitino KSEs along the light-cone directions.

# 3.2 The S components of the gravitino KSE

Substituting the Killing spinor  $\epsilon$ , (3.16), into the gravitino KSE and evaluating the resulting expression along the directions transverse to the light cone, one finds

$$\tilde{\nabla}_{i}\eta_{+} + \left(-\frac{i}{2}\Lambda_{i} - \frac{1}{4}h_{i} - \frac{i}{4}Y_{i\ell_{1}\ell_{2}}\Gamma^{\ell_{1}\ell_{2}} + \frac{i}{12}\Gamma_{i}^{\ell_{1}\ell_{2}\ell_{3}}Y_{\ell_{1}\ell_{2}\ell_{3}}\right)\eta_{+} \\ + \left(\frac{1}{16}\Gamma_{i}^{j}\Phi_{j} - \frac{3}{16}\Phi_{i} - \frac{1}{96}\Gamma_{i}^{\ell_{1}\ell_{2}\ell_{3}}H_{\ell_{1}\ell_{2}\ell_{3}} + \frac{3}{32}H_{i\ell_{1}\ell_{2}}\Gamma^{\ell_{1}\ell_{2}}\right)C * \eta_{+} = 0 , \quad (3.19)$$

and

$$\tilde{\nabla}_{i}\tau_{+} + \left(-\frac{i}{2}\Lambda_{i} - \frac{3}{4}h_{i} + \frac{i}{4}Y_{i\ell_{1}\ell_{2}}\Gamma^{\ell_{1}\ell_{2}} - \frac{i}{12}\Gamma_{i}^{\ell_{1}\ell_{2}\ell_{3}}Y_{\ell_{1}\ell_{2}\ell_{3}}\right)\tau_{+} \\ + \left(-\frac{1}{16}\Gamma_{i}^{\ j}\Phi_{j} + \frac{3}{16}\Phi_{i} - \frac{1}{96}\Gamma_{i}^{\ell_{1}\ell_{2}\ell_{3}}H_{\ell_{1}\ell_{2}\ell_{3}} + \frac{3}{32}H_{i\ell_{1}\ell_{2}}\Gamma^{\ell_{1}\ell_{2}}\right)C * \tau_{+} \\ + \left(-\frac{1}{4}dh_{ij}\Gamma^{j} - \frac{i}{12}X_{i\ell_{1}\ell_{2}\ell_{3}}\Gamma^{\ell_{1}\ell_{2}\ell_{3}}\right)\eta_{+} + \left(\frac{1}{32}\Gamma_{i}^{\ell_{1}\ell_{2}}L_{\ell_{1}\ell_{2}} - \frac{3}{16}L_{ij}\Gamma^{j}\right)C * \eta_{+} = 0. \quad (3.20)$$

Both the above equations are parallel transport equations along  $\mathcal{S}$ .

# 3.3 The dilatino KSE

It remains to evaluate the dilatino KSE (3.2) on the spinor (3.16). A direct substitution reveals that

$$\left(-\frac{1}{4}\Phi_{i}\Gamma^{i} + \frac{1}{24}H_{\ell_{1}\ell_{2}\ell_{3}}\Gamma^{\ell_{1}\ell_{2}\ell_{3}}\right)\eta_{+} + \xi_{i}\Gamma^{i}C * \eta_{+} = 0 , \qquad (3.21)$$

and

$$\left(-\frac{1}{4}\Phi_{i}\Gamma^{i} - \frac{1}{24}H_{\ell_{1}\ell_{2}\ell_{3}}\Gamma^{\ell_{1}\ell_{2}\ell_{3}}\right)\tau_{+} - \xi_{i}\Gamma^{i}C * \tau_{+} + \frac{1}{8}L_{ij}\Gamma^{ij}\eta_{+} = 0.$$
(3.22)

This concludes the evaluation of the KSEs in the IIB near-horizon geometries and their integration along the lightcone directions.

## 3.4 Independent KSEs

It is well known that the KSEs imply some of the Bianchi identities and field equations. Because of this, to find solutions, it is customary to solve the KSEs and then impose the remaining field equations and Bianchi identities. However, we shall not do this here because of the complexity of solving the KSEs (3.17), (3.18), (3.20), and (3.22) which contain the  $\tau$  spinor as expressed in (3.11) and (3.12). Instead, we shall show that all the KSEs which contain  $\tau$  are actually implied from those containing only  $\eta$ , i.e. (3.19) and (3.21), and some of the field equations and Bianchi identities.

#### 3.4.1 The (3.20) condition

The (3.20) component of KSEs is implied by (3.19) and (3.11) together with a number of bosonic field equations and Bianchi identities. To see this, first evaluate the LHS of (3.20) by substituting in (3.11) to eliminate  $\tau_+$ , and use (3.19) to evaluate the supercovariant derivatives of  $\eta_+$  and  $C * \eta_+$ . Also evaluate

$$\left(\frac{1}{4}\tilde{R}_{ij}\Gamma^{j} - \frac{1}{2}\Gamma^{j}(\tilde{\nabla}_{j}\tilde{\nabla}_{i} - \tilde{\nabla}_{i}\tilde{\nabla}_{j})\right)\eta_{+} = 0.$$
(3.23)

This vanishes identically, however a non-trivial identity is obtained by expanding out the supercovariant derivative terms again using (3.19). Then, on adding (3.23) to the LHS of (3.20), with  $\tau_+$  eliminated in favour of  $\eta_+$  using (3.11) and (3.19) as mentioned above, one obtains, after some calculation, a term proportional to (2.29).

Therefore, it follows that (3.20) is implied by (3.19) and (3.11) and the bosonic field equations and Bianchi identities. We remark that in addition to using (2.29) in establishing this identity, we also make use of (2.10), (2.7), (2.11), (2.14), (2.15), (2.19), (2.21) and (2.23).

### 3.4.2 The (3.22) condition

Next consider (3.21) and (3.22). On defining

$$\mathcal{A}_{1} = \left( -\frac{1}{4} \Phi_{i} \Gamma^{i} + \frac{1}{24} H_{\ell_{1}\ell_{2}\ell_{3}} \Gamma^{\ell_{1}\ell_{2}\ell_{3}} \right) \eta_{+} + \xi_{i} \Gamma^{i} C * \eta_{+} , \qquad (3.24)$$

and

$$\mathcal{A}_{2} = \left( -\frac{1}{4} \Phi_{i} \Gamma^{i} - \frac{1}{24} H_{\ell_{1}\ell_{2}\ell_{3}} \Gamma^{\ell_{1}\ell_{2}\ell_{3}} \right) \tau_{+} - \xi_{i} \Gamma^{i} C * \tau_{+} + \frac{1}{8} L_{ij} \Gamma^{ij} \eta_{+} , \qquad (3.25)$$

one obtains the following identity

$$\mathcal{A}_2 = -\frac{1}{2}\Gamma^i \tilde{\nabla}_i \mathcal{A}_1 + \left(\frac{3i}{4}\Lambda_i \Gamma^i + \frac{3}{8}h_i \Gamma^i - \frac{i}{12}Y_{\ell_1 \ell_2 \ell_3} \Gamma^{\ell_1 \ell_2 \ell_3}\right) \mathcal{A}_1 , \qquad (3.26)$$

where we have made use of (3.19) in order to evaluate the covariant derivative in the above expression. In addition, we also have made use of the field equations and Bianchi identities (2.14), (2.15), (2.17), (2.21), (2.23) and (2.25). It follows that these conditions, together with (3.21) imply (3.22).

### 3.4.3 The (3.17) condition

To show that (3.17) is also implied as a consequence of the KSEs (3.19) and (3.21), and the field equations and Bianchi identities, contract (3.20) with  $\Gamma^i$  and use (3.11) to rewrite the  $\tau_+$  terms in terms of  $\eta_+$ . Then subtract  $(\frac{3}{16}\bar{\Phi}_i\Gamma^i + \frac{1}{96}\bar{H}_{\ell_1\ell_2\ell_3}\Gamma^{\ell_1\ell_2\ell_3})\mathcal{A}_1$  from the resulting expression to obtain (3.17). In order to obtain (3.17) from these expressions, we also make use of (2.10), (2.7), (2.11), (2.23), (2.14), (2.21), and (2.27). It follows, from section 3.4.1 above, that (3.17) follows from the above mentioned Bianchi identities and field equations, together with (3.19) and (3.21).

#### 3.4.4 The (3.18) condition

The (3.18) condition is obtained from (3.17) as follows. First act on (3.17) with the Dirac operator  $\Gamma^i \tilde{\nabla}_i$ , and use the bosonic field equations and Bianchi identities to eliminate the  $d \star_8 dh$ , dL,  $d \star_8 L$ ,  $d \star_8 h$ , dY,  $d \star_8 Y$ , dH and  $d \star_8 H$  terms, and rewrite  $d\Phi$  in terms of L. Then use the algebraic conditions (3.21) and (3.22) to eliminate the  $\xi$ -terms from the resulting expression. The terms involving  $\Lambda$  then vanish as a consequence of (3.17).

Next consider the dh-terms; after some calculation, these can be rewritten as

$$\frac{1}{2}dh_{ij}\Gamma^{ij}\tau_{+} - \frac{7}{32}h_{\ell}\Gamma^{\ell}dh_{ij}\Gamma^{ij}\eta_{+} + \left(-\frac{1}{64}\Phi_{\ell}\Gamma^{\ell} + \frac{1}{384}H_{\ell_{1}\ell_{2}\ell_{3}}\Gamma^{\ell_{1}\ell_{2}\ell_{3}}\right)dh_{ij}\Gamma^{ij}C * \eta_{+} + \left(-\frac{1}{64}\Phi_{\ell}\Gamma^{\ell} + \frac{1}{384}H_{\ell_{1}\ell_{2}\ell_{3}}\Gamma^{\ell_{1}\ell_{2}\ell_{3}}\right)dh_{ij}\Gamma^{ij}C * \eta_{+} + \left(-\frac{1}{64}\Phi_{\ell}\Gamma^{\ell} + \frac{1}{384}H_{\ell_{1}\ell_{2}\ell_{3}}\Gamma^{\ell_{1}\ell_{2}\ell_{3}}\right)dh_{ij}\Gamma^{ij}C * \eta_{+}$$

The dh terms involving  $\eta_+$  and  $C * \eta_+$  in the above expression are then eliminated, using (3.17). On collating the remaining terms, one finds that those involving  $\Delta$  (but not  $d\Delta$ ) are

$$-\Delta h_j \Gamma^j \eta_+ . \tag{3.27}$$

It is also straightforward to note that the terms involving  $\overline{L}$  vanish, whereas the terms involving X and L can be rewritten as

$$-\frac{1}{2}L_{ij}\Gamma^{ij}C * \tau_{+} - \frac{i}{12}X_{\ell_{1}\ell_{2}\ell_{3}\ell_{4}}\Gamma^{\ell_{1}\ell_{2}\ell_{3}\ell_{4}}\tau_{+} , \qquad (3.28)$$

where the anti-self-duality of X has been used to simplify the expression. The remaining terms which are linear in  $\tau_+, C * \tau_+$  and quadratic in  $h, Y, \Phi, \overline{\Phi}, H, \overline{H}$  can be shown to vanish after some computation. After performing these calculations, the condition which is obtained is (3.18).

# 4 Solution of KSEs and geometry of horizons

## 4.1 The linear system

It is straightforward to derive the linear system associated with the (3.19) and (3.21) KSEs evaluated on the spinor

$$\eta_+ = p1 + qe_{1234} , \qquad (4.1)$$

where we have used a local  $Spin(8) \cdot U(1)$  transformation to arrange such that p, q are real functions on  $\mathcal{S}$ , and as required from the bilinear matching condition take

$$p^2 + q^2 = 1 {.} {4.2}$$

Substituting this spinor into (3.19) and (3.21), one obtains a linear system which is explicitly given in appendix A. This can be used to express some of the fluxes in terms of geometry and find the conditions on the geometry of the horizon sections S imposed by supersymmetry.

There are three different types of backgrounds which arise naturally in the investigation of the solutions of the linear system in appendix A:

- The generic horizons for which the Killing spinor is chosen such that  $p^2 + q^2 = 1$ ,  $p^2 q^2 \neq 0$ , and p and q not identically zero at the horizon though they may vanish at some points.
- The Spin(7) horizons for which  $p^2 + q^2 = 1$  and p = q.
- The pure SU(4) horizons for which either p = 1, q = 0 or p = 0, q = 1.

Clearly, the generic horizons include both the Spin(7) and pure SU(4) horizons. Typically, as the Killing spinor is parallel transported along the horizon section, it will change type as the only requirement is that  $p^2 + q^2 = 1$ . The solution of the linear system for all three cases is given in appendix A. In what follows, we shall explore the consequences which arise on the topology and geometry of horizon sections S from the solution of the linear system and hence from the solution of the KSEs.

# 4.2 Generic horizons

## 4.2.1 Solution of the linear system

The general solution of the linear system is described in appendix A under the assumption that the functions p and q may vanish at isolated points. The main properties of the solution are the following. All complex fluxes  $\xi, \Phi$  and H can be expressed in terms of the real fluxes Y and the geometry  $\Omega$ . Moreover, the solution of the linear system does *not* contain a condition which involves only the spin connection of S.

## 4.2.2 Topology and geometry of S

The gravitino KSE as reduced to S in (3.19) is a parallel transport equation of sections of a  $\lambda^{\frac{1}{2}} \otimes \Sigma^+$  bundle over S, where  $\lambda^{\frac{1}{2}} \otimes \Sigma^+$  is a  $Spin_c(8) = Spin(8) \cdot U(1)$  bundle and  $\Sigma^+$ is the complexified spin bundle over S associated with the positive chirality Majorana-Weyl representation of Spin(8). The real rank of  $\lambda^{\frac{1}{2}} \otimes \Sigma^+$  is 16. An application of [24] implies that the supercovariant connection as restricted on S given in (3.19) has holonomy contained in  $SL(16, \mathbb{R})$ . The existence of a solution to the gravitino KSE requires that the holonomy of the supercovariant connection reduces to  $SL(15, \mathbb{R})$ , and so  $\lambda^{\frac{1}{2}} \otimes \Sigma^+$ must admit a nowhere vanishing section spanned by the parallel spinor. The existence of nowhere vanishing sections typically impose a topological restriction on the spin bundle and so on a manifold. However, a priori this may not be the case here as  $\lambda^{\frac{1}{2}} \otimes \Sigma^+$  has rank much bigger than the dimension of S and so it always admits such a nowhere vanishing section.

Nevertheless, the existence of a solution to the KSEs requires that the structure group of  $\mathcal{S}$  reduces form SO(8) to a subgroup of U(4). To see this, recall that  $\mathcal{S}$  admits a nowhere vanishing spinor  $\eta_+ = p1 + qe_{1234}$  which is a section of  $\lambda^{\frac{1}{2}} \otimes \Sigma^+$ . One can consider the bilinears of  $\eta_+$ . There are two kinds of bilinears which can be constructed from  $\eta_+$ . First the bilinears constructed from  $\eta_+$  and itself with respect to the standard Hermitian inner product of Spin(8). These bilinears are forms on  $\mathcal{S}$ . In addition one can also consider the bilinears constructed from  $\eta_+$  and its complex conjugate  $C * \eta_+$ . Such bilinears are not forms on  $\mathcal{S}$ . Instead they are forms twisted by  $\lambda$  as they carry a U(1) charge.

Consider first the bilinears of  $\eta_+ = p1 + qe_{1234}$  which are forms on S. We find that a basis in the ring of these bilinears is

$$\omega = -i(p^2 + q^2)\delta_{\alpha\bar{\beta}}e^{\alpha} \wedge e^{\bar{\beta}} , \quad \psi = pq\operatorname{Re}\left(e^1 \wedge e^2 \wedge e^3 \wedge e^4\right) , \qquad (4.3)$$

with  $p^2 + q^2 = 1$ . Since  $\omega$  vanishes nowhere, it is an almost Hermitian form on S. In addition  $\psi$  defines a (4,0)-form on S. But as p and q can vanish at some points on S,  $\psi$  is not nowhere vanishing. As a result, the structure group of S reduces from SO(8) to a subgroup of U(4) rather than to a subgroup of SU(4). S is an almost Hermitian 8-dimensional manifold.

Next, the form bilinears of  $\eta_{+} = p1 + qe_{1234}$  which are twisted by  $\lambda$  are

$$\rho = -ipq\delta_{\alpha\bar{\beta}}e^{\alpha} \wedge e^{\bar{\beta}} , \quad \tau = p^2 e^1 \wedge e^2 \wedge e^3 \wedge e^4 + q^2 e^{\bar{1}} \wedge e^{\bar{2}} \wedge e^{\bar{3}} \wedge e^{\bar{4}} . \tag{4.4}$$

As a result  $\lambda \otimes (\Lambda^{4,0}(\mathcal{S}) \oplus \Lambda^{0,4}(\mathcal{S}))$  admits a nowhere vanishing section.

Furthermore, as has been mentioned, the KSEs solved in appendix A do not impose any additional geometric restrictions on S. As a result, the horizon sections S are  $\text{Spin}_c$ almost Hermitian 8-dimensional manifolds. The only remaining potential restrictions on Sare those imposed by the field equations and Bianchi identities which we shall investigate in some special cases below.

So far, we have performed our analysis without distinguishing whether the IIB scalars take values in the upper half plane or in the fundamental domain. Now we return to address this issue. If the IIB scalars take values in the upper half plane, then the line bundle  $\lambda$  is topologically trivial as it is the pull back of a trivial bundle over the upper half plane with respect to a smooth map. Nevertheless, if the IIB scalars are non-constant<sup>9</sup>  $\lambda$  will be geometrically twisted as it will have non-vanishing curvature.

In the case that the IIB scalars take values in the fundamental domain,  $\lambda$  may not be topologically trivial. This arises from the analysis of stringy cosmic string solutions in [25] where there are solutions which are smooth and the horizon section of the cosmic strings is a 2-sphere, e.g. the configuration of 24 cosmic strings. Of course, our case here is more general and S is not a complex manifold. Nevertheless, it is an indication that  $\lambda$ over S may not be topologically trivial in the fundamental domain case.

 $<sup>^{9}\</sup>text{It}$  is not apparent that there are smooth solutions for which  $\mathcal S$  is compact and the IIB scalars are non-trivial.

# **4.3** Spin(7) Horizons

#### 4.3.1 Solution of the linear system

The spinor  $\eta_+$  can be chosen as  $\eta_+ = \frac{1}{\sqrt{2}}(1 + e_{1234})$ . The solution of the linear system is given as in appendix A after setting  $p = q = 1/\sqrt{2}$  in the solution of the generic case. Furthermore, the expression for  $\mu$  can be simplified as

$$\frac{\mu_{\bar{\alpha}}}{\sqrt{2}} = -\frac{i}{2}\Lambda_{\bar{\alpha}} + iY_{\bar{\alpha}\lambda}{}^{\lambda} + \frac{i}{3}\epsilon_{\bar{\alpha}}{}^{\lambda_1\lambda_2\lambda_3}Y_{\lambda_1\lambda_2\lambda_3} , \qquad (4.5)$$

which is consistent with our previous results on IIB horizons with only 5-form flux.

#### 4.3.2 Topology and geometry of horizon sections

The analysis of the consequences for the existence of a parallel  $\eta_+$  spinor on the holonomy of the supercovariant connection as reduced on S is similar to the generic case. To find whether the structure group of S reduces to a subgroup of SO(8), we again compute the form bilinears associated with  $\eta_+$ . In this case, there is a single nowhere vanishing 4-form bilinear<sup>10</sup>

$$\phi = -\frac{1}{2}\omega \wedge \omega + 4\operatorname{Re}\chi\tag{4.6}$$

which is a representative of the fundamental form of Spin(7). Therefore the structure group of S reduces to Spin(7), where  $\chi = \frac{1}{4!} \epsilon_{\alpha_1 \alpha_2 \alpha_3 \alpha_4} e^{\alpha_1} \wedge e^{\alpha_2} \wedge e^{\alpha_3} \wedge e^{\alpha_4}$ . It is known that the structure group of an 8-dimensional compact spin manifold reduces from SO(8)to Spin(7) provided that [26]

$$\pm e - \frac{1}{2}p_2 + \frac{1}{8}p_1^2 = 0 , \qquad (4.7)$$

where e is the Euler class, and  $p_2$  and  $p_1$  are the second and first Pontryagin classes, respectively. This condition restricts the topology of S. Furthermore, a direct inspection of the solution of the linear system reveals that there are no additional geometric conditions on the Spin(7) structure of S. Thus S can be any manifold with a Spin(7) structure.

## 4.4 Pure SU(4) horizons

#### 4.4.1 Solution of the linear system

The  $\eta_+$  spinor in this case can be chosen as  $\eta_+ = 1$ . The solution of the linear system has been given in appendix A. In particular, many components of the complex fields, like H, can be expressed in terms of the components of the 5-form and components of the spin connection of spacetime. Again, the solution of the linear system in appendix A does not yield a condition which restricts the spin connection of S.

<sup>&</sup>lt;sup>10</sup>The almost hermitian form  $\omega$  and the (4,0)-form  $\chi$  may not be globally defined on S but  $\phi$  is. Observe that we have normalized  $\chi$  differently in [19].

## 4.4.2 Topology and geometry of horizon sections

A similar analysis to that presented for the generic case reveals that the structure group of  $\mathcal{S}$  reduces to a subgroup of U(4). In this case, it is instructive to also consider the mixed bilinears of  $\eta_+$  with  $\tilde{\eta}_+ = C * \eta_+ = e_{1234}$  in (4.4). In addition to the almost hermitian form  $\omega$  which arises as a bilinear of  $\eta_+$ , as in (4.3), there is a  $\lambda$ -twisted (4,0)-form bilinear given locally by  $\tau$  for p = 1, q = 0. Therefore the line bundle  $\lambda \otimes \Lambda^{4,0}$  on  $\mathcal{S}$  admits a nowhere vanishing section, and so it is topologically trivial. Thus  $\lambda$  can be identified with the anti-canonical bundle of  $\mathcal{S}$ . Furthermore, the solution of the linear system in appendix A does not reveal any additional geometric constraints on  $\mathcal{S}$ . As a result,  $\mathcal{S}$  is any almost Hermitian spin<sub>c</sub> manifold.

Next let us turn to examine the upper half plane and fundamental domain cases. As in the generic class, if the IIB scalars take values in the upper half plane,  $\lambda$  is topologically trivial. The structure group of S then reduces to a subgroup of SU(4). However,  $\lambda$ can be geometrically non-trivial as its curvature may not vanish provided that the IIB scalars are non-trivial functions<sup>11</sup>. On the other hand, if the IIB scalars take values in the fundamental domain, as has been explained in the generic case,  $\lambda$  may not be topologically trivial on S.

# 5 Special cases

# 5.1 Horizons with only 5-form flux

It is well known that IIB supergravity has two consistent truncations. One truncation leads to a sector with only 5-form fluxes active, while for the other truncation the 5-form field strength vanishes and both the 3-form and 1-form field strengths are real. The latter is the common sector. In both cases, the near-horizon geometries are compatible with both these truncations. It is straightforward to see this in the 5-form sector. The nearhorizon geometries with only 5-form fluxes can be recovered from the general IIB horizons that we have investigated by setting

$$P = G = 0 (5.1)$$

Conversely, the 5-form horizons investigated in [20] can be embedded into general IIB horizons exhibiting only non-vanishing 5-form fluxes. As a result, all the examples of IIB horizons constructed in [20] can be embedded into the general IIB horizons.

# 5.2 Common sector horizons

The embedding of the common sector horizons into general IIB horizons is not as straightforward as that described in the previous section for horizons with only 5-form fluxes. To describe the embedding of common sector horizons, let us recall how the common sector is obtained from the general IIB theory. For this set F = 0, and as has already been mentioned, choose both P and G to be real. Observe that Q = 0, and the Bianchi identities

<sup>&</sup>lt;sup>11</sup>It is not apparent that there exist smooth solutions with  $\mathcal{S}$  compact and non-trivial scalars.

of IIB supergravity can be solved after setting

$$P = \frac{1}{2}d\phi , \quad G = -e^{-\frac{1}{2}\phi}\mathcal{H} , \qquad (5.2)$$

provided that  $d\mathcal{H} = 0$ . These field definitions yield the common sector in the string frame provided that in addition we relate the IIB metric,  $ds_{\text{IIB}}^2$ , with the common sector metric,  $ds_{\text{CS}}^2$ , as

$$ds_{\rm IIB}^2 = e^{-\frac{1}{2}\phi} ds_{\rm CS}^2 , \qquad (5.3)$$

and identify  $\phi$  and  $\mathcal{H}$  with the dilaton and the NS-NS 3-form field strength, respectively.

Furthermore, the KSEs of the common sector are obtained from those of IIB provided that the IIB supersymmetry parameter is related to that of the common sector as

$$\epsilon_{\rm CS} = e^{\frac{1}{8}\phi} \epsilon_{\rm IIB} \ . \tag{5.4}$$

In particular, we find that the KSEs of the common sector are

$$\nabla_M^{\pm} \epsilon_{\rm CS}^{\pm} = 0 ,$$
  
$$(\Gamma^M \partial_M \phi \mp \frac{1}{12} \Gamma^{MNR} \mathcal{H}_{MNR}) \epsilon_{\rm CS}^{\pm} = 0 , \qquad (5.5)$$

where  $\nabla^{\pm} = \nabla \pm \frac{1}{2}\mathcal{H}$ ,  $\nabla$  the Levi-Civita connection of the common sector metric, and  $C * \epsilon_{\rm CS}^{\pm} = \pm \epsilon_{\rm CS}^{\pm}$ .

It remains to show how the common sector near-horizons are embedded into general IIB horizons. For this, one has to demonstrate that after making the field redefinitions (5.2) and (5.3), one can adapt a coordinate system such that the common sector near-horizon fields give rise to IIB near-horizon fields. Indeed consider first the metric. If the common sector metric is in near-horizon form, the IIB metric can also be put in near-horizon form provided that we make a coordinate transformation

$$r_{\rm IIB} = e^{-\frac{1}{2}\phi} r_{\rm CS} ,$$
 (5.6)

the 1-form  $h_{\rm CS}$  is replaced with  $h_{\rm IIB} = h_{\rm CS} + \frac{1}{2}d\phi$ , and the metric of the horizon section S is conformally scaled as  $g_{\rm IIB} = e^{-\frac{1}{2}\phi}g_{\rm CS}$ .

Next turn to the 3-form field strength. Suppose that  $\mathcal{H}$  is in a near-horizon form. G in (5.2) can also be put into a near-horizon form as well since the conformal factor in the definition of G is absorbed in the coordinate transformation for r in (5.6) and the redefinition of h. One also has to re-scale H as  $H_{\text{IIB}} = e^{-\frac{1}{2}\phi}H_{\text{CS}}$ . Observe now that  $H_{\text{CS}}$  is closed as expected. To make connection with our results for heterotic horizons, one can compensate the overall sign in the definition of  $\mathcal{H}$  by changing the sign of L,  $\Phi$  and H.

A consequence of our analysis above is that we can embed all the near-horizon geometries we have found in heterotic theory into IIB. We remark that all the horizons we had investigated in [18] have  $d\mathcal{H} = 0$  and so they can be thought as solutions of the common sector as well. In this way, all the explicit heterotic near-horizon geometries found in [18] can be utilized to give explicit examples of IIB horizons. Note also that all the IIB horizons with non-trivial fluxes that arise from the embedding of heterotic horizons in IIB preserve more than two supersymmetries and admit additional isometries.

#### **Pure** SU(4) horizons with constant scalars 6

#### 6.1Solution to the linear system

A special class of horizons are those for which the IIB scalars are constant. In this case, there is a significant simplification of the solution to the linear system because P = Q = 0.

The solution to the KSEs given in appendix A simplifies somewhat. In particular, one finds that the fluxes of pure SU(4) horizons with constant scalars can be expressed<sup>12</sup> as

$$Y = \frac{1}{4} (d\omega - \theta_{\text{Re}\chi} \wedge \omega) + \text{Im}\Phi \cdot \text{Re}\chi , \quad \omega \cdot H^{1,2} = H^{0,3} = \Phi^{0,1} = 0 ,$$
  
$$H^{3,0} = (\theta_{\text{Re}\chi} - \theta_{\omega}) \cdot \chi , \quad H_{\bar{\alpha}\lambda_1\lambda_2} = \frac{1}{2} \mathcal{N}_{\bar{\nu}_1\bar{\nu}_2\bar{\alpha}} \epsilon^{\bar{\nu}_1\bar{\nu}_2}{}_{\lambda_1\lambda_2} - 2\delta_{\bar{\alpha}[\lambda_1}\Phi_{\lambda_2]} , \quad h = \theta_{\text{Re}\chi} ,$$
  
(6.1)

where  $\theta_{\omega}$  and  $\theta_{\text{Re}\chi}$  are the Lee forms<sup>13</sup> of the almost Hermitian form  $\omega$  and the (4,0) form  $\chi = \frac{1}{4!} \epsilon_{\alpha_1 \alpha_2 \alpha_3 \alpha_4} e^{\alpha_1} \wedge e^{\alpha_2} \wedge e^{\alpha_3} \wedge e^{\alpha_4}$  of the SU(4) structure, respectively, and

$$\mathcal{N}_{jk}^{i} = I^{m}{}_{j} \nabla_{m} I^{i}{}_{k} - I^{m}{}_{k} \nabla_{m} I^{i}{}_{j} - I^{i}{}_{m} (\nabla_{j} I^{m}{}_{k} - \nabla_{k} I^{m}{}_{j}) , \qquad (6.2)$$

is the Nijenhuis tensor of the almost complex structure I. The fluxes  $\Phi^{1,0}$  and the traceless part of  $H^{1,2}$  are not restricted by the KSEs.

#### 6.2Topology and geometry of horizons

As the scalars are constant, the line bundle  $\lambda$  is trivial. As a consequence, the bilinears of both  $\eta_+$  and  $\tilde{\eta}_+ = e_{1234}$  are forms on  $\mathcal{S}$  and in particular  $\chi$  is a nowhere vanishing (4,0)-form on  $\mathcal{S}$ . Thus the structure group of  $\mathcal{S}$  reduces to a subgroup of SU(4).

As in the Spin(7) case, there are topological obstructions to reduce the structure group of  $\mathcal{S}$  from SO(8) to SU(4). Since Spin(7) and SU(4) have the same maximal torus<sup>14</sup> the obstruction (4.7) is also an obstruction in the SU(4) case. In particular using the relation between the Pontryagin and Chern classes  $p_1 = c_1^2 - 2c_2$  and  $p_2 = c_2^2 - 2c_1c_3 + 2c_4$  on almost complex manifolds, see e.g. [27], and the identification of the Euler class  $e = c_4$ , (4.7) is satisfied provided  $c_1 = 0$  which vanishes as a consequence of SU(4) structure. The orientation induced by the almost complex structure leads to the choice of the plus sign in (4.7). There may be additional obstructions to reduce the structure group from SO(8)to SU(4) which take values in cohomology with  $\mathbb{Z}_k$  coefficients.

The solution of the linear system does not impose any additional conditions on the geometry of  $\mathcal{S}$ . Therefore  $\mathcal{S}$  is an almost Hermitian manifold with an SU(4) structure.

#### 6.3Magnetic 3-form flux deformations

#### Complex deformation of horizons with 5-form fluxes 6.3.1

A class of horizons with 3-form flux can be constructed as a deformation of horizons with 5-form fluxes. Recall that the horizons with only 5-form fluxes are 2-SCYT manifolds.

<sup>&</sup>lt;sup>12</sup>If v is  $\ell$ -vector and  $\alpha$  a k-form, then  $(v \cdot \alpha)_{i_1...i_{k-\ell}} = v^{j_1...j_\ell} \alpha_{j_1...j_\ell i_1...i_{k-\ell}}$ . <sup>13</sup>We define  $(\theta_{\omega})_i = -I^j{}_i \nabla^k \omega_{kj}$  and  $(\theta_{\text{Re}\chi})_i = \frac{2}{3} \text{Re} \chi_i{}^{j_1 j_2 j_3} \nabla^k \text{Re} \chi_{kj_1 j_2 j_3}$ .

<sup>&</sup>lt;sup>14</sup>We thank Simon Salamon for discussions on this.

The deformation we shall investigate here lifts the 2-strong condition but the horizon remains a complex manifold. For this, we take<sup>15</sup>

$$\Phi = H^{0,3} = H^{3,0} = H^{2,1} = 0 , \quad Y^{0,3} = Y^{3,0} = 0 .$$
(6.3)

In this case, the conditions stated in appendix A imply that  $\mathcal{S}$  is a Hermitian manifold and in addition

$$Y = \frac{1}{4} (d\omega - \omega \wedge \theta_{\omega}) , \quad \omega \cdot H^{1,2} = 0 , \quad \theta_{\omega} = \theta_{\text{Re}\chi} , \quad h = \theta_{\omega} , \qquad (6.4)$$

where  $\theta_{\omega}$  and  $\theta_{\text{Re}\chi}$  are the Lee forms of the Hermitian form  $\omega$  and  $\text{Re}\chi$  is the real part of the (4,0)-form  $\chi$ . Observe that the traceless part of  $H^{1,2}$  is not restricted by the KSEs. Since S is a Hermitian manifold with an SU(4) structure, the condition  $\theta_{\omega} = \theta_{\text{Re}\chi}$  implies that S admits a connection  $\hat{\nabla} = \nabla + \frac{1}{2}\hat{H}$  with skew-symmetric torsion  $\hat{H}$  which has holonomy SU(4), i.e.  $\text{hol}(\hat{\nabla}) \subseteq SU(4)$ , where  $\hat{H}$  is uniquely determined in terms of the metric g and complex structure I of the horizon section as  $\hat{H} = -i_I d\omega$ . Thus S admits a CYT structure.

It remains to solve the field equations and Bianchi identities. One can show after a long but straightforward calculation that all these are satisfied provided that

$$d(\omega \wedge \mathring{H}) = \frac{i}{2}H \wedge \bar{H} , \quad dH = 0 , \quad h^{\bar{\gamma}}H_{\bar{\gamma}\alpha\bar{\beta}} + 2iH_{\alpha\bar{\gamma}_1\bar{\gamma}_2}Y_{\bar{\beta}}{}^{\bar{\gamma}_1\bar{\gamma}_2} = 0 .$$
 (6.5)

The first condition is a generalization of the 2-strong condition found in [20] for horizons with only 5-form fluxes. In particular, the 3-form flux appears as a source term in the rhs of the 2-strong condition. Observe also that the last condition together with  $\omega \cdot H^{1,2} = 0$  imply that  $H \cdot Y = 0$  and so H and Y must be orthogonal.

#### 6.3.2 Examples

To construct examples, one begins with an 8-dimensional CYT manifold and then imposes the conditions (6.5). Examples of CYT manifolds can be constructed as toric fibrations over Kähler manifolds [28, 29, 30]. This construction has been generalized in [20] and adapted to solve the 2-SCYT condition of IIB horizons with only 5-form fluxes. Here we shall use the analysis of [20] to find geometries that solve (6.5).

First consider S as a  $T^2$  fibration over a 6-dimensional Kähler manifold  $X^6$  with Kähler form  $\omega_{(6)}$ . Let  $(\lambda^1, \lambda^2)$  be a principal bundle connection on S. In order for S to admit an SU(4) structure compatible with a connection with skew-symmetric torsion, the curvature  $\mathcal{F}^1 = d\lambda^1$  must be identified with the Ricci form of  $X^6$ , i.e.

$$\rho_{(6)} = -\mathcal{F}^1 , \qquad (6.6)$$

and  $\mathcal{F}^2 = d\lambda^2$  must be (1,1) and traceless. In the latter case, such connections always exist on complex line bundles. The metric and Hermitian form on  $\mathcal{S}$  is chosen as

$$ds^{2} = \frac{2}{k} [(\lambda^{1})^{2} + (\lambda^{2})^{2}] + ds^{2} (X^{6}) , \quad \omega_{(8)} = -\frac{2}{k} \lambda^{1} \wedge \lambda^{2} + \omega_{(6)} , \qquad (6.7)$$

<sup>&</sup>lt;sup>15</sup>Note that since  $\Phi = 0$  it follows from (A.17), using (A.16), that  $\tau_+ = 0$  and therefore the Killing spinor in this case is just  $\epsilon = \eta_+$ , which is a considerable simplification.

where  $\mathcal{F}_{ij}^{1}\omega_{(6)}^{ij} = k$  and k is required to be constant<sup>16</sup>.

This choice of Hermitian structure specifies both H and Y. It remain to choose H. For this write

$$H = \bar{\tau} \wedge \alpha + \beta , \quad \tau = \frac{1}{\sqrt{2}} (\lambda^1 + i\lambda^2) , \qquad (6.8)$$

where  $\alpha$  and  $\beta$  are complex (1,1)- and (1,2)-forms on  $X^6$ . The requirement that H is traceless implies that

$$\omega_{(6)} \cdot \alpha = \omega_{(6)} \cdot \beta = 0 . \tag{6.9}$$

Furthermore dH = 0 implies that

$$d\alpha = 0 , \quad d\beta + \frac{1}{\sqrt{2}}\alpha \wedge [\mathcal{F}^1 - i\mathcal{F}^2] = 0 .$$
(6.10)

Next the first equation in (6.5) is solved provided

$$\alpha \wedge \bar{\alpha} = 0$$
,  $\alpha \wedge \bar{\beta} = 0$ ,  $\omega_{(6)} \wedge [(\mathcal{F}^1)^2 + (\mathcal{F}^2)^2] = \frac{ik}{4}\beta \wedge \bar{\beta}$ . (6.11)

The simplest case is to take  $\alpha = 0$ . Then since  $\theta_{\omega} = -\lambda^2$  and

$$Y = \frac{1}{2k} (\lambda^1 \wedge \mathcal{F}^2 - \lambda^2 \wedge \mathcal{F}^1) + \frac{1}{4} \lambda^2 \wedge \omega_{(6)}$$
(6.12)

the last condition in (6.5) is also satisfied.

Therefore to find examples of near-horizon geometries, one must find a Kähler manifold  $X^6$  with positive constant scalar curvature that admits a traceless (1,2) closed form  $\beta$  that satisfies the last equation in (6.11).

To find solutions, one possibility is to take  $X^6$  to either be a Kähler-Einstein manifold or products of Kähler-Einstein manifolds. For example, one can take  $X^6 = S^2 \times S^2 \times T^2$ . Normalizing the metric of  $S^2$ 's such that  $\mathcal{F}^1 = \omega^1 + \omega^2$  where  $\omega^1$  and  $\omega^2$  are the Kähler forms on the  $S^2$ 's, k = 4, and choosing

$$\begin{aligned}
\omega_{(6)} &= -d\varphi^1 \wedge d\varphi^2 + \omega^1 + \omega^2 , \quad H = m(d\varphi^1 - id\varphi^2) \wedge (\omega^1 - \omega^2) , \\
\mathcal{F}^2 &= \ell(\omega^1 - \omega^2) , \quad \ell \in \mathbb{Z} ,
\end{aligned}$$
(6.13)

all equations are solved provided that  $2|m|^2 = \ell^2 - 1$ , where  $\varphi^1$  and  $\varphi^2$  are the angular coordinates of  $T^2$  in  $X^6$ . This generalizes the examples of 2-SCYT near-horizon geometries with only 5-form flux in [20].

<sup>&</sup>lt;sup>16</sup>Since  $X^6$  is Kähler, observe that k is constant provided  $\mathcal{F}^1$  is co-closed. Equivalently  $X^6$  has positive constant scalar curvature.

# 7 Concluding remarks

We have solved the KSEs for all IIB horizons that admit at least one supersymmetry. This has been done by first integrating the KSEs along the lightcone directions and then identifying the independent equations using the Bianchi identities, field equations and bilinear matching condition. Then the independent KSEs are solved using spinorial geometry which leads to three different cases, the generic horizons, the Spin(7) horizons and the pure SU(4) horizons. We have found that the requirement of IIB horizons to admit one supersymmetry puts rather weak geometric restrictions on the horizon sections S. In particular for generic horizons and pure SU(4) horizons, the horizon sections can be any 8-dimensional almost Hermitian spin<sub>c</sub> manifold. A similar result also applies for Spin(7) horizons. We have also described some topological aspects of the horizon sections and how the various fluxes are expressed in terms of the geometry.

We have also explained how horizons with only 5-form fluxes and common sector horizons, which had been investigated previously, are included in our analysis. As a result all the examples constructed in these two special cases can be embedded in the full IIB theory. Furthermore, we give some examples for which, in addition to the KSEs, we also solve the field equations and Bianchi identities. In particular, we focus on horizons with constant scalars which have complex horizon sections, and we find a generalization of the 2-SCYT structure which had appeared for horizons with 5-form fluxes only.

The construction of examples with non-trivial scalars taking values on the upper halfplane after a  $SL(2, \mathbb{R})$  identification is natural within the context of 10-dimensional type of F-theory [31]. In particular, one may consider  $T^2$ -fibrations over S. However there are some differences. As we have mentioned, S are almost complex manifolds instead of Kähler, which mostly arise in the context of F-theory. In addition, the example of horizons with constant scalars that we have explicitly constructed indicates that S is not Kähler because of the presence of form fluxes. Nevertheless in the complex case, it may be possible to construct examples imitating techniques that have been employed in F-theory.

In most of our considerations, like the solution of the KSEs and the description of the geometry of the horizon sections, we have not used the compactness of  $\mathcal{S}$ . So our results apply to both black holes and brane horizons. For applications to black holes, it would be of interest to enforce compactness of  $\mathcal{S}$ . This has been done for M-horizons in [32], and after an application of the index theorem for the Dirac operator, it has led to the conclusion that all M-horizons preserve an even number of supersymmetries and admit an  $\mathfrak{sl}(2,\mathbb{R})$  symmetry. A similar application may be possible in IIB. However, there are some differences between M-theory and IIB. One of them is that unlike the M-horizon sections which are odd dimensional, the IIB near-horizon sections are even dimensional and so the index of the Dirac operator is not expected to vanish. As the vanishing of the Dirac index has been instrumental in proving supersymmetry enhancement for M-horizons, a similar application in IIB will require some modification. Nevertheless, it is expected that even if one cannot prove supersymmetry enhancement for IIB black hole horizons, it may be possible to relate the number of supersymmetries preserved in terms of the index of a Dirac operator on the horizon sections. Such a relation will generalize the classic formula N = index(D) which relates the number of parallel spinors N on irreducible holonomy Spin(7), SU(4) and Sp(2) 8-dimensional manifolds, for N = 1, 2 and 3 respectively, to the index of the Dirac operator. In turn such a formula will provide a topological criterion for 8-dimensional manifolds to admit Killing spinors.

### Acknowledgements

UG is supported by the Knut and Alice Wallenberg Foundation. JG is supported by the STFC grant, ST/1004874/1. GP is partially supported by the STFC rolling grant ST/J002798/1.

# Appendix A Solution of the linear system

# A.1 The linear system

It is straightforward to derive the linear system associated with the KSE (3.19) evaluated on the spinor  $\eta_+ = p1 + qe_{1234}$ , where p, q are real functions on S and  $p^2 + q^2 = 1$ . Substituting this spinor into (3.19), one obtains,

$$\partial_{\alpha}p + p\left(-\frac{i}{2}\Lambda_{\alpha} - \frac{1}{4}h_{\alpha} + \frac{1}{2}\Omega_{\alpha,\mu}{}^{\mu} - iY_{\alpha\mu}{}^{\mu}\right) + q\left(\frac{1}{4}H_{\alpha\mu}{}^{\mu} - \frac{1}{4}\Phi_{\alpha}\right) = 0 , \qquad (A.1)$$

$$p\left(\frac{1}{2}\Omega_{\alpha,\bar{\mu}_{1}\bar{\mu}_{2}} - iY_{\alpha\bar{\mu}_{1}\bar{\mu}_{2}} + i\delta_{\alpha[\bar{\mu}_{1}}Y_{\bar{\mu}_{2}]\lambda}^{\lambda} + \frac{1}{24}\delta_{\alpha[\bar{\mu}_{1}}\epsilon_{\bar{\mu}_{2}}^{\lambda} + \frac{1}{\lambda_{1}\lambda_{2}\lambda_{3}}H_{\lambda_{1}\lambda_{2}\lambda_{3}}\right) + q\left(-\frac{1}{4}\Omega_{\alpha,\lambda_{1}\lambda_{2}}\epsilon^{\lambda_{1}\lambda_{2}}_{\mu_{1}\bar{\mu}_{2}} + \frac{1}{4}H_{\alpha\bar{\mu}_{1}\bar{\mu}_{2}} - \frac{1}{8}\delta_{\alpha[\bar{\mu}_{1}}H_{\bar{\mu}_{2}]\lambda}^{\lambda} + \frac{1}{8}\delta_{\alpha[\bar{\mu}_{1}}\Phi_{\bar{\mu}_{2}}]\right) = 0 , \qquad (A.2)$$

$$\partial_{\alpha}q + q\left(-\frac{i}{2}\Lambda_{\alpha} - \frac{1}{4}h_{\alpha} - \frac{1}{2}\Omega_{\alpha,\mu}^{\mu} + \frac{1}{24}H_{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}}\epsilon^{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}}\alpha\right) + p\left(-\frac{1}{8}H_{\alpha\mu}^{\mu} - \frac{i}{3}Y_{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}}\epsilon^{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}}\alpha - \frac{1}{8}\Phi_{\alpha}\right) = 0 , \qquad (A.3)$$

$$\partial_{\bar{\alpha}}p + p\left(-\frac{i}{2}\Lambda_{\bar{\alpha}} - \frac{1}{4}h_{\bar{\alpha}} + \frac{1}{2}\Omega_{\bar{\alpha},\mu}^{\mu} + \frac{1}{24}H_{\lambda_{1}\lambda_{2}\lambda_{3}}\epsilon^{\lambda_{1}\lambda_{2}\lambda_{3}}\bar{\alpha}\right) + q\left(\frac{1}{8}H_{\bar{\alpha}\mu}^{\mu} - \frac{i}{3}Y_{\lambda_{1}\lambda_{2}\lambda_{3}}\epsilon^{\lambda_{1}\lambda_{2}\lambda_{3}}\bar{\alpha} - \frac{1}{8}\Phi_{\bar{\alpha}}\right) = 0 , \qquad (A.4)$$

$$p\left(\frac{1}{2}\Omega_{\bar{\alpha},\bar{\mu}_{1}\bar{\mu}_{2}} - \frac{1}{8}H_{\bar{\alpha}\lambda_{1}\lambda_{2}}\epsilon^{\lambda_{1}\lambda_{2}}_{\bar{\mu}_{1}\bar{\mu}_{2}} + \frac{1}{16}\epsilon^{\lambda}{}_{\bar{\alpha}\bar{\mu}_{1}\bar{\mu}_{2}}H_{\lambda\mu}{}^{\mu} + \frac{1}{16}\epsilon^{\lambda}{}_{\bar{\alpha}\bar{\mu}_{1}\bar{\mu}_{2}}\Phi_{\lambda}\right) + q\left(-\frac{1}{4}\Omega_{\bar{\alpha},\lambda_{1}\lambda_{2}}\epsilon^{\lambda_{1}\lambda_{2}}_{\bar{\mu}_{1}\bar{\mu}_{2}} + \frac{i}{2}Y_{\bar{\alpha}\lambda_{1}\lambda_{2}}\epsilon^{\lambda_{1}\lambda_{2}}_{\bar{\mu}_{1}\bar{\mu}_{2}} + \frac{1}{8}H_{\bar{\alpha}\bar{\mu}_{1}\bar{\mu}_{2}} - \frac{i}{2}\epsilon^{\lambda}{}_{\bar{\alpha}\bar{\mu}_{1}\bar{\mu}_{2}}Y_{\lambda\mu}{}^{\mu}\right) = 0, \quad (A.5)$$

$$\partial_{\bar{\alpha}}q + q\left(-\frac{i}{2}\Lambda_{\bar{\alpha}} - \frac{1}{4}h_{\bar{\alpha}} - \frac{1}{2}\Omega_{\bar{\alpha},\mu}{}^{\mu} + iY_{\bar{\alpha}\mu}{}^{\mu}\right) + p\left(-\frac{1}{4}H_{\bar{\alpha}\mu}{}^{\mu} - \frac{1}{4}\Phi_{\bar{\alpha}}\right) = 0.$$
(A.6)

Similarly, the linear system associated with (3.21) is

$$\frac{1}{12}p\epsilon_{\alpha}{}^{\bar{\gamma}_1\bar{\gamma}_2\bar{\gamma}_3}H_{\bar{\gamma}_1\bar{\gamma}_2\bar{\gamma}_3} - \frac{1}{4}qH_{\alpha\gamma}{}^{\gamma} - \frac{1}{4}q\Phi_{\alpha} + p\xi_{\alpha} = 0 , \qquad (A.7)$$

$$\frac{1}{12}q\epsilon_{\bar{\alpha}}{}^{\gamma_1\gamma_2\gamma_3}H_{\gamma_1\gamma_2\gamma_3} + \frac{1}{4}pH_{\bar{\alpha}\gamma}{}^{\gamma} - \frac{1}{4}p\Phi_{\bar{\alpha}} + q\xi_{\bar{\alpha}} = 0.$$
(A.8)

The above system can be solved to express some of the fluxes in terms of the geometry, and find the conditions on the geometry imposed by supersymmetry. In the analysis of the solutions, it is convenient to consider three different cases as described in section 4.1.

## A.2 Solution of the linear system

### A.2.1 Generic Horizons

The solution of the linear system can be arranged in different ways. The procedure which we adopt here is to solve first for the complex field strengths and express them in terms of the real fields and functions p, q, and then use the remaining equations to find the expression of the real fields in terms of the geometry, and to determine the conditions on the geometry. We shall demonstrate that, although the KSEs determine the complex fields in terms of the real fields and geometry, the real fields remain undetermined.

To solve the linear system, first recall that in the generic case  $p^2 - q^2 \neq 0$  and  $p^2 + q^2 = 1$ , and let us assume that in some open set  $p, q \neq 0$ . If either p, q vanish in an open set then the linear system will be solved as a special case. Next we take the trace of (A.2), and after a re-arrangement, that of (A.5) to find

$$q\left(-\Omega_{\bar{\lambda},\alpha}^{\bar{\lambda}} + iY_{\alpha\lambda}^{\lambda} - \frac{1}{8}\epsilon_{\alpha}^{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}}H_{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}}\right) + p\left(-\frac{1}{2}\epsilon_{\alpha}^{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}}\Omega_{\bar{\lambda}_{1},\bar{\lambda}_{2}\bar{\lambda}_{3}} + \frac{1}{8}H_{\alpha\lambda}^{\lambda} - \frac{3}{8}\Phi_{\alpha}\right) = 0 ,$$
(A.9)

$$p\left(\Omega_{\lambda,\bar{\alpha}}^{\lambda} + iY_{\bar{\alpha}\lambda}^{\lambda} + \frac{1}{8}\epsilon_{\bar{\alpha}}^{\lambda_{1}\lambda_{2}\lambda_{3}}H_{\lambda_{1}\lambda_{2}\lambda_{3}}\right) + q\left(\frac{1}{2}\epsilon_{\bar{\alpha}}^{\lambda_{1}\lambda_{2}\lambda_{3}}\Omega_{\lambda_{1},\lambda_{2}\lambda_{3}} + \frac{1}{8}H_{\bar{\alpha}\lambda}^{\lambda} + \frac{3}{8}\Phi_{\bar{\alpha}}\right) = 0 ,$$
(A.10)

respectively.

Next consider the equations (A.3) and (A.7), and (A.4) and (A.8) and solve them to express the components of the 3-form field strength H in terms of the rest of the fields yielding

$$\frac{1}{8}H_{\alpha\lambda}{}^{\lambda} = -\frac{1}{8}\Phi_{\alpha} + \frac{pq}{2}\xi_{\alpha} + p\partial_{\alpha}q - pq[\frac{i}{2}\Lambda_{\alpha} + \frac{1}{4}h_{\alpha} + \frac{1}{2}\Omega_{\alpha,\lambda}{}^{\lambda}] + \frac{i}{3}p^{2}\epsilon_{\alpha}{}^{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}}Y_{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}} ,$$
  
$$\frac{1}{24}\epsilon_{\alpha}{}^{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}}H_{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}} = -\frac{p^{2}}{2}\xi_{\alpha} + q\partial_{\alpha}q - q^{2}[\frac{i}{2}\Lambda_{\alpha} + \frac{1}{4}h_{\alpha} + \frac{1}{2}\Omega_{\alpha,\lambda}{}^{\lambda}] + \frac{i}{3}qp\epsilon_{\alpha}{}^{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}}Y_{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}} ,$$
  
$$(A.11)$$

$$\frac{1}{8}H_{\bar{\alpha}\lambda}{}^{\lambda} = \frac{1}{8}\Phi_{\bar{\alpha}} - \frac{pq}{2}\xi_{\bar{\alpha}} - q\partial_{\bar{\alpha}}p - pq[-\frac{i}{2}\Lambda_{\bar{\alpha}} - \frac{1}{4}h_{\bar{\alpha}} + \frac{1}{2}\Omega_{\bar{\alpha},\lambda}{}^{\lambda}] - \frac{i}{3}q^{2}\epsilon_{\bar{\alpha}}{}^{\lambda_{1}\lambda_{2}\lambda_{3}}Y_{\lambda_{1}\lambda_{2}\lambda_{3}} ,$$
  
$$\frac{1}{24}\epsilon_{\bar{\alpha}}{}^{\lambda_{1}\lambda_{2}\lambda_{3}}H_{\lambda_{1}\lambda_{2}\lambda_{3}} = -\frac{q^{2}}{2}\xi_{\bar{\alpha}} + p\partial_{\bar{\alpha}}p + p^{2}[-\frac{i}{2}\Lambda_{\bar{\alpha}} - \frac{1}{4}h_{\bar{\alpha}} + \frac{1}{2}\Omega_{\bar{\alpha},\lambda}{}^{\lambda}] + \frac{i}{3}qp\epsilon_{\bar{\alpha}}{}^{\lambda_{1}\lambda_{2}\lambda_{3}}Y_{\lambda_{1}\lambda_{2}\lambda_{3}} .$$
  
(A.12)

It remains to solve the trace conditions (A.9) and (A.10) and (A.1) and (A.6) of the linear system in terms of  $\phi$  and  $\xi$ . After some straightforward computation, one finds that

$$pq\Phi_{\alpha} = \partial_{\alpha}q^{2} - (2+q^{2})(i\Lambda_{\alpha} + \frac{1}{2}h_{\alpha}) - (-2+5q^{2})\Omega_{\alpha,\lambda}^{\lambda} - 2i(2-q^{2})Y_{\alpha\lambda}^{\lambda} + 2q^{2}\Omega_{\bar{\lambda},\bar{\lambda}_{\alpha}}^{\bar{\lambda}_{\alpha}} + 2ipq\epsilon_{\alpha}^{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}}Y_{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}} + pq\epsilon_{\alpha}^{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}}\Omega_{\bar{\lambda}_{1},\bar{\lambda}_{2}\bar{\lambda}_{3}} ,$$

$$p^{2}q^{2}\xi_{\alpha} = q^{2}\partial_{\alpha}q^{2} - (1+2q^{4})[\frac{i}{2}\Lambda_{\alpha} + \frac{1}{4}h_{\alpha}] - \frac{1}{2}(2q^{4} + 2q^{2} - 1)\Omega_{\alpha,\lambda}^{\lambda} - iY_{\alpha\lambda}^{\lambda} + q^{2}\Omega_{\bar{\lambda},\bar{\lambda}_{\alpha}} + pq(1+2q^{2})\frac{i}{3}\epsilon_{\alpha}^{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}}Y_{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}} + \frac{pq}{2}\epsilon_{\alpha}^{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}}\Omega_{\bar{\lambda}_{1},\bar{\lambda}_{2}\bar{\lambda}_{3}} ,$$

$$pq\Phi_{\bar{\alpha}} = \partial_{\bar{\alpha}}p^{2} - (2+p^{2})[i\Lambda_{\bar{\alpha}} + \frac{1}{2}h_{\bar{\alpha}}] + (-2+5p^{2})\Omega_{\bar{\alpha},\lambda}^{\lambda} + 2i(2-p^{2})Y_{\bar{\alpha}\lambda}^{\lambda} + 2p^{2}\Omega_{\lambda,\bar{\lambda}_{\alpha}} + 2ipq\epsilon_{\bar{\alpha}}^{\bar{\lambda}_{1}\lambda_{2}\lambda_{3}}Y_{\lambda_{1}\lambda_{2}\lambda_{3}} + pq\epsilon_{\bar{\alpha}}^{\bar{\lambda}_{1}\lambda_{2}\lambda_{3}}\Omega_{\lambda_{1},\lambda_{2}\lambda_{3}} ,$$

$$p^{2}q^{2}\xi_{\bar{\alpha}} = p^{2}\partial_{\bar{\alpha}}p^{2} - (1+2p^{4})[\frac{i}{2}\Lambda_{\bar{\alpha}} + \frac{1}{4}h_{\bar{\alpha}}] + \frac{1}{2}(2p^{4} + 2p^{2} - 1)\Omega_{\bar{\alpha},\lambda}^{\lambda} + iY_{\bar{\alpha}\lambda}^{\lambda} + p^{2}\Omega_{\lambda,\bar{\lambda}_{\bar{\alpha}}} + pq(1+2p^{2})\frac{i}{3}\epsilon_{\bar{\alpha}}^{\bar{\lambda}_{1}\lambda_{2}\lambda_{3}}Y_{\lambda_{1}\lambda_{2}\lambda_{3}} + \frac{pq}{2}\epsilon_{\bar{\alpha}}^{\bar{\lambda}_{1}\lambda_{2}\lambda_{3}}\Omega_{\lambda_{1},\lambda_{2}\lambda_{3}} .$$
(A.13)

Clearly, we can substitute the  $\Phi$ 's and the  $\xi$ 's into (A.11) and (A.12) to express the traces and (3,0) and (0,3) part of H in terms of the the Y fluxes and the geometry. However, we shall not do this here as for our conclusions this is not utilized.

Next using the expressions in (A.11) and (A.12), one can determine the (2,1) and (1,2) components of H as

$$\frac{p}{4}H_{\bar{\alpha}\mu_{1}\mu_{2}} = q\left[-\frac{1}{2}\Omega_{\bar{\alpha},\mu_{1}\mu_{2}} + iY_{\bar{\alpha}\mu_{1}\mu_{2}} + i\delta_{\bar{\alpha}[\mu_{1}}Y_{\mu_{2}]\lambda}^{\lambda}\right] + \frac{p}{4}\Omega_{\bar{\alpha},\bar{\nu}_{1}\bar{\nu}_{2}}\epsilon^{\bar{\nu}_{1}\bar{\nu}_{2}}{}_{\mu_{1}\mu_{2}} \\
-\delta_{\bar{\alpha}[\mu_{1}}\left[\partial_{\mu_{2}}\right]q - q\left(\frac{i}{2}\Lambda_{\mu_{2}}\right] + \frac{1}{4}h_{\mu_{2}}\right] + \frac{1}{2}\Omega_{\mu_{2}],\lambda}^{\lambda}\right) + \frac{i}{3}p\epsilon_{\mu_{2}}^{\lambda}\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}}Y_{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}}\right], \\
\frac{q}{4}H_{\alpha\bar{\mu}_{1}\bar{\mu}_{2}} = -p\left[\frac{1}{2}\Omega_{\alpha,\bar{\mu}_{1}\bar{\mu}_{2}} - iY_{\alpha\bar{\mu}_{1}\bar{\mu}_{2}} + i\delta_{\alpha[\bar{\mu}_{1}}Y_{\bar{\mu}_{2}]\lambda}^{\lambda}\right] + \frac{q}{4}\Omega_{\alpha,\nu_{1}\nu_{2}}\epsilon^{\nu_{1}\nu_{2}}{}_{\bar{\mu}_{1}\bar{\mu}_{2}} \\
-\delta_{\alpha[\bar{\mu}_{1}}\left[\partial_{\bar{\mu}_{2}}\right]p + p\left(-\frac{i}{2}\Lambda_{\bar{\mu}_{2}}\right] - \frac{1}{4}h_{\bar{\mu}_{2}}\right] + \frac{1}{2}\Omega_{\bar{\mu}_{2}],\lambda}^{\lambda}\right) + \frac{iq}{3}\epsilon_{\bar{\mu}_{2}}^{\lambda_{1}\lambda_{2}\lambda_{3}}Y_{\lambda_{1}\lambda_{2}\lambda_{3}}\right]. \tag{A.14}$$

Note that the expression for h in (3.10) is not independent, and so if one substitutes the solution for the fluxes given above, one gets identically zero. Conversely, any one of the solutions above can be used to express h in terms of the fluxes and geometry.

Furthermore, one can use the solution of the linear system to determine  $\mu$  in (3.13) in terms of the Y fluxes and geometry as

$$\sqrt{2} pq\mu_{\bar{\alpha}} = -\frac{i}{2}\Lambda_{\bar{\alpha}} + \frac{1}{4}(p^2 - q^2)h_{\bar{\alpha}} + \frac{1}{2}(p^2 - q^2)\Omega_{\bar{\alpha},\lambda}^{\lambda} + iY_{\bar{\alpha}\lambda}^{\lambda} + \frac{2i}{3}pq\epsilon_{\bar{\alpha}}^{\lambda_1\lambda_2\lambda_3}Y_{\lambda_1\lambda_2\lambda_3} .$$
(A.15)

Observe that the spin connection  $\tilde{\Omega}$  of the horizon sections  $\mathcal{S}$ , which is given by  $\tilde{\Omega}_{i,k}^{j} = \Omega_{i,k}^{j}$ , is not restricted. This indicates that there are no additional restrictions on the topology and geometry of  $\mathcal{S}$  apart from those required for the global existence of certain forms which we describe in section 4.2. This concludes the solution of the KSEs for the generic case.

The solution of the linear system for Spin(7) horizons can be derived from that of the generic case after setting  $p = q = 1/\sqrt{2}$ . This is straightforward to implement and we shall not carry out the substitution here.

## A.2.2 Pure SU(4) horizons

For pure SU(4) horizons, one has either p = 1, q = 0 or p = 0, q = 1. The two cases are symmetric and without loss of generality, we choose p = 1, q = 0. A direct computation reveals that the solution can be written as

$$\begin{split} \Phi_{\bar{\alpha}} &= H_{\bar{\alpha}\lambda}{}^{\lambda} = 0 , \quad \frac{1}{12} \epsilon_{\alpha}{}^{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}} H_{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}} + \xi_{\alpha} = 0 , \\ \Phi_{\alpha} &= -\epsilon_{\alpha}{}^{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}} \Omega_{\bar{\lambda}_{1},\bar{\lambda}_{2}\bar{\lambda}_{3}} + \frac{2i}{3} \epsilon_{\alpha}{}^{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}} Y_{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}} , \\ H_{\alpha\lambda}{}^{\lambda} &= e_{\alpha}{}^{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}} \Omega_{\bar{\lambda}_{1},\bar{\lambda}_{2}\bar{\lambda}_{3}} + 2i\epsilon_{\alpha}{}^{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}} Y_{\bar{\lambda}_{1}\bar{\lambda}_{2}\bar{\lambda}_{3}} \\ & \frac{1}{8} \epsilon_{\bar{\alpha}}{}^{\lambda_{1}\lambda_{2}\lambda_{3}} \Omega_{\bar{\lambda}_{1},\bar{\lambda}_{2}\bar{\lambda}_{3}} = -\Omega_{\lambda}{}^{\lambda}{}_{\bar{\alpha}} - iY_{\bar{\alpha}\lambda}{}^{\lambda} , \\ \frac{1}{2} H_{\bar{\alpha}\lambda_{1}\lambda_{2}} &= \frac{1}{2} \Omega_{\bar{\alpha},\bar{\nu}_{1}\bar{\nu}_{2}} \epsilon^{\bar{\nu}_{1}\bar{\nu}_{2}} \lambda_{1\lambda_{2}} - \frac{2i}{3} \delta_{\bar{\alpha}}[\lambda_{1}} \epsilon_{\lambda_{2}}]^{\bar{\nu}_{1}\bar{\nu}_{2}\bar{\nu}_{3}} Y_{\bar{\nu}_{1}\bar{\nu}_{2}\bar{\nu}_{3}} , \\ \cdot iY_{\alpha\bar{\lambda}_{1}\bar{\lambda}_{2}} &+ \frac{2i}{3} \delta_{\alpha}[\bar{\lambda}_{1}}Y_{\bar{\lambda}_{2}]\mu^{\mu}} + \frac{1}{2} \Omega_{\alpha,\bar{\lambda}_{1}\bar{\lambda}_{2}} - \frac{1}{3} \delta_{\alpha}[\bar{\lambda}_{1}\Omega_{|\mu|},{}^{\mu}\bar{\lambda}_{2}] = 0 , \\ &- \frac{1}{2} h_{\bar{\alpha}} + \frac{1}{3} \Omega_{\lambda}{}^{\lambda}{}_{\bar{\alpha}}} - \frac{2i}{3} Y_{\bar{\alpha}\lambda}{}^{\lambda} = 0 , \\ &i\Lambda_{\bar{\alpha}}} + h_{\bar{\alpha}} - \Omega_{\bar{\alpha},\lambda}{}^{\lambda} - \Omega_{\lambda}{}^{\lambda}{}_{\bar{\alpha}}} = 0 . \end{split}$$
(A.16)

Moreover, one finds that

$$\sqrt{2}\,\mu_{\bar{\alpha}} = \frac{1}{2}\epsilon_{\bar{\alpha}}{}^{\lambda_1\lambda_2\lambda_3}\Omega_{\lambda_1,\lambda_2\lambda_3} + \frac{i}{3}\epsilon_{\bar{\alpha}}{}^{\lambda_1\lambda_2\lambda_3}Y_{\lambda_1\lambda_2\lambda_3} \,. \tag{A.17}$$

Again notice that there is no condition which involves only the spin connection  $\hat{\Omega}$  of the horizon section S. This indicates that there are no additional conditions on the topology and geometry of the horizon sections apart from those required for the global existence of certain forms which we describe in section 4.4.

# A.3 Spinor conventions

For our spinor conventions, we use those of [19]. In addition to integrate the KSEs along the lightcone directions, we have decomposed the Spin(9,1) spinors  $\psi$  into positive and negative parts as

$$\psi = \psi_+ + \psi_-, \qquad \Gamma_\pm \psi_\pm = 0 , \qquad (A.18)$$

which is a lightcone, or equivalently Spin(1,1), chiral decomposition. Furthermore, we have found the following identities useful

$$\Gamma_{\ell_1...\ell_n}\psi_{\pm} = \pm (-1)^{[\frac{n}{2}]} \frac{1}{(8-n)!} \epsilon_{\ell_1...\ell_n}{}^{j_1...j_{8-n}} \Gamma_{j_1...j_{8-n}}\psi_{\pm} , \quad n \ge 4 , \qquad (A.19)$$

where  $\psi$  is a positive chirality Spin(9,1) spinor. While, one has

$$\Gamma_{\ell_1...\ell_n}\psi_{\pm} = \mp (-1)^{[\frac{n}{2}]} \frac{1}{(8-n)!} \epsilon_{\ell_1...\ell_n}{}^{j_1...j_{8-n}} \Gamma_{j_1...j_{8-n}}\psi_{\pm} , \quad n \ge 4 , \qquad (A.20)$$

provided that  $\psi$  is a negative chirality Spin(9,1) spinor.

# References

- J. Isenberg and V. Moncrief, "Symmetries of cosmological Cauchy horizons," Commun. Math. Phys. 89 (1983) 387.
- [2] H. Friedrich, I. Racz and R. M. Wald, "On the rigidity theorem for space-times with a stationary event horizon or a compact Cauchy horizon," Commun. Math. Phys. 204 (1999) 69; [gr-qc/9811021].
- [3] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, "Large N field theories, string theory and gravity," Phys. Rept. 323 (2000) 183; [hep-th/9905111].
- [4] W. Israel, "Event Horizons In Static Vacuum Space-Times," Phys. Rev. 164 (1967) 1776.
- [5] B. Carter, "Axisymmetric Black Hole Has Only Two Degrees of Freedom," Phys. Rev. Lett. 26 (1971) 331.
- [6] S. W. Hawking, "Black holes in general relativity," Commun. Math. Phys. 25 (1972) 152.
- [7] D. C. Robinson, "Uniqueness of the Kerr black hole," Phys. Rev. Lett. 34 (1975) 905.
- [8] W. Israel, "Event Horizons in Static, Electrovac Space-Times," Commun. Math. Phys. 8 (1968) 245.
- [9] P. O. Mazur, "Proof of Uniqueness of the Kerr-Newman Black Hole Solution," J. Phys. A 15 (1982) 3173.
- [10] D. Robinson, "Four decades of black hole uniqueness theorems," appeared in *The Kerr spacetime: Rotating black holes in General Relativity*, eds D. L. Wiltshire, M. Visser and S. M. Scott, pp 115-143, CUP 2009.
- [11] G. W. Gibbons, D. Ida and T. Shiromizu, "Uniqueness and non-uniqueness of static black holes in higher dimensions", Phys. Rev. Lett. 89 (2002) 041101; [hepth/0206049].
- [12] M. Rogatko, "Uniqueness theorem of static degenerate and non-degenerate charged black holes in higher dimensions", Phys. Rev. D67 (2003) 084025; [hep-th/0302091];
  "Classification of static charged black holes in higher dimensions," Phys. Rev. D73 (2006), 124027; [hep-th/0606116].

- [13] P. Figueras and J. Lucietti, "On the uniqueness of extremal vacuum black holes," Class. Quant. Grav. 27 (2010) 095001; [arXiv:0906.5565 [hep-th]].
- [14] R. Emparan, T. Harmark, V. Niarchos and N. Obers, "World-Volume Effective Theory for Higher-Dimensional Black Holes," Phys. Rev. Lett. **102** (2009) 191301;
   [arXiv:0902.0427 [hep-th]]; "Essentials of Blackfold Dynamics," JHEP **03** (2010) 063;
   [arXiv:0910.1601 [hep-th]].
- [15] H. S. Reall, "Higher dimensional black holes and supersymmetry", Phys. Rev. D68 (2003) 024024; [hep-th/0211290].
- [16] H. Elvang, R. Emparan, D. Mateos and H. S. Reall, "A Supersymmetric black ring," Phys. Rev. Lett. 93 (2004) 211302; [hep-th/0407065].
- [17] J. Gutowski and G. Papadopoulos, "M-Horizons," JHEP 12 (2012) 100; [arXiv:1207.7086 [hep-th]].
  J. Gutowski and G. Papadopoulos, "Static M-horizons," JHEP 01 (2012) 005; [arXiv:1106.3085 [hep-th]].
- [18] J. Gutowski and G. Papadopoulos, "Heterotic Black Horizons," JHEP 07 (2010) 011; [arXiv:0912.3472 [hep-th]].
- [19] U. Gran, J. Gutowski and G. Papadopoulos, "The spinorial geometry of supersymmetric IIB backgrounds," Class. Quant. Grav. 22 (2005) 2453; [hep-th/0501177].
  U. Gran, J. Gutowski and G. Papadopoulos, "The G<sub>2</sub> spinorial geometry of supersymmetric IIB backgrounds," Class. Quant. Grav. 23 (2006) 143; [hep-th/0505074].
  U. Gran, J. Gutowski, G. Papadopoulos and D. Roest, "Systematics of IIB spinorial geometry," Class. Quant. Grav. 23 (2006) 1617; [hep-th/0507087].
- [20] U. Gran, J. Gutowski and G. Papadopoulos, "IIB black hole horizons with five-form flux and KT geometry," JHEP 05 (2011) 050; [arXiv:1101.1247 [hep-th]].
  U. Gran, J. Gutowski and G. Papadopoulos, "IIB backgrounds with five-form flux," Nucl. Phys. B 798 (2008) 36 [arXiv:0705.2208 [hep-th]].
- [21] J. Gillard, U. Gran and G. Papadopoulos, "The Spinorial geometry of supersymmetric backgrounds," Class. Quant. Grav. 22 (2005) 1033; [hep-th/0410155].
- [22] J. H. Schwarz, P. C. West and , "Symmetries and Transformations of Chiral N=2 D=10 Supergravity," Phys. Lett. B 126 (1983) 301.
- [23] J. H. Schwarz, "Covariant Field Equations of Chiral N=2 D=10 Supergravity," Nucl. Phys. B 226 (1983) 269.
- [24] G. Papadopoulos and D. Tsimpis, "The Holonomy of IIB supercovariant connection," Class. Quant. Grav. 20 (2003) L253; [hep-th/0307127].
- [25] B. R. Greene, A. D. Shapere, C. Vafa and S. -T. Yau, "Stringy Cosmic Strings and Noncompact Calabi-Yau Manifolds," Nucl. Phys. B 337 (1990) 1.

- [26] C. J. Isham, C. N. Pope, "Nowhere Vanishing Spinors And Topological Obstructions To The Equivalence Of The Nsr And Gs Superstrings," Class. Quant. Grav. 5 (1988) 257.
- [27] J. W. Milnor and J. D. Stasheff, "Characteristic Classes," Princeton Univ. Press (1974).
- [28] K. Dasgupta, G. Rajesh, S. Sethi, "M theory, orientifolds and G flux," JHEP 08 (1999) 023; [hep-th/9908088].
- [29] E. Goldstein, S. Prokushkin, "Geometric model for complex nonKahler manifolds with SU(3) structure," Commun. Math. Phys. 251 (2004) 65; [hep-th/0212307].
- [30] D. Grantcharov, G. Grantcharov, Y. S. Poon, "Calabi-Yau connections with torsion on toric bundles," J. Diff. Geom 78 (2008) 13; [math.DG/0306207].
- [31] C. Vafa, "Evidence for F theory," Nucl. Phys. B 469 (1996) 403; [hep-th/9602022].
- [32] J. Grover, J. B. Gutowski, G. Papadopoulos and W. A. Sabra, "Index Theory and Supersymmetry of 5D Horizons;" [arXiv:1303.0853 [hep-th]].

J. B. Gutowski and G. Papadopoulos, "Index theory and dynamical symmetry enhancement of M-horizons;" [arXiv:1303.0869 [hep-th]].